Robust cone-mediated signaling persists late into rod

photoreceptor degeneration

Miranda L. Scalabrino[1], Mishek Thapa[1], Lindsey A. Chew[1], Esther Zhang[1], Jason Xu[2],

A.P. Sampath[3], Jeannie Chen[4], Greg D. Field[1*]

1. Department of Neurobiology, Duke University School of Medicine, Durham, NC

2. Department of Statistical Science, Duke University, Durham NC

3. Jules Stein Eye Institute, University of California Los Angeles, Los Angeles, CA.

4. Zilkha Neurogenetics Institute, Keck School of Medicine, University of Southern

California, Los Angeles, CA.

*Corresponding author: Greg D. Field, field@neuro.duke.edu

Abstract

Rod photoreceptor degeneration causes deterioration in the morphology and physiology

of cone photoreceptors along with changes in retinal circuits. These changes could

diminish visual signaling at cone-mediated light levels, thereby limiting the efficacy of

treatments such as gene therapy for rescuing normal, cone-mediated vision. However,

the impact of progressive rod death on cone-mediated signaling remains unclear. A

mouse model of rod degeneration was used to investigate the fidelity of retinal ganglion

cell (RGC) signaling throughout disease progression. Despite clear deterioration of cone

morphology with rod death, cone-mediated signaling among RGCs remained surprisingly

robust: spatiotemporal receptive fields changed little and the mutual information between

stimuli and spiking responses was relatively constant. This relative stability held until

nearly all rods had died and cones had completely lost well-formed outer segments.

Interestingly, RGC information rates were higher and more stable for natural movies than

checkerboard noise as degeneration progressed. The main change in RGC responses

with photoreceptor degeneration was a decrease in response gain. These results suggest

that gene therapies for rod degenerative diseases are likely to successfully prolong cone-
mediated vision even if there are changes to cone morphology and density.

Introduction

Rod photoreceptor degeneration frequently leads to cone photoreceptor
degeneration and death. Well before all rods have died, cones exhibit clear changes in
morphology and physiology (Hartong, Berson, & Dryja, 2006; Sahel, Bonnel, Mrejen, &
Paques, 2010). This is likely due to the loss of trophic factors and structural support
provided by rods to nearby cones (Campochiaro & Mir, 2018). The extent to which these
changes in cone structure and function compromise the ability of the retina to reliably
signal visual scenes at high light levels is not clear. One possibility is that rod
degeneration has an immediate impact on the ability of the retina to reliably signal visual
scenes at cone-mediated light levels. Alternatively, recent work has highlighted
homeostatic mechanisms in retinal circuits that help compensate for photoreceptor loss
(Care et al., 2020; Lee, Care, Santina, & Dunn, 2021; Shen, Wang, Soto, &
Kerschensteiner, 2020). Such mechanisms could facilitate reliable signaling at the level
of retinal output, despite deterioration in photoreceptor function. Resolving these
possibilities will inform treatment timepoints for gene therapies aimed at halting rod loss
to preserve cone-mediated vision.

To examine the impact of progressive rod loss on cone-mediated visual signaling,
we used a mouse line that models a human form of retinitis pigmentosa (RP), a blinding
disorder characterized by initial degeneration of rod photoreceptors that ultimately leads
to cone degeneration. This mouse line, $Cngb1^{neo/neo}$, contains an insertion that interrupts
translation of Cngb1, the beta subunit of the cyclic nucleotide gated cation channel in rods
(Chen et al., 2010). Without this subunit, normal channels fail to form, causing the rods
to be tonically hyperpolarized and ultimately resulting in rod death (Hüttl et al., 2005;
Zhang et al., 2009). This degeneration is relatively slow, with approximately 30% rod loss
at 1 month (M) postnatal, complete loss of rods by 7M postnatal, and complete cone loss
by 8-9M postnatal. Slow degeneration in this model provides a relatively large temporal
window in which to assay changes in retinal signaling to increasing amounts of rod loss
and accompanying changes in cone morphology and density. Slower forms of RP are

also more common among the human population (Grover et al., 1999; Hartong et al.,
2006), and thus slow degeneration models may be more therapeutically informative than
animal models with relatively rapid photoreceptor degeneration (e.g., *rd1* and *rd10*).

To determine the changes in cone-mediated retinal signaling induced by
progressive rod loss, we measured changes in the signaling of retinal ganglion cells
(RGCs), the 'output' neurons of the retina. Deterioration in the fidelity of RGC signals
captures net changes in retinal circuit function induced by rod degeneration: these include
changes in retinal circuits that compensate and/or exacerbate deteriorating photoreceptor
function. Large-scale multielectrode arrays (MEAs) were used to measure the visual
responses of hundreds of RGCs simultaneously in individual retinas while presenting a
variety of visual stimuli: e.g., checkerboard noise and natural movies. Three features of
RGC signaling were specifically investigated. First, we measured when and how the
spatiotemporal receptive fields (RFs) under cone-mediated conditions were altered by
rod degeneration. These RFs are indicative of the visual features that are being signaled
by the RGCs to the brain (Chichilnisky, 2001; Keat, Reinagel, Reid, & Meister, 2001; Yu,
Grzywacz, Lee, & Field, 2017), and thus changes in these RFs represent changes in the
kind of information being transmitted to the brain. Second, we probed how and when the
spontaneous activity of RGCs was altered by rod degeneration. Many previous studies
have noted the emergence of strong oscillatory activity in the spontaneous activity of
RGCs in animal models of RP. Such spontaneous activity could disrupt the ability of the
retina to encode stimuli. However, only *rd1* and *rd10* mice have been used to study when
these oscillations emerge relative to rod and cone photoreceptor death. These models
exhibit significantly more rapid photoreceptor degeneration than *Cngb1^{neo/neo}* mice and
may not be ideal for capturing aspects of human disease (Stasheff, 2008; Stasheff,
Shankar, & Andrews, 2011). Third, we explored when and how rod degeneration impacts
the fidelity of visual signals transmitted to the brain. Information theory was used to
quantify changes in the fidelity of signaling naturalistic and artificial stimuli as a function
of photoreceptor degeneration.

From 1 to 7M of age in *Cngb1^{neo/neo}* mice, there were marked and clear changes
in cone morphology and density resulting from rod degeneration. When assaying the
spatiotemporal RFs of RGCs as a function of degeneration, there were subtle changes to

the temporal RFs of RGCs as cone morphology changed. However, the spatiotemporal

RFs of RGCs were remarkably stable until the latest stages of degeneration (5-7M). The

primary change to cone-mediated RGC responses was a decrease in response gain as

photoreceptors were lost. Second, oscillations in the spontaneous activity of RGCs did

not emerge in this mouse model until after all rods and nearly all cones were dead (~9M).

This indicates that oscillatory activity emerges only in the very late states of photoreceptor

degenerative diseases, at least in animal models with relatively slow forms of retinal

degeneration. Third, as rods died and normal cone morphology deteriorated, the ability

of the retina to reliably signal the content of natural movies -- under cone-mediated

conditions -- was remarkably robust until the latest stages of degeneration (~7M; total rod

loss with severe cone morphology changes and death). Thus, the fidelity of information

transmission was relatively stable despite a reduction in response gain. This suggests

that homeostatic mechanisms in the retina serve to compensate for deteriorating

photoreceptors. Furthermore, these results suggest a broad therapeutic window in which

to treat RP, as cone-mediated visual signaling remains surprisingly robust despite clear

changes in cone morphology and density.

Results

Cngb1$^{neo/neo}$ **cones slowly degenerate lagging rod death**

Photoreceptors in *Cngb1$^{neo/neo}$* mice gradually die over a period of 7 to 8M as a

consequence of the *Cngb1* mutation (Figure 1A). While rods die first, shown by the

shrinking outer nuclear layer (ONL) over time, cones are present until late in disease

(Figure 1B). However, cone structure begins to deteriorate at 3M (Figure 1C, middle).

Cone outer segments gradually shorten and eventually disappear prior to total cell death

(Figure 1D). By 7M, nearly all rods are lost and only cone cell bodies remain; clear outer

segment structure is absent (Figure 1C, right). How does this deterioration in cone

morphology, and potential changes in cone function impact the ability of the retina to

transmit visual information to the brain? Answering this question is particularly important

given that humans primarily use cone vision for most tasks. One possibility is that cone-

mediated signaling at the level of the RGCs is robust to changes in cone morphology.

Alternatively, RGC signaling at cone-mediated light levels may rapidly deteriorate as rods

are lost and cone morphology becomes abnormal.

Figure 1. Cone morphology and density changes with rod degeneration. *A. Estimated fraction of surviving rods (black) and cones (cyan) relative to WT densities from 1 to 7 months of age in* Cngb1[neo/neo] *mice. B. Immunofluorescence of retinal cross sections in WT and* Cngb1[neo/neo] *mice (1-7 months). C. Cone morphologies imaged with cone arrestin (mCar) labeling in WT and* Cngb1[neo/neo] *mice. D. Whole mount view of cone density and morphology in WT and* Cngb1[neo/neo] *mice. Scale bars are 50 µm. ONL: outer nuclear layer; INL: inner nuclear layer; OS: outer segments.*

Identifying RGCs with space-time separable receptive fields

To determine the impact of changing cone morphology and density due to rod

death on retinal output, we began by measuring the RFs of RGCs from wild-type (WT)

and *Cngb1[neo/neo]* retinas using MEAs consisting of 512 or 519 electrodes (see Methods)

(Anishchenko et al., 2010; Field et al., 2010; Litke et al., 2004). The MEAs measured the

spiking activity in 186-560 RGCs in individual samples of retina from each animal (see

Methods). Animals were used at 1M intervals from 1-7M of age (post-natal). To estimate

the spatiotemporal RFs of RGCs, we used checkerboard noise and computed the spike-

triggered average stimulus (STA) for each RGC (Chichilnisky, 2001).

We measured RF structure at two light levels: a low mesopic light level (100

Rh*/rod/s) at which cones are just beginning to be activated, and a low photopic light level

(10,000 Rh*/rod/s). We chose the lower light level because we expected to see significant

changes in visual signaling between WT and *Cngb1^neo/neo^* animals as the latter has

severely compromised rod function even in surviving rods (Wang et al., 2019). We chose

the higher light level because it largely, if not completely, isolated cone-mediated

signaling, thereby allowing us to examine the net impact of rod dysfunction and death on

cone-mediated retinal output.

The STA estimates the linear component of each RF. We were particularly

interested in changes to the spatial or temporal integration of visual input, which are

Figure 2. Receptive field decomposition. *A. Example space-time separable STA from an RGC. Top is a cut through one spatial dimension and bottom is a cut through the orthogonal spatial dimension. D, V, A, and P indicate the dorsal, ventral, anterior, and posterior directions of the retina. B. Singular value decomposition (SVD) of STA in A: (left) distribution of variance explained by first six space-time vector pairs; (top right) spatial (left) and temporal (right) filters from the rank-one decomposition; (bottom right) spatial and temporal filters associated with the second singular value. C. Example STA from an RGC that was not space-time separable. D. SVD of STA in C; (left) distribution of variance explained by first six space-time vector pairs; (top right) spatial and temporal filters from the rank-one decomposition; (bottom right) spatial and temporal filters associated with the second singular value E. Distribution of variance explained by the rank-one decomposition for all WT RGCs (1654 cells from 5 mice) under mesopic (teal) and photopic (orange) conditions. Vertical line shows threshold for classifying cells as space-time separable. F. Fraction and quantity of RGCs measured at mesopic (teal) and photopic (orange) light levels that were classified as exhibiting space-time separable RFs.*

estimated by changes in the spatial and temporal RFs, respectively. However, some
RGC types (i.e., direction-selective RGCs) do not have a RF that can be decomposed
into unique spatial or temporal filters (Borghuis, Ratliff, Smith, Sterling, &
Balasubramanian, 2008; Devries & Baylor, 1997; Vaney, Sivyer, & Taylor, 2012), which
makes it an ill-posed problem to separately analyze changes in spatial or temporal
integration. Thus, we focused our analysis on RGCs with RFs that were well-
approximated by a space-time separable RF model: the outer product of two vectors, one
describing the spatial RF and the other the temporal RF (Figure 2) (see Methods).
Singular value decomposition (SVD) factorizes the STA into a linear combination of
spatial and temporal filter pairs: the outer product of these pairs, weighted by their
associated singular values, will reproduce the original STA (Golomb, Kleinfeld, Reid,
Shapley, & Shraiman, 1994; Wolfe & Palmer, 1998). A space-time separable STA will
yield one pair of spatial and temporal filters that captures most of the variance in the STA
(a.k.a., rank-1 approximation Figure 2A-B). Subsequent space-time filter pairs will exhibit
little to no structure and appear as 'noise' (Figure 2B, bottom-right). In contrast, non-
separable STAs will require multiple space-time vector pairs to reproduce the STA (Figure
2C-D) and each pair will exhibit clear structure in space and time (Figure 2D). RGCs for
which > 60% of the variance in the full space-time STA was captured by a separable RF
model were included in the analysis. This threshold was chosen because it captured a
clear mode in the distribution of all RGCs from WT retinas (Figure 2E). Across time-
points of degeneration, 8,028 out of 12,997 RGCs met this criterion. However, the fraction
passing this criterion changed as a function of degeneration, with more cells passing the
criterion early in degeneration and fewer passing at the latest timepoints (Figure 2F).
RGCs passing this criterion had a wide range of spatial RF sizes and temporal RF
durations and consisted of both ON and OFF RGC types. Thus, several, but not all, RGC
types met this criterion (see Discussion). Below we analyze changes in the spatial and
temporal RFs of RGCs at mesopic and photopic light levels as a function of photoreceptor
degeneration.
168
Changes in mesopic receptive fields with photoreceptor degeneration

At the mesopic light level, the distribution of spatial RF sizes between WT and 1M

Cngb1^neo/neo animals was relatively stable (Figure 3A-B). A small, but statistically

significant decrease in the mean relative to WT was observed at 1-4M (15% difference

between WT to 4M; p-value: 0.039). At 5M, the decrease was attenuated. By 7M, no

RGCs exhibited space-time separable RFs at the mesopic light level (Figure 2F) and only

35% of identified RGCs exhibited visual responses (see Methods). Note that corrupting a

rank- one STA with increasing amounts of noise will cause the rank-one approximation

to capture increasingly less total variance.

At mesopic conditions, we observed larger fractional changes in the temporal

integration of RGCs than in their spatial integration between WT and *Cngb1^neo/neo* mice

Figure 3. Changes in RF structure under mesopic conditions induced by rod death. *A. Example spatial RFs at mesopic light level showing smaller to larger RFs (left to right). B. Distributions of spatial RFs comparing WT and* Cngb1^neo/neo *mice from 1-5M of degeneration. C. Example temporal RFs at mesopic light level showing briefer to longer temporal integration. Grey arrows indicate the time to zero of RGC 1. Blue arrows on RGC 2 and RGC 3 show time to zero for RGCs 2 & 3. D. Distributions of temporal RF comparing WT and* Cngb1^neo/neo *mice from 1-5M of degeneration. E. Example STAs with lower and higher signal-to-noise (SNR) ratios. F. Distribution of SNRs comparing WT and* Cngb1^neo/neo *mice.*

(Figure 3C-D). At 1-3M the mean temporal integration and the variance was larger for

Cngb1^{neo/neo} animals than WT: mean of all RGCs increased by 14%, 17%, and 13% at

1M, 2M, and 3M respectively (p-value : 0.10, 0.08, 0.11) and the variance of all RGCs

increased by 38%, 34%, and 35% at 1, 2, and 3M of age respectively (p-value: 0.01,

0.02, 0.02). At 4M, the median temporal integration increased further by 88% relative to

WT (p-value: 0.001). However, at 5M the temporal integration decreased back toward

WT (9% increase in time-to-zero relative to WT; p-value: 0.14), with the variance in the

distribution substantially larger (37% increase in variance relative to WT; p-value:

0.009). Overall, spatial RF structure was relatively stable until the latest stages of

degeneration: largest change from WT in spatial integration was 15% (at 4M). Changes

in temporal integration were larger with the largest change being 88% (4M), but these

changes fluctuated over time and were non-monotonic with degeneration.

Given that photoreceptors are dying, it is potentially surprising that spatiotemporal

RF structure is relatively stable under mesopic conditions. Fewer photoreceptors ought

to result in diminished sensitivity, even if the area of spatial integration or the duration of

temporal integration are relatively stable. Thus, we analyzed the signal-to-noise ratio

(SNR) of the STAs which may be expected to be noisier with photoreceptor death. The

SNR of the STAs generally drifted down over time (Figure 3E-F); however, even these

changes were relatively small until 5M (compared to WT, there was a 5% and 15% decline

in SNR in the 4M and 5M experimental groups, respectively; p-value: 0.19 and 0.04). The

STA measurements were based on 30 min of checkerboard stimuli, which may be a

sufficiently long period of time to average away noise and/or compensate for diminished

gain. To more directly assess changes in response gain, we inspected the contrast

response functions (also called 'static nonlinearities') across RGCs, which capture the

relationship between the stimulus (filtered by the RF) and the spiking output of the RGC

(Chichilnisky, 2001). The gain of these contrast response functions steadily decreased as

a function of degeneration (Figure 4). The effect was most noticeable for stimuli that were

the most similar to the RF (Figure 4C-E). Thus, under mesopic conditions, gain steadily

decreased with photoreceptor degeneration, while spatial and temporal integration were

relatively stable until the latest stages of degeneration.

Figure 4. Gain of contrast response functions steadily decreases during RP. *A. Mean cumulative Gaussian fit to contrast response function across RGCs from WT and 7M* Cngb1[neo/neo] *mice. Four locations along the contrast response functions are highlighted by blue diamonds. B-E, Distribution of contrast response function values at contrast values indicated in A for WT and 1-7M retinas. Note, this analysis was not restricted to RGCs with space-time separable RFs, allowing the analysis to extend to 7M.*

Changes in photopic receptive fields with photoreceptor degeneration

Under photopic conditions, the distribution of spatial RF sizes was remarkably

stable throughout degeneration (Figure 5A-B). The largest change was at 7M with a 5%

decrease in spatial RF area relative to WT (p-value: 0.211). Note, unlike the mesopic

condition, RFs were detectable out to 7M under the photopic condition.

Temporal RFs exhibited greater changes than spatial RFs under photopic

conditions (Figure 5C-D) but were still surprisingly stable. Specifically, the time to zero

progressively decreased 10%, 12%, and 15% from 1-3M, respectively (relative to WT; p-

value: 0.32, 0.22, 0.04), indicating a shortening of temporal integration. At 4M the

temporal integration slowed somewhat, becoming 22% slower than 3M but only 4%

slower than WT (p-value: 0.423) (Figure 5D). At 5M and 7M, time to zero decreased by

12% and 13% (p-value: 0.22, 0.12), respectively, relative to 4M.

These changes in temporal and spatial RF structure under photopic conditions

were relatively small given the clear changes in cone morphology observed particularly

at 5 to 7 M (Figure 1): the largest deviation in mean temporal integration from WT occurred

Figure 5. Changes in RF structure under photopic conditions induced by rod death. *A. Example spatial RFs at the photopic light level showing small to larger RFs (left to right). B. Distributions of spatial RFs comparing WT and* Cngb1$^{neo/neo}$ *mice from 1-7M of degeneration. C. Example temporal RFs at photopic light level. Orange arrows indicates the time to zero for RGC 1; Red arrows show time to zero for RGCs 2 and 3. D. Distributions of temporal RFs comparing WT and* Cngb1$^{neo/neo}$ *mice from 1-7M of degeneration. E. Example STAs with lower and higher signal-to-noise (SNR) ratios. F. Distribution of SNRs comparing WT and* Cngb1$^{neo/neo}$ *mice.*

at 3Ms, and this was only a 15% slowing in the time-to-zero, while spatial RF sizes

changed by 5% at most (relative to WT) through 7M.

We next analyzed the SNR of the STAs (Figure 5E-F). We observed a decline in

SNR with degeneration under photopic conditions. Yet even at 7M, with 30% cone loss,

near complete rod loss, and abnormal cone morphologies, there was only a 6% decrease

in the median SNR (Figure 5F; p-value: 0.456). As with the mesopic conditions,

computing STAs over relatively long periods of checkerboard noise may mask decreases

in gain or increases in response noise. To inspect changes in gain, we analyzed the

contrast response functions (Figure 6). Like the mesopic condition, response gain steadily

decreased with degeneration, particularly for stimuli that increasingly matched the RF

(Figure 6D-E). Thus, the dominant effect of photoreceptor degeneration on photopic RF

Figure 6. Gain of the contrast response functions steadily decreases during RP under photopic conditions. *A. Mean cumulative Gaussian fit to contrast response function across RGCs from WT and 7M* Cngb1neo/neo *mice. Four locations along the contrast response functions are highlighted by diamonds. B-E, Distribution of contrast response function values at contrast values indicated in A for WT and 1-7M retinas.*

structure among RGCs in *Cngb1neo/neo* mice was a decrease in response gain, not a

change in spatial or temporal integration.

***Cngb1neo/neo* RGCs do not exhibit oscillatory activity while photoreceptors remain**

The results above indicate relatively small changes in spatial and temporal

integration of cone mediated responses as rods die and cones degenerate in *Cngb1neo/neo*

mice. They also indicate clear decreases in gain as a function of degeneration under

mesopic and photopic conditions (Figures 4 and 6). However, the analyses above do not

provide much insight into the extent that noise may be changing with degeneration. One

source of noise is signal-independent increases in spontaneous activity. Several mouse

models of retinal degeneration exhibit increased spontaneous activity arising as 5-10 Hz

oscillations in RGC spiking (Marc et al., 2007; Margolis, Newkirk, Euler, & Detwiler, 2008;

Stasheff, 2008). These oscillations are reported to occur throughout the degeneration

process (Stasheff et al., 2011). The presence of such oscillations may minimally impact

the RF measurements as it would be largely averaged away when computing the STA,

Figure 7. Retinal oscillations occur after vision loss in Cngb1[neo/neo] *mice. A. Spontaneous activity of three representative WT RGCs in total darkness. B. The power spectral density (PSD) of one example RGC. C-D. Example rasters (C) and PSD (D) of a representative 3M* Cngb1[neo/neo] *RGC. E-F. Example rasters (left) and the PSD of an example 9M* Cngb1[neo/neo] *RGC. G. Distributions of the PSD fluctuation ratio, defined as the maximum of PSD divided by the baseline PSD (see Methods). Mean indicated by orange diamond, gray bars indicate 100%, 80%, and 50% sections of the distribution around the median.*

but they could contribute to the lower SNR of the STAs (Figure 3F and 5F) along with the

decreased gain (Figures 4 and 6).

Thus, we analyzed the spontaneous activity of RGCs in WT and *Cngb1^{neo/neo}*

animals from 1-9M to check for oscillatory activity (Figure 7). Spontaneous activity was

measured in darkness as well as with static illumination set to the same mean intensity

as the checkerboard stimuli presented at the mesopic and photopic light levels.

Oscillations were absent in the spiking activity of all recorded RGCs in darkness and at

both mesopic and photopic light levels in *Cngb1^{neo/neo}* retinas from 1 – 7M (Figure 7A-D,

G). However, clear 5 Hz oscillatory activity did arise at 9M (Figure 7E-G). The oscillations

were present at all three light levels (not shown). At this last stage of degeneration, no

light response could be detected with 100% contrast flashes at the mesopic or photopic

light levels (not shown). Thus, oscillations in the spontaneous spiking activity of RGCs in

the *Cngb1^neo/neo^* mouse model are absent until all (or nearly-all) of the RGCs have lost

their light responses.

268 RGCs from *Cngb1^neo/neo^* retinas exhibit deteriorated visual signaling under mesopic

269 conditions

Thus far, we have observed a decrease in response gain among RGCs under

mesopic and photopic conditions that tracks progressive photoreceptor degeneration.

However, signal-independent noise, as assayed by changes in spontaneous activity,

appear constant until all (or nearly all) the photoreceptors have died in *Cngb1^neo/neo^* mice.

However, it is possible that the RGCs are signaling less reliably as photoreceptors die:

the STA method averages over many trials and is thus relatively insensitive to potential

changes in signal-dependent noise that could manifest as increased variability in either

the number or timing of spikes produced to a given stimulus. Because of this, it is

Figure 8. RGC signaling fidelity under mesopic conditions decreases with photoreceptor degeneration. *A. (Top) Rasters from three example RGC responding to a repeated mesopic checkerboard stimulus. (Bottom) distribution of information rates across all RGCs from 1 experiment; teal lines show where each example RGC lies in the distribution. B. Distributions of information rates of for all RGCs in each condition: WT and* Cngb1^neo/neo^ *mice. Mean shown by teal diamond, stars indicate significant changes from WT: ** and *** are p < 0.01 and 0.001, respectively. C. Distribution of information rates of 10% most informative RGCs across conditions: WT and* Cngb1^neo/neo^ *mice.*

important to identify potential changes in the fidelity of RGC signaling that may
accompany photoreceptor degeneration.

To measure the fidelity of RGC signaling, we used information theory (Shannon,
1948). Specifically, we calculated the mutual information between spike train and the
stimulus (see Methods). Mutual information indicates how much the uncertainty about the
visual stimulus is reduced by observing the spike train of an RGC (see review by
McDonnell, Ikeda, & Manton, 2011). If the RGC response is highly variable, observing a
single response will reduce uncertainty about the stimulus less than for an RGC with
highly reproducible responses.

We began by presenting a repeating checkerboard stimulus at the mesopic light
level used for the RF measurements above (~100 Rh*/ rod/ s). We segregated RGCs
based on their information rate for the checkerboard stimulus (Figure 8). Some RGCs
exhibited strongly modulated and reproducible responses across repetitions of the
stimulus (Figure 8A, RGC 1), while other RGCs were less reliably driven by the stimulus
(Figure 8A, RGC 2 & 3). Note all RGCs with a spike rate above 3 Hz (12,997 RGCs) were
used in this analysis: we included RGCs with and without space-time separable RFs.

For WT retinas, the distribution of information rates across RGCs was
approximately unimodal with a long tail skewed toward high rates (Figure 8A, bottom).
The peak of these distributions did not change much with degeneration (Figure 8B).
However, the peak primarily represented cells that were not particularly reliably driven by
the checkerboard stimulus even in WT retinas; in other words, these were not the cells
that were most informative about the stimulus. In contrast, the median of the information
rate distributions decreased as photoreceptors died (Figure 8B, teal diamonds). This
change in the median was largely driven by decreased information rates in the long tails
of these distributions (Figure 8B). We thus focused our analysis on the 10% of RGCs that
were most informative about the checkerboard stimulus: those with the highest
information rates (Figure 8C). For these RGCs, there was a clear drop in information rates
between WT and 1M *Cngb1^{neo/neo}* mice (mean of all RGCs changed from 5.7 bits/s to 4.3
bits/s; p-value: 0.003). In fact, *Cngb1^{neo/neo}* RGCs at all time points examined transmitted
significantly less information than WT RGCs at the mesopic light level (mean of all RGCs
ranged from 2.1-4.3 bits/s). Information rates between 1 and 2M did not change

significantly (difference of 0.09 bits/second was observed; p-value: 0.655), but there was

a steady decline at subsequent timepoints.

These data indicate that under low mesopic conditions, there is a relatively steady

decline in the reliability with which RGCs signal visual information in *Cngb1neo/neo* mice.

The difference between WT and 1-2M *Cngb1neo/neo* animals is likely because rods are

poorly responsive to light in *Cngb1neo/neo* animals due to a greatly reduced photocurrent

(Wang et al., 2019). The progressive decline in information rates after 2M is likely caused

by large-scale loss of rods (Figure 1A), and possibly a reduced ability of cones to signal

near their threshold for activation.

RGCs from *Cngb1neo/neo* retinas exhibit relatively stable visual signaling under

photopic conditions.

We next checked the reliability of RGC signaling under photopic conditions that

are well above the threshold of cone vision (10,000 R*/rod/s). It is possible that at higher

light levels, visual signaling is more (or less) stable. Similar to the lower light level, RGCs

Figure 9. RGC signaling fidelity under photopic conditions is relatively stable with rod degeneration. *A. (Top) Rasters from three example RGC responding to a repeated photopic checkerboard stimulus. (Bottom) distribution of information rates across all RGCs from 1 experiment; teal lines show where each example RGC lies in the distribution. B. Distributions of information rates of for all RGCs in each condition: WT and* Cngb1neo/neo *mice. Mean shown by teal diamond, stars designate significant changes from WT: ** and *** are p < 0.01 and 0.001, respectively. C. Distribution of information rates of 10% most informative RGCs across conditions: WT and* Cngb1neo/neo *mice.*

exhibited a wide range of information rates to checkerboard stimuli, with some exhibiting
very reliable responses to a repeated stimulus while others exhibited more variable
responses (Figure 9A). Information rates were much more stable among RGCs at these
higher light levels, with no significant differences in median information rates observed
between WT and 1M to 3Ms of degeneration (p-values of 0.56, 0.32, 0.29 between WT
and 1M, 2M, and 3M, respectively) (Figure 9B). Even when focusing on the top 10% of
most informative RGCs, 1 and 2M of degeneration were not significantly different from
WT (Figure 9C). Despite clear changes in cone morphology at 5Ms (Figure 1), decreases
in information rates were modest (mean of top 10% of RGCs changed from 23.1
bits/second in WT to 18.1 bits/s at 5M; p-value: 0.0009). At 7M, cone-mediated signaling
among the RGCs continued to transmit information, but with a relatively pronounced
decrease: mean rate was 8.6 bits/s among the top 10% of RGCs. Overall, these data
indicate that the fidelity of cone-mediated visual signaling among RGCs is quite stable
until ~2-3M of degeneration, corresponding to 50-70% rod loss.

Degenerating retina more robustly encodes natural than artificial stimuli

Given the wide range of information rates observed across RGCs for checkerboard
stimuli, we hypothesized that the results described above may depend on stimulus
choice. Thus, we repeated the experiments under photopic conditions (10,000 Rh*/rod/s)
using 2 natural movies (see Methods). As with the checkerboard stimuli, RGCs again
demonstrated a wide range of information rates when presented with the natural movies
(Figure 10A). However, for natural movies, the median information rates were much more
stable as photoreceptor degeneration progressed (Figure 10B): relative to WT, there was
only a 5% decline in MI at 4M; p-value: 0.543). Similarly, the RGCs with the highest
information rates (top 10%) did not show the same decline in information rate as observed
with the checkboard stimuli (Figure 10C): from WT to 4M, there was only a 9.5% decline
in information rate (p-value: 0.321).

Finally, we inspected how the information rates depended on the spike rates.
Lower information rates could be caused by stimuli producing fewer spikes in
degenerating retinas, and stimuli induced fewer spikes at later stages of degeneration
(Figure 6). Thus, we compared the information rate to the firing rate for checkerboard

Figure 10. **Cngb1**^{neo/neo} ***RGCs signal more information about natural movies than checkerboard movies.*** *A. (Top) Rasters from three example RGCs responding to a repeated photopic natural stimulus. (Bottom) distribution of information rates across all RGCs from 1 experiment; green lines show where each example RGC lies in the distribution. B. Distributions of information rates of for all RGCs in each condition for WT and* Cngb1^{neo/neo} *mice. Mean shown by green diamond. C. Distribution of information rates of 10% most informative RGCs across conditions: WT and* Cngb1^{neo/neo} *mice. D. Scatter plot of information rates and spike rates from RGC responses to photopic checkerboard stimulus (left) or natural stimuli (right). Grey dots are RGC responses from the total population, while black dots are specific to 7M* Cngb1^{neo/neo} *mutual responses to a checkerboard stimuli and green dots 7M* Cngb1^{neo/neo} *responses to a natural movie. E. (left) Mean +/- 2 standard error of information rates of RGC responses to natural movies (green), photopic checkerboard (open), and mesopic checkerboard (black) across WT and* Cngb1^{neo/neo} *mice. (right) Mean +/- 2 standard error of spike rates of RGC responses to natural movies (green), photopic checkerboard movies (open), and mesopic checkerboard (black) across WT and* Cngb1^{neo/neo} *mice. ρ is the linear correlation between the mean MI and the mean firing rate across retinas.*

stimuli and the natural movies (Figure 10D). When comparing information rates across

all RGCs at all timepoints, there is a clear correlation between spike rate and information

rate for the checkerboard stimuli and natural movies (Figure 10D, gray dots). However,

at 7M for checkerboard stimuli (Figure 10D, black dots), RGCs with the highest firing rates
exhibited suppressed information rates relative to RGCs with similar firing rates from
retinas with less degeneration. For natural movies, RGCs from 7M retinas exhibited much
higher information rates across a similar range of firing rates observed in checkerboard
stimuli (Figure 10D). Inspecting across degeneration timepoints, there was a sharp drop
in the information rates for mesopic checkerboard stimuli from WT to 1M *Cngb1^{neo/neo}*
animals (10% most informative cells) that was not present under photopic conditions
(Figure 10E). Under photopic conditions, information transmission was higher among the
10% most informative cells for natural movies than for checkerboard stimuli (Figure 10E).
These changes in information rates were not strongly correlated with the changes in spike
rates across conditions (Figure 10F, $\rho = 0.13$), suggesting that they are not simply a result
of changing spike rates.

Discussion

Determining the extent to which rod dysfunction, degeneration, and death impacts
cone-mediated visual signaling in the retina is important for diagnosing and treating RP.
It is possible that cone-mediated signals in RGCs, the 'output' neurons of the retina,
rapidly deteriorate with the loss of rods and concomitant changes in cone morphology.
Alternatively, it is conceivable that RGC signaling is robust to photoreceptor degeneration.
We have measured cone-mediated visual responses from thousands of RGCs from
Cngb1^{neo/neo} mice as rod and cone photoreceptors degenerate. We have spanned early
timepoints at which most rods remain and there is little to no cone death or changes in
cone morphology, to time points at which all rods are lost and cones exhibit abnormal
morphologies and are beginning to die. We find that despite clear changes in cone
morphology and density, RF structure and signal fidelity remain relatively stable until the
latest stages of degeneration. In this mouse line, oscillatory spontaneous activity among
RGCs did not arise until all or nearly all photoreceptors were lost in contrast to previous
results from *rd1* and *rd10* mice. These results suggest that early rescue of rods from
degeneration is likely to preserve relatively normal cone-mediated visual signaling among
RGCs.

Comparison to previous studies of RGC signaling in retinitis pigmentosa

Few studies track cone-mediated RGC signaling at many timepoints across RP. Focusing on RGC signaling has the advantage that it includes degradation in visual signaling caused by the degeneration of photoreceptors as well as downstream changes in retinal circuitry. Thus, it captures net changes in retinal processing that deteriorate signaling along with homeostatic mechanisms that may serve to preserve signaling (Care et al., 2020, 2019; Leinonen et al., 2020; Shen et al., 2020). In addition, examining cone-mediated vision, instead of rod vision, focuses on changes that are most likely to be relevant for humans. We did not distinguish among RGC types, other than identifying those with space-time separable RFs for the spatial and temporal RF analyses (Figures 2-6). This choice was largely because at 5M-7M it became increasingly difficult to reliably distinguish cell types, and at the earlier time points we did not observe systematic differences across RGCs types.

Previous work in a *P23H* rat model of RP indicated a monotonic decrease in RF size, a monotonic increase in the duration of temporal integration, and a rise in spontaneous activity as photoreceptors died (Sekirnjak et al., 2011). In contrast, using *Cngb1$^{neo/neo}$* mice we observed small shifts in RF size and temporal duration during degeneration, but these changes were not monotonic with disease progression. The only feature of visual signaling that we found to change monotonically was response gain (Figures 4 & 6). Both studies found that cone-mediated responses persisted longer than rod-mediated responses, which is unsurprising given that cones typically survive longer than rods in RP. We speculate that the differences between these studies arise from degeneration having different causes in *P23H* rats versus *Cngb1$^{neo/neo}$* mice. The rat model involves a mutation in rhodopsin which causes issues with protein folding and trafficking that are not thought to initially change the dark-current of rods or their resting glutamate release (Jones & Marc, 2005; Liu, Garriga, & Khorana, 1996; Machida et al., 2000; Sakami et al., 2011). However, the *Cngb1$^{neo/neo}$* rods exhibit a reduced dark current, are tonically hyperpolarized, and likely release less glutamate onto post-synaptic bipolar cells (Wang et al., 2019). These differences in rod physiology and how rods signal to downstream circuits are likely to have consequences for how the retina responds to rod

death. Fully understanding how different origins of rod degeneration impact retinal circuits
and RGC signaling is an important direction for future RP studies.

A novel feature of this study is the use of information theory to assay the reliability
of visual signaling during retinal degeneration (Figures 8-10). Information theory has been
used extensively in the vertebrate and invertebrate visual systems, as well as other
sensory systems to quantify encoding performance of sensory neurons (Fairhall et al.,
2006; K. Koch et al., 2006; Rieke, Bodnar, & Bialek, 1995; Van Steveninck, Lewen,
Strong, Koberle, & Bialek, 1997). In general, sensory neurons with more reliable and
selective responses exhibit higher information rates, measured in bits per second.
Information theory has not been used previously to assay the effects of neural
degeneration on sensory coding. Information theory has the advantage in that it provides
a way of comparing the signal fidelity across neurons that transmit information about very
different sorts of features (e.g., color vs. motion). It has the potential disadvantages that
it is a data-hungry analysis and can be prone to certain biases (Paninski, 2003). The
relatively long and stable nature of MEA experiments allowed us to overcome these
limitations (Figure 11). Consistent with previous results, we found a broad range of
information rates across the population of RGCs (K. Koch et al., 2004, 2006), even in
control (WT) retinas. What was perhaps surprising is how stable these information rates
were under photopic conditions until those latest stages of degeneration. This indicates
that while gain diminishes with degeneration (Figures 4 & 6), the precision and reliability
of RGC spiking remains relatively stable (Figures 8-10). Interestingly, this reliability of
signaling was more robust when using natural stimuli compared to checkerboard noise
(Figure 10). This difference may be the result of natural stimuli having spatiotemporal
correlations that are not present in checkerboard noise, and homeostatic mechanisms
that preserve retinal signaling during degeneration can potentially exploit these
correlations for higher fidelity signaling.

Visual compensation for cell loss begins in the retina

Human patients with RP45 (*Cngb1* mutations) maintain cone vision for many
years, but the assumption has been that vision was preserved primarily through cortical
compensation (Ferreira et al., 2017; Hickmott & Merzenich, 2002; Keck et al., 2013, 2008,

2011; Merabet & Pascual-Leone, 2010; Turrigiano, 2012). Indeed, studies in primary visual cortex using the *S334ter* rat model of RP support the notion that cortical plasticity bolsters visual processing during retinal degeneration (Chen et al., 2016). However, our findings indicate retinal signaling remains relatively robust under cone-mediated conditions despite large-scale rod loss, changes in cone morphology, and cone density. This suggests that at least some of the plasticity and homeostatic mechanisms required for vision during photoreceptor degeneration occur in the retina.

While we did not investigate mechanisms underlying this robust signaling, there is evidence that late-stage degenerated retinas maintain some photoreceptor to bipolar cell synapses (Beier et al., 2017; Beier, Palanker, & Sher, 2018; Care et al., 2019; Leinonen et al., 2020; Shen et al., 2020). Additionally, cone-mediated light responses are enhanced in a model with 50% rod loss compared to controls (Care et al., 2020). Mechanisms like these may help to explain why cone-mediated signaling was relatively stable until the latest stages of degeneration.

Oscillatory spontaneous activity appears late in *Cngb1^{neo/neo}* mice

A prominent feature of RP models is the emergence of abnormal spontaneous activity among RGCs (Trenholm & Awatramani, 2015). Previous studies on other models of RP, particularly *rd1* and *rd10* mouse models of Pde6b-RP, have shown abnormal spontaneous activity in two frequency bands: 1-2 Hz and 5-10 Hz (Biswas et al., 2014; Marc et al., 2007; Margolis et al., 2008; Menzler & Zeck, 2011; Poria & Dhingra, 2015; Stasheff et al., 2011; Tu, Chen, McQuiston, Chiao, & Chen, 2016; Ye & Goo, 2007). Abnormal horizontal cell activity is the source of the 1-2 Hz oscillations (Haq, Arango-Gonzalez, Zrenner, Euler, & Schubert, 2014), while AII amacrine cells are the likely source of the higher frequency oscillations (Borowska, Trenholm, & Awatramani, 2011; Choi et al., 2014; Ivanova, Yee, Baldoni, & Sagdullaev, 2016). Elevated and/or oscillatory spontaneous activity is a cause for concern in retinal degeneration because it is noise that competes with and deteriorates visual signals, particularly near threshold. It is unclear whether RGC oscillations also develop in humans with RP, although reports of phosphenes in RP patients could be explained by spontaneous activity (Gekeler, Messias, Ottinger, Ulrich Bartz-Schmidt, & Zrenner, 2006).

While previous studies have observed the emergence of oscillations in rodent

models of RP (e.g., *rd10*) prior to the loss of photoreceptors and visual signaling, we only

found evidence of spontaneous activity following the near total loss of photoreceptors in

Cngb1^{neo/neo} mice (9M, Figure 7). At 9M, we could not elicit visual responses from retinas

when the video display was switched from off (0 Rh*/rod/s) to on with a 'white' screen

(20,000 Rh*/rod/s). Furthermore, at 9M, there was no clear outer nuclear layer and cells

that labeled for cone arrestin were very sparse and dysmorphic and did not co-label for

M-opsin, indicating near total photoreceptor loss (data not shown). Even at 7M, when

nearly all the rods and ~30% of the cones have died, we observed no evidence for

changes in spontaneous activity or oscillations. This is a promising observation from the

perspective of gene therapy for rescuing photoreceptors: it suggests that in conditions

like RP45, elevated spontaneous activity and oscillation in RGC spiking will be avoided

given that gene therapy is delivered prior to total cell loss.

Implications for therapy

Our findings on the longevity of cone vision are particularly useful in determining

the time window for therapeutic intervention in patients with RP45. Both pre-clinical and

clinical studies have shown that late intervention does not ultimately halt photoreceptor

cell death despite improvements to visually guided behaviors (Bainbridge et al., 2015;

Cideciyan et al., 2013; Gardiner et al., 2020; Jacobson et al., 2015; S. Koch et al., 2012).

It is currently unclear if the continued degeneration is slowed by therapy or not, or what

the long-term implications are for vision restoration. Our study suggests that preserving

cone-mediated vision in RP is entirely feasible considering that cone-mediated signals

among RGCs are minimally perturbed by massive rod loss. However, it is not clear that

rod-directed gene therapy will be successful at preserving cone (or rod) vision if it is

delivered at a timepoint after which most of the rods have died. Thus, an important

direction for future work is to determine at what timepoints rod-directed gene therapies

need to be delivered to halt rod death such that cone vision can continue to function

normally throughout the remaining lifespan of the treated individual.

Materials and Methods

Mice

Mice were used according to Duke University Institutional Animal Care and Use Committee guidelines and the Association for Research in Vision and Ophthalmology guidelines for the use of animals in vision research. All mice were housed with 12h light/dark cycles and fed rodent chow ad libitum. $Cngb1^{neo/neo}$ mice of both sexes were used (12 male and 12 female). The $Cngb1^{neo/neo}$ line has a neomycin resistance cassette inserted at intron 19 to disrupt Cngb1 mRNA splicing (Chen et al., 2010; Wang et al., 2019). Control animals (WT) consisted of male and female heterozygous littermates between 1 and 9M postnatal age to account for aging (4 male and 3 female). No age-related differences were found between WT mice (data not shown), so results were pooled. Genotyping was performed by Transnetyx using the primers/probe for the neomycin insert FWD GGGCGCCCGGTTCTT, REV CCTCGTCCTGCAGTTCATTCA, PROBE ACCTGTCCGGTGCCC, and primers/probe for WT Cngb1 FWD TCCTTAGGCTCTGCTGGAAGA, REV CAGAGGATGAACAAGAGACAGGAA, PROBE CTGAGCTGGGTAATGTC. At least 3 mice were used per timepoint, except to assay spontaneous activity at 9M, in which 2 mice were used (Figure 7).

Immunohistochemistry and Confocal Microscopy

The sample of retina placed on the MEA and retina from the contralateral (unrecorded) eye were fixed for 30 minutes in 4% PFA (Thermo, 28908) at room temperature. Fixed eyes were hemisected with cornea and lens removed prior to immunolabeling.

For cryosections (Figure 1B-C), eye cups were placed in cold 30% sucrose for 3-12 hours, coated in Optimal Cutting Temperature Media (OCT; Tissue-Tek, 4583), placed in a microcentrifuge tube filled with more OCT, frozen using a bath of dry ice and 95% ethanol, and stored at -20°C for at least 24h. 12 µm sections were cut using a Leica cryostat (CM3050) and mounted onto frost free slides (VWR, 48311-703) stored at -20°C. To stain cryosections, slides were warmed to room temperature, rinsed 3x with 1x phosphate buffered saline (PBS; Santa Cruz, sc-296028), then incubated sequentially with 0.5% TritonX-100 (Sigma, X100) and 1% BSA (VWR, 0332) for 1h each. Primary antibodies were diluted with 0.3% TritonX-100 + 1% BSA, applied to slides at 4°C and

incubated overnight. The slide was rinsed 3x with 1x PBS before applying secondary

antibodies diluted with 1x PBS. After incubating at room temperature for 1h, slides were

again rinsed 3x with 1x PBS, covered with mounting media containing DAPI (Invitrogen,

P36935), cover slipped and sealed with clear nail polish.

For whole mounts (Figure 1D), retinal pieces were incubated with 4% normal

donkey serum (NDS, Jackson Immuno, C840D36) in 1x PBS for >12h. Primary antibodies

were diluted in 4% NDS. Tubes were shielded from light and placed on a rocker at 4°C

for 7 days. Retinas were rinsed 3x with 1x PBS, then secondary antibody diluted in 1xPBS

applied. Tubes were again placed on the 4°C rocker for 1 day, rinsed 3x, mounted onto

filter paper, mounting media applied, coverslipped, and sealed with nail polish.

All slides were kept at 4°C until imaged. Z-stack confocal images were taken using

a Nikon AR1 microscope using 20x air and 60x oil objectives and motorized stage. Images

were processed using FIJI software (Schindelin et al., 2012). Z-stacks were flattened

according to their standard deviation. Brightness and contrast were adjusted as

necessary. Cones were manually counted from 60x cross-sections labeled with

antibodies to cone arrestin. Rod counts were obtained by measuring the area of the ONL

and average DAPI nucleus using the 'Measure' function in FIJI, with cone counts

subtracted.

Antibodies:

Rabbit anti-mCar 1:500 (Millipore, AB15282)

Mouse anti-PCP2 1:500 (Santa Cruz, sc-137064)

Rabbit anti- M-opsin 1:500 (Sigma, AB5405)

Alexa-fluor secondaries 1:500 (Invitrogen, A31571 & A31572)

MEA Recordings

Mice were dark adapted overnight by placing their home cage in a light shielded

box fitted with an air pump for circulation. All dissection procedures the day of the

experiment were carried out in complete darkness using infrared converters and cameras.

Mice were decapitated, eyes enucleated and placed into oxygenated room-temperature

Ames solution (Sigma, A1420) during retinal dissection and vitrectomy as described

previously (Yao et al., 2018). A ~1x2 mm piece from dorsal retina was placed RGC side

down on a MEA with either 512 electrodes spaced 60-um apart, or 519 electrodes with

30-um spacing (Field et al., 2010; Frechette et al., 2005; Ravi, Ahn, Greschner,

Chichilnisky, & Field, 2018). Oxygenated Ames perfused the retina throughout the

experiment at a rate of 6-8 mL/min, heated to 32°C.

Spike Sorting

Raw voltage traces from the MEA were spike sorted using custom software

followed by manual curation as described previously (Field et al., 2007; Shlens et al.,

2006). Cells with a spike rate >0.1 Hz and with <10% contamination estimated from

refractory period violations were retained for further analysis.

Visual Stimuli

The image from a gamma-calibrated OLED display (Emagin, SVGA + XL Rev3) was

focused onto photoreceptors using an inverted microscope (Nikon, Ti-E) and 4x objective

(Nikon, CFI Super Fluor ×4). Checkerboard stimuli were created and presented using

custom Matlab code. Light from the OLED display was attenuated using neutral density

filters. Retina was allowed to settle in darkness for approximately 30 min prior to initiating

data collection. Then, spontaneous activity in darkness was recorded for 30 min. The

display was then switched to a grey screen emitting ~100 Rh*/rod/s and the retina was

left to adapt for 5 min, after which mesopic spontaneous activity was recorded for 30 min.

Repeated checkerboard noise was presented 200x, repeating every 10s. Additionally,

non-repeating checkerboard noise was presented for 30min to estimate spatial and

temporal RFs. For all mesopic checkerboard stimuli, the stimulus refreshed every 66 ms

and the size of each square was 150x150 μm. After mesopic stimuli were presented, a

static screen was presented and the NDF filter removed to allow the retina to adapt to a

photopic light level (~10,000 Rh*/rod/ s). Checkerboard noise repeats (200x, 10s) were

presented, followed by 30 minutes of non-repeated checkerboard stimuli. For the photopic

checkerboard movies, the stimulus refreshed every 33 ms and each square was 75x75

μm. Two 10s movie clips were presented 100x to estimate the mutual information

between RGC signals and naturalistic movies. Movie 1 was a black and white video from

a camera attached to a cat walking in the woods (Supplemental video 1) (Betsch,

Einhäuser, Körding, & König, 2004). Movie 2 was a black and white video from a camera

carried by a squirrel, depicting fast moving tree leaves (Supplemental video 2) (modified

from Freiheit, 2016).

Measuring receptive fields

RGC responses to checkerboard noise were used to estimate the spatial and

temporal components of the spike triggered average (STA) (Chichilnisky, 2001). The STA

estimates the spatial and temporal integration of visual signals within the RF of an RGC.

For many RGCs, the STA is not space-time separable, meaning it cannot be expressed

as the outer product of a function that depends only on space and a function that depends

only on time. This precludes separately analyzing changes in spatial or temporal

integration. To identify RGCs with space-time separable RFs, singular value

decomposition (SVD) was performed on each STA. RGCs with STAs that were well-

approximated by a rank-one factorization were kept for further analysis of their spatial

and temporal RFs (Figure 2): these were cells for which the rank-one approximation

captured >60% of the variance in the STA. For quantifying temporal integration, the time-

to-zero crossing was used. This approximates the time-to-peak in the spiking response

for a step increment (for ON RGCs) or step decrement (for OFF RGCs) of light

(Chichilnisky & Kalmar, 2002). The analysis required that the temporal filter identified by

SVD was biphasic because a well-defined zero-crossing was needed to estimate the

time-to-zero: 79% of space-time separable STAs met this criterion.

Mutual information

Mutual information (MI) was used to measure the fidelity of RGC signaling. In

particular, MI measures how much observing the spike train reduces uncertainty about

the stimulus. MI was estimated using the 'direct method' (Strong, Koberle, De Ruyter Van

Steveninck, & Bialek, 1998). MI between a response and stimulus was computed as

$I(S; R) = H(R) - H(R|S)$ (1)

$I(S;R)$ is the mutual information about the stimulus contained in the response. $H(R)$

here is defined as the Shannon entropy of the response distribution (Shannon, 1948).

This measure estimates the capacity of a neuron to convey information about the stimulus

space. H(R|S) is the conditional entropy, a measurement of how noisy neural responses

are across repeated trials of an individual stimulus.

$H(R|S) = P(s) \sum P(r|s)\log 2(r|s)$ (2)

$H(R) = - \sum P(r)\log 2(r)$ (3)

P(r) is the probability of a spike count occurring in a pattern across all trials of the

stimulus space, while P(r|s) is the probability of observing a response pattern when a

stimulus is presented. These probabilities are estimated by measuring the proportion of

the observed response patterns at a given epoch of time from all the response patterns

across all epochs of time.

Responses from 100 repeated trials of 10 second-white noise were used to

estimate mutual information across all recorded RGCs. For each RGC, spike trains were

binned to a time resolution that achieved the entropy estimates from the Ma upper bound

Figure 11. Determining stability of estimated information rates to trial number and trial duration. *A. Example rasters illustrating the sub-sampling for fewer trials. B. Information rates (mean +/- 2SE) of 4 RGCs from separate cohorts as a function of number of repeated stimuli. Information rates stabilized at ~60 trials (repeats). C. Example rasters illustrating the sub-sampling of trial duration. D. Information rates (mean +/- 2SE) of 3 RGCs from separate degeneration cohorts by sub-sampling briefer stimulus durations. The stimulus used in this analysis was movie 1. Similar results were obtained for the other stimuli (movie 2 and checkerboard noise).*

(Ma, 1981; Strong et al., 1998). The response pattern length (a.k.a. 'word' length) was

selected using the same procedure. The mutual information was calculated using the

'direct method' (Buračas, Zador, DeWeese, & Albright, 1998; Strong et al., 1998). Bins

that achieved the Ma upper bound ranged from 4-6 milliseconds and response patterns

ranged from 3-6 bins across all RGCs. The mutual information rate was computed as the

quotient of the mutual information and the time length of the response pattern. Analysis

of trial number and trial duration indicated that both were sufficiently large to produce

stable estimates of the information rate (Figure 11).

Spontaneous activity and power spectral analysis

The frequency spectra of spontaneous activity were calculated using the fast

Fourier transform (FFT) applied to spike times binned at 1 ms from 30 min of spontaneous

activity. Spectra were analyzed using a frequency range from 0.1 to 35 Hz. Peaks in the

spectra were quantified for every RGC by computing: $abs(b-a)/abs(a-b)$ where b

represents the power at a baseline frequency level and a represents the maximum

power. The baseline level was estimated by finding the average power in the 0.1 to 2 Hz

range because this range was consistently flat across experiments and cells.

Statistical tests

Significant changes across all assays were assessed using the two-way

Kolmogorov–Smirnov test, a non-parametric test used to determine whether two sets of

samples arise from the same distribution (Massey, 1952). P-values were corrected for

multiple comparisons by Bonferroni correction. Error bars indicate the range of values

within 2 standard errors (SE) of the mean, which was estimated by bootstrapping the

mean 2000 times.

To measure whether differences across time points could be produced by other

factors (e.g., experiment-to-experiment variability), a parametric linear mixed effects

model was used (Bates, Mächler, Bolker, & Walker, 2015). The mixed effects model

accounts for retina-to-retina variability by adding each experiment as a random effect.

This procedure permitted making broad level inferences about the RGC populations

without dependence on experimental variability. In addition, the sex of the animal was

considered by including it as an interaction term with the degeneration conditions. This
step enabled determining whether degeneration conditions were associated with
information rates and RF sizes in a sex-independent fashion. The model indicated that
conclusions about the impact of degeneration on RGC signaling were insensitive to both
sex and experiment-to-experiment variability.

Acknowledgements

We thank our funding sources: National Institute of Health (NIH) R01 EY024280-
01, NIH NEI core grant EY5722, Holland Trice Foundation, Whitehead Foundation, and
Research to Prevent Blindness unrestricted grant to Duke University. We also are grateful
to Dr. Jon Cafaro and Dr. Suva Roy for technical assistance and Erika Ellis for
discussions.

Competing Interests

None.

References

Anishchenko, A., Greschner, M., Elstrott, J., Sher, A., Litke, A. M., Feller, M. B., &
Chichilnisky, E. J. (2010). Receptive field mosaics of retinal ganglion cells are
established without visual experience. *Journal of Neurophysiology*, *103*(4), 1856–
1864.
https://doi.org/10.1152/JN.00896.2009/ASSET/IMAGES/LARGE/Z9K00310000900
04.JPEG
Bainbridge, J. W. B., Mehat, M. S., Sundaram, V., Robbie, S. J., Barker, S. E.,
Ripamonti, C., … Ali, R. R. (2015). Long-term effect of gene therapy on Leber's
congenital amaurosis. *The New England Journal of Medicine*, *372*(20), 1887–1897.
https://doi.org/10.1056/NEJMOA1414221
Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting Linear Mixed-
Effects Models Using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

https://doi.org/10.18637/JSS.V067.I01

Beier, C., Hovhannisyan, A., Weiser, S., Kung, J., Lee, S., Lee, D. Y., … Sher, A.

(2017). Deafferented adult rod bipolar cells create new synapses with

photoreceptors to restore vision. *Journal of Neuroscience*, *37*(17), 4635–4644.

https://doi.org/10.1523/JNEUROSCI.2570-16.2017

Beier, C., Palanker, D., & Sher, A. (2018). Stereotyped Synaptic Connectivity Is

Restored during Circuit Repair in the Adult Mammalian Retina. *Current Biology*,

28(11), 1818-1824.e2. https://doi.org/10.1016/j.cub.2018.04.063

Betsch, B. Y., Einhäuser, W., Körding, K. P., & König, P. (2004). The world from a cat's

perspective - Statistics of natural videos. *Biological Cybernetics*, *90*(1), 41–50.

https://doi.org/10.1007/s00422-003-0434-6

Biswas, S., Haselier, C., Mataruga, A., Thumann, G., Walter, P., & Müller, F. (2014).

Pharmacological analysis of intrinsic neuronal oscillations in rd10 retina. *PLoS*

ONE, *9*(6). https://doi.org/10.1371/journal.pone.0099075

Borghuis, B. G., Ratliff, C. P., Smith, R. G., Sterling, P., & Balasubramanian, V. (2008).

Design of a neuronal array. *The Journal of Neuroscience : The Official Journal of*

the Society for Neuroscience, *28*(12), 3178–3189.

https://doi.org/10.1523/JNEUROSCI.5259-07.2008

Borowska, J., Trenholm, S., & Awatramani, G. B. (2011). An Intrinsic Neural Oscillator

in the Degenerating Mouse Retina. *Journal of Neuroscience*, *31*(13), 5000–5012.

https://doi.org/10.1523/JNEUROSCI.5800-10.2011

Buračas, G. T., Zador, A. M., DeWeese, M. R., & Albright, T. D. (1998). Efficient

discrimination of temporal patterns by motion-sensitive neurons in primate visual

cortex. *Neuron*, *20*(5), 959–969. https://doi.org/10.1016/S0896-6273(00)80477-8

Campochiaro, P. A., & Mir, T. A. (2018). The mechanism of cone cell death in Retinitis

Pigmentosa. *Progress in Retinal and Eye Research*, *62*, 24–37.

https://doi.org/10.1016/J.PRETEYERES.2017.08.004

Care, R. A., Anastassov, I. A., Kastner, D. B., Kuo, Y.-M. M., Della Santina, L., & Dunn,

F. A. (2020). Mature Retina Compensates Functionally for Partial Loss of Rod

Photoreceptors. *Cell Reports*, *31*(10), 107730.

https://doi.org/10.1016/j.celrep.2020.107730

Care, R. A., Kastner, D. B., De la Huerta, I., Pan, S., Khoche, A., Della Santina, L., …

Dunn, F. A. (2019). Partial Cone Loss Triggers Synapse-Specific Remodeling and

Spatial Receptive Field Rearrangements in a Mature Retinal Circuit. *Cell Reports*,

27(7), 2171-2183.e5. https://doi.org/10.1016/j.celrep.2019.04.065

Chen, J., Woodruff, M. L., Wang, T., Concepcion, F. A., Tranchina, D., & Fain, G. L.

(2010). Channel modulation and the mechanism of light adaptation in mouse rods.

The Journal of Neuroscience : The Official Journal of the Society for Neuroscience,

30(48), 16232–16240. https://doi.org/10.1523/JNEUROSCI.2868-10.2010

Chen, K., Wang, Y., Liang, X., Zhang, Y., Ng, T. K., & Chan, L. L. H. (2016).

Electrophysiology Alterations in Primary Visual Cortex Neurons of Retinal

Degeneration (S334ter-line-3) Rats. *Scientific Reports*, *6*.

https://doi.org/10.1038/srep26793

Chichilnisky, E. J. (2001). A simple white noise analysis of neuronal light responses.

Network (Bristol, England), *12*(2), 199–213. https://doi.org/10.1088/0954-

898X/12/2/306

Chichilnisky, E. J., & Kalmar, R. S. (2002). Functional asymmetries in ON and OFF

ganglion cells of primate retina. *The Journal of Neuroscience : The Official Journal*

of the Society for Neuroscience, *22*(7), 2737–2747. https://doi.org/20026215

Choi, H., Zhang, L., Cembrowski, M. S., Sabottke, C. F., Markowitz, A. L., Butts, D. A.,

… Riecke, H. (2014). Intrinsic bursting of AII amacrine cells underlies oscillations in

the rd1 mouse retina. *Journal of Neurophysiology*, *112*(6), 1491–1504.

https://doi.org/10.1152/jn.00437.2014

Cideciyan, A. V, Jacobson, S. G., Beltran, W. a, Sumaroka, A., Swider, M., Iwabe, S.,

… Aguirre, G. D. (2013). Human retinal gene therapy for Leber congenital

amaurosis shows advancing retinal degeneration despite enduring visual

improvement. *Proceedings of the National Academy of Sciences of the United*

States of America, *110*(6), E517-25. https://doi.org/10.1073/pnas.1218933110

Devries, S. H., & Baylor, D. A. (1997). Mosaic arrangement of ganglion cell receptive

fields in rabbit retina. *Journal of Neurophysiology*, *78*(4), 2048–2060.

Fairhall, A. L., Burlingame, C. A., Narasimhan, R., Harris, R. A., Puchalla, J. L., & Berry,

M. J. (2006). Selectivity for multiple stimulus features in retinal ganglion cells.

Journal of Neurophysiology, *96*(5), 2724–2738.

https://doi.org/10.1152/JN.00995.2005/ASSET/IMAGES/LARGE/Z9K01106777000

09.JPEG

Ferreira, S., Pereira, A. C., Quendera, B., Reis, A., Silva, E. D., & Castelo-Branco, M. (2017). Primary visual cortical remapping in patients with inherited peripheral retinal degeneration. *NeuroImage: Clinical*, *13*, 428–438. https://doi.org/10.1016/j.nicl.2016.12.013

Field, G. D., Gauthier, J. L., Sher, A., Greschner, M., Machado, T. A., Jepson, L. H., … Chichilnisky, E. J. (2010). Functional connectivity in the retina at the resolution of photoreceptors. *Nature*, *467*(7316), 673–677. https://doi.org/10.1038/nature09424

Field, G. D., Sher, A., Gauthier, J. L., Greschner, M., Shlens, J., Litke, A. M., & Chichilnisky, E. J. (2007). Spatial properties and functional organization of small bistratified ganglion cells in primate retina. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *27*(48), 13261–13272. https://doi.org/10.1523/JNEUROSCI.3437-07.2007

Frechette, E. S., Sher, A., Grivich, M. I., Petrusca, D., Litke, A. M., & Chichilnisky, E. J. (2005). Fidelity of the ensemble code for visual motion in primate retina. *Journal of Neurophysiology*, *94*(1), 119–135. https://doi.org/10.1152/JN.01175.2004

Freiheit, D. (2016). *GoPro Awards: Squirrel Runs Off With GoPro*. YouTube.

Gardiner, K. L., Cideciyan, A. V., Swider, M., Dufour, V. L., Sumaroka, A., Komáromy, A. M., … Aguirre, G. D. (2020). Long-term Structural Outcomes of Late-stage RPE65 Gene Therapy. *Molecular Therapy*, *28*(1), 266–278. https://doi.org/10.1016/j.ymthe.2019.08.013

Gekeler, F., Messias, A., Ottinger, M., Ulrich Bartz-Schmidt, K., & Zrenner, E. (2006). Phosphenes Electrically Evoked with DTL Electrodes: A Study in Patients with Retinitis Pigmentosa, Glaucoma, and Homonymous Visual Field Loss and Normal Subjects. *IOVS*, *47*(11), 4966–4974. https://doi.org/10.1167/iovs.06-0459

Golomb, D., Kleinfeld, D., Reid, R. C., Shapley, R. M., & Shraiman, B. I. (1994). On temporal codes and the spatiotemporal response of neurons in the lateral geniculate nucleus. *Journal of Neurophysiology*, *72*(6), 2990–3003. https://doi.org/10.1152/JN.1994.72.6.2990

Grover, S., Fishman, G. A., Anderson, R. J., Tozatti, M. S. V., Heckenlively, J. R.,

Weleber, R. G., … Brown, J. (1999). Visual acuity impairment in patients with

retinitis pigmentosa at age 45 years or older. *Ophthalmology*, *106*(9), 1780–1785.

https://doi.org/10.1016/S0161-6420(99)90342-1

Haq, W., Arango-Gonzalez, B., Zrenner, E., Euler, T., & Schubert, T. (2014). Synaptic

remodeling generates synchronous oscillations in the degenerated outer mouse

retina. *Frontiers in Neural Circuits*, *8*, 108. https://doi.org/10.3389/fncir.2014.00108

Hartong, D. T., Berson, E. L., & Dryja, T. P. (2006). Retinitis pigmentosa. *Lancet*

(London, England), *368*(9549), 1795–1809. https://doi.org/10.1016/S0140-

6736(06)69740-7

Hickmott, P. W., & Merzenich, M. M. (2002). Local circuit properties underlying cortical

reorganization. *Journal of Neurophysiology*, *88*(3), 1288–1301.

https://doi.org/10.1152/jn.00994.2001

Hüttl, S., Michalakis, S., Seeliger, M., Luo, D.-G., Acar, N., Geiger, H., … Biel, M.

(2005). Impaired Channel Targeting and Retinal Degeneration in Mice Lacking the

Cyclic Nucleotide-Gated Channel Subunit CNGB1. *Journal of Neuroscience*, *21*(1),

130–138. https://doi.org/10.1523/JNEUROSCI.3764-04.2005

Ivanova, E., Yee, C. W., Baldoni, R., & Sagdullaev, B. T. (2016). Aberrant activity in

retinal degeneration impairs central visual processing and relies on Cx36-

containing gap junctions. *Experimental Eye Research*, *150*, 81–89.

https://doi.org/10.1016/j.exer.2015.05.013

Jacobson, S. G., Cideciyan, A. V, Roman, A. J., Sumaroka, A., Schwartz, S. B., Heon,

E., & Hauswirth, W. W. (2015). Improvement and decline in vision with gene

therapy in childhood blindness. *The New England Journal of Medicine*, *372*(20),

1920–1926. https://doi.org/10.1056/NEJMoa1412965

Jones, B. W., & Marc, R. E. (2005). Retinal remodeling during retinal degeneration.

Experimental Eye Research, *81*(2), 123–137.

https://doi.org/10.1016/J.EXER.2005.03.006

Keat, J., Reinagel, P., Reid, R. C., & Meister, M. (2001). Predicting every spike: a model

for the responses of visual neurons. *Neuron*, *30*(3), 803–817.

https://doi.org/10.1016/S0896-6273(01)00322-1

Keck, T., Keller, G. B., Jacobsen, R. I., Eysel, U. T., Bonhoeffer, T., & Hübener, M.

(2013). Synaptic scaling and homeostatic plasticity in the mouse visual cortex in

vivo. *Neuron*, *80*(2), 327–334. https://doi.org/10.1016/j.neuron.2013.08.018

Keck, T., Mrsic-Flogel, T. D., Vaz Afonso, M., Eysel, U. T., Bonhoeffer, T., & Hübener,

M. (2008). Massive restructuring of neuronal circuits during functional

reorganization of adult visual cortex. *Nature Neuroscience*, *11*(10), 1162–1167.

https://doi.org/10.1038/nn.2181

Keck, T., Scheuss, V., Jacobsen, R. I., Wierenga, C. J., Eysel, U. T., Bonhoeffer, T., &

Hübener, M. (2011). Loss of sensory input causes rapid structural changes of

inhibitory neurons in adult mouse visual cortex. *Neuron*, *71*(5), 869–882.

https://doi.org/10.1016/j.neuron.2011.06.034

Koch, K., McLean, J., Berry, M., Sterling, P., Balasubramanian, V., & Freed, M. A.

(2004). Efficiency of Information Transmission by Retinal Ganglion Cells. *Current*

Biology, *14*(17), 1523–1530. https://doi.org/10.1016/J.CUB.2004.08.060

Koch, K., McLean, J., Segev, R., Freed, M. A., Berry, M. J., Balasubramanian, V., &

Sterling, P. (2006). How Much the Eye Tells the Brain. *Current Biology*, *16*(14),

1428–1434.

Koch, S., Sothilingam, V., Garrido, M. G., Tanimoto, N., Becirovic, E., Koch, F., …

Michalakis, S. (2012). Gene therapy restores vision and delays degeneration in the

CNGB1 2/2 mouse model of retinitis pigmentosa. *Human Molecular Genetics*,

21(20), 4486–4496. https://doi.org/10.1093/hmg/dds290

Lee, J. Y., Care, R. A., Santina, L. Della, & Dunn, F. A. (2021). Impact of Photoreceptor

Loss on Retinal Circuitry. *Annual Review of Vision Science*, *7*, 105–128.

https://doi.org/10.1146/ANNUREV-VISION-100119-124713

Leinonen, H., Pham, N. C., Boyd, T., Santoso, J., Palczewski, K., & Vinberg, F. (2020).

Homeostatic plasticity in the retina is associated with maintenance of night vision

during retinal degenerative disease. *ELife*, *9*, 1–27.

https://doi.org/10.7554/ELIFE.59422

Litke, A. M. M., Bezayiff, N., Chichilnisky, E. J. J., Cunningham, W., Dabrowski, W.,

Grillo, A. A. A., … Sher, A. (2004). What does the eye tell the brain?: Development

of a system for the large-scale recording of retinal output activity. *IEEE*

862 *Transactions on Nuclear Science*, *51*(4 I), 1434–1440.

863 https://doi.org/10.1109/TNS.2004.832706

864 Liu, X., Garriga, P., & Khorana, H. G. (1996). Structure and function in rhodopsin:

865 correct folding and misfolding in two point mutants in the intradiscal domain of

866 rhodopsin identified in retinitis pigmentosa. *Proceedings of the National Academy*

867 *of Sciences of the United States of America*, *93*(10), 4554–4559.

868 https://doi.org/10.1073/PNAS.93.10.4554

869 Ma, S. keng. (1981). Calculation of entropy from data of motion. *Journal of Statistical*

870 *Physics 1981 26:2*, *26*(2), 221–240. https://doi.org/10.1007/BF01013169

871 Machida, S., Kondo, M., Jamison, J. A., Khan, N. W., Kononen, L. T., Sugawara, T., …

872 Sieving, P. A. (2000). P23H rhodopsin transgenic rat: Correlation of retinal function

873 with histopathology. *Investigative Ophthalmology and Visual Science*, *41*(10).

874 Marc, R. E., Jones, B. W., Anderson, J. R., Kinard, K., Marshak, D. W., Wilson, J. H., …

875 Lucas, R. J. (2007). Neural reprogramming in retinal degeneration. *Investigative*

876 *Ophthalmology & Visual Science*, *48*(7), 3364–3371.

877 https://doi.org/10.1167/iovs.07-0032

878 Margolis, D. J., Newkirk, G., Euler, T., & Detwiler, P. B. (2008). Functional stability of

879 retinal ganglion cells after degeneration-induced changes in synaptic input. *Journal*

880 *of Neuroscience*, *28*(25), 6526–6536. https://doi.org/10.1523/JNEUROSCI.1533-

881 08.2008

882 Massey, F. J. (1952). Distribution Table for the Deviation Between two Sample

883 Cumulatives. *The Annals of Mathematical Statistics*, *23*(3), 435–441.

884 https://doi.org/10.1214/aoms/1177729388

885 McDonnell, M. D., Ikeda, S., & Manton, J. H. (2011). An introductory review of

886 information theory in the context of computational neuroscience. *Biological*

887 *Cybernetics*, *105*(1), 55–70. https://doi.org/10.1007/s00422-011-0451-9

888 Menzler, J., & Zeck, G. (2011). Network oscillations in rod-degenerated mouse retinas.

889 *Journal of Neuroscience*, *31*(6), 2280–2291.

890 https://doi.org/10.1523/JNEUROSCI.4238-10.2011

891 Merabet, L. B., & Pascual-Leone, A. (2010). Neural reorganization following sensory

892 loss: The opportunity of change. *Nature Reviews Neuroscience*, *11*(1), 44–52.

893 https://doi.org/10.1038/nrn2758

894 Paninski, L. (2003). Estimation of Entropy and Mutual Information. *Neural Computation*,

895 (15), 1191–1253.

896 Poria, D., & Dhingra, N. K. (2015). Spontaneous oscillatory activity in rd1 mouse retina

897 is transferred from on pathway to off pathway via glycinergic synapse. *Journal of*

898 *Neurophysiology*, *113*(2), 420–425. https://doi.org/10.1152/jn.00702.2014

899 Ravi, S., Ahn, D., Greschner, M., Chichilnisky, E. J., & Field, G. D. (2018). Pathway-

900 Specific Asymmetries between ON and OFF Visual Signals. *The Journal of*

901 *Neuroscience : The Official Journal of the Society for Neuroscience*, *38*(45), 9728–

902 9740. https://doi.org/10.1523/JNEUROSCI.2008-18.2018

903 Rieke, F., Bodnar, D. A., & Bialek, W. (1995). Naturalistic stimuli increase the rate and

904 efficiency of information transmission by primary auditory afferents. *Proceedings.*

905 *Biological Sciences*, *262*(1365), 259–265. https://doi.org/10.1098/RSPB.1995.0204

906 Sahel, J., Bonnel, S., Mrejen, S., & Paques, M. (2010). Retinitis pigmentosa and other

907 dystrophies. *Developments in Ophthalmology*, *47*, 160–167.

908 https://doi.org/10.1159/000320079

909 Sakami, S., Maeda, T., Bereta, G., Okano, K., Golczak, M., Sumaroka, A., …

910 Palczewski, K. (2011). Probing mechanisms of photoreceptor degeneration in a

911 new mouse model of the common form of autosomal dominant retinitis pigmentosa

912 due to P23H opsin mutations. *The Journal of Biological Chemistry*, *286*(12),

913 10551–10567. https://doi.org/10.1074/JBC.M110.209759

914 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., …

915 Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis.

916 *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/NMETH.2019

917 Sekirnjak, C., Jepson, L. H., Hottowy, P., Sher, A., Dabrowski, W., Litke, A. M., &

918 Chichilnisky, E. J. (2011). Changes in physiological properties of rat ganglion cells

919 during retinal degeneration. *Journal of Neurophysiology*, *105*(5), 2560–2571.

920 https://doi.org/10.1152/jn.01061.2010

921 Shannon, C. E. (1948). A Mathmatical Theory of Cummunication. *The Bell System*

922 *Technical Journal*, *27*, 379–423, 623–656.

923 Shen, N., Wang, B., Soto, F., & Kerschensteiner, D. (2020). Homeostatic Plasticity

Shapes the Retinal Response to Photoreceptor Degeneration. *Current Biology*, *30*(10), 1916-1926.e3. https://doi.org/10.1016/j.cub.2020.03.033

Shlens, J., Field, G. D., Gauthier, J. L., Grivich, M. I., Petrusca, D., Sher, A., … Chichilnisky, E. J. (2006). The structure of multi-neuron firing patterns in primate retina. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *26*(32), 8254–8266. https://doi.org/10.1523/JNEUROSCI.1282-06.2006

Stasheff, S. F. (2008). Emergence of sustained spontaneous hyperactivity and temporary preservation of OFF responses in ganglion cells of the retinal degeneration (rd1) mouse. *Journal of Neurophysiology*, *99*(3), 1408–1421. https://doi.org/10.1152/jn.00144.2007

Stasheff, S. F., Shankar, M., & Andrews, M. P. (2011). Developmental time course distinguishes changes in spontaneous and light-evoked retinal ganglion cell activity in rd1 and rd10 mice. *Journal of Neurophysiology*, *105*(6), 3002–3009. https://doi.org/10.1152/jn.00704.2010

Strong, S. P., Koberle, R., De Ruyter Van Steveninck, R. R., & Bialek, W. (1998). Entropy and Information in Neural Spike Trains. *Physical Review Letters*, *80*(1), 197. https://doi.org/10.1103/PhysRevLett.80.197

Trenholm, S., & Awatramani, G. B. (2015). Origins of spontaneous activity in the degenerating retina. *Frontiers in Cellular Neuroscience*, *9*(277). https://doi.org/10.3389/fncel.2015.00277

Tu, H.-Y., Chen, Y.-J., McQuiston, A. R., Chiao, C.-C., & Chen, C.-K. (2016). A Novel Retinal Oscillation Mechanism in an Autosomal Dominant Photoreceptor Degeneration Mouse Model. *Frontiers in Cellular Neuroscience*, *9*, 513. https://doi.org/10.3389/fncel.2015.00513

Turrigiano, G. (2012). Homeostatic synaptic plasticity: Local and global mechanisms for stabilizing neuronal function. *Cold Spring Harbor Perspectives in Biology*, *4*(1). https://doi.org/10.1101/cshperspect.a005736

Van Steveninck, R. R. D. R., Lewen, G. D., Strong, S. P., Koberle, R., & Bialek, W. (1997). Reproducibility and variability in neural spike trains. *Science (New York, N.Y.)*, *275*(5307), 1805–1808. https://doi.org/10.1126/SCIENCE.275.5307.1805

Vaney, D. I., Sivyer, B., & Taylor, W. R. (2012). Direction selectivity in the retina:

symmetry and asymmetry in structure and function. *Nature Reviews. Neuroscience*,

13(3), 194–208. https://doi.org/10.1038/NRN3165

Wang, T., Pahlberg, J., Cafaro, J., Frederiksen, R., Cooper, A. J., Sampath, A. P., …

Chen, J. (2019). Activation of Rod Input in a Model of Retinal Degeneration

Reverses Retinal Remodeling and Induces Formation of Functional Synapses and

Recovery of Visual Signaling in the Adult Retina. *The Journal of Neuroscience*,

39(34), 6798 LP – 6810. https://doi.org/10.1523/JNEUROSCI.2902-18.2019

Wolfe, J., & Palmer, L. A. (1998). Temporal diversity in the lateral geniculate nucleus of

cat. *Visual Neuroscience*, *15*(4), 653–675.

https://doi.org/10.1017/S0952523898154068

Yao, X., Cafaro, J., McLaughlin, A. J., Postma, F. R., Paul, D. L., Awatramani, G., &

Field, G. D. (2018). Gap Junctions Contribute to Differential Light Adaptation across

Direction-Selective Retinal Ganglion Cells. *Neuron*, *100*, 216–228.

https://doi.org/10.1016/J.NEURON.2018.08.021

Ye, J. H., & Goo, Y. S. (2007). The slow wave component of retinal activity in rd/rd mice

recorded with a multi-electrode array. *Physiological Measurement*, *28*(9), 1079–

1088. https://doi.org/10.1088/0967-3334/28/9/009

Yu, W.-Q., Grzywacz, N. M., Lee, E.-J., & Field, G. D. (2017). Cell type-specific

changes in retinal ganglion cell function induced by rod death and cone

reorganization in rats. *Journal of Neurophysiology*, *118*(1), 434–454.

https://doi.org/10.1152/jn.00826.2016

Zhang, Y., Molday, L. L., Molday, R. S., Sarfare, S. S., Woodruff, M. L., Fain, G. L., …

Pittler, S. J. (2009). Knockout of GARPs and the β-subunit of the rod cGMP-gated

channel disrupts disk morphogenesis and rod outer segment structural integrity.

Journal of Cell Science, *122*, 1192–1200. https://doi.org/10.1242/JCS.042531