Integrated Genomics Approaches Identify Transcriptional Mediators and Epigenetic

Responses to Afghan Desert Particulate Matter in Small Airway Epithelial Cells

Arnav Gupta, Sarah K. Sasse, Reena Berman, Margaret A. Gruca, Robin D. Dowell, Hong Wei

Chu, Gregory P. Downey, Anthony N. Gerber

Abstract

Military Deployment to Southwest Asia and Afghanistan and exposure to toxic airborne

particulates has been associated with an increased risk of developing respiratory disease,

collectively termed deployment-related respiratory diseases (DRRD). Our knowledge about how

particulates mediate respiratory disease is limited, precluding the appropriate recognition or

management. Central to this limitation is the lack of understanding of how exposures translate

into dysregulated cell identity with dysregulated transcriptional programs. The small airway

epithelium is involved in both the pathobiology of DRRD and fine particulate matter deposition.

To characterize small airway epithelial cell epigenetic and transcriptional responses to Afghan

desert particulate matter (APM) and investigate the functional interactions of transcription

factors that mediate these responses, we applied two genomics assays, the assay for

transposase accessible chromatin with sequencing (ATAC-seq) and Precision Run-on

sequencing (PRO-seq). We identified activity changes in a series of transcriptional pathways as

candidate regulators of susceptibility to subsequent insults, including signal-dependent

pathways, such as loss of cytochrome P450 or P53/P63, and lineage-determining transcription

factors, such as GRHL2 loss or TEAD3 activation. We further demonstrated that TEAD3

activation was unique to APM exposure despite similar inflammatory responses when compared

with wood smoke particle exposure and that P53/P63 program loss was uniquely positioned at

the intersection of signal-dependent and lineage-determining transcriptional programs. Our

results establish the utility of an integrated genomics approach in characterizing responses to

exposures and identify genomic targets for the advanced investigation of the pathogenesis of

DRRD.

Introduction

The development of respiratory disease following post-9/11 deployment to Southwest Asia (SW

Asia) and Afghanistan and exposure to associated airborne hazards, collectively termed

deployment-related respiratory diseases (DRRD), is an important clinical problem that has had a

major adverse impact on the long-term health of Veterans. DRRD encompasses several

respiratory conditions, such as asthma, constrictive bronchiolitis and chronic rhinosinusitis (1),

which develop following deployment and exposure in individuals in the absence of known prior

respiratory illness or predisposition. Epidemiologic studies have previously associated exposure

with emergent respiratory disease, both in the context of SW Asia and Afghanistan deployment

(1, 2) and domestic air pollution (3). Additional studies of airborne exposures extend their impact

beyond inflammatory and remodeling diseases of the respiratory tract and implicate these

exposures in the development of lung cancer, cardiovascular disease and neuropsychiatric

disease (4–7). This raises additional concerns about long-term sequelae of deployment

exposures. However, our mechanistic understanding of the impacts of deployment exposures in

relationship to disease pathogenesis and long-term risk remains fragmentary.

Previous studies have used cell and animal-based models of particulate matter (PM) exposure

to identify potentially relevant cellular, physiologic and pathophysiologic responses. Although

such studies fail to capture the complexity of true inhalational particulate exposure, they have

been useful in defining PM-regulated pathways relevant to human exposures, such as Type I

inflammation. For example, wood smoke particle (WSP) exposure resulted in increased

inflammatory markers in the bronchoalveolar lavage (BAL) fluid of healthy human subjects (8)

and induced pro-inflammatory gene expression in submerged airway epithelial cell culture (9).

Building on these physiologic consistencies, other studies identified mediators that may

contribute to airways diseases, such as mucus abnormalities (10) or cilia dysfunction (11), as

well as long term sequelae, such as cellular dysplasia and malignancy (12). Additional studies

connected inflammatory responses to particulate exposure with development of fibrosis (13) or

allergic sensitization (14), underscoring the potential for particulate exposures to contribute to

new-onset disease. More recently, particulate matter models specific to Afghanistan deployment

(Afghan Desert Particulate matter or APM) have demonstrated physiologic responses relevant

to DRRD, such as inflammation or airway hyperresponsiveness (15), and induction of cytokines

such as CXCL8 and IL-33 (16). However, these studies did not identify the mechanistic basis for

these responses, nor did they examine factors that may distinguish the effects of APM in

comparison to other PMs that are associated with a different spectrum of disease risk.

Compounding the unknowns underlying particulate biology, the interactions between

transcription factors that respond to external stimuli and the underlying lineage-specific

transcriptional programming of the airway epithelia are poorly understood. We have previously

described that multiple transcription factor programs are responsible for the acute transcriptional

response to particulate exposure (17). To better conceptualize the interplay between

transcription factors in response to exposures, we have adopted a classification paradigm (18)

in which transcription factors are grouped based on their broad function in programming the

cellular transcriptome. In this paradigm, lineage-determining transcription factors (LDTFs)

establish and maintain cell-type specific chromatin architecture, which forms the genomic

context in which subsequent signals enact transcriptional responses. Signal-dependent

transcription factors (SDTFs) instead, but not mutually exclusively, direct transcriptional

responses to specific stimuli. However, this activity is frequently determined by the architecture

that arises from the activity of LDTFs (19). Although well established in other cell types (20), the

interplay between LDTFs and SDTFs in the airway epithelia has not been characterized in the

context of exposures. This mechanism is critical in understanding how exposures disrupt normal

airway epithelia cell function.

Our overall objective in this study was to identify acute responses to a deployment-relevant

exposure in small airway epithelial progenitor cells. This cell type features unique lineage-

specific transcriptional programming (21) and is a critical cell type in the development of

constrictive bronchiolitis. Although extrapolating acute responses to understand the

development of a chronic, exposure-associated disease is challenging, we interpreted our

results in the context of a multi-hit hypothesis, where acute responses mediate susceptibility to

further insults that are ultimately compounded into a dysregulated disease state. We developed

a comprehensive set of acute responses to APM exposure with both epigenetic and

transcriptional effects by performing ATAC-seq and PRO-seq on SAECs. Through comparison

with a non-deployment PM exposure, WSP, we identified responses unique to APM exposure.

Finally, we identified genomic regions that shared dynamic responses in chromatin accessibility

and nascent transcription to APM as candidate sites for interactions between SDTFs and

LDTFs. Genome-wide analysis of these regions identified the P53/P63 family as enriched within

this intersection and likely to play a significant role in intersecting small airway progenitor cell

fate and function in response to APM exposure.

Methods

Cell culture

Cell culture of primary small airway epithelial cells (SAECs) was performed as described

previously (22). Briefly, de-identified primary human small airway epithelial cells (< 2 mm

diameter) were obtained from the National Jewish Health Biobank from donors with no history of

deployment or chronic lung disease. Cells were first co-cultured with irradiated NIH/3T3 (ATCC)

fibroblasts in F-medium containing 1 uM Y-27632 (APEX Bio) and then passaged onto tissue

culture dishes double-coated with Type I bovine collagen solution (Advanced Biomatrix) and

grown to confluence in BronchiaLife Epithelial Medium with the complete LifeFactors

Supplement Kit (Lifeline Cell Technology). All exposures were performed in this medium. All

cells were maintained in 5% CO_2 at 37°C.

Afghan Desert Particulate Matter – Preparation and Exposure

Afghan Desert Particulate Matter (APM) was a generous gift from Drs. Gregory Downey and

Hong Wei Chu, who obtained this material from Drs. Brian Wong and Karen Mumy at the Naval

Medical Research Unit (NAMRU) in Dayton, Ohio. The composition and cell culture exposure

conditions of this material have been characterized previously (16). Briefly, topsoil from Bagram

Air Force Base in the Parwan Province of Afghanistan was collected in August 2009 and

shipped to the United States, where it was autoclaved and irradiated in accordance with

Department of Agriculture standards. The material was then aerosolized into an exposure

chamber using a Wright Dust Feeder (Mesa Labs, NJ) and size fractionated. Particle size was

monitored using an optical sensor, gravimetric filter, and cascade impactor sampler. Settled

particulates were collected from the bottom of the chamber and re-sterilized using gamma

irradiation. Scanning electron microscopy determined that 89.5% of the particles were < 2.5 um

in diameter, 10.4% between 2.5 and 10 um and 0.1% greater than 10 um. Dry APM was

resuspended in 1X Phosphate buffered saline (PBS, Dulbecco) and vortexed prior to addition to

cell culture media. After addition, media was gently agitated to ensure even distribution of the

particles.

Wood Smoke Particles – Preparation and Exposure

Wood Smoke Particles (WSP) were generously gifted by Dr. Andrew Ghio at the Environmental

Protection Agency Laboratory in North Carolina and generated as described previously (8).

WSP composition, preparation and exposure were described previously by our laboratory (22).

The mean diameter of the freshly generated wood smoke particles was 0.14 um. After receipt of

dry WSP, particles were suspended in 1X PBS and disaggregated by sonication and sterilized

under UV light for 20 minutes. WSP particle suspension was then added directly to cell culture

media at concentrations of 0.001 mg/ml to 1 mg/ml as indicated in the text and gently agitated to

evenly disperse particles.

Interleukin 8 Quantification by Enzyme linked Immunosorbent Assay (ELISA)

Quantification of IL-8 was performed by a commercial ELISA kit (R&D Systems). Briefly, plates

were prepared with anti-IL-8 capture antibody prior to addition of sample. Cell supernatant was

harvested following exposure and applied directly to prepared plates. Anti-IL-8 detection

antibody was then added, and IL-8 levels were quantified from streptavidin-Horseradish

peroxidase associated with the detection antibody using absorbance read by an Infinite M1000

plate reader (Tecan).

Precision Run-on Sequencing (PRO-seq)

Cells required treatment and harvest on two separate days to generate enough nuclei to

perform PRO-seq on two replicates per condition (untreated control (Ctrl) vs APM). For each set

of replicates, Sm1 cells (small airway epithelial cells from donor 1) were seeded on 5 x 15 cm

double collagen-coated plates per treatment and grown to confluence (3-4 days), then left

untreated (Ctrl) or treated with APM (50 μg/cm^2) for 20 hours. Cells were harvested and nuclei

prepared as reported previously (23). PRO-seq libraries for all replicates were then prepared

simultaneously by subjecting one aliquot of 1e7 nuclei/sample to 3-minute nuclear run-on

reactions in the presence of Biotin-11-CTP (PerkinElmer) following our previously detailed

protocol (24). Uniquely indexed libraries were pooled and sequenced on an Illumina NextSeq

instrument using 75 bp single-end reads by the BioFrontiers Sequencing Facility at the

University of Colorado Boulder.

PRO-seq Computational Analysis

PRO-seq data were processed using a standardized Nextflow pipeline

(https://github.com/Dowell-Lab/Nascent-Flow). A detailed quality control report including

trimming, mapping, coverage, and complexity metrics is included in Supplemental File SR1.

TDF coverage files output by the pipeline, normalized by reads per million mapped, were

visualized using the Integrative Genomics Viewer (IGV) genome browser (v. 2.8.0). FStitch (v.

1.0) and Tfit (v. 1.0) were used to identify regions with bidirectional transcriptional activity as

described (22). Reference Sequences for hg38 were downloaded from the UCSC genome

browser (May 18, 2018). DESeq2 (v. 1.20.0, Bioconductor release v. 3.7) was used to

determine differentially transcribed genes between control and APM exposure. Functional

annotation of differentially transcribed genes was performed using DAVID v. 6.8 (25). For

bidirectional comparisons, all predicted bidirectional Tfit calls were aggregated using mergeBed

(argument -d 60) from BEDTools (v. 2.28.0) to generate an annotation file. Counts were then

calculated for each sample using multicov (BEDTools v.2.28.0), and DESeq2 was used to

identify differentially transcribed bidirectionals between treatments.

Assay for Transposase Accessible Chromatin using Sequencing (ATAC-seq)

Sm1 cells were grown as described above for PRO-seq and aliquots of cells were taken from

confluent 15 cm tissue culture dishes after vehicle (PBS) or APM treatment. Sm2 and Sm3 cells

were grown to confluence in 6-well tissue culture dishes and treated with vehicle (PBS), APM or

WSP for 20 hours. Cells were rinsed and scraped in ice-cold PBS, then ~50K cells from each

treatment were pelleted and processed in duplicate for Omni-ATAC-seq using the protocol

developed by Corces et al (26). Uniquely indexed libraries were pooled and sequenced on an

Illumina NextSeq using 37 bp single- or paired-end reads by the BioFrontiers Sequencing

Facility at the University of Colorado Boulder or on an Illumina NovaSeq using 150 bp paired-

end reads at the University of Colorado Anschutz Medical Campus Shared Genomic Resource.

ATAC-seq Computational Analysis

Computational analysis of ATAC-seq data was performed as described previously (22). Briefly,

ATAC-seq reads were trimmed for adapters, minimum length, and minimum quality using the

bbduk tool from the BBMap Suite (v. 38.73) with arguments 'ref=adapters.fa ktrim=r qtrim=10

k=23 mink=11 hdist=1 maq=10 minlen=20'. Samples sequenced using 150 bp reads were

further trimmed to 37 bp in length for consistency using the additional argument 'ftr=36'.

Trimmed reads were mapped to the human genome (hg38; downloaded from the UCSC

genome browser on September 16, 2019, with corresponding hisat2 index files) using hisat2 (v.

2.1.0). Duplicate reads were removed using picard tools (v.2.26.6;

http://broadinstitute.github.io/picard/) MarkDuplicates and deduplicated and sorted BAM files

were converted to bedGraph coverage format using genomeCoverageBed from the BEDTools

suite (v. 2.29.2). Read coverage was normalized to reads per million mapped using a custom

python script and files converted to TDF format using igvtools (v. 2.5.3) for visualization in IGV.

MACS2 (v. 2.1.4) callpeak with '--SPMR' argument was applied to sorted BAM files for each

replicate pair of samples with parameters set to call peaks with $\log_2$ fold change > 1 above

baseline with q < 1e-5 as significant.

Venn Diagrams for Feature Intersection

ATAC-seq peaks called by MACS2 and PRO-seq regulatory elements identified by Tfit were

intersected using the BEDTools suite (v. 2.29.2) command "bedtools intersect." Briefly, this

operation took standard bed format input from two files and returned regions of the first file that

had at least 1 base pair overlap with regions of the second file.

Differential Transcription Factor Motif Enrichment

Transcription factor enrichment analysis for both PRO- and ATAC-seq datasets was performed

using TFEA (v.1.0; https://github.com/Dowell-Lab/TFEA; (27)). Briefly, sets of consensus

regions of interest (ROIs) are first defined for all Tfit-called bidirectionals (for PRO-seq) or all

MACS2-called peaks (for ATAC-seq) using the muMerge function of TFEA. TFEA then

calculates read coverage for each replicate within each ROI and applies DESeq2 to determine

differentially represented ROIs by treatment condition and assign rank as a function of p-value.

For each ranked ROI, the FIMO tool (28) from the MEME suite scans the 3 kb sequence

surrounding the ROI center (bidirectional origin or ATAC-seq peak summit) for positions of

transcription factor consensus motif matches (with p-value cutoff of 1e-5), represented by a

curated set of position weight matrices (29). Enrichment scores (E-scores) for each motif are

then calculated and corrected for sequence content to reduce known biases associated with

local GC enrichment and p-values are determined using Z-scores. Motif displacement

distributions are displayed as dot-plots where the location of all motif instances within $\pm$1500 bp

are indicated relative to the ROI center on the vertical axis, and the rank of each ROI based on

signal loss or gain is indicated on the horizontal axis.

Western Blotting

Western blotting and protein detection were performed using standard protocols (30) in SAECs

cultured to confluence in 6-well plates and treated with vehicle control or APM. Primary

antibodies purchased included anti-GRHL2 (Sigma HPA004820), anti-pan-TEAD (Cell Signaling

Technologies 13295S) and anti-GAPDH (Santa Cruz sc-25778). Secondary antibody was ECL

Donkey anti-Rabbit IgG, HRP-linked F(ab')$_2$ fragment (NA9340) from GE

Healthcare/Amersham.

RNA purification and quantitative reverse transcription-PCR (qRT-PCR)

RNA purification and qRT-PCR was performed as described previously (22). Briefly, SAECs

were grown to confluence in 12-well tissue culture dishes and treated with vehicle (PBS) or

APM for 20 hours. Cells were harvested in TRIzol (Life Technologies) and RNA purified by

PureLink RNA Mini Kit (Life Technologies) prior to qRT-PCR, performed with normalization to

RPL19 as previously detailed (23). Sequences of primers used for qRT-PCR analysis are

provided in Supplemental Table ST1.

Statistics

Statistical comparisons for IL-8 ELISA and qRT-PCR assays were made by 2-tailed t-test using

the Bonferroni correction when appropriate. These analyses were conducted using statistical

software embedded in Open Office (Apache OpenOffice v 4.1.7

http://www.openoffice.org/welcome/credits.html).

All High Throughput Sequencing Data is publicly available through the Gene Expression

Omnibus (GEO) with accession GSE201152. Subseries GSE201149 contains 20 ATAC-seq

libraries and subseries GSE201150 contains 4 PRO-seq libraries.

Results

Primary small airway epithelial cells (SAECs) secrete IL-8 in response to APM and WSP

exposure in submerged culture

To establish conditions to compare APM with other clinically relevant PM exposures, we

exposed SAECs to APM (50 µg/cm^2 culture surface area) for 20 hours, based on a previous

time course and dose response analysis (16). We demonstrated an increase in interleukin 8 (IL-

8) secretion in cells from three healthy non-smoking donors, consistent with prior studies (16)

(Figure 1A). We next sought to compare APM exposure with an additional particulate exposure

model that resulted in similar inflammatory responses but had different clinical associations. We

selected wood smoke particles (WSP), a highly relevant model for wildfire smoke exposure (31),

and measured IL-8 secretion in response to varied doses of WSP exposure after 20 hours

(Supplemental Figure SF1). Primary SAECs from the same three donors exhibited a similar

increase in IL-8 secretion in response to APM exposure at 50 µg/cm^2 and WSP exposure at 1

mg/ml (Figure 1B). Therefore, we selected these doses of APM (50 µg/cm^2) and WSP (1 mg/ml)

for subsequent genome-wide analysis and comparisons. Total particulate exposure volume was

similar but different metrics were used based on convention.

ATAC-seq (Assay for Transposase Accessible Chromatin with sequencing) identifies chromatin

accessibility changes induced by APM exposure

Changes in chromatin architecture in response to a stimulus are generally mediated by the

activity of transcription factors and the context of the underlying chromatin architecture of the

particular cell type (32). To determine and compare regulation of chromatin changes in

response to APM exposures, we performed ATAC-seq in SAECs following exposure to vehicle,

APM or WSP. Principle Component Analysis (PCA) showed that the ATAC-seq data clusters

predominantly by donor rather than exposure (Supplemental Figure SF2), reflecting the utility of

ATAC-seq to identify basal differences in chromatin architecture in airway epithelial cells

obtained from different donors. We next identified dynamic chromatin accessibility features in

the three SAEC lines following APM exposure (Figure 2A) and compared the control- and APM-

exposed ATAC-seq datasets using Transcription Factor Enrichment Analysis (TFEA).

Significantly enriched or depleted transcription factor binding motifs (p_{adj} < 1e-5) were compared

between SAEC lines (Figure 2B). Despite the divergent baseline chromatin accessibility profiles

between the three donor-derived lines, concordant changes in chromatin accessibility for

specific transcription factor binding motifs were identified across the three lines. A complete list

of these transcription factors is provided in Supplemental Table ST2. In Figure 2C, these data

are visualized for two specific shared motifs using dot plots, which depict the co-clustering of the

GRHL2 motif with depleted chromatin accessibility features and the TEAD3 motif with enriched

features, implying loss of GRHL2 and gain of TEAD3 activity. Using Western Blot, we confirmed

expression of GRHL2 and TEAD3 in all three SAECs and demonstrated that the changes in

factor binding motif enrichment or depletion described above were unrelated to expression

changes (Supplemental Figure SF3).

ATAC-seq distinguishes exposure- and donor-specific responses

To determine whether chromatin responses to particulate matter in SAECs are stereotyped or

instead may be influenced by particulate characteristics, we performed ATAC-seq on Sm2 and

Sm3 SAECs after 20 hours of WSP exposure and compared these responses to APM exposure

responses. We identified a high degree of overlap between ATAC-seq peaks from vehicle- or

WSP-treated samples and control- or APM-treated samples (Figure 3A). We again employed

TFEA to identify transcription factor binding motifs associated with dynamic chromatin

accessibility resulting from exposures and compared significant motifs (p_{adj} < 1e-5) between two

donors and two exposures (Figure 3B). Shared motifs reflected stereotyped chromatin

accessibility responses to exposure, but we also identified divergent motifs that represent

exposure- or donor-specific responses. Complete TFEA results for all significant transcription

factor binding motifs from all four samples are listed in Supplemental Table ST3. Depletion of

P53/P63 motifs, which are extremely similar, was shared between both SAECs and both

exposures, as demonstrated using dot plots (Figure 3C). In contrast to APM exposure, the

TEAD3 motif was not enriched in either Sm2 or Sm3 in response to WSP exposure (Figure 3D).

Thus, the SAEC response to particulate exposure may vary by donor and the underlying

particulate composition, but TEAD3 activity is implicated as a distinguishing feature in chromatin

responses to APM.

Nascent transcript sequencing uncovers transcriptional pathways responsive to APM exposure

We next employed PRO-seq, which identifies nascent transcripts, such as gene transcripts and

short-lived non-coding enhancer RNAs (eRNAs), with high temporal and spatial resolution, to

examine the direct transcriptional response to APM exposure in SAECs. Our previous study

demonstrated marked temporal variation in transcriptional responses within the first two hours of

particulate exposure (22). Therefore, we anticipated that nascent transcription after 20 hours of

exposure represents a steady-state transcriptional response to APM. We performed PRO-seq

on a primary culture of human SAECs, Sm1, after 20 hours of APM exposure and used DESeq2

to identify gene transcription changes induced by APM. We identified 946 genes with increased

transcription and 531 genes with decreased transcription in response to APM (Figure 4A).

Representative PRO-seq genomic tracks visualized in the IGV Genome Browser for *GAS1*

(downregulated) and *CXCL8* (upregulated) are demonstrated in Figure 4B. Gene transcription

data from PRO-seq correlated with mRNA-qPCR from Sm2 and Sm3 cells exposed to APM for

20 hours (Supplemental Figure SF4). DAVID functional annotation (25) identified common

pathways among APM-upregulated genes that included cell-cell adhesion (p = 0.0064) and

keratin-associated signaling (p = 0.00013). Pathways shared among downregulated genes

included cytochrome P450 (p = 2e-6) and glycolysis (p = 0.00064). The top 10 functional

annotation pathways for downregulated and upregulated genes are depicted in Figure 4C. Thus,

aligned with prior reports of particulate exposures, APM exposure causes significant changes in

gene expression associated with specific biologic processes.

Nascent enhancer RNA transcripts identify transcriptional regulatory elements and transcription

factors that mediate responses to APM

It is increasingly recognized that SDTFs integrate the cellular programming of lineage-

determining factors to exact cell-type specific responses to stimuli, such as inflammation or

exposures (33). To identify SDTFs underpinning changes in gene expression caused by APM in

small airway epithelia, we quantitated Transcriptional Regulatory Elements (TREs) using

Transcription-fit (Tfit) (34). Tfit interrogates PRO-seq transcripts for eRNA signatures, which

demarcate TREs, and TRE activity is estimated by quantification of eRNA transcripts at the

specific locus. Differential expression analysis of 20,190 TREs identified 207 TREs with

significantly increased eRNA transcription ($p_{adj} < 0.05$) and 310 TREs with decreased eRNA

transcription (Figure 5A). TFEA identified a series of depleted and enriched transcription factor

binding motifs associated with transcriptional changes at TREs (Figure 5B). We next defined the

intersection between PRO-seq TREs from APM-exposed Sm1 cells and ATAC-seq features

from APM-exposed SAECs based on genomic coordinates (Figure 5C). Simple Enrichment

Analysis (35) identified common motif occurrences within each set of intersected regions

(Supplemental Figure S4F). Representative genome tracks in Figures 5D-E show differentially

transcribed TREs neighboring *MAB21L4* and *AKR1C1*. These regions also exhibit dynamic

chromatin accessibility in select SAEC lines, suggesting that transcription factors mediating

APM exposure responses at these loci exhibit both signal-transduction and lineage-determining

properties.

Integrating genomics approaches delineate signal-transduction and lineage-determining

transcription factor mediators of APM exposure

We reasoned that regions of both dynamic transcription and chromatin accessibility in response

to APM exposure in small airway epithelial progenitor cells likely represent cooperative activity

of SDTFs and LDTFs. To interrogate these regions genome-wide, we re-applied TFEA to

ATAC-seq data from three SAEC lines within TREs called from PRO-seq data. A complete list

of significant motifs ($p_{adj} < 1e-5$) is provided in Supplemental table ST4. Using this approach, we

again examined the TEAD3 and P53/P63 motif changes previously identified in our study.

The TEAD3 motif was not enriched in SAEC ATAC-seq analysis using PRO-seq annotations or

Sm1 PRO-seq (Figure 6A). However, Sm2 and Sm3 cells exhibited increased gene expression

of four canonical TEAD3-target genes following APM exposure, including *CCN1*, *CCN2*, *MYC*

and *FGFBP1* (Figure 6B). Figure 6C demonstrates one example where, despite a significant

increase in *CCN1* gene transcription, neighboring genomic features containing canonical

TEAD3 binding sequences (JASPAR (36)) exhibit increased chromatin accessibility without

changes in eRNA transcription. This suggests that TEAD3 does not directly affect transcriptional

regulatory activity in the manner of a SDTF. Instead, TEAD3, in response to APM exposure,

changes the underlying chromatin landscape to promote expression of target genes, either

through cooperation with an unknown factor or inducing TRE-promoter looping, a function more

characteristic of a LDTF (19).

In contrast to TEAD3, we confirmed depletion of the P53/P63 motif in APM-exposed Sm2 and

Sm3 ATAC-seq samples when examined only within TREs identified by PRO-seq (from Sm1

cells) (Figure 6D). Depletion of the P53 motif was corroborated by a decrease in gene

expression of three canonical P53 target genes, *TP53INP1*, *SESN1* and *DDIT4*, following APM

exposure (Figure 6E). TREs regulating *DDIT4* and *TP53INP1*, with regulation suggested by

concordant transcriptional kinetics, contained canonical P53 binding sequences (JASPAR) and

demonstrated simultaneous transcriptional and chromatin accessibility changes in response to

APM exposure (Figure 6F). These data suggest a combined signal-transduction and lineage-

determining role for TREs harboring P53/P63 motifs, and strongly implicate a role of P53/P63 in

response to APM in airway epithelia.

Discussion

The mechanistic basis for the development of deployment- and exposure-associated respiratory

disease is poorly understood. Here, we approached this problem by exposing primary SAECs to

APM, a highly relevant deployment-specific exposure, and characterizing the genomic

responses. We identified several pathways that may be involved in the development of DRRD

and potential neoplastic implications of deployment exposures, such as loss of P53/P63,

GRHL2 or cytochrome P450 signaling. Furthermore, we identified transcription factors that

mediate transcriptional responses, such as the EGR family, factors that affect chromatin

accessibility change, such as the TEAD family, or factors with shared function, such as the

P53/P63 family. We identified pathways specific to APM exposure by comparing effects of APM

and WSP exposure, with TEAD activation, a pathway that previously had not been associated

with particulate exposures, as specific to APM. In aggregate, our data indicate that APM

exposure in human airway epithelial cells causes broad changes in transcription factor activity,

with certain effects specific to APM in comparison to WSP.

The combined application of PRO-seq and ATAC-seq allowed us to partition responses to APM

exposure into two categories. PRO-seq identifies direct transcriptional responses to a stimulus

while ATAC-seq profiles changes in chromatin architecture. Although both responses are

important to cell fate and function, we hypothesize that transcription factors that mediate these

changes fall under two distinct but overlapping categories. LDTFs regulate the underlying

chromatin architecture of the cell and, dependent on cell type or memory from previous stimuli,

establish the chromatin context for subsequent exposures (20). We interpreted responses

detected in ATAC-seq profiles to reflect the function of LDTFs. SDTFs directly control

transcriptional responses within the context of chromatin architecture established by LDTFs

(33). We interpreted PRO-seq data to reflect the functions of SDTFs. Using TFEA, we identified

transcription factors from ATAC-seq and PRO-seq data and empirically classified them under

these two categories.

We identified important SDTF pathways in SAECs exposed to APM that affect disease

susceptibility to additional insults and support a multi-hit hypothesis for the development of

DRRD. For example, APM exposure resulted in downregulation of cytochrome P450 signaling,

which is responsible for catabolism of polyaromatic compounds (37). Even transient loss of this

program can result in increased susceptibility to additional environmental exposures, as

increased polyaromatic hydrocarbon composition in particulate exposures is associated with

increased cytotoxicity (38) and several polyaromatic compounds are known carcinogens (39).

Thus, a loss of cytochrome activity may predispose to aberrant responses and additional injury

in response to repetitive exposures, which are typically a feature of DRRD.

Our data indicate that APM-mediated activation of the TEAD family of transcription factors is a

unique feature of APM that relates directly to small airway epithelia lineage. In contrast to APM,

WSP exposure did not result in TEAD enrichment, and with APM-exposure, TEAD enrichment

was demonstrated only in dynamic chromatin accessibility features identified by ATAC-seq and

not TREs identified by PRO-seq. TEAD transcription factors are classically identified as

downstream effectors of the Hippo-YAP-TAZ signal transduction pathway that promote

development and cell proliferation (40). Small molecules have been described that can activate

or block Hippo pathway signaling (41, 42), and the TEAD family may represent a potential

therapeutic target for the treatment of DRRD or mitigation of toxic responses to deployment-

related exposures. Ultimately, further investigation is necessary to characterize the impact of

TEAD-associated chromatin accessibility gain on subsequent exposures or cell fate.

P53/P63 signaling responses to APM were identified by interrogation of dynamic TREs from

PRO-seq and chromatin accessibility features from ATAC-seq, suggesting a combined signal-

dependent and lineage-determining role for P53/P63 in small airway epithelia progenitor cells.

This latter function aligns with numerous previous studies of the airway epithelia (43, 44, 46,

48), however, the integrated role of this family in exposure responses has not been reported to

our knowledge. The combined role may reflect differential activity of distinct P53/P63 family

members or the effects of transcription factors that cooperate with P53/P63. For example,

GRHL2, another LDTF identified by our study, has been demonstrated to work with P53/P63 in

determining epithelial cell fate at the epigenetic level (45).

The importance of P53/P63 signaling in APM exposure is further corroborated by a series of

studies that connect dysregulations in this family with bronchiolitis and small airways disease.

Reduced P53 activity has previously been implicated in distal bronchiolar pathology and a

failure of apoptosis in response to RSV infection (47), a common cause of acute bronchiolitis.

Moreover, P63 expression is reduced in patients with constrictive bronchiolitis arising from

mustard gas exposure (49). Although additional studies are needed to establish a causal

relationship, our work suggests that reduced P53/P63 activity may predispose to bronchiolitis in

the setting of APM exposure, either acting independently with multiple transcription factor roles

or in association with additional factors, such as GRHL2. Further investigation of dynamic

P53/P63 motif-containing TREs and chromatin accessibility features identified in our study may

delineate the specific factors responsible for these multiple P53/P63 roles and determine how

signal-dependent responses integrate with lineage factors in determining airway epithelia

response to exposures. Finally, increasing P53 activity via agents such as idasanutlin (50),

which is currently in clinical development for myeloid cancers (51), is feasible and may provide

another therapeutic avenue in DRRD.

The study of exposure biology has multiple layers of complexity, which contribute to the poor

understanding of DRRD. Factors such as concentration or duration of exposure are highly

variable in patients with DRRD and difficult to model. Even in model system studies such as

ours, the partition between homeostatic and pathophysiologic responses to exposure is not

clear. However, ATAC-seq is one genomic method that, given the low input cell number, can be

directly applied to patient-derived samples. In this study, we explored the utility of ATAC-seq to

identify mechanisms of APM exposure and identified the TEAD family of transcription factors, a

novel pathway not previously associated with particulate exposure. Furthermore, we

demonstrated that the intersection of ATAC-seq data from multiple SAEC lines with TREs

identified from a single PRO-seq experiment delineates a subclass of LDTFs that are shared

between SAECs. Future studies that apply ATAC-seq to patient samples, such as small airway

epithelial cells from distal bronchoscopic brush biopsy, can explore dysregulation in these core

lineage factors by integrating those data with the TRE set identified by PRO-seq in this study. In

addition, we used this set of TREs to investigate dynamic chromatin accessibility in multiple

SAEC lines. Through this approach, we were able to delineate which transcription factors

identified by ATAC-seq were consistently regulated by APM exposure or exhibited signal-

dependent properties. Thus, future studies applying ATAC-seq to a variety of dynamic samples

can use annotations from prior PRO-seq studies to classify transcription factors identified by the

analysis. Given the high input requirements for PRO-seq, such as large cell numbers, time and

sequencing cost, the annotation approach provides an easy-to-implement option when patient

or exposure sampling methods preclude paired PRO-seq and ATAC-seq library preparation.

Although our study was not specifically powered for detecting individual-level variations in

exposure responses, we demonstrated variation in genomic response patterns between SAECs

from different donors. For example, GRHL2 loss was present in Sm1 and Sm3 but not Sm2.

This variation may depend on several factors, such as age, sex, systemic disease or exposure

history. Future studies that incorporate donor-level data can investigate the effect of systemic

factors or genetic variants on responses to APM exposure. Finally, our approach also detected

differences between the genomic responses to APM versus WSP exposures. It is tempting to

speculate that these differences in our reductionist model portend the well-known clinical

differences in response to deployment-associated particulates and other more common sources

of PM.

Our model system had several limitations. SAECs were cultured and exposed in submerged cell

culture without differentiation into a cuboidal ciliated epithelium, which is characteristic of small

airways (52). The dose and composition of exposures may also diverge between our model and

true inhalational exposures experienced by military personnel deployed to Southwest Asia.

However, despite these limitations, our study identified several plausible and novel features of

APM exposure that could only be identified through the use of sophisticated genomics tools in

our model system. Extending our findings to *in vivo* models and even more sophisticated *in vitro*

systems, such as precision cut lung slices, will be crucial for validating the novel APM-

associated pathways we have reported here.

References

1.   **Krefft SD**, **Meehan R**, **Rose CS**. Emerging spectrum of deployment-related respiratory

diseases. *Curr. Opin. Pulm. Med.* 21 Lippincott Williams and Wilkins: 185–192, 2015.

2.   **Krefft SD**, **Rose CS**, **Nawaz S**, **Miller YE**. Deployment-Related Lung Disorders. [Online].

Fed Pract 32: 32–38, 2015. http://www.ncbi.nlm.nih.gov/pubmed/30766070 [24 Nov.

2020].

3.   **Kim KH**, **Kabir E**, **Kabir S**. A review on the human health impact of airborne particulate
matter. *Environ Int* 74: 136–143, 2015. doi: 10.1016/J.ENVINT.2014.10.005.

4.   **Li R**, **Zhou R**, **Zhang J**. Function of PM2.5 in the pathogenesis of lung cancer and
chronic airway inflammatory diseases. *Oncol. Lett.* 15 Spandidos Publications: 7506–
7514, 2018.

5.   **Hamanaka RB**, **Mutlu GM**. Particulate Matter Air Pollution: Effects on the Cardiovascular
System. *Front Endocrinol (Lausanne)* 9: 680, 2018. doi: 10.3389/FENDO.2018.00680.

6.   **Lee BJ**, **Kim B**, **Lee K**. Air Pollution Exposure and Cardiovascular Disease. *Toxicol Res*
30: 71, 2014. doi: 10.5487/TR.2014.30.2.071.

7.   **Calderón-Garcidueñas L**, **Stommel EW**, **Rajkumar RP**, **Mukherjee PS**, **Ayala A**.
Particulate Air Pollution and Risk of Neuropsychiatric Outcomes. What We Breathe,
Swallow, and Put on Our Skin Matters. *Int J Environ Res Public Health* 18: 11568, 2021.
doi: 10.3390/IJERPH182111568.

8.   **Ghio AJ**, **Soukup JM**, **Case M**, **Dailey LA**, **Richards J**, **Berntsen J**, **Devlin RB**, **Stone**
S, **Rappold A**. Exposure to wood smoke particles produces inflammation in healthy
volunteers. *Occup Environ Med* 69: 170–175, 2012. doi: 10.1136/oem.2011.065276.

9.   **Ghio AJ**, **Soukup JM**, **Dailey LA**, **Tong H**, **Kesic MJ**, **Budinger GRS**, **Mutlu GM**. Wood
Smoke Particle Sequesters Cell Iron to Impact a Biological Effect. *Chem Res Toxicol* 28:
2104–11, 2015. doi: 10.1021/acs.chemrestox.5b00270.

10.   **Memon TA**, **Nguyen ND**, **Burrell KL**, **Scott AF**, **Almestica-Roberts M**, **Rapp E**,
Deering-Rice CE, **Reilly CA**. Wood Smoke Particles Stimulate MUC5AC Overproduction
by Human Bronchial Epithelial Cells Through TRPA1 and EGFR Signaling. *Toxicol Sci*
174: 278–290, 2020. doi: 10.1093/toxsci/kfaa006.

11.   **Cao Y**, **Chen M**, **Dong D**, **Xie S**, **Liu M**. Environmental pollutants damage airway
epithelial cell cilia: Implications for the prevention of obstructive lung diseases. *Thorac*
Cancer 11: 505, 2020. doi: 10.1111/1759-7714.13323.

12.   **Kyung SY**, **Jeong SH**. Particulate-matter related respiratory diseases. *Tuberc. Respir.*
Dis. (Seoul). 83 Korean National Tuberculosis Association: 116–121, 2020.

13.   **Chen YC**, **Chuang TY**, **Liu CW**, **Liu CW**, **Lee TL**, **Lai TC**, **Chen YL**. Particulate matters
increase epithelial-mesenchymal transition and lung fibrosis through the ETS-1/NF-κB-
dependent pathway in lung epithelial cells. *Part Fibre Toxicol* 17: 1–14, 2020. doi:
10.1186/S12989-020-00373-Z/FIGURES/8.

14.   **Maes T**, **Provoost S**, **Lanckacker EA**, **Cataldo DD**, **Vanoirbeek JAJ**, **Nemery B**,
Tournoy KG, **Joos GF**. Mouse models to unravel the role of inhaled pollutants on

allergic sensitization and airway inflammation. *Respir Res 2010 111* 11: 1–16, 2010. doi:
10.1186/1465-9921-11-7.

15.   **Berman R**, **Kopf KW**, **Min E**, **Huang J**, **Downey GP**, **Alam R**, **Chu HW**, **Day BJ**. IL-
547   33/ST2 signaling modulates Afghanistan particulate matter induced airway
548   hyperresponsiveness in mice. *Toxicol Appl Pharmacol* 404: 115186, 2020. doi:
549   10.1016/j.taap.2020.115186.

550   16.   **Berman R**, **Downey GP**, **Dakhama A**, **Day BJ**, **Chu HW**. Afghanistan Particulate Matter
551   Enhances Pro-Inflammatory Responses in IL-13-Exposed Human Airway Epithelium via
552   TLR2 Signaling. *Toxicol Sci* 166: 345–353, 2018. doi: 10.1093/TOXSCI/KFY217.

553   17.   **A G**, **SK S**, **MA G**, **L S**, **RD D**, **AN G**. Deconvolution of multiplexed transcriptional
554   responses to wood smoke particles defines rapid aryl hydrocarbon receptor signaling
555   dynamics. .

556   18.   **Troutman TD**, **Kofman E**, **Glass CK**. Exploiting dynamic enhancer landscapes to
557   decode macrophage and microglia phenotypes in health and disease. *Mol Cell* 81: 3888–
558   3903, 2021. doi: 10.1016/J.MOLCEL.2021.08.004.

559   19.   **Miyazaki K**, **Miyazaki M**. The Interplay Between Chromatin Architecture and Lineage-
560   Specific Transcription Factors and the Regulation of Rag Gene Expression. *Front*
561   *Immunol* 12, 2021. doi: 10.3389/FIMMU.2021.659761/FULL.

562   20.   **Hao R-H**, **Guo Y**, **Wang C**, **Chen F**, **Di C-X**, **Dong S-S**, **Cao Q-L**, **Guo J**, **Rong Y**, **Yao**
563   **S**, **Zhu D-L**, **Chen Y-X**, **Chen H**, **Yang T-L**. Lineage-specific rearrangement of chromatin
564   loops and epigenomic features during adipocytes and osteoblasts commitment. .

565   21.   **Zuo W**, **Zhang T**, **Wu DZA**, **Guan SP**, **Liew AA**, **Yamamoto Y**, **Wang X**, **Lim SJ**,
566   **Vincent M**, **Lessard M**, **Crum CP**, **Xian W**, **McKeon F**. p63+Krt5+ distal airway stem
567   cells are essential for lung regeneration. *Nature* 517: 616, 2015. doi:
568   10.1038/NATURE13903.

569   22.   **Gupta A**, **Sasse SK**, **Gruca MA**, **Sanford L**, **Dowell RD**, **Gerber AN**. Deconvolution of
570   multiplexed transcriptional responses to wood smoke particles defines rapid aryl
571   hydrocarbon receptor signaling dynamics. *J Biol Chem* 0: 101147, 2021. doi:
572   10.1016/J.JBC.2021.101147.

573   23.   **Sasse SK**, **Gruca M**, **Allen MA**, **Kadiyala V**, **Song T**, **Gally F**, **Gupta A**, **Pufall MA**,
574   **Dowell RD**, **Gerber AN**. Nascent transcript analysis of glucocorticoid crosstalk with TNF
575   defines primary and cooperative inflammatory repression. *Genome Res* 29: 1753–1765,
576   2019. doi: 10.1101/gr.248187.119.

577   24.   **Gally F**, **Sasse SK**, **Kurche JS**, **Gruca MA**, **Cardwell JH**, **Okamoto T**, **Chu HW**, **Hou X**,

Poirion OB, **Buchanan J**, **Preissl S**, **Ren B**, **Colgan SP**, **Dowell RD**, **Yang I V.**, **Schwartz DA**, **Gerber AN**. The MUC5B-associated variant rs35705950 resides within an enhancer subject to lineage- and disease-dependent epigenetic remodeling. *JCI Insight* 6, 2021. doi: 10.1172/JCI.INSIGHT.144294.

25. **Huang DW**, **Sherman BT**, **Lempicki RA**. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44–57, 2009. doi: 10.1038/nprot.2008.211.

26. **Corces MR**, **Trevino AE**, **Hamilton EG**, **Greenside PG**, **Sinnott-Armstrong NA**, **Vesuna S**, **Satpathy AT**, **Rubin AJ**, **Montine KS**, **Wu B**, **Kathiria A**, **Cho SW**, **Mumbach MR**, **Carter AC**, **Kasowski M**, **Orloff LA**, **Risca VI**, **Kundaje A**, **Khavari PA**, **Montine TJ**, **Greenleaf WJ**, **Chang HY**. An improved ATAC-seq protocol reduces background and enablesinterrogation of frozen tissues. *Nat Methods* 14: 959, 2017. doi: 10.1038/NMETH.4396.

27. **Rubin JD**, **Stanley JT**, **Sigauke RF**, **Levandowski CB**, **Maas ZL**, **Westfall J**, **Taatjes DJ**, **Dowell RD**. Transcription factor enrichment analysis (TFEA) quantifies the activity of multiple transcription factors from a single experiment. *Commun Biol 2021 41* 4: 1–15, 2021. doi: 10.1038/s42003-021-02153-7.

28. **Grant CE**, **Bailey TL**, **Noble WS**. FIMO: Scanning for occurrences of a given motif. *Bioinformatics* 27: 1017–1018, 2011. doi: 10.1093/bioinformatics/btr064.

29. **Lambert SA**, **Jolma A**, **Campitelli LF**, **Das PK**, **Yin Y**, **Albu M**, **Chen X**, **Taipale J**, **Hughes TR**, **Weirauch MT**. The Human Transcription Factors. *Cell* 172 Cell Press: 650–665, 2018.

30. **Altonsy MO**, **Sasse SK**, **Phang TL**, **Gerber AN**. Context-dependent cooperation between nuclear factor κB (NF-κB) and the glucocorticoid receptor at a TNFAIP3 intronic enhancer: A mechanism to maintain negative feedback control of inflammation. *J Biol Chem* 289: 8231–8239, 2014. doi: 10.1074/jbc.M113.545178.

31. **Allen RW**, **Mar T**, **Koenig J**, **Liu LJS**, **Gould T**, **Simpson C**, **Larson T**. Changes in lung function and airway inflammation among asthmatic children residing in a woodsmoke-impacted urban area. *Inhal Toxicol* 20: 423–433, 2008. doi: 10.1080/08958370801903826.

32. **White SM**, **Snyder MP**, **Yi C**. Master lineage transcription factors anchor trans mega transcriptional complexes at highly accessible enhancer sites to promote long-range chromatin clustering and transcription of distal target genes. *Nucleic Acids Res* 49: 12196–12210, 2021. doi: 10.1093/NAR/GKAB1105.

612   33.   **Heinz S**, **Benner C**, **Spann N**, **Bertolino E**, **Lin YC**, **Laslo P**, **Cheng JX**, **Murre C**,
613   **Singh H**, **Glass CK**. Simple combinations of lineage-determining transcription factors
614   prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* 38:
615   576, 2010. doi: 10.1016/J.MOLCEL.2010.05.004.

616   34.   **Azofeifa JG**, **Dowell RD**. A generative model for the behavior of RNA polymerase.
617   *Bioinformatics* 33: 227–234, 2017. doi: 10.1093/bioinformatics/btw599.

618   35.   **Bailey TL**, **Grant CE**. SEA: Simple Enrichment Analysis of motifs. .

619   36.   **Castro-Mondragon JA**, **Riudavets-Puig R**, **Rauluseviciute I**, **Berhanu Lemma R**,
620   **Turchi L**, **Blanc-Mathieu R**, **Lucas J**, **Boddie P**, **Khan A**, **Perez NM**, **Fornes O**, **Leung**
621   **TY**, **Aguirre A**, **Hammal F**, **Schmelter D**, **Baranasic D**, **Ballester B**, **Sandelin A**,
622   **Lenhard B**, **Vandepoele K**, **Wasserman WW**, **Parcy F**, **Mathelier A**. JASPAR 2022:
623   The 9th release of the open-access database of transcription factor binding profiles.
624   *Nucleic Acids Res* 50: D165–D173, 2022. doi: 10.1093/NAR/GKAB1113.

625   37.   **Shimada T**, **Fujii-Kuriyama Y**. Metabolic activation of polycyclic aromatic hydrocarbons
626   to carcinogens by cytochromes P450 1A1 and 1B1. *Cancer Sci* 95: 1–6, 2004. doi:
627   10.1111/J.1349-7006.2004.TB03162.X.

628   38.   **Wang B**, **Li N**, **Deng F**, **Buglak N**, **Park G**, **Su S**, **Ren A**, **Shen G**, **Tao S**, **Guo X**. Human
629   bronchial epithelial cell injuries induced by fine particulate matter from sandstorm and
630   non-sandstorm periods: Association with particle constituents. *J Environ Sci (China)* 47:
631   201–210, 2016. doi: 10.1016/J.JES.2015.12.015.

632   39.   **Hecht SS**. Potential Carcinogenic Effects of Polynuclear Aromatic Hydrocarbons and
633   Nitroaromatics in Mobile Source Emissions - Air Pollution, the Automobile, and Public
634   Health - NCBI Bookshelf [Online]. [date unknown].
635   https://www.ncbi.nlm.nih.gov/books/NBK218134/ [11 Apr. 2022].

636   40.   **Lin KC**, **Park HW**, **Guan KL**. Regulation of the Hippo pathway transcription factor TEAD.
637   *Trends Biochem Sci* 42: 862, 2017. doi: 10.1016/J.TIBS.2017.09.003.

638   41.   **Aihara A**, **Iwawaki T**, **Abe-Fukasawa N**, **Otsuka K**, **Saruhashi K**, **Mikashima T**,
639   **Nishino T**. Small molecule LATS kinase inhibitors block the Hippo signaling pathway and
640   promote cell growth under 3D culture conditions. *J Biol Chem* 298: 101779, 2022. doi:
641   10.1016/J.JBC.2022.101779/ATTACHMENT/46711C20-8AFE-46B9-AC3A-
642   A429C72D720A/MMC1.PDF.

643   42.   **Crawford JJ**, **Bronner SM**, **Zbieg JR**. Hippo pathway inhibition by blocking the
644   YAP/TAZ-TEAD interface: a patent review. *Expert Opin Ther Pat* 28: 867–873, 2018. doi:
645   10.1080/13543776.2018.1549226.

43.   **Uddin MA**, **Barabutis N**. P53 in the impaired lungs. *DNA Repair (Amst)* 95: 102952,
2020. doi: 10.1016/J.DNAREP.2020.102952.

44.   **Garrido-Jimenez S**, **Barrera-Lopez JF**, **Diaz-Chamorro S**, **Mateos-Quiros CM**,
Rodriguez-Blanco I, **Marquez-Perez FL**, **Lorenzo MJ**, **Centeno F**, **Roman AC**,
Carvajal-Gonzalez JM. p53 regulation by MDM2 contributes to self-renewal and
differentiation of basal stem cells in mouse and human airway epithelium. *FASEB J* 35:
e21816, 2021. doi: 10.1096/FJ.202100638R.

45.   **Latil M**, **Nassar D**, **Beck B**, **Boumahdi S**, **Wang L**, **Brisebarre A**, **Dubois C**, **Nkusi E**,
Lenglez S, **Checinska A**, **Vercauteren Drubbel A**, **Devos M**, **Declercq W**, **Yi R**,
Blanpain C. Cell-Type-Specific Chromatin States Differentially Prime Squamous Cell
Carcinoma Tumor-Initiating Cells for Epithelial to Mesenchymal Transition. *Cell Stem Cell*
20: 191-204.e5, 2017. doi: 10.1016/J.STEM.2016.10.018.

46.   **Warner SMB**, **Hackett TL**, **Shaheen F**, **Hallstrand TS**, **Kicic A**, **Stick SM**, **Knight DA**.
Transcription factor p63 regulates key genes and wound repair in human airway epithelial
Basal cells. *Am J Respir Cell Mol Biol* 49: 978–988, 2013. doi: 10.1165/RCMB.2012-
0447OC/SUPPL_FILE/DISCLOSURES.PDF.

47.   **Eckardt-Michel J**, **Lorek M**, **Baxmann D**, **Grunwald T**, **Keil GM**, **Zimmer G**. The Fusion
Protein of Respiratory Syncytial Virus Triggers p53-Dependent Apoptosis. *J Virol* 82:
3236, 2008. doi: 10.1128/JVI.01887-07.

48.   **Bilodeau C**, **Shojaie S**, **Goltsis O**, **Wang J**, **Luo D**, **Ackerley C**, **M Rogers I**, **Cox B**,
Post M. TP63 basal cells are indispensable during endoderm differentiation into proximal
airway cells on acellular lung scaffolds. *npj Regen Med 2021 61* 6: 1–12, 2021. doi:
10.1038/s41536-021-00124-4.

49.   **Ghanei M**, **Chilosi M**, **Akbari HMH**, **Motiei-Langroudi R**, **Harandi AA**, **Shamsaei H**,
Bahadori M, **Tazelaar HD**. Use of Immunohistochemistry Techniques in Patients
Exposed to Sulphur Mustard Gas. *Patholog Res Int* 2011: 1–7, 2011. doi:
10.4061/2011/659603.

50.   **Khurana A**, **Shafer DA**. MDM2 antagonists as a novel treatment option for acute myeloid
leukemia: perspectives on the therapeutic potential of idasanutlin (RG7388). *Onco*
Targets Ther 12: 2903, 2019. doi: 10.2147/OTT.S172315.

51.   **Konopleva MY**, **Röllig C**, **Cavenagh J**, **Deeren D**, **Girshova L**, **Krauter J**, **Martinelli G**,
Montesinos P, **Schäfer JA**, **Ottmann OG**, **Petrini M**, **Pigneux A**, **Rambaldi A**, **Recher**
C, **Rodriguez-Veiga R**, **Taussig D**, **Vey N**, **Yoon S-S**, **Ott MG**, **Muehlbauer S**,
Beckermann BM, **Catalani O**, **Genevray M**, **Mundt KE**, **Jamois C**, **Fenaux P**, **Wei AH**.

Idasanutlin plus cytarabine in relapsed or refractory acute myeloid leukemia: results of

the MIRROS trial. *Blood Adv* 6: 4147–4156, 2022. doi:

10.1182/BLOODADVANCES.2021006303.

52.   **van den Bosch WB**, **James AL**, **Tiddens HAWM**. Structure and function of small

airways in asthma patients revisited. *Eur Respir Rev* 30: 1–18, 2021. doi:

10.1183/16000617.0186-2020.

40**.**   **Bushnell B.** BBMap: A Fast, Accurate, Splice-Aware Aligner [Online]. *United States N.*

p., 2014: [date unknown]. https://www.osti.gov/biblio/1241166-bbmap-fast-accurate-

splice-aware-aligner [9 Aug. 2021].

41.   **Andrews S.** FastQC: A Quality Control Tool for High Throughput Sequence Data [Online]

[Online]. 2010. http://www.bioinformatics.babraham.ac.uk/projects/fastqc/ [9 Aug. 2021].

42.   **Azofeifa J, Allen MA, Lladser M, Dowell R.** FStitch: A fast and simple algorithm for

detecting nascent RNA transcripts. In: *ACM BCB 2014 - 5th ACM Conference on*

Bioinformatics, Computational Biology, and Health Informatics. Association for Computing

Machinery, Inc, p. 174–183.

43.   **Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G,**

Mesirov JP. Integrative genomics viewer. *Nat. Biotechnol.* 29 NIH Public Access: 24–26,

2011.

44.   **Quinlan AR, Hall IM.** BEDTools: A flexible suite of utilities for comparing genomic

features. *Bioinformatics* 26: 841–842, 2010. doi: 10.1093/bioinformatics/btq033.

45.   **Kim D, Paggi JM, Park C, Bennett C, Salzberg SL.** Graph-based genome alignment

and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37: 907–915, 2019.

doi: 10.1038/s41587-019-0201-4.

46.   **Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,**

Durbin R. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078–

2079, 2009. doi: 10.1093/bioinformatics/btp352.

47.   **Liao Y, Smyth GK, Shi W.** FeatureCounts: An efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics* 30: 923–930, 2014. doi:

10.1093/bioinformatics/btt656.

48.   **Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C,**

Myers RM, Brown M, Li W, Liu XS. Model-based analysis of ChIP-Seq (MACS).

Genome Biol 9: R137, 2008. doi: 10.1186/gb-2008-9-9-r137.

49.   **Love MI**, **Huber W**, **Anders S**. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* 15, 2014. doi: 10.1186/S13059-014-0550-8.

Figure Legends

Figure 1 – Particulate exposure induces IL-8 secretion in SAECs derived from three

independent healthy donors. ELISA analysis of IL-8 concentration in the supernatant of Sm1,

Sm2 and Sm3 SAECs exposed to (A) APM (50 ug/cm^2 of well surface area) or (B) WSP (1

mg/ml of cell culture media) for 20 hours in submerged cell culture. Bars indicate mean IL-8

concentration ($\pm$SD); n $\geq$ 3/group, *p < 1e-3 for indicated comparisons.

Figure 2 – ATAC-seq identifies transcription factors associated with chromatin

accessibility changes in response to APM exposure. (A) Pyramid plots illustrate MACS2-

called ATAC-seq peaks (discrete dots) ranked by magnitude of accessibility gain (more red) or

loss (more green) in response to APM exposure, as output during Transcription Factor

Enrichment Analysis (TFEA). The total number (N) of unique loci for each SAEC line is listed

below each plot. (B) Venn diagrams depict TFEA-identified numbers of transcription factor

binding motifs for each SAEC line that were significantly (p_{adj} < 1e-5) depleted (motifs that

cluster with ATAC-seq features that lose accessibility in response to APM) or enriched (motifs

that cluster with ATAC-seq features that gain accessibility in response to APM) following APM

exposure and their overlap between SAEC lines. (C) Representative dot plots for one depleted

(GRHL2 – shared between Sm1 and Sm3, top right of Venn diagram in panel B) and one

enriched (TEAD3 – shared between Sm1, Sm2 and Sm3, center of Venn diagram in panel B)

motif identified by TFEA. Each dot represents the indicated motif encountered within $\pm$1500 bp

of an accessibility region, ranked by accessibility gain on the horizontal axis and distance to

accessibility region center (with midpoint = 0) on the vertical axis. Enrichment (E) scores (a

relative measure of motif enrichment between control vs APM exposure) and associated p_{adj}

values are listed, with significant p_{adj} values indicated in red.

Figure 3 – Chromatin accessibility profiles following APM vs WSP exposure distinguish

exposure- and donor-specific responses. (A) Venn diagrams delineate overlap of MACS2-

called ATAC-seq peaks identified in Sm2 or Sm3 cells exposed to APM vs WSP based on

genomic coordinates. Numbers below Venn diagrams indicate the fraction of peaks from each

indicated comparison found in the shared (pink) set. (B) Venn diagrams show TFEA-identified

numbers of significantly (p_{adj} < 1e-5) enriched or depleted transcription factor binding motifs

following APM or WSP exposure and their overlap, both by exposure and by donor, as

described for Fig. 2B. (C) Dot plots for P53 and P63 binding motifs that shared depletion in Sm2

and Sm3 in response to both APM and WSP exposures; significant depletion indicated in red

text. (D) Dot plots for the TEAD3 binding motif that was enriched in Sm2 and Sm3 exposed to

APM but not WSP; significant enrichment indicated in blue text.

Figure 4 – PRO-seq reveals sustained transcriptional programs associated with APM

exposure. (A) Volcano plot illustrates differentially regulated nascent gene transcripts in Sm1

cells treated with APM for 20 hours. Each point represents one gene, with red indicating p_{adj} <

0.05, light orange representing $\log_2$ fold change > 1, green indicating both p_{adj} < 0.05 and $\log_2$

fold change > 1 and black indicating neither condition was satisfied. The number of genes

meeting both criteria (green dots) that were down- or upregulated is shown. (B) Representative

examples of downregulated (left) and upregulated (right) genes shown as PRO-seq tracks

visualized in the Integrative Genomics Viewer (IGV) genome browser based on counts per

million mapped reads (vertical scales). Positive (blue) indicates reads annotated to the sense

strand while negative (red) data reflect reads annotated to the antisense strand. The TSS and

direction of transcription are indicated by arrows at the top of each panel. (C) Bar graphs display

top 10 most significantly enriched functional annotation terms output by DAVID Functional

Annotation Clustering applied to downregulated (left) and upregulated (right) transcripts.

Figure 5 – TREs defined by PRO-seq bidirectional signatures identify a distinct set of

transcription factors that respond to APM exposure. (A) Volcano plot illustrates differentially

regulated Transcriptional Regulatory Element (TRE) transcripts in Sm1 cells treated with APM

for 20 hours, as described for Fig. 4A. (B) Transcription factor binding motifs identified by TFEA

within PRO-seq data are represented as bar graphs. In this case, depleted motifs refers to those

encountered more frequently in TREs with loss of transcriptional signal, while enriched motifs

are those encountered more frequently in TREs with gain in transcriptional signal. Motifs listed

meet the threshold of p_{adj} < 1e-5 and are ranked by E score, here a relative measure of motif

enrichment within TREs. (C) Venn diagrams depict the total numbers of control and APM-

exposed ATAC-seq peaks from each SAEC line (ATAC-seq), control and APM-exposed PRO-

seq TREs from Sm1 cells (PRO-seq), and their overlap, based on genomic coordinates.

Numbers below each diagram represent the fraction of ATAC-seq or PRO-seq peaks found

within the intersection for each SAEC (pink). Representative examples of one upregulated TRE

near the *MAB21L4* promoter (D) and one downregulated TRE 5' to the *AKR1C1* gene locus (E)

shown as PRO-seq tracks from Sm1 exposed to APM and ATAC-seq tracks from three SAEC

lines exposed to APM, as visualized in IGV. ATAC-seq tracks are color-coded by SAEC line,

with green = Sm1, purple = Sm2, and cyan = Sm3. Vertical scales on all tracks indicate counts

per million mapped reads.

Figure 6 – APM-responsive transcription factors exhibit different classes of

transcriptional regulation. (A) Dot plots demonstrate no TEAD3 binding motif enrichment

when PRO-seq-defined TREs are used to define regions for TFEA analysis of control vs APM-

exposed SAECs. (B) qRT-PCR analysis of indicated gene expression in Sm2 or Sm3 cells

following APM exposure for canonical TEAD targets *CCN1*, *CCN2*, *MYC* and *FGFBP1*. Bars

represent mean normalized C_T values on a log_2 scale (±SD) relative to vehicle-treated controls;

n = 4/group, *p < 0.05 vs. vehicle. (C) IGV-visualized tracks of Sm1 control- and APM-exposed

PRO-seq samples and three SAEC control- and APM-exposed ATAC-seq samples for one

dynamic chromatin accessibility region 5' to *CCN1* (boxed in black and magnified). Specific

regions containing canonical TEAD3 binding sequences (JASPAR) are indicated as motif logos

under the genome tracks. (D) TFEA analysis of control vs APM-exposed Sm2 and Sm3 ATAC-

seq data within PRO-seq defined regions confirms significant (p_{adj} < 1e-5) depletion of the P53

binding motif, illustrated by dot plots. (E) qRT-PCR analysis of indicated gene expression in

Sm2 or Sm3 cells following APM exposure for canonical P53 targets *TP53INP1, SESN1* and

DDIT4; n = 4/group, *p < 0.05 vs. vehicle. (F) IGV-visualized tracks of Sm1 control- and APM-

exposed PRO-seq samples and three SAEC control- and APM-exposed ATAC-seq samples for

one TRE 5' to the *DDIT4* gene locus and one TRE 5' to the *TP53INP1* gene locus (boxed in

black and magnified). Both TREs contain canonical P53 binding sequences (JASPAR) as

indicated.

Supplemental Items are available at:

https://figshare.com/projects/Integrated_Genomics_Approaches_Identify_Mediators_of_Transcri

ptional_and_Epigenetic_Responses_to_Afghan_Desert_Particulate_Matter_in_Human_Small_

Airway_Epithelial_Cells/138039

A Downregulated = 531   Upregulated = 946

B PRO-seq

GAS1

1000 bp

Ctrl

Sm1

APM

0.25
-0.46

0.25
-0.46

CXCL8

1000 bp

Ctrl

Sm1

APM

0.38
-0.04

0.38
-0.04

C

Downregulated

- Cytochrome P450
- Aldo/keto reductase
- Oxidoreductase
- Histidine metabolism
- Retinol dehydrogenase activity
- Glycolysis/Gluconeogenesis
- Steroid hormone biosynthesis
- Lipid degradation
- Galactose detabolism
- EF-hand 2 domain

Enrichment Score

Upregulated

- Serine protease
- SH2 domain
- Cell-cell adhesion
- IL-1 receptor family
- LIM zinc-binding
- Serum Amyloid A protein
- Keratin-associated protein
- EF-hand-like domain
- Insulin family
- Positive regulation of cell division

Enrichment Score