Cocoa extract exerts sex-specific anti-diabetic effects in an aggressive type-2 diabetes

model: a pilot study

Kathryn C. Racine[a,1], Lisard Iglesias-Carres[a], Jacob A. Herring[b], Mario G. Ferruzzi[a,2], Colin D. Kay[a], Jeffery S. Tessem[b], Andrew P. Neilson[a]

[a]Plants for Human Health Institute, North Carolina State University, 600 Laureate Way, Kannapolis, NC 28081, USA; kcracine@ncsu.edu, liglesi@ncsu.edu, mferruz@ncsu.edu cdkay@ncsu.edu, aneilso@ncsu.edu

[b]Department of Nutrition, Dietetics, & Food Science, Brigham Young University, S-243 ESC Provo, UT 84042, USA; jeffery_tessem@byu.edu, herrin06@gmail.com

[1]Current affiliation: Mars, Inc., Chicago, IL, USA

[2]Current affiliation: Arkansas Children's Nutrition Center, U.S. Department of Agriculture and Department of Pediatrics, University of Arkansas Center for Medical Science, Little Rock, AR, USA; MFerruzzi@uams.edu

[*]**Corresponding Author:** Dr. Andrew P. Neilson, email: aneilso@ncsu.edu, phone: +1 (704) 250-5495

Abstract

Type 2 diabetes (T2D) is characterized by hyperglycemia and insulin resistance. Cocoa may

slow T2D development and progression. This study employed male and female BTBR.Cg-

$Lep^{ob/ob}$/WiscJ (ob/ob) and wild type (WT) controls to assess the potential for cocoa to

ameliorate progressive T2D and compare responses between sexes. Mice received diet without

(WT, ob/ob) or with cocoa extract (ob/ob + c) for 10 weeks. Acute cocoa reduced fasting

hyperglycemia in females, but not males, after 2 weeks. Chronic cocoa supplementation (6-10

weeks) ameliorated hyperinsulinemia in males and worsened hyperlipidemia and

hyperinsulinemia in females, yet also preserved and enhanced beta cell survival in females. The

underlying mechanisms of these differences warrant further study. If sex differences are apparent

in subsequent preclinical studies, clinical studies will be warranted to establish whether these

differences are relevant in humans. Sex differences may need to be considered when designing

human dietary interventions for T2D.

Keywords: BTBR, ob/ob, type-2 diabetes, flavanol, insulin, beta cell

1. Introduction

Diabetes is a global health crisis. The International Diabetes Federation reported in 2019 that ~463 million adults worldwide were living with diabetes [1]. It is estimated that 90% of these are type 2 (T2D) [1]. While beta cell dysfunction is traditionally viewed as a late-stage T2D event, growing data suggest it may be an early event [2–4]. Obesity is a major risk factor for T2D, as it can result in insulin resistance [5,6]. During insulin resistance, perceived insulin demand is heightened, resulting in stress on beta cells- ultimately increasing beta cell mass and disrupting the balance of proliferation and apoptosis [7]. This combination of hyperinsulinemia and dysregulated beta cell physiology leads to beta cell exhaustion and failure and ultimately hypoinsulinemia in late T2D [8].

Cocoa (*Theobroma cacao*) is one of the most concentrated dietary sources of flavanols [9–11]. Preclinical studies suggest a potential anti-diabetic role for cocoa flavanols [8]. Few studies have examined the impact of cocoa flavanols on beta cell function. Pancreatic beta cells secrete insulin and are responsible for maintenance of glucose homeostasis, along with glucagon-secreting alpha cells. Physiological conditions accompanying T2D, such as oxidative stress and inflammation, eventually induce beta cell failure and death and accelerate disease progression [5]. Promising studies suggest that flavanols protect beta cells *in vitro* [7,12–14] and *in vivo* [15,16], but this has not been investigated as thoroughly as effects in adipose tissue and skeletal muscle. We previously demonstrated that flavanols enhance beta cell function by increasing mitochondrial respiration [17]. Human studies of cocoa in the context of obesity and hyperglycemia have had mixed results [18,19]. Sex has been a largely ignored biological variable out of experimental convenience, however the potential differences due to sex have complicated translational research from male rodents to female humans. To harness the benefits

of cocoa in humans, more information from preclinical models is needed regarding the role of

disease state and sex in determining cocoa flavanol efficacy.

Rodent T2D models include insulin resistance and beta cell failure phenotypes. Monogenic

models include *ob/ob* and *db/db* mice that are deficient in leptin signaling, consuming excess

food and suppressing energy expenditure [20,21]. Environmental models include high fat feeding

and/or streptozotocin (STZ) administration. High fat feeding involves slow development of beta

cell damage, while STZ rapidly induces hyperglycemia by damaging beta cells. We employed an

aggressive T2D model with the potential to reach beta cell damage. To achieve this, the *ob/ob*

mutation on the BTBR (black and tan, brachyuric) background (BTBR.Cg-*Lep^{ob}*/WiscJ) was

used with wild type (WT) controls [22–24]. BTBR *ob/ob* mice progress rapidly to late-stage T2D

and experience a loss of specific microRNAs in pancreatic islets, potentially resulting in an

inability to increase beta cell replication in response to obesity [25].

Our objective was to perform a pilot study to identify sex differences in the response of

T2D mice to cocoa supplementation during aggressive T2D.

2. Materials and Methods

2.1 Reagents. Standards of (−)-epicatechin, (±)-catechin, and procyanidin B2 were obtained from

ChromaDex (Irvine, CA). Folin-Ciocalteu reagent and 4-dimethylaminocinnamaldehyde were

obtained from Sigma-Aldrich (St. Louis, MO).

2.2 Cocoa flavanol-rich extract production and characterization. Flavanol-rich cocoa extract

(CE) was prepared and analyzed by the Folin-Ciocalteu and 4-dimethylaminocinnamaldehyde

methods [26,27], thiolysis [26,28] and UPLC-MS/MS [29]. Methodological details are found in

Supplementary Material.

2.3 Animals. Approval was obtained from the Institutional Animal Care and Use Committee at

the David H. Murdock Research Institute (#20-011)**, and animal experiments followed** all

policies in the *Guide for the Care and Use of Laboratory Animals*. Four-week-old BTBR.Cg-

Lep^{ob}/WiscJ mice (stock # 004824) were obtained from Jackson (Bar Harbor, ME): N=12 males

and 12 females homozygous for the Lep^{ob} mutation (*ob/ob*), and N=6 male and 6 female wild-

type (WT) controls. Mice were housed under standard conditions and allowed access to food and

water *ad libitum* except where specified.

2.4 Diets. Mice were fed the 10% fat, 7% sucrose control diet from the Diet-Induced Obesity

series (D12450J, Research Diets, New Brunswick, NJ) alone or +0.8% CE (Supplementary Table

1). This dose was based on estimated 10% CE yield from cocoa powder and studies[30,31]

demonstrating that 8% cocoa powder in the diet blunts diet-induced obesity without altering food

intake (details in Supplementary Information). Based on body surface area [32], this corresponds

to 65 mg CE/kg/day in adult humans. For a 60 kg individual, this is 3900 mg CE or 39 g cocoa

powder/d, equivalent to 8 doses of cocoa powder (5 g/dose), or 2/3 of an 80 g chocolate bar with

70% cacao. Treatments: WT, *ob/ob*, *ob/ob* + cocoa extract (*ob*/ob + c) (*n*=3/sex/group).

2.5 Body weight and food intake. Body weight and food intake were measured weekly.

2.6 Blood glucose. Fasting blood glucose was measured from the tail vein at weeks 2 and 6 after

a 4 h fast using OneTouch Ultra Blue test strips (LifeScan, Inc., Milpitas, CA) and a glucometer

(upper limit was 600 mg/dL; values above that were noted and recorded as 600 mg/dL).

2.7 Euthanasia. Animals were fasted 12 h and euthanized by CO_2, followed by cervical

dislocation. Blood was collected via cardiac puncture and liver and pancreata were preserved in

10% neutral buffered formalin (Thermo) overnight and then 70% ethanol. Blood was collected in

serum tubes, clotted at room temperature (30 min) and serum prepared by centrifuging (2000 ×

g, 10 min). Samples were stored at −80°C.

2.8 Blood biomarkers. Fasting insulin and triglycerides from blood collected at euthanasia were

quantified using a mouse ultrasensitive insulin ELISA kit (Crystal Chem, Elk Grove Village, IL)

and a Triglyceride Quantification Colorimetric/Fluorometric Kit (Sigma).

2.9 Histology. Histological analysis of pancreata was conducted per established methods,

methodological details can be found in Supplementary Information.

2.10 Genotyping. Formalin fixed liver tissue was used to genotype each animal to confirm the

presence or absence of the *ob/ob* mutation. Samples were sent to Transnetyx (Cordova, TN) and

results can be found in Supplementary Information.

2.11 Statistics. Data were analyzed using GraphPad Prism v9.1.2 (La Jolla, CA). When values

were missing, a mixed-effects model was used. When initial one-way or two-way ANOVA

indicated a significant overall treatment effect or interaction (P<0.05), individual means were

compared using Tukey's or Sidak's *post-hoc* tests to control for multiple comparisons ($\alpha = 0.05$).

3. Results

3.1 Cocoa. The composition of cocoa extract is shown in Supplementary Table 2. Yield of CE

from powder was 18%.

3.2 Genotyping. We excluded one male (*ob/ob* + c) that did not carry the *ob* mutation

(Supplementary Table 3).

3.3 Body weight. No statistically significant differences were observed in weight gain (Figure 1),

likely due to significant variability in *ob/ob* mice on control diet. This could be due to weight

loss or lack of weight gain (despite *ob/ob*-induced hyperphagia) often exhibited in severe

diabetes [33]. WT mice had similar % weight gain compared to *ob/ob* controls, due to higher

initial weights of *ob/ob* mice (Supplementary Figure 1). *Ob/ob* mice ate more food than WT

controls (Supplementary Figure 1B, D). CE did not reduce food intake, and observed effects of

CE were not due to reduced caloric intake [30,31]. The average CE intake was ~1450 mg kg/d

(Supplementary Figure 2).

3.4 Fasting glucose. For males, fasting glucose was elevated in *ob/ob* and *ob/ob* + c mice

compared to WT controls at 2 and 6 weeks (Figure 2). In females, distinct effects were observed.

At week 2, *ob/ob* females had significantly elevated fasting glucose compared to WT controls,

but supplementation with CE prevented this increase. At week 6, fasting blood glucose had

worsened for both *ob/ob* and *ob/ob* + c, and the protective effect of CE was lost.

3.5 Insulin and triglycerides. Fasting blood insulin levels are shown in Figure 3A-B. In males,

fasting insulin was elevated in *ob/ob* compared to WT controls. This was reversed by CE. In

females, fasting insulin was elevated in ob/ob compared to WT controls, and CE further

increased levels. Triglyceride levels are shown in Figure 3C- D. In males, no effect of treatment

on serum triglycerides was detected. In females, serum triglycerides were not elevated in ob/ob

mice compared to WT controls (as in males), but CE resulted in significant increases.

3.6 Pancreatic beta cell area. The % insulin positive area was quantified to determine beta cell

survival (Figure 4). In males, the overall treatment effect was only borderline significant

(p=0.073), and the trend suggested that WT had slightly greater beta cell area overall compared

to *ob/ob* and *ob/ob* + c treatments. In females, *ob/ob* controls had significantly reduced beta cell

area compared to WT, but CE protected against beta cell loss.

4. Discussion

We compared effects of cocoa in male and female BTBR.Cg-*Lep[ob]*/WiscJ mice, an

aggressive hyperglycemia model. This is the first study of flavanols *ob/ob* T2D mice on the

BTBR background, and one of few studies to compare the effects of cocoa on both sexes [8,34].

Despite the small sample size, our results suggest significant differences between male and

female *ob/ob* mice when fed a diet supplemented with CE.

Sex differences in weight gain were unclear due to lack of statistical significance facilitated

by a small sample size. The speed of T2D progression was notable, as hyperglycemia and

hyperinsulinemia are accelerated with the *ob* mutation on the BTBR background compared to B6

ob/ob [6,22]. Cocoa exerted protective effects early in the intervention in females. *Ob/ob* + c

females exhibited significant protection against fasting hyperglycemia compared to *ob/ob* control

at week 2, but this was lost by week 6 (Figure 2). The data further suggest that cocoa effects

reverse based on sex as T2D progresses. Cocoa exerted a borderline significant protective effect

on fasting hyperinsulinemia in males but worsened it in females (Figure 3A-B). Cocoa protected

against beta cell loss in females (Figure 4) and together with fasting insulin expression, these

results potentially indicate that in females, CE protected beta cell health and function even

though it did not provide relief against fasting hyperglycemia. Conversely, CE reduced

hyperinsulinemia in males, but that protection was not carried across to beta cells. We previously

reported that cocoa monomers and cocoa flavanol microbial metabolites promote beta cell

stability and enhance/stimulate beta cell function *in vitro* [17,35]. These effects appear to

translate *in vivo* in female mice, but it remains unknown as to the mechanisms behind these

effects. If these effects are translatable to humans, cocoa may be useful at different stages of beta

cell function and insulin resistance based on sex.

CE did not affect fasting serum triglycerides in males but caused large increases in females,

with significantly higher levels than *ob/ob* control (Figure 3C-D). The mechanism for this, as

well as its implications, are unknown as these differences were not reflected in body weights

(Figure 1). The combination of elevated circulating insulin and serum triglycerides, particularly

in female *ob/ob* + c, can potentially be explained by more severe hyperglycemia and

hyperinsulinemia, which may have led to greater lipolysis in order to meet fuel demands. A

recent study reported worsening glucose metabolism in a mice fed purified polyphenols diet-

induced obese mouse model [36]. These data suggest that the potential anti-obesity and anti-

diabetic benefits of cocoa may differ in mice based on T2D stage, biological sex, and/or

hormonal status. Whether these sex-specific effects are seen in humans remains to be

established. Future studies should also capture metabolic responses in liver, skeletal muscle, and

adipose tissue to better understand these relationships between sex, beta cell function, and

glucose homeostatic perturbation.

This was a pilot study, with the limitation of small sample size due to mouse cost and

availability. Despite this, sex-specific effects were evident. Additionally, diet formulation was

based on WT mice, thus the actual dose of cocoa consumed was elevated due to *ob/ob*

hyperphagia. Future research is needed using more translatable doses. Given the preliminary

findings, larger studies with more statistical power are warranted.

These findings highlight the need to factor in sex-specific effects when studying cocoa and

diabetes in animals and humans [37,38]. Often, animal studies of a single strain and/or sex are

utilized to study dietary bioactives in chronic disease [39,40]. Studies should be performed to

elucidate whether sex differences are due to hormonal differences [41] and/or genetic

background. We previously reported that the anti-diabetic activity of quercetin was dependent on

genetic background in mice [37]. It is unknown whether the sex-specific effects of cocoa are

limited to *ob/ob* mice or the BTBR background. These findings need to be evaluated in multiple

preclinical mouse models of T2D. If the results are generalizable across strains, human work is

warranted. If the results are specific to narrow genetic contexts, studies can be performed to

identify the responsible genetic loci [42].

In conclusion, cocoa supplementation exerts sex-specific effects on T2D. Cocoa reduced

hyperglycemia in females, but not males, in the short term. Chronic supplementation ameliorated

hyperinsulinemia in males and worsen hyperlipidemia and hyperinsulinemia in females, yet also

preserve and enhance beta cell survival in females. The underlying mechanisms of these

differences warrant further study. These pilot data will inform experiments with larger sample

sizes. If sex differences are apparent in subsequent preclinical studies, clinical studies will be

warranted to establish whether these differences are relevant in humans. Sex differences may

need to be considered when designing human interventions for T2D.

Funding: This work was funded by the US Department of Agriculture by AFRI grant 2020-

67017-30846 (APN, JST, CDK, and MGF). The funding source had no role in in study design; in

the collection, analysis and interpretation of data; in the writing of the report; and in the decision

to submit the article for publication.

Acknowledgements. The authors acknowledge Dr. Glicerio Ignacio, DVM and Daniel Peralta

for contributions to animal care and husbandry, Lauren Essenmacher for assistance with animal

husbandry and blood glucose testing, and Lyric Ramsue for assistance with ELISAs.

References.

[1] IDF Diabetes Atlas 9th edition 2019, (n.d.). https://diabetesatlas.org/en/ (accessed March

13, 2020).

[2] E.U. Alejandro, B. Gregg, M. Blandino-Rosano, C. Cras-Méneur, E. Bernal-Mizrachi,

Natural history of β-cell adaptation and failure in type 2 diabetes, Mol. Aspects Med. 42

(2015) 19–41. https://doi.org/10.1016/j.mam.2014.12.002.

[3] M. Prentki, C.J. Nolan, Islet β cell failure in type 2 diabetes, J. Clin. Invest. 116 (2006)

1802–1812. https://doi.org/10.1172/JCI29103.

[4]   L.I. Hudish, J.E.B. Reusch, L. Sussel, β Cell dysfunction during progression of metabolic

syndrome to type 2 diabetes, J. Clin. Invest. 129 (n.d.) 4001–4008.

https://doi.org/10.1172/JCI129188.

[5]   M.E. Cerf, Beta Cell Dysfunction and Insulin Resistance, Front. Endocrinol. 4 (2013) 37.

https://doi.org/10.3389/fendo.2013.00037.

[6]   H. Lan, M.E. Rabaglia, J.P. Stoehr, S.T. Nadler, K.L. Schueler, F. Zou, B.S. Yandell, A.D.

Attie, Gene Expression Profiles of Nondiabetic and Diabetic Obese Mice Suggest a Role of

Hepatic Lipogenic Capacity in Diabetes Susceptibility, Diabetes. 52 (2003) 688–700.

https://doi.org/10.2337/diabetes.52.3.688.

[7]   L. Cedó, A. Castell-Auví, V. Pallarès, M. Blay, A. Ardévol, L. Arola, M. Pinent, Grape

seed procyanidin extract modulates proliferation and apoptosis of pancreatic beta-cells,

Food Chem. 138 (2013) 524–530.

[8]   S. Ramos, M.A. Martín, L. Goya, Effects of Cocoa Antioxidants in Type 2 Diabetes

Mellitus, Antioxidants. 6 (2017) 84. https://doi.org/10.3390/antiox6040084.

[9]   G. Schinella, S. Mosca, E. Cienfuegos-Jovellanos, M.A. Pasamar, B. Muguerza, D. Ramon,

J.L. Rios, Antioxidant properties of polyphenol-rich cocoa products industrially processed,

Food Res. Int. 43 (2010) 1614–1623.

[10] J. Oracz, D. Zyzelewicz, E. Nebesny, The Content of Polyphenolic Compounds in Cocoa

Beans (Theobroma cacao L.), Depending on Variety, Growing Region, and Processing

Operations: A Review, Crit. Rev. Food Sci. Nutr. 55 (2015) 1176–1192.

https://doi.org/10.1080/10408398.2012.686934.

[11] I. Andújar, M.C. Recio, R.M. Giner, J.L. Ríos, Cocoa polyphenols and their potential

benefits for human health, Oxid. Med. Cell. Longev. 2012 (2012).

[12] M.Á. Martín, E. Fernández-Millán, S. Ramos, L. Bravo, L. Goya, Cocoa flavonoid

epicatechin protects pancreatic beta cell viability and function against oxidative stress, Mol.

Nutr. Food Res. 58 (2014) 447–456. https://doi.org/10.1002/mnfr.201300291.

[13] M.Á. Martín, I. Cordero-Herrera, L. Bravo, S. Ramos, L. Goya, Cocoa flavanols show

beneficial effects in cultured pancreatic beta cells and liver cells to prevent the onset of type

2 diabetes, Food Res. Int. 63 (2014) 400–408.

https://doi.org/10.1016/j.foodres.2014.05.006.

[14] T. Kose, M. Vera-Aviles, P.A. Sharp, G.O. Latunde-Dada, Curcumin and (−)-

Epigallocatechin-3-Gallate Protect Murine MIN6 Pancreatic Beta-Cells against Iron

Toxicity and Erastin-Induced Ferroptosis, Pharmaceuticals. 12 (2019) 26.

https://doi.org/10.3390/ph12010026.

[15] E. Fernández-Millán, I. Cordero-Herrera, S. Ramos, F. Escrivá, C. Alvarez, L. Goya, M.A.

Martín, Cocoa-rich diet attenuates beta cell mass loss and function in young Zucker diabetic

fatty rats by preventing oxidative stress and beta cell apoptosis, Mol. Nutr. Food Res. 59

(2015) 820–824.

[16] Y. Yamashita, M. Okabe, M. Natsume, H. Ashida, Cinnamtannin A2, a Tetrameric

Procyanidin, Increases GLP-1 and Insulin Secretion in Mice, Biosci. Biotechnol. Biochem.

77 (2013) 888–891. https://doi.org/10.1271/bbb.130095.

[17] T.J. Rowley IV, B.F. Bitner, J.D. Ray, D.R. Lathen, A.T. Smithson, B.W. Dallon, C.J.

Plowman, B.T. Bikman, J.M. Hansen, M.R. Dorenkott, Monomeric cocoa catechins

enhance β-cell function by increasing mitochondrial respiration, J. Nutr. Biochem. 49

(2017) 30–41.

[18] H. Kord-Varkaneh, E. Ghaedi, A. Nazary-Vanani, H. Mohammadi, S. Shab-Bidar, Does

cocoa/dark chocolate supplementation have favorable effect on body weight, body mass

index and waist circumference? A systematic review, meta-analysis and dose-response of

randomized clinical trials, Crit. Rev. Food Sci. Nutr. 59 (2019) 2349–2362.

https://doi.org/10.1080/10408398.2018.1451820.

[19] L. Dicks, N. Kirch, D. Gronwald, K. Wernken, B.F. Zimmermann, H.-P. Helfrich, S.

Ellinger, Regular Intake of a Usual Serving Size of Flavanol-Rich Cocoa Powder Does Not

Affect Cardiometabolic Parameters in Stably Treated Patients with Type 2 Diabetes and

Hypertension—A Double-Blinded, Randomized, Placebo-Controlled Trial, Nutrients. 10

(2018) 1435. https://doi.org/10.3390/nu10101435.

[20] A.J. King, The use of animal models in diabetes research, Br. J. Pharmacol. 166 (2012)

877–894.

[21] J.P. Stoehr, J.E. Byers, S.M. Clee, H. Lan, I.V. Boronenkov, K.L. Schueler, B.S. Yandell,

A.D. Attie, Identification of Major Quantitative Trait Loci Controlling Body Weight

Variation in ob/ob Mice, Diabetes. 53 (2004) 245–249.

https://doi.org/10.2337/diabetes.53.1.245.

[22] S.M. Clee, S.T. Nadler, A.D. Attie, Genetic and genomic studies of the BTBR ob/ob mouse

model of type 2 diabetes, Am. J. Ther. 12 (2005) 491–498.

[23] T. Ranheim, C. Dumke, K.L. Schueler, G.D. Cartee, A.D. Attie, Interaction between BTBR

and C57BL/6J genomes produces an insulin resistance syndrome in (BTBR× C57BL/6J) F1

mice, Arterioscler. Thromb. Vasc. Biol. 17 (1997) 3286–3293.

[24] P.D. O'Brien, J. Hur, J.M. Hayes, C. Backus, S.A. Sakowski, E.L. Feldman, BTBR ob/ob

mice as a novel diabetic neuropathy model: Neurological characterization and gene

expression analyses, Neurobiol. Dis. 73 (2015) 348–355.

https://doi.org/10.1016/j.nbd.2014.10.015.

[25] E. Zhao, M.P. Keller, M.E. Rabaglia, A.T. Oler, D.S. Stapleton, K.L. Schueler, E.C. Neto,

J.Y. Moon, P. Wang, I.-M. Wang, P.Y. Lum, I. Ivanovska, M. Cleary, D. Greenawalt, J.

Tsang, Y.J. Choi, R. Kleinhanz, J. Shang, Y.-P. Zhou, A.D. Howard, B.B. Zhang, C.

Kendziorski, N.A. Thornberry, B.S. Yandell, E.E. Schadt, A.D. Attie, Obesity and genetics

regulate microRNAs in islets, liver, and adipose of diabetic mice, Mamm. Genome. 20

(2009) 476. https://doi.org/10.1007/s00335-009-9217-2.

[26] K.C. Racine, B.D. Wiersema, L.E. Griffin, L.A. Essenmacher, A.H. Lee, H. Hopfer, J.D.

Lambert, A.C. Stewart, A.P. Neilson, Flavanol Polymerization Is a Superior Predictor of α-

Glucosidase Inhibitory Activity Compared to Flavanol or Total Polyphenol Concentrations

in Cocoas Prepared by Variations in Controlled Fermentation and Roasting of the Same

Raw Cocoa Beans., Antioxid. Basel Switz. 8 (2019).

http://dx.doi.org.prox.lib.ncsu.edu/10.3390/antiox8120635.

[27] K.C. Racine, A.H. Lee, B.D. Wiersema, H. Huang, J.D. Lambert, A.C. Stewart, A.P.

Neilson, Development and Characterization of a Pilot-Scale Model Cocoa Fermentation

System Suitable for Studying the Impact of Fermentation on Putative Bioactive Compounds

and Bioactivity of Cocoa, Foods. 8 (2019) 102. https://doi.org/10.3390/foods8030102.

[28] M.R. Dorenkott, L.E. Griffin, K.M. Goodrich, K.A. Thompson-Witrick, G. Fundaro, L. Ye,

J.R. Stevens, M. Ali, S.F. O'Keefe, M.W. Hulver, A.P. Neilson, Oligomeric Cocoa

Procyanidins Possess Enhanced Bioactivity Compared to Monomeric and Polymeric Cocoa

Procyanidins for Preventing the Development of Obesity, Insulin Resistance, and Impaired

Glucose Tolerance during High-Fat Feeding, J. Agric. Food Chem. 62 (2014) 2216–2227.

https://doi.org/10.1021/jf500333y.

[29] K.M. Goodrich, A.P. Neilson, Simultaneous UPLC–MS/MS analysis of native catechins

and procyanidins and their microbial metabolites in intestinal contents and tissues of male

Wistar Furth inbred rats, J. Chromatogr. B. 958 (2014) 63–74.

https://doi.org/10.1016/j.jchromb.2014.03.011.

[30] Y. Gu, S. Yu, J.Y. Park, K. Harvatine, J.D. Lambert, Dietary cocoa reduces metabolic

endotoxemia and adipose tissue inflammation in high-fat fed mice, J. Nutr. Biochem. 25

(2014) 439–445. https://doi.org/10.1016/j.jnutbio.2013.12.004.

[31] Y. Gu, S. Yu, J.D. Lambert, Dietary cocoa ameliorates obesity-related inflammation in high

fat-fed mice, Eur. J. Nutr. 53 (2014) 149–158. https://doi.org/10.1007/s00394-013-0510-1.

[32] S. Reagan-Shaw, M. Nihal, N. Ahmad, Dose translation from animal to human studies

revisited, FASEB J. 22 (2008) 659–661. https://doi.org/10.1096/fj.07-9574LSF.

[33] C. Lee, H.J. Kim, T.I. Chang, E.W. Kang, Y.S. Joo, H.W. Kim, J.T. Park, T.-H. Yoo, S.-W.

Kang, S.H. Han, Synergic association of diabetes mellitus and chronic kidney disease with

muscle loss and cachexia: results of a 16-year longitudinal follow-up of a community-based

prospective cohort study, Aging. 13 (2021) 21941.

[34] L. Goya, M.Á. Martín, B. Sarriá, S. Ramos, R. Mateos, L. Bravo, Effect of Cocoa and Its

Flavonoids on Biomarkers of Inflammation: Studies of Cell Culture, Animals and Humans,

Nutrients. 8 (2016) 212. https://doi.org/10.3390/nu8040212.

[35] B.F. Bitner, J.D. Ray, K.B. Kener, J.A. Herring, J.A. Tueller, D.K. Johnson, C.M.T. Freitas,

D.W. Fausnacht, M.E. Allen, A.H. Thomson, Common gut microbial metabolites of dietary

flavonoids exert potent protective activities in β-cells and skeletal muscle cells, J. Nutr.

Biochem. 62 (2018) 95–107.

[36] H. Sanz-Lamora, P.F. Marrero, D. Haro, J. Relat, A Mixture of Pure, Isolated Polyphenols

Worsens the Insulin Resistance and Induces Kidney and Liver Fibrosis Markers in Diet-

Induced Obese Mice, Antioxidants. 11 (2022) 120. https://doi.org/10.3390/antiox11010120.

[37] L.E. Griffin, L. Essenmacher, K.C. Racine, L. Iglesias-Carres, J.S. Tessem, S.M. Smith,

A.P. Neilson, Diet-induced obesity in genetically diverse collaborative cross mouse founder

strains reveals diverse phenotype response and amelioration by quercetin treatment in

129S1/SvImJ, PWK/EiJ, CAST/PhJ, and WSB/EiJ mice, J. Nutr. Biochem. 87 (2021)

108521. https://doi.org/10.1016/j.jnutbio.2020.108521.

[38] A.Y. Sato, G.G. Pellegrini, M. Cregor, K. McAndrews, R.B. Choi, M. Maiz, O. Johnson,

L.D. McCabe, G.P. McCabe, M.G. Ferruzzi, M.A. Lila, M. Peacock, D.B. Burr, C.H.

Nakatsu, C.M. Weaver, T. Bellido, Skeletal Protection and Promotion of Microbiome

Diversity by Dietary Boosting of the Endogenous Antioxidant Response, J. Bone Miner.

Res. 36 (2021) 768–778. https://doi.org/10.1002/jbmr.4231.

[39] I. Preguiça, A. Alves, S. Nunes, R. Fernandes, P. Gomes, S.D. Viana, F. Reis, Diet-induced

rodent models of obesity-related metabolic disorders—A guide to a translational

perspective, Obes. Rev. 21 (2020) e13081. https://doi.org/10.1111/obr.13081.

[40] K. Kanasaki, D. Koya, Biology of Obesity: Lessons from Animal Models of Obesity, J.

Biomed. Biotechnol. 2011 (2011) e197636. https://doi.org/10.1155/2011/197636.

[41] J. Medina-Contreras, R. Villalobos-Molina, A. Zarain-Herzberg, J. Balderas-Villalobos,

Ovariectomized rodents as a menopausal metabolic syndrome model. A minireview, Mol.

Cell. Biochem. 475 (2020) 261–276. https://doi.org/10.1007/s11010-020-03879-4.

[42] L. Iglesias-Carres, A.P. Neilson, Utilizing preclinical models of genetic diversity to

improve translation of phytochemical activities from rodents to humans and inform

personalized nutrition, Food Funct. (2021). https://doi.org/10.1039/D1FO02782D.

Figure Legends

Figure 1. Weight gain (% of initial, mean ± SEM). Data were analyzed by 1-way ANOVA; due

to lack of observed overall treatment effects, no *post hoc* tests were performed.

Figure 2. Four hour fasting blood glucose for males and females (mean ± SEM). Data were

analyzed by 2-way ANOVA: treatment (Trt), time (T), and significance of main

effects/interactions shown. If a significant main effect or interaction was detected, Sidak (time)

or Tukey's (treatment) *post hoc* test for multiple comparisons were performed: *, **, *** and

**** indicate $P \leq 0.05$, $P \leq 0.01$, $P \leq 0.001$ and $P \leq 0.0001$, respectively. Lower-case and upper-

case superscripts indicate significance for week 2 and week 6, respectively.

Figure 3. Fasting blood insulin (A, B, mean ± SEM) and serum triglycerides (C, D, mean ± SEM).

Data were analyzed by 1-way ANOVA. If a significant treatment effect was detected, Tukey's

post hoc test was performed. Bars not sharing a common superscript are significantly different

($P<0.05$).

Figure 4. Representative pancreata images (males: A, females: C). Total % insulin positive area,

(males: B, females: D, mean ± SEM). Images were captured at 20x magnification and analyzed

using cellSens and ImageJ software for five section slides/animal. Data were analyzed by 1-way

ANOVA. If a significant treatment effect was detected, Tukey's *post hoc* test was performed. Bars

not sharing common superscripts are significantly different (P<0.05).

Figure 1

Figure 2

Figure 3

Figure 4

Male

A

	Insulin	Amylase	Dapi	Merged
WT				
ob/ob				
ob/ob + c				

B

Ins+ / pancreas area (%)

WT   *ob/ob*   *ob/ob* + c

p = 0.0734

Female

C

	Insulin	Amylase	Dapi	Merged
WT				
ob/ob				
ob/ob + c				

D

Ins+ / pancreas area (%)

a   b   a

WT   *ob/ob*   *ob/ob* + c