All-*trans* retinoic acid enhances the antitumor effects of fimaporfin-based photodynamic therapy

Judith Jing Wen Wong[1], Susanne Lorenz[2] and Pål Kristian Selbo[1],*

[1]Department of Radiation Biology, Institute for Cancer Research, The Norwegian Radium Hospital, Oslo University Hospital, Norway; judwon@rr-research.no (J.J.W.W.); selbo@rr-research.no (P.K.S.)

[2]Genomics Core Facility, Department of Core Facilities, The Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway; susanne.lorenz@rr-research.no (S.L.)

*Correspondence: Pål Kristian Selbo, selbo@rr-research.no; Tel.: +47-22781469

Abstract: The vitamin A metabolite all-*trans* retinoic acid (ATRA; tretinoin) has anticancer potential; however, lack of clinical success has prevented its approval for solid tumors. Herein, we propose combining low-dose ATRA preconditioning with fimaporfin-based photodynamic therapy (ATRA+PDT) for the treatment of solid cancers. Compared to monotherapies, ATRA+PDT induced synergistic cytotoxic responses including promotion of apoptosis in colon and breast carcinoma cell lines. In human HT-29 colon cancer cells, the effect of ATRA+PDT on gene expression was evaluated by RNA sequencing (RNA-seq). The unfolded protein response (UPR) and retinoic acid-mediated apoptosis signaling were predicted as strongly activated canonical pathways after ATRA+PDT compared to PDT. During the first 20 days post-ATRA+PDT, we obtained significant anti-tumor responses in HT-29 xenografts, including complete responses in 2/5 mice. In conclusion, ATRA+PDT represent a novel combination therapy for solid tumors that should be further tested in immunocompetent preclinical models.

Keywords: ATRA, photodynamic therapy, unfolded protein response, apoptosis, carcinoma

Abbreviations:

ALDH1A1: aldehyde dehydrogenase 1 family member A1

ALP: alkaline phosphatase

ATF3: activating transcription factor 3

ATRA: all-*trans* retinoic acid (tretinoin)

DEG: differentially expressed gene

EGR1: early growth response 1

GSEA: gene set enrichment analysis

HDAC9: histone deacetylase 9

IFN: interferon

IGFBP6: insulin-like growth factor-binding protein 6

IPA: Ingenuity pathway analysis

LCN2: lipocalin 2

LD_{50}: light-dose inducing 50% reduction of cell viability

LPS: lipopolysaccharide

PCI: photochemical internalisation

PDT: photodynamic therapy

RAR: retinoic acid receptor

REG4: regenerating islet-derived protein 4

RNA-seq: RNA sequencing

ROS: reactive oxygen species

UPR: unfolded protein response

TNF: tumor necrosis factor

TPCS2a: disulfonated tetraphenyl chlorin (fimaporfin)

TUNEL: Terminal deoxynucleotidyl transferase (TdT) dUTP nick-end labeling

Introduction

ATRA is derived from vitamin A and plays a crucial role in biological processes, including embryonic development, immune responses, and vision, through binding and activation of the nuclear retinoic acid receptors RARα, RARβ, and RARγ [1]. Transcriptional activation by RARs induce differentiation, cell cycle arrest, anti-proliferative, and apoptosis-inducing effects [2], which make retinoids attractive in cancer therapy. ATRA is successfully used in the differentiation and apoptosis induction therapy of acute promyelocytic leukemia [1, 3] and is currently being studied to prevent or treat carcinoma [4]. However, lack of effects in solid tumors has limited clinical translation of ATRA due to several factors: ATRA formulation and administration is challenging due to its low aqueous solubility. In addition, ATRA is oxidized when exposed to heat, UV light, or oxygen [4] and has a very short plasma half-life (~45 min) [5]. Moreover, RARβ expression is frequently lost in solid tumors, or the RAR promoter is epigenetically silenced, resulting in ATRA resistance [6, 7]. Thus, there is a high need for new ATRA drug delivery or combinatorial strategies to overcome resistance and improve the survival of patients with solid tumors.

PDT is used clinically for the treatment of different solid cancers and is based on three non-toxic components: photosensitizer, light, and oxygen. Upon light activation of the photosensitizer, energy is transferred to molecular oxygen, resulting in generation of cytotoxic reactive oxygen species (ROS), of which singlet oxygen is the most common [8]. The use of differentiation-promoting agents, e.g., preconditioning with vitamin D, to enhance PDT efficacy was first demonstrated by Ortel *et al.* to improve 5-aminolevulinic acid (ALA)-based PDT. However, ATRA was only shown to slightly enhance the photosensitizing compound PpIX production compared to ALA alone (non-significant) and no enhanced cytotoxic effects

was demonstrated [9]. Currently, there is a lack of mechanistic evidence showing that ATRA

preconditioning improves PDT of cancer.

In this study, we aimed to investigate the potential anti-cancer effects of ATRA+PDT using

fimaporfin (TPCS$_{2a}$), the core photosensitizer drug used for the drug delivery technology

photochemical internalization (PCI)[10]. Fimaporfin has been tested clinically in combination

with bleomycin [11] and vaccine peptide/protein-based antigens [12]. Recently, PCI in

combination with gemcitabine was shown to be safe and effective in a Phase I study for the

treatment of unresectable extrahepatic cholangiocarcinoma [13].

Here, we show for the first time that combining ATRA+PDT results in synergistic cytotoxic

responses in ATRA resistant colon and breast carcinoma cell lines compared to ATRA or

PDT monotherapy. Mechanistically, we demonstrate that ATRA+PDT results in cell cycle

arrest, increased apoptotic cell death, and a substantial reduction in cancer cell proliferation.

RNA-seq of the human colorectal adenocarcinoma cell line HT-29 identified 1129

differentially expressed genes (DEGs) after ATRA+PDT compared to PDT. Pathway analysis

of the RNA-seq data confirmed apoptosis signaling and identified activation of the unfolded

protein response (UPR). Quantitative RT-PCR and Western blotting revealed that

ATRA+PDT strongly elevated the stress-induced activating transcription factor 3 (ATF3). *In*

vivo, one single treatment with ATRA+PDT induced significant anti-tumor effects during the

first 20 days compared to no treatment in HT-29 tumor-bearing athymic nude mice, and was

the only group that achieved complete responses.

Results

ATRA+PDT enhances cytotoxic responses in breast and colon cancer cells

We evaluated the cytotoxic response of 0.01 to 100 µM ATRA monotherapy for 48 hours in

SKBR3, HCT116, and HT-29 cells. For all cell lines, the viability was > 50% at ATRA

concentrations < 50 µM (**Supplementary Fig. S1**). While the physiological concentration of

ATRA is in the nanomolar range in human plasma [14], pharmacological doses of ATRA

results in plasma concentrations in the micromolar range [15]. Thus, a therapeutic relevant

concentration of 0.1 µM ATRA was selected for the subsequent experiments and combined

with fimaporfin-PDT (ATRA+PDT). We examined the cytotoxic effects of ATRA+PDT in

five different cancer cell lines; two human colon cancer cell lines: HCT116 (ATRA-resistant)

and HT-29 (moderately ATRA-resistant) [16] and the MC-38 mouse colon cancer cell line.

Two human breast cancer cell lines were also included: SKBR3 (ATRA-sensitive) [17] and

MDA-MB-231 (ATRA-resistant) [18].

The cells were preconditioned with 0.1 µM ATRA and subjected to fimaporfin (TPCS2$_a$)-

PDT. In contrast to HCT116, enhanced cytotoxicity was observed in SKBR3 and HT-29 cells

(**Fig. 1a**) and in MC-38 and MDA-MB-231 (**Supplementary Fig. S2a**). The dual P13K and

mTOR inhibitor NVP-BEZ235 augments ATRA efficacy in leukemic cells [19] and was

therefore tested in combination with ATRA+PDT. However, no additional effect was found

when combining 0.1 µM ATRA+PDT with NVP-BEZ235 at concentrations ranging from 5-

100 nM in HT-29 and HCT116 (**data not shown**). Changing the treatment sequence where

PDT was performed prior to ATRA incubation did not have any effect on the cell viability,

indicating the importance of preconditioning (**Supplementary Fig. S2b**).

To further evaluate whether the effect of ATRA+PDT was synergistic in HT-29 and SKBR3,

the observed effect on cell viability was compared to the theoretical additive effect. A

significant difference was found indicating synergy between ATRA and PDT (**Fig. 1b**) and

124 supported with synergy calculations (**Supplementary Results**). The clonogenic assay

125 revealed that ATRA+PDT significantly induced a 4-fold and 6-fold reduction of the colony-

126 forming ability of HT-29 following 90- and 120-seconds light exposure, respectively, whereas

127 ATRA monotherapy failed to significantly affect survival (**Fig. 1c**). The impact of

128 ATRA+PDT on proliferation was also evaluated (**Fig 1d**). Strikingly, ATRA+PDT blocked

129 SKBR3 and HT-29 proliferation and attenuated proliferation of the ATRA-resistant HCT116

130 cells. ATRA monotherapy did not affect the proliferation of HT-29 and HCT116, in contrast

131 to the SKBR3 cells, where the proliferation was slightly reduced.

133 **Light exposure of ATRA alone is not cytotoxic, and *ex vivo* light exposure of ATRA and**

134 **fimaporfin does not lead to cytotoxic photoproducts**

135 ATRA is *per se* light-sensitive, and we, therefore, wanted to explore if blue light exposure of

136 ATRA would cause cytotoxic effects. However, ATRA+light did not affect cell viability

137 (**Supplementary Fig. S3a**). Furthermore, no cytotoxic photoproducts of ATRA+TPCS$_{2a}$ *ex*

138 *vivo* were observed (**Supplementary Results and Supplementary Fig. S3b**).

140 **ATRA+PDT induce cell cycle arrest and enhance apoptosis**

141 The cell cycle was analyzed using flow cytometry at 24 and 72 hours after light exposure

142 (**Fig. 2a**)**.** ATRA monotherapy did not induce any significant change in the cell cycle

143 distribution. In the PDT- or ATRA+PDT-treated HCT116 cells, a slight non-significant

144 increase in G1 was observed 24 hours post light-exposure. At 72 hours, a G2/M accumulation

145 was observed in PDT- or ATRA+PDT-treated HCT116 cells compared to the NT and ATRA

146 treatment (non-significant).

At 24 hours post-PDT or -ATRA+PDT, HT-29 cells significantly accumulated in G2/M, in

line with previous PDT-based studies [20, 21]. Specifically, 36.9-43.7% of PDT- or

ATRA+PDT-treated cells accumulated in G2/M compared to 12.6-12.9% in NT or ATRA-

treated cells ($p \leq$ 0.001-0.002). A non-significant increase in G2/M accumulation in both

PDT- and ATRA+PDT-treated HT-29 cells was also observed at 72 hours post-treatment. No

statistically significant differences were found when comparing PDT- and ATRA+PDT-

treated samples for HT-29 and HCT116. For SKBR3 cells, a slight increase in G2/M

accumulation was observed 24 hours after PDT, whereas a G2/M accumulation for

ATRA+PDT-treated cells was observed after 72-hour (non-significant).

As the anti-proliferation effect of ATRA+PDT increased with time (**Fig. 1d**), we decided to

assess the apoptotic response (TUNEL assay) at both 24 and 72 hours after a light-dose

inducing ~ 50% reduction of cell viability (LD_{50}). The apoptotic fraction 24 hours post-light

was low for both SKBR3 and HT-29. In contrast, an increased apoptotic fraction was found in

ATRA+PDT- and PDT-treated HCT116 cells; however, not significantly different (**Fig. 2a**)

At 72 hours post-ATRA+PDT, a significantly, 2-fold higher apoptotic response (~ 17%) was

detected in HT-29 ($p \leq$ 0.01) compared to PDT (~ 8.5%). A higher apoptotic response was

also detected in HCT116 and SKBR3 post-ATRA+PDT, although not significant compared to

PDT (**Fig. 2b**).

We, therefore, assessed cleaved caspase-3 in HT-29 after ATRA+PDT. In line with the

TUNEL data, no cleaved caspase-3 was detected 24 hours after light exposure (**data not**

shown). At 72 hours post-light, increased cleaved caspase-3 was observed in PDT- and

ATRA+PDT-treated cells (**Fig. 2c**). A tendency (non-significant) of higher expression of

cleaved caspase-3 was observed after ATRA+PDT compared to PDT across all three

independent experiments. Altogether this indicates that increased cell death after ATRA+PDT

is partly due to an enhanced apoptotic response.

Low-dose ATRA neither elevate intracellular ROS levels nor alkaline phosphatase

(ALP) activity and CD133 expression

ATRA induces ROS generation in leukemia [22]. The intracellular ROS was therefore

measured post-ATRA+PDT. As expected, an increase in DCF fluorescence, indicating

enhanced ROS generation, was observed in all PDT-treated samples (**Fig. 2d**). Compared to

PDT, ROS levels post ATRA+PDT showed a tendency to increase (~ 20%, non-significant)

only in the SKBR3 cells. However, low-dose ATRA (0.1 µM) did not influence the ROS

generation in all cell lines tested.

ALP is a marker of intestinal cell differentiation [23] and was used to evaluate treatment-

induced differentiation. Cells were short-term incubated with ATRA concentrations up to 1

µM for 42 hours, harvested, and assayed. The basal ALP activity was high in SKBR3 and low

in HCT116 and HT-29 (**Supplementary Fig. S4a**). ATRA did not significantly affect the

ALP activity. Intriguingly, we observed a tendency of reduced (non-significant) ALP activity

in ATRA-treated SKBR3 cells (**Supplementary Fig. S4a**).

The surface expression of the cancer stem cell marker CD133 was also evaluated using flow

cytometry. The CD133 expression was slightly reduced (non-significant) in HT-29 cells after

42 hours incubation of 0.1-1 µM ATRA (**Supplementary Fig. S4b**). Altogether, short-term

and low-dose ATRA treatment does not induce differentiation.

Differentially expressed genes (DEGs) after ATRA+PDT compared to PDT

As ATRA monotherapy did not have any effect on cell viability, we decided to perform DEGs

analysis of RNA-seq data comparing only PDT with PDT+ATRA. By this, we were able to

identify candidate genes and key pathways associated with the synergistic effect between

ATRA and PDT in HT-29 cells. Based on Prasmickaite *et al.* [24], RNA was harvested 3

hours post-light exposure using a photochemical light dose corresponding to ~ LD_{50}. We

identified 1129 genes that were significantly differentially expressed; 676 genes (59.9%)

upregulated and 453 (40.1%) downregulated (**Supplementary Table S1**). Among the

significant DEGs, 103 genes showed a log2-fold change of at least 1 (**Supplementary Table**

S2) including, 56 upregulated (54.4%) and 47 downregulated (45.6%) (**Fig. 3a**)**.** The

sequencing data are deposited at the European Nucleotide Archive (ENA,

https://www.ebi.ac.uk/ena) with the study number PRJEB49953.

Gene set enrichment analysis (GSEA)

To obtain specific, well-defined biological states or processes underlying ATRA+PDT, all

significant DEGs were first submitted to the GSEA software using the MSigDB hallmark

gene set collection. The analysis revealed ten significantly enriched gene sets (**Fig. 3b**).

Interestingly, a significant enrichment in immune-related gene sets were identified (allograft

rejection, inflammation, and IFN-α/ IFN-γ response). Moreover, apoptosis and upregulation

of KRAS signaling were positively correlated to ATRA+PDT, whereas late estrogen

response, fatty acid metabolism, and xenobiotic metabolism were negatively correlated.

All significant DEGs were further analyzed using the Ingenuity Pathway Analysis (IPA)

software for gene set enrichment analysis to gain mechanistic insight into the data on

biological processes, pathways, molecular networks, and upstream regulators. The IPA results

revealed that 68 canonical pathways were significantly enriched, of which 12 pathways had a

significant activation z-score (**Supplementary Table S3**). Five canonical pathways were

significantly activated, and seven were predicted to be significantly inhibited (**Fig. 3c**).

Strikingly, the UPR (**Fig. 3d**), IFN signaling and retinoic acid-mediated apoptosis signaling

were predicted as strongly activated canonical pathways. Although we show that ATRA alone

does not induce apoptosis, the ATRA-related genes CRABP2 and RARγ (**Supplementary**

Fig. S5a) are mapped to this pathway together with genes involved in apoptosis in IPA

(**Supplementary Table S3**). Superpathway of cholesterol biosynthesis and Huntington's

disease signaling were predicted as inhibited. Furthermore, top molecular and cellular

functions identified by IPA were associated with cellular movement, cell death and survival,

cellular development, cellular growth and proliferation, and cell-to-cell signaling interaction.

The upstream regulator analysis (URA) identified 1407 significant regulators ($p \leq 0.01$), of

which 229 were predicted as activated and 79 as inhibited (**Supplementary Table 4, Fig. 3c**).

Of relevance, strong activation of the pro-inflammatory and immune-related regulators tumor

necrosis factor (TNF), lipopolysaccharide (LPS), interferon-gamma (IFNG), poly rI:rC-RNA

(a synthetic dsRNA analogue known as poly (I:C)), and interferon-alpha (IFNA) were

predicted by the URA. As expected, the upstream regulator tretinoin (ATRA) had a

significant activation score (**Fig. 3c**).

Validation of gene expression by quantitative real-time PCR (qRT-PCR)

Quantitative RT-PCR was used to validate the RNA-seq data, including the following 6

DEGs: *REG4* (Regenerating islet-derived protein 4)*, IGFBP6* (Insulin-like growth factor-

binding protein 6)*, ATF3* (Activating transcription factor 3)*, EGR1* (Early growth response 1)*,

LCN2 (Lipocalin 2)*,* and *ALDH1A1* (Aldehyde dehydrogenase 1 family member A1). The

qRT-PCR data revealed that the selected genes' fold-change was comparable to the RNA-

sequencing results (**Fig. 4a**)

ATRA+PDT rapidly induces ATF3 and EGR1 expression

ATRA+PDT-treated samples were evaluated using qRT-PCR at 3 and 6 hours post-treatment

and compared with non-treated, PDT- and ATRA-treated cells. IGFBP6 and ALDH1A1 have

previously been described in the context of ATRA [25, 26], whereas REG4, ATF3, and EGR1

have, to the best of our knowledge, not been associated with ATRA treatment. ATF3 and

EGR1 were selected based on their association identified by the STRING network analysis

(**Fig. S5b**). Both the ATF3 and EGR1 gene expressions were upregulated synergistically after

ATRA+PDT. At 3 hours post-treatment, the EGR1 expression in the non-treated, PDT-, and

ATRA-treated samples were comparable (~ 1-fold). ATRA+PDT 3 hours post-treatment

induced a 2.5-fold increase in EGR1 expression. The ATF3 level in ATRA-treated cells was

comparable to non-treated control, whereas PDT induced a 63-fold increase 3 hours post-

light. Remarkably, a 282-fold increase of ATF3 expression was observed 3 hours post-

ATRA+PDT. This observation is in line with the literature, which indicates that ATF3 gene

expression is rapidly upregulated by various stress signals [27]. The mRNA expression of

ATF3 and EGR1 were strongly downregulated 6 hours post-PDT or -ATRA+PDT (**Fig. 4b**).

As expected, no difference in expression of LCN2, ALDH1A1, or IGFBP6 was observed

between ATRA and ATRA+PDT at 3 and 6 hours post-light. Strikingly, the REG4 expression

was entirely abolished in ATRA- or ATRA+PDT-treated cells, an observation that, to the best

of our knowledge, has not been reported before. The increased ATF3 and EGR1 expressions

were identified as unique for the ATRA+PDT combination.

The ATF3 protein expression was further validated by Western blotting of PDT- or

ATRA+PDT-treated samples 3 and 6 hours post-light exposure (**Fig. 4c**). At the 24 hour time-

point, the ATF3 expression was strongly downregulated, with slightly higher ATF3

expression in the ATRA+PDT samples compared to PDT. The basal level of ATF3 was low

in the non-treated samples and after ATRA monotherapy (**Fig 4c**).

PDT+ATRA induces initial tumor growth delay and CRs in HT-29 xenografts

The anti-tumor activities of ATRA+PDT were assessed in HT-29 xenografts in athymic nude

mice using a systemic protocol and a 652 nm laser dose of 15 J/cm^2. The ATRA+PDT

treatment had an initial strong anti-tumor effect; however, due to unexpected adverse effects

including, morbidity/mortality of the combination, the ATRA+PDT treatment group size was

too small to draw any conclusions (**Supplementary Results**). When a primary lesion or its

metastases are accessible, intratumoral (i.t.) injection of anti-cancer drugs, in particular,

immunostimulatory agents have been recommended [28]. The treatment protocol was

therefore changed to an intratumoral drug-injection strategy, and we also reduced the laser

dose to 10 J/cm^2 to achieve a more favorable adverse-effect profile.

In the second modified protocol, both ATRA and fimaporfin (TPCS$_{2a}$) were injected i.t. as a

single dose one day (18-24 hours) before light exposure. As anticipated, compared to systemic

delivery, i.t. caused only mild adverse effects including, edema and erythema in tumor areas

after PDT or ATRA+PDT (**Fig. 5a**). All ATRA+PDT-treated animals (5/5) had an initial

tumor growth delay compared to the other treatment groups (**Fig. 5b-c, Supplementary Fig.**

S7a). By comparing the mean tumor size in each treatment group at 5, 10, and 20 days post-

treatment, ATRA+PDT significantly inhibited tumor growth up to 20 days post-treatment

compared to non-treated (NT). The mean tumor volume in the ATRA+PDT group was 2-3

times smaller compared to the other treatment groups (not significant) (**Fig. 5b**). Although not

reflected in the tumor measurements, visual examination of the PDT- or ATRA+PDT-treated

tumors indicated they were flatter or had a necrotic center (**Fig. 5a**). The tumor volume

measurements are based on the assumption that tumors are spherical. As the PDT- and

ATRA+PDT-treated tumors are not spherical (but flat with a necrotic core), these groups'

tumor measurements are most likely overestimated. During laser illumination, fluorescence

was observed in tumors injected with fimaporfin ($TPCS_{2a}$) alone or in combination with

ATRA (ATRA+PDT) (**Fig. 5d**). In the PDT group, the fluorescence was limited to the tumor,

indicating that most of the photosensitizer is confined in the tumor. Intriguingly, in the

ATRA+PDT group, fluorescence in the tumor margin could also be observed.

Two out of five (40%) ATRA+PDT-treated animals achieved complete responses (CR) after

only one treatment (**Fig. 5e**). However, one of the cured animals was euthanized on day 21

due to weight loss, probably not associated with the treatment as one animal in the NT group

was also euthanized due to weight loss (**Fig. 5f**). None of the mice in any of the other

treatment groups achieved CR. The animals were followed up to at least 90 days post-

treatment. Surprisingly, in the ATRA group, 3/5 animals reached day 90 post-treatment. The

ATRA-treated tumors grew similar to the controls up to 20 days post-treatment. However,

tumor growth stagnated over time (**Supplementary Fig. S7b**), and surprisingly, the best

overall survival was obtained with ATRA monotherapy (p = 0.012 compared to NT (**Fig. 5c**).

The mean time to reach the endpoint (day 90, weight loss $\geq$ 20% and/or tumor size $\geq$ 1000

mm^3) was longest in the ATRA group (76.8 $\pm$ 8.1 days) (**Supplementary Fig. S7c**).

As the impact of the needle trauma and injection of fluids on the tumor microenvironment

was unknown, a sham procedure including an i.t. injection of PBS was included in addition to

the NT control group. No animals in the NT group reached day 90. However, 2/5 animals in

the PBS treatment group reached day 90, of which one animal was at endpoint (tumor volume

>1000 mm^3) (**Supplementary Fig. S7d**). The fact that PBS-treated animals also survived up

to day 90 indicates that i.t. injection of fluids itself affects tumor growth. One of five animals

in the PDT group (tumor volume >1000 mm^3) and in the ATRA+PDT group (CR) also

reached day 90 (**Supplementary Fig. S7d**).

328   Discussion

ATRA has not proven successful in the treatment of solid tumors due to intrinsic or acquired

drug resistance. It is therefore suggested that ATRA needs to be combined with other

therapies to improve therapeutic outcomes [4]. In this study, we present evidence

demonstrating that ATRA preconditioning combined with fimaporfin-PDT (ATRA+PDT) is a

promising anti-carcinoma strategy. Different experiments were conducted to identify the

possible mechanism behind the enhanced efficacy of ATRA+PDT, including assessment of

differentiation, ROS-generation, mTOR inhibition, cell cycle distribution, and cytotoxic

products of ATRA and fimaporfin, of which none were identified as the mechanism. Thus, in

an effort to provide a mechanistic rationale for how ATRA preconditioning is able to enhance

the cytotoxic efficacy of PDT, we decided to determine the differences of gene expression

levels between the two experimental conditions PDT and ATRA+PDT by RNA-seq. DEG

analysis of the RNA-seq data revealed a complex transcriptome profile post-ATRA+PDT.

The DEG analysis indicated that EGR1 and ATF3 were significantly upregulated in

ATRA+PDT-treated cells. In line with our observation, the stress response transcription

regulator ATF3 has previously shown to be induced after PDT [24, 29, 30], and implicated in

the UPR [31]. ATF3 was therefore selected for further analysis and validation on the mRNA

and protein level. Strikingly, the qRT-PCR results indicated a 282- and 63-fold increase in

ATF3 mRNA in the ATRA+PDT- and PDT-treated cells, respectively. Increased ATF3

protein expression after ATRA+PDT was also confirmed.

The IPA bioinformatics analysis revealed pathways and upstream regulators supporting our

observation related to enhanced cytotoxicity of ATRA+PDT, including apoptosis and the

UPR as one of the top pathways. Misfolded protein or the formation of protein aggregates due

to ROS-induced proteotoxic stress is a primary response to PDT independent of the cell type

and PDT strategy [32]. Upon endoplasmic reticulum (ER) stress, UPR is activated to reduce

accumulation and aggregation of misfolded or unfolded proteins playing an important role in

homeostasis and cell survival. However, if the ER stress is excessive and sustained, apoptosis

is induced [33]. Indeed, our observation of ATRA+PDT-enhanced apoptotic cell death by the

TUNEL assay was supported by both GSEA and IPA. Western blot analysis indicated

increased cleaved caspase-3 post-ATRA+PDT, but not significantly elevated compared to

PDT, which was confirmed by the DEG-analysis. However, compared to PDT monotherapy,

DEG-analysis revealed significant upregulation of caspase-4, -7, and -9 post-ATRA+PDT.

Furthermore, TNF-α, a potent pro-apoptotic trigger that may also induce necrosis [34], was

identified as an activated upstream regulator after ATRA+PDT [34].

Interestingly, the downstream effects of the p38 kinase inhibitor SB203580 were predicted to

be significantly inhibited suggesting that p38 kinase is involved in stress-induced death

signalling after ATRA+PDT. This is consistent with previous observations; Weyergang *et al.*

demonstrated that SB203580 in combination with $TPPS_{2a}$-based PDT resulted in increased

cell viability [35], while similar effects were shown by Olsen *et al.* where SB203580 was

combined with fimaporfin-PDT [21].

Of notice, HDAC was identified as one of the upstream regulators with a negative z-score

indicating HDAC inhibition (HDACi) despite that the class II deacetylase HDAC9 was 2.2-

fold upregulated by ATRA+PDT. HDAC9 negatively regulates the function of TRIM29 by

non-histone deacetylation of the protein [36], and thereby alters its ability to associate with

p53 and consequently inhibits its cell proliferation-promoting activity. Interestingly, our

RNA-seq data reveal that TRIM29 is 30% downregulated post-ATRA+PDT (p = 5.00E-05),

which may strengthen the concept of the HDACi effect of ATRA+PDT.

The ATRA+PDT combination strategy was further evaluated *in vivo* by using the HT-29

xenograft model in athymic nude mice. High morbidity (significant weight loss and

xeroderma) was observed when ATRA was combined with $TPCS_{2a}$-PDT systemically. In an

effort to enhance the safety profile and anti-tumor efficacy, ATRA and $TPCS_{2a}$ were co-

administered i.t. This technique was recently suggested by Marabelle *et al.* as a strategy to

deliver immunostimulatory agents [28]. This strategy also offers high drug concentration in

the tumor and has previously been explored by us using ribosome-inactivating protein toxins,

in combination with an amphiphilic photosensitizer that was delivered systemically [37, 38].

Remarkably, 40% (2/5) of the ATRA+PDT-treated animals achieved CR after only one

treatment, whereas no CRs were obtained in the other treatment groups. Moreover, all

ATRA+PDT-treated animals had significantly smaller tumors than NT during the first 20

days post-treatment. In the ATRA+PDT group, fimaporfin fluorescence was observed in the

tumors as well as in the tumor margins during light exposure, as opposed to the PDT

treatment group, where the fluorescence was confined to the tumors only. Although the exact

mechanism of this ATRA effect is not clear, the enhanced $TPCS_{2a}$ distribution in the tumor

margin may contribute to the improved overall PDT response and explain the initial tumor

growth inhibition in the ATRA+PDT group. In addition to direct cell killing, PDT with PCI-

based photosensitizers also targets the tumor vasculature [39-41]. Thus, light activation of

fimaporfin in the tumor periphery of the ATRA+PDT-treated animals may induce vascular

shutdown, which limits the oxygen and nutrition supply to the tumor, but this remains to be

tested experimentally. Despite the CRs obtained after ATRA+PDT, ATRA monotherapy

surprisingly gave the best overall survival, of which 60% of the mice were still alive at day

90.

In conclusion, our study shows that ATRA preconditioning is a promising strategy to improve

the efficacy of PDT in carcinomas and identify enhanced activation of apoptosis as a death

mechanism. DEG analysis predicted the unfolded protein response as the most significantly

upregulated pathway. To fully exploit the potential of this preconditioning approach,

preclinical studies of ATRA in combination with laser-activation of the clinical relevant

photosensitizer fimaporfin (PDT or PCI) in immunocompetent murine models are warranted.

Materials and Methods

Cell lines

All the cell lines were purchased from American Type Culture Collections (ATCC, Manassas,

VA, USA), except the MC-38 cell line, which was from Kerafast (Boston, MA, USA). A

detailed description of the cell lines and culture conditions are detailed in **Supplementary**

Materials and Methods.

Drugs and chemicals

ATRA (Sigma-Aldrich, Saint-Louis, MO, USA) was dissolved in DMSO was stored at -80

°C. The photosensitizer $TPCS_{2a}$ (in polysorbate 80, 2.8% mannitol, 50 mM Tris, pH 8.5) was

provided by PCI Biotech AS (Oslo, Norway) and stored at 4 °C. All work with ATRA or

TPCS$_{2a}$ was performed under subdued light.

***In vitro* ATRA+PDT treatment protocol**

To evaluate the treatment effects of ATRA+PDT, 3000 (HCT116 and HT-29) or 8000

(SKBR3) cells/well were seeded in 96-well plates (Nunc). The cells were first incubated with

0.1 µM ATRA for 24 hours and further incubated with 0.1 µM ATRA and/or 0.4 µg/mL

TPCS$_{2a}$ for 18 hours. Before light exposure, the cells were washed twice with PBS, chased for

4 hours in fresh media, and illuminated with LumiSource (PCI Biotech, Oslo, Norway) [41].

Cell viability was evaluated 48 hours post-treatment. The treatment protocol for MC-38 and

MDA-MB-231 is described in **Supplementary Materials and Methods**.

Evaluation of treatment-induced cytotoxicity and effect on proliferation

A description of the MTT assay, clonogenic assay, and the real-time monitoring of cell

proliferation can be found in **Supplementary Materials and Methods.**

Apoptosis and cell cycle analysis

The TUNEL assay and cell cycle analyses were performed as previously described [21] and

detailed in **Supplementary Materials and Methods**.

Detection of intracellular ROS

The cell permeant reagent 2',7'-dichlorofluorescin diacetate (DCF-DA) and its downstream

product 2',7'-dichlorofluorescein (DCF) was used to quantitatively assess ROS by flow

cytometry in live cell samples as previously described [21].

Flow cytometry

Flow cytometry was performed using an LSRII flow cytometer (Becton Dickinson, Franklin

Lakes, NJ, USA). Data were processed using the FlowJo software version 10 (Treestar, OR,

USA).

RNA extraction and qRT-PCR

Total cellular RNA was extracted from HT-29 cells at indicated time-points after treatments

prior to qRT-PCR as detailed in **Supplementary Materials and Methods**.

Western blot analysis

Western blot was performed as previously described [42] and detailed in **Supplementary**

Materials and Methods.

RNA-seq and differential gene analysis

Total RNA was isolated from three biological replicates of HT-29 cells treated with PDT or

ATRA+PDT. The samples were harvested 3 hours post-120 seconds blue light exposure,

which corresponds to ~LD_{50} measured by MTT. The RNA integrity number (RIN) was

evaluated using the 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA), where

all the samples had a RIN=10. Total RNA (500 ng) was processed using a TruSeq Stranded

mRNA kit (Illumina, San Diego, CA, USA) following the manufacturer's recommended

procedures. Final libraries were quality controlled using qRT-PCR and pooled. Pooled

libraries were sequenced on the NextSeq500 instrument (Illumina) with a HighOutput flow

cell, single read sequencing of 75 bp read length, and dual-indexes (IDT UDI's 2x8 bp). For

data analysis, FASTQ files were mapped using Star2 aligner (STAR v2.5.0b) and human

reference hg19. Counting, normalization, and differential gene expression analysis were

performed using Cufflinks2 (v2.2.1).

Gene set enrichment analyses

Broad Institute's GSEA software 4.1.0 was used to determine whether an *a priori* defined set

of genes is represented as DEGs in ATRA+PDT compared to PDT [43, 44]. The Hallmark

Gene sets were obtained from the Molecular Signature Database (MSigDb v7.4) [45]. False

discovery rate (FDR) of ≤ 25% and p-value ≤ 0.05 was considered significant. Further

analysis was performed using the IPA software (Qiagen, Hilden, Germany), which identify

canonical pathways and biological functions or diseases significantly associated with the

identified DEGs. The IPA software was used to analyze upstream regulators that are predicted

to be relevant for the biological mechanisms underlying the data based on information

gathered in the literature [46]. An overlap of p-value ≤ 0.01 and an activation z-score ≥ |2| was

considered significant. The z-score is used to deduce likely activation states based on

significant pattern matches of up- or downregulated genes. A positive z-score predicts

activation, and a negative z-score indicates inhibition.

Animal studies

All animal procedures were performed according to protocols approved by the Norwegian

Food Safety Authority, which is the national animal research authority, and were conducted

according to the Federation of European Laboratory Animal Science Association (FELASA)

regulations. A detailed description can be found in **Supplementary Materials and Methods.**

Statistical Analysis

Statistical analyses were performed using SigmaPlot 14.0 (Systat Software, Inc., San Jose,

CA, USA). Statistical significance was set at $p \leq 0.05$ unless stated otherwise. Synergy was

evaluated by comparing the difference in log (DL) of the calculated additive effect with the

observed effect, as previously described [47]. Survival analysis was performed by pairwise

log-rank (Mantel-Cox) comparison using SPSS Statistics 26 (IBM, Armonk, NY, USA).

Author Contributions: J.J.W.W. planned, performed, analyzed almost all experiments.

J.J.W.W. prepared the samples for RNA-seq. S.L. was involved in planning the RNA-seq

experiments, performed, and analyzed the RNA-seq data, and contributed to the gene set

enrichment analysis interpretation. J.J.W.W. and P.K.S. planned and performed the animal

studies. J.J.W.W. and P.K.S. wrote the manuscript with input from S.L. P.K.S. conceived the

original idea, acquired funding, provided resources, and supervised the project.

Funding: This research was funded by the South-Eastern Norway Regional Health Authority

(Helse-Sør Øst) (grant number: 2017068 (P.K.S.) and 2016023 (J.J.W.W.)) and The

Norwegian Radium Hospital Research Foundation (funding number: FU0803 (P.K.S.)).

Acknowledgments: We thank Drs. Trond Stokke and Idun Dale Rein at the Flow Cytometry

Core facility for advice on cell cycle analysis. We also thank Ane Sager Longva for excellent

technical assistance and advice on RNA extraction, qRT-PCR and Western blotting. We also

acknowledge Miriam Rosvold From for great assistance on the MDA-MB-231 cell

experiments and Ane Sofie Fremstedal for processing the tumor measurement data.

Conflicts of Interest: P.K.S. and J.J.W.W. has filed a patent (US 63/282,267 (2021)) on the

ATRA+PDT method. P.K.S. is a co-inventor on several patents and patent applications

related to the PCI technology. S.L. declares no conflict of interest.

References

1.   Tang XH, Gudas LJ: Retinoids, retinoic acid receptors, and cancer. *Annu Rev Pathol*
2011, 6:345-364.

2.   Altucci L, Leibowitz MD, Ogilvie KM, de Lera AR, Gronemeyer H: RAR and RXR
modulation in cancer and metabolic disease. *Nat Rev Drug Discov* 2007, 6(10):793-
810.

3.   Thomas X: Acute Promyelocytic Leukemia: A History over 60 Years-From the Most
Malignant to the most Curable Form of Acute Leukemia. *Oncol Ther* 2019, 7(1):33-
65.

4.   Ferreira R, Napoli J, Enver T, Bernardino L, Ferreira L: Advances and challenges in
retinoid delivery systems in regenerative and therapeutic medicine. *Nat Commun*
2020, 11(1):4265.

5.   Muindi JR, Frankel SR, Huselton C, DeGrazia F, Garland WA, Young CW, Warrell
RP, Jr.: Clinical pharmacology of oral all-trans retinoic acid in patients with acute
promyelocytic leukemia. *Cancer Res* 1992, 52(8):2138-2142.

6.   Connolly RM, Nguyen NK, Sukumar S: Molecular pathways: current role and future
directions of the retinoic acid pathway in cancer prevention and treatment. *Clin*
Cancer Res 2013, 19(7):1651-1659.

7.   Duong V, Rochette-Egly C: The molecular physiology of nuclear retinoic acid
receptors. From health to disease. *Biochim Biophys Acta* 2011, 1812(8):1023-1031.

8.   Agostinis P, Berg K, Cengel KA, Foster TH, Girotti AW, Gollnick SO, Hahn SM,
Hamblin MR, Juzeniene A, Kessel D *et al*: Photodynamic therapy of cancer: an
update. *CA Cancer J Clin* 2011, 61(4):250-281.

9.　Ortel B, Sharlin D, O'Donnell D, Sinha AK, Maytin EV, Hasan T: Differentiation

enhances aminolevulinic acid-dependent photodynamic treatment of LNCaP prostate

cancer cells. *Br J Cancer* 2002, 87(11):1321-1327.

10.　Selbo PK, Weyergang A, Hogset A, Norum OJ, Berstad MB, Vikdal M, Berg K:

Photochemical internalization provides time- and space-controlled endolysosomal

escape of therapeutic molecules. *J Control Release* 2010, 148(1):2-12.

11.　Sultan AA, Jerjes W, Berg K, Hogset A, Mosse CA, Hamoudi R, Hamdoon Z, Simeon

C, Carnell D, Forster M *et al*: Disulfonated tetraphenyl chlorin (TPCS2a)-induced

photochemical internalisation of bleomycin in patients with solid malignancies: a

phase 1, dose-escalation, first-in-man trial. *Lancet Oncol* 2016, 17(9):1217-1229.

12.　Otterhaug T, Janetzki S, Welters MJP, Hakerud M, Nedberg AG, Edwards VT,

Boekestijn S, Loof NM, Selbo PK, Olivecrona H *et al*: Photochemical Internalization

Enhanced Vaccination Is Safe, and Gives Promising Cellular Immune Responses to an

HPV Peptide-Based Vaccine in a Phase I Clinical Study in Healthy Volunteers. *Front

Immunol* 2020, 11:576756.

13.　Trojan J, Hoffmeister A, Neu B, Kasper S, Dechêne A, Jürgensen C, Schirra J, Jakobs

R, Palmer D, Selbo P *et al*: Photochemical Internalization of Gemcitabine Is safe and

Effective in Locally Advanced Inoperable Cholangiocarcinoma. *The Oncologist* 2022,

In press.

14.　Jing J, Isoherranen N, Robinson-Cohen C, Petrie I, Kestenbaum BR, Yeung CK:

Chronic Kidney Disease Alters Vitamin A Homeostasis via Effects on Hepatic RBP4

Protein Expression and Metabolic Enzymes. *Clin Transl Sci* 2016, 9(4):207-215.

15.　Adamson PC, Bailey J, Pluda J, Poplack DG, Bauza S, Murphy RF, Yarchoan R, Balis

FM: Pharmacokinetics of all-trans-retinoic acid administered on an intermittent

schedule. *J Clin Oncol* 1995, 13(5):1238-1241.

16.　Lee MO, Han SY, Jiang S, Park JH, Kim SJ: Differential effects of retinoic acid on

growth and apoptosis in human colon cancer cell lines associated with the induction of

retinoic acid receptor beta. *Biochem Pharmacol* 2000, 59(5):485-496.

17.   Paroni G, Fratelli M, Gardini G, Bassano C, Flora M, Zanetti A, Guarnaccia V,

574   Ubezio P, Centritto F, Terao M *et al*: Synergistic antitumor activity of lapatinib and

575   retinoids on a novel subtype of breast cancer with coamplification of ERBB2 and

576   RARA. *Oncogene* 2012, 31(29):3431-3443.

577   18.   Liu RZ, Graham K, Glubrecht DD, Germain DR, Mackey JR, Godbout R: Association

578   of FABP5 expression with poor survival in triple-negative breast cancer: implication

579   for retinoic acid therapy. *Am J Pathol* 2011, 178(3):997-1008.

580   19.   Ronan T. Swords P, Tino Schenk, Sven Stengel, Veronica Sylvina Gil, Kevin R.

581   Petrie, Aymee Perez, Rodriguez Ana, Justin M. Watts, Fernando Vargas, Roy Elias

582   and Arthut Zelent: Inhibition of the PI3K/AKT/mTOR Pathway Leads to Down-

583   Regulation of c-Myc and Overcomes Resistance to ATRA in Acute Myeloid

584   Leukemia. *Blood* 2015, 126(23):1363.

585   20.   Bostad M, Olsen CE, Peng Q, Berg K, Hogset A, Selbo PK: Light-controlled

586   endosomal escape of the novel CD133-targeting immunotoxin AC133-saporin by

587   photochemical internalization - A minimally invasive cancer stem cell-targeting

588   strategy. *J Control Release* 2015, 206:37-48.

589   21.   Olsen CE, Weyergang A, Edwards VT, Berg K, Brech A, Weisheit S, Hogset A, Selbo

590   PK: Development of resistance to photodynamic therapy (PDT) in human breast

591   cancer cells is photosensitizer-dependent: Possible mechanisms and approaches for

592   overcoming PDT-resistance. *Biochem Pharmacol* 2017, 144:63-77.

593   22.   Mathieu J, Giraudier S, Lanotte M, Besancon F: Retinoid-induced activation of NF-

594   kappaB in APL cells is not essential for granulocytic differentiation, but prolongs the

595   life span of mature cells. *Oncogene* 2005, 24(48):7145-7155.

596   23.   Shin J, Carr A, Corner GA, Togel L, Davalos-Salas M, Tran H, Chueh AC, Al-Obaidi

597   S, Chionh F, Ahmed N *et al*: The intestinal epithelial cell differentiation marker

598   intestinal alkaline phosphatase (ALPi) is selectively induced by histone deacetylase

599   inhibitors (HDACi) in colon cancer cells in a Kruppel-like factor 5 (KLF5)-dependent

600   manner. *J Biol Chem* 2014, 289(36):25306-25316.

601   24.   Prasmickaite L, Cekaite L, Hellum M, Hovig E, Høgset A, Berg K: Transcriptome
602   changes in a colon adenocarcinoma cell line in response to photochemical treatment as
603   used in photochemical internalisation (PCI). *FEBS Letters* 2006, 580(24):5739-5746.

604   25.   Kim EJ, Kang YH, Schaffer BS, Bach LA, MacDonald RG, Park JH: Inhibition of
605   Caco-2 cell proliferation by all-trans retinoic acid: role of insulin-like growth factor
606   binding protein-6. *J Cell Physiol* 2002, 190(1):92-100.

607   26.   Bhatlekar S, Viswanathan V, Fields JZ, Boman BM: Overexpression of HOXA4 and
608   HOXA9 genes promotes self-renewal and contributes to colon cancer stem cell
609   overpopulation. *J Cell Physiol* 2018, 233(2):727-735.

610   27.   Hai T, Wolford CC, Chang YS: ATF3, a hub of the cellular adaptive-response
611   network, in the pathogenesis of diseases: is modulation of inflammation a unifying
612   component? *Gene Expr* 2010, 15(1):1-11.

613   28.   Marabelle A, Andtbacka R, Harrington K, Melero I, Leidner R, de Baere T, Robert C,
614   Ascierto PA, Baurain JF, Imperiale M *et al*: Starting the fight in the tumor: expert
615   recommendations for the development of human intratumoral immunotherapy (HIT-
616   IT). *Ann Oncol* 2018, 29(11):2163-2174.

617   29.   Cekaite L, Peng Q, Reiner A, Shahzidi S, Tveito S, Furre IE, Hovig E: Mapping of
618   oxidative stress responses of human tumor cells following photodynamic therapy
619   using hexaminolevulinate. *BMC Genomics* 2007, 8:273.

620   30.   Weijer R, Broekgaarden M, van Golen RF, Bulle E, Nieuwenhuis E, Jongejan A,
621   Moerland PD, van Kampen AH, van Gulik TM, Heger M: Low-power photodynamic
622   therapy induces survival signaling in perihilar cholangiocarcinoma cells. *BMC Cancer*
623   2015, 15:1014.

624   31.   Jiang HY, Wek SA, McGrath BC, Lu D, Hai T, Harding HP, Wang X, Ron D,
625   Cavener DR, Wek RC: Activating transcription factor 3 is integral to the eukaryotic
626   initiation factor 2 kinase stress response. *Mol Cell Biol* 2004, 24(3):1365-1377.

627   32.   Broekgaarden M, Weijer R, van Gulik TM, Hamblin MR, Heger M: Tumor cell
628   survival pathways activated by photodynamic therapy: a molecular basis for
629   pharmacological inhibition strategies. *Cancer Metastasis Rev* 2015, 34(4):643-690.

33. Hetz C: The unfolded protein response: controlling cell fate decisions under ER stress

and beyond. *Nat Rev Mol Cell Biol* 2012, 13(2):89-102.

34. Chu WM: Tumor necrosis factor. *Cancer Lett* 2013, 328(2):222-225.

35. Weyergang A, Kaalhus O, Berg K: Photodynamic therapy with an endocytically

located photosensitizer cause a rapid activation of the mitogen-activated protein

kinases extracellular signal-regulated kinase, p38, and c-Jun NH2 terminal kinase with

opposing effects on cell survival. *Mol Cancer Ther* 2008, 7(6):1740-1750.

36. Yuan Z, Peng L, Radhakrishnan R, Seto E: Histone deacetylase 9 (HDAC9) regulates

the functions of the ATDC (TRIM29) protein. *J Biol Chem* 2010, 285(50):39329-

39338.

37. Berstad MB, Cheung LH, Berg K, Peng Q, Fremstedal AS, Patzke S, Rosenblum MG,

Weyergang A: Design of an EGFR-targeting toxin for photochemical delivery: in vitro

and in vivo selectivity and efficacy. *Oncogene* 2015, 34(44):5582-5592.

38. Selbo PK, Sivam G, Fodstad O, Sandvig K, Berg K: In vivo documentation of

photochemical internalization, a novel approach to site specific cancer therapy. *Int J*

Cancer 2001, 92(5):761-766.

39. Weyergang A, Cheung LH, Rosenblum MG, Mohamedali KA, Peng Q, Waltenberger

J, Berg K: Photochemical internalization augments tumor vascular cytotoxicity and

specificity of VEGF(121)/rGel fusion toxin. *J Control Release* 2014, 180:1-9.

40. Vikdal M, Weyergang A, Selbo PK, Berg K: Vascular endothelial cells as targets for

photochemical internalization (PCI). *Photochem Photobiol* 2013, 89(5):1185-1192.

41. Wong JJW, Berstad MB, Fremstedal ASV, Berg K, Patzke S, Sorensen V, Peng Q,

Selbo PK, Weyergang A: Photochemically-Induced Release of Lysosomal

Sequestered Sunitinib: Obstacles for Therapeutic Efficacy. *Cancers (Basel)* 2020,

12(2).

42. Berstad MEB, Cheung LH, Weyergang A: Production of Recombinant Gelonin Using

an Automated Liquid Chromatography System. *Toxins (Basel)* 2020, 12(8).

43. Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J,

Puigserver P, Carlsson E, Ridderstrale M, Laurila E *et al*: PGC-1alpha-responsive

659    genes involved in oxidative phosphorylation are coordinately downregulated in human

660    diabetes. *Nat Genet* 2003, 34(3):267-273.

661 44. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,

662    Paulovich A, Pomeroy SL, Golub TR, Lander ES *et al*: Gene set enrichment analysis:

663    a knowledge-based approach for interpreting genome-wide expression profiles. *Proc*

664    *Natl Acad Sci U S A* 2005, 102(43):15545-15550.

665 45. Liberzon A, Birger C, Thorvaldsdottir H, Ghandi M, Mesirov JP, Tamayo P: The

666    Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst*

667    2015, 1(6):417-425.

668 46. Kramer A, Green J, Pollard J, Jr., Tugendreich S: Causal analysis approaches in

669    Ingenuity Pathway Analysis. *Bioinformatics* 2014, 30(4):523-530.

670 47. Weyergang A, Kaalhus O, Berg K: Photodynamic targeting of EGFR does not predict

671    the treatment outcome in combination with the EGFR tyrosine kinase inhibitor

672    Tyrphostin AG1478. *Photochemical & Photobiological Sciences* 2008, 7(9):1032-

673    1040.

 Figure legends

Fig. 1. ATRA preconditioning enhances the cytotoxic efficacy of fimaporfin-PDT and

strongly impair proliferation in carcinoma cells.

a Viability assessed by MTT 48 hours post-light exposure in SKBR3, HT-29, and HCT116

cells treated with fimaporfin-PDT or the combination ATRA+PDT. Representative data of at

least three independent experiments. Data presented as mean of triplicates ± S.D. 60 seconds

light exposure ≈ 0.58 J/cm^2. **b** Synergistic cytotoxic effects induced by ATRA+PDT in HT-29

and SKBR3 cells. The observed effect of the combination in SKBR3 and HT-29 compared to

a theoretical additive effect at ~ LD$_{60}$ of at least three independent experiments. Data

presented as mean ± S.E. **c** Representative clonogenic assay of PDT compared to

ATRA+PDT in HT-29 cells. **d** Proliferation of SKBR3, HT-29 and HCT116 up to 110 hours

post-120 seconds light exposure (1.16 J/cm^2). Representative experiment of three

experiments, data presented as mean of triplicates ± S.E. Statistical significance calculated

using Student's two-tailed t-test (*** p ≤ 0.001, ** p ≤ 0.01, * p ≤ 0.05).

Fig. 2. ATRA preconditioning increases PDT-induced cell cycle arrest and apoptosis

The cells were subjected to 120 seconds (SKBR3) and 90 seconds (HT-29 and HCT116) light

exposure, which corresponds to ≈ 1.16 J/cm^2 and 0.87 J/cm^2, respectively. 4 Gy-treated Reh

cells harvested 24 hours post-radiotherapy was included as a positive control. **a** Cell cycle

analysis at 24- and 72-hours post-light exposure. Left panels, percent distribution in G2/M, S,

and G1 phase after the different treatments. Data presented as mean of three independent

experiments ± S.E. Right panels, representative cell cycle histograms. **b** Apoptosis was

evaluated by the TUNEL assay at 24- and 72-hours post-treatment. Data presented as mean of

three experiments ± S.E. **c** Western blot of cleaved caspase-3 in HT-29 cells harvested 72

hours post-treatment. Cells treated with 10 Gy were included as a positive control, and γ-

tubulin was included as a loading control. Representative blot of three experiments. **d**

Intracellular ROS was measured using the DCFH-DA assay immediately after light exposure.

Data are normalized to non-treated samples and presented as mean of three experiments $\pm$

S.E. Statistical significance calculated with one-way ANOVA. (*** $p \leq 0.001$, ** $p \leq 0.01$

and * $p \leq 0.05$, n.s.: not significant).

Fig. 3. Differentially expressed genes (DEGs) and gene set enrichment analyses (GSEA)

in ATRA+PDT-treated HT-29 cells compared to PDT-treatment.

a Heat map of DEGs of fold change $\geq |2|$ at 3 hours post-light exposure in PDT and

ATRA+PDT-treated samples. **b** All significantly enriched biological states or processes in

ATRA+PDT compared to PDT determined by GSEA analysis using the Hallmark gene set. A

p-value of ≤ 0.05 and an FDR of $\leq 25\%$ were considered significant. **c** Visualization of IPA

analysis of top five canonical pathways based on p-value, top five molecular and cellular

functions, and top ten activated and inhibited based on z-value. A p-value of ≤ 0.01 and a

predicted activation, z-score of $\geq |2|$ were considered significant. **d** Heat map of DEGs

mapped to the unfolded protein response.

Fig. 4. Post-RNA-seq validation of six selected DEGs by qRT-PCR and ATF3 protein

expression by Western blotting.

a Validation of mRNA expression of six selected DEGs (ALDH1A1, ATF3, EGR1, IGFBP6,

LCN2 and REG4) by RT-qPCR of samples harvested 3 hours post-treatment. Quantitative

RT-PCR data are presented as log-2-fold change (FC) (ratio of ATRA+PDT/PDT) and

compared to the RNA-seq data. **b** Relative normalized expression of DEGs in samples

obtained 3 or 6 hours post-ATRA+PDT treatment and compared to non-treated (NT), ATRA-

and PDT-treated controls. The data are normalized to two reference genes (RPLP0 and

GADPH) and presented as mean of three experiments ± S.E. Statistical significance

calculated with one-way ANOVA. **c** Relative ATF3 signal in ATRA+PDT to PDT samples.

The data were normalized to loading control. Data are presented as mean of four experiments

± S.E. One-sample t-test. Representative western blot of ATF3 at 3-, 6-, and 24-hours post-

treatment. γ-tubulin was included as a loading control. (*** $p \leq 0.001$, ** $p \leq 0.01$ and * $p \leq$

0.05, n.s.: not significant).

Fig. 5. Treatment response in HT-29 xenograft-tumor bearing mice following i.t.

delivery of ATRA and fimaporfin, and photochemical treatment.

a Representative images of one animal in each treatment group at day 0 (before light

exposure), 1 and 7 post-treatment. **b** Tumor volume (mm^3) of each treatment group at day 5,

10, and 20 post-treatment. Data presented as mean ± S.E. Statistical significance calculated

with one-way ANOVA. (*** $p \leq 0.001$, ** $p \leq 0.01$ and * $p \leq 0.05$, n.s.: not significant). **c**

Tumor growth of individual animals in each treatment group up to 20 days post-treatment.

Each line represents one animal. **d** Images of fimaporfin fluorescence in the tumors during

light exposure in the PDT- and ATRA+PDT-group. The yellow frames indicate animals that

achieved complete responses. **e** Images of the two mice that were cured in the ATRA+PDT

group before light exposure (day 0) and post-treatment at a later time-point (day 18 and 102).

f Relative weight of each individual animal, relative to day -1, in each treatment group. Each

line represents one animal. For all treatment groups, n = 5, except for NT where n = 6.