Recent exposure to environmental stochasticity does not determine the resilience of natural populations

James Cant[1,2] [*], Pol Capdevila[3,4], Maria Beger[2,5], and Roberto Salguero-Gómez[4, 5, 6]

[1] Centre for Biological Diversity, University of St Andrews, St Andrews, KY16 9TH, United Kingdom.

[2] School of Biology, Faculty of Biological Sciences, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, United Kingdom.

[3] School of Biological Sciences, University of Bristol, 24 Tyndall Ave, BS8 1TQ, Bristol, UK

[4] Department of Zoology, University of Oxford, 11a Mansfield Road, Oxford, OX1 3SZ, United Kingdom.

[5] Centre for Biodiversity and Conservation Science, School of Biological Sciences, University of Queensland, Brisbane, QLD 4072, Australia.

[6] Max Planck Institute for Demographic Research, Konrad Zuße Straße 1, 18057 Rostock, Germany.

* Corresponding author:

       James Cant : jic2@st-andrews.ac.uk

A manuscript for publication as a Research Article in PNAS

Word Count:

Abstract: 197; *Main text (including Methods)*: 4,521; *Figures & Tables*: 6, *References*: 97

Significance statement (119/120)

Populations that currently endure more variable abiotic conditions are often expected to be less vulnerable to future increases in climatic variability. However, without defining the link between abiotic variability and the capacity for populations to resist and recover following disturbances (*i.e.*, their resilience), we cannot predict the consequences of ongoing community reassembly. Evaluating the association between measures of abiotic variability

and the resilience attributes of 2,242 animal, plant, and algae populations, we discredit the

assumption that contemporary exposure to more frequent environmental shifts confers a

greater ability to cope with future global change. Instead, the resilience attributes of natural

populations appear to have been moulded over longer-term evolutionary timeframes and are

thus not a response to more recent experiences.

Abstract

Escalating climatic and anthropogenic pressures expose ecosystems worldwide to

increasingly frequent disturbances. Yet, our ability to forecast the responses of natural

populations to these disturbances is impeded by a limited understanding for how exposure to

stochastic environments shapes population resilience. Instead, the resilience, and

vulnerability, of natural populations to ongoing global change is often presumed based on

their contemporary exposure to environmental stochasticity. To test the validity of this

assumption, we investigated the association between the resilience attributes (e.g., resistance

and recovery) of natural animal and plant populations, and measures of local environmental

stochasticity (e.g., spectral frequency and abiotic range); collating data from 2,242

populations across 369 animal, plant, and algal species. Unexpectedly, recent abiotic

stochasticity regimes from the past 50 years do not predict the inherent ability of populations

to resist or recover from disturbances. Instead, population resilience is strongly affected by

phylogenetic relationships among species, with survival and developmental investments

shaping their responses to stochastic regimes. Contrary to the classical assumption that

exposure to recent environmental shifts confers a greater ability to cope with current and

future global change, our findings suggest that population resilience is a consequence of

evolutionary processes and/or deep-time environmental regimes.

Introduction

Ongoing global shifts in the timing and magnitude of environmental variation warrant an understanding for the processes underlying the capacity for populations to resist and recover from disturbances (1, 2) (*i.e.*, their resilience [3]). However, half a century since Holling first defined resilience in ecological systems (3), we still do not know whether and how past environmental regimes shape the resilience of extant species (4). Resolving this knowledge gap is pivotal for identifying those species most vulnerable to future increases in environmental stochasticity (5, 6), and thus for designing effective ecosystem management strategies (7).

While the resilience of ecological systems has attracted much attention for decades (8, 9), approaches to evaluate the resilience of natural populations and communities often overlook its short-term nature (10, 11) (but see (12)). Classic life-history theory posits that organisms operate under a strong trade-off coordinating their investments across different vital rates (13, 14). Indeed, considerable attention has been directed at using the vital rates of survival, progression (*e.g.*, growth, development), retrogression (*e.g.*, shrinkage [15], rejuvenation [16]), and reproduction to describe how energetic trade-offs shape population performance when exposed to disturbances (17–21) (*i.e.,* external (a)biotic forces that can modify the composition of a system). Yet, attention has largely focused on the use of stochastic derivatives of long-term (*i.e.,* asymptotic) population characteristics (such as stochastic population growth rate, λ_s) as measures of population performance in the face of recurrent disturbances (22). However, disturbances can and oftentimes displace populations into a transient (*i.e.*, short-term) phase, where their dynamics can vary considerably from their asymptotic trajectories (23) (Fig. 1A). Long-term measures of population performance are, thus, unlikely to reveal how environmental stochasticity influences the resilience of populations (10). Instead, transient population metrics, describing how the dynamics of

populations can change following disturbance relative to their long-term characteristics (24, 25), can offer greater insight.

Natural populations exposed to frequent disturbances are expected to experience a selection for traits, and vital rates, that enhance their resilience (26, 27). Within more stochastic environments populations endure more frequent (a)biotic fluctuations, each with the capacity to impose shifts in their structural composition. Following these disturbances, characteristics that inhibit transient declines in population growth, or those promoting short-term increases are likely more beneficial than those maximising long-term performance (12, 28). Logically, therefore, increased stochasticity should select for increased *resistance* (*i.e.*, a population's ability to avoid a decline in size after disturbance) and/or *compensation* (the ability to increase in size after disturbance), and shorter *recovery time* (henceforth recovery, the time needed for converging back to a stationary equilibrium after disturbance) (29, 30) (Fig. 1A). Accordingly, evaluating variation in the transient characteristics of populations across a gradient in environmental stochasticity presents a unique opportunity for exploring how population resilience is shaped by stochastic environments (11, 29–31). Moreover, this approach represents a space-for-time substitute (32) for predicting the performance of populations under future scenarios of increased stochasticity and more frequent recurrent disturbances.

Here, we provide a global assessment of the environmental determinants of population resilience, examining how short-term environmental drivers mediate long-term evolutionary outputs. Specifically, we quantify the inherent capacity for resistance, compensation, and recovery of 2,242 natural populations across 61 animal, 305 plant, and 3 algae species from two global demographic databases (33, 34) (Fig. 1B & Supplementary Table S1). Building on the expectation that past disturbance regimes have shaped current resilience (27, 35), we anticipated that (H1a) exposure to higher frequency environmental

stochasticity will select for faster recovery, whilst (H1b) exposure to broader spectra in

environmental stochasticity selects for increased resistance and reduced compensation. With

the vital rates of survival, progression, retrogression, and reproduction describing how trade-

offs in individual-level fitness translates into population-level performance (36), we expected

(H2) patterns in population resilience to correspond with the underlying vital rates of

populations; with investments into individual survival associated with enhanced resistance

and greater reproductive investments aligned with faster recovery.

Results and Discussion

Sourcing climate records at the necessary temporal and spatial resolution for quantifying

deep-time environmental legacies presents a considerable challenge (37, 38). Thus, a

population's resilience to future global change is more often inferred from its exposure to

more contemporary conditions (4, 39). Here, however, using phylogenetically-corrected

partial least squares regression (pPLS) and Pearson's tests of correlation, we found that no

measure of environmental stochasticity constitutes a strong predictor of variation in the

resistance, compensation, and recovery attributes of populations (Fig. 2 & Table 1). This

finding was insensitive to both phylogenetic imputation (40) (Fig. S3) and the length of time

used to quantify recent-past exposure to environmental stochasticity (Fig. S6 & S7).

120   Using a novel framework, based on punctual disturbances altering population

structure, to quantify demographic resilience (30), we estimated the six transient metrics

comprising resistance (first-step attenuation, $\underline{\rho}$ & maximal attenuation, $\underline{\rho}_{max}$), compensation

(reactivity, $\overline{\rho}$ & maximal amplification, $\overline{\rho}_{max}$), and recovery (damping ratio, ρ & period of

oscillation, ψ) in each population following disturbance (25, 30, 41) (Fig. 1A). Next, we

quantified the exposure of these populations to environmental stochasticity using measures of

mean thermal range (m), as well as the spectral frequency and autocorrelation of temperature

(β_T & a_T) and precipitation regimes (β_P & a_P) they experienced during the 50-years preceding

the beginning of each study (Supplementary S4). Over any given timeframe, shorter-lived

populations (mean life expectancy $\leq$ 10 years) are likely to have had a larger number of

generations exposed to local stochasticity, thereby offering greater opportunity for adaptive

change. Yet, our finding persists irrespective of mean life expectancy, with recent-past

exposure to environmental stochasticity having an equally negligible influence on the

resilience attributes of populations of both long- and short-lived species (Fig. S5).

Resistance and compensation, but not recovery, are determined by energetic

investments across somatic maintenance/development and reproduction. Using pPLS across

our 2,242 populations, we evaluated the relationship between our six measures of

demographic resilience (Fig. 1A) and their sensitivities to each of the vital rates of survival

(σ), progression (γ), retrogression (τ), and fecundity (φ; Fig. 3). These vital rate sensitivities

reflect how much each transient metric would change following perturbations in each vital

rate (41). Thus, these sensitivities highlight how investments into any one vital rate influence

a populations' capacity to resist, compensate, and recover following a disturbance, and so

provide a measure of the absolute importance of each vital rate in shaping demographic

resilience. We focused on sensitivities here, rather than elasticities (proportional sensitivities

[42]), as the former provide a closer representation of selection gradients (43). Patterns across

the vital rate sensitivities of each transient characteristic corroborate recent evidence that the

resistance and compensation attributes of populations are constrained by individual-level

investments across survival, progression, retrogression, and reproduction (44). Meanwhile,

we found no evidence of demographic selection pressures on the attribute of recovery (Fig. 3

& Table 2). Again, these findings are insensitive to phylogenetic imputations (Fig. S4).

A   strong   phylogenetic   signal   across   energetic   investments   into   resistance,

compensation, and recovery suggests that species demographic resilience is not shaped by

recent environmental patterns, but by phylogenetic ancestry. Using estimates of phylogenetic

signal (Pagel's λ [45]; not to be confused with long-term population growth rate, λ) we

demonstrate a strong phylogenetic signal (Pagel's $\lambda > 0.940$, p <0.001) across the transient

metrics of resistance (first-timestep attenuation & maximal attenuation) and compensation

(reactivity & maximal amplification), and their corresponding vital rate sensitivities (Table

3). Phylogenetic signal reflects the proportion of variation in a trait across different species

that can be explained by their shared evolutionary history (45, 46). As such, this analysis

quantifies the extent to which evolution may have shaped the observed resilience attributes of

natural populations. We observed little association between the phylogenetic patterns of

resilience and the environmental stochasticity regimes to which our examined natural

populations were exposed. Selection pressures constrain how individuals allocate finite

resources across survival, somatic development, and reproduction, thus mediating the

capacity for populations to exploit and prevail within their local environments (14, 26). Over

time, these selective forces have moulded the resilience attributes of populations, but not in

response to the stochasticity regimes they currently endure (47, 48). It is necessary, therefore,

to highlight the limitations associated with using the recent exposure of populations to

environmental stochasticity as a predictor of their continued resilience to future disturbances.

Caution is necessary when interpreting our results regarding the selection pressures

maintained by environmental stochasticity. Our exploration into the environmental drivers of

demographic resilience focuses only on terrestrial populations. Compared with marine

systems, terrestrial environments afford organisms with more opportunities for seeking out

tolerable microclimates (49), thereby reducing the susceptibility of terrestrial taxa to

environmental shifts. By contrast, marine species inhabit conditions closer to their

physiological limits (50), rendering them more sensitive to abiotic shifts. More importantly,

though, here we quantify past exposure to environmental stochasticity using maximum

exposure legacies of 100-years. It is plausible, however, that deep-time environmental regimes would offer greater predictive potential. Indeed, deep-time studies have demonstrated how the latitudinal diversity gradient, a prominent pattern underlying contemporary ecological understanding, has only persisted during the past 30 million years (51). Whilst environmental stochasticity is known to influence population dynamics (52), its observable effects on population characteristics can remain negligible until compounded by external factors such as changing habitat configurations (53). Any direct impacts of stochasticity on the resilience of natural populations may, therefore, become more detectable with time.

Despite the potential influence of historical climate legacies on the resilience of natural populations, our results show a robust lack of influence of recent-past climate on population resilience, even after explicitly accounting for lifespan and legacy time. Adaptation is the accumulation of beneficial, heritable characteristics within a population over multiple generations. Therefore, populations exposed to specific conditions over extended timeframes, or those that turn over multiple generations within a short period of time, are more likely to display trait characteristics adapted to those conditions (47, 48, 54). To evaluate the implications of differing exposure periods on our observations, we first repeated our pPLS analyses exploring the relationship between demographic resilience and environmental stochasticity focusing on only populations of short-lived species (Supplementary S6). We determined population longevity using estimates of mean life expectancy (η_e), with populations of η_e ≤10-years defined as short-lived (n = 1,606 populations). Next, we repeated our analyses using only populations for which 100-year abiotic legacies could be sourced (Supplementary S7). Neither approach improved our ability to predict the demographic resilience attributes of natural populations using measures of environmental stochasticity (Figs. S5, S6 & S7).

Conclusions

Global change is changing the periodicity of phenological drivers (55, 56), reducing return

times between severe disturbance events (57, 58). Natural populations worldwide are thus

being exposed to increasingly stochastic environments, with many facing imminent

collapse(59). Here, we illustrate, however, that the recent-past exposure (last 50-100 years) of

natural populations to environmental stochasticity does not constrain, nor guarantee, their

resilience towards future climatic shifts. This realisation that population resilience appears

constrained by the adaptation of species over longer-term evolutionary timeframes, not the

conditions experienced by populations following their dispersal, is not trivial. Rather, this key

insight can help to focus current debates on the concept of resilience (9, 29, 60), enabling our

attention to be directed towards identifying populations with high adaptive potential, and

maintaining the high genetic diversity necessary to enhance the resilience of natural

populations. Reliant on preserving and enhancing the earth's natural capital, initiatives such

as the United Nations Sustainable Development Agenda are interwoven with the

conservation-orientated goals of our responsibility for environmental stewardship (61, 62).

Accordingly, a detailed understanding of the mechanisms that do and do not promote the

resilience of natural populations is imperative for not only achieving global conservation

success, but also for securing our socioeconomic future.

Methods

Demographic data extraction & transient indices

To evaluate the selection pressures on the resilience attributes of natural populations, we

extracted Matrix Population Models (MPMs) from the open-source COMPADRE Plant

Matrix Database (34) (v. 5.0.1) and COMADRE Animal Matrix Database (33) (v. 3.0.1).

MPMs are discrete time, structured population models where the lifecycle is categorised into

discrete state classes (*i.e.*, age, size, and/or developmental stages) (41). Combined,

COMPADRE and COMADRE contain over 12,000 MPMs from more than 1,100 animal and

plant species. However, here we only retained MPMs satisfying the following six criteria to

test our hypotheses: 1) MPMs reflecting the demographic characteristics of individual

populations recorded across a single time period (*Individual* MPMs). We did, however, select

MPMs consisting of demographic information averaged across multiple populations and/or

time periods (*mean* MPMs), for populations for which no individual matrices were available

(330 populations after applying the additional criteria below); 2) MPMs based on annual

surveys to ensure all subsequent metrics obtained reflected identical units of time, thus

allowing for their comparability; 3) MPMs representing wild, un-manipulated populations, to

guarantee investigating the selection pressures underpinning the resilience of natural

populations and the possibility to link their dynamics to their local environmental regimes; 4)

MPMs comprised of three or more life stages, as lower dimension MPMs typically lack the

necessary resolution for estimating vital rates (63) and transient dynamics (64); 5) MPMs

from populations with known latitude and longitude information to allow us to link their

demographic properties to local environmental regimes; and finally 6) MPMs describing full

life cycles (e.g., no missing data on survival, progression, retrogression, and reproduction) to

ensure the correct calculation of vital rates and transient metrics. Following these criteria, we

retained 3,890 MPMs corresponding with 3,204 populations across 438 plant species, 665

populations across 112 animal species, and 21 populations across six algae species

(Supplementary Table S1).

We further refined our list of MPMs according to their transient, asymptotic, and

species-specific properties. All MPMs were tested for irreducibility (*i.e.,* all life cycle stages

are either directly, or indirectly connected to one another), ergodicity (*i.e.,* asymptotic

dynamics are independent of the initial population structure), and primitivity (*i.e.,* MPMs

consist of non-negative elements) (41). We carried out these tests using the corresponding

isErgodic, *isIrreducible*, and *isPrimitive*, diagnostic functions from the R package '*popdemo*'

(65). A total of 1,203 reducible, imprimitive, and/or non-ergodic MPMs were excluded from

further analyses on the basis that they represent untenable life cycles that defy logical

biological processes (66). MPMs with population growth rates $\lambda > 2$, indicating that the

population is projected to increase two-fold or more every year, were also rejected as they

represent unlikely realisations of population performance in our experience. Equally, MPMs

from highly migratory (e.g., > 1,000 km) species were discarded, since their vital rate

schedules are unlikely to be mostly shaped by the environment in which they were measured.

We also note here that, across our initial population sample, the vital rate of clonality (κ) was

rare, with only 140 populations across 37 plant species, and two populations from one animal

species (*Amphimedon compressa* [67]) explicitly exhibiting this demographic process. Thus,

to focus our analyses on common demographic currencies, we excluded all populations

exhibiting clonality. Overall, our strict selection criteria resulted in a final sample of 2,242

MPMs, corresponding with 369 species: 402 populations from 61 animal species, 1,830

populations from 305 plant species, and 10 populations from three species of algae

(Supplementary Table S1).

For each retained MPM, we calculated six transient metrics quantifying each

population's potential for demographic resistance (first-timestep attenuation, $\underline{\rho}$ & and

maximal attenuation, $\underline{\rho}_{max}$), compensation (reactivity, $\overline{\rho}$ & maximal amplification, $\overline{\rho}_{max}$), and

recovery (damping ratio, ρ & period of oscillation, ψ), following a disturbance (30) (Fig. 1).

Firstly, with estimates of transient dynamics known to be contingent on the reproductive

strategies of populations, it was necessary to convert all post-reproductive matrices into a pre-

reproductive format by adjusting patterns of reproduction to include a measure of adult

survival (68). All MPMs were then standardised to separate their transient and asymptotic

properties by dividing each matrix element by the MPM's dominant eigenvalue, λ (24, 41).

Following standardisation, from each MPM we obtained estimates of reactivity and first-

timestep attenuation, as the upper and lower bounds returned by the function *reac*, and

calculated maximal amplification and maximal attenuation indices using their respective

functions *maxamp* and *maxatt*, all within the R package '*popdemo*' (65). Meanwhile, we

calculated the damping ratio of each MPM using *eigen.analysis* from the '*popbio*' package

(69). Finally, we calculated the period of oscillation (ψ) for each MPM using its subdominant

eigenvalue (λ_2) (equation 1) (41).

$$\psi = \frac{2\pi}{\theta_2} \qquad \text{Where: } \theta_2 = \tan^{-1}\left(\frac{I(\lambda_2)}{R(\lambda_2)}\right) \tag{1}$$

To explore how the fitness components of individuals mediate the selection gradients

placed on demographic resilience by environmental variability, we calculated the sensitivity

of each transient metric towards the vital rates of survival (σ), progression (γ), retrogression

(τ), and fecundity (φ). For each MPM, we first estimated all vital rate sensitivities from their

element-level constituents. Individual elements a_{ij} within the MPM **A** typically describe

combinations of multiple vital rates (70). Subsequently, calculating the sensitivity of each

transient metric (s_x) with respect to underlying vital rates requires the decomposition of

element-level sensitivities into their vital rate components (70). Briefly, this decomposition

requires the estimation of stage-specific survival probabilities (σ_j) for each MPM. These

estimates of σ_j are then used to determine the proportion of each matrix element a_{ij}

corresponding with survival (σ), progression (γ), retrogression (τ), and fecundity (φ) (70). We

initially calculated the sensitivity of each transient metric at the matrix element-level (s_{ij}).

The sensitivities of the damping ratio (ρ) and period of oscillation (ψ) were determined using

equations 2 & 3, with s_{ij}^1 and s_{ij}^2 as the matrix element-level sensitivity matrices of the

dominant and subdominant eigenvalues, respectively (41).

$$s_{\rho_{ij}} = \frac{1}{|\lambda_2|}\left(s_{ij}^1 - \frac{\rho}{|\lambda_2|}\left[R\left(s_{ij}^2\right) + I\left(s_{ij}^2\right)\right]\right) \tag{2}$$

$$s_{\psi_{ij}} = \frac{-2\pi}{\theta_2|\lambda_2|^2}\left[I\left(s_{ij}^2\right) - R\left(s_{ij}^2\right)\right] \tag{3}$$

The sensitivities of reactivity, amplification, and attenuation (s_{ij}) with respect to

element a_{ij} were then estimated as the magnitude of change (δ) in each transient metric (x)

following a small change (here 0.01) in a_{ij} (Equation 4) (71).

$$s_{ij} = \frac{\delta x}{\delta a_{ij}} = \frac{x_{Perturbed} - x_{Original}}{a_{ij}^{Perturbed} - a_{ij}^{Original}} \tag{4}$$

The distributions of each transient metric and its corresponding vital rate sensitivities

were checked prior to subsequent regression analyses. To ensure normality across each

distribution, outliers, defined as values outside the 95% confidence intervals of the

distribution, were omitted, with the remaining estimates then transformed if necessary. For

each transient metric, power transformations (y^x) were used to achieve approximate normality

using the Box-Cox transformation functions of the R package '*caret*' (72) to estimate *x*. The

distributions of damping ratio, period of oscillation, reactivity, and maximal amplification

raised negative *x* values, and so their transformations took the form $1/y^{|x|}$. Inverse and log

transformations were also necessary for several of the vital rate sensitivity variables (See

Table S2 for further details).

Phylogenetic correction

Evaluating the selection pressures exerted on attributes of demographic resilience across

multiple species requires an explicit consideration for how traits are expected to covary due

to phylogenetic relationships (46, 73, 74). To account for such relationships in our analyses,

we constructed a population-within-species level phylogenetic tree using phylogenetic data

extracted from the Open tree of Life for the animal, plant, and algae species across our

population sample (OTL [75]; Supplementary S3). Our approach here also allowed us to

accommodate studies that included multiple, separate populations for the same species (*see*

below). Firstly, the scientific names of each species associated with our extracted MPMs

were checked against current taxonomy records using the R package '*taxize*' (76). Next, we

extracted information regarding the taxonomic classification and phylogeny of each species

from the OTL database with the R package '*rotl*' (77). Subsequently, using the '*ape*' (78) and

'*phytools*´ packages (79), this phylogenetic information was used to construct a species-level

phylogenetic tree corresponding with the 369 unique species within our MPM list.

Beyond accounting for phylogenetic signals in trait variance-covariance across our

population sample, it was necessary to ensure that our phylogenetic tree reflected the

influence of spatial signals in the development of traits within species. Thus, we expanded

our phylogenetic tree by adding branch tips to incorporate multiple population entries per

species (*sensu* [80]), generating a population-level tree comprising our full sample of 2,242

populations (Supplementary S3). Finally, we computed the branch lengths for our

phylogenetic tree using the function *compute.brlen* in the R package '*ape*' (78). These branch

lengths were estimated using Grafen's arbitrary branch lengths (81), assuming a Brownian

motion model with the variance between species directly proportional to time since

divergence (82). Importantly, we constrained branch lengths between populations of the same

species to approximately zero (0.0000001) under the assumption of negligible phylogenetic

distance between species replicate populations.

Quantifying environmental variability

To investigate the role of environmental selection pressures on the resistance, compensation,

and recovery components of resilience in natural populations, we used a pPLS regression

exploring the association between transient characteristics and metrics of environmental

variability. We quantified the magnitude and frequency of environmental variation to which

each population was exposed using the geographical location information extracted with each

MPM from COMPADRE & COMADRE (33, 34) (Supplementary S4). Since temperature

and precipitation rates are universal drivers of biological community assembly across terrestrial environments (83), we selected data describing temporal trends in thermal and precipitation regimes as a measure of the environmental variability experienced by each population. Crucially, however, with precipitation not directly influencing marine environments (although see [84]), we excluded marine populations (29 populations from six animal species, and 10 populations from three algal species) from this portion of our analyses.

We quantified environmental variance through the metrics of autocorrelation, abiotic range, and frequency spectrum using long-term temperature and precipitation records sourced from the CHELSA climate database (85). For each population, we extracted monthly records of maximum and minimum temperatures (°C) and mean precipitation rates (kg m^{-2}) corresponding with the specific period during which the population was surveyed as detailed in COMPADRE and COMADRE, plus an additional 50-years prior to the onset of censusing to account for environmental legacy effects (86). We condensed maximum and minimum temperature readings into monthly estimates of mean temperatures and thermal range. Next, as a gauge of disturbance magnitude (*m*), we estimated the mean thermal range experienced by each population across their associated temporal records. We then arranged our monthly mean temperature and precipitation estimates into time series depicting the temporal environmental regimes to which each population was exposed. Using the '*colorednoise*' package (87), we calculated the autocorrelation of each temperature (a_T) and precipitation (a_P) time series as a measure of environmental predictability (88).

The colour of environmental variation is depicted on a red to blue colour scale, from lower to higher frequencies, respectively (88). The frequency spectrum of a time series is expressed by its spectral exponent (*β*), which is calculated as the negative slope of the linear regression between the log spectral density and log frequency of the time series (89). We

calculated the frequency spectrum of each precipitation (β_P) and temperature (β_T) time series

as an indicator of the colour of environmental variation experience by each population (89).

The spectral exponent for each time series was estimated using the spectrum function from

the R package '*stats*' (90).

Partial Least Squares Regression

We utilised a phylogenetically corrected Partial Least Squares regression (pPLS) framework

to test our hypothesis that the resistance, compensatory, and recovery of natural populations

correspond with gradients in environmental variability and evaluate how this is mediated by

fitness investments. Using a pPLS, we evaluated the relationship between our estimates of

population resilience, and both associated environmental variability regimes and their vital

rate sensitivities. The pPLS technique is considered a more powerful comparative tool than

other available multivariate regression methods (91), as it simultaneously condenses the

variation among numerous predictors whilst maximising the variance explained among

response variables. Subsequently, we investigated the selection pressures on the resistance,

compensation, and recovery attributes of natural populations, and therefore the capacity for

environmental legacies, and vital rate characteristics, to serve as predictors of resilience

attributes.

We first applied a phylogenetically-corrected Pearson's test of correlation and pPLS

to analyse the correlation between environmental variability and transient demographic

characteristics. This approach enabled us to test for covariation between the transient

characteristics of populations and gradients in their exposure to environmental variability.

pPLS tests were carried out for each transient measure with the predictor variable set

comprised of our five metrics of environmental variability. From each test, we then extracted

component scores and loadings, which describe the arrangement of the environmental

predictor variables within a multivariate space. We also obtained the percentage variance

(*%var*) among the predictors explained by each regression component and the proportion of

variance in the transient response variable explained by each component (r^2) to estimate the

strength of any association between environmental variability and the transient dynamics of

our population sample.

Phylogenetically corrected correlation tests and pPLS analyses were used to examine

for patterns between each transient characteristics and its associated vital rate sensitivities.

Again, test coefficients (*r*), component scores, loadings, *%var*, and r^2 values were calculated

to quantify the influence of the fitness components of survival, progression, retrogression,

and reproduction towards the transient characteristics of natural populations. All pPLS

analyses were conducted using the '*pls*' R package (92), with modifications included to

ensure our analyses accounted for any evolutionary covariance between the sensitivity

patterns and transient characteristics of different populations (79, 93, 94).

We carried out all pPLS analyses using only complete entries, omitting populations

missing estimates for any one variable. To provide further clarity regarding any patterns we

observed between our measures of population resilience, their environmental legacies, and

their vital-rate sensitivities, we repeated each analysis twice. During these repeated tests, we

first evaluated whether considering the life expectancies of populations influenced any

observed patterns (Supplementary S6). Within a given time period, long-lived species will

likely experience fewer generations than shorter-lived species diminishing the relative impact

of existing selection pressures on their trait characteristics (95). Thus, it was necessary to

ensure that the inclusion of long-lived species within our population sample did not limit our

capacity for exploring environmental selection pressures. We categorised each population

within our sample as either long- or short-lived according to their associated mean life

expectancy (η_e), calculated from each extracted MPM using the R package '*IPMpack*' (96).

Next, we repeated our pPLS analyses using only 'short-lived' populations for which $\eta_e \leq 10$

years (n = 1606 populations). This threshold was selected as a balance between maximising

the number of generations experienced by populations during the multi-year time series used

in calculating environmental legacies and maximising our sample size.

Understanding patterns in the responses of natural populations to environmental

change is often impeded by gaps in available data linking population characteristics to abiotic

regimes (97). To ensure that missing data was not obscuring any correlation between our

measures of population resilience and environmental stochasticity we repeated our analyses

using phylogenetic imputation to estimate any demographic measures of resilience missing

across our population sample (40) (Supplementary S5). We calculated the phylogenetic signal

(Pagel's λ [45]) of each transient and sensitivity variable using the *phylosig* function from the

'*phytools*' package (79). Pagel's λ exists on the scale $0 < \lambda > 1$, with 0 indicating traits have

evolved independently of phylogeny, and 1 representing a high phylogenetic signal (45). For

any variable exhibiting a strong phylogenetic signal (*i.e.,* Pagel's $\lambda \geq 0.65$) that differs

significantly from zero ($p < 0.05$), we imputed all missing values assuming a Brownian

motion evolutionary model, before repeating our pPLS analyses.

Acknowledgments

Funding for this research was provided by a DTP Natural Environment Research Council

Scholarship to JC, a NERC IRF (NE/M018458/1) to RS-G, a Ramon Areces Foundation

Postdoctoral Fellowship to PC, and Winifred Violet Scott Estate funding to MB. We

acknowledge the University of Leeds High Performance Computing (ARC) team, who

maintain the facilities used for the extraction of extensive climate records from CHELSA

(https://chelsa-climate.org/).

Author Contributions

JC, RS-G, MB, and PC conceived the ideas and methodology for the project. JC, PC, and RS-

G carried out data extraction and analysis. JC led the writing of the manuscript, with close

support from RS-G, and all authors contributing critically to the writing and giving final

approval for publication.

Data Accessibility

All Matrix Projection Models used within this study are available from the COMADRE and

COMPADRE demographic databases (https://compadre-db.com/). Equally, all taxonomic

information used in the construction of our phylogenetic trees can be sourced from the Open

Tree of Life (https://tree.opentreeoflife.org/), and all temperature, and precipitation records

used in this study can be downloaded from the CHELSA climate database (https://chelsa-

climate.org/). Finally, the R code used for all the data extraction and analyses presented

throughout this manuscript have been uploaded to GitHub and will be made publicly

available following publication.

References

1.   R. J. Standish, *et al.*, Resilience in ecology: Abstraction, distraction, or where the

action is? *Biol Conserv* **177**, 43–51 (2014).

2.   D. G. Angeler, C. R. Allen, Quantifying resilience. *Journal of Applied Ecology* **53**,

617–624 (2016).

3.   C. S. Holling, Resilience and stability of ecological systems. *Annu Rev Ecol Syst* **4**, 1–

23 (1973).

4.   B. H. Walker, Resilience: what it is and is not. *Ecology and Society* **25**, 1–3 (2020).

5.   J. C. Gaillard, Vulnerability, capacity and resilience: Perspectives for climate and

development policy. *J Int Dev* **22**, 218–232 (2010).

6. E. McLeod, E. C. Shaver, M. Beger, J. Koss, G. Grimsditch, Using resilience

assessments to inform the management and conservation of coral reef ecosystems. *J*

Environ Manage **277**, 1–8 (2021).

7. R. L. Pressey, M. Cabeza, M. E. Watts, R. M. Cowling, K. A. Wilson, Conservation

planning in a changing world. *Trends Ecol Evol* **22**, 583–592 (2007).

8. P. Capdevila, *et al.*, Reconciling resilience across ecological systems, species and

subdisciplines. *Journal of Ecology* **109**, 3102–3113 (2021).

9. S. Kéfi, *et al.*, Advancing our understanding of ecological stability. *Ecol Lett* **22**,

1349–1356 (2019).

10. J. Cant, R. Salguero-Gómez, M. Beger, Transient demographic approaches can

drastically expand the toolbox of coral reef science. *Coral Reefs* **41**, 885–896 (2022).

11. A. Hastings, *et al.*, Transient phenomena in ecology. *Science (1979)* **361**, 1–9 (2018).

12. J. L. McDonald, I. Stott, S. Townley, D. J. Hodgson, Transients drive the demographic

dynamics of plant populations in variable environments. *Journal of Ecology* **104**, 306–

314 (2016).

13. S. C. Stearns, *The Evolution of Life Histories* (Oxford University Press, 1992).

14. S. C. Stearns, Trade-Offs in Life-History Evolution. *Funct Ecol* **3**, 259–268 (1989).

15. R. Salguero-Gómez, B. B. Casper, Keeping plant shrinkage in the demographic loop.

Journal of Ecology **98**, 312–323 (2010).

16. R. Salguero-Gómez, R. P. Shefferson, M. J. Hutchings, Plants do not count... or do

they? New perspectives on the universality of senescence. *Journal of Ecology* **101**,

545–554 (2013).

17. M. S. Boyce, *et al.*, Demography in an increasingly variable world. *Trends Ecol Evol*

21, 141–148 (2006).

18.   N. Jonzén, T. Pople, J. Knape, M. Sköld, Stochastic demography and population

500   dynamics in the red kangaroo Macropus rufus. *Journal of Animal Ecology* **79**, 109–116

501   (2010).

502   19.   S. D. Tuljapurkar, Population dynamics in variable environments. II. Correlated

503   environments, sensitivity analysis and dynamics. *Theor Popul Biol* **21**, 114–140

504   (1982).

505   20.   S. D. Tuljapurkar, S. H. Orzack, Population dynamics in variable environments I.

506   Long-run growth rates and extinction. *Theor Popul Biol* **18**, 314–342 (1980).

507   21.   S. Tuljapurkar, J. M. Gaillard, T. Coulson, From stochastic environments to life

508   histories and back. *Philosophical Transactions of the Royal Society B* **364**, 1499–1509

509   (2009).

510   22.   E. E. Crone, *et al.*, How do plant ecologists use matrix population models? *Ecol Lett*

511   **14**, 1–8 (2011).

512   23.   A. Hastings, Transient dynamics and persistence of ecological systems. *Ecol Lett* **4**,

513   215–220 (2001).

514   24.   D. N. Koons, J. B. Grand, B. Zinner, R. F. Rockwell, Transient population dynamics:

515   Relations to life history and initial population state. *Ecol Modell* **185**, 283–297 (2005).

516   25.   I. Stott, S. Townley, D. Hodgson, A framework for studying transient dynamics of

517   population projection matrix models. *Ecol Lett* **14**, 959–970 (2011).

518   26.   R. Lande, S. J. Arnold, The measurement of selection on correlated characters.

519   *Evolution (N Y)* **37**, 1210–1226 (1983).

520   27.   P. W. Boyd, *et al.*, Biological responses to environmental heterogeneity under future

521   ocean conditions. *Glob Chang Biol* **22**, 2633–2650 (2016).

28. J. Cant, *et al.*, Transient amplification enhances the persistence of tropicalising coral

populations in marginal high latitude environments. *Ecography* (2022)

https:/doi.org/10.1101/2021.04.23.441133.

29. D. Hodgson, J. L. McDonald, D. J. Hosken, What do you mean, "resilient"? *Trends*

Ecol Evol **30**, 503–506 (2015).

30. P. Capdevila, I. Stott, M. Beger, R. Salguero-Gómez, Towards a comparative

framework of demographic resilience. *Trends Ecol Evol* **35**, 776–786 (2020).

31. A. Hastings, Transients: The key to long-term ecological understanding? *Trends Ecol*

Evol **19**, 39–45 (2004).

32. S. T. A. Pickett, "Space-for-time substitutions as an alternative to long-term studies."

in *Long-Term Studies in Ecology*, G. E. Likens, Ed. (Springer, 1989), pp. 110–135.

33. R. Salguero-Gómez, *et al.*, COMADRE: A global data base of animal demography.

Journal of Animal Ecology **85**, 371–384 (2016).

34. R. Salguero-Gómez, *et al.*, The COMPADRE Plant Matrix Database: An open online

repository for plant demography. *Journal of Ecology* **103**, 202–218 (2015).

35. E. B. Rivest, S. Comeau, C. E. Cornwall, The role of natural variability in shaping the

response of coral reef organisms to climate change. *Curr Clim Change Rep* **3**, 271–281

(2017).

36. C. J. E. Metcalf, S. Pavard, Why evolutionary biologists should be demographers.

Trends Ecol Evol **22**, 205–212 (2007).

37. A. M. Dixon, P. M. Forster, M. Beger, Coral conservation requires ecological climate-

change vulnerability assessments. *Front Ecol Environ* **19**, 243–250 (2021).

38. L. Kwiatkowski, P. R. Halloran, P. J. Mumby, D. B. Stephenson, What spatial scales

are believable for climate model projections of sea surface temperature? *Clim Dyn* **43**,

1483–1496 (2014).

39.   W. B. Foden, *et al.*, Climate change vulnerability assessment of species. *Wiley*

Interdiscip Rev Clim Change **10**, 1–36 (2019).

40.   T. D. James, R. Salguero-Gómez, O. R. Jones, D. Z. Childs, A. P. Beckerman,

Bridging gaps in demographic analysis with phylogenetic imputation. *Conservation*

Biology **35**, 1210–1221 (2021).

41.   H. Caswell, *Matrix population models: Construction, analysis and interpretation*, 2nd

Edition (Oxford University Press Inc, 2001).

42.   H. de Kroon, J. van Groenendael, J. Ehrlén, Elasticities: a review of methods and

model limitations. *Ecology* **81**, 607–618 (2000).

43.   P. H. van Tienderen, Elasticities and the link between demographic and evolutionary

dynamics. *Ecology* **81**, 666–679 (2000).

44.   P. Capdevila, *et al.*, Life history mediates the trade-offs among different components

of demographic resilience. *Ecol Lett* **00**, 1–14 (2022).

45.   M. Pagel, Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884

(1999).

46.   R. P. Freckleton, P. H. Harvey, M. Pagel, Phylogenetic analysis and comparative data:

A test and review of evidence. *American Naturalist* **160**, 712–726 (2002).

47.   G. Bell, Evolutionary rescue and the limits of adaptation. *Philosophical Transactions*

of the Royal Society B: Biological Sciences **368**, 1–6 (2013).

48.   G. Bell, A. Gonzalez, Adaptation and evolutionary rescue in metapopulations

experiencing environmental deterioration. *Science (1979)* **332**, 1327–1330 (2011).

49.   J. M. Sunday, *et al.*, Thermal-safety margins and the necessity of thermoregulatory

behavior across latitude and elevation. *Proc Natl Acad Sci U S A* **111**, 5610–5615

(2014).

50.   M. L. Pinsky, A. M. Eikeset, D. J. McCauley, J. L. Payne, J. M. Sunday, Greater

vulnerability to warming of marine versus terrestrial ectotherms. *Nature* **569**, 108–111

(2019).

51.   P. D. Mannion, P. Upchurch, R. B. J. Benson, A. Goswami, The latitudinal

biodiversity gradient through deep time. *Trends Ecol Evol* **29**, 42–50 (2014).

52.   R. Lande, S. Engen, B. E. Sther, *Stochastic population dynamics in ecology and*

conservation (Oxford University Press, 2003).

53.   J. M. Fraterrigo, S. M. Pearson, M. G. Turner, Joint effects of habitat configuration

and temporal stochasticity on population dynamics. *Landsc Ecol* **24**, 863–877 (2009).

54.   A. Compagnoni, *et al.*, Herbaceous perennial plants with short generation time have

stronger responses to climate anomalies than those with longer generation time. *Nat*

Commun **12**, 1–8 (2021).

55.   M. Paniw, N. Maag, G. Cozzi, T. Clutton-Brock, A. Ozgul, Life history responses of

meerkats to seasonal changes in extreme environments. *Science (1979)* **363**, 631–635

(2019).

56.   L. S. Cordes, *et al.*, Contrasting effects of climate change on seasonal survival of a

hibernating mammal. *Proc Natl Acad Sci U S A* **117**, 18119–18126 (2020).

57.   P. K. Thornton, P. J. Ericksen, M. Herrero, A. J. Challinor, Climate variability and

vulnerability to climate change: A review. *Glob Chang Biol* **20**, 3313–3328 (2014).

58.   T. P. Hughes, *et al.*, Spatial and temporal patterns of mass bleaching of corals in the

Anthropocene. *Science (1979)* **359**, 80–83 (2018).

59.   T. M. Lenton, *et al.*, Climate tipping points — too risky to bet against. *Nature* **575**,

592–595 (2019).

60.   J. Ingrisch, M. Bahn, Towards a comparable quantification of resilience. *Trends Ecol*

Evol **33**, 251–259 (2018).

61.   N. Seddon, *et al.*, Understanding the value and limits of nature-based solutions to

climate change and other global challenges. *Philosophical Transactions of the Royal*

Society B: Biological Sciences **375** (2020).

62.   B. Walker, "Conserving Biological Diversity through Ecosystem Resilience" (1995).

63.   R. Salguero-Gómez, J. B. Plotkin, Matrix dimensions bias demographic inferences:

Implications for comparative plant demography. *American Naturalist* **176**, 710–722

(2010).

64.   B. Tenhumberg, A. J. Tyre, R. Rebarber, Model complexity affects transient

population dynamics following a dispersal event: a case study with pea aphids.

Ecology **90**, 1878–1890 (2009).

65.   I. Stott, D. J. Hodgson, S. Townley, Popdemo: An R package for population

demography using projection matrix analysis. *Methods Ecol Evol* **3**, 797–802 (2012).

66.   I. Stott, S. Townley, D. Carslake, D. J. Hodgson, On reducibility and ergodicity of

population projection matrix models. *Methods Ecol Evol* **1**, 242–252 (2010).

67.   A. E. Mercado-Molina, A. M. Sabat, P. M. Yoshioka, Demography of the demosponge

Amphimedon compressa: Evaluation of the importance of sexual versus asexual

recruitment to its population dynamics. *J Exp Mar Biol Ecol* **407**, 355–362 (2011).

68.   K. Jelbert, *et al.*, Demographic amplification is a predictor of invasiveness among

plants. *Nat Commun* **10**, 1–6 (2019).

69.   C. Stubben, B. Milligan, Estimating and analyzing demographic models using the

popbio package in R. *J Stat Softw* **22**, 1–23 (2007).

70.   M. Franco, J. Silvertown, A comparative demography of plants based upon elasticities

of vital rates. *Ecology* **85**, 531–538 (2004).

71.   W. F. Morris, D. F. Doak, *Quantitative conservation biology: Theory and practice of*

population viability analysis (Sinauer Associates, Inc., 2002).

72.   M. Kuhn, caret: Classification and regression training (2020).

73.   R. P. Freckleton, The seven deadly sins of comparative analysis. *J Evol Biol* **22**, 1367–
1375 (2009).

74.   L. J. Revell, Phylogenetic signal and linear regression on species data. *Methods Ecol*
Evol **1**, 319–329 (2010).

75.   C. E. Hinchliff, *et al.*, Synthesis of phylogeny and taxonomy into a comprehensive tree
of life. *Proceedings of the National Academy of Sciences* **112**, 12764–12769 (2015).

76.   S. Chamberlain, *et al.*, taxize: Taxonomic information from around the web (2020).

77.   F. Michonneau, J. Brown, D. Winter, rotl: An R package to interact with the Open
Tree of Life data. *Methods Ecol Evol* **7**, 1–17 (2016).

78.   E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35** (2018).

79.   L. J. Revell, phytools: An R package for phylogenetic comparative biology (and other
things). *Methods Ecol Evol* **3**, 217–223 (2012).

80.   R. P. Freckleton, W. Jetz, Space versus phylogeny: Disentangling phylogenetic and
spatial signals in comparative data. *Proceedings of the Royal Society B: Biological*
Sciences **276**, 21–30 (2009).

81.   A. Grafen, The phylogenetic regression. *Philos Trans R Soc Lond B Biol Sci* **326**, 119–
157. (1989).

82.   L. J. Revell, L. J. Harmon, D. C. Collar, Phylogenetic signal, evolutionary process, and
rate. *Syst Biol* **57**, 591–601 (2008).

83.   C. Howard, C. H. Flather, P. A. Stephens, A global assessment of the drivers of
threatened terrestrial species richness. *Nat Commun* **11**, 1–10 (2020).

84.   J. Haapkylä, *et al.*, Seasonal rainfall and runoff promote coral disease on an inshore
reef. *PLoS One* **6** (2011).

85. D. N. Karger, *et al.*, Climatologies at high resolution for the earth's land surface areas.

Sci Data **4**, 1–20 (2017).

86. S. Evers, *et al.*, Lagged and dormant-season climate better predict plant vital rates than

climate during the growing season. *Glob Chang Biol* **27**, 1927–1941 (2021).

87. J. Pilowsky, colorednoise: Simulate temporally autocorrelated populations (2019).

88. L. Ruokolainen, A. Lindén, V. Kaitala, M. S. Fowler, Ecological and evolutionary

dynamics under coloured environmental variation. *Trends Ecol Evol* **24**, 555–563

(2009).

89. D. A. Vasseur, P. Yodzis, The color of environmental noise. *Ecology* **85**, 1146–1152

(2004).

90. R Core Team, R: A language and environment for statistical computing. (2019).

91. L. M. Carrascal, I. Galván, O. Gordo, Partial least squares regression as an alternative

to current regression methods used in ecology. *Oikos* **118**, 681–690 (2009).

92. B. Mevik, R. Wehrens, K. Hovde Liland, pls: Partial least squares and principal

components regression (2019).

93. D. C. Adams, R. N. Felice, Assessing trait covariation and morphological integration

on phylogenies using evolutionary covariance matrices. *PLoS One* **9** (2014).

94. L. J. Revell, Size-correction and principal components for interspecific comparative

studies. *Evolution (N Y)* **63**, 3258–3268 (2009).

95. A. Robert, F. Sarrazin, D. Couvet, S. Legendre, Releasing adults versus young in

reintroductions: Interactions between demography and genetics. *Conservation Biology*

18, 1078–1087 (2004).

96. C. J. E. Metcalf, S. M. McMahon, R. Salguero-Gómez, E. Jongejans, IPMpack: An R

package for integral projection models. *Methods Ecol Evol* **4**, 195–200 (2013).

97.   D. A. Conde, *et al.*, Data gaps and opportunities for comparative and conservation

671   biology. *PNAS* **116**, 9658–9664 (2019).

Table 1. The resilience attributes of resistance (green), compensation (blue), and
recovery (orange) of natural populations do not correlate with their relative exposure to
environmental stochasticity. Using phylogenetically-corrected Pearson's tests of
correlation, we explored the association between transient metrics of demographic resistance
(first-timestep attenuation, $\underline{\rho}$ & and maximal attenuation, $\underline{\rho}_{max}$), compensation (reactivity, $\bar{\rho}$ &
maximal amplification, $\bar{\rho}_{max}$), and recovery (damping ratio, ρ & period of oscillation, ψ; Fig.
1A) and five metrics of environmental stochasticity: temperature frequency spectrum (β_T),
temperature autocorrelation (a_T), thermal range (m), precipitation frequency spectrum (β_P),
and precipitation autocorrelation (a_P). Correlation displayed using Pearson's correlation
coefficient (r).

Transient metric	Resilience attribute	β_T	a_T	m	β_P	a_P
$\underline{\rho}$	Resistance	-0.0106	0.0060	-0.0094	-0.0024	0.0042
$\underline{\rho}_{max}$		-0.0071	0.0031	-0.0121	-0.0093	0.0068
$\bar{\rho}$	Compensation	-0.0026	0.0007	0.0070	-0.0080	0.0041
$\bar{\rho}_{max}$		-0.0016	<0.0001	0.0069	-0.0026	0.0009
ρ	Recovery	0.0012	-0.0007	0.0026	-0.0124	0.0147
ψ		-0.0071	0.0133	0.0052	0.0037	-0.0070

Table 2. Variation across resistance (green) and compensation (blue) of natural
populations corresponds with the energetic investments of their individuals, but
characteristics of recovery (orange) do not. Using phylogenetically-corrected Pearson's
tests of correlation, we explored the association between the transient metrics of resistance
(first-timestep attenuation, $\underline{\rho}$ & and maximal attenuation, $\underline{\rho}_{max}$), compensation (reactivity, $\overline{\rho}$ &
maximal amplification, $\overline{\rho}_{max}$), and recovery (damping ratio, ρ & period of oscillation, ψ; Fig.
1A) and their sensitivities to the vital rates of survival (σ), progression (γ), retrogression (τ),
and reproduction (ϕ). Correlation displayed using Pearson's correlation coefficient (r), with
the colour gradation highlighting the strength of the absolute correlation between each
transient metric and its vital rate sensitivities, with darker shades representing stronger
associations.

Transient metric	Resilience attribute	Survival σ	Progression γ	Retrogression τ	Reproduction ϕ
$\underline{\rho}$	Resistance	-0.70	-0.07	0.28	-0.39
$\underline{\rho}_{max}$		-0.17	0.07	0.31	-0.34
$\overline{\rho}$	Compensation	-0.11	-0.49	0.39	-0.04
$\overline{\rho}_{max}$		-0.02	-0.49	-0.13	-0.14
ρ	Recovery	-0.06	-0.06	0.11	-0.01
ψ		<0.01	0.06	-0.08	0.08

Table 3. Variation in resistance (green), compensation (blue), and recovery (orange) is

strongly predicted by phylogenetic association. However, whilst a strong phylogenetic

signal is also evident across the vital rate sensitivities of measures of resistance and

compensation, there is a negligible phylogenetic signal across the vital rate sensitivities

of measures of recovery. To quantify the strength of statistical non-independence in the

resilience attributes of natural populations due to species' common ancestry, we estimated the

phylogenetic signal (Pagel's λ(45)) across our transient metrics of demographic resistance

(first-timestep attenuation, $\underline{\rho}$ & and maximal attenuation, $\underline{\rho}_{max}$), compensation (reactivity, $\bar{\rho}$ &

maximal amplification, $\bar{\rho}_{max}$), and recovery (damping ratio, ρ & period of oscillation, ψ; Fig.

1A), as well as their sensitivities to the vital rates of survival (σ), progression (γ),

retrogression (τ), and reproduction (ϕ). Pagel's λ(45) ranges between 0, indicating that traits

have evolved independently of phylogeny, and 1, representing a high phylogenetic signal.

Colour gradation highlights the likelihood that the phylogenetic signal observed across each

transient metric and its vital rate sensitivities differs significantly from 0 (p <0.05), with

darker shades representing stronger signals.

Transient metric	Resilience attribute	Pagel's λ	Survival σ	Progression γ	Retrogression τ	Reproduction ϕ
$\underline{\rho}$	Resistance	0.997 p <0.001	0.997 p <0.001	0.989 p <0.001	0.967 p <0.001	0.985 p <0.001
$\underline{\rho}_{max}$		0.997 p <0.001	0.993 p <0.001	0.985 p <0.001	0.952 p <0.001	0.989 p <0.001
$\bar{\rho}$	Compensation	0.998 p <0.001	0.990 p <0.001	0.999 p <0.001	0.995 p <0.001	0.996 p <0.001
$\bar{\rho}_{max}$		0.998 p <0.001	0.988 p <0.001	0.998 p <0.001	0.946 p <0.001	0.996 p <0.001
ρ	Recovery	0.996 p <0.001	0.003 p = 0.344	<0.001 p = 0.686	0.005 p = 0.094	<0.001 p >0.999
ψ		0.992 p <0.001	<0.001 p >0.999	0.179 p <0.001	<0.001 p >0.999	<0.001 p >0.999

Figure captions

Figure 1. Demographic resistance, compensation, and time of recovery across 2,242

natural populations show how population resilience corresponds with exposure to

environmental stochasticity. (A) Within stochastic environments, the dynamics of

populations can vary considerably from long-term (*i.e.*, asymptotic) expectations. Under

stable stationary conditions, populations display asymptotic growth trajectories whereby the

size of a population changes at a constant rate (λ). However, following a disturbance (*),

populations can enter a transient state during which their growth rate can differ significantly

from asymptotic expectations. The duration and form of this transient phase depends, in

addition to key moments of disturbances (*e.g.*, magnitude, frequency, *etc.*), on a population's

resilience attributes of resistance, compensation, and time to recovery. Here, *resistance* is the

ability for a population to avoid a decline in size following a disturbance, whereas

compensation is the extent to which a population may increase in size following a

disturbance. Meanwhile, time of recovery (henceforth *recovery*) is the duration needed for a

population to converge back to its stationary equilibrium following a disturbance. Transient

demographic theory offers the opportunity to quantify the short-term dynamics of natural

populations, thus unlocking the potential for macroecological studies exploring patterns and

plausible mechanisms of demographic resilience. Transient increases in population size (N)

can be evaluated using metrics of population reactivity ($\overline{\rho}$; increase in N within a one-time

step following a disturbance) and maximal amplification ($\overline{\rho}_{max}$; maximum increase in N

during the transient period). Equally, the magnitude of transient declines in population size

can be assessed using the metrics of first-step attenuation ($\underline{\rho}$; decrease in N within a one-time

step following a disturbance) and maximal attenuation ($\underline{\rho}_{max}$; maximum decrease in N during

transient period). Finally, the damping ratio (ρ; rate of convergence back to stationary

stability), and period of oscillation (ψ; time between corresponding phases of the largest

oscillatory cycle in N) of populations offer insights into their capacity to go back to a

stationary stable equilibrium. **(B)** We quantified the aforementioned transient metrics of

demographic resilience for 2,242 populations from 369 species for whom Matrix Population

Models (MPMs) are available from the global COMPADRE[25] (Blue; plants & algae) and

COMADRE[24] databases (Orange; animals). Point size illustrates the relative length of time

over which each population was surveyed to parameterise the MPMs deposited within

COMPADRE and COMADRE.

Figure 2. Variation across metrics of demographic resilience: resistance (green),

compensation (blue), and recovery (orange) does not correspond with patterns in the

exposure of populations to environmental stochasticity. Scores and loadings of a

phylogenetically-weighted Partial Least Squares (pPLS) regression analysis exploring the

correlation between patterns in the variation of the six transient metrics of **(A)** first-step

attenuation ($\underline{\rho}$), **(B)** maximal attenuation ($\underline{\rho}_{max}$), **(C)** reactivity ($\overline{\rho}$), **(D)** maximal

amplification ($\overline{\rho}_{max}$), **(E)** damping ratio (ρ), and **(F)** period of oscillation (ψ), and the five

metrics of environmental stochasticity: temperature frequency spectrum (β_T), temperature

autocorrelation (a_T), thermal range/magnitude (m), precipitation frequency spectrum (β_P), and

precipitation autocorrelation (a_P). The component scores along each axis display the

percentage variance in the environmental stochasticity variables captured by each component,

with the first two components alone explaining >50% of the variance across all models. The

gradation in point colour then reflects patterns in the relative magnitude of each transient

metric recorded from each population, with darker shades indicating higher estimates. Insert

barplots are the standardised regression coefficients (b) highlighting the relative weighting of

each abiotic variable in the overall capacity of each pPLS model to explain variation in a

given transient metric (r^2).

Figure 3. The resilience attributes of resistance (green), compensation (blue), and recovery (orange), in natural populations are determined by the relative energetic investments of their individuals. Scores and loadings of a phylogenetically weighted Partial Least Squares regression analysis exploring the sensitivity patterns of the six transient metrics of **(A)** first-step attenuation ($\underline{\rho}$), **(B)** maximal attenuation ($\underline{\rho}_{max}$), **(C)** reactivity ($\overline{\rho}$), **(D)** maximal amplification ($\overline{\rho}_{max}$), **(E)** damping ratio (ρ), and **(F)** period of oscillation (ψ), towards the vital rates of survival (σ), progression (γ), retrogression (τ), and reproduction ($\square$). The component scores along each axis display the percentage variance in the environmental stochasticity variables captured by each component, with the first two components alone explaining >47% of the variance across all models. The gradation in point colour then reflects patterns in the relative magnitude of each transient metric recorded from each population, with darker shades indicating higher estimates. Insert barplots are the standardised regression coefficients (*b*) highlighting the relative weighting of each vital rate in the overall capacity of each pPLS model to explain variation in a given transient metric (r^2).

     Figure 1

 Figure 2.

Figure 3.