Deficiency in pericyte remodeling as a basis for impaired capillary flow and structure during brain aging

Andrée-Anne Berthiaume[1,2], Franca Schmid[3,4#], Stefan Stamenkovic[1#], Vanessa Coelho-Santos[1#], Cara D. Nielson[1,5], Bruno Weber[4,6], Mark W. Majesky[1,7], and Andy Y. Shih[1,2,8,9,*]

[1] Center for Developmental Biology and Regenerative Medicine, Seattle Children's Research Institute, Seattle, WA, USA.
[2] Department of Neuroscience, Medical University of South Carolina, Charleston, SC, USA.
[3] Institute of Fluid Dynamics, ETH Zurich, Sonneggstrasse 3, Zurich, Switzerland.
[4] Institute of Pharmacology and Toxicology, University of Zurich, Winterthurerstrasse 190, Zurich, Switzerland.
[5] Graduate Program in Neuroscience, University of Washington, Seattle, WA, USA.
[6] Neuroscience Center Zurich, University and ETH Zurich, Winterthurerstrasse 190, Zurich, Switzerland.
[7] Department of Pathology, University of Washington, Seattle, WA, USA.
[8] Department of Pediatrics, University of Washington, Seattle, WA, USA.
[9] Department of Bioengineering, University of Washington, Seattle, WA, USA.

[#]These authors share second authorship
*Corresponding author

Abbreviated title: Pericyte remodeling diminished in aging brain

Abstract: 150 words
Narrative: 4500 words (Introduction, Results, Discussion)
References: 56 (15 online methods)
Figures: 9 main (all color); 13 supplementary
Tables   2 supplementary

Abbreviations:
AD   Alzheimer's disease
BBB   Blood-brain barrier
RBC   Red blood cell
VCID   Vascular contributions to cognitive impairment and dementia
vSMC   Venular smooth muscle cell

Correspondence: Andy Y. Shih
                        Center for Developmental Biology and Regenerative Medicine
                        Seattle Children's Research Institute
                        1900 9th Ave. M/S JMB-5
                        Seattle, WA 98101
                        Office: 206-884-1314
                        Fax: 206-884-1407
                        Email: Andy.Shih@SeattleChildrens.org

Deterioration of brain capillary flow and architecture is a hallmark of aging and dementia. It remains unclear how loss of brain pericytes in these conditions contributes to capillary dysfunction. We conducted cause-and-effect studies by optically ablating pericytes in adult and aged mice *in vivo*. Focal pericyte loss induced capillary dilation without blood-brain barrier disruption. These abnormal dilations were exacerbated in the aged brain, and resulted in increased flow heterogeneity in capillary networks. A subset of affected capillaries reduced in perfusion due to flow steal. Some capillaries stalled in flow and regressed, leading to loss of capillary connectivity. Remodeling of neighboring pericytes restored endothelial coverage and vascular tone within days. Pericyte remodeling was slower in the aged brain, resulting in regions of persistent capillary dilation. These findings link pericyte loss to disruption of capillary flow and structure. They also identify pericyte remodeling as a therapeutic target to preserve capillary flow dynamics.

INTRODUCTION

Alterations in brain capillary architecture commonly occur with aging. Loss of capillary density, increased tortuosity and twisting of vessels, formation of non-patent string capillaries, and abnormal lumen diameter are consistent observations across decades of neuropathological research on the aged brain.[1] When these defects accumulate in the capillary network, they hinder the distribution of oxygen carrying blood cells, leading to tissue hypometabolism and heightened sensitivity to disease processes. In Vascular Contributions to Cognitive Impairment and Dementia (VCID) and Alzheimer's disease (AD), these capillary defects are seen at higher frequencies and contribute to neurodegeneration.[1] Despite a wealth of knowledge on their association with cognitive decline, the etiology of age-related capillary changes remain poorly understood, due in part to the cross-sectional nature of human autopsy studies and limited resolution of *in vivo* clinical imaging. Lacking mechanistic insight, therapeutic access points to mitigate age-related capillary insufficiency are missing.

Pericytes are mural cells that line the capillary beds of the brain.[2] They serve essential roles in the formation and maturation of blood vessels[3], establishment of blood-brain barrier (BBB) integrity[4, 5], and regulation of capillary diameter and blood flow[6, 7]. Recent studies suggest that pericytes are also sensitive to neuronal activity and participate in neurovascular coupling.[8, 9] Pericytes form a near-continuous cellular chain along the entire capillary network in healthy microvasculature, leaving little of the capillary bed uncontacted by their extensive processes. Each pericyte occupies a distinct territory, which does not overlap with the territories of its immediate neighbors.[10, 11] Mounting evidence indicates that pericytes are hyper-sensitive to cerebrovascular pathology, and loss of pericyte number and endothelial coverage occurs in

normal aging[12] and is accelerated in both VCID and AD.[13-17] Pericyte loss has been predominantly linked to deficits in BBB integrity.[13, 17, 18] However, it is unclear how this loss is related to deficiencies in capillary structure and perfusion in the aged and diseased brain.[19]

In prior work, we demonstrated the necessity of pericyte coverage to maintain capillary tone and flow resistance *in vivo*.[6] Optical ablation of individual capillary pericytes led to aberrant dilation of uncovered capillaries and increased blood flow. This pericyte loss also triggered the growth of processes from neighboring pericytes to re-establish endothelial contact and restore vascular tone, revealing an intrinsic repair mechanism.[10] These studies raised several questions. First, the consequence of focal pericyte loss on broader capillary flow dynamics remained unknown. Second, it was unclear if structural remodeling was the primary means to restoring endothelial coverage with greater degrees of pericyte loss. Third, the capacity for pericyte remodeling was not examined in the aging brain, the context in which pericyte remodeling is most important. Here, we address these questions in brain capillary networks of healthy adult and aged mice using *in vivo* two-photon single cell ablation and longitudinal imaging.

RESULTS

No difference in capillary landscape in upper cortex of adult and aged mice.

We compared capillary structure and pericyte abundance at baseline between adult and aged PDGFRβ-tdTomato mice using *in vivo* two-photon imaging (**Fig. 1a,b, Supplementary Fig. 1a,b**). Our analyses focused on capillary networks within the first 10-100 μm from the cortical pial surface, as subsequent pericyte ablation studies were limited to these depths. Capillary networks of aged mice (18-24 months) did not differ in total length, number of junctions, or abundance of pericyte somata when compared to adult mice (3-6 months)(**Supplementary Fig. 1c, d, e**). Further, the limited capillary regions that lacked pericyte coverage did not differ between ages (**Supplementary Fig. 1f**). Thus, capillary structure and pericyte coverage in upper layers of cortex is comparable at baseline between adult and aged mice.

Acute consequence of triple pericyte ablation.

In vivo optical ablation of capillary pericytes allowed us to observe the response to focal pericyte loss. We ablated three contiguous pericytes from capillary regions in upper layers of cortex in adult and aged mice (**Fig 1c-f**). Across both age groups, triple pericyte ablations uncovered capillary lengths of 643.75 ± 26.43 μm (mean ± SEM; 475-809 μm range)(**Fig. 1d,g**), which encompassed ~15 total capillary segments, *i.e.*, sections between branch-points (**Fig. 1h**). The

affected individual capillary segments varied greatly in length (4.79 to 236.6 µm)(**Fig. 1i**). Triple pericyte ablation resulted in loss of pericyte-pericyte contact at 7-10 process termini from an average of 8 neighboring pericytes, each with potential to grow into the uncovered territory (**Fig. 1d,j,k**). We detected no difference in these basal vascular or pericyte metrics between adult and aged mice.

Pericyte remodeling is efficient in the adult mouse brain, but impaired with aging

To measure pericyte remodeling, we re-examined capillary regions with pericyte loss at 3, 7, 14, and 21 days post-ablation **(Fig. 2)**. In adult mice, neighboring pericytes remodeled their processes to achieve contact with nearly the entire uncovered region within 21 days (**Fig. 2a,b**). In a representative example, endothelial contact was regained through simultaneous growth of 10 neighboring pericytes, each extending a single process toward the uncovered region **(Fig. 2c)**. Remodeling led to eventual contact with another pericyte process terminus, *i.e.*, pericyte-pericyte contact, upon which no further growth was possible (**Fig. 2d**).[10] In the aged group, a similar number of remodeling processes contributed to endothelial re-coverage, but a greater amount of capillary length remained uncovered at 21 days (**Fig 2e-h**).

In the adult brain, 84% of processes had made pericyte-pericyte contact within the 21-day timeframe (**Fig. 3a,b**). In the aged brain, less than 40% of processes had made contact. To assess the dynamics of pericyte process growth, we measured process length and growth rate for each interval of imaging, *i.e.*, day 3 to 7 (**Fig 3c,d**). "Maximum growth rate" was the time interval with the fastest growth, whereas "average growth rate" was the mean rate across all time intervals before pericyte-pericyte contact was achieved. We observed a significant decrease in both maximum and average growth rates in aged mice (**Fig. 3e,f**). After 21 days, however, the final length of extension achieved and total process length (from soma to process terminus) was not statistically different between age groups (**Fig. 3g,h**). An age-related effect in these metrics may be masked by the high variance of remodeling between pericytes, and because pericytes were studied broadly across the arterio-venous axis, as will be discussed below.

To better understand the outcome of synergistic remodeling, we calculated the length of endothelium uncovered over time. In adult mice, uncovered capillary length declined steeply in the first 7 days following ablation, and then slowly approached zero over the next 14 days (**Fig. 3i, left**). The aged group displayed a slower decline of uncovered territory, with a portion of the endothelium still left uncovered after 21 days (**Fig. 3i, right**). In a subset of experiments in aged mice, ablation areas were re-visited 49 days post-ablation and sizeable stretches of capillary remained uncovered (**Fig. 3j, Supplementary Fig. 2**). Together, these data show that the

interconnected structure of the capillary network facilitates restoration of endothelial coverage, but the capacity for pericyte remodeling is reduced with aging.

Since tdTomato expression was driven under constitutive *pdgfrb* promotor activity, new pericytes arising from proliferation or migration would be fluorescently labeled. However, we did not observe the emergence of new pericyte somata in either age group, suggesting that remodeling of existing pericytes is the main reparative strategy at this scale of pericyte loss.

Contributors to the heterogeneity of pericyte process growth.
We explored what parameters might contribute to variability in pericyte process growth. Many remodeling processes met one or more capillary bifurcations as they entered the uncovered territory, when the processes would split and extended in both directions, unless the terminus of another pericyte impeded its growth (**Supplementary Fig. 3a**). A single process could split up to three times, resulting in four growing termini. Given this parallel growth, processes that split had significantly greater final extension lengths than those that did not (**Supplementary Fig. 3b**). In adult mice, 49% of processes split, as opposed to 35% in aged mice, a difference attributed to their slower growth rather than differences in the number of capillary bifurcations available (**Supplementary Fig. 1d)**. There was no difference in the number of bifurcations within uncovered regions between age groups (7.3 ± 2.7 and 7.9 ± 2.9 bifurcations uncovered in adult and aged mice, respectively; p=0.4035, t-test; mean ± SEM).

We further examined if the maximum process length achieved was limited by baseline process length. Baseline lengths did not differ between age groups, consistent with normal pericyte density at baseline (**Supplementary Fig. 3c**). We therefore pooled age groups and plotted baseline process length as a function of length extended post-ablation. This revealed a negative correlation suggesting limited growth potential in individual processes longer than ~150 μm (**Supplementary Fig. 3d**). It also showed that short basal processes that split during growth had the greatest final added length.

In both age groups, ~25% growing pericyte processes created small vessel distortions ("microbends") when they tethered to a point further along the capillary and pulled the anchored point closer (**Supplementary Fig. 4**). Microbends were transient and capillary shape returned to normal once the process terminus had extended further. Processes that created microbends grew at a faster rate and gained more contact distance (**Supplementary Fig. 4e,f**). Thus, both intrinsic pericyte properties (baseline process length, ability to distort vessel) and extrinsic factors (capillary branching) can influence pericyte growth potential.

Mural cells across microvascular zones can contribute to capillary re-coverage

The cortical microvasculature is divided into arteriole-to-capillary transition, capillary, and venular zones, across which mural cells vary in their transcriptional profile, morphology and contractile dynamics.[20, 21] Although pericytes were exclusively ablated in the capillary zone, they were often adjacent to arteriole-capillary transition and post-capillary zones, allowing us to assess remodeling of mural cells from these regions into the capillary zone (**Fig. 4a**).

Mural cells from all zones were able to extend processes into the uncovered capillary territory. Interestingly, regardless of the morphological characteristics of their existing processes, mural cells extended into the capillary zone with a "thin-strand" morphology typical of capillary pericytes (**Fig 4b-d, Supplementary Fig. 5a-d**). Striking examples are ensheathing pericytes and venular smooth muscle cells (vSMCs), whose pre-existing processes were more complex (**Fig. 4b,d**) This suggests that pericyte process morphology is dictated by the endothelial zone it contacts. In further support of this concept, we also ablated ensheathing pericytes and found that remodeling neighbors retained their circumferentially-oriented processes when extending within the arteriole-capillary transition zone (**Supplementary Fig. 5e,f**).

We found no difference in rate of process growth across microvascular zones in adult mice (**Fig. 4e**). The aged group exhibited selective impairment in remodeling of ensheathing and capillary pericytes (mesh and thin-strand morphology) of the arteriole-capillary transition and capillary zones, respectively. In contrast, vSMCs on ascending venules maintained their remodeling capacity (**Fig. 4e, Supplementary Fig. 5c,d**). Comparing maximal process extension between adult and aged mice for each vascular zone yielded a similar outcome (**Fig. 4f**). This points to vSMCs as a retained source of repair capacity within the aged brain and suggests that arteriole-capillary transition and capillary zones may be more susceptible to prolonged loss of pericyte coverage.

Overt BBB leakage is very rare following focal pericyte ablation.

Pericytes are well-established as custodians of BBB integrity.[5] In recent studies, BBB permeability occurring from adult-induced pericyte loss could be detected by extravasation of a 70 kDa dextran dye[22] and even larger IgG proteins[23]. In both adult and aged mice, we detected no extravasation of 70 kDa dye after focal loss of pericyte coverage loss (**Supplementary Fig. 6a,b**). However, in a rare occurrence in aged mice (1 out of 7 triple ablation experiments; 1 of 98 uncovered capillary segments inspected) leakage was observed at days 3 and 7 post-ablation (**Supplementary Fig. 6c**). We also assessed a low molecular weight dextran dye in aged mice (10 kDa FITC-dextran),

which stays within the blood plasma for several minutes following a single i.v. bolus. No increase in dye extravasation was seen after pericyte loss (3 days post-ablation), compared to regions receiving off-target sham irradiations where the ablative laser path was placed away from a pericyte soma but of similar distance to the vessel wall (**Fig. 5a-d**).

We considered the possibility that BBB leakage was too slow or subtle to be detected by imaging of dye extravasation. Since microglia are highly sensitive to vascular pathology[24, 25], cluster around leaky vessels[26], and react to BBB leakage in models with more extensive pericyte loss[22], we reasoned that they would be sensitive indicators of any BBB leakage. In adult and aged double transgenic mice with co-labeled mural cells and microglia, microglia reacted within minutes to engulf the soma of the ablated pericyte (**Fig. 5e,g,i**). Critically, this reaction was wholly focused on the soma of the dying pericyte, and not the broader capillary regions previously contacted by their processes. Following off-target sham irradiations, microglial processes occasionally extended to investigate the area in the 30 minutes following laser exposure, but this reaction was comparatively mild (**Fig. 5f,h,j**). Three days post-ablation or off-target irradiation, we observed no aggregation of microglia or their processes along peri-vascular regions suggesting no delayed BBB leakage.

We further examined endothelial cell junctions after focal pericyte ablation. Using Claudin5-eGFP fusion protein mice, prior studies showed the development of protrusions and small gaps within tight junction strands during stroke-induced BBB leakage, indicative of endothelial remodeling and BBB damage, respectively.[27] After pericyte ablation in adult mice with co-labeled pericytes and Claudin-eGFP, we observed a transient but non-significant increase in small protrusions, and no increase in tight junction gaps (**Supplementary Fig. 7**). We also used adeno-associated virus to label astrocytes in aged mice. No overt changes in astrocyte endoot apposition to the capillary wall following focal pericyte ablation were seen (**Supplementary Fig. 8**). Thus, the degree of pericyte loss in our paradigm does not reach the threshold needed to cause overt BBB disruption, except with one rare observation in an aged mouse.

Aberrant capillary dilation with loss of pericyte coverage is exacerbated with aging

In prior studies, we reported that loss of pericyte coverage led to abnormal capillary dilation in adult mice, indicating a role in regulation of basal capillary tone.[6, 10] We compared this effect between adult and aged mice with triple pericyte ablations. At baseline, we detected no difference in the diameter of capillaries across age groups (**Fig. 6a**). For each ablation experiment, we measured the diameters of 4-5 capillaries at the following time-points: *i*) Pre-ablation, when the capillary was covered, *ii*) 3 days post-ablation, when the endothelium was uncovered, and *iii*) 14-

21 days post-ablation, when the endothelium was re-covered by process growth from neighboring pericytes (**Fig. 6b**). In adult mice, we observed average dilations of 0.6 μm (21% increase from baseline) when the capillary was uncovered (**Fig. 6c,d,g**). Dilation occurred exclusively where pericyte contact was lacking (**Supplementary Fig. 9**). With coverage regained by pericyte remodeling, capillary diameters returned to levels no different from baseline (**Fig. 6h**).

In aged mice, uncovered capillary segments dilated to a significantly greater extent than in adult mice, *i.e.,* an average of 1.1 μm (36% increase from baseline), with some individual capillaries dilating as much as ~1.6 μm (**Fig. 6e-g**). Further, slight dilations persisted even after pericyte coverage had returned (**Fig. 6h**), suggesting inefficient re-establishment of vascular tone. Capillaries that continued to lack coverage beyond 21 days post-ablation maintained their dilated states (**Fig. 6i**). We confirmed that the dilations were not due to the laser damage alone, as off-target sham irradiations led to no change in capillary diameter (**Supplementary Fig. 10**). Further, these results were not an effect of isoflurane anesthesia, as dilations persisted when mice were imaged in the awake state (**Supplementary Fig. 11**). Altogether, this indicates that pericyte contact is key to maintaining basal capillary tone *in vivo*, and dilation with pericyte loss is exacerbated in the aged brain.

Local dilations alter flow distribution and increase flow heterogeneity in capillary networks
The flow and oxygenation of red blood cells (RBCs) in brain capillaries is heterogeneous at rest.[6, 28] Homogenization of blood flow among capillaries is necessary to efficiently extract and distribute oxygen (O_2) to the tissue.[29-32] Therefore, increased flow heterogeneity is a barrier to achieving flux homogenization and O_2 extraction. To examine whether focal capillary dilations increased flow heterogeneity, we performed triple pericyte ablations (**Fig. 7a**) and used line-scans to measure blood cell flux (cells/s) and flow directionality in capillaries both within and immediately surrounding the uncovered region (**Fig. 7b,c**). Flux was measured in the same capillaries before and 3 days post-ablation, when pericyte remodeling had not yet restored coverage and tone.

In an aged mouse, the already high variance of basal flux among capillaries, 18 to 231 cells/s, further broadened to 37 to 453 cells/s post-ablation (**Fig. 7d**). The increased flux was predominantly observed in dilated capillary segments lacking pericyte coverage (**Fig. 7b,e,f**). However, a fraction of sampled capillaries surrounding these segments decreased in flux (**Fig. 7b,e,g**). When pericyte contact was lost in only one of the daughter branches of a divergent bifurcation, blood cells flowed preferentially through the dilated branch. This left the alternate branch under-perfused relative to baseline, even when overall flow entering the bifurcation was

higher post-ablation (**Fig. 7h-j**). This non-uniform partitioning of blood cell flow at capillary bifurcations reflects the Zweifach-Fung effect[33], where increased flow rate in a dilated capillary pulls a larger fraction of RBCs into the dilated daughter branch. These local upstream changes can have propagating effects within the capillary network[34], as we also observed decreased flow at downstream, convergent bifurcations (**Fig. 7k-m**).

Focal capillary dilation increases flow heterogeneity in silico.

To understand capillary flow changes on a broader network level, we performed blood flow simulations in realistic microvascular networks derived from mouse parietal cortex (**Fig. 8a**).[35] Four cases were studied in two microvascular networks, each involving dilation of 13-14 contiguous capillary segments with similar characteristics to those targeted during triple pericyte ablations *in vivo* (**Fig. 8b**)(**Supplementary Table 1**). The capillaries were each dilated by 0.6 µm or 1.1 µm from their baseline diameters to mimic average dilations measured in the adult and aging brain, respectively (**Fig. 6g**). An additional group with 1.6 µm dilation was included to examine the effect of the most extreme dilations observed in aged mice. Consistent with observations *in vivo*, focal dilations produced both increases and decreases in capillary flow (defined as >10% change from baseline)(**Fig. 8c**). Flow perturbations were detected in hundreds of capillaries surrounding the dilated region. The number of affected capillaries increased commensurately with the extent of dilation, with ~1/3 of the affected capillaries decreasing in flow irrespective of the extent of dilation. Capillaries with increased flow localized generally to the epicenter of dilation, while those with decreased flow resided in surrounding areas (**Fig. 8d**).

To understand how capillary architecture related to blood flow change, we categorized capillaries into three groups: Dilated, Gen1-Neighbors (1 branch from dilated), and Gen2-Neighbors (2 branches from a dilated segment). Consistent with *in vivo* observations, dilated capillaries generally increased in flow, while Gen1 and Gen2 neighbors showed both increases and decreases from baseline (**Fig. 8e**). Further, capillary steal was observed at divergent capillary bifurcations, where one daughter branch was selectively dilated and the undilated branch experienced decreased flow (**Fig. 8f**). As predicted by the Zweifach-Fung effect[33], the steal effect is even more pronounced with RBC flux (**Supplementary Figure 12a**). Again consistent with *in vivo* data, the variance in absolute capillary flow increased substantially with focal capillary dilation (**Supplementary Figure 12b**). To assess whether capillary flow heterogeneity deviated from a normal range, we defined the expected baseline flow heterogeneity among a multitude of undilated capillaries within the two microvascular networks (**Fig. 8g**). This revealed that capillary

dilation in the adult brain led to abnormally elevated regional flow heterogeneity, and that heterogeneity further increases with magnitudes of dilation seen in the aged brain (**Fig. 8h**).

Local dilations produce blood flow stalls and capillary regression in vivo

In surveying volumetric data collected across all *in vivo* triple ablation experiments, we found a higher frequency of blood flow stalls in experiments involving pericyte ablation, compared to off-target irradiations (**Fig. 9a-d, Supplementary Fig. 13**). These stalls appeared as vessels with no moving blood cell shadows within the dye-labeled plasma, and could be detected in images without line-scanning.[36] Stalls occurred with greatest likelihood one branch-point away from an uncovered capillary segment (**Fig. 9e**), suggesting blood steal by an adjacent dilated capillary. Many of the stalls occurred at divergent bifurcations, which occur closer to the arteriole-capillary transition zone (**Supplementary Fig. 13b**).

Most capillary stalls were transient, and presumably resolved by restoration of capillary tone in dilated neighbors. Transient stalls were defined as capillary segments with no blood flow on one day of imaging, but with flow observed in a subsequent imaging session. We could not determine the precise duration over which flow was stalled, due to limited sampling frequency. However, we suspect that some capillaries experienced prolonged loss of flow, as a subset of stalling events were later associated with regression of the non-flowing capillary segment (13% of total stalls). Regressions left behind a pericyte bridge without an endothelial lumen, reminiscent of a string capillary (**Fig. 9b,d,f, Supplementary Fig. 13c**). These outcomes were based on pooling data across adult and aged mice, as their frequency in focal ablation experiments was insufficient for age comparison. Additionally, we did not detect increases in stalled or low-flow capillaries in the *in silico* data (**Supplementary Figure 12c,d**). This is not surprising since biological processes such as blood cell adhesion to the endothelium and potential vasoconstrictive events were not captured in our modeling. Collectively, these data reveal how altered capillary flow distribution due to pericyte loss can lead to cessation of blood flow in capillary segments, and lead to enduring loss of capillary structure.

DISCUSSION

We have ablated pericytes focally in capillary networks of the adult and aged mouse brain to understand how pericyte loss affects capillary function *in vivo*, and to study pericyte remodeling in the context of brain aging. Our experiments show that restoration of endothelial coverage

following pericyte loss is a synergistic effort from multiple neighboring pericytes, leveraging highly interconnected capillary networks to create access points for remodeling pericytes. This suggests that pericyte remodeling would be less efficient in aging or disease, particularly in sparser capillary networks of white matter and hippocampus.[37-39] Resilience would be further diminished in the arteriole-capillary transition and capillary zones of the aged brain due to impaired pericyte growth.

Abnormal capillary dilations caused by pericyte loss lead to marked increases in local blood flow.[6] However, uneven dilation among capillaries results in steal of flow from capillary segments due to abnormal partitioning of blood cell flux at bifurcations. This finding is supported by concordant data from *in silico* and *in vivo* experiments showing reduction of flow in ~1/3 of affected capillaries, and increased flow heterogeneity. The brain's ability to homogenize flux among capillaries is critical for proper allocation of O_2 to brain tissue.[30, 31] Broadening of capillary flux heterogeneity by pericyte loss creates a larger barrier for flux homogenization, limiting tissue oxygenation at rest and during functional hyperemia. Indeed, greater capillary flow heterogeneity has been associated with age-related cerebrovascular diseases underlying stroke and dementia.[40, 41] In future studies, it will be important to examine the consequence of these capillary abnormalities on tissue oxygen content.[42]

Pericyte loss also led to increased blood flow stalling. Stalls were observed in more than half of our experiments, and we estimate that an average of 3%, and maximum of 9%, of capillaries in the region (including dilated vessels, Gen1 and Gen2 neighbors) experienced a stall in flow. Flow stalls are typically seen in only 0.4% of brain capillaries of healthy adult mice, and an increase to 1.8% in a mouse model of Alzheimer's disease was associated with worsened cognitive performance.[36] In agreement with this idea, a recent study using an adult-induced genetic pericyte depletion model also observed increased capillary stalling.[43] This effect was attributed to enhanced leukocyte-endothelial interaction, but may also involve flow steal in capillary networks.

Capillary networks of adult mouse cortex are also remarkably stable in structure.[44] Our observation of capillary regression with ablations, but not shams, suggests that flow disruptions due to pericyte loss can promote enduring reductions in capillary connectivity. Collectively, these data establish a link between pericyte loss and cerebral blood flow deficiency, wherein aberrant network flow dynamics are a key facet of the impairment. The effects we report here are based on a restricted scale of pericyte loss, and we expect capillary defects to compound with the severity of pericyte loss reported in AD (30-60% loss of total pericyte numbers).[13, 16] Further, we suspect that string capillaries and overall reduced capillary density seen in aging and dementia

may result, in part, from capillary flow abnormalities induced by pericyte loss.[19, 45]

Our findings help to explain previously observed changes in capillary structure and function with brain aging. While we did not detect basal differences in capillary density or pericyte abundance in superficial somatosensory cortex, broader examinations by other groups have reported lower capillary density in cortical gray and white matter[37, 38, 46], and reduction in both pericyte number and coverage[12] in the aged rodent brain. Two-photon imaging studies in aged rats[47] and mice[48] have revealed average increases in capillary diameter and RBC velocity, which in combination with reduced capillary density and hematocrit, become associated with microscopic pockets of under-perfused tissues.[48] Recent microvascular angiography studies in the aged mouse brain have also revealed greater heterogeneity in flow velocity across cortical capillaries under basal conditions.[49] It will be important to establish how spontaneous, age-related pericyte loss affects capillary perfusion and structure, particularly in the white matter where capillary rarefaction in aging is widely observed.

Studies using global pericyte depletion have shown marked decreases in neurovascular coupling responses[50, 51], BBB disruption and circulatory failure due to vasogenic edema.[23] Our studies provide a clearer view on how pericyte loss affects capillary flow, since the focal ablation approach affects blood flow without disrupting the BBB or inducing overt vascular pathology. However, the lack of BBB leakage in our model does not contradict prior work showing BBB leakage with more extensive pericyte loss.[22, 23] It is likely that pericytes use paracrine signaling pathways to ensure overall endothelial health, and ablation of a few isolated pericytes does not compromise overall barrier function.

PDGF-B/PDGFRβ signaling is involved in pericyte coverage during adulthood since genetic ablation of endothelial PDGF-BB in adult mice leads to age-dependent pericyte loss and BBB defects.[22] Augmenting PDGF-B/PDGFRβ activity in adulthood may therefore be a strategy to increase pericyte coverage and improve capillary homeostasis. *In vitro,* administration of PDGF-BB leads to proliferation of human brain pericytes and protection from apoptosis.[52] Intravenous treatment with PDGF-BB peptide was sufficient to promote pericyte coverage and restore capillary tone and flow rate in a model of epilepsy involving mural cell loss.[53] Interestingly, the immunomodulatory drug, thalidomide, can increase expression of PDGF-BB from endothelial cells, promoting pericyte coverage and reducing bleeding in models exhibiting vascular pathology in the CNS[54] and peripheral tissues.[55, 56]. While deeper mechanistic studies are needed, these studies form the logic to develop therapeutics that can promote pericyte coverage and preserve normal capillary flow dynamics in age-related cerebrovascular diseases.

METHODS

Animals. PDGFRβ-tdTomato mice[57] were bred from Pdgfrb-Cre[58] and Ai14[59] mice and aged in-house for the study. The adult mouse group (n = 6; 3 female) included animals between 3 to 6 months of age. The aged mouse group (n = 6; 3 female) included animals between 18 and 24 months of age. Age group brackets were determined by the Jax "Life Span as a Biomarker" criteria, which defines 3-6 months as "mature adult" stage, and 18-24 months as "old" stage in mice. Aged mice were visually checked for general health and any overt signs of illness such as tumors or weight loss resulted in exclusion from the study. For a subset of experiments, we crossed PDGFRβ-tdTomato mice with CX3CR1-GFP[60, 61] or Claudin-eGFP fusion protein[27] mice to simultaneously label mural cells alongside microglia or endothelial tight junctions, respectively. PDGFRβ-tdTomato;CX3CR1 mice (n=5; 2 female) were aged to 3-6 months (adult) and 18-24 months (aged) before the start of experiments. PDGFRβ-tdTomato;Claudin-eGFP mice (n=3; 1 female) were aged to 7-12 months. The blood cell flux example (**Fig. 7**) is from a 19-month-old male PDGFRβ-tdTomato mouse. The Institutional Animal Care and Use Committee at the Seattle Children's Research Institute approved all procedures used in this study.

Surgery. Chronic cranial windows (skull removed, dura intact) were implanted in the skulls of all mice for triple pericyte ablation experiments. Briefly, surgical plane anesthesia was induced with a cocktail consisting of fentanyl citrate (0.05 mg/kg), midazolam (5 mg/kg) and dexmedetomidine hydrochloride (0.5 mg/kg)(all from Patterson Veterinary). Dexamethasone (40 µL; Patterson Veterinary) was given 4-6 hours prior to surgery to reduce brain swelling during the craniotomy. Cicular craniotomies ~3 mm in diameter were generated under sterile conditions, and sealed with a glass coverslip consisting of a round 3 mm glass coverslip (Warner Instruments; 64-0720 (CS-3R)) glued to a round 4 mm coverslip (Warner Instruments; 64-0724 (CS-4R)) with UV-cured optical glue (Norland Products; 7110). The coverslip was positioned with the 3 mm side placed directly over the craniotomy, while the 4 mm coverslip laid on the skull surface at the edges of the craniotomy. An instant adhesive (Loctite Instant Adhesive 495) was carefully dispensed along the edge of the 4 mm coverslip to secure it to the skull. Lastly, the area around the cranial window was sealed with dental cement. This two-coverslip "plug" fits precisely into the craniotomy and helps to inhibit skull regrowth, thereby preserving the optical clarity of the window over months. Mice recovered for a minimum of 3 weeks following surgery.

In a subset of experiments investigating the response of microglia to the pericyte ablation, thinned-skull windows (PoRTs window technique) were used to minimize disruption of the brain and best preserve resting state microglia.[62] This window involved using a dental drill to thin the

skull for optical penetrance, but without breaching the calvaria. These windows were imaged one day following surgery for a maximum of three days.

Two-photon imaging. *In vivo* two-photon imaging system was performed with a Bruker Investigator (run by Prairie View 5.5 software) coupled to a Spectra-Physics Insight X3. Green, red and far red fluorescence emission was collected through 515/30 nm, 615/60 nm, 700/75 nm bandpass filters, respectively, and detected by gallium arsenide phosphide photomultiplier tubes. Low-resolution maps of the cranial window were first collected for navigational purposes using a 4-X (0.16 NA) objective (Olympus; UPlanSAPO). We then switched to a 20-X (1.0 NA) water-immersion objective (Olympus; XLUMPLFLN) and used 975 nm for imaging of mural cells (tdTomato) and vasculature (FITC-dextran). For 3 channel imaging of tdTomato, GFP and Alexa 680-dextran we used 920 nm excitation. All imaging with the water-immersion lens was done with room temperature distilled water.

In the majority of studies, imaging was performed while the mice were anesthetized with 0.75-1.5% (MAC) isoflurane delivered in medical grade air through a custom nose cone. In a subset of studies, we imaged mice in the awake state, while head-fixed and free to run on a low-resistance treadmill equipped with a motion sensor. In these studies that compared capillary diameters between anesthetized and awake state, the mice were first imaged under isoflurane, and then isoflurane was stopped and the mouse was allowed to breathed medical air for a minimum of 15 minutes. Awake status of the mouse was evident by movement on the treadmill, and imaging data was captured during periods of rest between movements.

Pericyte ablation. Pericyte ablations were conducted with focused two-photon line-scan to specifically target individual pericyte somata.[10, 63] This approach is effective only for pericyte ablation in the upper ~200 μm of cerebral cortex. The ablative scan was performed at 725 nm. Ablative line-scans were applied in 20-30 second increments, for a total of 60-90 seconds per soma. The power ranged from 30 to 50 mW and was adjusted on a case-by-case basis taking into account the depth of the soma from the pial surface. To limit power input required for successful pericyte ablation, all ablations were done within the first 100 μm of cortical depth beneath the pial surface (adult mice, 32.76 ± 15.66 μm; aged mice, 40.33 ± 16.51 μm from the pial surface. $p=0.1352$ by unpaired t test for n=21 pericytes from 6 adult mice, 21 pericytes from 6 aged mice). A successful ablation was confirmed on subsequent imaging days, if the targeted pericyte did not regain fluorescence. This indicated that cell death was more likely than temporary photobleaching. Fluorescence recovery after photobleaching can occur within seconds to minutes after light irradiation, but loss of pericyte fluorescence caused by our ablations was permanent.

Even with successful ablation, a dim imprint of the soma may remain for several days, as the dead cellular material is slowly cleared away. In unsuccessful ablations, the pericyte re-gains fluorescence over time in its soma and processes, as protein synthesis machinery replenishes cytosolic tdTomato. These cases, as well as cases where the ablation caused an immediate breach in the capillary wall, were excluded from the study.

For triple pericyte ablations, the day of ablation was considered "Day 0," and took place a minimum of 3 weeks after implantation of cranial window. Images of the area of interest were captured "pre-ablation" as well as "5 minutes post-ablation" on Day 0. The ablation regions were revisited and imaged at post-ablation days 3, 7, 14, and 21. At each imaging time point, a 70kDa FITC-dextran i.v. dye was injected to simultaneously label the vasculature and test BBB permeability.

Mural cells of differing microvascular zones. These studies relied on mural cell morphology and vascular architecture in mouse cerebral cortex to define microvascular zones, as previously described.[6, 20, 21] We considered penetrating arterioles as branch order zero, and the 1st to 3rd branch order of penetrating arteriole offshoots as the "arteriole-capillary transition zone", which we previously called the "pre-capillary arteriole". We expected a greater likelihood of ensheathing pericytes in the transition zone, which are known to express high levels of α-SMA, and have protruding ovoid cell bodies and circumferential processes that completely cover the majority of the vessel wall. At 4th branch order and beyond, referred to as the "capillary zone", we expected occasional mesh pericytes and an abundance of thin-strand pericytes. This zone is α-SMA low/negative and thin-strand pericytes have fine processes that trail longitudinally on the endothelium. Since the number of branch orders within the arteriole-capillary transition zone varies slightly across penetrating arteriole offshoots[21], it was possible that mesh pericytes, when present, fell into either "transitional" and "capillary" groups. To preserving objectivity and consistency, we relied on both vascular branch order and mural cell morphology to categorize zones. On the venule side, we considered the ascending venule itself as the "venule zone," or "V0," which are covered by stellate-shaped vSMCs.

10 kDa Dye extravasation imaging and analysis. Aged PDGFRβ-tdTomato mice (35-40 g in weight) were implanted with chronic, skull-removed cranial windows. Three weeks after window implantation, double or triple pericyte ablations and sham irradiations were performed. Three days later, a single 30 μL bolus of 10 kDa FITC-dextran (5% w/v in sterile PBS; Sigma, FD10S) was injected retro-orbitally. Immediately following injection, image stacks were collected continuously from the affected area with an interval of 2 minutes over a period of 18 minutes. Fluorescence

intensity over time was measured from ROIs using ImageJ/Fiji. Rates of parenchymal fluorescence intensity change were calculated using linear regression fitting over the first 10 minutes of data in MATLAB software.

Microglia experiments. We bred PDGFRβ-tdTomato;CX3CR1-eGFP[+/-] mice to co-label mural cells in red and microglia in green. Mice were implanted with thinned skull windows and imaged 1-2 days post implantation.[62] During imaging, Alexa 680-dextran (2 MDa, custom conjugated)[64] was injected retro-orbitally to label the vasculature and imaging was performed at 920 nm excitation for 3 channel imaging. We examined the reaction of microglia to both the on-target pericyte ablations and off-target sham irradiations in adult and aged mice. Ablations and irradiations were performed one day after surgery (Day 0). The volume was imaged before ablation/sham, and again 30 minutes and 3 days after. Microglial reactivity was quantified as GFP intensity at the site of pericyte soma ablation or irradiation (focal ROI), as well as along the capillary territory occupied by the pericyte (territory ROI).

Claudin-eGFP imaging and analysis. We bred PDGFRβ-tdTomato;Claudin5-eGFP mice to co-label mural cells in red and endothelial tight junctions in green. Mice were implanted with chronic, skull-removed cranial windows. During imaging, Alexa 680-dextran was injected retro-orbitally to label the vasculature and imaging was performed at 920 nm excitation. Images were captured prior to, and 5 minutes, 3 days, 7 days, and 14 days following on-target pericyte ablations or sham irradiations. The control group consisted of 2 sham irradiations and 1 failed pericyte ablation. Tight junction disassembly was analyzed by calculating the percentage of eGFP+ strands that contained gaps, and tight junction remodeling was analyzed by calculating the percentage of eGFP+ strands that contained protrusions or varicosities, following prior studies.[27] All data were normalized to pre-ablation. Only the tight junction strands on the vessel segment adjacent to the irradiation site were analyzed. Quantification was performed using image stacks in ImageJ/Fiji.

Astrocyte endfeet imaging and analysis. Aged PDGFRβ-tdTomato mice received intracortical adeno-associated virus (AAV) injections during chronic, skull removed window implantation. A Nanoject III (Drummond) was used to inject 50 nL of pAAV.GfaABC1D.PI.Lck-GFP.SV40 (105598-AAV5)[65] at ~ 100 µm below the cortical surface. Mice were allowed to incubate with the virus for 3-weeks before imaging. Alexa 680-dextran was injected retro-orbitally, and imaging was performed at 920 nm excitation. The apposition of astrocyte endfeet to the capillary wall was examined before and 3 days after pericyte ablation. Analysis of fluorescence intensity profiles was performed using ImageJ/Fiji.

Quantification of 3-D vascular architecture. Totally capillary lengths and branchpoints were measured from 350 μm (x), 350 μm (y), and 90 μm (z; ~10-100 μm from the pial surface) image stacks collected from adult and aged mice using Imaris software.

Quantification of pericyte remodeling. Pericyte process length over time was measured in 3-D using the Simple Neurite Tracer plugin for ImageJ/Fiji. To measure baseline process length, the full span of processes from soma to terminus was measured. All growth over time was plotted as a change in process length from baseline. Maximum growth rate was the greatest change in process length, in micrometers per day, between two imaging days. Average growth rate was calculated as the average of all post-ablation imaging timepoints. For average growth rate, only rates before pericyte-pericyte contact were included in the calculation. Pericyte-pericyte contact indicated completion of pericyte process remodeling. This was defined as the terminus of a process being <10 μm from another pericyte process.

Quantification of capillary diameter. Capillary segment diameters were measured with the ImageJ/Fiji macro Vasometrics, which provides unbiased diameter measurements across a vessel segment of interest (full-width at half-max fluorescence intensity) and reports an average diameter.[66]

Acquisition and quantification of capillary flow data. Line-scans were performed with 2MDa FITC-dextran, at a wavelength of 800 nm. Line-scans were obtained for each capillary of interest with a single line transecting the center of the vessel. We calculated RBC flux by manually counting the number of blood cell streaks (over ~1.3 s of scan time) and normalizing all numbers to represent flux in a period of 1 second. The angle of the streaks in relation to the direction of the scan was used to determine RBC flow direction. Line-scans were presented to an independent rater in a blinded fashion (S.S.), to reduce bias.

Statistics for in vivo data. All statistical analyses were performed with GraphPad Prism version 8 software or MATLAB (R2018a). Details of specific tests can be found in the Figure legends. For all unpaired t-tests, F tests for equality of variance were performed. When statistically significant F tests were observed, unpaired t-test with Welch's correction for unequal variance was used.

Microvascular networks for in silico modeling. The vectorized microvascular networks used for blood flow modeling were acquired in a previous study by Blinder et al.[67] In these data, each vessel is represented by an edge with a given length and diameter. The edges are connected at bifurcation (graph vertices), such that a fully connected microvascular network is obtained.

Both microvascular networks used for in silico modeling in this work were acquired from the vibrissa primary sensory (vS1) cortex of C57/BL6 male mice. Microvascular network 1 (MVN1) and 2 (MVN2) were within a tissue volume of ~1.6 mm^3 and ~2.2 mm^3 and contained ~12,100 and ~19,300 vessels, respectively. The vessels are labeled as pial arteries, descending arterioles, capillaries, ascending venules and pial veins; 96% (MVN1) and 94% (MVN2) of all vessels were capillaries.

Blood flow modeling with discrete RBC tracking. The numerical model to simulate blood flow with discrete RBC tracking in realistic microvascular networks was previously described.[35] We briefly summarize the key aspects of the modeling approach here, but more detailed descriptions are available in prior studies.[35, 68] The modeling approach is based on the small Reynolds number (Re < 1.0 for all vessels) in the microvasculature. As such the flow is laminar and mostly even in the Stokes regime, which allows description of the flow in individual vessels by Poiseuille's law. The flow rate q_{ij} in vessel *ij* between node *i* and *j* is computed by:

$$\text{Equation (1)} \qquad q_{ij} = \frac{p_i - p_j}{R_{ij}^e} = \frac{\pi\, D_{ij}^4}{128\, L_{ij}\, \mu\, \mu_{rel}^e}\, (p_i - p_j)$$

where D_{ij} and L_{ij} are the vessel diameter and the length and p_i and p_j are the pressure at node *i* and *j*, respectively. μ is the dynamic plasma viscosity and μ_{rel}^e the relative effective viscosity, which accounts for the presence of RBCs and is computed as a function of local hematocrit and vessel diameter as described in Pries et al.[69]

To compute the local hematocrit we track individual RBCs through the MVNs. Hereby, we need to account for the Fahraeus effect (RBC velocity is larger than bulk flow velocity) and the phase separation (RBCs partition with a different ration than the bulk flow at divergent bifurcations).[69] The phase separation effect is also called Zweifach-Fung effect[33] and describes the phenomenon that at divergent bifurcations the daughter vessel with the larger flow rate receives a larger fraction of RBCs. At divergent bifurcations with a diameter >10 µm the phase separation is described by empirical equations.[69] At smaller diameters RBCs propagate in single file flow and consequently the RBC motion can be approximated by assuming that the RBCs follow the path of the largest pressure force.[33, 35] It is important to note that the RBC distribution and therewith the flow field fluctuate in time and that the RBC distribution impacts the local flow field.[68, 70] In this study, we use the time-averaged flow field to compare changes in response to locally altered capillary diameters. The time-averaged flow field is computed by averaging over 15.4 s. The RBC flux per vessel is computed from the product of the time-averaged flow rate and discharge hematocrit,

where the discharge hematocrit is a function of the local tube hematocrit and the vessel diameter.[69] The pressure boundary conditions and the inflow hematocrit of 0.3 are kept constant for all simulations. Further details on assigning suitable pressure boundary conditions are available in Schmid *et al.*[35]

Analyses of in silico data. The capillaries selected for dilation were selected to mimic the *in vivo* scenario as closely as possible. Capillaries chosen for dilation were in the upper 100 µm of cortex and were beyond 4 branch orders from the main branch of the penetrating arteriole (0th order), and beyond one branch from the ascending venule main branch. *In vivo,* between 12-15 capillaries were affected by ablating 3 contiguous pericytes, which corresponds to an uncovered capillary length of 688 ± 70 µm. The pre-ablation characteristics of these capillaries as measured *in vivo* were: 1) diameter: 3.04 ± 0.59 µm (range: 2.17-4.81 µm), 2) RBC flux: 114 ± 73 RBCs/s (range: 17-381 RBCs/s) and 3) RBC velocity: 1.11 ± 0.63 mm/s (range: 0.19-2.70 mm/s).

We used these pre-ablation characteristics to choose representative "base capillaries" around which dilations seen after *in vivo* pericyte ablation was mimicked. To avoid being too close to the cortical surface *in silico* the base capillary had to be at a cortical depth between 20-120 µm. In addition to these selection criteria, we added that the base capillary needed to be at least two branches apart from the boundary of the simulation domain and that the Euclidean distance of the base capillary be smaller than 0.9 times the mean Euclidean distance of all vessels to the center of the MVN in the x-y-plane. The latter two constraints ensured that the simulation results were not affected by the pressure boundary conditions at the domain boundary. The selection criteria for the base capillary are summarized in **Supplementary Table 1**. We found that 193 and 382 capillaries fulfilled the selection criteria in MVN1 and MVN2, respectively. From each subset we randomly chose seven base capillaries.

Subsequently 12-15 capillaries around the base capillary need to be selected to mimic the extent of dilations observed in response to the ablation of 3 pericyte somata. To attain this goal, we first added the capillaries directly adjacent to the upstream bifurcation of the base capillary (generation 1 neighbors) to the set of affected capillaries. Subsequently, the capillaries connected to the generation 1 neighbors were added. The procedure was continued until >= 6 capillaries had been added to the set of affected capillaries. This sequential approach was also initiated at the downstream node of the base capillary until at least 13 capillaries were selected in total. Vessels that were: 1) labeled as main branch (0th order), 2) less than two branches apart from a descending arteriole main branch and 3) less than two branches apart from the domain boundary could not be added to the set of affected capillaries. Thus, the total number of affected capillaries

and the number of capillaries added on the upstream and downstream side of the base capillary varied for different base capillaries. **Supplementary Table 2** summarizes the baseline characteristics for seven distinct base capillaries in MVN1 and MVN2. Two cases per MVN were chosen for further analysis (referred to as Cases 1-4 in the Figures).

To mimic the alterations in response to pericyte ablation *in vivo*, the set of affected capillaries were dilated. Three scenarios were tested: 1) Dilation by 0.6 μm (average response to pericyte ablation in adult mice), 2) Dilation by 1.1 μm (average response in aged mice) and 3) Dilation by 1.6 μm (extreme scenario in aged mice). For all four cases and for each dilation scenario the blood flow simulation was repeated and the time-averaged flow field computed.

Comparing pre- and post-ablation characteristics in silico. In the analyses of the simulation results, we focused on relative flow changes in response to dilation of the capillaries affected by pericyte ablation. As previously stated, due to the presence of RBCs, the flow field fluctuates in time.[70] These fluctuations can also affect the time-averaged flow field. To compute representative relative changes and to avoid large relative changes in capillaries with low baseline flow rate, we employ an absolute threshold of 0.1 μm^3/ms. Changes below this threshold were set to 0. These vessels represented a minority, as 2.6% (MVN1) and 0.7% (MVN2) of all vessels had a time-averaged flow rate < 0.1 μm^3/ms. More details on the definition of a suitable threshold are available in.[71]

To analyze the impact of the local capillary dilations across larger regions of the MVN, we computed the relative flow changes in analysis spheres around the center of the dilated region (**Figure 8d**). The center of the dilated region was defined as the averaged coordinate of all bifurcations of the dilated capillaries. Subsequently, we computed the Euclidean distance of each vessel to the center of the dilated capillaries and grouped the vessels based on the resulting distance to center. Additionally, we separated vessels with flow increase and decrease. Only relative changes >10% from baseline were considered for this analysis.

To more closely inspect relative changes in the vicinity of the dilated vessels we compare the relative flow change in: 1) the dilated capillaries, 2) the capillaries adjacent to the dilated capillaries (Gen1-Neighbors) and 3) the capillaries adjacent to the Gen1-Neighbors (Gen2-Neighbors)(**Figure 8e**). We further examined flow changes at divergent bifurcations with one dilated and one un-dilated outflow vessels (**Figure 8f**). Across the four cases, six such configurations could be identified.

To compare flow heterogeneity in the set of affected capillaries to the expected baseline heterogeneity, we calculate the standard deviation of flow for 193 and 382 different sets of capillaries for MVN1 and MVN2, respectively. These sets of capillaries contain 12-15 capillaries each and were defined similarly as the set of affected capillaries to mimic pericyte ablation. The distribution of standard deviations for MVN1 and MVN2 are shown in **Figure 8g**. The median standard deviation of such a set of capillaries is used as reference for the expected baseline heterogeneity.

To examine for changes in the number of stalled or no-flow capillaries after focal dilations, we set a threshold value at the lowest 5% of all time-averaged flow rates in the upper 200 µm at baseline. For MVN1 and MVN2 this corresponds to a flow rate of 0.37 µm^3/ms and 0.92 µm^3/ms, respectively. For all scenarios (baseline, diameter change = 0.6 µm, 1.1 µm and 1.6 µm) we count the number of stalled/low flow capillaries (*i.e.*, with a time-averaged flow rate smaller than the lowest 5% threshold) in the dilated, generation 1 (Gen-1) and generation 2 (Gen-2) neighbors (**Supplementary Figure 12c**). In a similar manner we quantify the number of low flow stalled capillaries within 200 µm distance from the center of focal dilation (**Supplementary Figure 12d**).

DATA AVAILABILITY

Source data used to generate graphs for this study are available in the Supplementary Materials. Any additional data are available from the corresponding author upon request.

CODE AVAILABILITY.

All details regarding the simulation code used to produce the in silico results are available in the original publication of the in silico model.[35] Further details and explanations are available from F.S. upon request.

REFERENCES

1.   Brown, W.R. & Thore, C.R. Review: cerebral microvascular pathology in ageing and neurodegeneration. *Neuropathology and applied neurobiology* **37**, 56-74 (2011).
2.   Sweeney, M.D., Ayyadurai, S. & Zlokovic, B.V. Pericytes of the neurovascular unit: key functions and signaling pathways. *Nature Neuroscience* **19**, 771-783 (2016).
3.   Gerhardt, H. & Betsholtz, C. Endothelial-pericyte interactions in angiogenesis. *Cell and Tissue Research* **314**, 15-23 (2003).
4.   Daneman, R., Zhou, L., Kebede, A.A. & Barres, B.A. Pericytes are required for blood-brain barrier integrity during embryogenesis. *Nature* **468**, 562-566 (2010).
5.   Armulik, A.*, et al.* Pericytes regulate the blood-brain barrier. *Nature* **468**, 557-561 (2010).
6.   Hartmann, D.A.*, et al.* Brain capillary pericytes exert a substantial but slow influence on blood flow. *Nature neuroscience* **24**, 633-645 (2021).
7.   Hall, C.N.*, et al.* Capillary pericytes regulate cerebral blood flow in health and disease. *Nature* **508**, 55-60 (2014).
8.   Glück, C.*, et al.* Distinct signatures of calcium activity in brain mural cells. *Elife* **10**, e70591.
9.   Rungta, R.L., Chaigneau, E., Osmanski, B.F. & Charpak, S. Vascular Compartmentalization of Functional Hyperemia from the Synapse to the Pia. *Neuron* **99**, 362-375 (2018).
10.   Berthiaume, A.A.*, et al.* Dynamic remodeling of pericytes in vivo maintains capillary coverage in the adult mouse brain. *Cell Reports* **22**, 8-16 (2018).
11.   Berthiaume, A.A., Hartmann, D.A., Majesky, M.W., Bhat, N.R. & Shih, A.Y. Pericyte Structural Remodeling in Cerebrovascular Health and Homeostasis. *Frontiers in Aging Neuroscience* **10**, 210 (2018).
12.   Soto, I.*, et al.* APOE Stabilization by Exercise Prevents Aging Neurovascular Dysfunction and Complement Induction. *Plos Biology* **13**, e1002279 (2015).
13.   Sengillo, J.D.*, et al.* Deficiency in mural vascular cells coincides with blood-brain barrier disruption in Alzheimer's disease. *Brain Pathology* **23**, 303-310 (2013).
14.   Schultz, N.*, et al.* Amyloid-beta 1-40 is associated with alterations in NG2+ pericyte population ex vivo and in vitro. *Aging Cell* **17**, e12728 (2018).
15.   Miners, J.S., Schulz, I. & Love, S. Differing associations between Aβ accumulation, hypoperfusion, blood-brain barrier dysfunction and loss of PDGFRB pericyte marker in the precuneus and parietal white matter in Alzheimer's disease. *Journal of Cerebral Blood Flow & Metabolism* **38**, 103-115 (2017).
16.   Halliday, M.R.*, et al.* Accelerated pericyte degeneration and blood-brain barrier breakdown in apolipoprotein E4 carriers with Alzheimer's disease. *Journal of Cerebral Blood Flow & Metabolism* **36**, 216-227 (2016).
17.   Ding, R.*, et al.* Loss of capillary pericytes and the blood-brain barrier in white matter in poststroke and vascular dementias and Alzheimer's disease. *Brain Pathology* **Epub ahead of print** (2020).
18.   Montagne, A.*, et al.* Blood-brain barrier breakdown in the aging human hippocampus. *Neuron* **85**, 296-302 (2015).
19.   Hunter, J.M.*, et al.* Morphological and pathological evolution of the brain microcirculation in aging and Alzheimer's disease. *PLoS One* **7**, e36893 (2012).
20.   Hartmann, D.A., Coelho-Santos, V. & Shih, A.Y. Pericyte control of blood flow across microvascular zones in the central nervous system. *Annual Review of Physiology* **84**, 331-354 (2021).
21.   Grant, R.I.*, et al.* Organizational hierarchy and structural diversity of microvascular pericytes in adult mouse cortex. *Journal of Cerebral Blood Flow & Metabolism* **39**, 411-425 (2017).

22.   Vazquez-Liebanas, E.*, et al.* Adult-induced genetic ablation distinguishes PDGFB roles in blood-brain barrier maintenance and development. *Journal of Cerebral Blood Flow & Metabolism* **2**, 264-279 (2021).

23.   Nikolakopoulou, A.M.*, et al.* Pericyte loss leads to circulatory failure and pleiotrophin depletion causing neuron loss. *Nature Neuroscience* **22**, 1089-1098 (2019).

24.   Lou, N.*, et al.* Purinergic receptor P2RY12-dependent microglial closure of the injured blood-brain barrier. *Proceedings of the National Academy of Sciences* **113**, 1074-1079 (2016).

25.   Halder, S.K. & Milner, R. A critical role for microglia in maintaining vascular integrity in the hypoxic spinal cord. *Proc Natl Acad Sci U S A* **116**, 26029-26037 (2019).

26.   Davalos, D.*, et al.* Fibrinogen-induced perivascular microglial clustering is required for the development of axonal damage in neuroinflammation. *Nature Communications* **3**, 1227 (2012).

27.   Knowland, D.*, et al.* Stepwise recruitment of transcellular and paracellular pathways underlies blood-brain barrier breakdown in stroke. *Neuron* **82**, 603-617 (2014).

28.   Lyons, D.G., Parpaleix, A., Roche, M. & Charpak, S. Mapping oxygen concentration in the awake mouse brain. *Elife* **5**, e12024 (2016).

29.   Li, Y., Wei, W. & Wang, R.K. Capillary flow homogenization during functional activation revealed by optical coherence tomography angiography based capillary velocimetry. *Scientific Reports* **8**, 4107 (2018).

30.   Li, B.*, et al.* More homogeneous capillary flow and oxygenation in deeper cortical layers correlate with increased oxygen extraction. *Elife* **8**, e42299 (2019).

31.   Jespersen, S.N. & Østergaard, L. The roles of cerebral blood flow, capillary transit time heterogeneity, and oxygen tension in brain oxygenation and metabolism. *Journal of Cerebral Blood Flow & Metabolism* **32**, 264-277 (2012).

32.   Stefanovic, B.*, et al.* Functional reactivity of cerebral capillaries. *Journal of Cerebral Blood Flow & Metabolism* **28**, 961-972 (2007).

33.   Fung, Y.C. Stochastic flow in capillary blood vessels. *Microvascular Research* **5**, 34-48 (1973).

34.   Merlo, A.*, et al.* A few upstream bifurcations drive the spatial distribution of red blood cells in model microfluidic networks. *Soft Matter* **18**, 1463-1478 (2022).

35.   Schmid, F., Tsai, P.S., Kleinfeld, D., Jenny, P. & Weber, B. Depth-dependent flow and pressure characteristics in cortical microvascular networks. *PloS Computational Biology* **13**, e1005392 (2017).

36.   Cruz Hernández, J.C.*, et al.* Neutrophil adhesion in brain capillaries reduces cortical blood flow and impairs memory function in Alzheimer's disease mouse models. *Nature Neuroscience* **22**, 413-420 (2019).

37.   Schager, B. & Brown, C.E. Susceptibility to capillary plugging can predict brain region specific vessel loss with aging. *Journal of Cerebral Blood Flow & Metabolism* **40**, 2475-2490 (2020).

38.   Murugesan, N., Demarest, T.G., Madri, J.A. & Pachter, J.S. Brain regional angiogenic potential at the neurovascular unit during normal aging. *Neurobiology of Aging* **33**, e1-16 (2012).

39.   Shaw, K.*, et al.* Neurovascular coupling and oxygenation are decreased in hippocampus compared to neocortex because of microvascular differences. *Nature Communications* **12**, 3190 (2021).

40.   Østergaard, L.*, et al.* Capillary dysfunction: its detection and causative role in dementias and stroke. *Current neurology and neuroscience reports* **15**, 37 (2015).

41.   Østergaard, L.*, et al.* Cerebral small vessel disease: Capillary pathways to stroke and cognitive decline. *Journal of Cerebral Blood Flow & Metabolism* **36**, 302-325 (2016).

42.   Esipova, T.V.*, et al.* Oxyphor 2P: A high-performance probe for deep-tissue longitudinal oxygen imaging. *Cell Metabolism* **S1550-4131**, 30759-30759 (2019).

43. Choe, Y.G.*, et al.* Pericyte Loss Leads to Capillary Stalling Through Increased Leukocyte-Endothelial Cell Interaction in the Brain. *Frontiers in Cellular Neuroscience* **Epub ahead of print** (2022).

44. Cudmore, R.H., Dougherty, S.E. & Linden, D.J. Cerebral vascular structure in the motor cortex of adult mice is stable and is not altered by voluntary exercise. *Journal of Cerebral Blood Flow & Metabolism* **37**, 3725-3743 (2017).

45. Challa, V.R., Thore, C.R., Moody, D.M., Anstrom, J.A. & Brown, W.R. Increase of white matter string vessels in Alzheimer's disease. *Journal of Alzheimer's Disease* **6**, 379-383 (2004).

46. Reeson, P., Choi, K. & Brown, C.E. VEGF signaling regulates the fate of obstructed capillaries in mouse cortex. *Elife* **7** (2018).

47. Desjardins, M., Berti, R., Lefebvre, J., Dubeau, S. & Lesage, F. Aging-related differences in cerebral capillary blood flow in anesthetized rats. *Neurobiology of Aging* **37**, 1947-1955 (2014).

48. Moeini, M.*, et al.* Compromised microvascular oxygen delivery increases brain tissue vulnerability with age. *Scientific Reports* **8**, 8219 (2018).

49. Li, Y.*, et al.* Aging-associated changes in cerebral vasculature and blood flow as determined by quantitative optical coherence tomography angiography. *Neurobiology of Aging* **70**, 148-159 (2018).

50. Kisler, K.*, et al.* Pericyte degeneration leads to neurovascular uncoupling and limits oxygen supply to brain. *Nature Neuroscience* **20**, 406-416 (2017).

51. Kisler, K.*, et al.* Acute Ablation of Cortical Pericytes Leads to Rapid Neurovascular Uncoupling. *Frontiers in Cellular Neuroscience* **14**, 27 (2020).

52. Smyth, L.C.D.*, et al.* Characterisation of PDGF-BB:PDGFRβ signalling pathways in human brain pericytes: evidence of disruption in Alzheimer's disease. *Communications Biology* **5**, 235 (2022).

53. Arango-Lievano, M.*, et al.* Topographic Reorganization of Cerebrovascular Mural Cells under Seizure Conditions. *Cell reports* **23**, 1045-1059 (2018).

54. Zhu, W.*, et al.* Thalidomide Reduces Hemorrhage of Brain Arteriovenous Malformations in a Mouse Model. *Stroke* **49**, 1232-1240 (2018).

55. Bauditz, J. Effective treatment of gastrointestinal bleeding with thalidomide--Chances and limitations. *World Journal of Gastroenterology* **22**, 158-164 (2016).

56. Lebrin, F.*, et al.* Thalidomide stimulates vessel maturation and reduces epistaxis in individuals with hereditary hemorrhagic telangiectasia. *Nature Medicine* **16**, 420-428 (2010).

METHODS REFERENCES

57. Hartmann, D.A.*, et al.* Pericyte structure and distribution in the cerebral cortex revealed by high-resolution imaging of transgenic mice. *Neurophotonics*, 041402 (2015).

58. Cuttler, A.S.*, et al.* Characterization of Pdgfrb-Cre transgenic mice reveals reduction of ROSA26 reporter activity in remodeling arteries. *Genesis* **49**, 673-680 (2011).

59. Madisen, L.*, et al.* A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nature Neuroscience* **13**, 133-140 (2010).

60. Davalos, D.*, et al.* ATP mediates rapid microglial response to local brain injury *in vivo*. *Nature Neuroscience* **8**, 752-758 (2005).

61. Nimmerjahn, A., Kirchhoff, F. & Helmchen, F. Resting microglial cells are highly dynamic surveillants of brain parenchyma *in vivo*. *Science* **308**, 1314-1318 (2005).

62. Drew, P.J.*, et al.* Chronic optical access through a polished and reinforced thinned skull. *Nature Methods* **7**, 981-984 (2010).

63. Nielson, C.D. & Shih, A.Y. In vivo single cell optical ablation of brain pericytes. *Frontiers in Neuroscience* **in press** (2022).

64.   Li, B.*, et al.* Two-photon microscopic imaging of capillary red blood cell flux in mouse brain reveals vulnerability of cerebral white matter to hypoperfusion. *Journal of Cerebral Blood Flow & Metabolism* **40**, 501-512 (2019).

65.   Shigetomi, E.*, et al.* Imaging calcium microdomains within entire astrocyte territories and endfeet with GCaMPs expressed using adeno-associated viruses. *Journal of General Physiology* **141**, 633-647 (2013).

66.   Watson, A.N.*, et al.* Mild pericyte deficiency is associated with aberrant brain microvascular flow in aged PDGFRβ+/- mice. *Journal of Cerebral Blood Flow & Metabolism* (2020).

67.   Blinder, P.*, et al.* The cortical angiome: An interconnected vascular network with noncolumnar patterns of blood flow. *Nature Neuroscience* **16**, 889-897 (2013).

68.   Schmid, F., Reichold, J., Weber, B. & Jenny, P. The impact of capillary dilation on the distribution of red blood cells in artificial networks. *American Journal of Physiology: Heart and Circulatory Physiology* **308**, H733-H742 (2015).

69.   Pries, A.R. & Secomb, T.W. Microvascular blood viscosity *in vivo* and the endothelial surface layer. *American Journal of Physiology - Heart and Circulation Physiology* **289**, H2657-2664 (2005).

70.   Schmid, F., Barrett, M.J.P., Obrist, D., Weber, B. & Jenny, P. Red blood cells stabilize flow in brain microvascular networks. *PloS Computational Biology* **15**, e1007231 (2019).

71.   Schmid, F., Conti, G., Jenny, P. & Weber, B. The severity of microstrokes depends on local vascular topology and baseline perfusion. *Elife* **10**, e60208 (2021).

ACKNOWLEDGMENTS

Our work is supported by grants to A. Shih from the NIH/NINDS (NS106138, NS097775) and NIH/NIA (AG063031, AG062738), the American Heart Association (14GRNT20480366), Alzheimer's Association NIRG award (2016-NIRG-397149), a scholarship for Research in the Biology of Aging to A.A. Berthiaume from the American Federation for Aging Research, and by a grant from the Swiss National Science Foundation (310030_182703) to B. Weber. We appreciate the helpful comments and discussion of B. Gumbiner, C. Cowan, N. Bhat, L. Faulhaber, and J. Gust. We thank D. Agalliu for generously providing Claudin5-eGFP mice.

AUTHOR CONTRIBUTION

Conceptualization and experimental design by AAB and AYS. Experiments were conducted by AAB, SS, VCS, CN, and AYS. Data analysis and statistics were performed by AAB, SS, VSC, CN and AYS. *In silico* studies and analyses were performed by FS and BW. The manuscript was written by AAB, FS and AYS with contributions from SS, VCS, CN, BW and MWM.

COMPETING INTERESTS STATEMENT

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. Acute consequence of contiguous multiple pericyte ablation

(a) In vivo two-photon imaging through a chronic cranial window in anesthetized PDGFRβ-tdTomato mice.

(b) Capillary bed with vessels labeled green with i.v. FITC-dextran (70 kDa), and pericytes genetically labeled in red. Inset shows the structure of capillary pericytes.

(c) A capillary pericyte targeted for two-photon ablation encircled in yellow. Ablative line-scanning is restricted to the pericyte soma. Post-ablation image shows loss of fluorescence from targeted pericyte, including its processes, within minutes post-ablation. Neighboring pericytes are unaffected.

(d) Schematic showing metrics collected following ablation of 3 contiguous pericytes. Left, targeted pericytes are marked with asterisks. Right, metrics include total capillary length uncovered, number of remodeling pericytes, and number of remodeling pericyte processes.

(e,f) Before and 5 minutes after triple pericyte ablation in adult and aged animal. Arrowheads indicate targeted pericyte somata, and dotted magenta line outlines approximate regions of uncovered endothelium. Inset shows a magnified region with pericyte ablation and uncovered territory 5 minutes afterwards.

(g-i) Comparison of vascular metrics between adult and aged mice after triple pericyte ablation. (g) Total capillary length uncovered; $t(12)=1.790$, $p=0.0988$. (h) Number of fully and partially uncovered capillary segments; $t(12)=0.1936$, $p=0.8497$. (i) Range of capillary segment lengths; $t(195)=0.2325$, $p=0.8164$. All are unpaired t tests, for n=7 triple pericyte ablations (6 adult mice), 7 triple pericyte ablations (6 aged mice) for panels (g) and (h) and n=99 capillary segments (6 adult mice), n=capillary 98 segments (6 aged mice) for panel (i). Data shown as mean ± SD. ns= non-significant.

(j,k) Comparison of pericyte metrics between adult and aged mice after triple pericyte ablation. (j) Number of remodeling pericytes; $t(12)=0.6030$, $p=0.5577$. (k) Number of remodeling pericyte processes; $t(12)=0.7206$, $p=0.4850$. Unpaired t tests for n=7 triple pericyte ablation (6 adult mice), 7 triple pericyte ablations (6 aged mice). Data shown as mean ± SD.

Figure 2. Pericyte remodeling ensures capillary coverage following loss of pericyte contact.

(a) Representative example of pericyte remodeling from an adult mouse (4 months old at pre-ablation), showing the area before triple pericyte ablation, and at select time-points after ablation. Arrowheads indicate pericytes targeted for ablation. The lower panels show mural cells only. Insets show magnified view of region with pericyte ablation followed with remodeling of neighboring pericyte processes.

(b) Schematic of capillary coverage by pericytes on each imaging day.

(c) Schematic representation of growth contributed by different pericyte processes over 21 days. Each color represents an individual pericyte process, with location of original terminus of the process as a dashed line.

(d) Plot of added pericyte process length over time from Day 0. Each color corresponds to an individual pericyte process from (c). Closed circles mark the day of pericyte-pericyte contact, after which growth stops (dashed lines).

(e-h) Representative example from an aged mouse (21 months old at pre-ablation), with same layout as for adult mouse.

Figure 3. Inefficient pericyte remodeling in aged mice fails to regain complete capillary coverage.

(a) *In vivo* two-photon images of a pericyte process growing over time (white arrow), until it is inhibited by pericyte-pericyte contact.

(b) Percent of total pericyte processes making pericyte-pericyte contact on each imaging day across age groups. n=60 processes from 6 adult mice, n=57 processes from 6 aged mice.

(c) Pericyte process extension over time in each age group, including all processes examined. n=60 processes from 6 adult mice, n=57 processes from 6 aged mice.

(d) Plot showing the extension of pericyte processes, excluding those that have made pericyte-pericyte contact, at each post-ablation imaging day. Mixed-effects analysis with Sidak's multiple comparisons, $F_{(1,115)}=9.979$; overall effect **p=0.002, Day 3, *p=0.0187; Day 7, **p=0.0056; Day 14, **p=0.0087; Day 21, p=0.6574. Data shown as mean ± SEM

(e,f) Maximum and average rate of pericyte process growth in aged compared to adult mice. (e) Unpaired t tests with Welch's correction for unequal variances: $t(109.1)=3.313$; **$p=0.0013$; (f) $t(90.50)=4.977$; ****$p<0.0001$. n=60 processes from 6 adult mice, 57 processes from 6 aged mice.

(g) Maximum process extension achieved in 21 days. Unpaired t test $t(115)=0.9370$; $p=0.3507$. n=60 processes from 6 adult mice, 57 processes from 6 aged mice.

(h) Total process length at day 21. $t(112)=1.066$; $p=0.2889$. Data in panels e-h shown as mean ± SD. Statistics performed with unpaired t tests for n=60 processes from 6 adult mice, n=57 processes from 6 aged mice.

(i) Uncovered vessel length remaining over time. Each line represents one triple pericyte ablation region. Adult, n=7 triple pericyte ablations; 6 mice; Aged n=7 triple pericyte ablations; 6 mice.

(j) Comparison of average vessel length remaining on each imaging day across age groups. Two-way ANOVA with Sidak's multiple comparisons test, $F(1, 12)=16.72$; overall effect **$p=0.0015$, Day 0, $p=0.4196$. Day 3, *$p=0.0118$. Day 7, **$p=0.0098$. Day 14, **$p=0.0052$. Day 21, *$p=0.0490$. Adult, n=7 regions, 6 mice; Aged, n=7 regions, 6 mice. Day 49 is from a subset of triple pericyte ablation experiments in aged animals only (5 regions from 4 mice), not included in statistical analyses. Data shown as mean ± SEM.

Figure 4. Mural cell remodeling capacity varies with age across microvascular zones.

(a) *In vivo* two-photon image of a microvascular network from penetrating arteriole to ascending venule, with branch orders in white. Bottom image highlights mural cell subtypes within zones.

(b) Example of an ensheathing pericyte from an adult mouse remodeling into the capillary zone. Arrow indicates position of leading process terminus over time. Bottom row shows high-resolution image of the cell at 21 days post-ablation.

(c) Example of a mesh and thin-strand pericyte from an adult mouse growing within the capillary bed. Arrows show growing terminal ends of processes.

(d) Example of a venule smooth muscle cell process from an adult mouse growing into the capillary bed, as indicated by arrows.

(e) Average process growth by mural cells in different microvascular zones. Two-way ANOVA with Tukey's multiple comparisons test, For age comparison, $F(1,102)=21.45$; overall effect,

****p=0.0001; Arteriole-capillary transition (adult vs. aged) ****p<0.0001; capillary (adult vs. aged) **p=0.0016; Venule (adult vs. aged) p>0.9999. For vessel type comparison, $F_{(2,102)}=0.1838$; overall effect, p=0.8324, For interaction between age and vessel type, $F_{(2,102)} = 6.029$, overall effect, **p=0.0033. Arteriole-capillary transition vs. capillary (adult), p=0.5833; arteriole-capillary transition vs. venule (adult), **p<0.4125; capillary vs. venule (adult), p=0.9533. Arteriole-capillary transition vs. capillary (aged), p=0.4855; arteriole-capillary transition vs. venule (aged), **p<0.0398; capillary vs. venule (aged), p=0.4516. Adult, n=7 regions, 6 mice; Aged, n=7 regions, 6 mice.

(f) Maximum process extension by mural cells in different microvascular zones. Two-way ANOVA. For age comparison, $F_{(1,97)}=0.1455$; overall effect, p=0.7037. For vessel type comparison, $F_{(2,97)}=2.613$; overall effect, p=0.0784. For interaction between age and vessel type, $F_{(2,97)}=5.532$; **p=0.0053. Arteriole-capillary transition vs. capillary (adult), p>0.9999; arteriole-capillary transition vs. venule (adult), p<0.9897; capillary vs. venule (adult), p=0.3511. Arteriole-capillary transition vs. capillary (aged), p=0.8750; arteriole-capillary transition vs. venule (aged), **p<0.0014; capillary vs. venule (aged), **p=0.0044. Adult, n=7 regions, 6 mice; Aged, n=7 regions, 6 mice.

Figure 5. Lack of overt BBB disruption or perivascular inflammation with focal pericyte loss.

(a) Imaging time course to examine for extravasation of 10 kDa FITC-dextran dye in aged mice. Pre-ablation image shows location of targeted pericytes and sham irradiation controls. At day 3, T0 shows regions of interest (ROI) from which fluorescence intensity measurements were collected over time.

(b) Intensity of FITC-dextran fluorescence within the capillary lumen as a function of time post-injection.

(c) Intensity of FITC-dextran fluorescence in parenchymal ROI between ablation and sham regions.

(d) Rate of fluorescence intensity change between ablation and sham ROI regions. Wilcoxon rank sum test, p = 0.904. N = 7 ablation regions and n = 7 sham region from 4 aged mice.

(e,f) In vivo two-photon images from adult Pdgfrβ-tdtomato::Cx3Cr1-GFP mice, showing microglia reaction to (e) pericyte ablation or (f) sham irradiation after 30 minutes and 3 days. ROIs are

drawn in the region directly exposed to laser ablation or irradiation (focal point), or the capillary segments covered by pericyte processes (territory)

(g,h) GFP intensity in focal point (g) and territory (h) after pericyte ablation and sham, shown in arbitrary units (a.u.). Two-way repeated measures ANOVA with Sidak's multiple comparisons. Focal point: $F_{(1.45, 7.252)} = 24.41$, overall effect ***$p=0.0009$. Pericyte ablation vs. sham irradiation *$p=0.0216$ at 30 min; $p=0.5504$ at 3 days. Territory: $F_{(1.149, 5.744)} = 3.170$, $p=0.1263$.

(i,j) GFP intensity in focal point (i) and territory (j) in aged mice. Two-way repeated measures ANOVA. Focal point: $F_{(1.03, 7.183)} = 3.214$, overall effect $p=0.1146$. Territory: $F_{(1.761, 12.33)} = 0.2922$, $p=0.7248$. For (g) and (h), n= 3 pericyte ablations, 4 sham irradiations from 5 mice between 3-6 months of age. For (i) and (j), n= 5 pericyte ablations, 4 sham irradiations from 3 mice between18-24 months of age. Data presented as mean ± SEM.

Figure 6. Pericyte coverage loss leading to augmented capillary dilations in aged brain.

(a) Comparison of baseline capillary diameters across age. Unpaired t test, $t_{(60)}=0.7245$; $p=0.4716$, unpaired t test for n=30 capillaries from 6 adult mice, n=32 capillaries from 6 aged mice.

(b) Capillary dilation occurs in region with loss of pericyte coverage, but not in adjacent capillaries that maintain pericyte coverage. Arrowhead points to pericyte targeted for ablation.

(c) Example capillary region from adult mouse following triple pericyte ablation. Arrowheads show two targeted cells. Bottom row shows i.v. dye alone, with capillary segments measured over time identified by blue lines.

(d) Plot of capillary diameters over time from adult mice. Repeated measures one-way ANOVA with Tukey's multiple comparisons test, $F_{(1.824, 45.60)} = 35.95$, ****$p<0.0001$. Baseline vs uncovered, ****$p<0.0001$; baseline vs. re-covered, $p=0.7393$; uncovered vs. re-covered, ****$p<0.0001$. N=26 capillaries from 6 mice.

(e) Example capillary region from aged mouse following triple pericyte ablation. Arrowheads point to targeted pericytes. Bottom row shows i.v. dye alone, with capillary segments measured over time identified by purple lines.

(f) Plot of capillary diameters over time from aged mice. Repeated measures one-way ANOVA with Tukey's multiple comparisons test, $F(1.914, 53.59)= 111.3$, ****$p<0.000$. Baseline vs uncovered, ****$p<0.0001$; baseline vs. re-covered, ** $p=0.0018$; uncovered vs. re-covered, ****$p<0.0001$. N=29 capillaries from 6 mice.

(g) Change in diameter from baseline at 3 days post ablation, when capillaries lack pericyte coverage. Unpaired t test, $t(54)=3.501$, ***$p=0.0009$. N=26 capillaries in 6 adult, n=30 capillaries in 6 aged mice.

(h) Difference between baseline and re-covered capillary diameter detected in adult and aged groups. Adult: One sample t-test, $t(25)=0.7451$; $p=0.4631$, for n=26 (6 mice). Aged: One sample t-test, $t(33)=4.172$; ***$p=0.0002$, for n=34 capillaries (6 mice). Difference in re-covered diameter between adult and aged mice. Unpaired t-test with Welch's correction, $t(41.40)=2.751$; **$p=0.0088$.

(I) Dilations persist for capillaries that do no regain pericyte coverage at 21 days in aged mice. Repeated measures one-way ANOVA with Tukey's multiple comparisons test, $F(1.967, 19.67)=31.35$; overall effect ****$p<0.0001$. Pre-ablation vs 3 days, ****$p<0.0001$; pre-ablation vs 21 days, ***$p=0.0002$; 3 days vs 21 days, $p=0.5976$. N=11 capillaries from 6 mice.

Figure 7. Blood flow changes in ablation and surrounding capillary areas.

(a) Image showing capillaries uncovered by a triple ablation (purple).

(b) Three days post-ablation, changes in red blood cell (RBC) flux from baseline are color coded in a subset of vessels.

(c) Overlay of blood flow direction on vascular image. Ascending venules are marked in blue, and penetrating arterioles in red.

(d) Blood cell flux before and 3 days post ablation for segments in and immediately around the ablation area. Paired t test, $t(24)=3.880$; ***$p=0.0007$ for n=25 capillaries from an aged mouse. Significantly higher variance was detected at 3 days compared to baseline, F test to compare variances, $F(24, 24)=3.545$; ***$p=0.0029$. Data shown as mean ± SD.

(e) Proportion of vessels in the area that experienced an increase vs. decrease in flux on Day 3

(f) Blood cell flux in capillaries that increased in flux from baseline to 3 days post-ablation. Paired t test, $t(17)=6.273$; ****$p<0.0001$, for n=18 capillaries from an aged mouse.

(g) A subset of capillaries decreased in flux following pericyte ablation. Paired t test, $t(6)=3.229$; *$p=0.0179$, n=6 capillaries from an aged mouse. Data in (d,f,g) are represented as mean ± SD.

(h-j) A divergent bifurcation that loses pericyte coverage in one downstream branch, while the alternate route remains covered. RBCs passing through i.v. dye captured by line scans are shown for each vessel (*i, ii, iii*) over time.

(k-m) A convergent bifurcation which loses pericyte coverage to one upstream vessel. RBCs passing through i.v. dye captured by line scans are shown for each vessel (*i, ii, iii*) over time.

Figure 8. *In silico* study on the impact of local capillary dilations in response to pericyte ablation.

(a) Microvascular network 1 (MVN1) from mouse parietal cortex

(b) Example of a typical region affected by pericyte ablation in MVN1. The color bar shows the flow rates in individual capillaries pre-ablation.

(c) Relative flow changes >10% of baseline in the entire MVN1 for an increasing extent of dilation. Vessels with a relative flow change between -10% and +10% are not colored and depicted by the light grey lines.

(d) Relative flow change >10% of baseline at different distances to the center of the dilated capillaries.

(e) Relative flow changes in the dilated vessels and their direct neighbors (Gen1-Neighbors and Gen2-Neighbors, see schematic).

(f) Flow steal at divergent bifurcations with one dilated outflow capillary. The scatter plots (d-f) show the relative flow change of individual vessels. The data of all four cases has been combined.

(g) Distribution of standard deviations for numerous undilated capillary sets in upper cortex. The expected baseline heterogeneity of capillaries is defined as the median of all standard deviations. Left: MVN1 (n(sets)= 170, average set size = 14.5 ±1.6), Right: MVN2 (n(sets)= 346, average set size = 14.6±1.8).

(h) Standard deviation of flow in the dilated capillaries at baseline and with increasing extent of dilation. Dashed lines show expected baseline heterogeneities at 0.50 and 0.75 quantile (see panel g).

Figure 9. Blood flow stalls and regression in capillaries neighboring dilation.

(a) A divergent bifurcation close to the arteriole-capillary transition zone. One branch of the bifurcation is dilated from pericyte loss 3 days post-ablation, and the resulting re-direction of flow creates a stall in the alternate, covered branch.

(b) Example of blood flow defects at a capillary junction near the arteriole-capillary transition zone. At 21 days post-ablation, a central capillary is uncovered, dilated and flowing. The other branches have stalled in flow or regressed.

(c) Occurrence of stalls in triple off-target sham irradiation and triple pericyte ablation experiments. Unpaired t-test with Welch's correction for unequal variances, t(16.10)=2.698; overall effect *p=0.0158, for n=6 off-target irradiation experiments (3 in adult mice, 3 in aged mice), n= 14 pericyte ablation experiments (7 in adult mice, 7 in aged mice).

(d) Additional example of a capillary regression occurring after a prolonged stall.

(e) Plot of the relative location of stall and regression events following pericyte ablation, by branch order from a dilated segment.

(f) Schematic summarizing findings of blood flow interruption following pericyte coverage loss.

Berthiaume et al. Fig 1

Berthiaume et al. Fig 2

Berthiaume et al. Fig 3

Berthiaume et al. Fig 4

Berthiaume et al. Fig 5

Berthiaume et al. Fig 6

a Region lacking pericyte contact following ablation

Uncovered capillaries

50 μm

b Change in RBC flux (3 days post-ablation)

Increased flux
Decreased flux

c Direction of blood flow

A
k-m
h-j
V
A
V

d RBC flux

Cells/s

***

Baseline   3 day post

e Change from baseline

Increased flux
Decreased flux

28%
72%

n = 25 capillaries

f Capillaries with RBC flux increase

Cells/s

****

Baseline   3 day post

g Capillaries with RBC flux decrease

Cells/s

*

Baseline   3 day post

h Divergent bifurcation

Baseline   3 days post

10 μm

i
iii
ii

i

135
70
54

253
246
37

RBC flux (cells/s)

j

i   *ii*   *iii*
pre post pre post pre post

0.1s
10 μm

k Convergent bifurcation

Baseline   3 days post

i
iii
ii

l

196
207
64

135
276
242

RBC flux (cells/s)

m

i   *ii*   *iii*
pre post pre post pre post

0.1s
10 μm

Berthiaume et al. Fig 7

Berthiaume et al. Fig 8

Berthiaume et al. Fig 9

Supplementary Material:

Deficiency in pericyte remodeling as a basis for impaired capillary flow and structure during brain aging

By Andrée-Anne Berthiaume, Franca Schmid, Stefan Stamenkovic, Vanessa Coelho-Santos, Cara D. Nielson, Bruno Weber, Mark W. Majesky, and Andy Y. Shih

Supplementary Fig. 1. Comparison of basal vascular features in superficial cortex of adult and aged mice. (a) Representative image of a superficial capillary bed volume (10-100 μm from pial surface) in adult mice (3-6 months) and **(b)** in aged mice (18-24 months). **(c)** (Top) Representation of 3D reconstruction of the capillary vascular network. (Bottom) No difference in capillary length per imaging volume (350x350x90 μm regions) in adult versus aged animals, t(31)=0.7711, p=0.4465 by unpaired t test. N= 15 regions from 9 adult mice, n=18 regions from 8 aged mice. **(d)** Example image of capillary junctions (white circles), and quantification showing no difference in capillary junctions per imaging volume (350x350x90 μm) in adult and aged mice, t(31)=0.03123, p=0.9753 by unpaired t test for n=15 volumes from 9 adult mice, n=18 volumes from 8 aged mice. **(e)** Example image of pericyte somata (cyan circles), and plot showing no difference in abundance per imaging volume (350x350x90 μm) in adult versus aged mice, t(31)=0.5517, p=0.5851 by unpaired t-test for n=15 volumes from 9 adult mice, n=18 volumes from 8 aged mice. **(f)** Exent of the capillary network not covered by pericytes, t(32) = 0.8138, p=0.4216 by unpaired t-test for n=15 volumes from 9 adult mice, n=18 volumes from 8 aged mice. Left image from panel (f) adapted from Berthiame et al. Frontiers in Aging Neuroscience, 2018 (PMID: 30065645). All data are shown as mean ± SD.

Supplementary Fig. 2. Additional examples of triple pericyte ablations in adult and aged mice. (a) Full time course of 21 day triple pericyte ablation in an adult mouse. Targeted pericytes marked with yellow asterisks. Bottom rows show pericytes only. **(b)** Schematic of capillary coverage over time. **(c)** Example of triple pericyte ablation experiment in an aged mouse, with additional chronic timepoint at 49 days post-ablation and **(d)** corresponding capillary coverage over time. **(e)** Second example of ablation from an aged mouse including 49-day timepoint. and **(f)** pericyte coverage of this capillary network over time.

Supplementary Fig. 3. Contributors to pericyte process growth potential. (a) Pericyte process growing onto a capillary over time, with two occurrences of process splitting during extension. **(b)** Amount of pericyte process growth for processes that split during extension versus those that did not. T(73.79)=7.824, ****p<0.0001 by unpaired t test with Welch's correction for n=50 processes (no split), and 66 processes (split) from 12 mice (6 adult, 6 aged). Presented as mean ± SD. **(c)** Baseline process lengths are no different across age groups. T(113)=0.0847, p=0.9327 by unpaired t test. N=57 processes from 6 adult mice, 58 processes from 6 aged mice. Shown as mean ± SD. **(d)** Correlating baseline process length to length added during remodeling reveals a strong negative correlation. Processes that split versus those that did not are also noted. Linear regression F(1, 40)=11.63, **p=0.0015. Pearson correlation r= -0.4746, for n=42 processes from 11 mice pooled across age groups (5 adult, 6 aged).

Supplementary Fig. 4. Pericyte processes distort capillary shape to gain endothelial coverage (a) Tracking a remodeling pericyte growing over a capillary segment reveals the formation of a "microbend" near the growing process tip. **(b)** Examples of multiple microbends forming post-ablation in a triple ablation area in an adult mouse. **(c)** Multiple microbends created by remodeling processes in a triple ablation area in an aged mouse. **(d)** Plot of the proportion of processes which form a microbend in the underlying capillary during their extension. n=10 triple pericyte ablation experiments from 8 mice (4 adult, 4 aged). **(e)** Greater average rate of growth of processes that form microbends compared to those that do not. T(24.21)=2.922, **p=0.0074, by unpaired t test with Welch's correction. **(f)** Greater overall final growth at Day 21 for processes that formed microbends compared to those that did not. T(25.54)=3.325, **p=0.0027 by unpaired t test with Welch's correction. For both (e) and (f), n=20 pericyte processes creating microbends, 57 pericyte processes that don't create microbends, across 8 mice (4 adult, 4 aged). Data represented as mean ± SD.

Supplementary Fig. 5. Pericyte growth across microvascular zones in the adult and aged brain. (a) An ensheathing pericyte from an adult mouse remodeling into the capillary zone. Ablated cells marked with a yellow asterisk, and growth front of processes marked with magenta dashed lines. **(b)** An ensheathing pericyte failing to remodel into the capillary zone over 7 days in an aged mouse. **(c)** Venule SMC from an adult brain growing into the capillary zone. **(d)** Venule SMC extending onto a capillary in an aged brain. **(e)** Image of an arteriole-capillary transition zone, with branch orders from the penetrating arteriole (0) marked. **(f)** Inset from (e), showing a 3rd order vessel with an ensheathing pericyte targeted for ablation. Over the next 7 days, neighboring ensheathing pericytes extend their processes into the area and their processes maintain an ensheathing morphology. Scale bars = 10μm.

Site of pericyte ablation
Perivascular region of interest

70 kDa i.v. dye

Supplementary Fig. 6. Blood-brain barrier leakage is very rare with focal loss of pericyte coverage.
(a) Three examples of vasculature before and at two time points after triple pericyte ablation in adult mice. Sites of ablation are denoted with yellow asterisks, and perivascular regions of interest (ROI) for quantification of i.v. dye extravasation are in blue dashed lines. **(b)** Three examples of the vascular channel before after triple pericyte ablation in aged mice. Asterisks indicate location of pericyte ablation and ROIs for leakage assessment in dashed lines. Note the lack of 70 kDa i.v. dye permeating outside of the vessel walls across all examples. **(c)** (Top) A singular case of active leakage of 70 kDa i.v. dye noted at 3 days post-ablation in an aged mouse. No leakage occurred at 5 minutes post-ablation, suggesting that this was not caused by acute laser damage. (Bottom) Insets showing high resolution images of the capillary segment of interest. Sites of ablation are marked with asterisks, and dash ROI shows where parenchymal signal intensity measurements were taken. Scale bars = 50 μm. **(d)** Quantification of i.v. dye extravasation from perivascular locations lacking pericyte coverage at 3 days post ablation compared to pre-ablation. No parenchymal intensity change detected after pericyte loss in adult group, T(6)=0.8406, p=0.4328, by one-sample t test. No perivascular leakage detected on average in aged group, T(6)=1.070, p=0.3257 by one-sample t test. Singular example of leakage in this group can be seen as outlier driving large SD. No difference in intensity change between groups, T(6.345)=1.196, p=0.2746, by t test with Welch's correction for unequal variances. Data represented as mean ±SD.

Supplementary Fig. 7. No significant change in tight junction structure following the loss of individual pericytes. (a,b) In vivo two-photon images from adult Pdgfrb-tdtomato;Claudin5-eGFP mice, showing pericytes and tight junction (TJ) structure before and 5 min, 3 days, and 7 days following (a) pericyte ablation or (b) off-target sham irradiation. Orange arrowheads denote gaps in the TJ strands, and pink arrowheads denote protrusions. **(c)** No significant difference in the percentage of TJ strands containing protrusions between the ablation and off-target sham irradiation groups at any post-irradiation timepoint compared to baseline. However, at day 7 there is a trend towards an increased number of protrusions in the ablation group compared to the control group. Two-way repeated measures ANOVA; no interaction detected between time and experiment type, $F_{(4,20)}=2.317$, p=0.0925. **(d)** No significant difference in the percentage of TJ strands containing gaps between the ablation and control groups at any post-irradiation timepoint compared to baseline. Two-way repeated measures ANOVA; no interaction detected between time and experiment type, $F_{(4,20)}=0.5185$, p=0.7231. For (c) and (d), n=4 pericyte ablations and 3 control irradiations (2 off-target irradiations and 1 failed pericyte ablation) from 3 mice aged 7-12 months. Data presented as mean ± SEM.

Supplementary Fig. 8. Apposition of astrocyte endfeet to the capillary wall is not overtly disrupted by focal pericyte loss. (a,b) In vivo two-photon images from aged Pdgfrb-tdtomato mice expressing Lck-GFP in astrocytes, before and 3 days after pericyte ablation. Left images show astrocytes (GFP) together with i.v. dye (Alexa 680-dextran) and right images show astrocyte together with pericytes. Here, a single pericyte has been ablated and the fluorescence intensity of astrocyte and i.v. dye signal is taken across a region affected by the pericyte ablation (white square). **(c,d)** Fluorescence intensity of astrocyte and i.v. dye signals plotted as a function of distance across the capillary cross-section at pre-ablation and post-ablation time-points. Note how sharp GFP peaks flanking the capillary lumen are still present after pericyte loss, suggesting intact endfoot coverage. **(e)** The distance between the intravascular wall (measured by FWHM) and the peak of the GFP signal does not change with pericyte loss. N=4 separate pericyte ablations were performed in 2 aged mice. Repeated-measures t-test, t(3)=1.131, p=0.3402.

Supplementary Fig. 9. Dilations are localized to portions of the capillary segment lacking pericyte coverage. (a) Example of a capillary region before and 3 days after a pericyte ablation in an 18 month-old mouse. The edge of new process growth is marked with an orange arrow at 3 days post-ablation. **(b)** Corresponding i.v. dye channel, showing crosslines at which full-width at half-maximum intensity measurements were taken for diameter. Three individual capillary segments are examined. The edge of new pericyte process growth along Capillary 2 is marked at 3 days post-ablation. **(c)** Diameter of a fully uncovered capillary at Day 3 shows consistent dilations across its entire length. **(d)** A partially re-covered segment is dilated only beyond the end of a remodeling process. **(e)** A capillary segment not uncovered by ablation shows no change in diameter at 3 days compared to its baseline diameter. Data shown as mean ± SEM.

Supplementary Fig. 10. No dilations detected 3 days following off-target sham irradiation. (a) Example of a triple off-target irradiation in an adult mouse. Areas of irradiation are marked with yellow asterisks. Capillary diameters collected over time at segments marked with blue lines. **(b)** Example of a triple off-target irradiation in an aged mouse, as shown with yellow asterisks. Purple lines show segments measured for diameter. **(c)** No change in diameter in either age group at 3 days post-irradiation compared to baseline. For adults, t(14)=0.6475, p=0.5278 for 15 capillaries from 4 adult mice by one-sample t test. For aged, t(8)=0.6732, p=0.5198 for 9 capillaries from 3 aged mice by one-sample t test. Data shown as mean ± SD.

Supplementary Figure 11. Comparison of capillary diameters in awake versus isoflurane anesthesia conditions in aged mice.
(a) In vivo two-photon images of the same capillaries from an aged mouse, taken while under (left) isoflurane anesthesia (1.5% MAC) and (right) awake state. Pericyte targeted for ablation is marked with an asterisk. **(b)** Overlay of capillary area from (a), with image from anesthetized state in red and image from awake state in green. Note the presence of an artery within the field of view, which dilates under isoflurane and therefore serves as a positive control. **(c)** Capillary images 3 days post-ablation of the target pericyte, under both isoflurane and awake imaging conditions. Capillary segments lacking pericyte coverage are marked by arrowheads. **(d)** Overlay of capillary images obtained under isoflurane or awake 3 days post-ablation. **(e)** (Left) Quantification of capillary diameters in 3 separate regions from 2 aged mice (1 female) having undergone single pericyte ablation 3 days prior. Unpaired t-test t(26) = 1.819, p=0.0804. Diameters from uncovered segments are circled in black. (Right) Scatter plot showing diameter change in individual capillaries going from anesthetized to awake state. **(f)** Correlation of diameters in isoflurane vs awake 3 days post-ablation, with segments uncovered and dilated circled in black. Pearson correlation, r=0.9382, ****p<0.0001. Theoretical perfect correlation is shown as a red line for visual comparison.

Supplementary Fig. 12. Capillary flow changes after focal capillary dilation in silico. (a) Flow steal at divergent bifurcations with one dilated outflow capillary shown with change in RBC flux. **(b)** Flow rate in dilated vessels for baseline (diameter increase = 0) and with increasing extent of dilation (diameter increase 0.6 µm, 1.1 µm and 1.6 µm). Number of dilated vessels: 13-14 (see Supplementary Table 1). Case 2 and Case 3 are in the center of the capillary bed, whereas Case 1 is closer to the venule side and Case 4 is closer to the arteriole side. **(c,d)** Number of low flow or stalled capillaries in different vessel categories (c) or within a distance of 200 µm from the center of dilated vessels (c) does not increase with greater levels of focal capillary dilation. The threshold for low flow capillaries for the in silico study is defined as the lowest 5% of the capillary flow rates in the upper 200 um of the microvascular networks.

Supplementary Fig. 13. Additional examples of capillary regressions after prolonged blood flow stalls. (a) Two additional images from aged mice of blood flow stalls in capillary segments immediately adjacent to capillaries uncovered by pericyte ablation. **(b)** Detailed view of blood flow redirection (arrows) in capillaries downstream from the arteriole-capillary transition zone. The pericyte loss-induced dilation in one segment 3 days post-ablation results in the complete steal of flow away from the alternate, pericyte-covered segment. Example from an adult mouse. **(c)** The development of a capillary regression in an adult mouse over time following prolonged capillary stalling. A blood flow stall is observed 3 days post-ablation, in a capillary adjacent to a pericyte-uncovered, dilated segment. This capillary is undergoing regression by 7 days post-ablation, a process which is fully complete by 21 days post-ablation.