Sex-Specific Effects of Psychedelic Drug Exposure on Central Amygdala Reactivity and

Behavioral Responding

DP Effinger, B.S.[1,2], SG Quadir, Ph.D[1,2], MC Ramage[1,2], MG Cone[1,2], MA Herman, Ph.D[1,2]

[1] Department of Pharmacology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599

[2] Bowles Center for Alcohol Studies, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599

Corresponding Author: M.A. Herman melissa_herman@unc.edu

ABSTRACT

Psilocybin, and its active metabolite psilocin, have been shown to elicit rapid and long-lasting symptom

improvements in a variety of affective psychiatric illnesses. However, the region-specific alterations underlying

these therapeutic effects remain relatively unknown. The central amygdala (CeA) is a primary output region

within the extended amygdala that is dysregulated in affective psychiatric disorders. Here, we measured CeA

activity using the activity marker c-Fos and CeA reactivity using fiber photometry paired with an aversive air-

puff stimulus. We found that psilocin administration acutely increased CeA activity in both males and females

and increased stimulus specific CeA reactivity in females, but not males. In contrast, psilocin produced time-

dependent decreases in reactivity in males, but not females as early as 2-days and lasting to 28-days post

administration. We also measured behavioral responses to the air-puff stimulus and found sex-dependent

changes in threat responding but not exploratory behavior or general locomotion. Repeated presentations of

the auditory component of the air-puff were also performed and sex-specific effects of psilocin on CeA

reactivity to the auditory-alone stimulus were also observed. This study provides new evidence that a single

dose of psilocin produces sex-specific, time-dependent, and enduring changes in CeA reactivity and behavioral

responding to specific components of an aversive stimulus.

INTRODUCTION

Clinical studies suggest that psilocybin, and active metabolite psilocin [1-3], produce rapid, long-lasting

improvements in psychiatric disorders including depression, anxiety, and substance use disorder [4-12].

Sustained improvements in symptoms were seen as long as 4 years following administration [13]. Research on

psychedelic compounds has focused primarily on human imaging studies and receptor pharmacology. Human

fMRI studies have shown that psychedelics produce robust alterations in network connectivity [8, 14-19].

Receptor studies have demonstrated that these compounds promote synaptogenesis [20-22] and identified the

serotonin 5-hydroxytryptamine (5-HT)$_{2A}$ receptor to be instrumental in mediating the hallucinogenic actions of

psychedelic compounds including psilocybin and lysergic acid diethylamide (LSD) [16, 23-28]. Despite the

observed clinical outcomes in psychedelic studies, the underlying neural correlates of these therapeutic effects

remain relatively unknown.

The central amygdala (CeA) is a primary output region within the extended amygdala that receives input

regarding internal and external arousal state and coordinates appropriate behavioral responses [29, 30]. CeA

dysregulation has been implicated in anxiety, depression, and PTSD [29, 31-36]. Preclinical studies

demonstrate that the psychedelic compounds LSD and 2,5-dimethoxy-4-iodoamphetamine (DOI) increase Fos

expression in the amygdala [37, 38] and that systemic and local DOI administration in the amygdala promotes

suppression of fear responses [38]. Psilocybin produces alterations in reactivity and connectivity of the

amygdala that correlate with positive therapeutic outcome [12, 15, 17, 39-41]. One clinical imaging study

reported significantly decreased amygdala reactivity 1-week following psilocybin administration, with lasting

changes in affect at one-month follow-up [41].

The current study utilized a preclinical approach to determine how the active compound, psilocin, alters CeA

activity, reactivity, and behavioral responding to an aversive stimulus in male and female Sprague Dawley rats.

Psilocin was used as it is the active metabolite of psilocybin in the central nervous system (CNS), mediating

effects of psilocybin at 5-HT$_{2AR}$ [2, 28]. We focused on the CeA as it's the primary output region of the

amygdala and as amygdala hyperreactivity is associated with many psychiatric disorders that psilocybin shows

promise in treating [42-45]. We first measured expression of c-Fos, an immediate early gene that is

upregulated following neuronal activity, in the CeA of male and female rats administered psilocin (2 mg/kg) or

vehicle. Next, we measured CeA reactivity using an air-puff stimulus involving a burst of air directed towards

the face. The air-puff stimulus has been validated as an aversive stimulus in passive avoidance tests [46] and,

due to known CeA involvement in pain modulation [47, 48], it was essential to utilize a stimulus that would

evoke a response without the addition of potential confounding effects of pain. Utilizing fiber photometry in

conjunction with the aversive air-puff stimulus, we were able to demonstrate a robust and reliable CeA

reactivity response. We then used this model to determine how administration of psilocin altered CeA reactivity

during acute exposure and at more prolonged timepoints. We observed sex-specific effects of psilocin on CeA

activity as measured by c-Fos expression and in-vivo reactivity and behavioral responding. Specifically, we

found that psilocin elicited acute increases in reactivity in females, but not males. Additionally, we found long-

term decreases in CeA reactivity in males that were not seen in females or in the vehicle control. To test

whether these effects would contribute to the development of a conditioned response to the sound of the air-

puff, additional assays were performed utilizing an auditory-only air-puff stimulus, wherein an air-puff was

administered with the nozzle facing outside of the box to isolate the auditory component of the air-puff. In doing

so, we show differential reactivity of the CeA suggesting that the observed changes were stimulus-specific and

dynamic in nature. Finally, we found that decrease in reactivity in males were primarily driven by animals that

exhibited darting vs, freezing behavior in response to the air puff stimulus.

METHODS

Animals/Stereotaxic Surgery

Adult male and female Sprague Dawley rats (200-400 g, ~7-weeks old at arrival, Envigo, Indianapolis, IN) were

group-housed in a humidity- and temperature-controlled (22°C) vivarium on a 12hr light/dark cycle with *ad*

libitum access to food and water. Bilateral infusions of pGP-AAV-syn-jGCaMP7f-WPRE (Addgene plasmid#

104488-AAV9) were performed at a rate of 100nl/minute into the CeA (AP:-2.0mm; ML:+/- 3.9mm; DV:-

8.0mm). Directly following injection, Doric 200um NA 0.37 silica optic fiber cannulas with a 9mm tip were

inserted at DV -7.7mm and secured in place using Metabond (Parkell, Brentwood, NY).

Following intracranial surgery at ~8 weeks of age, animals were single-housed in flat lid cages with waterspout

access, and food bowls to avoid damage to implants.

Drug Administration

On the day of injection, animals received a subcutaneous (s.c.) injection of either vehicle (0.9% Saline/2%

glacial acetic acid) or a 2 mg/kg psilocin (Cayman Chemical, graciously provided by Dr. Bryan Roth) dissolved

in 2% glacial acetic acid, as previously described [49]. Solutions were prepared fresh on the day of testing and

were stored at 4ºC for 30min prior to first injection. Injections were given at a 1mg/ml volume 40min prior to the

beginning of fiber photometry recordings. In the c-Fos cohort, injections were performed 2hr prior to perfusion.

For this study, a dose of 2 mg/kg was chosen as previous work has shown that administration of psilocin at 2

mg/kg in rats produced increases in BOLD signaling in the amygdala, an effect not seen when utilizing a lower

dose [50].

Fiber Photometry Recording

All animals were ~11 weeks of age at the beginning of testing. Animals were habituated to testing room for 1

hour prior to recording sessions. Rats were individually placed in a clear plexiglass chamber [50cm

(l)×50cm(w)×38cm(h)] in a red-light sound-attenuated behavioral cabinet and allowed to acclimate for 10min.

Fiber placement and GCaMP injection were performed bilaterally, however, recordings were collected

unilaterally. Baseline recordings were taken from both hemispheres and analyzed to determine which

hemisphere to use for the remainder of the experiment. Ca^{2+} signals from subjects, with 200µm optic fibers

placed .3mm above the injection site in the CeA, were recorded using a TDT RZ5 real time processor

equipped with the TDT Synapse program (Tucker-Davis Technologies, Alachua, FL). For each recording,

jGCaMP7f was excited using a 465nm calcium-dependent signal and a 405nm signal was used as an

isosbestic control. MATLAB script [51] was then used to analyze raw signal. Changes in fluorescence as a

function of baseline (ΔF/F) showing fluctuations in fluorescence compared to the overall baseline fluorescence

throughout the session were calculated through smoothing isosbestic control channel signal, scaling the

isosbestic signal by regressing it onto smoothed GCaMP signal, and then generating a predicted 405nm from

the regression to remove calcium-independent signaling from the raw GCaMP signal and control for artifact

from movement, photo-bleaching, and fiber bending artifacts. Baseline fluorescence was calculated through a

least-squares linear fitting of the 405nm isosbestic control to the 465nm GCaMP signal to create a resting

GCaMP signal over the entire session not containing event-dependent fluctuations in GCaMP signal [52]. Peri-

event plots were then generated by averaging changes in fluorescence (ΔF/F) across two different air-puff

trials to generate a mean change in reactivity [51]. For all photometry experiments, the same male

experimenter handled animals and recording. Additionally, female animals were not checked for estrus cycle to

avoid potential confounding effects of an added stressor. All subjects were video recorded during stimulus

presentations and behavior was scored using the Behavioral Observation Research Interactive Software

(BORIS) program [53]. Behavioral analysis was performed by an experimenter blind to the experimental status

of each animal.

Air-Puff Stimulus

An aversive air-puff stimulus was designed using house air supply (85psi) controlled by a Parker solenoid (Part

#003-0868-900), powered by a custom made 12V electrical circuit box connected to the TDT RZ5 system to

trigger 500ms openings of the solenoid and simultaneous timestamps through the Synapse program. Each

session consisted of a 10min habituation period before recording sessions that consisted of two air-puff

administrations with a 5min inter-stimulus interval (ISI) between them. Air-puffs were directed at the face and

within close proximity of the animal to maintain similar conditions and level of aversion to the stimulus between

subjects.

Perfusion/Tissue Collection

Rats were transcardially perfused with phosphate buffered saline (PBS) followed by 4% paraformaldehyde

(w/v). Brains were extracted, stored in 4% paraformaldehyde for 24h, then transferred to 30% sucrose (w/v)

and stored at 4ºC until sectioning. Using a freezing microtome, tissue was sliced into 40µm thick sections and

stored in cryoprotectant (30% v/v ethylene glycol + 30% w/v sucrose in phosphate buffered saline) at 4ºC until

use.

Immunofluorescence Staining

For site verification on the fiber photometry cohorts, 6 sections/subject (anterior-posterior (AP) coordinates -

1.75 to -2.75) from bregma were washed in phosphate buffered saline (PBS) prior to 1hr blocking in 5% normal

donkey serum + 0.3% Triton X in PBS. Sections were then incubated in chicken anti-GFP (Abcam 13970;

1:1000) in 0.5% normal donkey serum/0.3% Triton X/PBS overnight at 4ºC. Next, samples were triple-washed

and incubated in blocking solution (1hr) followed by goat anti-chicken 488 (Jackson 703-545-155; 1:700) in

0.5% normal donkey serum/0.3% Triton X/PBS for 2hr at room temperature.

For the c-Fos cohort, sections (AP coordinates -1.92mm to -2.16mm) were washed in PBS, incubated in 50%

methanol/PBS solution for 30min, washed in 3% hydrogen peroxide/PBS for 5min, placed in blocking solution

(0.3% Triton X-100; Thermo Fisher), and then 1% bovine serum albumin (BSA; Sigma) for 1hr, at room

temperature (RT). Slices were then incubated with rabbit anti-*cFos* (1:3000, Millipore Sigma; ABE457) for 24hr

at 4°C. Next, slices were washed with 0.1% Tween-20 in tris-buffered saline (TNT) before being transferred

into TNB blocking buffer (Perkin-Elmer FP1012) for 30min. After blocking, slices were incubated in goat anti-

rabbit horseradish peroxidase (HRP; 1:200, Abcam ab6721) for 2hr followed by another round of TNT washes.

Finally, slices were incubated in tyramide conjugated fluorescein (1:50) in TSA amplification diluent (Akoya

Biosciences, NEL741001KT) for 10min at RT. Slices were washed with TNT buffer, mounted, cover slipped

with Vectashield ® HardSet™ Antifade Mounting Medium with DAPI (H1500, Vector Laboratories, Burlingame,

CA) and stored at 4ºC before being imaged with the Keyence BZ-X800 fluorescence microscope. Images were

then analyzed by an experimenter blinded to the animal's condition. Images were taken at 20X and the BZ-

X800 Analyzer program was used to stitch images together. Randomly selected hemispheres were used, and

the area of interest was outlined and manually counted using the ImageJ [54] multi-point counter tool. Cell

counts were averaged per animal.

Statistical Analysis

Raw data from the TDT RZ5 machine were imported and processed for artifact removal, down sampling, and

detrending using MATLAB code [51]. Changes in fluorescence ($\Delta F/F$) were then calculated and collected 5s

before and 10s following air-puff administration. The mean of two traces were averaged together for each

subject and combined using custom MATLAB script (https://doi.org/10.5061/dryad.pnvx0k6q9) to sort

individual data into groups. Group 465nm signal data were then converted into 500ms bins for analysis to

quantify differences between experimental conditions. All binned trace plots were normalized by subtracting

the mean of the signal pre-air-puff from each time point in order to correct for differences in basal activity prior

to air puff administration. All comparisons were made using time bins post air puff, corresponding to 0-20 on

the x-axis of binned analysis plots. All statistical analyses were conducted using GraphPad Prism 7. Five

subjects were removed from the study due to lack of signal/cannula removal. Repeated-measures one-way

analysis of variance (ANOVA) with Dunnett's post hoc test was used to determine differences between

baseline and follow-up time points within each group in the air-puff condition. Two-way ANOVAs with Šidák's

post hoc tests were used to compare time bins between groups and time points. Paired t-tests were used to

determine differences within groups between time points in the binned trace and peak point. Fisher's exact

tests were used to compare percentages in darting behavior. Mixed-effects two-way ANOVA were used to

compare time spent walking and rearing. All data including Matlab structures and scripts can be found at

https://doi.org/10.5061/dryad.pnvx0k6q9.

RESULTS

Acute Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus

At ~8 weeks, male and female Sprague Dawley rats received bilateral injection of pGP-AAV-syn-jGCaMP7f-

WPRE into the CeA immediately followed by installation of fiber optic cannulae above the injection site (**Fig.**

1A-B). Following 3-weeks for virus transduction, animals underwent a series of fiber photometry experiments

testing the effects of psilocin on fluctuations in calcium dynamics within the CeA time-locked to an aversive air-

puff stimulus (**Fig. 1A**). For both baseline and injection recording sessions, each animal received two air-puffs

with an ISI of 5min and trace plots were generated showing change in fluorescence as a function of baseline

($\Delta F/F$). Traces were analyzed 5s before and 10s following administration of air-puff (**Fig. 1C**). Following

injection of the drug, animals were not assessed for any behavioral indices of hallucination (i.e. head twitch,

wetback shake).

Sub-Region-Specific Differences in CeA Basal Activity Following Psilocin Administration

To assess the effects of psilocin on basal activity in the CeA, male and female Sprague Dawley rats were

injected with psilocin (2 mg/kg, s.c.) or vehicle (1 ml/kg, s.c.) 2hr prior to transcardial perfusion.

Immunohistochemistry (IHC) assays were performed using CeA tissue collected within the anterior-posterior

bregma range of -1.92mm to -2.16mm. In females, there was a significantly increased c-Fos expression in the

psilocin group as compared to vehicle across regions (2-way ANOVA: $F_{Interaction}(2,24)=1.597$, $p=0.22$;

$F_{Region}(1,24)=30.84$, p<0.0001; $F_{injection}(1,24)=9.149$ $p=0.006$; **Fig.2A-B**) with greater increases in the capsular

division of the central amygdala (CeC) (Šidák's: $p=0.02$). In males, there was also significantly increased c-Fos

expression in the psilocin group across regions (2-way ANOVA: $F_{Interaction}(2,27)=1.241$, $p=0.30$;

$F_{Region}(2,27)=12.02$, p=0.0002; $F_{injection}(1,27)=5.543$, $p=0.02$; **Fig.2C-D**).

Acute Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus Following Psilocin

Administration

To test the acute effects of psilocin on CeA reactivity, we measured changes in CeA reactivity during time-

locked stimuli presentation at baseline and 40min post administration of vehicle or psilocin (2 mg/kg, s.c.).

Following a 10min habituation period, air-puffs were delivered as described above. Mean traces were taken

from two separate air-puff responses within a session (**Fig. 2E, 2I**). These traces were then collapsed into

500ms bins for statistical analysis (**Fig. 2G, 2H, 2K, 2L**). To compare baseline to injection, paired t-tests were

performed comparing the individual time bins following the air-puff collapsed across subjects (Fig. **2F**, **2J**). In

the female vehicle control group (n=9), there were no changes between baseline and injection (t(19)=1.556,

p=0.13; **Fig.2F-G**). In the female psilocin group (n=7), there was a significant increase between baseline and

injection (t(19)=12.38, p<0.0001; **Fig.2F, 2H**). In the male vehicle control group (n=10), there was an increase

in reactivity between baseline and injection (t(19)=7.841, p<0.0001; **Fig.2J-K**). In the male psilocin group

(n=11), there was no significant difference between baseline and injection (t(19)=1.372, p=0.18; **Fig.2J, 2L**).

Prolonged Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus Following Psilocin

Administration

To assess long-term alterations in CeA reactivity following a single dose of psilocin, fiber photometry

recordings were performed on days 2, 6, and 28 post-administration of vehicle or psilocin. On each follow-up

recording day, animals received two air-puffs with a 5-minute ISI. Given the interest in long-term effects of

psychedelics, a comparison between baseline and the 28-day follow-up was made. In the vehicle control

females, there was a significant increase in reactivity at the 28-day follow-up compared to baseline

(t(19)=4.762, p=0.0001; **Fig.3B**). In the psilocin treated females, there was also an increase in reactivity seen

at the 28-day follow-up (t(19)=2.832, p=0.01; **Fig.3B**), though the effect was less significant. Next, we

compared all the follow-up time points to baseline. In the vehicle control females there was an effect of

injection (F(1.558,29.61)=29.32, p<0.0001; **Fig.3C**). Dunnett's post hoc analysis revealed a significant increase

in reactivity at the 2-day (p<0.0001) and 28-day follow-up (p=0.0005). In the female psilocin group (**Fig. 3A,**

3D), there was an effect of injection (F(3.178,60.38)=50.25, p<0.0001; **Fig.3D**). Dunnett's post hoc found

significant increases in reactivity at the 2-day (p=0.01), 6-day (p<0.0001), and 28-day follow-up (p=0.03).

In the male vehicle control group, there was no difference between baseline and the 28-day follow-up

(t(19)=0.07061, p=0.94; **Fig.3F**). In the male psilocin group, there was a significant decrease in reactivity seen

at the 28-day follow-up compared to baseline (t(19)=7.234, p<0.0001; **Fig.3F**). Comparing all follow-up time

points to baseline, in the male vehicle group there was an effect of injection ($F(1.797,34.14)=14.48$, $p<0.0001$;

Fig.3G). Dunnett's post hoc showed a significant increase in reactivity at the 2-day follow-up ($p<0.0001$). In the

male psilocin group, there was a significant effect of injection ($F(2.163,41.10)=63.78$, $p<0.0001$; **Fig.3H**).

Dunnett's post hoc analysis revealed significant decreases in reactivity at the 2-day, 6-day, and 28-day follow-

up ($p<0.0001$).

To compare changes in reactivity between vehicle and psilocin groups, 2-way ANOVA revealed differential

effects of injection across time points between groups in females ($F_{Interaction}(4,152)=53.46$, $p<0.0001$;

$F_{Injection}(1,38)=2.015$, $p=0.16$; $F_{Time}(2.343,89.04)=18.82$, $p<0.0001$; **Fig.3J**) and males ($F_{Interaction}(4,152)=23.96$,

$p<0.0001$; $F_{Injection}(1,38)=2.310$, $p=0.13$, $F_{Time}(2.603,98.90)=50.45$, $p<0.0001$; **Fig.3I**). Šidák post hoc analysis

revealed significant increases in reactivity during injection ($p<0.0001$) and at the 2-day follow-up ($p=0.005$) in

the female psilocin group compared to vehicle control (**Fig.3J**). In males, there was a significant decrease in

reactivity at the 2-day follow-up in the psilocin group compared to vehicle control ($p=0.003$; **Fig.3J**)

Acute and Prolonged Behavioral Effects of Psilocin

To determine if psilocin elicited changes in locomotion and exploratory behavior, video recordings were scored

for total time spent walking and time spent rearing, respectively. All behavior was scored during the 5min ITI

between the first and second air-puff. Rearing was defined as standing on the two hindlegs, regardless of any

balancing on the walls of the box. In females, there were no differences in overall locomotion between vehicle

and psilocin groups ($F_{Interaction}(3,42)=0.2118$, $p=0.89$; $F_{Injection}(1,14)=2.238$, $p=0.16$; $F_{Time}(2.497,34.95)=0.6065$,

$p=0.59$; **Fig.4A**). There were also no differences in rearing between female vehicle and psilocin groups

($F_{Interaction}(4,56)=1.345$, $p=0.26$; $F_{Injection}(1,14)=4.433$, $p=0.05$; $F_{Time}(4,56)=2.486$, $p=0.05$; **Fig. 4B**). In the males,

there were no differences in locomotion ($F_{Interaction}(3,57)=0.1054$, $p=0.96$; $F_{Injection}(1,19)=2.563$, $p=0.12$;

$F_{Time}(3,57)=1.255$, $p=0.29$; **Fig.4C**) or rearing ($F_{Interaction}(4,76)=0.5546$, $p=0.69$; $F_{Injection}(1, 19)=0.7576$, $p=0.39$;

$F_{Time}(4,76)=1.190$, $p=0.32$; **Fig.4D**) between vehicle and psilocin groups. Collectively, no significant alterations

in locomotion were seen resulting from psilocin administration (**Fig.4A, 4C**), although females appeared to

display increased locomotion overall as compared to males, regardless of injection type.

To probe potential differences in threat responding to the air-puff stimulus, we separated groups based on

active or passive coping strategy. Subgroups were created containing animals that remained immobile

(passive) following air-puff administration versus animals that exhibited darting behavior (active) following air-

puff administration. Consistent with previous literature [55], females primarily employed an active darting

response (**Fig.4E**), while active vs. passive coping behavior was more evenly split in males (**Fig.4F**). In

females, there were no differences in the percentage of animals that darted or that stayed immobile between

groups at any of the time points (**Fig. 4E**). In contrast, the male psilocin group displayed nonsignificant

increases in darting and reductions in immobility in the vehicle control group that were not seen in the psilocin

group under injection conditions (Fisher's exact: $p=0.08$; **Fig. 4F**). At the 2-day follow-up, there was a

significantly higher percentage of darting in the vehicle control group compared to the psilocin group ($p=0.03$;

Fig.4F, left) and a significantly higher percentage immobile in the psilocin group (Fisher's exact: $p=0.03$;

Fig.4F, right). Though differences were not significant at every time point, psilocin seemed to prevent a

trending increase in darting behavior seen in the male vehicle control group and increase in overall immobility

in male but not female rats (**Fig.4F**).

To explore the CeA reactivity correlates to the observed behavioral differences in threat responding, fiber

photometry traces were split into the corresponding active vs. passive responder groups. Since females

primarily employed active coping strategies that were not affected by psilocin, we only examined CeA reactivity

between the male active and passive responders. To assess changes in the amplitude of CeA response to the

stimulus, comparisons looking at the peak point, or highest $\Delta F/F$ value following air-puff, were made. In the

male passive responder, there were no significant differences in peak point in the vehicle ($F(4,20)=1.019$,

$p=0.42$; **Fig.4G**) or psilocin group ($F(4,20)=1.278$, $p=0.31$; **Fig.4H**). In the active responder vehicle males, no

differences were seen in peak point compared to baseline ($F(4,12)=1.959$, $p=0.16$; **Fig.4I**). However, in the

active responder psilocin males, there was a main effect of injection ($F(4,16)=6.398$, $p=0.002$; **Fig.4J**) with a

significant decrease in peak point at day 2 compared to baseline ($p=0.003$), suggesting that psilocin-induced

decreases in the amplitude of CeA reactivity were more pronounced in males employing an active vs. passive

threat response at baseline.

Changes in CeA Reactivity in Response to an Auditory Stimulus Following Psilocin Administration

To test whether the animals developed a conditioned response to the tone associated with the air-puff

stimulus, and whether psilocin treatment altered the response to the auditory stimulus alone, two separate

auditory-only sessions were conducted immediately following baseline recordings and again at the 28-day

recording session. These auditory only sessions followed the same parameters as the air-puff with the

exceptions being that the ISI was reduced to 3min and the air-puff apparatus was facing outside of the box to

maintain the same auditory tone without the physical sensation of the air-puff.

In vehicle control females (**Fig.5A**), there was a significant decrease in peak point reactivity from the initial to

the final auditory condition (t(8)=2.806, *p*=0.02; **Fig.5B**) and a significant decrease in binned analysis from the

initial auditory to the final auditory condition (t(29)=6.727, *p*<0.0001; **Fig.5C-D**). In psilocin females (**Fig.5E**),

there were no significant differences between the initial and final auditory session in either peak point

(t(6)=0.5642, *p*=0.59; **Fig.5F)** or binned analysis (t(29)=0.09698, *p*=0.92; **Fig.5G-H**).

In the vehicle males (**Fig.6A**), there weren't any differences in peak point (t(9)=0.8811, *p*=0.40; **Fig. 6B**) or

binned analysis from the initial auditory to the final auditory condition (t(19)=0.6978, *p*=0.49; **Fig.6C-D**). In the

psilocin-treated males (**Fig.6E**), there was no difference in peak point from the initial to the final auditory

session (t(10)=0.5758, *p*=0.57; **Fig.6F**). However, there was a significant decrease in binned analysis of the

initial auditory and final auditory condition (t(19)=7.327, *p*<0.0001; **Fig.6G-H**). While there was a decrease

seen in the psilocin group binned analysis, there was no decrease in the amplitude of response (**Fig.6B, 6F**).

DISCUSSION

The present study explored the effects of the psychedelic compound psilocin on CeA activity using an

immunohistological activity marker and on CeA reactivity to an aversive stimulus using *in-vivo* fiber photometry

recordings during acute drug exposure and at 2- , 6- , and 28-days post drug administration. The study

examined multi-level neurobiological and behavioral effects of psilocin by looking at acute drug effects, long-

term alterations in activity following drug administration, and by probing for potential sex-specific effects of both

injection and time. Psilocin administration produced increases in c-Fos expression in both males and females

and sub-region analysis showed that increases in the female CeC were significantly greater in psilocin

compared to the vehicle control (**Fig. 2B**). CeA reactivity in response to the air-puff stimulus following psilocin

administration also displayed sex- and time-specific effects. In females, psilocin increased CeA reactivity

acutely as compared to vehicle controls (**Fig. 2E-H**). Interestingly, both vehicle and psilocin groups showed

increases in reactivity at specific follow-up timepoints compared to baseline. In the vehicle control females,

there were large increases in the duration of CeA response at the 2- and 28-day follow-up. In the psilocin

treated females, there were small increases at the 2- and 28-day follow-up, with a much larger response seen

at the 6-day follow-up. In contrast, the male psilocin group did not show the same increased acute reactivity

seen in the females (**Fig. 2H**) and instead displayed decreases in CeA reactivity at 2-day, 6-day, and 28-day

follow-up recordings (**Fig. 3F, 3H**) that were not seen in vehicle control males (**Fig. 3G**). These long-term

reductions in reactivity seen in males could be due to plasticity dependent changes in innervating circuitry.

Future studies will investigate a potential upstream mediator of these observed effects.

In this study, we have replicated previous work demonstrating an increase in basal activity following

administration of a 5-HT_{2AR} agonist psychedelic [37]. This could in part be due to local 5-HT_{2AR} excitatory

activation in the CeA. However, the CeA contains a high proportion of 5-HT_{1AR}, a presynaptic receptor with

inhibitory mechanisms [56]. Another potential explanation lies in more of a circuit/systems level effect of the

drug. For instance, the dorsal raphe nucleus is a region within the brainstem that is involved in serotonin

synthesis that also has a high density of 5-HT_{1AR}. Previous work has shown that administration of psychedelics

produces a local decrease in activity within the DRN that is likely driven by 5-HT_{1AR} [57]. Additionally, it has

been shown that there are dense projections of serotonergic neurons from the DRN to the locus coeruleus

(LC) [58], and that serotonin release within the LC produces a tonic inhibition of LC neurons [59]. Thus, given

previously reported noradrenergic projections from LC to CeA [60], it is possible that disinhibition of the LC

through inhibition of the DRN via activation of DRN 5-HT_{1ARs} could result in increased basal activity within the

CeA. Future work will look at potential circuit-based mediators of both the observed changes in basal activity

and changes in stimulus reactivity within the CeA.

Clinical studies have illustrated that psilocybin produces alterations in amygdala reactivity and connectivity [8,

12, 15, 17, 39-41]. Here, we utilized a preclinical approach to examine the actions of psilocin, the psychoactive

metabolite of psilocybin, on CeA activity in a rodent model. As most clinical studies report a reduction in

amygdala reactivity or connectivity that correlated with therapeutic improvements [8, 15, 17, 39-41], we

hypothesized that psilocin would produce similar decreases in reactivity in our model. Here we found that a

single dose of psilocin produced persistent reductions in CeA reactivity to an aversive stimulus in male but not

female rats. These findings lend to recent studies in both clinical [11, 13, 61] and preclinical [62] literature

showing persistent reductions in depressive behavior after a single dose of psilocybin. However, given the

increases seen in the female group, these data do not perfectly align with clinical findings with samples that

include representation from female participants. One potential explanation could be that human fMRI lacks the

resolution needed to truly parse apart amygdalar sub-nuclei activation. It is possible that effects seen in the

CeA would not match effects seen in the basolateral amygdala (BLA). Additionally, the amygdala has been

shown to be involved in both positive and negative valence, and therefore functioning within this region seems

to be very dynamic and stimulus-dependent. A surprise air puff to a rat may be an entirely different experience

subjectively than seeing an emotional face, and therefore could result in far different functioning within the

amygdala complex. Additional experiments comparing drug effects on whole brain connectivity/activity will be

the addressed in future studies. While the specific mechanism behind these therapeutic effects remains

unknown, this study provides evidence that changes in CeA activity may play a role in observed therapeutic

effects.

Lateralization of amygdala response to fear processing has been noted in both preclinical [63] and clinical

literature [64, 65], however there are contradictions in these data and therefore no clear, resolute consensus.

For instance, one study in human imaging showed that males had greater activation in the right amygdala as

compared to females during an emotional face perception fMRI task utilizing angry facial expressions [64].

Contrarily, another human imaging study employing a similar task showed that amygdala activation differed

based on the valence of the stimuli. Additionally, while males had more laterality, both male and female

subjects showed greater activation in the left amygdala to fearful faces [65]. This is particularly relevant to the

current study given that recordings were taken from a single hemisphere. Initially, hemispheres were chosen

based on consistency and quality of signal. Within each group for both males and females, there is

representation from both left and right hemispheres. While this does offer some protection from potentially

confounding effects of amygdala lateralization. Future work will be conducted to determine if there is an effect

of amygdala lateralization on psilocin-induced changes in reactivity.

We report sex-specific effects of psilocin on basal activity and stimulus-induced reactivity within the CeA.

Changes in basal activity were measured by c-Fos, an immediate early gene shown to be upregulated

following neuronal activation [66]. We observed an increase in CeA c-Fos expression in both males and

females following administration of psilocin, with region specific increases found in the CeC subregion in

females. CeA reactivity was assessed using fiber photometry to measure in-vivo stimulus-induced fluctuations

in calcium signaling. In the female psilocin group, we observed an acute increase in reactivity to the air-puff

stimulus while on the drug that was not seen in the males. Additionally, we saw increases in reactivity at all of

the follow-up recording days, while we saw reductions in CeA reactivity in the males on follow-up recording

days (**Fig. 3B, 3D, 3E, 3G**). A recent study looking at cerebral blood flow and the expression of the serotonin

transporter (5-HTT) showed that, in response to an aversive predator odor, there were increases in blood flow

and c-Fos expression in the CeA that were related to 5-HTT expression in males but not females [67]. These

findings suggest potential sex-specific expression of serotonin receptors within the amygdala. Given the affinity

of psilocin for the serotonin system, differential expression of serotonin receptors within the CeA or surrounding

nuclei could explain the sex-specific effects seen in this study. Additionally, a previous study showed sex-

specific differences in 5-HT$_{2AR}$ expression at baseline, wherein females had lower expression in the

orbitofrontal cortex (OFC) than males [68]. Another potential explanation for the differential effects of psilocin

between males and females may be due to differences in CeA circuitry. For example, the OFC has been

shown, in rodents and non-human primates, to have direct connections to the CeA as well as to the

intercalated nuclei, a local GABAergic population within the amygdala complex [69, 70]. Lack of activation of

this circuit due to a reduced expression of OFC 5-HT$_{2AR}$ in females could prevent an inhibition of CeA firing,

thus causing the increase in reactivity seen while on psilocin in females but not males. Future experiments are

needed to examine the effects of psilocin on CeA circuits innervating distinct brain regions.

In the presence of a threatening stimulus, animals will employ either active or passive coping strategies. In this

study, the active threat response was characterized by darting, or immediate fleeing from the air-puff stimulus,

while the passive threat response was characterized by immobility following the air-puff. Here, we show that

females predominantly exhibited an active response, while males were more evenly split between active and

passive responses. In male psilocin-treated animals who exhibited an active response at baseline, there was a

significant decrease in the amplitude of CeA reactivity following exposure to the air-puff at the 2-day follow-up,

with no changes in vehicle control. This decrease in reactivity was not seen in the passive-responder vehicle or

psilocin groups. While further interpretation regarding adaptive vs. maladaptive strategies in this context is

outside the scope of the current study, these results suggest that there may be differences in CeA reactivity

and threat responding strategy that are implicated in the efficacy of psychedelics.

To determine the effects of psilocin on associative learning, we utilized an auditory-only stimulus recording

taken after the initial baseline recordings and then again after the 28-day recording. Interestingly, in females

the vehicle control group displayed reductions in CeA response magnitude to the auditory-only stimulus while

reactivity in the males did not change. In the psilocin group however, reductions in CeA reactivity to the

auditory stimulus were not observed in females, while in males, reductions were seen in binned analysis only

with no changes seen in the magnitude of response (**Fig. 6**). Compared to the changes in reactivity seen with

the standard air-puff stimulus, the changes in reactivity to the auditory stimulus were contradictory in females

and less pronounced in males. These data suggest a more dynamic alteration in CeA reactivity that is stimulus

specific. One potential interpretation of the sustained activation in the psilocin females vs. vehicle control is that

psilocin may influence associative learning to an aversive stimulus. Several studies have shown that $5-HT_{2A}$

agonist psychedelics produce improvements in various dimensions of cognition, including associative learning

[71-73]. Another potential interpretation of these findings is that psilocin created a more robust/meaningful

memory of the air-puff. Many of the clinical studies cite participants rating their experience during a psilocybin

treatment as one of the most meaningful experiences of their life [13, 74, 75]. Additionally, increases in

plasticity could contribute to differences in memory reconsolidation during air-puff administration while on the

drug.

Our findings demonstrate that the psychedelic compound, psilocin, produces dynamic, sex-specific changes in

the CeA, a key nucleus within the amygdala complex. Here we showed that administration of psilocin produces

increases in CeA c-Fos expression and that, in females, these increases were primarily driven by increases in

expression in the CeC. We have also shown that there are sex-specific changes in CeA reactivity while on

psilocin. Specifically, we show that a single administration of psilocin produced acute increases in CeA

reactivity in females, but not males. Additionally, we show that persistent reductions in CeA reactivity are seen

as early as 2 and as long as 28 days following administration of the drug in males, but not females. Given that

dysregulation of the amygdala is a hallmark in many different psychiatric disorders, including anxiety, MDD,

and PTSD [29, 31-36] and that psychedelic compounds show promise in treating many of these illnesses [4, 6-

11], these findings provide important information on subregion-specific alterations in CeA function that may

play a part in observed therapeutic effects of psychedelics. One limitation of this work is that no behavioral

indices of hallucination (head twitch, wet back shakes, etc.) were collected following administration of the drug.

Therefore, no claims can be made regarding the effects of hallucinations on amygdala activity following

psychedelic drug exposure. Additionally, without the use of antagonists or known 5-HT_{2AR} mediated behavioral

effects, we cannot claim any sort of cell type specificity in our findings. Future work will be required to explore

potential circuit-based mediators contributing to these observed alterations in CeA reactivity, but these findings

provide important evidence for CeA alterations underlying the potential therapeutic effects of psilocin and

psilocybin.

ACKNOWLEDGEMENTS

This work was supported by National Institute of Health grants AA026858 (M.A.H.), GM135095 (D.P.E.),

AA007573 (S.G.Q.), and by the Brain and Behavior Research Foundation NARSAD Young Investigator Award

(M.A.H.)

<u>**COMPETING INTERESTS**</u>

The authors have no competing interests to report.

REFERENCES

1. Nichols, D.E., *Hallucinogens.* Pharmacology & therapeutics, 2004. **101**(2): p. 131-181.
2. Dinis-Oliveira, R.J., *Metabolism of psilocybin and psilocin: clinical and forensic toxicological relevance.* Drug metabolism reviews, 2017. **49**(1): p. 84-91.
3. Passie, T., et al., *The pharmacology of psilocybin.* Addiction biology, 2002. **7**(4): p. 357-364.
4. Davis, A.K., et al., *Effects of psilocybin-assisted therapy on major depressive disorder: a randomized clinical trial.* JAMA psychiatry, 2021. **78**(5): p. 481-489.
5. Bogenschutz, M.P. and M.W. Johnson, *Classic hallucinogens in the treatment of addictions.* Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2016. **64**: p. 250-258.
6. Johnson, M.W., A. Garcia-Romeu, and R.R. Griffiths, *Long-term follow-up of psilocybin-facilitated smoking cessation.* The American journal of drug and alcohol abuse, 2017. **43**(1): p. 55-60.
7. Johnson, M.W. and R.R. Griffiths, *Potential therapeutic effects of psilocybin.* Neurotherapeutics, 2017. **14**(3): p. 734-740.
8. Carhart-Harris, R.L., et al., *Psilocybin for treatment-resistant depression: fMRI-measured brain mechanisms.* Scientific reports, 2017. **7**(1): p. 1-11.
9. Griffiths, R.R., et al., *Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: A randomized double-blind trial.* Journal of psychopharmacology, 2016. **30**(12): p. 1181-1197.
10. Bogenschutz, M.P., et al., *Psilocybin-assisted treatment for alcohol dependence: a proof-of-concept study.* Journal of psychopharmacology, 2015. **29**(3): p. 289-299.
11. Ross, S., et al., *Rapid and sustained symptom reduction following psilocybin treatment for anxiety and depression in patients with life-threatening cancer: a randomized controlled trial.* Journal of psychopharmacology, 2016. **30**(12): p. 1165-1180.
12. Roseman, L., et al., *Increased amygdala responses to emotional faces after psilocybin for treatment-resistant depression.* Neuropharmacology, 2018. **142**: p. 263-269.
13. Agin-Liebes, G.I., et al., *Long-term follow-up of psilocybin-assisted psychotherapy for psychiatric and existential distress in patients with life-threatening cancer.* Journal of Psychopharmacology, 2020. **34**(2): p. 155-166.
14. Barrett, F.S., et al., *Psilocybin acutely alters the functional connectivity of the claustrum with brain networks that support perception, memory, and attention.* NeuroImage, 2020. **218**: p. 116980.
15. Grimm, O., et al., *Psilocybin modulates functional connectivity of the amygdala during emotional face discrimination.* Eur Neuropsychopharmacol, 2018. **28**(6): p. 691-700 DOI: 10.1016/j.euroneuro.2018.03.016.
16. Preller, K.H., et al., *Psilocybin induces time-dependent changes in global functional connectivity.* Biological psychiatry, 2020. **88**(2): p. 197-207.
17. Mertens, L.J., et al., *Therapeutic mechanisms of psilocybin: changes in amygdala and prefrontal functional connectivity during emotional processing after psilocybin for treatment-resistant depression.* Journal of Psychopharmacology, 2020. **34**(2): p. 167-180.
18. Roseman, L., et al., *The effects of psilocybin and MDMA on between-network resting state functional connectivity in healthy volunteers.* Frontiers in human neuroscience, 2014. **8**: p. 204.
19. Carhart-Harris, R.L., et al., *Neural correlates of the psychedelic state as determined by fMRI studies with psilocybin.* Proceedings of the National Academy of Sciences, 2012. **109**(6): p. 2138-2143.
20. Shao, L.-X., et al., *Psilocybin induces rapid and persistent growth of dendritic spines in frontal cortex in vivo.* Neuron, 2021. **109**(16): p. 2535-2544. e4.

21. Jones, K.A., et al., *Rapid modulation of spine morphology by the 5-HT2A serotonin receptor through kalirin-7 signaling.* Proceedings of the National Academy of Sciences, 2009. **106**(46): p. 19575-19580.

22. Ly, C., et al., *Psychedelics promote structural and functional neural plasticity.* Cell reports, 2018. **23**(11): p. 3170-3182.

23. Lee, H.-M. and B.L. Roth, *Hallucinogen actions on human brain revealed.* Proceedings of the National Academy of Sciences, 2012. **109**(6): p. 1820-1821.

24. Kometer, M., et al., *Activation of serotonin 2A receptors underlies the psilocybin-induced effects on α oscillations, N170 visual-evoked potentials, and visual hallucinations.* Journal of Neuroscience, 2013. **33**(25): p. 10544-10551.

25. Kometer, M., et al., *Psilocybin biases facial recognition, goal-directed behavior, and mood state toward positive relative to negative emotions through different serotonergic subreceptors.* Biological psychiatry, 2012. **72**(11): p. 898-906.

26. Halberstadt, A.L., et al., *Differential contributions of serotonin receptors to the behavioral effects of indoleamine hallucinogens in mice.* Journal of psychopharmacology, 2011. **25**(11): p. 1548-1561.

27. Halberstadt, A.L., S.B. Powell, and M.A. Geyer, *Role of the 5-HT2A receptor in the locomotor hyperactivity produced by phenylalkylamine hallucinogens in mice.* Neuropharmacology, 2013. **70**: p. 218-227.

28. Madsen, M.K., et al., *Psychedelic effects of psilocybin correlate with serotonin 2A receptor occupancy and plasma psilocin levels.* Neuropsychopharmacology, 2019. **44**(7): p. 1328-1334 DOI: 10.1038/s41386-019-0324-9.

29. Gilpin, N.W., M.A. Herman, and M. Roberto, *The central amygdala as an integrative hub for anxiety and alcohol use disorders.* Biological psychiatry, 2015. **77**(10): p. 859-869.

30. Ciocchi, S., et al., *Encoding of conditioned fear in central amygdala inhibitory circuits.* Nature, 2010. **468**(7321): p. 277-282.

31. Sandu, A.L., et al., *Amygdala and regional volumes in treatment-resistant versus nontreatment-resistant depression patients.* Depression and anxiety, 2017. **34**(11): p. 1065-1071.

32. Armony, J.L., et al., *Amygdala response in patients with acute PTSD to masked and unmasked emotional facial expressions.* American Journal of Psychiatry, 2005. **162**(10): p. 1961-1963.

33. Frodl, T., et al., *Enlargement of the amygdala in patients with a first episode of major depression.* Biological Psychiatry, 2002. **51**(9): p. 708-714 DOI: https://doi.org/10.1016/S0006-3223(01)01359-2.

34. Rauch, S.L., et al., *Exaggerated amygdala response to masked facial stimuli in posttraumatic stress disorder: a functional MRI study.* Biological psychiatry, 2000. **47**(9): p. 769-776.

35. Semple, W.E., et al., *Higher brain blood flow at amygdala and lower frontal cortex blood flow in PTSD patients with comorbid cocaine and alcohol abuse compared with normals.* Psychiatry, 2000. **63**(1): p. 65-74.

36. Siegle, G.J., et al., *Increased Amygdala and Decreased Dorsolateral Prefrontal BOLD Responses in Unipolar Depression: Related and Independent Features.* Biological Psychiatry, 2007. **61**(2): p. 198-209 DOI: https://doi.org/10.1016/j.biopsych.2006.05.048.

37. Gresch, P., L. Strickland, and E. Sanders-Bush, *Lysergic acid diethylamide-induced Fos expression in rat brain: role of serotonin-2A receptors.* Neuroscience, 2002. **114**(3): p. 707-713.

38. Pędzich, B.D., et al., *Effects of a psychedelic 5-HT2A receptor agonist on anxiety-related behavior and fear processing in mice.* Neuropsychopharmacology, 2022: p. 1-11.

39. Kraehenmann, R., et al., *Psilocybin-Induced Decrease in Amygdala Reactivity Correlates with Enhanced Positive Mood in Healthy Volunteers.* Biol Psychiatry, 2015. **78**(8): p. 572-81 DOI: 10.1016/j.biopsych.2014.04.010.

40. Kraehenmann, R., et al., *The mixed serotonin receptor agonist psilocybin reduces threat-induced modulation of amygdala connectivity.* NeuroImage: Clinical, 2016. **11**: p. 53-60.

41. Barrett, F.S., et al., *Emotions and brain function are altered up to one month after a single high dose of psilocybin.* Scientific Reports, 2020. **10**(1): p. 1-14.

42. Bas-Hoogendam, J.M., et al., *Amygdala hyperreactivity to faces conditioned with a social-evaluative meaning–a multiplex, multigenerational fMRI study on social anxiety endophenotypes.* NeuroImage: Clinical, 2020. **26**: p. 102247.

43. Donegan, N.H., et al., *Amygdala hyperreactivity in borderline personality disorder: implications for emotional dysregulation.* Biological psychiatry, 2003. **54**(11): p. 1284-1293.

44. Yang, T.T., et al., *Adolescents with major depression demonstrate increased amygdala activation.* Journal of the American Academy of Child & Adolescent Psychiatry, 2010. **49**(1): p. 42-51.

45. Joos, A.A., et al., *Amygdala hyperreactivity in restrictive anorexia nervosa.* Psychiatry Research: Neuroimaging, 2011. **191**(3): p. 189-195.

46. Moriarty, O., et al., *Validation of an air-puff passive-avoidance paradigm for assessment of aversive learning and memory in rat models of chronic pain.* Journal of neuroscience methods, 2012. **204**(1): p. 1-8.

47. Veinante, P., I. Yalcin, and M. Barrot, *The amygdala between sensation and affect: a role in pain.* Journal of molecular psychiatry, 2013. **1**(1): p. 1-14.

48. Wilson, T.D., et al., *Dual and opposing functions of the central amygdala in the modulation of pain.* Cell reports, 2019. **29**(2): p. 332-346. e5.

49. Horsley, R.R., et al., *Psilocin and ketamine microdosing: effects of subchronic intermittent microdoses in the elevated plus-maze in male Wistar rats.* Behavioural pharmacology, 2018. **29**(6): p. 530-536.

50. Spain, A., et al., *Neurovascular and neuroimaging effects of the hallucinogenic serotonin receptor agonist psilocin in the rat brain.* Neuropharmacology, 2015. **99**: p. 210-220.

51. Barker, D.J., et al., *Lateral preoptic control of the lateral habenula through convergent glutamate and GABA transmission.* Cell reports, 2017. **21**(7): p. 1757-1769.

52. Lerner, T.N., et al., *Intact-brain analyses reveal distinct information carried by SNc dopamine subcircuits.* Cell, 2015. **162**(3): p. 635-647.

53. Friard, O. and M. Gamba, *BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations.* Methods in ecology and evolution, 2016. **7**(11): p. 1325-1330.

54. Schneider, C.A., W.S. Rasband, and K.W. Eliceiri, *NIH Image to ImageJ: 25 years of image analysis.* Nature methods, 2012. **9**(7): p. 671-675.

55. Gruene, T.M., et al., *Sexually divergent expression of active and passive conditioned fear responses in rats.* Elife, 2015. **4**: p. e11352.

56. Albert, P.R. and F. Vahid-Ansari, *The 5-HT1A receptor: signaling to behavior.* Biochimie, 2019. **161**: p. 34-45.

57. Sprouse, J.S. and G.K. Aghajanian, *Electrophysiological responses of serotoninergic dorsal raphe neurons to 5-HT1A and 5-HT1B agonists.* Synapse, 1987. **1**(1): p. 3-9.

58. Morgane, P.J. and M. Jacobs, *Raphe projections to the locus coeruleus in the rat.* Brain research bulletin, 1979. **4**(4): p. 519-534.

59. Kaehler, S.T., N. Singewald, and A. Philippu, *Dependence of serotonin release in the locus coeruleus on dorsal raphe neuronal activity.* Naunyn-Schmiedeberg's archives of pharmacology, 1999. **359**(5): p. 386-393.

60. Gu, Y., et al., *A brainstem-central amygdala circuit underlies defensive responses to learned threats.* Molecular psychiatry, 2020. **25**(3): p. 640-654.

61. Carhart-Harris, R.L., et al., *Psilocybin with psychological support for treatment-resistant depression: six-month follow-up.* Psychopharmacology, 2018. **235**(2): p. 399-408.

62. Hibicke, M., et al., *Psychedelics, but not ketamine, produce persistent antidepressant-like effects in a rodent experimental system for the study of depression.* ACS chemical neuroscience, 2020. **11**(6): p. 864-871.

63. Baker, K.B. and J.J. Kim, *Amygdalar lateralization in fear conditioning: evidence for greater involvement of the right amygdala.* Behavioral neuroscience, 2004. **118**(1): p. 15.

64. Schneider, S., et al., *Boys do it the right way: sex-dependent amygdala lateralization during face processing in adolescents.* Neuroimage, 2011. **56**(3): p. 1847-1853.

65. Killgore, W.D. and D.A. Yurgelun-Todd, *Sex differences in amygdala activation during the perception of facial affect.* Neuroreport, 2001. **12**(11): p. 2543-2547.

66. Morgan, J. and T. Curran, *Calcium as a modulator of the immediate-early gene cascade in neurons.* Cell calcium, 1988. **9**(5-6): p. 303-311.

67. Kolter, J.F., et al., *Serotonin transporter genotype modulates resting state and predator stress-induced amygdala perfusion in mice in a sex-dependent manner.* PloS one, 2021. **16**(2): p. e0247311.

68. Pitychoutis, P., et al., *5-HT1A, 5-HT2A, and 5-HT2C receptor mRNA modulation by antidepressant treatment in the chronic mild stress model of depression: sex differences exposed.* Neuroscience, 2012. **210**: p. 152-167.

69. REMPEL-CLOWER, N.L., *Role of orbitofrontal cortex connections in emotion.* Annals of the New York Academy of Sciences, 2007. **1121**(1): p. 72-86.

70. Barbas, H., *Flow of information for emotions through temporal and orbitofrontal pathways.* Journal of anatomy, 2007. **211**(2): p. 237-249.

71. Gimpl, M., I. Gormezano, and J. Harvey, *Effects of LSD on learning as measured by classical conditioning of the rabbit nictitating membrane response.* Journal of Pharmacology and Experimental Therapeutics, 1979. **208**(2): p. 330-334.

72. Doss, M.K., et al., *Psilocybin therapy increases cognitive and neural flexibility in patients with major depressive disorder.* Translational psychiatry, 2021. **11**(1): p. 1-10.

73. Roberts, M. and P. Bradley, *Studies on the effects of drugs on performance of a delayed discrimination.* Physiology & Behavior, 1967. **2**(4): p. 389-397.

74. Griffiths, R.R., et al., *Mystical-type experiences occasioned by psilocybin mediate the attribution of personal meaning and spiritual significance 14 months later.* Journal of psychopharmacology, 2008. **22**(6): p. 621-632.

75. Griffiths, R.R., et al., *Psilocybin can occasion mystical-type experiences having substantial and sustained personal meaning and spiritual significance.* Psychopharmacology, 2006. **187**(3): p. 268-283.

Figure 1. (**A**) Experimental timeline. Animals arrived at 6-7 weeks old and were allowed to habituate for 1-week

prior to surgeries. Bilateral injection of calcium sensor (jGCaMP7f) and implantation of fiber optic cannulas were

conducted. After 3 weeks for viral transduction, animals underwent a series of fiber photometry experiments.

(**B**) Representative coronal section showing fiber placement and GCaMP expression. (**C**) Mean air-puff traces

showing consistency of CeA signal in response to the air-puff stimulus.

Figure 2. *Acute Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus Following Psilocin*

Administration (**A**) Females: representative images showing c-Fos+ cells tagged with a green fluorescence

protein (GFP) in the CeA. Scale bar = 100 **μm.** (**B**) Female vehicle vs. psilocin: histogram showing the number

of c-Fos+ cells in each subregion of the CeA. CeC = capsular CeA, CeM =. Medial CeA, CeL = lateral CeA.

Data points correspond to the mean of 2-4 hemispheres for each individual subject. (**C**) Males: representative

images showing c-Fos+ cells tagged with a green fluorescence protein (GFP) in the CeA. Scale bar = 100 **μm.**

(**D**) Male vehicle vs. psilocin: histogram showing the number of c-Fos+ cells in each subregion of the CeA.

CeC = capsular CeA, CeM =. Medial CeA, CeL = lateral CeA. Data points correspond to the mean of 2-4

hemispheres for each individual subject. (**E**) Female psilocin group: representative raw ΔF/F traces showing an

individual at baseline and while on psilocin. Blue line = air-puff onset, Grey lines = individual traces, Green line

= mean trace, Std= standard deviation (**F**) Females: Estimation plots for binned trace plots comparing baseline

to treatment. Each data point corresponds to a 500ms bin averaged across subjects +/- SEM (**G**) Female

vehicle group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85

psi. Data points represent group averages within 500ms binned window +/- S.E.M. (**H**) Female psilocin group:

air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data

points represent group averages within 500ms binned window +/- S.E.M. (**I**) Male psilocin group:

representative raw ΔF/F traces showing an individual at baseline and while on psilocin. Blue line = air-puff

onset, Grey lines = individual traces, Green line = mean trace, Std= standard deviation (**J**) Males: Estimation

plots for binned trace plots comparing baseline to treatment. Each data point corresponds to a 500ms bin

averaged across subjects +/- SEM. (**K**) Male vehicle group: air-puff trace plots of changes in CeA fluorescence

following exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned

window +/- S.E.M. (**L**) Male psilocin group: air-puff trace plots of changes in CeA fluorescence following

exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned window +/-

S.E.M. *$p<0.05$, **$p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure 3. *Prolonged Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus Following Psilocin*

Administration **(A)** Female psilocin group: representative raw ΔF/F traces showing an individual at baseline and

at the 28-day follow-up. Blue line = air-puff onset, Grey lines = individual traces, Green line = mean trace, Std=

standard deviation **(B)** Females: Estimation plots comparing baseline to the 28-day follow-up recording. Data

points represent each individual time bin post air-puff collapsed across subjects **(C)** Female vehicle group: air-

puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points

represent group averages within 500ms binned window +/- S.E.M. **(D)** Female psilocin group: air-puff trace plots

of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group

averages within 500ms binned window +/- S.E.M. **(E)** Male psilocin group: representative raw ΔF/F traces

showing an individual at baseline and while on psilocin. Blue line = air-puff onset, Grey lines = individual traces,

Green line = mean trace, Std= standard deviation **(F)** Males: estimation plots comparing baseline to the 28-day

follow-up recording. Data ppoints represent each individual time bin post air-puff collapsed across subjects. **(G)**

Male vehicle group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff

at 85 psi. Data points represent group averages within 500ms binned window +/- S.E.M. **(H)** Male psilocin group:

air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points

represent group averages within 500ms binned window. **(I)** Females: Summary histogram comparing the mean

of all 20 time bins following air-puff administration. Each data point represents an individual time bin within that

condition collapsed across all subjects. +/- S.E.M. **(J)** Males: Summary histogram comparing the mean of all 20

time bins following air-puff administration +/- S.E.M. Each data point represents an individual time bin within that

condition collapsed across all subjects. +/- S.E.M. *$p<0.05$, **$p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure 4. *Acute and Prolonged Behavioral Effects of Psilocin*. **(A)** Females: locomotion plots comparing vehicle

and psilocin groups at each time point. Data points are mean time in seconds +/- SEM. **(B)** Females: time spent

rearing comparing vehicle and psilocin groups at each time point. Data points are mean time in seconds +/-

SEM. **(C)** Males: locomotion plots comparing vehicle and psilocin groups at each time point. Data points are

mean time in seconds +/- SEM. **(D)** Males: time spent rearing comparing vehicle and psilocin groups at each

time point. Data points are mean time in seconds +/- SEM. **(E)** Females threat responding behavior. First figure

is showing percentage of subjects that darted following air-puff administration. Second figure is showing the

percentage of animals that remained immobile following air-puff. Third and fourth figure are pie charts showing

the percentage of animals that darted vs. remained immobile **(F)** Males threat responding behavior: First figure

is showing percentage of subjects that darted following air-puff administration. Second figure is showing the

percentage of subject that remained immobile following air-puff. Third and fourth figure are pie charts showing

the percentage of animals that darted vs. remained immobile **(G)** Male vehicle passive responders (remained

immobile) group: First figure shows air-puff trace plots of changes in CeA fluorescence following exposure to a

500ms air-puff at 85 psi. Data points represent group averages within 500ms binned window +/- S.E.M.; Second

figure is a histogram showing differences in the mean peak $\Delta F/F$ value for each group +/- S.E.M following air-

puff administration. Data points reflect each individual subject within the corresponding subgroup. **(H)** Male

Psilocin passive responders (remained immobile) group: First figure shows air-puff trace plots of changes in CeA

fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group averages within

500ms binned window +/- S.E.M. Second figure is a histogram showing differences in the mean peak $\Delta F/F$ value

for each group +/- S.E.M following air-puff administration. Data points reflect each individual subject within the

corresponding subgroup. **(I)** Male vehicle active responders (darted) group: First figure shows air-puff trace plots

of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group

averages within 500ms binned window +/- S.E.M. Second figure is a histogram showing differences in the mean

peak $\Delta F/F$ value for each group +/- S.E.M following air-puff administration. Data points reflect each individual

subject within the corresponding subgroup. **(J)** Male Psilocin active responders (darted) group: First figure shows

air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points

represent group averages within 500ms binned window +/- S.E.M. Second figure is a histogram showing

differences in the mean peak $\Delta F/F$ value for each group +/- S.E.M following air-puff administration. Data points

reflect each individual subject within the corresponding subgroup.

Figure 5. *Changes in CeA Reactivity in Response to an Auditory Stimulus Following Psilocin Administration in*

Female Subjects. **(A)** Female vehicle group: representative mean ΔF/F traces showing an individual at initial

auditory compared to final auditory. Blue line = air-puff onset, Grey lines = individual traces, Green line = mean

trace, Std= standard deviation **(B)** Female vehicle group: peak point of binned mean ΔF/F traces compared

between initial and final auditory recordings. Data points represent each individual subject's peak ΔF/F value

+/- SEM. **(C)** Female vehicle group: air-puff trace plots of changes in CeA fluorescence following exposure to a

500ms air-puff at 85 psi. Data points represent group averages within 500ms binned window +/- S.E.M **(D)**

Female vehicle group: Estimation plots for binned trace plots comparing baseline to treatment. Each data point

corresponds to a 500ms bin averaged across subjects +/- SEM. **(E)** Female psilocin group: representative

mean ΔF/F traces showing an individual at initial auditory compared to final auditory. Blue line = air-puff onset,

Grey lines = individual traces, Green line = mean trace, Std= standard deviation **(F)** Female psilocin group:

peak point of binned mean ΔF/F traces compared between initial and final auditory recordings. Data points

represent individual subject's peak ΔF/F value +/- SEM. **(G)** Female psilocin group: air-puff trace plots of

changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group

averages within 500ms binned window +/- S.E.M. **(H)** Female psilocin group: Estimation plots for binned trace

plots comparing baseline to treatment. Each data point corresponds to a 500ms bin averaged across subjects

+/- SEM. *$p<0.05$, **$p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure 6. *Changes in CeA Reactivity in Response to an Auditory Stimulus Following Psilocin Administration in*

Male Subjects. **(A)** Male vehicle group: representative mean ΔF/F traces showing an individual at initial auditory

compared to final auditory. Blue line = air-puff onset, Grey lines = individual traces, Green line = mean trace,

Std= standard deviation **(B)** Male vehicle group: peak point of binned mean ΔF/F traces compared between

initial and final auditory recordings. Data points represent each individual subject's peak ΔF/F value +/- SEM.

(C) Male vehicle group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-

puff at 85 psi. Data points represent group averages within 500ms binned window +/- S.E.M. **(D)** Male vehicle

group: Estimation plots for binned trace plots comparing baseline to treatment. Each data point corresponds to

a 500ms bin averaged across subjects +/- SEM. **(E)** Male psilocin group: representative mean ΔF/F traces

showing an individual at initial auditory compared to final auditory. Blue line = air-puff onset, Grey lines =

individual traces, Green line = mean trace, Std= standard deviation **(F)** Male psilocin group: peak point of binned

mean ΔF/F traces compared between initial and final auditory recordings. Data points represent individual

subject's peak ΔF/F value +/- SEM. **(G)** Male psilocin group: air-puff trace plots of changes in CeA fluorescence

following exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned

window +/- S.E.M. **(H)** Male psilocin group: Estimation plots for binned trace plots comparing baseline to

treatment. Each data point corresponds to a 500ms bin averaged across subjects +/- SEM. *$p<0.05$, **$p<0.01$,

*** $p<0.001$, **** $p<0.0001$.

Female   # Males

A Locomotion   **B** Exploratory   **C** Locomotion   **D** Exploratory

E Female Threat Responding

F Males Threat Responding

Male Passive Responders

G Vehicle   **H** Psilocin

Male Active Responders

I Vehicle   **J** Psilocin