Sex-Specific Effects of Psychedelic Drug Exposure on Central Amygdala Reactivity and

Behavioral Responding

DP Effinger, B.S.[1,2], SG Quadir, Ph.D[1,2], MC Ramage[1,2], MG Cone[1,2], MA Herman, Ph.D[1,2]

[1] Department of Pharmacology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599

[2] Bowles Center for Alcohol Studies, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599

Corresponding Author: M.A. Herman melissa_herman@unc.edu

ABSTRACT

Psilocybin, and its active metabolite psilocin, have been shown to elicit rapid and long-lasting symptom

improvements in a variety of affective psychiatric illnesses. However, the region-specific alterations underlying

these therapeutic effects remain relatively unknown. The central amygdala (CeA) is a primary output region

within the extended amygdala that is dysregulated in affective psychiatric disorders. Here, we measured CeA

activity using the activity marker *c-Fos* and CeA reactivity using fiber photometry paired with an aversive air-

puff stimulus. We found that psilocin administration acutely increased CeA activity in both males and females

and increased stimulus-specific CeA reactivity in females, but not males. In contrast, psilocin produced time-

dependent decreases in reactivity in males, but not females as early as 2-days and lasting to 28-days post

administration. We also measured behavioral responses to the air-puff stimulus and found sex-dependent

changes in threat responding but not exploratory behavior or general locomotion. Repeated presentations of

the auditory component of the air-puff were also performed and sex-specific effects of psilocin on CeA

reactivity to the auditory-alone stimulus were also observed. This study provides new evidence that a single

dose of psilocin produces sex-specific, time-dependent, and enduring changes in CeA reactivity and behavioral

responding to specific components of an aversive stimulus.

INTRODUCTION

Clinical studies suggest that psilocybin, and active metabolite psilocin [1-3], produce rapid, long-lasting improvements in psychiatric disorders including depression, anxiety, and substance use disorder [4-12]. Sustained improvements in symptoms were seen as long as 4 years following administration [13]. Research on psychedelic compounds has focused primarily on human imaging studies and receptor pharmacology. Human fMRI studies have shown that psychedelics produce robust alterations in network connectivity [8, 14-19]. Receptor studies have demonstrated that these compounds promote synaptogenesis [20-22] and identified the serotonin 5-hydroxytryptamine (5-HT)$_{2A}$ receptor to be instrumental in mediating the hallucinogenic actions of psychedelic compounds including psilocybin and lysergic acid diethylamide (LSD) [16, 23-28]. Despite the observed clinical outcomes in psychedelic studies, the underlying neural correlates of these therapeutic effects remain relatively unknown.

The central amygdala (CeA) is a primary output region within the extended amygdala that receives input regarding internal and external arousal state and coordinates appropriate behavioral responses [29, 30]. CeA dysregulation has been implicated in anxiety, depression, and PTSD [29, 31-36]. Preclinical studies demonstrate that the psychedelic compounds LSD and 2,5-dimethoxy-4-iodoamphetamine (DOI) increase Fos expression in the amygdala [37, 38] and that systemic and local DOI administration in the amygdala promotes suppression of fear responses [38]. Psilocybin produces alterations in reactivity and connectivity of the amygdala that correlate with positive therapeutic outcome [12, 15, 17, 39-41]. One clinical imaging study reported significantly decreased amygdala reactivity 1-week following psilocybin administration, with lasting changes in affect at one-month follow up [41].

The current study utilized a preclinical approach to determine how the active compound, psilocin, alters CeA activity, reactivity, and behavioral responding to an aversive stimulus in male and female Sprague Dawley rats. Psilocin was used as it is the active metabolite of psilocybin in the central nervous system (CNS), mediating effects of psilocybin at 5-HT$_{2AR}$ [2, 28]. We focused on the CeA as the primary output region of the amygdala and as amygdala hyperreactivity is associated with many psychiatric disorders that psilocybin shows promise

in treating [42-45]. We first measured expression of *c-Fos*, an immediate early gene that is upregulated

following neuronal activity, in the CeA of male and female rats administered psilocin (2 mg/kg) or vehicle. Next,

we measured CeA reactivity using an air-puff stimulus involving a burst of air directed towards the face. The

air-puff stimulus has been validated as an aversive stimulus in passive avoidance tests [46] and, due to known

CeA involvement in pain modulation [47, 48], it was essential to utilize a stimulus that would evoke a response

without the addition of potential confounding effects of pain. Utilizing fiber photometry in conjunction with the

aversive air-puff stimulus, we were able to demonstrate a robust and reliable CeA reactivity response. We then

used this model to determine how administration of psilocin altered CeA reactivity during acute exposure and

at more prolonged timepoints. We observed sex-dependent effects of psilocin on CeA activity as measured by

c-Fos expression and in-vivo reactivity and behavioral responding. Additionally, we found that psilocin elicited

long-term decreases in CeA reactivity in males that were not seen in females or in the vehicle control. To test

whether these effects would contribute to the development of a conditioned response to the sound of the air-

puff, additional assays were performed utilizing an auditory-only air-puff stimulus, wherein an air-puff was

administered with the nozzle facing outside of the box to isolate the auditory component of the air-puff. In doing

so, we show differential reactivity of the CeA suggesting that the observed changes were stimulus-specific and

dynamic in nature.

METHODS

Animals/Stereotaxic Surgery

Adult male and female Sprague Dawley rats (200-400 g, 6-weeks old at arrival, Envigo, Indianapolis, IN) were

group-housed in a humidity- and temperature-controlled (22°C) vivarium on a 12hr light/dark cycle with *ad*

libitum access to food and water. Following intracranial surgery, animals were single-housed in flat lid cages

with waterspout access, and food bowls to avoid damage to implants. Bilateral infusions of pGP-AAV-syn-

jGCaMP7f-WPRE (Addgene plasmid# 104488-AAV9) were performed at a rate of 100nl/minute into the CeA

(AP:-2.0mm; ML:+/- 3.9mm; DV:-8.0mm). Directly following injection, Doric 200um NA 0.37 silica optic fiber

cannulas with a 9mm tip were inserted at DV -7.7mm and secured in place using Metabond (Parkell,

Brentwood, NY).

Drug Administration

On the day of treatment, animals received a subcutaneous (s.c.) injection of either vehicle (0.9% Saline/2%

glacial acetic acid) or a 2 mg/kg psilocin (Cayman Chemical, graciously provided by Dr. Bryan Roth) dissolved

in 2% glacial acetic acid, as previously described [62]. Solutions were prepared fresh on the day of testing and

were stored at 4ºC for 30min prior to first injection. Injections were given at a 1mg/ml volume 30min prior to the

beginning of fiber photometry recordings. In the *c-Fos* cohort, injections were performed 2hr prior to perfusion.

Fiber Photometry Recording

Animals were habituated to testing room for 1 hour prior to recording sessions. Rats were individually placed in

a clear plexiglass chamber [50cm (l)×50cm(w)×38cm(h)] in a red-light sound-attenuated behavioral cabinet

and allowed to acclimate for 10min. Fiber placement and GCaMP injection were performed bilaterally,

however, recordings were collected unilaterally. Baseline recordings were taken from both hemispheres and

analyzed to determine which hemisphere to use for the remainder of the experiment. Ca^{2+} signals from

subjects, with 200µm optic fibers placed .3mm above the injection site in the CeA, were recorded using a TDT

RZ5 real time processor equipped with the TDT Synapse program (Tucker-Davis Technologies, Alachua, FL).

For each recording, jGCaMP7f was excited using both a 465nm calcium-dependent signal and a 405nm

isosbestic control. MATLAB script [63] was then used to analyze raw signal. Changes in fluorescence as a

function of baseline (ΔF/F) showing fluctuations in fluorescence compared to the overall baseline fluorescence

throughout the session were calculated through smoothing isosbestic control channel signal, scaling the

isosbestic signal by regressing it onto smoothed GCaMP signal, and then generating a predicted 405nm from

the regression to remove calcium-independent signaling from the raw GCaMP signal and control for artifact

from movement, photo-bleaching, and fiber bending artifacts. Additionally baseline fluorescence was

calculated through a least-squares linear fitting of the 405nm isosbestic control to the 465nm GCaMP signal to

create a resting GCaMP signal over the entire session not containing event-dependent fluctuations in GCaMP

signal [64]. Peri-event plots were then generated by averaging changes in fluorescence (ΔF/F) across two

different air-puff trials to generate a mean change in reactivity [63]. Area under curve (AUC) and peak point

(PP) were used to capture characteristics of amplitude and duration of CeA response. All subjects were video

recorded during stimulus presentations and behavior was scored using the Behavioral Observation Research

Interactive Software (BORIS) program [65]. Behavioral analysis was performed by an experimenter blind to the

experimental status of each animal.

Air-Puff Stimulus

An aversive air-puff stimulus was designed using house air supply (85psi) controlled by a Parker solenoid (Part

#003-0868-900), powered by a custom made 12V electrical circuit box connected to the TDT RZ5 system to

trigger 500ms openings of the solenoid and simultaneous timestamps through the Synapse program. Each

session consisted of a 10min habituation period before recording sessions that consisted of two air-puff

administrations with a 5min inter-stimulus interval (ISI) between them. Air-puffs were directed at the face and

within close proximity of the animal to maintain similar conditions and level of aversion to the stimulus between

subjects.

Perfusion/Tissue Collection

Rats were transcardially perfused with phosphate buffered saline (PBS) followed by 4% paraformaldehyde

(w/v). Brains were extracted, stored in 4% paraformaldehyde for 24h, then transferred to 30% sucrose (*w/v*)

and stored at 4⁰C until sectioning. Using a freezing microtome, tissue was sliced into 40μm thick sections and

stored in cryoprotectant (30% *v/v* ethylene glycol + 30% *w/v* sucrose in phosphate buffered saline) at 4⁰C until

use.

Immunofluorescence Staining

For site verification on the fiber photometry cohorts, 6 sections/subject anterior-posterior (AP) coordinates -

1.75 to -2.75 from bregma were washed in phosphate buffered saline (PBS) prior to 1hr blocking in 5% normal

donkey serum + 0.3% Triton X in PBS. Sections were then incubated in chicken anti-GFP (Abcam 13970;

1:1000) in 0.5% normal donkey serum/0.3% Triton X/PBS overnight at 4^{o}C. Next, samples were triple-washed

and incubated in blocking solution (1hr) followed by goat anti-chicken 488 (Jackson 703-545-155; 1:700) in

0.5% normal donkey serum/0.3% Triton X/PBS for 2hr at room temperature.

For the *c-Fos* cohort, sections AP coordinates -1.92mm to -2.16mm were washed in PBS, incubated in 50%

methanol/PBS solution for 30min, washed in 3% hydrogen peroxide/PBS for 5min, and placed in blocking

solution (0.3% Triton X-100; Thermo Fisher), 1% bovine serum albumin (BSA; Sigma) for 1hr, at room

temperature (RT). Slices were then incubated with rabbit anti-*cFos* (1:3000, Millipore Sigma; ABE457) for 24hr

at 4°C. Next, slices were washed with 0.1% Tween-20 in tris-buffered saline (TNT) before being transferred

into TNB blocking buffer (Perkin-Elmer FP1012) for 30min. After blocking, slices were incubated in goat anti-

rabbit horseradish peroxidase (HRP; 1:200, Abcam ab6721) for 2hr followed by another round of TNT washes.

Finally, slices were incubated in tyramide conjugated fluorescein (1:50) in TSA amplification diluent (Akoya

Biosciences, NEL741001KT) for 10min at RT. Slices were washed with TNT buffer, mounted, cover slipped

with Vectashield ® HardSet™ Antifade Mounting Medium with DAPI (H1500, Vector Laboratories, Burlingame,

CA) and stored at 4ºC before being imaged with the Keyence BZ-X800 fluorescence microscope. Images were

then analyzed by an experimenter blinded to the animal's condition. Images were taken at 20X and the BZ-

X800 Analyzer program was used to stitch images together. Randomly selected hemispheres were used, and

the area of interest was outlined and manually counted using the ImageJ multi-point counter tool. Cell counts

were averaged per animal.

Statistical Analysis

Raw data from the TDT RZ5 machine were imported and processed for artifact removal, down sampling, and

detrending using MATLAB code [63]. Changes in fluorescence (ΔF/F) were then calculated and collected 5s

before and 10s following air-puff administration. The mean of two traces were averaged together for each

subject and combined using custom MATLAB script (https://doi.org/10.5061/dryad.pnvx0k6q9) to sort

individual data into groups. Group 465nm signal data were then converted into 500ms bins for analysis to

quantify differences between experimental conditions. Area under the curve (AUC) analysis was calculated by

summing all raw ΔF/F values following the air puff administration. This would correspond to the timepoints 0-20

on the binned analysis plots. This calculation was captured for each animals mean trace within each condition

and then collapsed across subjects to determine group means in AUC following air puff. All statistical analyses

were conducted using GraphPad Prism 7. Five subjects were removed from the study due to lack of

signal/cannula removal. Repeated-measures one-way analysis of variance (ANOVA) with Dunnett's post hoc

test was used to determine differences between conditions within each groups in the air-puff condition. Two-

way ANOVAs with Šidák's post hoc tests were used to compare area under the curve, peak point, and time

bins between groups and time points. Paired t-tests were used to determine differences within groups between

time points in the binned trace and auditory-only analyses. Fisher's exact tests were used to compare

percentages in darting behavior. Mixed-effects two-way ANOVA were used to compare time spent walking and

rearing. All data including Matlab structures and scripts can be found at

https://doi.org/10.5061/dryad.pnvx0k6q9.

RESULTS

Acute Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus

Male and female Sprague Dawley rats received bilateral injection of pGP-AAV-syn-jGCaMP7f-WPRE into the

CeA immediately followed by installation of fiber optic cannulae above the injection site (**Fig. 1A-B**). Following

3-weeks for virus incubation, animals underwent a series of fiber photometry experiments testing the effects of

psilocin on fluctuations in calcium dynamics within the CeA time-locked to an aversive air-puff stimulus (**Fig.**

1A). For both baseline and treatment recording sessions, each animal received two air-puffs with an ISI of

5min and trace plots were generated showing change in fluorescence as a function of baseline (ΔF/F). Traces

were analyzed 5s before and 10s following administration of air-puff (**Fig. 1C**).

Sub-Region-Specific Differences in CeA Basal Activity Following Psilocin Administration

To assess the effects of psilocin on basal activity in the CeA, male and female Sprague Dawley rats were injected with psilocin (2 mg/kg, s.c.) or vehicle (1 ml/kg, s.c.) 2hr prior to transcardial perfusion. Immunohistochemistry (IHC) assays were performed using CeA tissue collected within the anterior-posterior bregma range of -1.92mm to -2.16mm. In females, there was a significantly increased *c-Fos* expression in the psilocin group as compared to vehicle across regions ($F_{Interaction}(2,24)=1.597$, $p=0.22$; $F_{Region}(1,24)=30.84$, $p<0.0001$; $F_{Treatment}(1,24)=9.149$ $p=0.006$; **Fig.2A-B**) with greater increases in the capsular division of the central amygdala (CeC) (Šidák's: $p=0.02$). In males, there was also significantly increased *c-Fos* expression in the psilocin group across regions ($F_{Interaction}(2,27)=1.241$, $p=0.30$; $F_{Region}(2,27)=12.02$, $p=0.0002$; $F_{Treatment}(1,27)=5.543$, $p=0.02$; **Fig.2C-D**).

Acute Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus Following Psilocin Administration

To test the acute effects of psilocin on CeA reactivity, we measured changes in CeA reactivity during time-locked stimuli presentation at baseline and 30min post administration of vehicle or psilocin (2 mg/kg, s.c.). Following a 10min habituation period, air-puffs were delivered as described above. Mean traces were taken from two separate air-puff responses within a session (**Fig. 2E, 2I**). These traces were then collapsed into 500ms bins for statistical analysis (**Fig. 2G, 2H, 2K, 2L**). To compare baseline to treatment, paired t-tests were performed comparing the individual time bins following the air-puff collapsed across subjects (Fig. **2F**, **2J**). In the female vehicle control group (n=9), there was a significant increase between baseline and treatment ($t(19)=2.780$, $p=0.01$; **Fig.2F-G**). In the female psilocin group (n=7), there was a significant increase between baseline and treatment ($t(19)=12.63$, $p<0.0001$; **Fig.2F, 2H**). Though changes were seen in both vehicle and psilocin, the increase in mean of differences in reactivity was greater in the psilocin group compared to the vehicle control (**Fig.2F**). In the male vehicle control group (n=10), there was an increase in reactivity between baseline and treatment, though the effect was small ($t(19)=4.452$, $p=0.0003$; **Fig.2J-K**). In the male psilocin

group (n=11), there was no significant difference between baseline and treatment (t(19)=0.6231, *p*=0.54;

Fig.2J, 2L).

Prolonged Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus Following Psilocin

Administration

To assess long-term alterations in CeA reactivity following a single dose of psilocin, fiber photometry

recordings were performed on days 2, 6, and 28 post-administration of vehicle or psilocin. On each follow up

recording day, animals received two air-puffs with a 5-minute ISI. In females, there was no difference between

vehicle and psilocin groups in the area under curve (AUC) of the group-averaged individual mean traces

between baseline and the 28-day follow ($F_{Interaction}$(1,14)=0.003383, *p*=0.95; $F_{Treatment}$(1,14)=0.09635, *p*=0.76;

F_{Time}(1,14)=0.2126, *p*=0.65; **Fig.3A-B**). Additionally, no differences were found between vehicle and psilocin

groups at any of the follow-up timepoints ($F_{Interaction}$(2,28)=1.437, *p*=0.25; $F_{Treatment}$(1,14)=0.0003490, *p*=0.99;

F_{Time}(2,28)=3.067, *p*=0.06; **Fig.3B**). However, in the vehicle control females, 1-way ANOVA found an effect of

treatment (F(3,57)=50.02, *p*<0.0001; **Fig.3C**) and a significant increase in reactivity at the 2-day and 6-day

follow up (Dunnett's: *p*<0.0001). In the female psilocin group (**Fig. 3A, 3D**), 1-way ANOVA found an effect of

treatment (F(3,57)=54.13, *p*<0.0001; **Fig.3D**) with significant increases in reactivity at the 2-day (Dunnett's:

p<0.0001) and 6-day follow up (Dunnett's: *p*<0.0001).

In the males, there was decreased reactivity between the vehicle and psilocin group from baseline to the 28-

day follow up ($F_{Interaction}$(1,19)=5.068, *p*=0.03; $F_{Treatment}$(1,19)=0.2503, *p*=0.62; F_{Time}(1,19)=1.361, *p*=0.26; **Fig.3F**)

with a significant reduction in reactivity at the 28-day follow up in the psilocin group (Šidák's: *p*=0.04). There

were also reductions in the AUC at the follow up recordings in the psilocin group compared to the vehicle

control ($F_{Interaction}$(2,38)=0.2113, *p*=0.81; $F_{Treatment}$(1,19)=4.926, *p*=0.03; F_{Time}(2,38)=0.7906, *p*=0.46; **Fig.3E-F**).

In the male vehicle group, 1-way ANOVA found an effect of treatment (F(3,57)=5.554, *p*=0.002; **Fig.3G**) with a

significant increase in reactivity at the 28-day follow up (Dunnett's: *p*=0.01). In the male psilocin group, 1-way

ANOVA found a significant effect of treatment (F(3,57)=72.60, *p*<0.0001; **Fig.3H**) revealing significant

decreases in reactivity at the 2-day, 6-day, and 28-day follow up (Dunnett's: *p*<0.0001). A 2-way ANOVA found

differential effects of treatment across time points between groups in females ($F_{Interaction}(4,152)=35.02$,

$p<0.0001$; $F_{Treatment}(1,38)=0.9089$, $p=35$; $F_{Time}(2.343,89.04)=68.58$, $p<0.0001$; **Fig.3J**) and males

($F_{Interaction}(4,152)=38.35$, $p<0.0001$; $F_{Treatment}(1,38)=1.059$, $p=0.31$, $F_{Time}(2.603,98.90)=42.25$, $p<0.0001$; **Fig.3I**)

There were increases in reactivity during treatment and reductions at the 2-day follow up in the female psilocin

group compared to vehicle control (Šidák: $p=0.001$; **Fig.3J**). In males, there was a significant decrease in

reactivity at the 28-day follow up in the psilocin group compared to vehicle control (Šidák: $p=0.01$; **Fig.3J**)

Acute and Prolonged Behavioral Effects of Psilocin

To determine if psilocin elicited changes in locomotion and exploratory behavior, video recordings were scored

for total time spent walking and time spent rearing, respectively. In females, there were no differences in

overall locomotion between vehicle and psilocin groups ($F_{Interaction}(3,42)=0.2118$, $p=0.89$; $F_{Treatment}(1,14)=2.238$,

$p=0.16$; $F_{Time}(2.497,34.95)=0.6065$, $p=0.59$; **Fig.4A**). There were also no differences in rearing between female

vehicle and psilocin groups ($F_{Interaction}(4,56)=1.345$, $p=0.26$; $F_{Treatment}(1,14)=4.433$, $p=0.05$; $F_{Time}(4,56)=2.486$,

$p=0.05$; **Fig. 4B**). In the males, there were no differences in locomotion ($F_{Interaction}(3,57)=0.1054$, $p=0.96$;

$F_{Treatment}(1,19)=2.563$, $p=0.12$; $F_{Time}(3,57)=1.255$, $p=0.29$; **Fig.4C**) or rearing ($F_{Interaction}(4,76)=0.5546$, $p=0.69$;

$F_{Treatment}(1, 19)=0.7576$, $p=0.39$; $F_{Time}(4,76)=1.190$, $p=0.32$; **Fig.4D**) between vehicle and psilocin groups.

Collectively, no significant alterations in locomotion were seen resulting from psilocin administration (**Fig.4A,**

4C), although females appeared to display increased locomotion overall as compared to males, regardless of

treatment.

To probe potential differences in threat responding to the air-puff stimulus, we separated groups based on

active or passive coping strategy. Subgroups were created containing animals that remained immobile

(passive) following air-puff administration versus animals that exhibited darting behavior (active) following air-

puff administration. Consistent with previous literature [49], females primarily employed an active darting

response (**Fig.4E**), while active vs passive coping behavior was more evenly split in males (**Fig.4F**). In

females, there were no differences in the percentage of animals that darted or that stayed immobile between

groups at any of the time points (**Fig. 4E**). In contrast, the male psilocin group displayed nonsignificant

increases in darting and reductions in immobility in the vehicle control group that were not seen in the psilocin

group under treatment conditions ($p=0.08$; **Fig. 4F**). At the 2-day follow up, there was a significantly higher

percentage of darting in the vehicle control group compared to the psilocin group ($p=0.03$; **Fig.4F, left**) and a

significantly higher percentage immobile in the psilocin group ($p=0.03$; **Fig.4F, right**). Though differences were

not significant at every time point, psilocin seemed to prevent a trending increase in darting behavior seen in

the male vehicle control group and increase in overall immobility in male but not female rats (**Fig.4F**).

To explore the CeA reactivity correlates to the observed behavioral differences in threat responding, fiber

photometry traces were split into the corresponding active vs. passive responder groups. Since females

primarily employed active coping strategies that were not affected by psilocin, we only examined CeA reactivity

between the male active and passive responders. To assess changes in the amplitude of CeA response to the

stimulus, comparisons looking at the peak point, or highest ΔF/F value following air-puff, were made. In the

male passive responder, there were no significant differences in peak point in the vehicle ($F(4,20)=1.219$,

$p=0.33$; **Fig.4G**) or psilocin group ($F(4,20)=1.276$, $p=0.31$; **Fig.4H**). In the active responder vehicle males, no

differences were seen in peak point compared to baseline ($F(4,12)=1.046$, $p=0.42$; **Fig.4I**). However, in the

active responder psilocin males, there was a main effect of treatment ($F(4,16)=6.566$, $p=0.003$; **Fig.4J**) with a

significant decrease in peak point at day 2 compared to baseline ($p=0.003$), suggesting that psilocin-induced

decreases in the amplitude of CeA reactivity were more pronounced in males employing an active vs. passive

threat response at baseline.

Changes in CeA Reactivity in Response to an Auditory Stimulus Following Psilocin Administration

To test whether the animals developed a conditioned response to the tone associated with the air-puff

stimulus, and whether psilocin treatment altered the response to the auditory stimulus alone, two separate

auditory-only sessions were conducted immediately following baseline recordings and again at the 28-day

recording session. These auditory only sessions followed the same parameters as the air-puff with the

exceptions being that the ISI was reduced to 3min and the air-puff apparatus was facing outside of the box to

maintain the same auditory tone without the physical sensation of the air-puff.

In vehicle control females (**Fig.5A**), there was a significant decrease in peak point reactivity from the initial to

the final auditory condition (t(8)=3.934, *p*=0.004; **Fig.5B**) and a significant decrease in binned analysis from the

initial auditory to the final auditory condition (t(19)=18.28, *p*<0.0001; **Fig.5C-D**). In psilocin females (**Fig.5E**),

there were no significant differences between the initial and final auditory session in either peak point

(t(6)=0.4346, *p*=0.67; **Fig.5F)** or binned analysis (t(19)=0.01624, *p*=0.98; **Fig.5G-H**).

In the vehicle males (**Fig.6A**), there were no significant differences in peak point between initial and final

auditory condition (t(9)=2.136, *p*=0.06; **Fig. 6B**). However, there was a significant decrease in binned analysis

of the final auditory session compared to the initial (t(19)=9.719, *p*<0.0001; **Fig.6C-D**). In the psilocin-treated

males (**Fig.6E**), there was no difference in peak point from the initial to the final auditory session

(t(10)=0.09932, *p*=0.92; **Fig.6F**). However, there was a significant decrease in binned analysis of the initial

auditory and final auditory condition (t(19)=4.943, *p*<0.0001; **Fig.6G-H**). While there was a decrease in both

the vehicle and psilocin groups, the decrease in peak point in the vehicle group was of greater overall

magnitude (**Fig.6B, 6F**) and the estimation plots showed a much greater reduction in the vehicle (**Fig.6D**) as

compared to the psilocin group (**Fig.6H**).

DISCUSSION

The present study explored the effects of the psychedelic compound psilocin on CeA activity using an

immunohistological activity marker and on CeA reactivity to an aversive stimulus using *in-vivo* fiber photometry

recordings during acute drug exposure and at 2- , 6- , and 28-days post drug administration. The study

examined multi-level neurobiological and behavioral effects of psilocin by looking at acute drug effects, long-

term alterations in activity following drug administration, and by probing for potential sex-dependent effects of

both treatment and time. Psilocin administration produced increases in *c-Fos* expression in both males and

females and sub-region analysis showed that increases in the female CeC were significantly greater in psilocin

compared to the vehicle control (**Fig. 2B**). CeA reactivity in response to the air-puff stimulus following psilocin

administration also displayed sex- and time-dependent effects. In females, psilocin increased CeA reactivity as

compared to vehicle controls (**Fig. 2E-H**). Interestingly, both vehicle and psilocin groups showed increases in

reactivity at the 2- and 6-day follow up compared to baseline. In contrast, the male psilocin group did not show

the same increased reactivity seen in the females (**Fig. 2H**) and instead displayed decreases in CeA reactivity

at 2-day, 6-day, and 28-day follow-up recordings (**Fig. 3F, 3H**) that were not seen in vehicle control males (**Fig.**

3G).

Clinical studies have illustrated that psilocybin produces alterations in amygdala reactivity and connectivity [8,

12, 15, 17, 39-41]. Here, we utilized a preclinical approach to examine the actions of psilocin, the psychoactive

metabolite of psilocybin, on central amygdala activity in a rodent model. As most clinical studies report a

reduction in amygdala reactivity or connectivity that correlated with therapeutic improvements [8, 15, 17, 39-

41], we hypothesized that psilocin would produce similar decreases in reactivity in our model. Here we found

that a single dose of psilocin produced persistent reductions in CeA reactivity to an aversive stimulus in male

but not female rats. These findings lend to recent studies in both clinical [11, 13, 50] and preclinical [51]

literature showing persistent reductions in depressive behavior after a single dose of psilocybin. Although the

mechanism behind these therapeutic effects remains unknown, this study provides evidence that changes in

CeA activity may play a role in observed therapeutic effects.

We report sex-specific effects of psilocin on basal activity and stimulus-induced reactivity within the CeA.

Changes in basal activity were measured by *c-Fos*, an immediate early gene shown to be upregulated

following neuronal activation [52]. We observed an increase in *c-Fos* expression in both males and females

following administration of psilocin, with region specific increases found the CeC subregion in females. CeA

reactivity was assessed using fiber photometry to measure in-vivo stimulus-induced fluctuations in calcium

signaling. In the female psilocin group, we observed an acute increase in reactivity to the air-puff stimulus while

on the drug that was not seen in the males. Additionally, we saw increases in reactivity at 2- and 6-day follow

up recording, while we saw reductions in CeA reactivity in the males at each of the follow up recording days

(**Fig. 3B, 3D, 3E, 3G**). A recent study looking at cerebral blood flow and the expression of the serotonin

transporter (5-HTT) showed that, in response to an aversive predator odor, there were increases in blood flow

and *c-Fos* expression in the CeA that were related to 5-HTT expression in males but not females [53]. These

findings suggest potential sex-specific expression of serotonin receptors within the amygdala. Given the affinity

of psilocin for the serotonin system, differential expression of serotonin receptors within the CeA or surrounding

nuclei could explain the sex-specific effects seen in this study. Additionally, a previous study showed sex-

specific differences in 5-HT$_{2AR}$ expression at baseline, wherein females had lower expression in the

orbitofrontal cortex (OFC) than males [54]. Another potential explanation for the differential effects of psilocin

between males and females may be due to differences in CeA circuitry. For example, the OFC has been

shown, in rodents and non-human primates, to have direct connections to the CeA as well as to the

intercalated nuclei, a local GABAergic population within the amygdala complex [55, 56]. Lack of activation of

this circuit due to a reduced expression of OFC 5-HT$_{2AR}$ in females could prevent an inhibition of CeA firing,

thus causing the increase in reactivity seen while on psilocin in females but not males. Future experiments are

needed to examine the effects of psilocin on CeA circuits innervating distinct brain regions.

In the presence of a threatening stimulus, animals will employ either active or passive coping strategies. In this

study, the active threat response was characterized by darting, or immediate fleeing from the air-puff stimulus,

while the passive threat response was characterized by freezing, or immobility following the air-puff. Here, we

show that females predominantly exhibited an active response, while males were more evenly split between

active and passive responses. In male psilocin-treated animals who exhibited an active response at baseline,

there was a significant decrease in the amplitude of CeA reactivity following exposure to the air-puff at the 2-

day follow up, with no changes in vehicle control. This decrease in reactivity was not seen in the passive-

responder vehicle or psilocin groups. While further interpretation regarding adaptive vs. maladaptive strategies

in this context is outside the scope of the current study, these results suggest that there may be differences in

CeA reactivity and threat responding strategy that are implicated in the efficacy of psychedelics.

To determine the effects of psilocin on associative learning, we utilized an auditory-only stimulus recording

taken after the initial baseline recordings and then again after the 28-day recording. Interestingly, in both males

and females the vehicle control group displayed reductions in CeA response to the auditory-only stimulus. In

the psilocin group however, reductions in CeA reactivity to the auditory stimulus were not observed in females,

while in males, reductions were of a much lesser degree than the vehicle control group (**Fig. 6**). Given the

reductions in reactivity seen with the standard air-puff stimulus, the sustained reactivity to the auditory stimulus

in the psilocin group suggests that the decreases in reactivity we found under the air-puff conditions were not

due to an overall blunting of CeA function. Instead, these data suggest a more dynamic alteration in CeA

reactivity in response to different types of stimuli. One potential interpretation of these data is that psilocin may

influence associative learning to an aversive stimulus. Several studies have shown that $5\text{-}HT_{2A}$ agonist

psychedelics produce improvements in various dimensions of cognition, including associative learning [57-59].

Another potential interpretation of these findings is that psilocin created a more robust/meaningful memory of

the air-puff. Many of the clinical studies cite participants rating their experience during a psilocybin treatment as

one of the most meaningful experiences of their life [13, 60, 61]. Additionally, increases in plasticity could

contribute to differences in memory reconsolidation during air-puff administration while on the drug.

Our findings demonstrate that the psychedelic compound, psilocin, produces dynamic, sex-specific changes in

the CeA, a key nucleus within the amygdala complex. Here we showed that administration of psilocin produces

increases in CeA *c-Fos* expression and that, in females, these increases were primarily driven by increases in

expression in the CeC. We have also shown that there are sex-dependent changes in CeA reactivity while on

psilocin. Specifically, we show that a single administration of psilocin produced acute increases in CeA

reactivity in females, but not males. Additionally, we show that persistent changes in CeA reactivity are seen as

early as 2 and as long as 28 days following administration of the drug in males, but not females. Given that

dysregulation of the amygdala is a hallmark in many different psychiatric disorders, including anxiety, MDD,

and PTSD [29, 31-36] and that psychedelic compounds show promise in treating many of these illnesses [4, 6-

11], these findings provide important information on subregion-specific alterations in CeA function that may

play a part in observed therapeutic effects of psychedelics. Additionally, the long-term reductions in amygdala

reactivity seen with the air-puff stimulus do not seem to be the result of overall desensitization of the CeA,

given the sustained response in the auditory-only stimulus response seen in the psilocin group for both males

and females. Future work will be required to explore potential circuit-based mediators contributing to these

observed alterations in CeA reactivity, but these findings provide important evidence for CeA alterations

underlying the potential therapeutic effects of psilocin and psilocybin.

ACKNOWLEDGEMENTS

This work was supported by National Institute of Health grants AA026858 (M.A.H.), GM135095 (D.P.E.), AA007573 (S.G.Q.), and by the Brain and Behavior Research Foundation NARSAD Young Investigator Award (M.A.H.)

COMPETING INTERESTS

The authors have no competing interests to report.

REFERENCES

1. Nichols, D.E., *Hallucinogens.* Pharmacology & therapeutics, 2004. **101**(2): p. 131-181.

2. Dinis-Oliveira, R.J., *Metabolism of psilocybin and psilocin: clinical and forensic toxicological relevance.* Drug metabolism reviews, 2017. **49**(1): p. 84-91.

3. Passie, T., et al., *The pharmacology of psilocybin.* Addiction biology, 2002. **7**(4): p. 357-364.

4. Davis, A.K., et al., *Effects of psilocybin-assisted therapy on major depressive disorder: a randomized clinical trial.* JAMA psychiatry, 2021. **78**(5): p. 481-489.

5. Bogenschutz, M.P. and M.W. Johnson, *Classic hallucinogens in the treatment of addictions.* Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2016. **64**: p. 250-258.

6. Johnson, M.W., A. Garcia-Romeu, and R.R. Griffiths, *Long-term follow-up of psilocybin-facilitated smoking cessation.* The American journal of drug and alcohol abuse, 2017. **43**(1): p. 55-60.

7. Johnson, M.W. and R.R. Griffiths, *Potential therapeutic effects of psilocybin.* Neurotherapeutics, 2017. **14**(3): p. 734-740.

8. Carhart-Harris, R.L., et al., *Psilocybin for treatment-resistant depression: fMRI-measured brain mechanisms.* Scientific reports, 2017. **7**(1): p. 1-11.

9. Griffiths, R.R., et al., *Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: A randomized double-blind trial.* Journal of psychopharmacology, 2016. **30**(12): p. 1181-1197.

10. Bogenschutz, M.P., et al., *Psilocybin-assisted treatment for alcohol dependence: a proof-of-concept study.* Journal of psychopharmacology, 2015. **29**(3): p. 289-299.

11. Ross, S., et al., *Rapid and sustained symptom reduction following psilocybin treatment for anxiety and depression in patients with life-threatening cancer: a randomized controlled trial.* Journal of psychopharmacology, 2016. **30**(12): p. 1165-1180.

12. Roseman, L., et al., *Increased amygdala responses to emotional faces after psilocybin for treatment-resistant depression.* Neuropharmacology, 2018. **142**: p. 263-269.

13. Agin-Liebes, G.I., et al., *Long-term follow-up of psilocybin-assisted psychotherapy for psychiatric and existential distress in patients with life-threatening cancer.* Journal of Psychopharmacology, 2020. **34**(2): p. 155-166.

14. Barrett, F.S., et al., *Psilocybin acutely alters the functional connectivity of the claustrum with brain networks that support perception, memory, and attention.* NeuroImage, 2020. **218**: p. 116980.

15. Grimm, O., et al., *Psilocybin modulates functional connectivity of the amygdala during emotional face discrimination.* Eur Neuropsychopharmacol, 2018. **28**(6): p. 691-700 DOI: 10.1016/j.euroneuro.2018.03.016.

16. Preller, K.H., et al., *Psilocybin induces time-dependent changes in global functional connectivity.* Biological psychiatry, 2020. **88**(2): p. 197-207.

17. Mertens, L.J., et al., *Therapeutic mechanisms of psilocybin: changes in amygdala and prefrontal functional connectivity during emotional processing after psilocybin for treatment-resistant depression.* Journal of Psychopharmacology, 2020. **34**(2): p. 167-180.

18. Roseman, L., et al., *The effects of psilocybin and MDMA on between-network resting state functional connectivity in healthy volunteers.* Frontiers in human neuroscience, 2014. **8**: p. 204.

19. Carhart-Harris, R.L., et al., *Neural correlates of the psychedelic state as determined by fMRI studies with psilocybin.* Proceedings of the National Academy of Sciences, 2012. **109**(6): p. 2138-2143.

20. Shao, L.-X., et al., *Psilocybin induces rapid and persistent growth of dendritic spines in frontal cortex in vivo.* Neuron, 2021. **109**(16): p. 2535-2544. e4.

21. Jones, K.A., et al., *Rapid modulation of spine morphology by the 5-HT2A serotonin receptor through kalirin-7 signaling.* Proceedings of the National Academy of Sciences, 2009. **106**(46): p. 19575-19580.

22. Ly, C., et al., *Psychedelics promote structural and functional neural plasticity.* Cell reports, 2018. **23**(11): p. 3170-3182.

23. Lee, H.-M. and B.L. Roth, *Hallucinogen actions on human brain revealed.* Proceedings of the National Academy of Sciences, 2012. **109**(6): p. 1820-1821.

24. Kometer, M., et al., *Activation of serotonin 2A receptors underlies the psilocybin-induced effects on α oscillations, N170 visual-evoked potentials, and visual hallucinations.* Journal of Neuroscience, 2013. **33**(25): p. 10544-10551.

25. Kometer, M., et al., *Psilocybin biases facial recognition, goal-directed behavior, and mood state toward positive relative to negative emotions through different serotonergic subreceptors.* Biological psychiatry, 2012. **72**(11): p. 898-906.

26. Halberstadt, A.L., et al., *Differential contributions of serotonin receptors to the behavioral effects of indoleamine hallucinogens in mice.* Journal of psychopharmacology, 2011. **25**(11): p. 1548-1561.

27. Halberstadt, A.L., S.B. Powell, and M.A. Geyer, *Role of the 5-HT2A receptor in the locomotor hyperactivity produced by phenylalkylamine hallucinogens in mice.* Neuropharmacology, 2013. **70**: p. 218-227.

28. Madsen, M.K., et al., *Psychedelic effects of psilocybin correlate with serotonin 2A receptor occupancy and plasma psilocin levels.* Neuropsychopharmacology, 2019. **44**(7): p. 1328-1334 DOI: 10.1038/s41386-019-0324-9.

29. Gilpin, N.W., M.A. Herman, and M. Roberto, *The central amygdala as an integrative hub for anxiety and alcohol use disorders.* Biological psychiatry, 2015. **77**(10): p. 859-869.

30. Ciocchi, S., et al., *Encoding of conditioned fear in central amygdala inhibitory circuits.* Nature, 2010. **468**(7321): p. 277-282.

31. Sandu, A.L., et al., *Amygdala and regional volumes in treatment-resistant versus nontreatment-resistant depression patients.* Depression and anxiety, 2017. **34**(11): p. 1065-1071.

32. Armony, J.L., et al., *Amygdala response in patients with acute PTSD to masked and unmasked emotional facial expressions.* American Journal of Psychiatry, 2005. **162**(10): p. 1961-1963.

33. Frodl, T., et al., *Enlargement of the amygdala in patients with a first episode of major depression.* Biological Psychiatry, 2002. **51**(9): p. 708-714 DOI: https://doi.org/10.1016/S0006-3223(01)01359-2.

34. Rauch, S.L., et al., *Exaggerated amygdala response to masked facial stimuli in posttraumatic stress disorder: a functional MRI study.* Biological psychiatry, 2000. **47**(9): p. 769-776.

35. Semple, W.E., et al., *Higher brain blood flow at amygdala and lower frontal cortex blood flow in PTSD patients with comorbid cocaine and alcohol abuse compared with normals.* Psychiatry, 2000. **63**(1): p. 65-74.

36. Siegle, G.J., et al., *Increased Amygdala and Decreased Dorsolateral Prefrontal BOLD Responses in Unipolar Depression: Related and Independent Features.* Biological Psychiatry, 2007. **61**(2): p. 198-209 DOI: https://doi.org/10.1016/j.biopsych.2006.05.048.

37. Gresch, P., L. Strickland, and E. Sanders-Bush, *Lysergic acid diethylamide-induced Fos expression in rat brain: role of serotonin-2A receptors.* Neuroscience, 2002. **114**(3): p. 707-713.

38. Pędzich, B.D., et al., *Effects of a psychedelic 5-HT2A receptor agonist on anxiety-related behavior and fear processing in mice.* Neuropsychopharmacology, 2022: p. 1-11.

39. Kraehenmann, R., et al., *Psilocybin-Induced Decrease in Amygdala Reactivity Correlates with Enhanced Positive Mood in Healthy Volunteers.* Biol Psychiatry, 2015. **78**(8): p. 572-81 DOI: 10.1016/j.biopsych.2014.04.010.

40. Kraehenmann, R., et al., *The mixed serotonin receptor agonist psilocybin reduces threat-induced modulation of amygdala connectivity.* NeuroImage: Clinical, 2016. **11**: p. 53-60.

41. Barrett, F.S., et al., *Emotions and brain function are altered up to one month after a single high dose of psilocybin.* Scientific Reports, 2020. **10**(1): p. 1-14.

42. Bas-Hoogendam, J.M., et al., *Amygdala hyperreactivity to faces conditioned with a social-evaluative meaning–a multiplex, multigenerational fMRI study on social anxiety endophenotypes.* NeuroImage: Clinical, 2020. **26**: p. 102247.

43. Donegan, N.H., et al., *Amygdala hyperreactivity in borderline personality disorder: implications for emotional dysregulation.* Biological psychiatry, 2003. **54**(11): p. 1284-1293.

44. Yang, T.T., et al., *Adolescents with major depression demonstrate increased amygdala activation.* Journal of the American Academy of Child & Adolescent Psychiatry, 2010. **49**(1): p. 42-51.

45. Joos, A.A., et al., *Amygdala hyperreactivity in restrictive anorexia nervosa.* Psychiatry Research: Neuroimaging, 2011. **191**(3): p. 189-195.

46. Moriarty, O., et al., *Validation of an air-puff passive-avoidance paradigm for assessment of aversive learning and memory in rat models of chronic pain.* Journal of neuroscience methods, 2012. **204**(1): p. 1-8.

47. Veinante, P., I. Yalcin, and M. Barrot, *The amygdala between sensation and affect: a role in pain.* Journal of molecular psychiatry, 2013. **1**(1): p. 1-14.

48. Wilson, T.D., et al., *Dual and opposing functions of the central amygdala in the modulation of pain.* Cell reports, 2019. **29**(2): p. 332-346. e5.

49. Gruene, T.M., et al., *Sexually divergent expression of active and passive conditioned fear responses in rats.* Elife, 2015. **4**: p. e11352.

50. Carhart-Harris, R.L., et al., *Psilocybin with psychological support for treatment-resistant depression: six-month follow-up.* Psychopharmacology, 2018. **235**(2): p. 399-408.

51. Hibicke, M., et al., *Psychedelics, but not ketamine, produce persistent antidepressant-like effects in a rodent experimental system for the study of depression.* ACS chemical neuroscience, 2020. **11**(6): p. 864-871.

52. Morgan, J. and T. Curran, *Calcium as a modulator of the immediate-early gene cascade in neurons.* Cell calcium, 1988. **9**(5-6): p. 303-311.

53. Kolter, J.F., et al., *Serotonin transporter genotype modulates resting state and predator stress-induced amygdala perfusion in mice in a sex-dependent manner.* PloS one, 2021. **16**(2): p. e0247311.

54. Pitychoutis, P., et al., *5-HT1A, 5-HT2A, and 5-HT2C receptor mRNA modulation by antidepressant treatment in the chronic mild stress model of depression: sex differences exposed.* Neuroscience, 2012. **210**: p. 152-167.

55. REMPEL-CLOWER, N.L., *Role of orbitofrontal cortex connections in emotion.* Annals of the New York Academy of Sciences, 2007. **1121**(1): p. 72-86.

56. Barbas, H., *Flow of information for emotions through temporal and orbitofrontal pathways.* Journal of anatomy, 2007. **211**(2): p. 237-249.

57. Gimpl, M., I. Gormezano, and J. Harvey, *Effects of LSD on learning as measured by classical conditioning of the rabbit nictitating membrane response.* Journal of Pharmacology and Experimental Therapeutics, 1979. **208**(2): p. 330-334.

58. Doss, M.K., et al., *Psilocybin therapy increases cognitive and neural flexibility in patients with major depressive disorder.* Translational psychiatry, 2021. **11**(1): p. 1-10.

59. Roberts, M. and P. Bradley, *Studies on the effects of drugs on performance of a delayed discrimination.* Physiology & Behavior, 1967. **2**(4): p. 389-397.

60. Griffiths, R.R., et al., *Mystical-type experiences occasioned by psilocybin mediate the attribution of personal meaning and spiritual significance 14 months later.* Journal of psychopharmacology, 2008. **22**(6): p. 621-632.

61. Griffiths, R.R., et al., *Psilocybin can occasion mystical-type experiences having substantial and sustained personal meaning and spiritual significance.* Psychopharmacology, 2006. **187**(3): p. 268-283.

62. Horsley, R.R., et al., *Psilocin and ketamine microdosing: effects of subchronic intermittent microdoses in the elevated plus-maze in male Wistar rats.* Behavioural pharmacology, 2018. **29**(6): p. 530-536.

63.   Barker, D.J., et al., *Lateral preoptic control of the lateral habenula through convergent*

glutamate and GABA transmission. Cell reports, 2017. **21**(7): p. 1757-1769.

64.   Lerner, T.N., et al., *Intact-brain analyses reveal distinct information carried by SNc dopamine*

subcircuits. Cell, 2015. **162**(3): p. 635-647.

65.   Friard, O. and M. Gamba, *BORIS: a free, versatile open-source event-logging software for*

video/audio coding and live observations. Methods in ecology and evolution, 2016. **7**(11): p.

1325-1330.

Figure 1. (**A**) Experimental timeline. Animals arrived at 6-7 weeks old and were allowed to habituate for 1-week prior to surgeries. Bilateral injection of calcium sensor (jGCaMP7f) and implantation of fiber optic cannulas were conducted. After 3 weeks for viral incubation, animals underwent a series of fiber photometry experiments. (**B**) Representative coronal section showing fiber placement and GCaMP expression. (**C**) Mean air-puff traces showing consistency of CeA signal in response to the air-puff stimulus.

Figure 2. *Acute Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus Following Psilocin Administration* (**A**) Females: representative images showing c-Fos+ cells tagged with a green fluorescence protein (GFP) in the CeA. (**B**) Female vehicle vs. psilocin: histogram showing the number of c-Fos+ cells in each subregion of the CeA. CeC = capsular CeA, CeM =. Medial CeA, CeL = lateral CeA. Data points correspond to the mean of 2-4 hemispheres for each individual subject. 2-way ANOVA: Treatment x time: $F_{(2, 24)}$ = 1.597, p = 0.22; Treatment: $F_{(1, 24)}$ = 9.149, p = 0.006; Time: $F_{(1, 24)}$ = 9.149, p = 0.006 (**C**) Males: representative images showing c-Fos+ cells tagged with a green fluorescence protein (GFP) in the CeA. (**D**) Male vehicle vs. psilocin: histogram showing the number of c-Fos+ cells in each subregion of the CeA. CeC = capsular CeA, CeM =. Medial CeA, CeL = lateral CeA. Data points correspond to the mean of 2-4 hemispheres for each individual subject. 2-way ANOVA: Treatment x time: $F_{(2, 27)}$ = 1.241, p = 0.30; Treatment: $F_{(1, 27)}$ = 5.543, p = 0.03; Time: $F_{(2, 27)}$ = 12.02, p = 0.0002; Dunnett's: CeC (p = 0.01) (**E**) Female psilocin group: representative raw ΔF/F traces showing an individual at baseline and while on psilocin. Blue line = air-puff onset, Grey lines = individual traces, Green line = mean trace, Std= standard deviation (**F**) Females: Estimation plots for binned trace plots comparing baseline to treatment. Each data point corresponds to a 500ms bin averaged across subjects +/- SEM, Vehicle: p = 0.01, t (19) =2.780, mean of differences = 0.34; Psilocin: p = 0.01, t (19) =2.780, mean of differences = 0.34 (**G**) Female vehicle group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned window +/- S.E.M. (**H**) Female psilocin group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned window +/- S.E.M. (**I**) Male psilocin group: representative raw ΔF/F traces showing an individual at baseline and while on psilocin. Blue line = air-puff onset, Grey lines = individual traces, Green line = mean trace, Std= standard deviation (**J)** Males: Estimation plots for binned trace plots comparing baseline to

treatment. Each data point corresponds to a 500ms bin averaged across subjects +/- SEM. Vehicle: $p =$

0.0003, t (19) =4.452, mean of differences = 0.33; Psilocin: $p = 0.54$, t (19) =0.6231, mean of differences = -

0.06 **(K)** Male vehicle group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms

air-puff at 85 psi. Data points represent group averages within 500ms binned window +/- S.E.M. **(L)** Male

psilocin group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85

psi. Data points represent group averages within 500ms binned window +/- S.E.M. *p<0.05, **p<0.01, ***

*p<0.001, **** p<0.0001.*

Figure 3. *Prolonged Changes in CeA Reactivity in Response to an Aversive Air-Puff Stimulus Following*

Psilocin Administration **(A)** Female psilocin group: representative raw ΔF/F traces showing an individual at

baseline and at the 28-day follow up. Blue line = air-puff onset, Grey lines = individual traces, Green line =

mean trace, Std= standard deviation **(B)** First graph: area under curve (AUC) histogram plots comparing

baseline to the 28-day follow up. 2-way ANOVA: Treatment x Time: F (1, 14) = 0.003383, $p = 0.95$; Treatment:

F (1, 14) = 0.09635, $p = 0.76$; Time: F (1, 14) = 0.2126, $p = 0.65$; Second graph: changes in AUC at each of

the follow up time points between the vehicle and psilocin group. Mean AUC +/- S.E.M., 2-way ANOVA:

Treatment x Time: F (2, 28) = 1.437, $p = 0.25$; Treatment: F (1, 14) = 0.0003490, $p = 0.99$; Time: F (2, 28) =

3.067, $p = 0.06$ **(C)** Female vehicle group: air-puff trace plots of changes in CeA fluorescence following

exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned window +/-

S.E.M., 1-way ANOVA: F (3, 57) = 50.02, $p < 0.0001$; Dunnett's: 2- day ($p < 0.0001$) and 6-day ($p < 0.0001$**).**

(D) Female psilocin group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms

air-puff at 85 psi. Data points represent group averages within 500ms binned window +/- S.E.M., (3, 57) =

54.13, $p < 0.0001$; Dunnett's: 2-Day ($p < 0.0001$) and 6-day ($p < 0.0001$). **(E)** Male psilocin group:

representative raw ΔF/F traces showing an individual at baseline and while on psilocin. Blue line = air-puff

onset, Grey lines = individual traces, Green line = mean trace, Std= standard deviation **(F)** First graph: AUC

histogram plots comparing baseline to the 28-day follow up between male vehicle and psilocin groups. 2-way

ANOVA: Treatment x Time: F (1, 19) = 5.068, $p = 0.03$; Treatment: F (1, 19) = 0.2503, $p = 0.62$; Time: F (1,

19) = 1.361, $p = 0.26$, Šidák: 28-day ($p = 0.04$). Second graph: changes in AUC at each of the follow up time

points between the male vehicle and psilocin group. Mean AUC +/- S.E.M. 2-way ANOVA: Treatment x Time:

$F_{(2, 38)}$ = 0.2113, p = 0.81; Treatment: $F_{(1, 19)}$ = 4.926, p = 0.03; Time: $F_{(2, 38)}$ = 0.7906, p =0.46 **(G)** Male

vehicle group: air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85

psi. Data points represent group averages within 500ms binned window +/- S.E.M. 1-way ANOVA: $F_{(3, 57)}$ =

5.554, p = 0.002; Dunnett's: 28-day follow up (p = 0.01) **(H)** Male psilocin group: air-puff trace plots of changes

in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group averages

within 500ms binned window. 1-way ANOVA: $F_{(3, 57)}$ = 72.60, p < 0.0001; Dunnett's: 2-day (p < 0.0001), 6-

day (p < 0.0001), and 28-day (p < 0.0001) **(I)** Females: Summary histogram comparing the mean of all 20 time

bins following air-puff administration. Each data point represents an individual time bin within that condition

collapsed across all subjects. +/- S.E.M.; 2-way ANOVA: Treatment x time: $F_{(4, 152)}$ = 35.02, p < 0.0001;

Treatment: **(J)** Males: Summary histogram comparing the mean of all 20 time bins following air-puff

administration +/- S.E.M. Each data point represents an individual time bin within that condition collapsed

across all subjects. +/- S.E.M. *p<0.05, **p<0.01, *** p<0.001, **** p<0.0001.

Figure 4. *Acute and Prolonged Behavioral Effects of Psilocin.* **(A)** Females: locomotion plots comparing

vehicle and psilocin groups at each time point. Data points are mean time in seconds +/- SEM. 2-way ANOVA:

Treatment x Time: $F_{(3, 42)}$ = 0.2118, p = 0.89; Treatment: $F_{(1, 14)}$ = 2.238, p = 0.16; Time: $F_{(2.497, 34.95)}$

= 0.6065, p = 0.59 **(B)** Females: time spent rearing comparing vehicle and psilocin groups at each time point.

Data points are mean time in seconds +/- SEM. 2-way ANOVA: Treatment x Time: $F_{(4, 56)}$ = 1.345, p = 0.26;

Treatment: $F_{(1, 14)}$ = 4.433, p = 0.05 Time: $F_{(4, 56)}$ = 2.486, p = 0.05 **(C)** Males: locomotion plots comparing

vehicle and psilocin groups at each time point. Data points are mean time in seconds +/- SEM. 2-way ANOVA:

Treatment x Time: $F_{(3, 57)}$ = 0.1054, p = 0.96; Treatment: $F_{(1, 19)}$ = 2.563, p = 0.12; Time: $F_{(3, 57)}$ = 1.255,

p =0.29 **(D)** Males: time spent rearing comparing vehicle and psilocin groups at each time point. Data points

are mean time in seconds +/- SEM. 2-way ANOVA: Treatment x Time: $F_{(4, 76)}$ = 0.5546, p = 0.69; Treatment:

$F_{(1, 19)}$ = .7576, p = 0.39; Time: $F_{(4, 76)}$ = 1.190, p =0.32 **(E)** Females threat responding behavior. First

figure is showing percentage of subjects that darted following air-puff administration. Second figure is showing

the percentage of animals that remained immobile following air-puff. Third and fourth figure are pie charts

showing the percentage of animals that darted vs. remained immobile **(F)** Males threat responding behavior:

First figure is showing percentage of subjects that darted following air-puff administration. Fisher's exact test:

2-day (p = 0.03); Second figure is showing the percentage of subject that remained immobile following air-puff.

Fisher's exact test: 2-day (p = 0.03);; Third and fourth figure are pie charts showing the percentage of animals

that darted vs. remained immobile **(G)** Male vehicle passive responders (remained immobile) group: First figure

shows air-puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi.

Data points represent group averages within 500ms binned window +/- S.E.M.; Second figure is a histogram

showing differences in the mean peak ΔF/F value for each group +/- S.E.M following air-puff administration.

Data points reflect each individual subject within the corresponding subgroup. 1-way ANOVA: Treatment: $F_{(4,}$

$_{20)}$ = 1.219, p = 0.33; **(H)** Male Psilocin passive responders (remained immobile) group: First figure shows air-

puff trace plots of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points

represent group averages within 500ms binned window +/- S.E.M. Second figure is a histogram showing

differences in the mean peak ΔF/F value for each group +/- S.E.M following air-puff administration. Data points

reflect each individual subject within the corresponding subgroup. 1-way ANOVA: Treatment: $F_{(4, 20)}$ = 1.276,

p = 0.31 **(I)** Male vehicle active responders (darted) group: First figure shows air-puff trace plots of changes in

CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group averages

within 500ms binned window +/- S.E.M. Second figure is a histogram showing differences in the mean peak

ΔF/F value for each group +/- S.E.M following air-puff administration. Data points reflect each individual subject

within the corresponding subgroup. 1-way ANOVA: Treatment: $F_{(4, 12)}$ = 1.046, p = 0.42 **(J)** Male Psilocin

active responders (darted) group: First figure shows air-puff trace plots of changes in CeA fluorescence

following exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned

window +/- S.E.M. Second figure is a histogram showing differences in the mean peak ΔF/F value for each

group +/- S.E.M following air-puff administration. Data points reflect each individual subject within the

corresponding subgroup. 1-way ANOVA: Treatment: $F_{(4, 16)}$ = 6.566, p = 0.003

Figure 5. *Changes in CeA Reactivity in Response to an Auditory Stimulus Following Psilocin Administration in*

Female Subjects. **(A)** Female vehicle group: representative mean ΔF/F traces showing an individual at initial

auditory compared to final auditory. Blue line = air-puff onset, Grey lines = individual traces, Green line = mean

trace, Std= standard deviation **(B)** Female vehicle group: peak point of binned mean ΔF/F traces compared

between initial and final auditory recordings. Data points represent each individual peak ΔF/F value +/- SEM.

Paired t-test: $p = 0.004$, t (8) =3.934, mean of differences = -3.17; **(C)** Female vehicle group: air-puff trace plots

of changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group

averages within 500ms binned window +/- S.E.M **(D)** Female vehicle group: Estimation plots for binned trace

plots comparing baseline to treatment. Each data point corresponds to a 500ms bin averaged across subjects

+/- SEM.; Paired t-test: $p < 0.0001$, t (19) =18.28, mean of differences = -2.30; **(E)** Female psilocin group:

representative mean ΔF/F traces showing an individual at initial auditory compared to final auditory. Blue line =

air-puff onset, Grey lines = individual traces, Green line = mean trace, Std= standard deviation **(F)** Female

psilocin group: peak point of binned mean ΔF/F traces compared between initial and final auditory recordings.

Data points represent individual peak ΔF/F value +/- SEM. Paired t-test: $p = 0.67$, t (6) =0.4346, mean of

differences = -0.49; **(G)** Female psilocin group: air-puff trace plots of changes in CeA fluorescence following

exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned window +/-

S.E.M. **(H)** Female psilocin group: Estimation plots for binned trace plots comparing baseline to treatment.

Each data point corresponds to a 500ms bin averaged across subjects +/- SEM. Paired t-test: $p = 0.98$, t (19)

=0.01624, mean of differences = 0.002; *$p<0.05$, **$p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure 6. *Changes in CeA Reactivity in Response to an Auditory Stimulus Following Psilocin Administration in*

Male Subjects. **(A)** Male vehicle group: representative mean ΔF/F traces showing an individual at initial

auditory compared to final auditory. Blue line = air-puff onset, Grey lines = individual traces, Green line = mean

trace, Std= standard deviation **(B)** Male vehicle group: peak point of binned mean ΔF/F traces compared

between initial and final auditory recordings. Data points represent each individual peak ΔF/F value +/- SEM.

Paired t-test: $p = 0.06$, t (9) =2.136, mean of differences = -1.58 **(C)** Male vehicle group: air-puff trace plots of

changes in CeA fluorescence following exposure to a 500ms air-puff at 85 psi. Data points represent group

averages within 500ms binned window +/- S.E.M. **(D)** Male vehicle group: Estimation plots for binned trace

plots comparing baseline to treatment. Each data point corresponds to a 500ms bin averaged across subjects

+/- SEM. Paired t-test: $p < 0.0001$, t (19) =9.719, mean of differences = -0.85 **(E)** Male psilocin group:

representative mean ΔF/F traces showing an individual at initial auditory compared to final auditory. Blue line =

air-puff onset, Grey lines = individual traces, Green line = mean trace, Std= standard deviation **(F)** Male

psilocin group: peak point of binned mean ΔF/F traces compared between initial and final auditory recordings.

Data points represent individual peak ΔF/F value +/- SEM. Paired t-test: $p = 0.92$, t (10) =0.09932, mean of

differences = 0.09 **(G)** Male psilocin group: air-puff trace plots of changes in CeA fluorescence following

exposure to a 500ms air-puff at 85 psi. Data points represent group averages within 500ms binned window +/-

S.E.M. **(H)** Male psilocin group: Estimation plots for binned trace plots comparing baseline to treatment. Each

data point corresponds to a 500ms bin averaged across subjects +/- SEM. Paired t-test: $p < 0.0001$, t (19)

=4.943, mean of differences = -0.39; *$p<0.05$, **$p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Female

Males

A Locomotion

B Exploratory

C Locomotion

D Exploratory

E Female Threat Responding

F Males Threat Responding

Male Passive Responders

G Vehicle

H Psilocin

Male Active Responders

I Vehicle

J Psilocin

Females Vehicle

Females Vehicle

Females Psilocin

Females Psilocin

Males Vehicle

A

Initial Auditory

Final Auditory

B

C Vehicle Group

**** INITIAL FINAL

D Estimation Plot: Time Bins Post Air Puff

Males Psilocin

E

Initial Auditory

Final Auditory

F

G Psilocin Group

**** INITIAL FINAL

H Estimation Plot: Time Bins Post Air Puff

1