Divergence of trafficking and polarity mechanisms for PIN auxin

transporters during land plant evolution

Han Tang[1#], Kuan-Ju Lu[2#], You-Liang Cheng[3], Shih-Long Tu[3], and Jiří Friml[1*]

1. Institute of Science and Technology Austria (ISTA), Am Campus 1, 3400

Klosterneuburg, Austria

2. Graduate Institute of Biotechnology, National Chung Hsing University, No.

145, Xingda Rd., South Dist., Taichung 40227, Taiwan, R.O.C.

3. Institute of Plant and Microbial Biology, Academia Sinica, 128 Sec. 2,

Academia Rd, Nankang, Taipei 11529, Taiwan, R.O.C.

# Equal contribution, * Correspondence: jiri.friml@ista.ac.at

Abstract (~200 words)

The phytohormone auxin and its directional transport through tissues play a

fundamental role in plant development. This polar auxin transport relies on the

PIN auxin exporters, whose polarization at the plasma membrane determines

the overall direction of auxin flow. Hence, PIN polarization is crucial for

development, but its evolution during the rise of developmental complexity in

land plants remains unclear. Here, we investigated the evolution of PIN

trafficking and polarization by comparing two model

bryophytes, *Physcomitrella patens* and *Marchantia polymorpha*, with the

angiosperm model *Arabidopsis thaliana*. All examined PINs show a conserved

auxin export function, which is reflected by comparable bryophytic growth

defects in overexpression lines. In bryophytes, bryophytic PINs polarize to

filamentous apices, while Arabidopsis PINs distribute symmetrically on the

membrane with additional intracellular puncta. In the Arabidopsis root

epidermis, bryophytic PINs show no defined polarity. Pharmacological

interference revealed a strong cytoskeleton dependence of bryophytic but not

Arabidopsis PIN polarization, which, in contrast, strongly relies on Brefeldin

A-sensitive trafficking. The divergence of PIN polarization and trafficking is

also observed within the bryophyte clade and between tissues in individual

species. These results collectively reveal a divergence of PIN trafficking and

polarity mechanisms throughout land plant evolution and suggest co-evolution

of PIN sequence-based and cell-based polarity mechanisms.

Introduction

Auxin is a crucial regulator of polarity and morphogenesis in land plants (Mockaitis and Estelle, 2008; Smit and Weijers, 2015; Kato et al., 2018; Leyser, 2018; Yu et al., 2022). The auxin concentration gradient through tissues coordinates a broad spectrum of plant development, ranging from embryogenesis to organ formation and tropic growth (Vanneste and Friml, 2009; Friml, 2021). The establishment of the auxin gradient relies on particular auxin transporters, including the efflux carriers PIN-FORMED (PINs). In different tissues, specific PINs are polarized at different domains of the cells, thus directly driving the directional auxin flow across tissues (Adamowski and Friml, 2015). Therefore, given the essential impact of auxin flow in various developmental processes, the function and polarization of PIN proteins are crucial for maintaining the correct patterns of plant growth (Sauer and Kleine-Vehn, 2019).

PINs are found in all land plants and can be traced back to charophytic green algae (Krecek et al., 2009; Skokan et al., 2019). The functional conservation of PIN proteins in the auxin transport has also been demonstrated by exogenously expressing PINs from the green algae *Klebsormidium flaccidum*, the moss *Physcomitrium (Physcomitrella) patens*, and the angiosperm *Arabidopsis thaliana* in transgenic plants and heterologous systems (Zourelidou et al., 2014; Skokan et al., 2019). Additionally, when exogenous PINs from charophytes or Arabidopsis are overexpressed in Physcomitrella, the transgenic plants show growth inhibition that resembles auxin deprivation phenotypes (Viaene et al., 2014; Lavy et al., 2016; Tao and Estelle, 2018; Skokan et al., 2019). Therefore, these observations support the idea that PIN-mediated polar auxin transport may

govern plant development since the emergence of land plants.

The mechanistic studies of PIN polarity regulation have been extensively

investigated in the angiosperm Arabidopsis model. Canonical PINs, featuring a

long hydrophilic loop between two transmembrane domains, are first delivered

via the ER-Golgi vesicle trafficking pathway to the plasma membrane (PM).

De-polymerization of actin filaments induces the accumulation of PIN-labelled

small intracellular puncta near the PM, but with no apparent PIN polarity defect

(Geldner et al., 2001; Glanc et al., 2018, 2019). Microtubules are involved in

the cytokinetic trafficking of PINs, but are not required for the polarity

establishment or maintenance at the PM of non-dividing cells (Geldner et al.,

2001; Kleine-Vehn et al., 2008b; Glanc et al., 2019). Notably, the disruption of

both cytoskeleton networks delays, but does not abolish AtPIN2 polarization in

Arabidopsis epidermal cells (Glanc et al., 2019). This suggests that the

cytoskeleton networks are involved, but not exclusively essential for PIN polar

trafficking, while other mechanisms contribute to the PIN polar localization.

Once PINs are present on the PM, different processes, such as reducing

lateral diffusion and enhanced protein clustering, coordinate to maintain the

polar localization of PINs (Kleine-Vehn et al., 2011; Li et al., 2021). Recycling

of PINs from and to the PM also reinforces their polarity (Sauer and

Kleine-Vehn, 2019). During the PM recycling, PINs are first internalized

through the clathrin-dependent endocytosis (Dhonukshe et al., 2007), and

endocytic PINs are subsequently sorted out via a GNOM ARF-GEF pathway

(guanine nucleotide exchange factor) (Steinmann et al., 1999; Kleine-Vehn et

al., 2008b; Narasimhan et al., 2021). The GNOM ARF–GEF was shown to

interact with *trans*-Golgi network/early endosome (TGN/EE) and thereby

regulate the recycling of PINs (Naramoto et al., 2014). In Arabidopsis, the

fungal toxin Brefeldin A (BFA) inhibits the return of PINs from endocytic

vesicles by blocking GNOM functions through unknown mechanisms. It thus

leads to the accumulation of PINs in an enlarged endomembrane system, the

so-called BFA bodies (Geldner et al., 2001).

PIN polarization was also observed in the moss model *Physcomitrella*

patens. Physcomitrella grows as filamentous protonemata with tip cell growth,

and PpPINA- and PpPINB-GFP show polar localization at the tip of protonema

cells (Bennett et al., 2014; Viaene et al., 2014). However, their polar

localization is not robust in other species. When PpPINA and PpPINB are

expressed in Arabidopsis epidermal cells, where AtPIN2 presents clear apical

localization, both Physcomitrella PIN proteins are mislocalized at basal and

apical sites (Zhang et al., 2019). Furthermore, PINs from the liverwort

Marchantia polymorpha and the green algae *Klebsormidium flaccidum* are also

mislocalized in epidermal cells of Arabidopsis (Zhang et al., 2019). These

distinctive PIN localizations in different species suggest that mechanisms for

PINs trafficking and polarization may have diversified after the emergence of

land plants.

Despite the profound significance of PIN polarization and the resulting

directional auxin transport for land plant development, PIN trafficking and

polarization mechanisms are mainly derived from the angiosperm model

Arabidopsis thaliana. This research conducted a cross-species investigation to

elucidate mechanisms and evolution of PIN trafficking and polarization by

comparing two model bryophytes, *Physcomitrella patens* and *Marchantia*

polymorpha with the model angiosperm *Arabidopsis thaliana*. In bryophytes

and Arabidopsis, endogenous PINs show different trafficking and polarization

patterns in a range of tissues, suggesting that tissues may utilize diverse

trafficking pathways for PIN localization. Pharmacological interference of

known Arabidopsis trafficking mechanisms further suggests that bryophytes

employ a different trafficking machinery for PIN polarization. Furthermore,

Arabidopsis PINs cannot be recruited by bryophyte trafficking pathways and

vice versa, confirming a species specificity for PIN polarity regulations. Overall,

this study revealed the divergence of PIN trafficking and polarity mechanisms

between bryophytes and angiosperms.

Results

Physcomitrella PINs are polarized via different cytoskeleton-dependent

trafficking pathways

Physcomitrella has a filamentous protonema stage and a leafy gametophore

stage in its life cycle. The polarization of PpPINA and PpPINB at the tips of

filamentous protonema cells has been demonstrated before (Viaene et al.,

2014). However, the underlying mechanism and whether it engages the same

trafficking pathways for the polarization of PINs in angiosperms, such as

Arabidopsis, remains unclear.

A stable Physcomitrella transgenic line carrying PpPINA genomic DNA

fused with GFP driven by its promoter (PpPINA-GFP) was used as the

standard model line throughout this research. However, we used a

constitutively over expressor maize actin promoter to drive PpPINB-GFP since

its expression under the *PpPINB* promoter is undetectable (Viaene et al.,

2014).

We observed the PpPINA-GFP localized at the PM of the protonema tip

with a clear polarity (Fig. 1A). Interestingly, PpPINB-GFP showed different

polarization patterns from PpPINA-GFP. In addition to a PM localization at the

filament tip, most of the PpPINB-GFP signal was accumulated at the dome

area of the protonema cell with a cytosolic diffusion (Fig. 1B). To examine

whether the polar localization of PpPINs appears in multicellular tissues, we

also observed their localizations in gametophores. In gametophytic leaves,

both PpPINA- and PpPINB-GFP were non-polarly distributed on the PM with

additional cytosolic puncta (Fig. S1).

Next, we examined the role of the cytoskeleton networks in the

polarization of PpPINs. To disrupt actin filaments or microtubules,

Physcomitrella tissues were treated with Latrunculin B (LatB) or Oryzalin (Ory),

respectively. Surprisingly, the abolishment of actin filaments resulted in the

hyper-polarization of PpPINA-GFP (Fig. 1A, 1C), accumulating at a focal locus

to the very tip of the cell (Fig. 1C-D). The disruption of microtubules resulted in

less accumulation of PpPINA-GFP at the tip of protonemata, and the

PpPINA-GFP appeared to be detached from the PM (Fig. 1A, 1C). Notably,

none of the drug treatments affected the polar localization of PpPINB-GFP (Fig.

1B). This result suggests that PpPINA and PpPINB are polarized through

different trafficking pathways in moss.

Arabidopsis AtPIN1 and AtPIN2 are not polarized in Physcomitrella

The intriguing divergence between PpPINA-GFP and PpPINB-GFP

polarization mechanisms prompted us to test whether it is a species-specific

mechanism. To achieve this, we generated Physcomitrella transgenic lines,

carrying Arabidopsis AtPIN1- and AtPIN2-GFP driven by *XVE* inducible

promoter (Fig. S2) (Kubo et al., 2013). In protonemata, both AtPIN1-GFP and

AtPIN2-GFP showed an even distribution on the PM without apparent polarity,

but they also displayed numerous small intracellular puncta (Fig. 1E, S3). The

dependency of the cytoskeletons was tested with LatB or Ory treatments,

however, no obvious effects on AtPIN1-GFP and AtPIN2-GFP PM localization

and cytosolic puncta were observed (Fig. 1E, S3). These observations indicate

that Arabidopsis PINs cannot be recruited into the moss polar trafficking

pathways and suggest a divergence of polarity mechanisms during the

evolution of land plants.

Different modes of Brefeldin A action on trafficking in Physcomitrella

BFA is a commonly used inhibitor (Geldner et al., 2001) to disrupt the PM

recycling of Arabidopsis PINs, resulting in the large intracellular BFA bodies

(Narasimhan et al., 2021). Therefore, we opted BFA to examine the PM

recycling of PpPINs in Physcomitrella. We first tested whether BFA has similar

effects in Physcomitrella by treating wild-type plants with BFA, followed by

labelling with the membrane dye FM4-64. Unlike in Arabidopsis, no clear BFA

bodies were observed (Fig. S4). Instead, the BFA-treated plants showed much

less internal FM4-64 particles compared to the DMSO control (Fig. S4), which

suggests an inhibitory effect of BFA on endocytosis in Physcomitrella.

Accordingly, PpPINA-GFP and PpPINB-GFP showed no striking changes in

subcellular distribution after BFA treatment (Fig. 2A). In contrast, AtPIN1-GFP

appeared to form larger intracellular puncta in response to BFA, resembling

BFA bodies in Arabidopsis (Fig. 2A).

The response of AtPIN1-GFP to BFA treatment suggests the presence of

a BFA-sensitive PM recycling in Physcomitrella. To further test whether

PpPINs and AtPINs undergo constitutive BFA-sensitive PM recycling, we

conducted fluorescence recovery after photobleaching (FRAP) experiments in

combination with BFA treatment. The PpPINA-GFP line was treated with the

protein synthesis inhibitor cycloheximide (CHX) with or without BFA present.

We performed FRAP analysis on a specific PM region near the filamentous tip

(Fig. 2B, white arrowheads). The recovery rate of PM-localized PpPINA-GFP

showed no difference in the presence or absence of BFA, suggesting that

PpPINA-GFP might not use the BFA-sensitive PM recycling pathway (Fig. 2B,

2D). In contrast to PpPINA-GFP, AtPIN1-GFP showed a slower recovery rate

when endocytosis was inhibited by BFA (Fig. 2C-2D), which is consistent with

the idea that AtPIN1-GFP might undergo the BFA-sensitive PM recycling in

Physcomitrella.

These observations show that Physcomitrella PINs do not use

BFA-sensitive PM recycling pathways, whereas heterologous expressed

Arabidopsis PINs do. This suggests that the functional domain of Arabidopsis

PINs can be recognized and recruited to a heterologous BFA-sensitive PM

recycling machinery in Physcomitrella.

Overexpression of PINs inhibits the chloronema-caulonema transition in

Physcomitrella

Because polarized PIN proteins act as auxin exporters to directly guide polar

auxin transport, they mediate many aspects of plant growth and development

(Friml, 2021). We therefore tested whether PpPINA- and PpPINB-GFP with

distinct polar localizations generate distinct gain-of-function phenotypes. To

investigate this, we generated overexpression lines by driving PpPINA- and

PpPINB-GFP under the control of the maize actin promoter. However, the

PpPINA-GFP over-expression line showed a robust cytoplasmic signal and

almost no signal on the PM (Fig. S5). To overcome this, we used the

pro*PINA*::PpPINA-GFP transgenic line, which is generated in a wild-type

background and carries two copies of *PpPINA*, and is thus an overexpression

plant.

During early development, Physcomitrella gradually transits its

filamentous protonemata from thicker/shorter chloronema cells with

perpendicular division planes to thinner/longer caulonema cells with oblique

division planes (Rensing et al., 2020). Overexpression of PpPINA- and

PpPINB-GFP led to similar defects in this chloronema-caulonema transition

with a shorter cell length of the subapical cell and a smaller division angle (Fig.

3A-C). No significant differences were observed between PpPINA- and

PpPINB-GFP overexpression. The inhibitory effect on the protonema transition

was also reflected in a smaller colony size in both PpPINA- and PpPINB-GFP

overexpression lines (Fig. 3D) (Viaene et al., 2014). This data indicates that

despite different localization patterns, Physcomitrella PINs produced

comparable gain-of-function phenotypes by inhibiting chloronema-caulonema

transition.

To test whether non-polar localized AtPIN1- and AtPIN2-GFP also result in

the same growth phenotypes in Physcomitrella, we analyzed the protonema

transition of inducible AtPIN1-GFP and AtPIN2-GFP lines. The

chloronema-caulonema transition was inhibited by overexpressing either

AtPIN1-GFP or AtPIN2-GFP (Fig. 3A-C). Furthermore, after induction, the

colony size of AtPIN1- and AtPIN2-GFP was much smaller than the wild-type

control, which showed a similar size to the DMSO treated wild-type plants (Fig.

3D). This result demonstrates the inhibitory effect of AtPINs overexpression on

the chloronema-caulonema transition. In addition to the protonema transition,

the overexpression of AtPIN1-GFP and AtPIN2-GFP led to wavy filaments and

multiple branching phenotypes, which was not observed in PpPINA- and

PpPINB-GFP overexpression lines (Fig. 3E).

These results thus indicate that regardless of their polarization, the

overexpression of PM-localized PINs likely leads to auxin depletion in

Physcomitrella filaments and thereby inhibits the chloronema-caulonema

transition.

Marchantia MpPINZ is polarized at rhizoid tips but not in gemma cells

The divergence of PIN polarity and trafficking between Physcomitrella and

Arabidopsis suggests an evolution deviation between bryophytes and

angiosperms. To confirm this notion, we looked at another bryophyte, the

liverwort *Marchantia polymorpha*. Gemmae are the asexual reproductive

progenies of Marchantia. After water imbibition, rhizoids emerge from the large

rhizoid precursor cells on the epidermis of the gemma. To analyze the

subcellular localization of the sole canonical PIN in Marchantia, MpPINZ, we

generated a pro*35S*::MpPINZ-GFP transgenic Marchantia line. Interestingly,

MpPINZ-GFP was localized on the PM with small intracellular puncta and no

apparent polarity in all gemma epidermal cells (Fig. 4A). However, when

gemmae were stimulated to grow rhizoids, the signal of MpPINZ-GFP

accumulated at the protrusion site of emerging rhizoids in the rhizoid precursor

cells (Fig. 4B, blue arrows). Later after their emergence, MpPINZ-GFP was

polarized at the tip of young rhizoids (Fig. 4C), whereas the signal diminished

when the rhizoids elongated.

We used the same pharmacological interference as in Physcomitrella to

investigate whether the MpPINZ-GFP PM localization or polarization relies on

the cytoskeleton networks and BFA-sensitive PM recycling. In gemmae, the

disruption of either microtubules or actin filaments did not affect the PM

localization, but the number and the size of intracellular puncta increased (Fig.

4D-F). In young rhizoids, MpPINZ-GFP remains polarized at the tip when actin

filaments were disrupted, while the disruption of microtubules resulted in the

dislocation of the polarized MpPINZ-GFP (Fig. 4D).

To test the BFA effect in Marchantia, we applied FM4-64 labeling to Tak-1

wild-type plants. The endocytic FM4-64 accumulated in significantly larger

intracellular compartments after BFA treatment (Fig. S6), suggesting that BFA

inhibits the return pathway of PM recycling in Marchantia as it does in

Arabidopsis. The BFA treatment of MpPINZ-GFP resulted in a small increase

of particles with larger size (Fig. 4D-F), indicating that MpPINZ-GFP does not

intensively go through the BFA-sensitive PM recycling. Consistent with the

notion, in young rhizoids, MpPINZ-GFP remained its polarization at the tip of

the filament with BFA treatment (Fig. 4D). Overall, these results reveal that

BFA-sensitive PM recycling and PIN polarity mechanisms also diverge within

the bryophyte clade.

Arabidopsis PINs are not polarized in Marchantia

We demonstrated that AtPIN1- and AtPIN2-GFP appeared non-polarized and

revealed cytosolic puncta in Physcomitrella. To investigate whether these

features are conserved in bryophytes, we introduced AtPIN1- and AtPIN2-GFP

driven by *35S* promoter into Marchantia. In gemma epidermal cells, the

AtPIN1-GFP showed non-polar PM localization with a high cytosolic signal (Fig.

5A). In emerging rhizoids, AtPIN1-GFP did not show visible polar localization

in rhizoid precursor cells (Fig. 5B). In young rhizoids, AtPIN1-GFP was evenly

distributed on the PM with homogenized cytosolic signals (Fig. 5C). The signal

of AtPIN2-GFP was much weaker but showed similar patterns as AtPIN1-GFP

in gemma epidermal cells (Fig. S7). However, we could not detect any signal of

AtPIN2-GFP in rhizoid cells, we therefore only used AtPIN1-GFP for further

study.

Next, we applied the same cytoskeleton-disruptive drugs to Marchantia

transgenic plants but we did not observe any effect on AtPIN1-GFP localization

in all examined tissues. We then tested whether the inhibition of BFA-sensitive

PM recylcing affects the localization of AtPIN1-GFP. With a range of different

concentrations and durations of BFA treatment, AtPIN1-GFP showed the same

PM and cytosolic signals as the non-treated control (Fig. S8), which indicates

that the non-polar localization of AtPIN1-GFP in Marchantia is insensitive to

BFA. These results demonstrate that Arabidopsis PINs in Marchantia do not

employ the same trafficking machinery as MpPINZ, and thus reinforce the

notion of divergence in PINs trafficking and polarity regulations between

bryophytes and angiosperms. Furthermore, the distinct localization patterns of

AtPIN-GFPs in moss and liverwort also verify the divergence of PIN trafficking

mechanisms within the bryophyte clade.

Overexpression of PINs leads to a vertical growth of the thallus in

Marchantia

Different localization patterns of MpPINZ- and AtPIN1-GFP in Marchantia

prompted us to investigate whether these differences result in divergent

growth phenotypes. The wild-type Marchantia has prostrate thalli; however,

with the overexpression of either MpPINZ-GFP or AtPIN-GFPs, the thallus

grew more vertically, as apparent from the side view (Fig. 6A). The angles

between the thallus and the horizontal agar medium were measured to

quantify the vertical growth (Fig. 6B). The overexpression of MpPINZ-GFP

caused the most striking phenotype, but the overexpression of AtPIN-GFPs

also resulted in a vertical growth that showed significant differences from the

wild-type. Notably, despite low expression levels, the overexpression of

AtPIN2-GFP still resulted in the vertical growth of the thallus. This data shows

that, similar to results in Physcomitrella, MpPINZ and AtPINs with different

subcellular localization, can trigger similar physiological responses.

Bryophytic PINs utilize different trafficking mechanisms in Arabidopsis

Since our results indicated that bryophytic PINs in their original species are

delivered to the PM through different trafficking pathways from the known

routes in Arabidopsis, we next addressed whether bryophytic PINs can utilize

Arabidopsis trafficking machinery. To analyze bryophytic PINs, we observed

the protein localization at epidermal cells of transgenic Arabidopsis plants

expressing pro*PIN2*::AtPIN2-, MpPINZ-, PpPINA-, and PpPINB-GFP (Zhang

et al., 2019). As previously described, AtPIN2-GFP showed apical localization

in the epidermal cells, while MpPINZ-GFP mainly localized on the basal side.

PpPINA-GFP localized on both the apical and basal sides, and PpPINB-GFP

presented not only PM localization but also intracellular signals (Fig. 7) (Zhang

et al., 2019).

The disruption of actin filaments by LatB resulted in an increasing

appearance of AtPIN2-GFP signals in the cell interior, while all bryophytic

PIN-GFPs showed hypersensitivity reflected by their accumulation in larger

intracellular compartments (Fig. 7). Ory was used to depolymerize

microtubules and caused a minor effect on AtPIN2-GFP and PpPINs, while the

MpPINZ-GFP in turn accumulated in large intracellular compartments (Fig. 7).

Following the inhibition of ARF GEF-mediated trafficking by the BFA, all three

bryophytic PINs accumulated in BFA bodies similar to AtPIN2-GFP. To verify

whether the PIN-GFP signal in the BFA bodies is recycled from the PM, we

inhibited the protein synthesis with CHX application prior to BFA treatment.

Surprisingly, the accumulated MpPINZ-GFP signal was not visible when CHX

was supplied, indicating that the intracellular accumulated signals of

MpPINZ-GFP was originated mainly from the pool of newly synthesized

proteins. Our results therefore suggest that the bryophytic PINs traffic to the

PM through divergent trafficking routes in Arabidopsis, and not all of them

undergo intensive BFA-sensitive PM recycling.

Bryophytic PINs are polarized at the tip of initial root hairs

As the polarity of bryophytic PINs in their original species mainly appeared at

the tip of filamentous tissues, we questioned whether the bryophytic PINs are

polarized in similar filamentous tissues in Arabidopsis; e.g. root hairs. We

observed bryophytic PIN-GFP signals in the initial and elongated root hairs.

Notably, all examined PIN-GFPs, including AtPIN2-GFP and bryophytic

PIN-GFPs, revealed a polar localization at the tip of initial root hairs. However,

the polarity of PIN signals disappeared in the elongated hairs (Fig. 8).

Next we tested whether this polar localization relies on the cytoskeleton

networks or is mediated through BFA-sensitive PM recycling. Surprisingly, all

treatments that disrupt cytoskeleton networks or PM recycling did not affect the

polarization of PINs on the initial root hairs (Fig. 8), which suggests that

different trafficking machinery are engaged in Arabidopsis epidermis and root

hairs.

In summary, these observations also confirm the divergence of trafficking

and PM localization mechanisms between bryophytes and angiosperms. In

addition, different trafficking pathways operate in distinct tissues, even within

individual species.

Discussion

PIN polarity mechanisms display tissue-specific modifications

In Arabidopsis, despite the functional redundancy of PIN proteins in auxin

transport, PINs show different polarization in different tissues (Vieten et al.,

2005). For example, AtPIN2 is apicalized in epidermal cells, while it is localized

on the basal side of young cortex cells (Kleine-Vehn et al., 2008a). The

observations imply that different tissues possess specific trafficking pathways

for the same PIN protein. In line with this, our results show that in filamentous

cells, e.g., protonemata in Physcomitrella, rhizoids in Marchantia, and root

hairs in Arabidopsis, endogenous PIN proteins are polarized at the tip of apical

cells. Nevertheless, the different dependency of cytoskeletons suggests that

the polarized trafficking mechanisms diversified between the three examined

species (Fig. 1, 4, 8). On the contrary, in multicellular tissues, e.g., leaves in

moss, thalli in Marchantia, and primary roots in Arabidopsis, the bryophytic PIN

proteins tend to distribute evenly on the PM (Fig. S1, 4, 7). Unlike in

filamentous tissues, the disruption of cytoskeleton networks shows minor

effect on the PINs distribution in multicellular tissues (Fig. S1, 4, 7). This data

shows that PIN polarity and trafficking mechanisms have evolved with specific

modifications in different tissues and cell types, both in angiosperms and

early-diverging land plants. This likely reflects different requirements for

directional auxin transport in different developmental contexts and implies a

co-evolution of PIN sequence-based signals and cell type-specific polar sorting

and trafficking mechanisms.

Diversification of PIN polarity mechanisms during land plant evolution

The core mechanisms for auxin biosynthesis, auxin signaling, and PIN-mediated auxin transport are evolutionary conserved across land plants (Kato et al., 2018; Sauer and Kleine-Vehn, 2019; Blazquez et al., 2020). Our cross-species studies addressed to what extent the mechanism of PINs trafficking is conserved during the evolution of land plants. It revealed that exogenous PINs, such as Arabidopsis PINs in bryophytes and bryophytic PINs in Arabidopsis, could be delivered to the PM but they cannot develop a correct polarity and also show different sensitivities to interference with trafficking or cytoskeletons (Fig. 1, 5, 7). These observations clearly show a divergence of polarity mechanisms between bryophytic and vascular PINs. Notably, endogenous PINs showed different responses to cytoskeletal disruption within the bryophyte clade, indicating that trafficking pathways have already been diversified between moss and liverwort (Fig. 1 and 4). Our study thus demonstrates that PINs trafficking and polarity mechanisms are not conserved throughout plant evolution. Instead, different plant species evolved specialized pathways to deliver PINs and maintain their polarity at the PM. This is likely linked to an increasing repertoire of auxin transport developmental roles adopted with increasing morphological complexity during land plant evolution.

Diversification of PIN trafficking mechanisms during land plant evolution

Our comparative and interspecies complementation studies revealed different cytoskeleton requirements for PIN polarization between bryophytes and angiosperms. In Physcomitrella, both microtubules and actin filaments are involved in the polarization of PpPINA-GFP to the tip of filaments (Fig. 1). In

Marchantia, only the disruption of microtubules disturbed the polar localization

of MpPINZ-GFP at the tip of young rhizoids (Fig. 4). In contrast, in Arabidopsis,

disruption of neither microtubules nor actin filaments would abolish the

AtPIN2-GFP apical localization (Glanc et al., 2019; Fig. 7). Instead, the

polarization of Arabidopsis PINs relies on several cellular processes, including

PM recycling, reduced lateral diffusion, and clustering (Kleine-Vehn et al., 2011;

Sauer and Kleine-Vehn, 2019). In line with this, we found that AtPIN1-GFP is

accumulated in larger intracellular puncta in BFA-treated Physcomitrella (Fig.

2), demonstrating the engagement of BFA-sensitive PM recycling routes in

AtPIN1 trafficking.

BFA was shown to preferentially cause intracellular accumulation of the

basal AtPIN2 but not the apical AtPIN2 (Kleine-Vehn et al., 2008b), which

supports that delivery to different polar domains at the PM may engage diverse

trafficking pathways. Notably, in the Arabidopsis epidermis, PpPINA-GFP

shows the same mislocalization patterns as the MpPINZ-GFP, but they display

distinct responses to the disruption of BFA-sensitive PM recycling (Fig. 7). This

observation implies that the distinct responses to BFA are probably due to not

only different make-up of trafficking pathways in different species, but also to

the differences in PIN molecules *per se*.

Overall, our data collectively show a strong sensitivity of bryophytic

PINs to the cytoskeletal disturbance and of Arabidopsis PINs to the BFA

treatment in different species. This finding suggests a gradual shift in the

dependence of PIN polarity and trafficking from the cytoskeleton-dependent

pathways towards the ARF GEF-mediated trafficking along the land plant

evolution.

Co-evolution of polar trafficking mechanisms and PIN sequence-based determinants

One of the polarity maintenance mechanisms of PIN proteins is the ability to recycle from and to the membrane (Kleine-Vehn et al., 2011). In Physcomitrella, FRAP analysis demonstrated that PpPINA-GFP does not use the BFA-sensitive PM recycling pathway, while AtPIN1-GFP does (Fig. 2). This data demonstrates the BFA-sensitive PM recycling machinery already existed in Physcomitrella. In Marchantia, BFA treatment led to the accumulation of internalized FM4-64 dye in intracellular compartments (Fig. S6), which resembles the BFA body in Arabidopsis, and thus again implicates that the BFA-sensitive PM recycling is present in Marchantia. However, neither MpPINZ-GFP nor AtPIN1-GFP accumulated in the BFA bodies in Marchantia (Fig. 4, S8). These results demonstrate that BFA-sensitive, ARF GEF-dependent PM recycling pathways emerged since bryophytes; however, neither PpPINs nor MpPINZ employs these machineries in the original species, despite the fact that PpPIN-GFPs can utilize them in Arabidopsis (Fig. 7).

These observations further confirm that both trafficking mechanisms and relevant PIN sequences underwent a complex co-evolution during the gradual rise of morphological complexity in land plants.

Methods

Plant growth and transformation

Arabidopsis seeds were surface sterilized and grown on 1/2 MS plates. After a two-day stratification at 4°C, seedlings were grown under the long-day condition (16 hours light, 8 hours dark) at 22°C with 100-120 μmol photons $m^{-2}s^{-1}$ white light. For moss, all transgenic and WT plants were cultured on

BCD medium plates in a growth chamber at 24°C under the long-day condition

(16 hours light, 8 hours dark) with 35μmol photons $m^{-2}s^{-1}$ white light. For

Marchantia, WT and all transgenic plants were cultured on 1/2 B5 plates in a

growth chamber under the long-day condition (16 hours light, 8 hours dark) at

22°C with 50-60 μmol photons $m^{-2}s^{-1}$ white LED light.

All transgenic plants used in this study are listed in Supplemental Table 1.

Physcomitrella transgenic plants pro*PINA*::PINA-GFP and pro*ACT*::PINB-GFP

have been generated and verified as previously reported (Viaene et al., 2014),

and the inducible overexpression AtPIN1-GFP and AtPIN2-GFP lines were

generated as described in previous studies (Yamada et al., 2016; Tang et al.,

2020). In brief, the Gransden 2004 WT moss plants were freshly propagated

mosses and transformed via the PEG-mediated transformation (Nishiyama et

al., 2000). The transformants were selected under Hygromycin (20μg/mL), and

two independent transgenic lines were selected for imaging and analysis.

pro*35S*::MpPINZ-, AtPIN1-, and AtPIN2-GFP Marchantia plants were

generated via the Agrobacterium transformation method described before

(Kubota et al., 2013). In brief, the apical meristemic region of each

two-week-old Takaragaike-1 (Tak-1) thallus was removed and further cut into

eight pieces. After culturing on 1/2 B5 with 1% sucrose agar plates for 3 days,

the cut thalli were transferred to 50ml 0M51C medium with 200 μM

acetosyringone (4'-Hydroxy-3',5'-dimethoxyacetophenone) in 200ml flasks

with agitation at 130 rpm to coculture with OD_{600} = 1 density agrobacteria

harboring the target construct for another 3 days. The transformed thalli were

washed and plated on 1/2 B5 plates with proper antibiotic selection.

Independent T1 lines were isolated, and G1 lines from independent T1 lines

were generated by sub-cultivating single gemmalings, which emerged

asexually from a single initial cell (Shimamura, 2016). The next generation of

G1, called the G2 generation, was used for analyses.

Arabidopsis transgenic lines bearing bryophytic PIN-GFPs under *AtPIN2*

promoter control were generated and used as in a previous study (Zhang et al.,

2019). For root imaging, seeds were sown on 1/2 Murashige–Skoog (MS)

medium plates and kept at 4°C for 2 days and moved to the growth chamber to

culture vertically for another 4 days.

Plasmid construction

Plasmids and primers for construction and genotype confirmation are listed in

Supplemental Table 2. For transgenic lines with inducible overexpression by

moss, the cDNA of *AtPIN1* and *AtPIN2*, which inserted the *GFP* gene into the

hydrophilic loop at 461bp in *AtPIN1* and 736bp in *AtPIN2*, were amplified from

previously generated plasmids (Zhang et al., 2019) by PCR and cloned into

the gateway entry plasmid pENTR/D-TOPO. The fragments were sub-cloned

into the pPGX8 vector, which contains a pro*35S* driven β-estradiol inducible

XVE cassette (Nakaoka et al., 2012; Kubo et al., 2013; Floriach-Clark et al.,

2021) via a Gateway® LR reaction (Invitrogen) as the manufacture's

recommendation.

To generate a pro*35S*::MpPINZ-, AtPIN1-, and AtPIN2-GFP construct, the

cDNA of the target fragment containing proper *GFP* insertion were amplified

with the primers listed in Supplemental Table 2. The amplified fragments were

cloned into the pENTR/D-TOPO® vector (Invitrogen) with the protocol supplied

by the manufacturer. Plasmids with target genes were further sub-cloned into

the pMpGWB102 vector containing a *35S* promoter (Ishizaki et al., 2015) by

the   Gateway®   LR   reaction   (Invitrogen)   as   the   manufacturer's

recommendation.

Microscopy

For moss observation, protonemata were cultured in glass-bottom dishes

covered by BCD agar medium for 6 – 7d. The live-cell imaging was then

performed on a Leica SP8X-SMD confocal microscope equipped with a hybrid

single-molecule detector (HyD) and an ultrashort pulsed white-light laser (WLL;

50%; 1 ps at 40 MHz frequency). The Leica Application Suite X was used as a

software platform, and imaging was conducted with a HC PL APO CS2

40x/1.20 water immersion objective. The following settings were used: scan

speed of 400 Hz, resolution of 1024 x 1024 pixels and standard acquisition

mode of the hybrid detector. The time-gating system was activated to avoid the

autofluorescence emitted from chloroplasts. For the filament growth assay, the

imaging dish was applied with FM4–64 (Invitrogen) solution for 10-30 mins,

and the 10x objective lens was used.

The FRAP analysis was performed under similar conditions with some

modifications. After capturing the first image as a "before" image, the HyD

detector was closed, and 2 lasers with full power were turned on. We

performed 8 – 10 times imaging capture with a 15-fold zoom-in that focused on

the bleaching area to ensure the bleaching efficiency. The bleaching area was

selected to cover at least one-third of the enriched signal on the membrane of

apical cells. After bleaching, a time series imaging was carried out in the same

setting as the "before" image with 1 min interval for another 8 minutes.

For Marchantia observation, gemmae were picked from a gemma cup and

transferred into a 24-well plate with 500 µl liquid 1/2 B5 medium. After culturing

in the growth chamber for 24 hours, gemmae from each sample were

transferred onto a 24mm x 50mm coverslip sandwich and observed on a slide

under Leica Stellaris 8 system with HyD detectors and the ultrashort pulsed

white-light laser (WLL; 70%; 1 ps at 40 MHz frequency). The Leica Application

Suite X was used as a software platform, and imaging was conducted with an

HC PL APO CS2 40x/1.20 water immersion objective. The following settings

were used: scan speed of 400 Hz and resolution of 1024 x 1024 pixels. For

GFP-containing images, a 488nm white light laser was selected, and the

detection range was set between 500nm to 525nm. For FM4-64 imaging, the

white light laser was set at 515nm for excitation, and the detection range was

set between 600nm to 700nm. The tau-gating model, harvesting photons with

1.0-10.0ns lifetime, was used for all Marchantia imaging to avoid

autofluorescence emitted from the chloroplasts. For the surface section

(rhizoid precursor cell observation), a 5μm-thick section was set using the

z-section method with auto-optimization spacing to capture rhizoids'

protrusion.

For Arabidopsis root imaging, 4-day-old seedlings of each indicated genotype

were used for fluorescence imaging. After treatment in the liquid MS medium

supplied with indicated chemicals, seedlings were carefully mounted on a slice

with growth medium and then placed into a chambered coverslip (Lab-Tek) for

imaging. For the root hair imaging, a 3μm Z-projection image with 1μm step

was taken around the medium plan of the root hair. All fluorescence imaging

was performed using a laser scanning confocal microscopy (Zeiss LSM800,

20x air lens). The default setting for GFP detection was applied with 488nm

excitation and 495-545nm emission.

Images quantification

All images were analyzed by Fiji (ImageJ, https://imagej.net/software/fiji/). For the moss PpPINA-GFP polarization patterns at the tip of filaments, a line with 5px thickness was plotted along the PM from the blue to the green point, as shown in Fig. 1. The representative images for the DMSO control and drug treatments, obtained from the same imaging settings, were used to draw the line, and the mean intensity along the line was shown. For the moss phenotype analysis, mosses bearing inducible AtPIN1-GFP and AtPIN2-GFP were cultivated in the imaging dish for 5 days, followed by 1μM β-estradiol induction for another 3 days. For WT, pro*PINA*::PpPINA-GFP, and pro*ACT*::PpPINB-GFP, tissues were cultivated for 8 days and cell outlines were stained with FM4-64 for 10-30 mins. The line drawing function in Fiji was used to measure the length of subapical cell, and the line was depicted between the middle points of the two cell division planes. The angle measurement function was applied to examine the angle between the first cell division plane and horizontal cell outline for the division angle measurement.

For the Marchantia particle analysis, an individual cell was first selected by the polygon selection tool and was further analyzed by the ComDet plugin (https://imagej.net/plugins/spots-colocalization-comdet). We used 3 images for each transgenic line and selected 20 cells for each image. We separately detect particle sizes >3 pixels and >6 pixels for each cell. Other settings are default.

Genetic verification

The genomic DNA of transformants was isolated by the cetyltrimethylammonium bromide (CTAB) gDNA extraction (Schlink and Reski, 2002). In brief, moss tissues were harvested from one full plate and were

ground in liquid nitrogen. The ground tissues were then mixed and incubated

with a CTAB buffer, followed by the addition of chloroform. After centrifugation,

the supernatant was collected and precipitated with isopropanol at -20°C for

one hour. The PCR of the genomic DNA was performed with specific primers

listed in Supplemental Table S2.

Pharmacological treatments

To depolymerize microtubules and actin filaments, we used oryzalin (Sigma)

and latrunculin B (Sigma) to treat plants. The brefeldin A and cycloheximide

(Sigma) were used in the indicated conditions (Kleine-Vehn et al., 2008b). The

concentration and duration of treatment for different plants have been given in

numbers, and the conditions we used have been shown to efficiently

depolymerize cytoskeleton networks in different species (Baskin et al., 1994;

Baluska et al., 2001; Oda et al., 2009; Vidali et al., 2009; Glanc et al., 2019).

For Physcomitrella, the chemicals were diluted in a liquid BCD medium and

applied to the imaging dishes before imaging.

For Marchantia, G2 gemmae from transgenic plants were transferred into each

well of a 24-well plate containing 500μl liquid 1/2 B5 medium and cultured in

the growth chamber for 16 hours. The chemicals were diluted directly into the

medium prior to imaging at the time indicated in the text. Based on previous

studies, the concentration of Ory (Era et al., 2013) and LatB (Otani et al., 2018)

were selected.

For Arabidopsis, 4-days-old seedlings were submerged into the liquid MS

medium supplied with chemicals and then transferred to another agar medium

for imaging. 0.1% DMSO (Duchefa; 10 mM dimethylsulfoxide) was used as

control for all treatments.

Moss colony growth assay

For WT, overexpression of PpPINA- and PpPINB-GFP, the diameter of 20-day-old colonies cultivated on the same BCD medium agar plate was measured. For the inducible AtPIN1- and AtPIN2-GFP lines, 10-day-old colonies were transferred to another plate either with or without 1µM β-estradiol (Sigma) for another 10 days. The diameter of each colony was then measured. AtPIN1- and AtPIN2-GFP protonema phenotypes were imaged using the Nikon Optiphot-2 Trinocular Transmitted light microscope after 6 days of induction.

Marchantia thallus growth assay

G2 gemmae were transferred onto 1/2 B5 agar plates and grew for 10 days. Gemmae were imaged under a dissecting microscope (SZN71, LiWeng, Taiwan) with a CCD camera (Polychrome M, LiWeng, Taiwan). For measuring the vertical growth angle, an agar cube with an individual plant was cut out from the plate and placed in the middle of a slide. The slide was put on the surface of a laminar flow at a fixed distance to the edge, and images were taken by an HTC U11 cell phone camera. The growth angle was further measured by using ImageJ (https://imagej.net/software/fiji/).

FRAP analysis

For the FRAP analysis, images were captured at the medium plane and the target regions were selected to focus on the PM signal and avoid intracellular puncta. The intensity measurement was carried out with the Fiji program. The mean intensity of the bleached area in the before image was measured and

normalized to 100%. The intensity of the same area at the time of serious

imaging after bleaching was measured and normalized according to the before

image to calculate the recovery rate.

Author contributions and Acknowledgments

H. T., K. J. L., and J. F. designed experiments. H. T. performed and analyzed

all the Physcomitrella and Arabidopsis related experiments with the help of Y.

L. C. and S. L. T.. K. J. L. performed and analyzed all the Marchantia related

experiments. H. T., K. J. L, and J. F. wrote the manuscript.

The authors would like to thank Dr A. Johnson and Ms Alexandra Mally for

their careful proofreading and Dr Y.Z. Zhang for kindly sharing At-, Pp-, and

MpPIN constructs. We also thank Dr T. Kohchi for sharing the pMpGWB102

vector (Addgene #68556; http://n2t.net/addgene:68556;

RRID:Addgene_68556).

References

Adamowski, M., and Friml, J. (2015). PIN-dependent auxin transport: action,
regulation, and evolution. Plant Cell 27, 20-32.
Baluska, F., Jasik, J., Edelmann, H.G., Salajova, T., and Volkmann, D. (2001).
Latrunculin B-induced plant dwarfism: Plant cell elongation is
F-actin-dependent. Dev Biol 231, 113-124.
Baskin, T.I., Wilson, J.E., Cork, A., and Williamson, R.E. (1994). Morphology
and microtubule organization in Arabidopsis roots exposed to oryzalin
or taxol. Plant Cell Physiol 35, 935-942.
Bennett, T.A., Liu, M.M., Aoyama, T., Bierfreund, N.M., Braun, M., Coudert, Y.,
Dennis, R.J., O'Connor, D., Wang, X.Y., White, C.D., Decker, E.L.,
Reski, R., and Harrison, C.J. (2014). Plasma membrane-targeted PIN
proteins drive shoot development in a moss. Curr Biol 24, 2776-2785.

Blazquez, M.A., Nelson, D.C., and Weijers, D. (2020). Evolution of Plant
Hormone Response Pathways. Annu Rev Plant Biol 71, 327-353.

Dhonukshe, P., Aniento, F., Hwang, I., Robinson, D.G., Mravec, J., Stierhof,
Y.D., and Friml, J. (2007). Clathrin-mediated constitutive endocytosis of
PIN auxin efflux carriers in Arabidopsis. Curr Biol 17, 520-527.

Era, A., Kutsuna, N., Higaki, T., Hasezawa, S., Nakano, A., and Ueda, T.
(2013). Microtubule stability affects the unique motility of F-actin in
Marchantia polymorpha. J Plant Res 126, 113-119.

Floriach-Clark, J., Tang, H., and Willemsen, V. (2021). Mosses: Accessible
Systems for Plant Development Studies. In Model Organisms in Plant
Genetics (IntechOpen.

Friml, J. (2021). Fourteen Stations of Auxin. Cold Spring Harb Perspect Biol 14,
a039859.

Geldner, N., Friml, J., Stierhof, Y.D., Jurgens, G., and Palme, K. (2001). Auxin
transport inhibitors block PIN1 cycling and vesicle trafficking. Nature
413, 425-428.

Glanc, M., Fendrych, M., and Friml, J. (2018). Mechanistic framework for
cell-intrinsic re-establishment of PIN2 polarity after cell division. Nat
Plants 4, 1082-1088.

Glanc, M., Fendrych, M., and Friml, J. (2019). PIN2 Polarity Establishment in
Arabidopsis in the Absence of an Intact Cytoskeleton. Biomolecules 9,
222.

Ishizaki, K., Nishihama, R., Ueda, M., Inoue, K., Ishida, S., Nishimura, Y.,
Shikanai, T., and Kohchi, T. (2015). Development of Gateway Binary
Vector Series with Four Different Selection Markers for the Liverwort
Marchantia polymorpha. PLoS One 10, e0138876.

Kato, H., Nishihama, R., Weijers, D., and Kohchi, T. (2018). Evolution of
nuclear auxin signaling: lessons from genetic studies with basal land
plants. J Exp Bot 69, 291-301.

Kleine-Vehn, J., Langowski, L., Wisniewska, J., Dhonukshe, P., Brewer, P.B.,
and Friml, J. (2008a). Cellular and molecular requirements for polar PIN
targeting and transcytosis in plants. Mol Plant 1, 1056-1066.

Kleine-Vehn, J., Dhonukshe, P., Sauer, M., Brewer, P.B., Wisniewska, J.,
Paciorek, T., Benkova, E., and Friml, J. (2008b). ARF GEF-dependent
transcytosis and polar delivery of PIN auxin carriers in Arabidopsis. Curr
Biol 18, 526-531.

Kleine-Vehn, J., Wabnik, K., Martiniere, A., Langowski, L., Willig, K., Naramoto,
S., Leitner, J., Tanaka, H., Jakobs, S., Robert, S., Luschnig, C.,

Govaerts, W., Hell, S.W., Runions, J., and Friml, J. (2011). Recycling,
clustering, and endocytosis jointly maintain PIN auxin carrier polarity at
the plasma membrane. Mol Syst Biol 7, 540.

Krecek, P., Skupa, P., Libus, J., Naramoto, S., Tejos, R., Friml, J., and
Zazimalova, E. (2009). The PIN-FORMED (PIN) protein family of auxin
transporters. Genome Biol 10, 249.

Kubo, M., Imai, A., Nishiyama, T., Ishikawa, M., Sato, Y., Kurata, T., Hiwatashi,
Y., Reski, R., and Hasebe, M. (2013). System for stable
beta-estradiol-inducible gene expression in the moss Physcomitrella
patens. PLoS One 8, e77356.

Kubota, A., Ishizaki, K., Hosaka, M., and Kohchi, T. (2013). Efficient
Agrobacterium-mediated transformation of the liverwort Marchantia
polymorpha using regenerating thalli. Biosci Biotechnol Biochem 77,
167-172.

Lavy, M., Prigge, M.J., Tao, S., Shain, S., Kuo, A., Kirchsteiger, K., and Estelle,
M. (2016). Constitutive auxin response in Physcomitrella reveals
complex interactions between Aux/IAA and ARF proteins. Elife 5,
e13325.

Leyser, O. (2018). Auxin Signaling. Plant Physiol 176, 465-479.

Li, H., von Wangenheim, D., Zhang, X., Tan, S., Darwish-Miranda, N.,
Naramoto, S., Wabnik, K., De Rycke, R., Kaufmann, W.A., Gutl, D.,
Tejos, R., Grones, P., Ke, M., Chen, X., Dettmer, J., and Friml, J. (2021).
Cellular requirements for PIN polar cargo clustering in Arabidopsis
thaliana. New Phytol 229, 351-369.

Mockaitis, K., and Estelle, M. (2008). Auxin receptors and plant development:
a new signaling paradigm. Annu Rev Cell Dev Biol 24, 55-80.

Nakaoka, Y., Miki, T., Fujioka, R., Uehara, R., Tomioka, A., Obuse, C., Kubo,
M., Hiwatashi, Y., and Goshima, G. (2012). An inducible RNA
interference system in Physcomitrella patens reveals a dominant role of
augmin in phragmoplast microtubule generation. Plant Cell 24,
1478-1493.

Naramoto, S., Otegui, M.S., Kutsuna, N., de Rycke, R., Dainobu, T.,
Karampelias, M., Fujimoto, M., Feraru, E., Miki, D., Fukuda, H., Nakano,
A., and Friml, J. (2014). Insights into the localization and function of the
membrane trafficking regulator GNOM ARF-GEF at the Golgi apparatus
in Arabidopsis. Plant Cell 26, 3062-3076.

Narasimhan, M., Gallei, M., Tan, S., Johnson, A., Verstraeten, I., Li, L.,
Rodriguez, L., Han, H., Himschoot, E., Wang, R., Vanneste, S.,

Sanchez-Simarro, J., Aniento, F., Adamowski, M., and Friml, J. (2021).
Systematic analysis of specific and nonspecific auxin effects on
endocytosis and trafficking. Plant Physiol 186, 1122-1142.

Nishiyama, T., Hiwatashi, Y., Sakakibara, I., Kato, M., and Hasebe, M. (2000).
Tagged mutagenesis and gene-trap in the moss, Physcomitrella patens
by shuttle mutagenesis. DNA Res 7, 9-17.

Oda, Y., Hirata, A., Sano, T., Fujita, T., Hiwatashi, Y., Sato, Y., Kadota, A.,
Hasebe, M., and Hasezawa, S. (2009). Microtubules regulate dynamic
organization of vacuoles in Physcomitrella patens. Plant Cell Physiol 50,
855-868.

Otani, K., Ishizaki, K., Nishihama, R., Takatani, S., Kohchi, T., Takahashi, T.,
and Motose, H. (2018). An evolutionarily conserved NIMA-related
kinase directs rhizoid tip growth in the basal land plant Marchantia
polymorpha. Development 145, dev154617.

Rensing, S.A., Goffinet, B., Meyberg, R., Wu, S.Z., and Bezanilla, M. (2020).
The Moss Physcomitrium (Physcomitrella) patens: A Model Organism
for Non-Seed Plants. Plant Cell 32, 1361-1376.

Sauer, M., and Kleine-Vehn, J. (2019). PIN-FORMED and PIN-LIKES auxin
transport facilitators. Development 146, dev.168088.

Schlink, K., and Reski, R. (2002). Preparing high-quality DNA from moss
(Physcomitrella patens). Plant Molecular Biology Reporter 20, 423-423.

Shimamura, M. (2016). Marchantia polymorpha: Taxonomy, Phylogeny and
Morphology of a Model System. Plant Cell Physiol 57, 230-256.

Skokan, R., Medvecka, E., Viaene, T., Vosolsobe, S., Zwiewka, M., Muller, K.,
Skupa, P., Karady, M., Zhang, Y., Janacek, D.P., Hammes, U.Z., Ljung,
K., Nodzynski, T., Petrasek, J., and Friml, J. (2019). PIN-driven auxin
transport emerged early in streptophyte evolution. Nat Plants 5,
1114-1119.

Smit, M.E., and Weijers, D. (2015). The role of auxin signaling in early embryo
pattern formation. Curr Opin Plant Biol 28, 99-105.

Steinmann, T., Geldner, N., Grebe, M., Mangold, S., Jackson, C.L., Paris, S.,
Galweiler, L., Palme, K., and Jurgens, G. (1999). Coordinated polar
localization of auxin efflux carrier PIN1 by GNOM ARF GEF. Science
286, 316-318.

Tang, H., Duijts, K., Bezanilla, M., Scheres, B., Vermeer, J.E.M., and
Willemsen, V. (2020). Geometric cues forecast the switch from two- to
three-dimensional growth in Physcomitrella patens. New Phytol 225,
1945-1955.

Tao, S., and Estelle, M. (2018). Mutational studies of the Aux/IAA proteins in
Physcomitrella reveal novel insights into their function. New Phytol 218,
1534-1542.
Vanneste, S., and Friml, J. (2009). Auxin: a trigger for change in plant
development. Cell 136, 1005-1016.
Viaene, T., Landberg, K., Thelander, M., Medvecka, E., Pederson, E., Feraru,
E., Cooper, E.D., Karimi, M., Delwiche, C.F., Ljung, K., Geisler, M.,
Sundberg, E., and Friml, J. (2014). Directional auxin transport
mechanisms in early diverging land plants. Curr Biol 24, 2786-2791.
Vidali, L., Rounds, C.M., Hepler, P.K., and Bezanilla, M. (2009).
Lifeact-mEGFP reveals a dynamic apical F-actin network in tip growing
plant cells. PLoS One 4, e5744.
Vieten, A., Vanneste, S., Wisniewska, J., Benkova, E., Benjamins, R.,
Beeckman, T., Luschnig, C., and Friml, J. (2005). Functional
redundancy of PIN proteins is accompanied by auxin-dependent
cross-regulation of PIN expression. Development 132, 4521-4531.
Yamada, M., Miki, T., and Goshima, G. (2016). Imaging Mitosis in the Moss
Physcomitrella patens. Methods Mol Biol 1413, 263-282.
Yu, Z., Zhang, F., Friml, J., and Ding, Z. (2022). Auxin signaling: Research
advances over the past 30 years. J Integr Plant Biol 64, 371-392.
Zhang, Y., Xiao, G., Wang, X., Zhang, X., and Friml, J. (2019). Evolution of fast
root gravitropism in seed plants. Nat Commun 10, 3480.
Zourelidou, M., Absmanner, B., Weller, B., Barbosa, I.C., Willige, B.C., Fastner,
A., Streit, V., Port, S.A., Colcombet, J., de la Fuente van Bentem, S.,
Hirt, H., Kuster, B., Schulze, W.X., Hammes, U.Z., and Schwechheimer,
C. (2014). Auxin efflux by PIN-FORMED proteins is activated by two
different protein kinases, D6 PROTEIN KINASE and PINOID. Elife 3,
e02860.

Figure legends

Figure 1. The divergence of PpPINs- and AtPIN1-GFP localization and

trafficking in Physcomitrella.

A, PpPINA-GFP is polarized to the tip of protonema, and the disruption of actin

filaments by Latrunculin B (LatB) caused its hyperpolarization. The disruption

of microtubules by Oryzalin (Ory) disturbed the polarization of PpPINA-GFP.

The blue dotted curves indicate the selected areas for intensity plot shows in

D.

B, PpPINB-GFP is polarized at the dome area of the protonema, but none of

the treatments influenced this localization.

C, Enlarged pictures derived from A with the focus on protonema tips. White

arrowheads indicate the hyperpolarization or the absence of PpPINA-GFP.

D, Relative intensity of PpPINA-GFP along the cell membrane between blue

and green dots as indicated in A.

E, After the induction of expression, AtPIN1-GFP is localized on the PM with

cytosolic puncta. The disruption of cytoskeletons did not affect the localization

of AtPIN1-GFP.

All images show the z-projection of 10 sections with a 1µm step. FM4-64 was

used as a PM marker. Treatments were conducted as indicated. All conditions

were verified in at least 2 independent lines and 3 independent experiments.

Scale bar = 10 µm.

Figure 2. PpPINs- and AtPIN1-GFP show different responses to BFA in

Physcomitrella**.**

A, Representative images of PpPINA-, PpPINB-, and AtPIN1-GFP localization

at the tips of protonema cells. The BFA treatment did not affect PpPINA- and

PpPINB-GFP. The white arrowheads indicate the accumulation of

AtPIN1-GFP in cytosolic puncta after the BFA treatment. All images show the

z-projection of 10 sections with a 1µm step.

B and C, Example images of FRAP analyses of PpPINA- and AtPIN1-GFP.

The white arrowheads depict the FRAP target regions. Single image was

captured with the medium focus on cell apices. Imaging details are descried in

Methods.

D, Quantification of the FRAP analysis. The recovery of AtPIN1-GFP signals

was reduced by BFA treatment, while PpPINA-GFP showed no recovery.

N=11 for each condition; each data point was collected from 3 independent

experiments. Scale bar = 10 µm, ***P<0.001; Student's *t*-test.

Figure 3. Overexpression of PpPIN- and AtPIN-GFPs inhibits the

chloronema-caulonema transition in Physcomitrella.

A, Representative protonema cells of the indicated genetic background. The

cell outline was stained with FM4-64. White arrowheads indicate the first cell

division plane, and the yellow double arrow indicates the cell length of the

subapical cell. Scale bar = 100 µm.

B and C, Quantitative results of the subapical cell length and the division angle

(Ө) of indicated lines, respectively.

D, The colony size of indicated lines. For inducible AtPIN-GFP lines, the

colony size under induction (β-estr) or mock (DMSO) condition was measured.

Four colonies were measured for each indicated line. ***P<0.001; Student's

t-test.

E, Growth phenotypes caused by the overexpression of AtPIN1- or

AtPIN2-GFP. White arrowheads indicate the multiple branches and wavy

filaments.

Figure 4. MpPINZ-GFP is only polarized at the tip of young rhizoids in

Marchantia.

A – C, The localization of MpPINZ-GFP in gemmae, emerging rhizoids, and

young rhizoids.

A, A middle section image as illustrated showing the even distribution of

MpPINZ-GFP on the PM of gemmae.

B, Representative image with a 5 µm thick Z-section at the gemma surface as

illustrated. Blue arrows indicate the accumulation of MpPINZ-GFP at tips of

emerging rhizoids in rhizoid precursor cells.

C, MpPINZ-GFP is polarized at the tip of young rhizoids.

D, The responses of MpPINZ-GFP to indicated treatments. MpPINZ-GFP is

localized on the PM with cytosolic puncta signals in gemmae.

De-polymerization of cytoskeleton networks caused more and larger cytosolic

puncta; BFA treatment resulted in the increase of puncta only with larger size.

MpPINZ-GFP is polarized at the tip of young rhizoids, and its polarization was

only abolished by Oryzalin treatment (white arrowhead).

E and F, Quantification of intracellular puncta larger than 3 pixels and 6 pixels.

20 cells were quantified in each image, and three images were used to

quantify for each treatment (N=60 in total). $*P<0.05$, $**P<0.01$, $***P<0.001$;

Student's *t*-test.

Figure 5. Arabidopsis AtPIN1-GFP is not polarized in Marchantia.

A – C, The localization of AtPIN1-GFP in Marchantia gemmae, emerging

rhizoids, and young rhizoids. Imaging sections were obtained as illustrated in A

and B. The AtPIN1-GFP is localized on the PM and in the cytoplasm with no

apparent puncta signals in all examined tissues.

Figure 6. Overexpression of MpPINZ- and AtPIN-GFPs causes vertical growth

of Marchantia thalli.

A, Marchantia Tak-1 (WT) expands its thallus horizontally on the agar surface.

The overexpression of MpPINZ-, AtPIN1-, and AtPIN2-GFP showed vertical

thallus growth, which generated a large angle between the lower surface of the

thallus and the surface of the agar (θ).

B, The quantification of the thallus growth angle (θ) showed in A. N= 16 – 18

thali for each line. ***P<0.001; Student's *t*-test.

Figure 7. Bryophytic PINs and AtPIN2-GFP utilize different trafficking

mechanisms in Arabidopsis epidermal cells.

All PINs were translationally fused with GFP and expressed by the *AtPIN2*

promoter. De-polymerization of cytoskeleton networks caused mild

internalization of AtPIN2- and PpPINA-GFP, while PpPINB- and MpPINZ-GFP

showed strong responses with larger and more intracellular compartments.

BFA bodies were observed in all PIN-GFP lines with BFA treatment, while only

MpPINZ-GFP showed no BFA bodies in CHX+BFA treatment. Imaging details

are described in Methods.

Figure 8. Bryophytic PINs and AtPIN2-GFP are only polarized to the tip of

initial but not elongated root hairs.

All PIN-GFP lines show polarized signals at the tip of the initial root hair and

none of the indicated treatments disrupted their polarization. Note that the

polarization of the PIN-GFP signals were not observed in elongated root hairs.

All imaging details are described in Methods. The pixel values ranging from 0

to 255 are represented by the rainbow color.

Figure 1. The divergence of PpPINs- and AtPIN1-GFP localization and trafficking in Physcomitrella.

Figure 1. The divergence of PpPINs- and AtPIN1-GFP localization and trafficking in Physcomitrella.

A, PpPINA-GFP is polarized to the tip of protonema, and the disruption of actin filaments by Latrunculin B (LatB) caused its hyperpolarization. The disruption of microtubules by Oryzalin (Ory) disturbed the polarization of PpPINA-GFP. The blue dotted curves indicate the selected areas for intensity plot shows in D.

B, PpPINB-GFP is polarized at the dome area of the protonema, but none of the treatments influenced this localization.

C, Enlarged pictures derived from A with the focus on protonema tips. White arrowheads indicate the hyperpolarization or the absence of PpPINA-GFP.

D, Relative intensity of PpPINA-GFP along the cell membrane between blue and green dots as indicated in A.

E, After the induction of expression, AtPIN1-GFP is localized on the PM with cytosolic puncta. The disruption of cytoskeletons did not affect the localization of AtPIN1-GFP.

All images show the z-projection of 10 sections with a 1μm step. FM4-64 was used as a PM marker. Treatments were conducted as indicated. All conditions were verified in at least 2 independent lines and 3 independent experiments. Scale bar = 10 μm.

Figure 2. PpPINs- and AtPIN1-GFP show different responses to BFA in Physcomitrella.

Figure 2. PpPINs- and AtPIN1-GFP show different responses to BFA in Physcomitrella.

A, Representative images of PpPINA-, PpPINB-, and AtPIN1-GFP localization at the tips of protonema cells. The BFA treatment did not affect PpPINA- and PpPINB-GFP. The white arrowheads indicate the accumulation of AtPIN1-GFP in cytosolic puncta after the BFA treatment. All images show the z-projection of 10 sections with a 1μm step.

B and C, Example images of FRAP analyses of PpPINA- and AtPIN1-GFP. The white arrowheads depict the FRAP target regions. Single image was captured with the medium focus on cell apices. Imaging details are descried in Methods.

D, Quantification of the FRAP analysis. The recovery of AtPIN1-GFP signals was reduced by BFA treatment, while PpPINA-GFP showed no recovery. N=11 for each condition; each data point was collected from 3 independent experiments. Scale bar = 10 μm, ***P<0.001; Student's *t*-test.

Figure 3. Overexpression of PpPIN- and AtPIN-GFPs inhibits the chloronema-caulonema transition in Physcomitrella.

Figure 3. Overexpression of PpPIN- and AtPIN-GFPs inhibits the chloronema-caulonema transition in Physcomitrella.

A, Representative protonema cells of the indicated genetic background. The cell outline was stained with FM4-64. White arrowheads indicate the first cell division plane, and the yellow double arrow indicates the cell length of the subapical cell. Scale bar = 100 μm.

B and C, Quantitative results of the subapical cell length and the division angle (Θ) of indicated lines, respectively.

D, The colony size of indicated lines. For inducible AtPIN-GFP lines, the colony size under induction (β-estr) or mock (DMSO) condition was measured. Four colonies were measured for each indicated line. ***P<0.001; Student's *t*-test.

E, Growth phenotypes caused by the overexpression of AtPIN1- or AtPIN2-GFP. White arrowheads indicate the multiple branches and wavy filaments.

Figure 4. MpPINZ-GFP is only polarized at the tip of young rhizoids in Marchantia.

Figure 4. MpPINZ-GFP is only polarized at the tip of young rhizoids in Marchantia.

A – C, The localization of MpPINZ-GFP in gemmae, emerging rhizoids, and young rhizoids.

A, A middle section image as illustrated showing the even distribution of MpPINZ-GFP on the PM of gemmae.

B, Representative image with a 5 µm thick Z-section at the gemma surface as illustrated. Blue arrows indicate the accumulation of MpPINZ-GFP at tips of emerging rhizoids in rhizoid precursor cells.

C, MpPINZ-GFP is polarized at the tip of young rhizoids.

D, The responses of MpPINZ-GFP to indicated treatments. MpPINZ-GFP is localized on the PM with cytosolic puncta signals in gemmae. De-polymerization of cytoskeleton networks caused more and larger cytosolic puncta; BFA treatment resulted in the increase of puncta only with larger size. MpPINZ-GFP is polarized at the tip of young rhizoids, and its polarization was only abolished by Oryzalin treatment (white arrowhead).

E and F, Quantification of intracellular puncta larger than 3 pixels and 6 pixels. 20 cells were quantified in each image, and three images were used to quantify for each treatment (N=60 in total). *P<0.05, **P<0.01, ***P<0.001; Student's *t*-test.

Figure 5. Arabidopsis AtPIN1-GFP is not polarized in Marchantia.

Figure 5. Arabidopsis AtPIN1-GFP is not polarized in Marchantia.

A – C, The localization of AtPIN1-GFP in Marchantia gemmae, emerging rhizoids, and young rhizoids. Imaging sections were obtained as illustrated in A and B. The AtPIN1-GFP is localized on the PM and in the cytoplasm with no apparent puncta signals in all examined tissues.

Figure 6. Overexpression of MpPINZ- and AtPIN-GFPs causes vertical growth of Marchantia thalli.

Figure 6. Overexpression of MpPINZ- and AtPIN-GFPs causes vertical growth of Marchantia thalli.

A, Marchantia Tak-1 (WT) expands its thallus horizontally on the agar surface. The overexpression of MpPINZ-, AtPIN1-, and AtPIN2-GFP showed vertical thallus growth, which generated a large angle between the lower surface of the thallus and the surface of the agar (Ө).
B, The quantification of the thallus growth angle (Ө) showed in A. N= 16 – 18 thali for each line. ***P<0.001; Student's *t*-test.

Figure 7. Bryophytic PINs and AtPIN2-GFP utilize different trafficking mechanisms in Arabidopsis epidermal cells.

Figure 7. Bryophytic PINs and AtPIN2-GFP utilize different trafficking mechanisms in Arabidopsis epidermal cells.

All PINs were translationally fused with GFP and expressed by the *AtPIN2* promoter. De-polymerization of cytoskeleton networks caused mild internalization of AtPIN2- and PpPINA-GFP, while PpPINB- and MpPINZ-GFP showed strong responses with larger and more intracellular compartments. BFA bodies were observed in all PIN-GFP lines with BFA treatment, while only MpPINZ-GFP showed no BFA bodies in CHX+BFA treatment. Imaging details are described in Methods.

Fig. 8 Bryophytic PINs and AtPIN2-GFP are only polarized to the tip of initial but not elongated root hairs.

Figure 8. Bryophytic PINs and AtPIN2-GFP are only polarized to the tip of initial but not elongated root hairs.

All PIN-GFP lines show polarized signals at the tip of the initial root hair and none of the indicated treatments disrupted their polarization. Note that the polarization of the PIN-GFP signals were not observed in elongated root hairs. All imaging details are described in Methods. The pixel values ranging from 0 to 255 are represented by the rainbow color.

Parsed Citations

Adamowski, M., and Friml, J. (2015). PIN-dependent auxin transport: action, regulation, and evolution. Plant Cell 27, 20-32.
Google Scholar: Author Only Title Only Author and Title

Baluska, F., Jasik, J., Edelmann, H.G., Salajova, T., and Volkmann, D. (2001). Latrunculin B-induced plant dwarfism: Plant cell elongation is F-actin-dependent. Dev Biol 231, 113-124.
Google Scholar: Author Only Title Only Author and Title

Baskin, T.I., Wilson, J.E., Cork, A., and Williamson, R.E. (1994). Morphology and microtubule organization in Arabidopsis roots exposed to oryzalin or taxol. Plant Cell Physiol 35, 935-942.
Google Scholar: Author Only Title Only Author and Title

Bennett, T.A, Liu, M.M., Aoyama, T., Bierfreund, N.M., Braun, M., Coudert, Y., Dennis, R.J., O'Connor, D., Wang, X.Y., White, C.D., Decker, E.L., Reski, R., and Harrison, C.J. (2014). Plasma membrane-targeted PIN proteins drive shoot development in a moss. Curr Biol 24, 2776-2785.
Google Scholar: Author Only Title Only Author and Title

Blazquez, M.A, Nelson, D.C., and Weijers, D. (2020). Evolution of Plant Hormone Response Pathways. Annu Rev Plant Biol 71, 327-353.
Google Scholar: Author Only Title Only Author and Title

Dhonukshe, P., Aniento, F., Hwang, I., Robinson, D.G., Mravec, J., Stierhof, Y.D., and Friml, J. (2007). Clathrin-mediated constitutive endocytosis of PIN auxin efflux carriers in Arabidopsis. Curr Biol 17, 520-527.
Google Scholar: Author Only Title Only Author and Title

Era, A, Kutsuna, N., Higaki, T., Hasezawa, S., Nakano, A, and Ueda, T. (2013). Microtubule stability affects the unique motility of F-actin in Marchantia polymorpha. J Plant Res 126, 113-119.
Google Scholar: Author Only Title Only Author and Title

Floriach-Clark, J., Tang, H., and Willemsen, V. (2021). Mosses: Accessible Systems for Plant Development Studies. In Model Organisms in Plant Genetics (IntechOpen.
Google Scholar: Author Only Title Only Author and Title

Friml, J. (2021). Fourteen Stations of Auxin. Cold Spring Harb Perspect Biol 14, a039859.
Google Scholar: Author Only Title Only Author and Title

Geldner, N., Friml, J., Stierhof, Y.D., Jurgens, G., and Palme, K. (2001). Auxin transport inhibitors block PIN1 cycling and vesicle trafficking. Nature 413, 425-428.
Google Scholar: Author Only Title Only Author and Title

Glanc, M., Fendrych, M., and Friml, J. (2018). Mechanistic framework for cell-intrinsic re-establishment of PIN2 polarity after cell division. Nat Plants 4, 1082-1088.
Google Scholar: Author Only Title Only Author and Title

Glanc, M., Fendrych, M., and Friml, J. (2019). PIN2 Polarity Establishment in Arabidopsis in the Absence of an Intact Cytoskeleton. Biomolecules 9, 222.
Google Scholar: Author Only Title Only Author and Title

Ishizaki, K., Nishihama, R., Ueda, M., Inoue, K., Ishida, S., Nishimura, Y., Shikanai, T., and Kohchi, T. (2015). Development of Gateway Binary Vector Series with Four Different Selection Markers for the Liverwort Marchantia polymorpha. PLoS One 10, e0138876.
Google Scholar: Author Only Title Only Author and Title

Kato, H., Nishihama, R., Weijers, D., and Kohchi, T. (2018). Evolution of nuclear auxin signaling: lessons from genetic studies with basal land plants. J Exp Bot 69, 291-301.
Google Scholar: Author Only Title Only Author and Title

Kleine-Vehn, J., Langowski, L., Wisniewska, J., Dhonukshe, P., Brewer, P.B., and Friml, J. (2008a). Cellular and molecular requirements for polar PIN targeting and transcytosis in plants. Mol Plant 1, 1056-1066.
Google Scholar: Author Only Title Only Author and Title

Kleine-Vehn, J., Dhonukshe, P., Sauer, M., Brewer, P.B., Wisniewska, J., Paciorek, T., Benkova, E., and Friml, J. (2008b). ARF GEF-dependent transcytosis and polar delivery of PIN auxin carriers in Arabidopsis. Curr Biol 18, 526-531.
Google Scholar: Author Only Title Only Author and Title

Kleine-Vehn, J., Wabnik, K., Martiniere, A, Langowski, L., Willig, K., Naramoto, S., Leitner, J., Tanaka, H., Jakobs, S., Robert, S., Luschnig, C., Govaerts, W., Hell, S.W., Runions, J., and Friml, J. (2011). Recycling, clustering, and endocytosis jointly maintain PIN auxin carrier polarity at the plasma membrane. Mol Syst Biol 7, 540.
Google Scholar: Author Only Title Only Author and Title

Krecek, P., Skupa, P., Libus, J., Naramoto, S., Tejos, R., Friml, J., and Zazimalova, E. (2009). The PIN-FORMED (PIN) protein family of auxin transporters. Genome Biol 10, 249.
  Google Scholar: Author Only Title Only Author and Title

Kubo, M., Imai, A., Nishiyama, T., Ishikawa, M., Sato, Y., Kurata, T., Hiwatashi, Y., Reski, R., and Hasebe, M. (2013). System for stable beta-estradiol-inducible gene expression in the moss Physcomitrella patens. PLoS One 8, e77356.
  Google Scholar: Author Only Title Only Author and Title

Kubota, A., Ishizaki, K., Hosaka, M., and Kohchi, T. (2013). Efficient Agrobacterium-mediated transformation of the liverwort Marchantia polymorpha using regenerating thalli. Biosci Biotechnol Biochem 77, 167-172.
  Google Scholar: Author Only Title Only Author and Title

Lavy, M., Prigge, M.J., Tao, S., Shain, S., Kuo, A., Kirchsteiger, K., and Estelle, M. (2016). Constitutive auxin response in Physcomitrella reveals complex interactions between Aux/IAA and ARF proteins. Elife 5, e13325.
  Google Scholar: Author Only Title Only Author and Title

Leyser, O. (2018). Auxin Signaling. Plant Physiol 176, 465-479.
  Google Scholar: Author Only Title Only Author and Title

Li, H., von Wangenheim, D., Zhang, X., Tan, S., Darwish-Miranda, N., Naramoto, S., Wabnik, K., De Rycke, R., Kaufmann, W.A, Gutl, D., Tejos, R., Grones, P., Ke, M., Chen, X., Dettmer, J., and Friml, J. (2021). Cellular requirements for PIN polar cargo clustering in Arabidopsis thaliana. New Phytol 229, 351-369.
  Google Scholar: Author Only Title Only Author and Title

Mockaitis, K., and Estelle, M. (2008). Auxin receptors and plant development: a new signaling paradigm. Annu Rev Cell Dev Biol 24, 55-80.
  Google Scholar: Author Only Title Only Author and Title

Nakaoka, Y., Miki, T., Fujioka, R., Uehara, R., Tomioka, A, Obuse, C., Kubo, M., Hiwatashi, Y., and Goshima, G. (2012). An inducible RNA interference system in Physcomitrella patens reveals a dominant role of augmin in phragmoplast microtubule generation. Plant Cell 24, 1478-1493.
  Google Scholar: Author Only Title Only Author and Title

Naramoto, S., Otegui, M.S., Kutsuna, N., de Rycke, R., Dainobu, T., Karampelias, M., Fujimoto, M., Feraru, E., Miki, D., Fukuda, H., Nakano, A., and Friml, J. (2014). Insights into the localization and function of the membrane trafficking regulator GNOM ARF-GEF at the Golgi apparatus in Arabidopsis. Plant Cell 26, 3062-3076.
  Google Scholar: Author Only Title Only Author and Title

Narasimhan, M., Gallei, M., Tan, S., Johnson, A, Verstraeten, I., Li, L., Rodriguez, L., Han, H., Himschoot, E., Wang, R., Vanneste, S., Sanchez-Simarro, J., Aniento, F., Adamowski, M., and Friml, J. (2021). Systematic analysis of specific and nonspecific auxin effects on endocytosis and trafficking. Plant Physiol 186, 1122-1142.
  Google Scholar: Author Only Title Only Author and Title

Nishiyama, T., Hiwatashi, Y., Sakakibara, I., Kato, M., and Hasebe, M. (2000). Tagged mutagenesis and gene-trap in the moss, Physcomitrella patens by shuttle mutagenesis. DNA Res 7, 9-17.
  Google Scholar: Author Only Title Only Author and Title

Oda, Y., Hirata, A, Sano, T., Fujita, T., Hiwatashi, Y., Sato, Y., Kadota, A, Hasebe, M., and Hasezawa, S. (2009). Microtubules regulate dynamic organization of vacuoles in Physcomitrella patens. Plant Cell Physiol 50, 855-868.
  Google Scholar: Author Only Title Only Author and Title

Otani, K., Ishizaki, K., Nishihama, R., Takatani, S., Kohchi, T., Takahashi, T., and Motose, H. (2018). An evolutionarily conserved NIMA-related kinase directs rhizoid tip growth in the basal land plant Marchantia polymorpha. Development 145, dev154617.
  Google Scholar: Author Only Title Only Author and Title

Rensing, S.A, Goffinet, B., Meyberg, R., Wu, S.Z., and Bezanilla, M. (2020). The Moss Physcomitrium (Physcomitrella) patens: A Model Organism for Non-Seed Plants. Plant Cell 32, 1361-1376.
  Google Scholar: Author Only Title Only Author and Title

Sauer, M., and Kleine-Vehn, J. (2019). PIN-FORMED and PIN-LIKES auxin transport facilitators. Development 146, dev.168088.
  Google Scholar: Author Only Title Only Author and Title

Schlink, K., and Reski, R. (2002). Preparing high-quality DNA from moss (Physcomitrella patens). Plant Molecular Biology Reporter 20, 423-423.
  Google Scholar: Author Only Title Only Author and Title

Shimamura, M. (2016). Marchantia polymorpha: Taxonomy, Phylogeny and Morphology of a Model System. Plant Cell Physiol 57, 230-256.
  Google Scholar: Author Only Title Only Author and Title

Skokan, R., Medvecka, E., Viaene, T., Vosolsobe, S., Zwiewka, M., Muller, K., Skupa, P., Karady, M., Zhang, Y., Janacek, D.P., Hammes, U.Z., Ljung, K., Nodzynski, T., Petrasek, J., and Friml, J. (2019). PIN-driven auxin transport emerged early in streptophyte evolution. Nat Plants 5, 1114-1119.
Google Scholar: Author Only Title Only Author and Title

Smit, M.E., and Weijers, D. (2015). The role of auxin signaling in early embryo pattern formation. Curr Opin Plant Biol 28, 99-105.
Google Scholar: Author Only Title Only Author and Title

Steinmann, T., Geldner, N., Grebe, M., Mangold, S., Jackson, C.L., Paris, S., Galweiler, L., Palme, K., and Jurgens, G. (1999). Coordinated polar localization of auxin efflux carrier PIN1 by GNOM ARF GEF. Science 286, 316-318.
Google Scholar: Author Only Title Only Author and Title

Tang, H., Duijts, K., Bezanilla, M., Scheres, B., Vermeer, J.E.M., and Willemsen, V. (2020). Geometric cues forecast the switch from two- to three-dimensional growth in Physcomitrella patens. New Phytol 225, 1945-1955.
Google Scholar: Author Only Title Only Author and Title

Tao, S., and Estelle, M. (2018). Mutational studies of the Aux/IAA proteins in Physcomitrella reveal novel insights into their function. New Phytol 218, 1534-1542.
Google Scholar: Author Only Title Only Author and Title

Vanneste, S., and Friml, J. (2009). Auxin: a trigger for change in plant development. Cell 136, 1005-1016.
Google Scholar: Author Only Title Only Author and Title

Viaene, T., Landberg, K., Thelander, M., Medvecka, E., Pederson, E., Feraru, E., Cooper, E.D., Karimi, M., Delwiche, C.F., Ljung, K., Geisler, M., Sundberg, E., and Friml, J. (2014). Directional auxin transport mechanisms in early diverging land plants. Curr Biol 24, 2786-2791.
Google Scholar: Author Only Title Only Author and Title

Vidali, L., Rounds, C.M., Hepler, P.K., and Bezanilla, M. (2009). Lifeact-mEGFP reveals a dynamic apical F-actin network in tip growing plant cells. PLoS One 4, e5744.
Google Scholar: Author Only Title Only Author and Title

Vieten, A., Vanneste, S., Wisniewska, J., Benkova, E., Benjamins, R., Beeckman, T., Luschnig, C., and Friml, J. (2005). Functional redundancy of PIN proteins is accompanied by auxin-dependent cross-regulation of PIN expression. Development 132, 4521-4531.
Google Scholar: Author Only Title Only Author and Title

Yamada, M., Miki, T., and Goshima, G. (2016). Imaging Mitosis in the Moss Physcomitrella patens. Methods Mol Biol 1413, 263-282.
Google Scholar: Author Only Title Only Author and Title

Yu, Z., Zhang, F., Friml, J., and Ding, Z. (2022). Auxin signaling: Research advances over the past 30 years. J Integr Plant Biol 64, 371-392.
Google Scholar: Author Only Title Only Author and Title

Zhang, Y., Xiao, G., Wang, X., Zhang, X., and Friml, J. (2019). Evolution of fast root gravitropism in seed plants. Nat Commun 10, 3480.
Google Scholar: Author Only Title Only Author and Title

Zourelidou, M., Absmanner, B., Weller, B., Barbosa, I.C., Willige, B.C., Fastner, A., Streit, V., Port, S.A., Colcombet, J., de la Fuente van Bentem, S., Hirt, H., Kuster, B., Schulze, W.X., Hammes, U.Z., and Schwechheimer, C. (2014). Auxin efflux by PIN-FORMED proteins is activated by two different protein kinases, D6 PROTEIN KINASE and PINOID. Elife 3, e02860.
Google Scholar: Author Only Title Only Author and Title