Divergence of trafficking and polarization mechanisms for PIN auxin
transporters during land plant evolution

Han Tang[1,5], Kuan-Ju Lu[2,5], YuZhou Zhang[3], You-Liang Cheng[4], Shih-Long Tu[4], and
Jiří Friml[1]*

1. Institute of Science and Technology Austria (ISTA), Am Campus 1, 3400
Klosterneuburg, Austria
2. Graduate Institute of Biotechnology, National Chung Hsing University, No. 145,
Xingda Rd., South Dist., Taichung 40227, Taiwan, R.O.C.
3. College of Life Sciences, Northwest A&F University, Shaanxi, Yangling, China.
4. Institute of Plant and Microbial Biology, Academia Sinica, 128 Sec. 2, Academia Rd,
Nankang, Taipei 11529, Taiwan, R.O.C.

5. These authors contribute equally to this work.
* Correspondence: jiri.friml@ist.ac.at

23 **Running title: Evolution of PIN trafficking and polarization**

Abstract

The phytohormone auxin and its directional transport through tissues play a fundamental role in development of higher plants. This polar auxin transport predominantly relies on PIN-FORMED (PIN) auxin exporters. Hence, PIN polarization is crucial for development, but its evolution during the rise of morphological complexity in land plants remains unclear. Here, we performed a cross-species investigation by observing the trafficking and localization of endogenous and xenologous PINs in two bryophytes, *Physcomitrium patens* and *Marchantia polymorpha*, and in the flowering plant *Arabidopsis thaliana.* We confirmed that the GFP fusion did not compromise the auxin export function of all examined PINs by using radioactive auxin export assay and by observing the phenotypic changes in transgenic bryophytes. Endogenous PINs polarize to filamentous apices, while xenologous Arabidopsis PINs distribute symmetrically on the membrane in both bryophytes. In Arabidopsis root epidermis, bryophytic PINs show no defined polarity. Pharmacological interference revealed a strong cytoskeleton dependence of bryophytic but not Arabidopsis PIN polarization. The divergence of PIN polarization and trafficking is also observed within the bryophyte clade and between tissues in individual species. These results collectively reveal a divergence of PIN trafficking and polarity mechanisms throughout land plant evolution and a co-evolution of PIN sequence-based and cell-based polarity mechanisms.

Key words: Arabidopsis, evolution, Marchantia, Physcomitrium, PINs, polarization, protein trafficking

Introduction

Auxin is a crucial regulator of polarity and morphogenesis in land plants (Kato et al., 2018; Leyser, 2018; Mockaitis and Estelle, 2008; Smit and Weijers, 2015; Yu et al., 2022). The auxin gradients and local maxima within tissues coordinate a broad spectrum of plant development, ranging from embryogenesis to organ formation and tropisms (Friml, 2021; Vanneste and Friml, 2009). The establishment of the auxin gradient relies dominantly on directional auxin transport driven by the efflux carriers, PIN-FORMEDs (PINs). In different tissues, specific PINs are polarized at different plasma membrane (PM) domains, thus directly driving the directionality of auxin flow (Adamowski and Friml, 2015). Therefore, given the essential impact of auxin flow in various developmental processes, the function and polarization of PIN proteins are crucial for maintaining the correct pattern of plant growth and patterning (Sauer and Kleine-Vehn, 2019).

PINs are found in all land plants and can be traced back to charophytic green algae (Skokan et al., 2019; Viaene et al., 2013). The functional conservation of PINs in the auxin transport has also been demonstrated by exogenously expressing PINs from the green algae *Klebsormidium flaccidum* (*K. flaccidum*), the moss *Physcomitrium patens* (*P. patens*), and the angiosperm *Arabidopsis thaliana* in transgenic plants and in heterologous systems (Skokan et al., 2019; Zourelidou et al., 2014). Additionally, when xenologous PINs from charophytes or Arabidopsis are overexpressed in *P. patens*, the transgenic plants show growth inhibition that resembles auxin deprivation phenotypes (Lavy et al., 2016; Skokan et al., 2019; Tao and Estelle, 2018; Viaene et al., 2014). Therefore, these observations support the idea that PIN-mediated polar auxin transport may govern plant development since the emergence of land plants.

PIN polarity regulations have been extensively investigated in the angiosperm Arabidopsis. Canonical PINs, featuring a long central hydrophilic loop (HL) between two transmembrane domains, are delivered via the ER-Golgi vesicle trafficking pathway to the plasma membrane (PM). De-polymerization of actin filaments induces the accumulation of PIN-labelled small intracellular puncta near the PM, but with no apparent PIN polarity defect (Geldner et al., 2001; Glanc et al., 2018; 2019). Microtubules are involved in the cytokinetic trafficking of PINs, but are not required for the polarity establishment or maintenance at the PM of non-dividing cells (Geldner et al., 2001; Glanc et al., 2019; Kleine-Vehn et al., 2008b). Notably, the disruption of both

cytoskeletal networks delays but does not abolish AtPIN2 polarization in Arabidopsis

epidermal cells (Glanc et al., 2019). This suggests that the cytoskeletal networks are

involved, but not strictly essential for PIN polar trafficking, while other mechanisms

contribute to the PIN polar localization. PINs are known to undergo constitutive cycles

of endocytosis and recycling, which is modulated by auxin itself (Narasimhan et al.,

2021; Narasimhan et al., 2020); this process is essential for their polar distribution

(Doyle et al., 2015; Kleine-Vehn et al., 2011).

The Phosphorylation of specific sites within the HL region has been

demonstrated to be a critical determinant for PINs' apical-basal polarization in

Arabidopsis epidermis. A serine/threonine kinase, PINOID (PID), phosphorylates

specific sites on AtPIN2 and leads to its apical localization (Friml et al., 2004). In

contrast, when phosphatase 2A (PP2A) de-phosphorylates AtPIN2, which counteracts

the PID-dependent phosphorylation, and thus guides the delivery of AtPIN2 to the

basal domain of epidermal cells (Michniewicz et al., 2007). The phosphorylation sites

targeted by different kinase families are crucial for polar localizations and functions of

PINs, and most sites are highly conserved within Arabidopsis canonical PINs

(Zwiewka et al., 2019). Since PINs are present in all land plants, one can hypothesize

that the phosphorylation-based polarity regulation may have been established since

the emergence of early land plants. However, it has never been clarified whether these

phosphorylation sites are evolutionary conserved in early land plants.

The PIN polarization has been observed in the early-divergent moss, *P. patens*,

which grows as filamentous protonemata, and its endogenous PpPINA-GFP shows a

polar localization at the tip of protonema cells (Bennett et al., 2014; Viaene et al., 2014).

However, the polar localization of PpPINA-GFP is not robust in other species. When

PpPINA-GFP is expressed in Arabidopsis root epidermal cells, where AtPIN2-GFP

presents a clear apical localization, PpPINA-GFP is mislocalized at both basal and

apical sites (Zhang et al., 2019). Furthermore, PINs from the liverwort, *Marchantia*

polymorpha (*M. polymorpha*), and from the green algae, *K. flaccidum*, are also

mislocalized in root epidermal cells of Arabidopsis (Zhang et al., 2019). These

distinctive PIN localizations in different species suggest that mechanisms for PINs

trafficking and polarization may have diversified after the emergence of land plants.

Despite the profound significance of PIN polarization and the resulting directional

auxin transport for land plant development, PIN trafficking and polarization

mechanisms are mainly derived from the angiosperm model *Arabidopsis thaliana*.

Here we investigated PIN trafficking/polarization mechanisms from an
evolutionary perspective. We show that canonical PINs from bryophytes *P. patens*, *M.*
polymorpha, and Arabidopsis present high conservation in their transmembrane
domains and phosphorylation sites. Endogenous PIN-GFPs show different localization
patterns in various developmental contexts, suggesting tissue-specific PIN
polarization mechanisms. Cross-species investigation revealed that xenologous PINs
can traffic to the PM but fail to enrich at the polar domains revealing species-specific
mechanisms for PIN polarization. This notion was further verified by different
dependency of cytoskeletons for the polarization of Arabidopsis PINs and bryophytic
PINs. Overall, our results highlight that PINs trafficking and polarization mechanisms
underwent complex evolution during the gradual rise of morphological complexity in
land plants.
145

Results
Phosphorylation sites are highly conserved between bryophytic and
Arabidopsis PINs
To dissect the conservation between bryophytic and Arabidopsis PINs, we first
performed a phylogenetic analysis. The coding sequence of the single canonical PIN,
MpPINZ, from *M. polymorpha*, three canonical PINs, PpPINA-PpPINC, from *P. patens*,
and five canonical PINs, AtPIN1-AtPIN4 and AtPIN7, from Arabidopsis were subjected
to MEGA X program (Kumar et al., 2018) for the phylogenetic study. Bryophytic PINs
are categorized in one group, and the MpPINZ is closer to Arabidopsis PINs than
PpPINs are (Figure 1A). Arabidopsis AtPIN1 and AtPIN2 were identified with clear
polarity at the PM that plays crucial role in embryogenesis, organ formation, and tropic
growth (Krecek et al., 2009; Omelyanchuk et al., 2016). The phylogenetic tree showed
that both AtPIN1 and AtPIN2 are equally close to bryophytic PINs, while AtPIN2 was
extensively investigated for its polarization in roots (Abas et al., 2006; Glanc et al.,
2018; Kleine-Vehn et al., 2008a), thus we first picked AtPIN2 as our reference for
further alignment analyses. The identity index of coding amino acid sequences
showed that full-length AtPIN2 has around 50% identity with each bryophytic PIN
(Figure 1B). Since the transmembrane domains show a high similarity between all
examined PINs, we suspected that central HLs are more divergent. Surprisingly, the
identity index for only the HL region of AtPIN2 presented over 40% identity with

MpPINZ (45%), PpPINA (42%), and PpPINB (43%) (Figure 1B). The identity index of

full-length AtPIN1 and HLs showed similar results as AtPIN2 (Supplemental Figure 1).

168   The overview of coding sequences for all examined PINs revealed highly

conserved transmembrane domains at N-terminus and C-terminus, connected by a

less conserved HL region (Figure 1C). Because the polarization of AtPIN2 is tightly

associated with its phosphorylation status on the HL, we used AtPIN2 as a reference

to search for and highlight these experimental-identified phosphorylation sites

(Barbosa et al., 2018; Sukumar et al., 2009; Zhang et al., 2010). Comparing to

transmembrane domains, despite of a relatively lower conservation in their HLs, the

four identified phosphorylation sites are fully conserved between Arabidopsis PINs and

bryophytic PINs (Figure 1D), which suggests that the PIN phosphorylation might be

evolutionarily conserved for the regulation of their intracellular localization.

The hydrophilic loops in AtPIN2, PpPINA, and MpPINZ are less conserved

To dissect how conserved between Arabidopsis PINs and bryophytic PINs is in their

protein structures, we next performed structural prediction and alignment. Structures

predicted by the Alphafold2 program resemble newly resolved structures of AtPIN1

and AtPIN8 (Supplemental Figure 2) (Jumper et al., 2021; Ung et al., 2022; Yang et

al., 2022), supporting that the structure prediction is reliable. Therefore, to compare

the structures of well-investigated polarized AtPINs, we predicted the structures of

AtPIN1 and AtPIN2 by Alphafold2 and aligned their predicted structures in ChimeraX

(Jumper et al., 2021; Pettersen et al., 2021). The structural conservation was

presented in almost perfect alignment of transmembrane domains while their HL

regions showed similar patterns (Figure 2A, close arrowheads) with two extra random

loops in AtPIN2 (Figure 2A, open arrowheads). We next aligned the structure of AtPIN2

with PpPINA and MpPINZ. The transmembrane domains are highly conserved with

nearly perfect alignments, but the HLs are less conserved that only one loop showed

partial similarity (Figure 2B, close arrowheads). Note that to reveal the conservation

and phosphorylation sites clearly, the orientation of protein structures presented in

Figure 2A and 2B-E were rotated in different angles. In general, bryophytic PINs

possess loosen and larger loops compared to AtPIN2 (Figure 2B-2E). The predicted

structures were shown individually with the annotated phosphorylation sites that are

indicated in Figure 1D (Figure 2C-2E). The structural conservations in transmembrane

domains imply that ancestry bryophytic PINs may traffic to the PM through the same

delivery pathways as Arabidopsis PINs, whereas the loosen loops with conserved
phosphorylation sites suggest a deviation from conserved regulation for their
polarization.
203

GFP-fused PIN proteins possess auxin export activity

The sequence and structural analyses between Arabidopsis PINs and bryophytic PINs
revealed high conservation in sequence, phosphorylation sites, and structures of
transmembrane domains. Thus, we next would like to investigate whether bryophytic
PINs are delivered to the PM with polar domain enrichment as Arabidopsis PINs. To
achieve this, we first generated PINs with GFP-fused into an equivalent position of
AtPIN1, AtPIN2, PpPINA, and MpPINZ as shown in Figure 3A (Zhang et al., 2019). To
verify the auxin export function of these GFP-fused PINs, we subcloned each GFP-
fused *PIN* gene into a moss vector, where each *PIN-GFP* gene was driven by an
inducible *XVE* promoter (Kubota et al., 2013). We generated transgenic moss plants
carrying different *XVE::PIN-GFPs* separately, and we used these transgenic lines to
perform the auxin export assay. In brief, the overexpression of *PIN-GFPs* were
induced for 3 days, followed by a radioactive auxin H^3-IAA treatment for 24 hours. The
radioactive tissues were then washed twice and were incubated in fresh growth
mediums for another 24 hrs. The culture medium was collected for H^3 scintillation
detection (Lewis and Muday, 2009). The wild-type moss plants were taken along as
an internal control to show the basal exportation of H^3-IAA by endogenous PpPINs. In
comparison to the wild-type, all examined PIN-GFP plants showed higher amount of
radioactive auxin in the culture medium, indicating their auxin export activities (Figure
3B).
Additionally, the function of PIN-GFPs was also confirmed in the growth
changes caused by *PIN-GFPs* overexpression in both *P. patens*. and *M. polymorpha*.
During early development, *P. patens* gradually transits its filamentous protonemata
from thicker/shorter chloronema cells with perpendicular division planes to
thinner/longer caulonema cells with oblique division planes (Rensing et al., 2020). In
the previous study, PpPINA, PpPINB, and PpPIND overexpression has been
demonstrated to inhibit the protonema transition (Viaene et al., 2014). Here we
showed that overexpression of *PpPINA-, AtPIN1-*, and *AtPIN2-GFP* led to similar
defects in this chloronema-caulonema transition with a shorter cell length of the
subapical cell and a smaller division angle (Figure 3C-3E). The wild-type *M.*

polymorpha has prostrate thalli; however, with the overexpression of either *MpPINZ-*
GFP or *AtPIN-GFPs*, the thallus grew more vertically, as apparent from the side view
(Figure 3F), phenocopying the auxin deficient phenotype (Kato et al., 2017). The
angles between the thallus and the horizontal agar were measured to quantify the
vertical growth (Figure 3G). The overexpression of *MpPINZ-GFP* caused the most
striking phenotype, but the overexpression of *AtPIN-GFPs* also resulted in a vertical
growth that showed significant differences from the wild-type. Our results show that all
PIN-GFPs can export auxin in the moss system and cause phenotypic changes in both
P. patens and *M. polymorpha* upon overexpression.
243
Endogenous PINs present different localization patterns in different types of
tissue
To observe the PpPINA-GFP localization in *P. patens*, we used a stable moss
transgenic line carrying the native promoter controlled *PpPINA* genomic DNA-GFP
fusion (*pPINA::PpPINA-GFP*) where GFP was inserted in the same position as we
verified. The moss *P. patens* has a filamentous protonema stage and a leafy
gametophore stage in its life cycle. The initial protonema cell was newly regenerated
from a detached leaf, and the elongated protonema was imaged from a 6-days-old
moss colony. PpPINA-GFP localized at the PM of the protonema tip with a clear
polarity in both initial and elongated protonemata (Figure 4A). To examine whether the
polar localization of PpPINA-GFP appears in complex tissues composed of multiple
cell layers as well, we observed its localization in gametophytic leaves. In young
gametophytic leaves, PpPINA-GFP showed clear basal-apical polarization along the
leaf axis with notable corner enrichment (Figure 4A), whereas in mature leaves
PpPINA-GFP was evenly distributed on the PM (Supplemental Figure 3). We also
analyzed another endogenous *P. patens* PpPINB, fused with GFP and driven by its
endogenous promoter. Notably, PpPINB-GFP did not show visible polarity at the tip of
protonema, but more even PM distribution with an increased intracellular signal
(Supplemental Figure 4). The difference between the localization of PpPINA-GFP and
PpPINB-GFP suggests different trafficking and polarization pathways for different PINs
co-existing in the same cell already in bryophytes.
The divergence of PpPINA-GFP polarity in filamentous protonema cells and in
gametophytic leaves made us wondering if this tissue-specific polarization of PINs is
conserved in other bryophytes. To achieve this, we looked at another bryophyte, the

liverwort *M. polymorpha* that produces gemmae as the asexual reproductive progenies
consisted of multiple cell layers. After water imbibition, single-cellular rhizoids emerge
from the large rhizoid precursor cells on epidermis of a gemma (Shimamura, 2016).
To analyze the subcellular localization of the sole canonical PIN, MpPINZ, in *M.*
polymorpha, we generated a *p35S::MpPINZ-GFP* transgenic line. Interestingly,
MpPINZ-GFP localized on the PM with small intracellular puncta and with no apparent
polarity in all gemma epidermal cells (Figure 4B, right panel). However, when gemmae
were stimulated to grow rhizoids, the signal of MpPINZ-GFP accumulated at the
protrusion site of emerging rhizoids in the rhizoid precursor cells (Figure 4B, yellow
arrowheads). Later after their emergence, MpPINZ-GFP was polarized at the tip of
young rhizoids (Figure 4B), whereas the signal diminished when the rhizoids
elongated.
The localization of PpPINA- and MpPINZ-GFP in filamentous and complex
tissues suggests distinct polarity recognition mechanisms for bryophytic PINs in
different tissue types. To examine if Arabidopsis PINs show similar polar patterns as
bryophytic PINs in different types of tissues, we observed *pPIN2::AtPIN2-GFP* line in
filamentous root hairs and complex epidermal cells. As previously described, AtPIN2-
GFP showed apical localization in epidermal cells (Figure 4C) (Zhang et al., 2019).
Since the polarity of bryophytic PINs in their original species mainly appeared at the
tip of filamentous tissues, we focused on the localization of AtPIN2-GFP in initial and
elongated root hairs, respectively. AtPIN2-GFP revealed a polar localization at the tip
of initial root hairs (Figure 4C). However, the polarity of AtPIN2-GFP signals diminished
in the elongated hairs (Figure 4C). The different polarity patterns of PINs in different
types of tissue support the notion that regulations of PIN polarization are specialized
in different cellular profiles and in different developmental contexts.
293

Xenologous PINs are PM localized with no defined polarity
Since bryophytic and Arabidopsis PINs demonstrated tissue- and development-
specific polarity regulations, we wonder if the conserved phosphorylation sites in the
HL are sufficient to drive the polarization of xenologous PINs in other species. To
achieve this, we observed the same moss *XVE::PIN-GFPs* transgenic lines we
generated for the auxin export assay, carrying PpPINA-, AtPIN1-, and AtPIN2-GFP,
respectively. In protonemata, AtPIN1-GFP showed an even distribution on the PM and
cell division plane without apparent polarity at the tips, and displayed numerous small

intracellular puncta, whereas the PpPINA-GFP signal presented the same polarization
as driven by its endogenous promoter (Figure 4A, 5A, and 5D). AtPIN2-GFP under the
same induction condition showed much lower intensity but resembled AtPIN1-GFP
localization patterns, we thus took AtPIN1-GFP as representative images
(Supplemental Figure 5). A short-term induction of *XVE::AtPIN1-GFP* was performed,
and it showed no difference in the localization, which verified that the localization
patterns of AtPIN1-GFP were not caused by overexpression (Supplemental Video 1).
To investigate whether these features are conserved in bryophytes, we
introduced AtPIN1-GFP driven by the *35S* promoter into *M. polymorpha*. In gemma
epidermal cells, the AtPIN1-GFP showed a non-polar PM localization with high
cytosolic signals (Supplemental Figure 6A). In emerging rhizoids, AtPIN1-GFP did not
show visible polar localization in rhizoid precursor cells (Supplemental Figure 6B). In
young rhizoids, where MpPINZ-GFP showed clear tip polarization, AtPIN1-GFP was
evenly distributed on the PM with homogenized cytosolic signals (Figure 5B and 5D).
Since our results indicated that Arabidopsis AtPIN1-GFP in examined
bryophytes are delivered to the PM with no visible polarity, we next addressed whether
Arabidopsis trafficking machinery can polarize bryophytic PINs. To analyze bryophytic
PINs, we observed the protein localization at epidermal cells of transgenic Arabidopsis
lines expressing *pPIN2::AtPIN2-, PpPINA-, and MpPINZ-GFP* (Zhang et al., 2019). As
previously described, AtPIN2-GFP showed apical localization in the epidermal cells,
while PpPINA-GFP localized at both the apical and basal sides and MpPINZ-GFP
mainly localized at the basal side with some lateral residence. (Figure 5C and 5E)
(Zhang et al., 2019). These data thus support that the machinery driving PIN proteins
to the PM through a generally conserved cellular trafficking pathway, whereas the PIN
polarization is specialized in different species.
327
Cytoskeletal networks are important for the polarization of bryophytic PINs
The diversification of PIN polarities in different plant species and tissues suggests that
plant cells might utilize various machineries to deliver and maintain PIN proteins to the
target side on the PM. To verify this, we examined the necessity of the cytoskeletal
networks in the polarization of bryophytic PINs and AtPIN2 in their original species.
We depolymerized actin filaments or microtubules by treating plant tissues with
Latrunculin B (LatB) or Oryzalin (Ory), respectively. In *P. patens*, the abolishment of
actin filaments resulted in the hyper-polarization of PpPINA-GFP, where PpPINA-GFP

accumulated at a focal locus to the very tip of the cell (Figure 6A and 6C). The
disruption of microtubules resulted in less accumulation of PpPINA-GFP at the tip of
protonemata, and the PpPINA-GFP appeared to be detached from the PM (Figure 6A
and 6C). The changes of PpPINA-GFP localization in response to drug treatments
demonstrated the requirement of cytoskeletal networks for PpPINA-GFP polarization
in filamentous tissues.

In *M. polymorpha*, we used the same pharmacological interference to
investigate whether the MpPINZ-GFP polarization in young rhizoids relies on the
cytoskeletal networks. MpPINZ-GFP remains polarized at the tip when actin filaments
were disrupted, while the disruption of microtubules resulted in the dislocation of the
polarized MpPINZ-GFP (Figure 6B and 6C). These results collectively demonstrate
the diversification on cytoskeletal dependency for PINs polarization within the
bryophyte clade.

In Arabidopsis, the disruption of actin filaments by LatB resulted in an increasing
appearance of AtPIN2-GFP signals in the cell interior, while the Ory treatment caused
a minor effect on AtPIN2-GFP localization (Figure 6D). Upon cytoskeleton disruption,
the apical polarization of AtPIN2-GFP in epidermal cells was still evident; suggesting
that in contrast to the dependency of cytoskeletal networks for the bryophytic PINs,
the polarization of Arabidopsis PINs mainly relies on other trafficking or polarity
retention mechanisms. Again, these results reinforce the notion that mechanisms
underlying PIN polarization have been expanded and specifically diversified in
different species.

Discussion

Evolution of sequence-specific determinants of PIN polarity

In flowering plants, as exemplified by Arabidopsis, the five PM-localized canonical
PINs show different polarities in different developmental and tissue contexts, thus
mediating directional auxin fluxes and generating asymmetric auxin distribution for
plethora of developmental processes ultimately shaping the plant form. Originating
from a single PIN auxin transporter, such as found in simple filamentous streptophyte
algae (Skokan et al., 2019), PIN family radiated during evolution into PINs showing
different expression and localization patterns and mediating many developmental and

physiological functions.

Ectopic co-expression of PINs in the same cell type such as AtPIN1 and AtPIN2
in root epidermis can lead to diverse polarity patterns (AtPIN1 – basal; AtPIN2 – apical),
thus demonstrating simultaneous presence of multiple polarity mechanisms in those
cells (Wisniewska et al., 2006). In this study, we showed that in bryophytes, as
exemplified by the moss *P. patens,* PpPINA and PpPINB endogenously expressed in
protonemata filaments, present different localization patterns (Figure 4 and
Supplemental Figure 4). PpPINA shows tip-focused PM localization, whereas PpPINB
can be found more spread on the PM and also intracellularly. Whereas, the functional
importance of this difference remains unclear, it clearly shows that already in
bryophytes, parallel polarity/trafficking mechanisms exists in the same cells, to which
different PINs can be recruited, presumably based on specific sequence-based signals.

Notably, the sequence alignment between the HL regions of PpPINA and
PpPINB showed 90.31% identity, suggesting that these signals are presumably
encoded within the 10% difference regions. Further analysis of the differences
between PpPINA and PpPINB or AtPIN1 and AtPIN2 would help to identify the
sequence signals for PIN polarity regulations. Despite the identity of the sequence-
based signals remain unclear, our observations show that cellular polarity mechanisms
and PIN sequence-based polarity signals, which are crucial for diverse developmental
roles in flowering plants, started to diversify already in bryophytes.

Context-specific determinants of PIN polarity
It is well known from Arabidopsis that the same PINs show different localization
patterns in different contexts (Vieten et al., 2005). For example, AtPIN2 is apicalized
in epidermal cells, while it is localized on the basal side of young cortex cells (Kleine-
Vehn *et al.*, 2008a). The observations imply that different cell types possess specific
trafficking pathways for the same PIN protein. In line with this, our results show that in
filamentous cells, e.g., protonemata in *P. patens*, rhizoids in *M. polymorpha*, and root
hairs in Arabidopsis, endogenous PIN proteins are polarized at the tip of apical cells
(Figure 4A-4C). Notably, MpPINZ-GFP and AtPIN2-GFP signals were diminished
when the rhizoids or root hairs elongated, and the PpPINA-GFP was polarized only in
young gametophytic leaves but the pattern dispersed in mature leaves. These data
collectively demonstrate that PIN polarity is differentially regulated in different
developmental contexts.

In complex tissues with multiple cell layers, unlike in filamentous tissues, the
PM-localized MpPINZ-GFP did not show polarity in thalli, while PpPINA-GFP and
AtPIN2-GFP presented polar localization at the apical-basal domain of the cells
(Figure 4A-4C). This data shows that PIN polarity and trafficking mechanisms have
evolved with specific modifications in different tissues and cell types, in both
angiosperms and early-diverging land plants. This likely reflects different requirements
for directional auxin transport in different developmental contexts, and it implies a co-
evolution of PIN sequence-based signals and cell type-specific polar sorting and
trafficking mechanisms.
413

Diversification of PIN trafficking and polarity mechanisms during land plant
evolution
The core mechanisms for auxin biosynthesis, auxin signaling, and PIN-mediated auxin
transport are evolutionary conserved across land plants (Blazquez et al., 2020; Kato
et al., 2018; Sauer and Kleine-Vehn, 2019). However, bryophytes and vascular plants
diverged around 450 million years ago and emerged different types of tissues and
organs, it is not clear how conserved the PIN polarity regulation is under such drastic
changes during land plant evolution. Our cross-species studies revealed that
xenologous PINs, such as Arabidopsis PINs in bryophytes and bryophytic PINs in
Arabidopsis, are able to traffic to the PM (Figure 5), which suggests that all canonical
PINs can be recognized by the general protein transport machinery in other species.
Interestingly, when the Arabidopsis PINs are ectopically expressed in
bryophytes, they fail to form any specific polarity, whereas bryophytic PINs remain in
apical-basal domains in Arabidopsis epidermis (Figure 5). These observations may
hint an evolutionary loss of some regulatory motifs for PIN polarization that are
presence in bryophytic PINs but absence in Arabidopsis PINs. In line with this, the
overall coding sequences of bryophytic PINs are longer than Arabidopsis PINs, and
the extra sequences are positioned in their HLs, the main regulatory regions for PIN
polarization. Our study thus demonstrates that PINs polarity mechanisms are not
conserved throughout plant evolution.
This notion was also verified by the differences in cytoskeleton requirements
for PIN polarization between bryophytes and angiosperms. The polarity of bryophytic
PINs was disrupted when cytoskeletal networks are depolymerized, while Arabidopsis
AtPIN2 was not affected (Figure 6). This finding suggests a gradual shift in the

dependence of PIN polarity and trafficking from the cytoskeleton-dependent pathways
towards the cytoskeleton-independent pathways along the land plant evolution.

Overall, our results demonstrate that different plant species evolved specialized
pathways to deliver PINs and maintain their polarity at the PM. This is likely linked to
an increasing repertoire of auxin transport developmental roles adopted with
increasing morphological complexity during land plant evolution.

Methods

Plant growth and transformation
Arabidopsis seeds were surface sterilized and grown on 1/2 MS plates. After a two-
day stratification at 4°C, seedlings were grown under the long-day condition (16 hours
light, 8 hours dark) at 22°C with 100-120 µmol photons $m^{-2}s^{-1}$ white light. For *P. patens*,
all transgenic and WT plants were cultured on BCD medium plates in a growth
chamber at 24°C under the long-day condition (16 hours light, 8 hours dark) with
35µmol photons $m^{-2}s^{-1}$ white light. For *M. polymorpha*, WT and all transgenic plants
were cultured on 1/2 B5 plates in a growth chamber under the long-day condition (16
hours light, 8 hours dark) at 22°C with 50-60 µmol photons $m^{-2}s^{-1}$ white LED light.

All transgenic plants used in this study are listed in Supplemental Table 1. *P.*
patens transgenic plants *pPINA::PINA-GFP* has been generated and verified as
previously reported (Viaene et al., 2014), and the inducible overexpression
XVE::AtPIN1-GFP, *XVE::AtPIN2-GFP*, and *XVE::MpPINZ-GFP* lines were generated
as described in previous studies (Tang et al., 2020; Yamada et al., 2016). In brief, the
Gransden 2004 WT moss plants were freshly propagated and transformed via the
PEG-mediated transformation (Nishiyama et al., 2000). The transformants were
selected under Hygromycin (20µg/mL), and two independent transgenic lines were
selected for imaging and analysis.

p35S::MpPINZ-, AtPIN1-, and *AtPIN2-GFP M. polymorpha* plants were
generated via the Agrobacterium transformation method described before (Kubota et
al., 2013). In brief, the apical meristemic region of each two-week-old Takaragaike-1
(Tak-1) thallus was removed and further cut into four pieces. After culturing on 1/2 B5
with 1% sucrose agar plates for 3 days, the cut thalli were transferred to 50ml 0M51C
medium with 200 µM acetosyringone (4'-Hydroxy-3',5'-dimethoxyacetophenone) in

200ml flasks with 130 rpm agitation to coculture with $OD_{600} = 1$ density agrobacteria

harboring the target construct for another 3 days. The transformed thalli were washed

and plated on 1/2 B5 plates with proper antibiotic selection. Independent T1 lines were

isolated, and G1 lines from independent T1 lines were generated by sub-cultivating

single gemmalings, which emerged asexually from a single initial cell (Shimamura,

2016). The next generation of G1, called the G2 generation, was used for analyses.

Arabidopsis transgenic lines bearing bryophytic PIN-GFPs under *AtPIN2* promoter

control were generated and used as in a previous study (Zhang et al., 2019). For root

imaging, seeds were sown on 1/2 Murashige–Skoog (MS) medium plates and kept at

4°C for 2 days and moved to the growth chamber to culture vertically for another 4

days.

Plasmid construction

Plasmids and primers for construction and genotype confirmation are listed in

Supplemental Table 2. For transgenic lines with inducible overexpression by moss, the

insertion site of the *GFP* gene into the HL was indicated in Figure 3A, and the *PIN-*

GFP regions were amplified from previously generated plasmids (Zhang et al., 2019)

by PCR and cloned into the gateway entry plasmid pENTR/D-TOPO® as the

manufacturer suggested. The fragments were sub-cloned into the pPGX8 vector,

which contains a p*35S* driven β-estradiol inducible XVE cassette (Floriach-Clark et al.,

2021; Kubo et al., 2013; Nakaoka et al., 2012) via a Gateway® LR reaction (Invitrogen)

as the manufacture's recommendation.

To generate a *p35S::MpPINZ-, AtPIN1-*, and *AtPIN2-GFP* construct, the cDNA

of the target fragment containing proper *GFP* insertion were amplified with the primers

listed in Supplemental Table 2. The amplified fragments were cloned into the

pENTR/D-TOPO® vector (Invitrogen) with the protocol supplied by the manufacturer.

Plasmids with target genes were further sub-cloned into the pMpGWB102 vector

containing a *35S* promoter (Ishizaki et al., 2015) by the Gateway® LR reaction

(Invitrogen) as the manufacturer's recommendation.

Microscopy

For moss observation, protonemata were cultured in glass-bottom dishes covered by

BCD agar medium for 6 – 7d. The live-cell imaging was then performed on a Leica

SP8X-SMD confocal microscope equipped with a hybrid single-molecule detector

(HyD) and an ultrashort pulsed white-light laser (WLL; 50%; 1 ps at 40 MHz frequency).
The Leica Application Suite X was used as a software platform, and imaging was
conducted with a HC PL APO CS2 40x/1.20 water immersion objective. The following
settings were used: scan speed of 400 Hz, resolution of 1024 x 1024 pixels and
standard acquisition mode of the hybrid detector. The time-gating system was
activated to avoid the autofluorescence emitted from chloroplasts. For the filament
growth assay, the imaging dish was applied with FM4-64 (Invitrogen) solution for 10-
30 mins, and the 10x objective lens was used.

For Marchantia observation, gemmae were picked from a gemma cup and
transferred into a 24-well plate with 500 µl liquid 1/2 B5 medium. After culturing in the
growth chamber for 24 hours, gemmae from each sample were transferred and
observed on a slide under Leica Stellaris 8 system with HyD detectors and the
ultrashort pulsed white-light laser (WLL; 70%; 1 ps at 40 MHz frequency). The Leica
Application Suite X was used as a software platform, and imaging was conducted with
an HC PL APO CS2 40x/1.20 water immersion objective. The following settings were
used: scan speed of 400 Hz and resolution of 1024 x 1024 pixels. For GFP-containing
images, a 488nm white light laser was selected, and the detection range was set
between 500nm to 525nm. The tau-gating model, harvesting photons with 1.0-10.0ns
lifetime, was used for all Marchantia imaging to avoid autofluorescence emitted from
the chloroplasts. For the surface section (rhizoid precursor cell observation), a 5 µm-
thick section was set using the z-section method with auto-optimization spacing to
capture rhizoids' protrusion.

For Arabidopsis root imaging, 4-day-old seedlings of each indicated genotype
were used for fluorescence imaging. After treatment in the liquid MS medium supplied
with indicated chemicals, seedlings were carefully mounted on a slice with growth
medium and then placed into a chambered coverslip (Lab-Tek) for imaging. For the
root hair imaging, a 3 µm Z-projection image with 1 µm step was taken around the
medium plan of the root hair. All fluorescence imaging was performed using a laser
scanning confocal microscopy (Zeiss LSM800, 20x air lens). The default setting for
GFP detection was applied with 488nm excitation and 495-545nm emission.

Images quantification

All images were analyzed by Fiji (ImageJ, https://imagej.net/software/fiji/). For the

polarization patterns at the tip of filaments in *P. patens*, a line with 5 px thickness was

plotted along the PM as depicted in Figure 4 and Figure 5. The representative images

for the DMSO control and drug treatments, obtained from the same imaging settings,

were used to draw the line, and the mean intensity along the line was shown. For the

moss phenotype analysis, mosses bearing inducible *PIN-GFP* were cultivated in the

imaging dish for 5 days, followed by 1 µM β-estradiol induction for another 3 days. The

cell outlines were stained with FM4-64 for 10-30 mins. The line drawing function in Fiji

was used to measure the length of subapical cell, and the line was depicted between

the middle points of the two cell division planes. The angle measurement function was

applied to examine the angle between the first cell division plane and horizontal cell

outline for the division angle measurement.

Genomic DNA isolation

The genomic DNA of transformants was isolated by the cetyltrimethylammonium

bromide (CTAB) gDNA extraction (Schlink and Reski, 2002). In brief, moss tissues

were harvested from one full plate and were ground in liquid nitrogen. The ground

tissues were then mixed and were incubated with a CTAB buffer, followed by the

addition of chloroform. After centrifugation, the supernatant was collected and was

precipitated with isopropanol at -20°C for one hour.

Pharmacological treatments

To depolymerize microtubules and actin filaments, we used oryzalin (Sigma) and

latrunculin B (Sigma) to treat plants. The concentration and duration of treatment for

different plants have been given in numbers, and the conditions we used have been

shown to efficiently depolymerize cytoskeletal networks in different species (Baluska

et al., 2001; Baskin et al., 1994; Glanc et al., 2019; Oda et al., 2009; Vidali et al., 2009).

For *P. patens*, the chemical was diluted in a liquid BCD medium and was applied to

the imaging dishes before imaging.

For *M. polymorpha*, G2 gemmae from transgenic plants were transferred into

each well of a 24-well plate containing 500 µl liquid 1/2 B5 medium and cultured in the

growth chamber for 16 hours. The chemicals were diluted directly into the medium

prior to imaging at the time indicated in the text. Based on previous studies, the

concentration of Ory (Era et al., 2013) and LatB (Otani et al., 2018) were selected.

For Arabidopsis, 4-days-old seedlings were submerged into the liquid MS medium

supplied with chemicals and then transferred to another agar medium for imaging. 0.1% DMSO (Duchefa; 10 mM dimethylsulfoxide) was used as control for all treatments.

P. patens auxin export assay

The auxin export assay performed with transgenic moss plants was modified based on the protocol developed for Arabidopsis seedlings (Lewis and Muday, 2009). In brief, 7-days-old fresh tissues were transfer to liquid BCDAT growth medium with 1µM β-estradiol induction for another 4d with gentle shaking, followed by ^{3}H-IAA treatment with a final concentration of 10 nM for 24 hrs. The radioactive tissues were then washed twice with sterile H_2O, and were cultivated in fresh BCDAT medium for another 24 hrs. The cultivated medium was then collected to mix with ScintiVerse BD cocktail (Fisher, SX18-4) in 1:30 (v:v), and the export of auxin was measured by the Scintillation counter (Beckman, LS6500).

M. polymorpha thallus growth assay

G2 gemmae were transferred onto 1/2 B5 agar plates and grew for 10 days. Gemmae were imaged under a dissecting microscope (SZN71, LiWeng, Taiwan) with a CCD camera (Polychrome M, LiWeng, Taiwan). For measuring the vertical growth angle, an agar cube with an individual plant was cut out from the plate and placed in the middle of a slide. The slide was put on the surface of a laminar flow at a fixed distance to the edge, and images were taken by an HTC U11 cell phone camera. The growth angle was further measured by using ImageJ (https://imagej.net/software/fiji/).

Phylogenetic analysis

The phylogenetic analysis for full length amino acid sequences of all examined PINs was carried out in MEGA X program (Kumar et al., 2018) and the results were imported into iTOL (https://itol.embl.de/) for visual illustration. The evolutionary history was inferred by using the Maximum Likelihood method and JTT matrix-based model with default settings (Jones et al., 1992). The alignment and identity index were produced by online CLUSTAL alignment program with default settings.

Funding

This work was supported by the ERC grant (PR1023ERC02) to H. T. and J. F., and by

606   the ministry of science and technology (grant number 110-2636-B-005-001) to K. J. L.

Author contributions

608   H. T., K. J. L., and J. F. designed experiments. Y.Z. C. initiated the project and
609   provided key constructs and materials. H. T. performed and analyzed all the *P. patens*
610   and Arabidopsis related experiments with the help of Y. L. C. and S. L. T.. K. J. L.
611   performed and analyzed all the *M. polymorpha* related experiments. H. T., K. J. L, and
612   J. F. wrote the manuscript.

Acknowledgments

614   The authors would like to thank Dr. A. Johnson and Ms. Alexandra Mally for their
615   careful proofreading. We also thank Dr. T. Kohchi for sharing the pMpGWB102 vector
616   (Addgene #68556; http://n2t.net/addgene:68556; RRID:Addgene_68556).

References

Abas, L., Benjamins, R., Malenica, N., Paciorek, T., Wisniewska, J., Moulinier-Anzola, J.C.,
Sieberer, T., Friml, J., and Luschnig, C. (2006). Intracellular trafficking and proteolysis of the
Arabidopsis auxin-efflux facilitator PIN2 are involved in root gravitropism. Nat Cell Biol **8**:249-
256. 10.1038/ncb1369.
Adamowski, M., and Friml, J. (2015). PIN-dependent auxin transport: action, regulation, and
evolution. Plant Cell **27**:20-32. 10.1105/tpc.114.134874.
Baluska, F., Jasik, J., Edelmann, H.G., Salajova, T., and Volkmann, D. (2001). Latrunculin B-
induced plant dwarfism: Plant cell elongation is F-actin-dependent. Dev Biol **231**:113-124.
10.1006/dbio.2000.0115.
Barbosa, I.C.R., Hammes, U.Z., and Schwechheimer, C. (2018). Activation and Polarity
Control of PIN-FORMED Auxin Transporters by Phosphorylation. Trends Plant Sci **23**:523-538.
10.1016/j.tplants.2018.03.009.
Baskin, T.I., Wilson, J.E., Cork, A., and Williamson, R.E. (1994). Morphology and
microtubule organization in Arabidopsis roots exposed to oryzalin or taxol. Plant Cell Physiol
35:935-942.
Bennett, T.A., Liu, M.M., Aoyama, T., Bierfreund, N.M., Braun, M., Coudert, Y., Dennis, R.J.,
O'Connor, D., Wang, X.Y., White, C.D., et al. (2014). Plasma membrane-targeted PIN proteins
drive shoot development in a moss. Curr Biol **24**:2776-2785. 10.1016/j.cub.2014.09.054.
Blazquez, M.A., Nelson, D.C., and Weijers, D. (2020). Evolution of Plant Hormone Response
Pathways. Annu Rev Plant Biol **71**:327-353. 10.1146/annurev-arplant-050718-100309.
Doyle, S.M., Haeger, A., Vain, T., Rigal, A., Viotti, C., Langowska, M., Ma, Q., Friml, J.,
Raikhel, N.V., Hicks, G.R., et al. (2015). An early secretory pathway mediated by GNOM-LIKE
1 and GNOM is essential for basal polarity establishment in Arabidopsis thaliana. Proc Natl
Acad Sci U S A **112**:E806-815. 10.1073/pnas.1424856112.

Era, A., Kutsuna, N., Higaki, T., Hasezawa, S., Nakano, A., and Ueda, T. (2013). Microtubule
stability affects the unique motility of F-actin in Marchantia polymorpha. J Plant Res
126:113-119. 10.1007/s10265-012-0496-4.
Floriach-Clark, J., Tang, H., and Willemsen, V. (2021). Mosses: Accessible Systems for Plant
Development Studies. In Model Organisms in Plant Genetics, (IntechOpen.
Friml, J. (2021). Fourteen Stations of Auxin. Cold Spring Harb Perspect Biol **14**:a039859.
10.1101/cshperspect.a039859.
Friml, J., Yang, X., Michniewicz, M., Weijers, D., Quint, A., Tietz, O., Benjamins, R.,
Ouwerkerk, P.B., Ljung, K., Sandberg, G., et al. (2004). A PINOID-dependent binary switch in
apical-basal PIN polar targeting directs auxin efflux. Science **306**:862-865.
10.1126/science.1100618.
Geldner, N., Friml, J., Stierhof, Y.D., Jurgens, G., and Palme, K. (2001). Auxin transport
inhibitors block PIN1 cycling and vesicle trafficking. Nature **413**:425-428. 10.1038/35096571.
Glanc, M., Fendrych, M., and Friml, J. (2018). Mechanistic framework for cell-intrinsic re-
establishment of PIN2 polarity after cell division. Nat Plants **4**:1082-1088. 10.1038/s41477-
018-0318-3.
Glanc, M., Fendrych, M., and Friml, J. (2019). PIN2 Polarity Establishment in Arabidopsis in
the Absence of an Intact Cytoskeleton. Biomolecules **9**:222. 10.3390/biom9060222.
Ishizaki, K., Nishihama, R., Ueda, M., Inoue, K., Ishida, S., Nishimura, Y., Shikanai, T., and
Kohchi, T. (2015). Development of Gateway Binary Vector Series with Four Different
Selection Markers for the Liverwort Marchantia polymorpha. PLoS One **10**:e0138876.
10.1371/journal.pone.0138876.
Jones, D.T., Taylor, W.R., and Thornton, J.M. (1992). The rapid generation of mutation data
matrices from protein sequences. Comput Appl Biosci **8**:275-282.
10.1093/bioinformatics/8.3.275.
Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate
protein structure prediction with AlphaFold. Nature **596**:583-589. 10.1038/s41586-021-
03819-2.
Kato, H., Nishihama, R., Weijers, D., and Kohchi, T. (2018). Evolution of nuclear auxin
signaling: lessons from genetic studies with basal land plants. J Exp Bot **69**:291-301.
10.1093/jxb/erx267.
Kato, H., Kouno, M., Takeda, M., Suzuki, H., Ishizaki, K., Nishihama, R., and Kohchi, T.
(2017). The Roles of the Sole Activator-Type Auxin Response Factor in Pattern Formation of
Marchantia polymorpha. Plant Cell Physiol **58**:1642-1651. 10.1093/pcp/pcx095.
Kleine-Vehn, J., Langowski, L., Wisniewska, J., Dhonukshe, P., Brewer, P.B., and Friml, J.
(2008a). Cellular and molecular requirements for polar PIN targeting and transcytosis in
plants. Mol Plant **1**:1056-1066. 10.1093/mp/ssn062.
Kleine-Vehn, J., Dhonukshe, P., Sauer, M., Brewer, P.B., Wisniewska, J., Paciorek, T.,
Benkova, E., and Friml, J. (2008b). ARF GEF-dependent transcytosis and polar delivery of PIN
auxin carriers in Arabidopsis. Curr Biol **18**:526-531. 10.1016/j.cub.2008.03.021.
Kleine-Vehn, J., Wabnik, K., Martiniere, A., Langowski, L., Willig, K., Naramoto, S., Leitner,
J., Tanaka, H., Jakobs, S., Robert, S., et al. (2011). Recycling, clustering, and endocytosis
jointly maintain PIN auxin carrier polarity at the plasma membrane. Mol Syst Biol **7**:540.
10.1038/msb.2011.72.
Krecek, P., Skupa, P., Libus, J., Naramoto, S., Tejos, R., Friml, J., and Zazimalova, E. (2009).
The PIN-FORMED (PIN) protein family of auxin transporters. Genome Biol **10**:249.

10.1186/gb-2009-10-12-249.

Kubo, M., Imai, A., Nishiyama, T., Ishikawa, M., Sato, Y., Kurata, T., Hiwatashi, Y., Reski, R.,
and Hasebe, M. (2013). System for stable beta-estradiol-inducible gene expression in the
moss Physcomitrella patens. PLoS One **8**:e77356. 10.1371/journal.pone.0077356.

Kubota, A., Ishizaki, K., Hosaka, M., and Kohchi, T. (2013). Efficient Agrobacterium-
mediated transformation of the liverwort Marchantia polymorpha using regenerating thalli.
Biosci Biotechnol Biochem **77**:167-172. 10.1271/bbb.120700.

Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol **35**:1547-1549.
10.1093/molbev/msy096.

Lavy, M., Prigge, M.J., Tao, S., Shain, S., Kuo, A., Kirchsteiger, K., and Estelle, M. (2016).
Constitutive auxin response in Physcomitrella reveals complex interactions between Aux/IAA
and ARF proteins. Elife **5**:e13325. 10.7554/eLife.13325.

Lewis, D.R., and Muday, G.K. (2009). Measurement of auxin transport in Arabidopsis
thaliana. Nat Protoc **4**:437-451. 10.1038/nprot.2009.1.

Leyser, O. (2018). Auxin Signaling. Plant Physiol **176**:465-479. 10.1104/pp.17.00765.

Michniewicz, M., Zago, M.K., Abas, L., Weijers, D., Schweighofer, A., Meskiene, I., Heisler,
M.G., Ohno, C., Zhang, J., Huang, F., et al. (2007). Antagonistic regulation of PIN
phosphorylation by PP2A and PINOID directs auxin flux. Cell **130**:1044-1056.
10.1016/j.cell.2007.07.033.

Mockaitis, K., and Estelle, M. (2008). Auxin receptors and plant development: a new
signaling paradigm. Annu Rev Cell Dev Biol **24**:55-80.
10.1146/annurev.cellbio.23.090506.123214.

Nakaoka, Y., Miki, T., Fujioka, R., Uehara, R., Tomioka, A., Obuse, C., Kubo, M., Hiwatashi,
Y., and Goshima, G. (2012). An inducible RNA interference system in Physcomitrella patens
reveals a dominant role of augmin in phragmoplast microtubule generation. Plant Cell
24:1478-1493. 10.1105/tpc.112.098509.

Narasimhan, M., Johnson, A., Prizak, R., Kaufmann, W.A., Tan, S., Casillas-Perez, B., and
Friml, J. (2020). Evolutionarily unique mechanistic framework of clathrin-mediated
endocytosis in plants. Elife **9**10.7554/eLife.52067.

Narasimhan, M., Gallei, M., Tan, S., Johnson, A., Verstraeten, I., Li, L., Rodriguez, L., Han,
H., Himschoot, E., Wang, R., et al. (2021). Systematic analysis of specific and nonspecific
auxin effects on endocytosis and trafficking. Plant Physiol **186**:1122-1142.
10.1093/plphys/kiab134.

Nishiyama, T., Hiwatashi, Y., Sakakibara, I., Kato, M., and Hasebe, M. (2000). Tagged
mutagenesis and gene-trap in the moss, Physcomitrella patens by shuttle mutagenesis. DNA
Res **7**:9-17. 10.1093/dnares/7.1.9.

Oda, Y., Hirata, A., Sano, T., Fujita, T., Hiwatashi, Y., Sato, Y., Kadota, A., Hasebe, M., and
Hasezawa, S. (2009). Microtubules regulate dynamic organization of vacuoles in
Physcomitrella patens. Plant Cell Physiol **50**:855-868. 10.1093/pcp/pcp031.

Omelyanchuk, N.A., Kovrizhnykh, V.V., Oshchepkova, E.A., Pasternak, T., Palme, K., and
Mironova, V.V. (2016). A detailed expression map of the PIN1 auxin transporter in
Arabidopsis thaliana root. BMC Plant Biol **16 Suppl 1**:5. 10.1186/s12870-015-0685-0.

Otani, K., Ishizaki, K., Nishihama, R., Takatani, S., Kohchi, T., Takahashi, T., and Motose, H.
(2018). An evolutionarily conserved NIMA-related kinase directs rhizoid tip growth in the
basal land plant Marchantia polymorpha. Development **145**:dev154617.
10.1242/dev.154617.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H.,
and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. Protein Sci **30**:70-82. 10.1002/pro.3943.
Rensing, S.A., Goffinet, B., Meyberg, R., Wu, S.Z., and Bezanilla, M. (2020). The Moss
Physcomitrium (Physcomitrella) patens: A Model Organism for Non-Seed Plants. Plant Cell
32:1361-1376. 10.1105/tpc.19.00828.
Sauer, M., and Kleine-Vehn, J. (2019). PIN-FORMED and PIN-LIKES auxin transport
facilitators. Development **146**:dev.168088. 10.1242/dev.168088.
Schlink, K., and Reski, R. (2002). Preparing high-quality DNA from moss (Physcomitrella
patens). Plant Molecular Biology Reporter **20**:423-423.
Shimamura, M. (2016). Marchantia polymorpha: Taxonomy, Phylogeny and Morphology of a
Model System. Plant Cell Physiol **57**:230-256. 10.1093/pcp/pcv192.
Skokan, R., Medvecka, E., Viaene, T., Vosolsobe, S., Zwiewka, M., Muller, K., Skupa, P.,
Karady, M., Zhang, Y., Janacek, D.P., et al. (2019). PIN-driven auxin transport emerged early
in streptophyte evolution. Nat Plants **5**:1114-1119. 10.1038/s41477-019-0542-5.
Smit, M.E., and Weijers, D. (2015). The role of auxin signaling in early embryo pattern
formation. Curr Opin Plant Biol **28**:99-105. 10.1016/j.pbi.2015.10.001.
Sukumar, P., Edwards, K.S., Rahman, A., Delong, A., and Muday, G.K. (2009). PINOID kinase
regulates root gravitropism through modulation of PIN2-dependent basipetal auxin
transport in Arabidopsis. Plant Physiol **150**:722-735. 10.1104/pp.108.131607.
Tang, H., Duijts, K., Bezanilla, M., Scheres, B., Vermeer, J.E.M., and Willemsen, V. (2020).
Geometric cues forecast the switch from two- to three-dimensional growth in Physcomitrella
patens. New Phytol **225**:1945-1955. 10.1111/nph.16276.
Tao, S., and Estelle, M. (2018). Mutational studies of the Aux/IAA proteins in Physcomitrella
reveal novel insights into their function. New Phytol **218**:1534-1542. 10.1111/nph.15039.
Ung, K.L., Winkler, M., Schulz, L., Kolb, M., Janacek, D.P., Dedic, E., Stokes, D.L., Hammes,
U.Z., and Pedersen, B.P. (2022). Structures and mechanism of the plant PIN-FORMED auxin
transporter. Nature **609**:605-610. 10.1038/s41586-022-04883-y.
Vanneste, S., and Friml, J. (2009). Auxin: a trigger for change in plant development. Cell
136:1005-1016. 10.1016/j.cell.2009.03.001.
Viaene, T., Delwiche, C.F., Rensing, S.A., and Friml, J. (2013). Origin and evolution of PIN
auxin transporters in the green lineage. Trends Plant Sci **18**:5-10.
10.1016/j.tplants.2012.08.009.
Viaene, T., Landberg, K., Thelander, M., Medvecka, E., Pederson, E., Feraru, E., Cooper,
E.D., Karimi, M., Delwiche, C.F., Ljung, K., et al. (2014). Directional auxin transport
mechanisms in early diverging land plants. Curr Biol **24**:2786-2791.
10.1016/j.cub.2014.09.056.
Vidali, L., Rounds, C.M., Hepler, P.K., and Bezanilla, M. (2009). Lifeact-mEGFP reveals a
dynamic apical F-actin network in tip growing plant cells. PLoS One **4**:e5744.
10.1371/journal.pone.0005744.
Vieten, A., Vanneste, S., Wisniewska, J., Benkova, E., Benjamins, R., Beeckman, T.,
Luschnig, C., and Friml, J. (2005). Functional redundancy of PIN proteins is accompanied by
auxin-dependent cross-regulation of PIN expression. Development **132**:4521-4531.
10.1242/dev.02027.
Wisniewska, J., Xu, J., Seifertova, D., Brewer, P.B., Ruzicka, K., Blilou, I., Rouquie, D.,
Benkova, E., Scheres, B., and Friml, J. (2006). Polar PIN localization directs auxin flow in
plants. Science **312**:883. 10.1126/science.1121356.

Yamada, M., Miki, T., and Goshima, G. (2016). Imaging Mitosis in the Moss Physcomitrella
patens. Methods Mol Biol **1413**:263-282. 10.1007/978-1-4939-3542-0_17.
Yang, Z., Xia, J., Hong, J., Zhang, C., Wei, H., Ying, W., Sun, C., Sun, L., Mao, Y., Gao, Y., et al.
(2022). Structural insights into auxin recognition and efflux by Arabidopsis PIN1. Nature
609:611-615. 10.1038/s41586-022-05143-9.
Yu, Z., Zhang, F., Friml, J., and Ding, Z. (2022). Auxin signaling: Research advances over the
past 30 years. J Integr Plant Biol **64**:371-392. 10.1111/jipb.13225.
Zhang, J., Nodzynski, T., Pencik, A., Rolcik, J., and Friml, J. (2010). PIN phosphorylation is
sufficient to mediate PIN polarity and direct auxin transport. Proc Natl Acad Sci U S A
107:918-922. 10.1073/pnas.0909460107.
Zhang, Y., Xiao, G., Wang, X., Zhang, X., and Friml, J. (2019). Evolution of fast root
gravitropism in seed plants. Nat Commun **10**:3480. 10.1038/s41467-019-11471-8.
Zourelidou, M., Absmanner, B., Weller, B., Barbosa, I.C., Willige, B.C., Fastner, A., Streit, V.,
Port, S.A., Colcombet, J., de la Fuente van Bentem, S., et al. (2014). Auxin efflux by PIN-
FORMED proteins is activated by two different protein kinases, D6 PROTEIN KINASE and
PINOID. Elife **3**:e02860. 10.7554/eLife.02860.
Zwiewka, M., Bilanovicova, V., Seifu, Y.W., and Nodzynski, T. (2019). The Nuts and Bolts of
PIN Auxin Efflux Carriers. Front Plant Sci **10**:985. 10.3389/fpls.2019.00985.

Figure legends

Figure 1. Phosphorylation sites are highly conserved between bryophytic and Arabidopsis
PINs.

(A) Phylogenetic analysis of canonical PINs from early-divergent plants *P. patens (Pp), M.*
polymorpha (Mp), and representative angiosperm *A. thaliana (At)*. The unrooted tree
shows the relationships of different PINs in two representative bryophytes and
Arabidopsis. The scale bar represents the number of changes per site.

(B) Identity indexes of all PINs compared to AtPIN2 with full length or only the hydrophilic
loop region of coding amino acid sequences. Identity indexes of all PINs compared to
AtPIN1 is shown in Supplemental Figure 1.

(C) Alignment of PIN amino acid sequences. Identical amino acids are highlighted with
blue column. The rea and green boxes show the hydrophilic loop regions bearing
conserved phosphorylation sites, which are enlarged in (D).

(D) Four conserved phosphorylation sites, labeled by S1 to S4, were verified in previous
AtPIN2 studies and are depicted with black frames. Note that S1 to S4 are conserved
in every examined PINs.

Figure 2. The hydrophilic loops in AtPIN2, PpPINA, and MpPINZ are less conserved.

(A) Structural alignment of AtPIN1 and AtPIN2. The protein structures are predicted by

Alphafold2 and aligned in ChimeraX software. The structural conserved regions are

indicated by black arrowheads, and the non-conserved regions are indicated by empty

arrowheads.

(B) Structural alignment of AtPIN2 with PpPINA and MpPINZ. Only transmembrane

domains and one loop aligned with each other, while the majority of hydrophilic loop

regions are not conserved. The four conserved phosphorylation sites are labeled with

atom details in ball-and-stick style.

(C) – (E) Individual protein structure retrieved from (B). The four conserved

phosphorylation sites are indicated by arrowheads.

Figure 3. GFP-fused PIN proteins possess auxin export activity.

(A) The insertion site of GFP in indicated PIN proteins. Numbers represent the amino acid

position in respective proteins.

(B) The auxin export assay with *P. patens* transgenic plants. Fresh tissues were incubated

with radioactive H^3 IAA, followed by washing and the radiative H^3-IAA exported in the

new culture medium was measured with the scintillation detector after one day.

(C) Representative protonema cells of the indicated genetic background. The cell outline

was stained with FM4-64. White arrowheads indicate the first cell division plane, and

the yellow double arrow indicates the cell length of the subapical cell. Scale bar = 100

μm.

(D) and (E), Quantitative results of the subapical cell length and the division angle (Ɵ) of

indicated lines, respectively. Bold horizontal lines indicate the median and whiskers

indicate the first and the third quarter of the data value. ***P<0.001; Student's *t*-test.

(F) Above (upper panels) and side (lower panels) views of indicated *M. polymorpha* lines.

Tak-1 (WT) expands its thallus horizontally on the agar surface. The overexpression of

MpPINZ-, AtPIN1-, and AtPIN2-GFP showed vertical thallus growth, which generated a

large angle between the lower surface of the thallus and the surface of the agar (Ɵ).

Scale bar = 0.5 cm.

(G) The quantification of the thallus growth angle (Ɵ) showed in A. Bold horizontal lines

indicate the median and whiskers indicate the first and the third quarter of the data

value. ***P<0.001; Student's *t*-test.

Figure 4. Endogenous PINs present different localization patterns in different tissues.

(A) The localization of PpPINA-GFP in tissues with different complexities. PpPINA-GFP is

polarized to the tip of initial and elongated protonema cells. The initial protonema cell

is regenerated from a detached leaf and the representative image shows the

maximum projection with a 5 µm thick Z-section. Auto-fluorescence of the

chloroplasts is indicated by asterisks. In elongated protonema cell, the polarity of

PpPINA-GFP is plotted by a intensity measurement along the PM as the yellow arrows

represented. The same measurement is applied to (B) and (C). In complex tissues

composed of multiple cell layers e.g. young leaves in *P. patens*, PpPINA-GFP is polarized

at both apical and basal domains (white arrowheads). Scale bars = 10 µm.

(B) The localization of MpPINZ-GFP in emerging rhizoids, young rhizoids, and gemma

epidermal cells. Representative image with a 5 µm thick Z-section at the gemma

surface shows the accumulation of MpPINZ-GFP at tips of emerging rhizoids as

indicated by yellow arrowheads in rhizoid precursor cells (asterisks). MpPINZ-GFP is

polarized at the tip of young rhizoids (middle). A middle section image showing the

even distribution of MpPINZ-GFP on the PM of gemmae composed of multiple cell

layers. Scale bars = 10 µm

(C) AtPIN2-GFP shows a polarized signal at the tip of the initial root hair but the

polarized signal is not observed in elongated root hairs. All imaging details are

described in Methods. The pixel values ranging from 0 to 255 are represented by the

rainbow color. In Arabidopsis epidermal cells, AtPIN2-GFP shows apical polarization as

white arrowheads indicated. Scale bar, 10 µm for root hairs, and 1 cm for epidermis.

Figure 5. Xenologous PINs are PM localized with no defined polarity.

(A) Overexpressed PpPINA-GFP is polarized to the tip of protonema, but overexpressed

AtPIN1-GFP is evenly distributed on the PM with intracellular puncta. The occurrence

frequency is indicated in the left lower corner in each image. Scale bars = 10 µm.

(B) Overexpressed MpPINZ-GFP is polarized to the tip of young rhizoids, while

overexpressed AtPIN1-GFP shows strong cytosolic signals and weak PM localization

with no polarity. Scale bars = 10 µm.

(C) AtPIN2-GFP under its native promoter is apicalized in epidermal cells, but PpPINA-

GFP and MpPINZ-GFP are mislocalized to the basal and lateral sites as white

arrowheads indicated. Scale bar = 1 cm.

(D) The intensity plots of the representative images for PpPINA-GFP, AtPIN1-GFP in *P.*

patens protonema cells and MpPINZ-GFP and AtPIN1-GFP in *M. polymorpha* young

rhizoids.

(E) Polarity index (ratio of signal intensity at the apical plasma membrane/ signal

intensity at the lateral plasma membrane) for apicalization of the indicated PIN-GFP

in Arabidopsis epidermal cells. PpPINA-GFP and MpPINZ-GFP are significantly lower

in the polarity index. 12-20 cells from 3 roots in three independent experiments for

each line. ***$P<0.001$; Student's *t*-test.

Figure 6. Cytoskeletal networks are important for the polarization of mainly bryophytic PINs.

(A) PpPINA-GFP is polarized to the tip of protonema, and the disruption of actin filaments

by Latrunculin (LatB) caused its hyperpolarization (white arrowhead). The disruption

of microtubules by Oryzalin (Ory) disturbed the polarization of PpPINA-GFP. Scale bars

= 10 µm.

(B) MpPINZ-GFP is polarized at the tip of young rhizoids, and its polarization was only

abolished by Oryzalin treatment (white arrowhead). Scale bars = 10 µm.

(C) AtPIN2-GFP is polarized at the apical site of epidermal cells, and disruption of either

actin filaments or microtubules did not change its polarization (white arrowheads).

Scale bars = 1 cm.

(D) Relative intensity of PpPINA-GFP and MpPINZ-GFP along the cell membrane with

indicated treatment.

(E) Polarity index of AtPIN2-GFP in indicated treatment, no significance is shown. The

concentration and duration of treatments for each species are described as in

Methods.

Figure 1. Phosphorylation sites are highly conserved between bryophytic and Arabidopsis PINs.

Figure 1. Phosphorylation sites are highly conserved between bryophytic and Arabidopsis PINs.

(A) Phylogenetic analysis of canonical PINs from early-divergent plants *P. patens (Pp), M. polymorpha (Mp),* and representative angiosperm *A. thaliana (At)*. The unrooted tree shows the relationships of different PINs in two representative bryophytes and Arabidopsis. The scale bar represents the number of changes per site.

(B) Identity indexes of all PINs compared to AtPIN2 with full length or only the hydrophilic loop region of coding amino acid sequences. Identity indexes of all PINs compared to AtPIN1 is shown in Supplemental Figure 1.

(C) Alignment of PIN amino acid sequences. Identical amino acids are highlighted with blue column. The rea and green boxes show the hydrophilic loop regions bearing conserved phosphorylation sites, which are enlarged in (D).

(D) Four conserved phosphorylation sites, labeled by S1 to S4, were verified in previous AtPIN2 studies and are depicted with black frames. Note that S1 to S4 are conserved in every examined PINs.

Figure 2. The hydrophilic loops in AtPIN2, PpPINA, and MpPINZ are less conserved.

Figure 2. The hydrophilic loops in AtPIN2, PpPINA, and MpPINZ are less conserved.

(A) Structural alignment of AtPIN1 and AtPIN2. The protein structures are predicted by Alphafold2 and aligned in ChimeraX software. The structural conserved regions are indicated by black arrowheads, and the non-conserved regions are indicated by empty arrowheads.

(B) Structural alignment of AtPIN2 with PpPINA and MpPINZ. Only transmembrane domains and one loop aligned with each other, while the majority of hydrophilic loop regions are not conserved. The four conserved phosphorylation sites are labeled with atom details in ball-and-stick style.

(C) – (E) Individual protein structure retrieved from (B). The four conserved phosphorylation sites are indicated by arrowheads.

Figure 3. GFP-fused PIN proteins possess auxin export activity.

Figure 3. GFP-fused PIN proteins possess auxin export activity.

(A) The insertion site of GFP in indicated PIN proteins. Numbers represent the amino acid position in respective proteins.

(B) The auxin export assay with *P. patens* transgenic plants. Fresh tissues were incubated with radioactive H^3 IAA, followed by washing and the radiative H^3-IAA exported in the new culture medium was measured with the scintillation detector after one day.

(C) Representative protonema cells of the indicated genetic background. The cell outline was stained with FM4-64. White arrowheads indicate the first cell division plane, and the yellow double arrow indicates the cell length of the subapical cell. Scale bar = 100 μm.

(D) and (E), Quantitative results of the subapical cell length and the division angle (Θ) of indicated lines, respectively. Bold horizontal lines indicate the median and whiskers indicate the first and the third quarter of the data value. ***P<0.001; Student's *t*-test.

(F) Above (upper panels) and side (lower panels) views of indicated *M. polymorpha* lines. Tak-1 (WT) expands its thallus horizontally on the agar surface. The overexpression of MpPINZ-, AtPIN1-, and AtPIN2-GFP showed vertical thallus growth, which generated a large angle between the lower surface of the thallus and the surface of the agar (Θ). Scale bar = 0.5 cm.

(G) The quantification of the thallus growth angle (Θ) showed in A. Bold horizontal lines indicate the median and whiskers indicate the first and the third quarter of the data value. ***P<0.001; Student's *t*-test.

Figure 4. Endogenous PINs present different localization patterns in different tissues.

Figure 4. Endogenous PINs present different localization patterns in different tissues.

(A) The localization of PpPINA-GFP in tissues with different complexities. PpPINA-GFP is polarized to the tip of initial and elongated protonema cells. The initial protonema cell is regenerated from a detached leaf and the representative image shows the maximum projection with a 5 μm thick Z-section. Auto-fluorescence of the chloroplasts is indicated by asterisks. In elongated protonema cell, the polarity of PpPINA-GFP is plotted by a intensity measurement along the PM as the yellow arrows represented. The same measurement is applied to (B) and (C). In complex tissues composed of multiple cell layers e.g. young leaves in *P. patens*, PpPINA-GFP is polarized at both apical and basal domains (white arrowheads). Scale bars = 10 μm.

(B) The localization of MpPINZ-GFP in emerging rhizoids, young rhizoids, and gemma epidermal cells. Representative image with a 5 μm thick Z-section at the gemma surface shows the accumulation of MpPINZ-GFP at tips of emerging rhizoids as indicated by yellow arrowheads in rhizoid precursor cells (asterisks). MpPINZ-GFP is polarized at the tip of young rhizoids (middle). A middle section image showing the even distribution of MpPINZ-GFP on the PM of gemmae composed of multiple cell layers. Scale bars = 10 μm

(C) AtPIN2-GFP shows a polarized signal at the tip of the initial root hair but the polarized signal is not observed in elongated root hairs. All imaging details are described in Methods. The pixel values ranging from 0 to 255 are represented by the rainbow color. In Arabidopsis epidermal cells, AtPIN2-GFP shows apical polarization as white arrowheads indicated. Scale bar, 10 μm for root hairs, and 1 cm for epidermis.

Figure 5. Xenologous PINs are PM localized with no defined polarity.

Figure 5. Xenologous PINs are PM localized with no defined polarity.

(A) Overexpressed PpPINA-GFP is polarized to the tip of protonema, but overexpressed AtPIN1-GFP is evenly distributed on the PM with intracellular puncta. The occurrence frequency is indicated in the left lower corner in each image. Scale bars = 10 μm.

(B) Overexpressed MpPINZ-GFP is polarized to the tip of young rhizoids, while overexpressed AtPIN1-GFP shows strong cytosolic signals and weak PM localization with no polarity. Scale bars = 10 μm.

(C) AtPIN2-GFP under its native promoter is apicalized in epidermal cells, but PpPINA-GFP and MpPINZ-GFP are mislocalized to the basal and lateral sites as white arrowheads indicated. Scale bar = 1 cm.

(D) The intensity plots of the representative images for PpPINA-GFP, AtPIN1-GFP in *P. patens* protonema cells and MpPINZ-GFP and AtPIN1-GFP in *M. polymorpha* young rhizoids.

(E) Polarity index (ratio of signal intensity at the apical plasma membrane/ signal intensity at the lateral plasma membrane) for apicalization of the indicated PIN-GFP in Arabidopsis epidermal cells. PpPINA-GFP and MpPINZ-GFP are significantly lower in the polarity index. 12-20 cells from 3 roots in three independent experiments for each line. ***$P<0.001$; Student's *t*-test.

Figure 6. Cytoskeletal networks are important for the polarization of mainly bryophytic PINs.

Figure 6. Cytoskeletal networks are important for the polarization of mainly bryophytic PINs.

(A) PpPINA-GFP is polarized to the tip of protonema, and the disruption of actin filaments by Latrunculin (LatB) caused its hyperpolarization (white arrowhead). The disruption of microtubules by Oryzalin (Ory) disturbed the polarization of PpPINA-GFP. Scale bars = 10 µm.

(B) MpPINZ-GFP is polarized at the tip of young rhizoids, and its polarization was only abolished by Oryzalin treatment (white arrowhead). Scale bars = 10 µm.

(C) AtPIN2-GFP is polarized at the apical site of epidermal cells, and disruption of either actin filaments or microtubules did not change its polarization (white arrowheads). Scale bars = 1 cm.

(D) Relative intensity of PpPINA-GFP and MpPINZ-GFP along the cell membrane with indicated treatment.

(E) Polarity index of AtPIN2-GFP in indicated treatment, no significance is shown. The concentration and duration of treatments for each species are described as in Methods.