The "LINC" in between Δ40p53-miRNA axis in the regulation of cellular processes

Apala Pal[1] and Saumitra Das[1*]

1   [1]Department of Microbiology and Cell Biology, Indian Institute of Science,

2   Bangalore-560012, India.

*Address for correspondence:

Dr. Saumitra Das, Professor,

Department of Microbiology and Cell Biology,

Indian Institute of Science, Bangalore-560012, India

Email: sdas@iisc.ac.in

Phone: 91 80 2293 2886

Fax: 91 80 2360 2697

ABSTRACT

Recently, we showed that Δ40p53, the translational isoform of p53, can inhibit cell growth independently of p53 by regulating microRNAs. Here, we explored the role of Δ40p53 in regulating the lncRNA-miRNA axis. Based on early research from our laboratory, we selected LINC00176 for further exploration. LINC00176 levels were affected by the overexpression and knockdown of Δ40p53. Further, under DNA damage, ER stress, and glucose deprivation, LINC00176 was upregulated in HCT116 p53-/- cells (harboring only Δ40p53) compared to HCT116 p53+/+ cells. Additionally, ChIP and RNA stability assays revealed that Δ40p53 transactivates LINC00176 transcriptionally and stabilizes it post-transcriptionally. We ectopically overexpressed and knocked down LINC00176 in HCT116 p53-/- cells, which affected proliferation, cell viability, and the expression of epithelial markers. Finally, RNA pulldown revealed that LINC00176 sequesters several putative miRNA targets. These results provide important insights into the pivotal role of Δ40p53 in regulating the lncRNA-miRNA axis.

Keywords: p53 isoform; Δ40p53; LINC00176; miRNA; cellular effect of Δ40p53

INTRODUCTION

The tumor suppressor p53 is a master transcriptional regulator that regulates several critical events during cell fate determination, including transcriptional regulation of multiple genes involved in cell survival, cell cycle regulation, apoptosis, and metabolism (1). p53 has 12 isoforms created by alternate promoters, splicing sites, and translation initiation sites. These different isoforms have unique functions and are capable of modulating p53 activity (2). The Δ40p53 isoform is the only translational isoform of p53. Our laboratory has shown that translation initiation from the first IRES (Internal Ribosomal Entry Site) of *TP53* generates full-length p53 protein, whereas translation initiation from the second IRES generates Δ40p53 (3). Various ITAFs (IRES *trans* acting factors) are required for the IRES-mediated translation of the two isoforms under different stress conditions (4-6). We have also shown how a protein and miRNA crosstalk at the TP53 3'-UTR influences differential expression of p53 and Δ40p53 (7).

These two isoforms have a robust affinity toward each other as both contain the oligomerization domain, and they can therefore form homo- and hetero-tetramers (8). Δ40p53 can both positively and negatively regulate p53 activity: it can alter the recruitment of p53

co-activators, increasing its activity (9), but it can negatively regulate the transactivation of p53 target genes, such as p21 (10). Δ40p53 also has p53-independent functions. It controls the switch from pluripotency to differentiation in somatic cells and helps maintain pluripotency in embryonic stem cells (11). It also regulates proliferation and glucose homeostasis in mice (12). Moreover, it uniquely regulates genes involved in apoptosis (13) and senescence (14). In certain cancers, full-length p53 is insufficient to protect against tumorigenesis, and Δ40p53 is essential for mediating apoptosis through the selective activation of specific target genes (15). Recent observations also suggest a role for Δ40p53 as a tumor suppressor in hepatocellular carcinoma (16). One of our latest studies showed that Δ40p53 can specifically regulate miR-186-5p (17). However, although some studies have examined the functional role of Δ40p53, there has been little detailed investigation regarding intermediate molecules (such as RNAs or proteins) through which Δ40p53 exerts its influence on cells.

Some RNAs in the p53 response pathway have been shown to be involved in tumor suppression (18). Among the cellular pool of RNAs, lncRNAs (long non-coding RNAs) have gained attention in cancer because of their transactivation functions. LncRNAs are a heterogeneous class of noncoding RNAs >200 nucleotides in length with very low or no coding potential. Several lncRNAs crosstalk with p53, thereby regulating cellular processes including cell proliferation, differentiation and development, chromosomal imprinting, and genomic stability (19-21). Therefore, we were interested in exploring the regulation of long noncoding RNAs by Δ40p53.

To understand the potential role of Δ40p53-mediated regulation of lncRNAs, we investigated a cellular lncRNA profile (unpublished observation) in our laboratory in H1299 cells (a p53-null cell line) with overexpression of full-length p53, Δ40p53, or both isoforms. Preliminary results suggested that several lncRNAs are differentially regulated under these conditions. Interestingly, LINC00176 was predominantly regulated by Δ40p53. LINC00176 is a long noncoding RNA that is differentially expressed in cancer and normal tissues and may have a tumor-suppressive role. Recent studies have demonstrated that LINC00176 regulates the expression of microRNAs in hepatocellular carcinoma (22) and that it promotes ovarian cancer progression by downregulating ceruloplasmin (23). By contrast, a report on genome-scale CRISPR screening has emphasized its possible role as a tumor suppressor lncRNA (24). LINC00176 is regulated through several pathways involving different upstream regulators and downstream targets. Based on the results obtained from our preliminary experiments, we decided to focus on LINC00176 and its differential regulation by p53 isoforms. We found

75   that LINC00176 is predominantly regulated by Δ40p53. Therefore, in this study, we explored

76   the role of the Δ40p53/p53-LINC00176 axis in microRNA-mediated regulation of cell

77   growth and proliferation.

MATERIALS AND METHODS

Bioinformatic Analysis

GEPIA (http://gepia.cancer-pku.cn/detail.php?gene=LINC00176) was used for analysing

differential LINC00176 levels in cancer. LINC00176 was queried in the database and the

results were generated by selecting Cancer versus Normal Analysis (25). starBase v2.0

database (https://starbase.sysu.edu.cn/starbase2/browseNcRNA.php) was used to identify

potential miRNA targets of LINC00176 (26). catRAPID fragments software was used to

predict possible protein binding regions in LINC00176

(http://service.tartaglialab.com/page/catrapid_group_old), and the interaction score between

LINC00176 and the Δ40p53 was predicted by RPISeq software

(http://pridb.gdcb.iastate.edu/RPISeq/results.php). PROMOALGEN has been used to predict

transcription factor binding site in *LINC00176* gene (http://alggen.lsi.upc.es/cgi-

bin/promo_v3/promo/promoinit.cgi?dirDB=TF_8.3). dbDEMC2.0 has been used to perform

a meta profiling of selected miRNAs (https://www.biosino.org/dbDEMC/help#help00) (27).

Cell lines and transfections

We have used three cell lines in the current study: H1299 (lung adenocarcinoma cell line

lacking p53/Δ40p53), HCT116 p53+/+ cells (colon carcinoma cell line harboring wild-type

p53; hereafter called HCT116+/+); and HCT116 p53-/- cells (colon carcinoma cell line

harboring only Δ40p53; hereafter called HCT116-/-). These cells were maintained in DMEM

(Sigma) with 10% Foetal Bovine Serum (GIBCO, Invitrogen). 70%-80% confluent

monolayer of cells was transfected with various plasmid constructs using Lipofectamine 2000

(Invitrogen) and Turbofectamine in Opti-MEM (Invitrogen). The medium was replaced with

DMEM (with antibiotic) and 10% FBS four hours later. The cells were harvested and

processed as required at the desired time point. For the stress-induced experiments, the cells

were treated with doxorubicin (2μM) for 16 h, Thapsigargin (0.1 μM, 1 μM) for 16 h,

DMEM minus Glucose medium for 8 h. For stability experiments cells were transfected with

siRNA directed to Δ40p53, post 40 h of transfection cells were treated with Actinomycin D

(1 mg/ml) and cells were harvested at 0 h, 4 h and 8 h post treatment.

Plasmids and constructs

pGFP-hp-p53- 5'UTR-cDNA (14A): It expresses both p53FL and Δ40p53 (a generous gift from Dr. Robin Fahraeus, INSERM, France). pGFP-hp-p53-5'UTR (A135T)-cDNA which expresses onlyΔ40p53 and pGFP-hp-p53- 5'UTR (A251G/T252 C/G253T)-cDNA which expresses only full-length p53(p53FL) (17). Two constructs were used for Tagged RNA Affinity Purification: Plasmid pMS2 and pMS2-GST. (Both the constructs were a generous gift from Je-Hyun Yoon and Myriam Gorospe Laboratory of Genetics and Genomics, National Institutes of Health, Baltimore, MD 21224, USA) (28,29). LINC00176 (3495–4302) construct obtained in the study (a generous gift from Dr. T Tamura Institut fuer Biochemie, Hannover, Germany) has been used to generate the overexpression construct. LINC00176 overexpression construct was generated by cloning the insert into pMS2 plasmid between HindIII and EcoRI sites. 5'CCGGTGTCGTCTTGCAAACATTAAACTCGAGTTTAATGTTTGCAAGACGAC ATTTTTG-3' sequence of shRNA was used to target LINC00176.

siRNA transfections

HCT116+/+ and HCT116-/- cells were transfected with 30 nM si p53 RNA (IDT). 5'-AACCUCUUGGUGAACCUUAGUACCU-3' is the sequence for p53 siRNA, directed against the 3'UTR of p53; therefore, it targets both p53 as well as Δ40p53. A non-specific siRNA (Dharmacon) was used in the partial silencing of p53/Δ40p53 in the experiments as control. si-LINC00176 (5'-CUCGUUCUGUAGACUUGUU-3') was used for partial silencing of LINC00176 (a generous gift from Dr. T Tamura, Institut fuer Biochemie, Hannover, Germany) (22).

Western Blot analysis

Protein concentrations of the extracts were assayed by Bradford (Bio-Rad) and equal amounts of cell extracts were separated by SDS 12% PAGE and transferred to nitrocellulose membrane (Bio-Rad). Samples were then analyzed by Western blotting using rabbit-raised anti-p53 polyclonal antibody (CM1, kind gift from Dr. Robin Fahraeus, INSERM, France and Prof. J.C. Bourdon, of University of Dundee, UK), followed by secondary antibody (horseradish peroxidase-conjugated anti-mouse or anti-rabbit IgG; Sigma). Mouse-monoclonal anti-β-actin antibody (Sigma) was used as a control for equal loading of total cell extracts. 1801 antibody (Santa Cruz Biotechnology, Cat. No. SC-98) was used to detect p53 and Δ40p53. E-cadherin antibody (Santa Cruz Biotechnology, Cat. No. SC-8426), Slug

antibody (Santa Cruz Biotechnology, Cat. No. SC-166476), Bip antibody (Abclonal, Cat. No. A4908) and GST Antibody (B-14) (Santa Cruz Biotechnology, Cat. No. SC-138) was used to detect E-cadherin, Slug, Bip and GST proteins respectively.

RNA isolation and Real time PCR

Total RNA was isolated from cells with TRI Reagent ^{TM} (Sigma), according to manufacturer's protocol. The isolated RNA was treated with 10 units of DNase I (Promega), extracted with acidic phenol-chloroform, and precipitated with 3 M sodium acetate (pH 5.2) and 2.5 volumes of absolute alcohol. RNA amount was quantified with Nano-spectrophotometer and cDNAs were synthesized using specific reverse primers and MMLV RT (Revertaid ^{TM} Thermo Scientific) at 42°C for 1 h, according to standard protocol. While 2-5µg of total RNA has been used for checking the expression of mRNAs and lncRNAs, 50ng of total RNA has been used to check the expression of miRNAs.

SYBR green Assay System was used for mRNA, lncRNA and miRNA detection and quantification. We have used actin as endogenous control for mRNA, lncRNA and 5SrRNA as endogenous control for miRNAs. The thermocycling conditions for SYBR green Assay system include 1 cycle of 95°C for 10 min, and 40 cycles of 95°C for 15 s, and 60°C for 30 s and 72°C for 30 s (Applied Biosystems). $2^{-\Delta\Delta Ct}$ method algorithm was used to analyse the relative changes in expressions, where actin/5S served as an endogenous control. The fold change was calculated using $2^{-\Delta\Delta Ct}$. ΔCt = Ct (target gene) −Ct (endogenous control) and $\Delta\Delta Ct$ = ΔCt target sample) −ΔCt (control sample). Melting curve analysis of every (q)PCR was conducted after each cycle.

Tagged RNA Affinity Pull down Assay (TRAP Assay)

The method is based on the addition of MS2 RNA hairpin loops to a target RNA of interest, followed by co-expression of the MS2-tagged RNA together with the protein MS2 (which recognizes the MS2 RNA elements) fused to an affinity tag (29). After purification of the MS2 RNP complex, the miRNAs present in the complex are identified. In this study we have tagged the LINC00176 with MS2 hairpins and have co-expressed it in Human colon carcinoma cells (HCT116-/-) along with the chimeric protein MS2-GST (glutathione S-transferase). After affinity purification using glutathione SH beads, the microRNAs present in the RNP complex were identified by reverse transcription (RT) and real-time, quantitative (q)PCR.

MTT Assay

HCT116-/- cells were transfected with either LINC00176 o/e or sh-LINC00176 construct, 36 h post transfection these cells were re-seeded in 96 well plate. After 20 h of reseeding 10 µl of MTT (50 mg/ml) reagent was added and cells were incubated for 3-4 h, following which the supernatant was removed and cells were dissolved in 200µl of DMSO. Cell viability was measured as a spectrophotometry reading of amount of MTT incorporated in the cell at 560 nm wavelength. The reading taken on the first day is considered as Day 0. Similarly, the reading was measured on Day 2 and Day 4 and Day 6 post re-seeding to check the effect on cell viability over a period of time.

Cell cycle Analysis

HCT116-/- cells were transfected with either LINC00176 o/e or sh-LINC00176 construct, 48 h post transfection the cells were harvested and fixed with methanol. The fixed cells were treated with RNase A (10 mg/ml) for 1 h at 37°C, following which PI (Propidium iodide-1mg/ml) was added to the cells and taken for Flow cytometric analysis.

Colony formation Assay

HCT116-/- cells were transfected with either LINC00176 o/e or sh-LINC00176 construct. 36 h post transfection 1000 cells from each condition were reseeded on 6 well plate and was incubated for 10-14 days to observe the number of colonies obtained. Post 10-14 days, cells were harvested by directly adding methanol to the plates for cell fixation, followed by which the cells were stained with Crystal Violet solution. After the cells got stained, the excess colour was washed with water 2-3 times, following which the colonies were quantified for any differences in number.

BrdU cell proliferation assay

BrdU cell proliferation assay was performed as described in BrdU Cell Proliferation ELISA Kit (ab126556). Briefly, HCT116-/- cells were transfected with si-LINC00176. 36 h post transfection these cells were reseeded in 96 well plate. After 12 h of reseeding BrdU solution was added and cells were incubated for 12 h, followed by the protocol as mentioned in the manual.

Statistical analysis

The data were expressed as mean ± SD. Statistical significance was determined using two-sided Student's t-test. The criterion for statistical significance was $p \leq 0.05$ (*) or $p \leq 0.01$ (**) or $p \leq 0.001$(***)

RESULTS

Selection and validation of LINC00176 for further studies

We identified LINC00176 based on a preliminary screen (unpublished data from our laboratory) of lncRNA expression in cells expressing different p53 isoforms. As mentioned earlier, LINC00176 is known to have multiple regulators that govern important cancer-driven processes. We assessed the expression of LINC00176 using the GEPIA database. LINC00176 was differentially expressed in different cancer types **(Figure S1A).** Box plot analysis suggested that the expression of LINC00176 was lower in tumor tissues (lung and colon) than in normal tissues, which is analogous to WT p53 and Δ40p53 levels in cancer **(Figure S1B)**. To draw a direct association with p53, we assessed the correlation between p53 and LINC00176 expression and found that in normal tissues, LINC00176 was positively correlated with p53 levels **(Figure S1C, E, G).** However, in tumor tissues, they were negatively correlated **(Figure S1D, F, H)**. This observation suggests that LINC00176 may be positively regulated by WT p53 in normal conditions and negatively regulated by mutant p53 in tumor conditions. Since Δ40p53 was not annotated in the GEPIA database, we could not analyze the correlation between Δ40p53 and LINC00176. However, we hypothesize that Δ40p53 is one of the most important isoforms of p53 and could be a potential regulator of LINC00176.

Given the distinctive functions of LINC00176 reported in the literature, we investigated its regulation by p53 isoforms. To directly investigate the role of p53 isoforms on LINC00176, we examined the expression of LINC00176 in three cancer cell lines: H1299 cells with ectopic overexpression of p53, Δ40p53, or both (14A construct); HCT116+/+ and HCT116-/- cells. LINC00176 levels were first validated in H1299 cells after overexpressing p53, Δ40p53, or both p53 and Δ40p53. It was upregulated under all conditions compared to control cells **(Figure 1A, B)**. However, the fold change was highest in cells with Δ40p53 overexpression, suggesting that Δ40p53 might be a more important regulator than p53. Since it would be more physiologically relevant to examine LINC00176 in a cell line with endogenous expression of p53, we also examined the levels of LINC00176 in HCT116+/+

and HCT116-/- cells. Interestingly, we found that the levels of LINC00176 were much higher

in HCT116-/- cells than in HCT116+/+ cells **(Figure 1C, D)**. To confirm the ability of p53 to

regulate LINC00176, an siRNA targeting to the 3'-UTR of *TP53* was transfected into

HCT116+/+ and HCT116-/- cells **(Figure 1F, H)**. The levels of LINC00176 decreased in

both cell lines after siRNA transfection **(Figure 1E, G)**. However, we did not observe a

significant decrease in LINC00176 in either cell line, which may be because p53/Δ40p53 are

not the only regulators of LINC00176. Nevertheless, silencing Δ40p53 in HCT116-/- cells

(Figure 1G) led to a greater reduction in LINC00176 levels than silencing p53 in

HCT116+/+ cells **(Figure 1E)**. To further examine whether both isoforms could modulate the

levels of LINC00176, p53 was overexpressed in HCT116-/- cells **(Figure 1J)**, which reduced

LINC00176 levels similar to those of 14A construct (expressing both p53 and Δ40p53)

observed earlier **(Figure 1I)**. These preliminary results suggest that LINC00176 is regulated

by both p53 translational isoforms, but the levels differ with respect to isoform abundance.

When p53 or Δ40p53 is present alone, either protein can upregulate LINC00176; however,

there is higher expression of LINC00176 when only Δ40p53 is present.

Regulation of LINC00176 under different stress conditions

p53 and Δ40p53 are translated by an IRES-mediated mechanism under stress conditions, such

as DNA damage, ER stress, and glucose starvation (4-6). Cells were treated with doxorubicin

and thapsigargin for 16 h to induce DNA damage and ER stress, respectively, and cultured in

glucose-deprived DMEM for 8 h to induce glucose starvation **(Figure 2A)**. In HCT116-/-

cells, Δ40p53 levels increased under all stress conditions **(Figure 2C**, **E, G).** LINC00176

levels also increased, with the highest increase observed in cells treated with thapsigargin

(Figure 2B, D, F). In HCT116+/+ cells, LINC00176 was upregulated in the doxorubicin-

treated and glucose-deprived cells; however, in cells treated with thapsigargin, there was no

significant change **(Figure S2 A-F)**. It is unclear whether p53 or Δ40p53 was responsible for

the increase in LINC00176 in HCT116+/+ cells as both were found increased upon induction

of stress. However, in line with our earlier observation of dominant regulation of LINC00176

by Δ40p53, the results in HCT116-/- cells were more significant and consistent.

Next, we wanted to confirm whether Δ40p53 alone was involved in regulating LINC00176

levels under stress or whether another factor could also regulate LINC00176. We selected ER

stress as it maximally induced Δ40p53 in this model, and we observed significant

upregulation of LINC00176 in cells treated with thapsigargin **(Figure 2D)**. We knocked

down Δ40p53 in HCT116-/- cells, followed by induction of ER stress **(Figure 2H)**.

Compared to control cells that were untreated and not transfected with siΔ40p53, there was

no change in LINC00176 levels in the cells transfected with siΔ40p53 and treated with

thapsigargin **(Figure 2I-K),** indicating that Δ40p53 is a major regulator of LINC00176.

These results suggest that the expression of LINC00176 is closely linked to Δ40p53 levels.

However, we cannot exclude the possibility that other factors might also be involved in

regulating LINC00176 expression.

Mechanism of LINC00176 regulation by Δ40p53/p53

Next, we investigated whether LINC00176 is directly regulated by Δ40p53/p53 **(Figure 3A)**.

Using PROMO ALGGEN, we found that there are several Δ40p53/p53 binding sites in the

LINC00176 gene, which suggests that Δ40p53/p53 may regulate LINC00176 at the

transcriptional level. An earlier study on LINC00176 revealed that it has three putative

promoter regions in different cell types (22,24). We used ChIP experiments to examine the

association of Δ40p53/p53 with these three probable promoter regions, one located at the

proximal region of LINC00176 exon 1 (62665399–62666196) and two located at the

proximal region of E2 (62667411–62667655 and 62667984–62668233). We

immunoprecipitated HCT116-/- and HCT116+/+ crosslinked lysates with 1801 antibody

which can detect both Δ40p53 and p53 proteins **(Figure S3A, B).** The results showed an

association between Δ40p53 and the *LINC00176* promoters in HCT116-/- cells. However, no

such association was found for p53 in HCT116+/+ cells **(Figure 3B, C)**. Because we

observed a decrease in LINC00176 levels upon p53 overexpression in HCT116-/- cells, we

also examined the effect of ectopic p53 on the binding of Δ40p53 to *LINC00176* promoters in

HCT116-/- cells. Interestingly, we found that the binding of Δ40p53 to all three *LINC00176*

promoters decreased with p53 overexpression in HCT116-/- cells **(Figure 3D, E)**. This

observation is in accordance with the reported literature that p53 and Δ40p53 regulate each

other's activity through the formation of hetero-oligomers (4). Hence, these results strengthen

the observation that Δ40p53 controls LINC00176 regulation more than p53.

We also wanted to further assess the ability of Δ40p53/p53 to directly regulate LINC00176

at the RNA level. p53 is known to directly affect certain RNAs through its C-terminal domain

(30) which is shared with Δ40p53. We aimed to determine whether Δ40p53 could regulate

LINC00176 RNA directly or indirectly. Analysis using CATRAPID and RPI-Seq software

was used to identify potential interactions between mature LINC00176 and Δ40p53 **(Figure**

S3 C-F). Both the databases predicted probable binding regions and positive interaction score

for LINC00176 and the p53/Δ40p53. Because the protein domains of Δ40p53, which might

bind to LINC00176, are common with p53, we cannot exclude the possibility that there could

be an interaction between LINC00176 and p53. However, based on our initial results, we

focused on the Δ40p53-LINC00176 interaction. Based on the probable RNA binding, we

treated cells with actinomycin D after transfection with siΔ40p53 to assess LINC00176

stability **(Figure 3F).** The results indicate that knockdown of Δ40p53 decreased LINC00176

levels after treatment with actinomycin D, indicating that the stability of LINC00176 is

increased by Δ40p53 **(Figure 3G, H)**. Therefore, the increased LINC00176 levels observed

in the presence of Δ40p53 in comparison to p53 could be due to multistep regulation at both

the transcriptional and RNA stability levels.

Deciphering the mechanism of action of LINC00176

Thus far, the results obtained in this study indicate that Δ40p53 can regulate long noncoding

RNAs in addition to miRNAs (17). Therefore, it is important to understand how LINC00176

functions in cells and what its mechanism of action could be **(Figure 4A)**. Long noncoding

RNAs are known to interact with miRNAs, acting as a decoy or promoting their degradation,

thereby impacting cellular processes (31). Therefore, miRNA-interacting partners of

LINC00176 were predicted using the Starbase V.2.0 database **(Figure 4B),** and their levels

were measured in HCT116+/+ and HCT116-/- cells **(Figure 4C)**. The expressions of these

miRNAs (miR-761, miR-184, miR-503-5p, miR-15b-5p, miR-138, and miR-9-5p) varied.

Since most had lower expression in HCT116-/- cells than in HCT116+/+ cells, they could be

possible targets of LINC00176, as LINC00176 levels are very high in HCT116-/- cells. In

particular, miR-9-5p, which is a known target of LINC00176, was downregulated in

HCT116-/- cells compared to HCT116+/+ cells. To assess the direct effect of LINC00176 on

these miRNAs, LINC00176 was silenced in HCT116-/- cells, and the corresponding changes

in miRNA levels were assessed. Upon silencing LINC00176, the levels of LINC00176 were

reduced by 60% compared to that in control cells **(Figure 4D)**. The levels of the miRNAs

increased from 30% to 80% **(Figure 4E)**.

To investigate whether LINC00176 sequesters target miRNAs, tagged RNA affinity

pulldown assays were performed **(Figure 4G).** miRNA levels were quantified in the

pulldown fraction and normalized to input levels (**Figure 4F**). miR-184, miR-138, miR-503,

miR-15b-5p, and miR-9-5p were significantly enriched in the pulldown fraction, indicating

that they are direct targets of LINC00176. However, miR-761 did not show any enrichment

in the pulldown fraction, implying that it might have an indirect interaction with LINC00176

or could be degraded in a complex. These results indicate that LINC00176 is capable of

titrating miRNA levels in cells, which may affect cellular processes.

The impact of the Δ40p53-LINC00176 axis on cellular behavior

LncRNAs regulate a wide variety of cellular processes (32). To understand the effect of

LINC00176 on cellular behavior, we overexpressed or knocked down LINC00176 in

HCT116-/- cells and examined the effects **(Figure 5A-C)**. First, we examined the effect of

LINC00176 on EMT marker expression. LINC00176 overexpression increased the levels of

E-cadherin (epithelial marker) and decreased the levels of Slug and vimentin (mesenchymal

markers) **(Figure 5D, E)**. Interestingly, LINC00176 overexpression also increased Δ40p53

levels. By contrast, knocked down LINC00176 decreased the mRNA levels of E-cadherin but

increased those of vimentin and Slug **(Figure 5F);** however, we did not observe the same

trend for the protein levels of E-cadherin and Slug **(Figure S5A)**. We further analyzed cell

viability after LINC00176 overexpression or knockdown and observed that LINC00176

overexpression decreased cell viability **(Figure 5G, S5 B, C)**, whereas LINC00176

knockdown increased cell viability **(Figure 5H, S5 D, E)**. However, these effects on cell

viability may not be very significant, because LINC00176 might contribute to only a small

percentage of the vast majority of cellular processes. We also found that LINC00176

knockdown increased proliferation using the BrdU assay **(Figure 5I)**. Δ40p53 abundance is

closely linked with cell cycle phases (3); therefore, we also assessed the effect of LINC00176

on cell cycle progression. We observed a slight increase in the proportion of cells in G1 phase

and a slight decrease in the proportion of cells in S phase with LINC00176 overexpression

(Figure S5F). Furthermore, to check for any effect on cell growth, we performed a colony

formation assay, wherein we found a decrease in the number of colonies after LINC00176

overexpression **(Figure S5G)** and an increase in the number of colonies after LINC00176

knockdown **(Figure S5H)**. Together, these results indicate that LINC00176 promotes the

epithelial phenotype and suppresses the mesenchymal phenotype, cell growth, viability, and

proliferation, all of which suggest that this lncRNA contributes to tumor suppression in cells.

DISCUSSION

In cancer, lncRNAs are key players in the p53-dependent transcriptional pathway, wherein

lncRNAs and p53 can regulate each other (33). Therefore, to identify important cancer

signaling pathways, it is extremely important to identify novel lncRNAs that are regulated

through the p53 pathway. *TP53* is the most frequently mutated gene in human cancer (34);

however, these mutations are not found in all cancers, indicating that there could be other factors interfering with WT p53 function (35). These factors, which have complicated our understanding of p53 signaling, include more recently discovered p53 isoforms (36). Owing to the lack of detailed functional characterization, little information is available regarding p53 isoforms. However, one isoform that has recently gained attention in cancer research is Δ40p53, the only translational isoform of p53 (37). Although a few reports have shown the transactivation capacity of Δ40p53, there are no studies on the regulation of lncRNAs by Δ40p53. We selected LINC00176 for our current study because of its variable regulation by different transcription factors and differential expression in cancer.

Significant changes in the levels of LINC00176 were observed in H1299 cells expressing p53, Δ40p53, and both isoforms. However, the maximum upregulation was observed in cells expressing only Δ40p53. The upregulation of LINC00176 by Δ40p53 was also observed in other cell lines expressing only Δ40p53 (HCT116-/-) as compared to cells expressing both isoforms (HCT116+/+). Furthermore, LINC00176 was upregulated under different stress conditions, and its expression correlated with the abundance of Δ40p53. However, we cannot rule out the possibility that there could be other factors involved in the regulation of LINC00176 because of certain observations. First, we did not observe a significant decrease in LINC00176 levels upon knockdown of Δ40p53. Second, upon induction of ER stress in cells with Δ40p53 knockdown, LINC00176 levels did not change with respect to control cells. This indicates that although Δ40p53 could be a major regulator of LINC00176, there are other factors that regulate LINC00176 abundance. This regulation could be at the transcriptional or processing stages; therefore, we investigated any direct binding between the molecules.

Δ40p53 was previously thought to be devoid of transactivation capacity because it lacks TAD1 (38); however, subsequent studies reported that it can regulate some p53 target genes (13,39). It has also been reported that Δ40p53 is sufficient to induce cell death independent of p53 through a specific subset of genes (Fas, Dr5, Api1, and Pig3) (15). More recently, a study suggested an FLp53-independent, pro-tumoral role for Δ40p53 because of its ability to transactivate the transcription of antiapoptotic proteins, such as netrin-1 (40). Therefore, we examined the association of Δ40p53 and p53 with LINC00176. We observed binding of Δ40p53 to *LINC00176* promoters; surprisingly, p53 did not bind to *LINC00176* promoters. However, overexpression of p53 in HCT116-/- cells decreased the association of Δ40p53 with *LINC00176* promoters, suggesting that the formation of hetero-oligomers between p53 and Δ40p53 interfered with the transactivation function of Δ40p53, which is consistent with

the results of earlier reports (8). We also observed that Δ40p53 regulates LINC00176 not

only at the transcriptional level but also post-transcriptionally, since the stability of

LINC00176 also decreased after Δ40p53 knockdown. p53 is known to bind to the lncRNA

MEG3 (28); however, there are no previous reports of Δ40p53 binding to RNA. Whether

Δ40p53 can directly or indirectly bind to LINC00176 RNA remains unknown and requires

further investigation. Although we also observed LINC00176 upregulation with p53

expression, it is possible that p53 might regulate it through an unrelated mechanism.

The significance of LINC00176 upregulation by Δ40p53 was determined by examining

different miRNA targets. We identified different miRNA targets for LINC00176, from which

we selected a few targets based on the number of binding sites and their functions in different

cancers **(Figure S4)**. The majority of the selected miRNAs are differentially expressed in

cancer owing to their diverse regulatory pathways. Their levels were mostly decreased in

HCT116-/- cells compared to those in HCT116+/+ cells, and their expressions negatively

correlated with LINC00176 levels. Upon further analyzing their levels after knockdown of

LINC00176 and TRAP assays, we found that most of these miRNAs were sequestered by

LINC00176. However, although the level of miR-761 increased upon LINC00176

knockdown, it was not sequestered by LINC00176. As lncRNAs are known to serve as

miRNA decoys, LINC00176 may be degrading these miRNAs; however, the mechanism of

miRNA regulation by LINC00176 needs to be further examined.

Finally, we examined the direct effects of LINC00176 on cellular behavior through ectopic

overexpression or silencing. Based on the different hallmark processes of cancer (41) we

examined EMT, cell proliferation, cell viability, cell cycle progression, and colony formation.

LINC00176 overexpression enhanced the epithelial phenotype and upregulated Δ40p53,

suggesting a possible feed-forward loop between Δ40p53 and LINC00176 through which

they regulate the other's levels. However, LINC00176 knockdown did not give consistent

results, as the expression of EMT markers also decreased upon LINC00176 knockdown. We

speculate that the decrease in the protein levels of EMT markers could be due to a decrease in

Δ40p53, which might also directly regulate EMT markers. Therefore, the effect of

LINC00176 knockdown is inconclusive. Δ40p53 function is known to be cell context-

specific, and Δ40p53 can have differential effects when overexpressed or silenced (35). This

strengthens our hypothesis that, since there is a feed-forward loop between LINC00176 and

Δ40p53, many of the lncRNA effects will be dependent on the levels of Δ40p53 and therefore

will vary. Furthermore, we observed a decrease in cell viability and colony formation and a

slight decrease in the proportion of cells in S phase with LINC00176 overexpression.

In conclusion, our results indicated that LINC00176 contributes to the tumor-suppressive function of Δ40p53, independent of p53FL function **(Graphical Abstract)**. However, it would be interesting to determine whether the function of the Δ40p53-LINC00176 axis is universal or cell type-specific. Taken together, this study opens up avenues for mechanistic insights into how Δ40p53 governs cellular processes through lncRNAs.

ACKNOWLEDGEMENTS

We are grateful to Prof. J.C. Bourdon, University of Dundee, UK and Dr. Robin Fahraeus, INSERM, France, for providing us the anti-p53 antibody. We would like to thank Dr. Je-Hyun Yoon and Dr. Myriam Gorospe, National Institutes of Health, USA, for providing us with the TRAP constructs. We are grateful to Dr. T Tamura, Institut fuer Biochemie, Germany, for providing us with LINC00176 overexpression construct and siRNA. We would like to thank Prof. Sathees C. Raghavan for his valuable suggestions. We also thank the present and past SD lab-members for helpful discussion of the work. This work is supported by a research grant to SD from the Department of Biotechnology (DBT), Govt. of India. SD also acknowledges J.C. Bose fellowship for research support. This study was also supported by the DBT-IISc partnership program. DST Fund for Improvement of Science and Technology Infrastructure (FIST) level II infrastructure and University Grants Commission Centre of Advanced Studies support is acknowledged. We would like to thank Editage (www.editage.com) for English language editing.

AUTHOR CONTRIBUTIONS

AP and SD: Conception and design of studies, analysis and interpretation, and writing the article. AP for performing the experiments.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCE

1.   Somasundaram, K. (2000) Tumor suppressor p53: regulation and function. *Front Biosci*, **5**,
D424-437.
2.   Bourdon, J.C., Fernandes, K., Murray-Zmijewski, F., Liu, G., Diot, A., Xirodimas, D.P., Saville,
M.K. and Lane, D.P. (2005) p53 isoforms can regulate p53 transcriptional activity. *Genes Dev*,
19, 2122-2137.
3.   Ray, P.S., Grover, R. and Das, S. (2006) Two internal ribosome entry sites mediate the
translation of p53 isoforms. *EMBO Rep*, **7**, 404-410.
4.   Grover, R., Ray, P.S. and Das, S. (2008) Polypyrimidine tract binding protein regulates IRES-
mediated translation of p53 isoforms. *Cell Cycle*, **7**, 2189-2198.
5.   Khan, D., Katoch, A., Das, A., Sharathchandra, A., Lal, R., Roy, P., Das, S., Chattopadhyay, S.
and Das, S. (2015) Reversible induction of translational isoforms of p53 in glucose
deprivation. *Cell Death Differ*, **22**, 1203-1218.
6.   Sharathchandra, A., Lal, R., Khan, D. and Das, S. (2012) Annexin A2 and PSF proteins interact
with p53 IRES and regulate translation of p53 mRNA. *RNA Biol*, **9**, 1429-1439.
7.   Katoch, A., George, B., Iyyappan, A., Khan, D. and Das, S. (2017) Interplay between PTB and
miR-1285 at the p53 3'UTR modulates the levels of p53 and its isoform Delta40p53alpha.
Nucleic Acids Res, **45**, 10206-10217.
8.   Ghosh, A., Stewart, D. and Matlashewski, G. (2004) Regulation of human p53 activity and cell
localization by alternative splicing. *Mol Cell Biol*, **24**, 7987-7997.
9.   Candeias, M.M., Malbert-Colas, L., Powell, D.J., Daskalogianni, C., Maslon, M.M., Naski, N.,
Bourougaa, K., Calvo, F. and Fahraeus, R. (2008) P53 mRNA controls p53 activity by
managing Mdm2 functions. *Nat Cell Biol*, **10**, 1098-1105.
10. Powell, D.J., Hrstka, R., Candeias, M., Bourougaa, K., Vojtesek, B. and Fahraeus, R. (2008)
Stress-dependent changes in the properties of p53 complexes by the alternative translation
product p53/47. *Cell Cycle*, **7**, 950-959.
11. Ungewitter, E. and Scrable, H. (2010) Delta40p53 controls the switch from pluripotency to
differentiation by regulating IGF signaling in ESCs. *Genes Dev*, **24**, 2408-2419.
12. Hinault, C., Kawamori, D., Liew, C.W., Maier, B., Hu, J., Keller, S.R., Mirmira, R.G., Scrable, H.
and Kulkarni, R.N. (2011) Delta40 Isoform of p53 controls beta-cell proliferation and glucose
homeostasis in mice. *Diabetes*, **60**, 1210-1222.
13. Ohki, R., Kawase, T., Ohta, T., Ichikawa, H. and Taya, Y. (2007) Dissecting functional roles of
p53 N-terminal transactivation domains by microarray expression analysis. *Cancer Sci*, **98**,
189-200.
14. Maier, B., Gluba, W., Bernier, B., Turner, T., Mohammad, K., Guise, T., Sutherland, A.,
Thorner, M. and Scrable, H. (2004) Modulation of mammalian life span by the short isoform
of p53. *Genes Dev*, **18**, 306-319.
15. Phang, B.H., Othman, R., Bougeard, G., Chia, R.H., Frebourg, T., Tang, C.L., Cheah, P.Y. and
Sabapathy, K. (2015) Amino-terminal p53 mutations lead to expression of apoptosis
proficient p47 and prognosticate better survival, but predispose to tumorigenesis. *Proc Natl*
Acad Sci U S A, **112**, E6349-6358.
16. Ota, A., Nakao, H., Sawada, Y., Karnan, S., Wahiduzzaman, M., Inoue, T., Kobayashi, Y.,
Yamamoto, T., Ishii, N., Ohashi, T. *et al.* (2017) Delta40p53alpha suppresses tumor cell
proliferation and induces cellular senescence in hepatocellular carcinoma cells. *J Cell Sci*,
130, 614-625.
17. Katoch, A., Tripathi, S.K., Pal, A. and Das, S. (2021) Regulation of miR-186-YY1 axis by the p53
translational isoform 40p53: implications in cell proliferation. *Cell Cycle*, **20**, 561-574.
18. Hunten, S., Kaller, M., Drepper, F., Oeljeklaus, S., Bonfert, T., Erhard, F., Dueck, A., Eichner,
N., Friedel, C.C., Meister, G. *et al.* (2015) p53-Regulated Networks of Protein, mRNA, miRNA,
and lncRNA Expression Revealed by Integrated Pulsed Stable Isotope Labeling With Amino
Acids in Cell Culture (pSILAC) and Next Generation Sequencing (NGS) Analyses. *Mol Cell*
Proteomics, **14**, 2609-2629.

19. Paralkar, V.R. and Weiss, M.J. (2013) Long noncoding RNAs in biology and hematopoiesis.
Blood, **121**, 4842-4846.

20. Knoll, M., Lodish, H.F. and Sun, L. (2015) Long non-coding RNAs as regulators of the
endocrine system. *Nat Rev Endocrinol*, **11**, 151-160.

21. Chaudhary, R. and Lal, A. (2017) Long noncoding RNAs in the p53 network. *Wiley Interdiscip*
Rev RNA, **8**.

22. Tran, D.D.H., Kessler, C., Niehus, S.E., Mahnkopf, M., Koch, A. and Tamura, T. (2018) Myc
target gene, long intergenic noncoding RNA, Linc00176 in hepatocellular carcinoma
regulates cell cycle and cell survival by titrating tumor suppressor microRNAs. *Oncogene*, **37**,
75-85.

23. Dai, L., Niu, J. and Feng, Y. (2020) Knockdown of long non-coding RNA LINC00176 suppresses
ovarian cancer progression by BCL3-mediated down-regulation of ceruloplasmin. *J Cell Mol*
Med, **24**, 202-213.

24. Zhu, S., Li, W., Liu, J., Chen, C.H., Liao, Q., Xu, P., Xu, H., Xiao, T., Cao, Z., Peng, J. *et al.* (2016)
Genome-scale deletion screening of human long non-coding RNAs using a paired-guide RNA
CRISPR-Cas9 library. *Nat Biotechnol*, **34**, 1279-1286.

25. Tang, Z., Li, C., Kang, B., Gao, G., Li, C. and Zhang, Z. (2017) GEPIA: a web server for cancer
and normal gene expression profiling and interactive analyses. *Nucleic Acids Res*, **45**, W98-
W102.

26. Li, J.H., Liu, S., Zhou, H., Qu, L.H. and Yang, J.H. (2014) starBase v2.0: decoding miRNA-
ceRNA, miRNA-ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data.
Nucleic Acids Res, **42**, D92-97.

27. Yang, Z., Wu, L., Wang, A., Tang, W., Zhao, Y., Zhao, H. and Teschendorff, A.E. (2017)
dbDEMC 2.0: updated database of differentially expressed miRNAs in human cancers.
Nucleic Acids Res, **45**, D812-D818.

28. Yoon, J.H. and Gorospe, M. (2016) Identification of mRNA-Interacting Factors by MS2-TRAP
(MS2-Tagged RNA Affinity Purification). *Methods Mol Biol*, **1421**, 15-22.

29. Yoon, J.H., Srikantan, S. and Gorospe, M. (2012) MS2-TRAP (MS2-tagged RNA affinity
purification): tagging RNA to identify associated miRNAs. *Methods*, **58**, 81-87.

30. Zhu, J., Liu, S., Ye, F., Shen, Y., Tie, Y., Zhu, J., Wei, L., Jin, Y., Fu, H., Wu, Y. *et al.* (2015) Long
Noncoding RNA MEG3 Interacts with p53 Protein and Regulates Partial p53 Target Genes in
Hepatoma Cells. *PLoS One*, **10**, e0139790.

31. Yoon, J.H., Abdelmohsen, K. and Gorospe, M. (2014) Functional interactions among
microRNAs and long noncoding RNAs. *Semin Cell Dev Biol*, **34**, 9-14.

32. Statello, L., Guo, C.J., Chen, L.L. and Huarte, M. (2021) Gene regulation by long non-coding
RNAs and its biological functions. *Nat Rev Mol Cell Biol*, **22**, 96-118.

33. Lin, T., Hou, P.F., Meng, S., Chen, F., Jiang, T., Li, M.L., Shi, M.L., Liu, J.J., Zheng, J.N. and Bai, J.
(2019) Emerging Roles of p53 Related lncRNAs in Cancer Progression: A Systematic Review.
Int J Biol Sci, **15**, 1287-1298.

34. Goldstein, I., Marcel, V., Olivier, M., Oren, M., Rotter, V. and Hainaut, P. (2011)
Understanding wild-type and mutant p53 activities in human cancer: new landmarks on the
way to targeted therapies. *Cancer Gene Ther*, **18**, 2-11.

35. Zhang, X., Groen, K., Morten, B.C., Steffens Reinhardt, L., Campbell, H.G., Braithwaite, A.W.,
Bourdon, J.C. and Avery-Kiejda, K.A. (2022) Effect of p53 and its N-terminally truncated
isoform, Delta40p53, on breast cancer migration and invasion. *Mol Oncol*, **16**, 447-465.

36. Vieler, M. and Sanyal, S. (2018) p53 Isoforms and Their Implications in Cancer. *Cancers*
(Basel), **10**.

37. Steffens Reinhardt, L., Zhang, X., Wawruszak, A., Groen, K., De Iuliis, G.N. and Avery-Kiejda,
K.A. (2020) Good Cop, Bad Cop: Defining the Roles of Delta40p53 in Cancer and Aging.
Cancers (Basel), **12**.

38. Courtois, S., Verhaegh, G., North, S., Luciani, M.G., Lassus, P., Hibner, U., Oren, M. and
Hainaut, P. (2002) DeltaN-p53, a natural isoform of p53 lacking the first transactivation
domain, counteracts growth suppression by wild-type p53. *Oncogene*, **21**, 6722-6728.

39. Yin, Y., Stephen, C.W., Luciani, M.G. and Fahraeus, R. (2002) p53 Stability and activity is
regulated by Mdm2-mediated induction of alternative p53 translation products. *Nat Cell*
Biol, **4**, 462-467.
40. Sun, Y., Manceau, A., Frydman, L., Cappuccio, L., Neves, D., Basso, V., Wang, H., Fombonne,
J., Maisse, C., Mehlen, P. *et al.* (2021) Delta40p53 isoform up-regulates netrin-1/UNC5B
expression and potentiates netrin-1 pro-oncogenic activity. *Proc Natl Acad Sci U S A*, **118**.
41. Hanahan, D. and Weinberg, R.A. (2011) Hallmarks of cancer: the next generation. *Cell*, **144**,
646-674.

FIGURE LEGENDS

Figure 1. Selection and validation of LINC00176 for further studies

(A) Quantitative PCR for validation of LINC00176 in H1299 cells expressing control, p53 only, Δ40p53 only and 14A construct. (B) Western blot analysis of cell extracts from H1299 cells expressing control, p53 only, Δ40p53 only and 14A construct, probed with CM1 after 48 h. (C) Quantitative PCR for validations of LINC00176 in HCT116+/+ and HCT116-/- cells. (D) Western blot analysis of cell extracts from HCT116+/+ and HCT116-/-cells, probed with CM1. (E) Quantitative PCR of LINC00176 in HCT116+/+ cells transfected with si p53 (30nM) and non-specific si (Nsp si). (F) Western blot analysis of cell extracts from HCT116+/+ cells transfected with either si p53 (30nM) and non-specific si (Nsp si), probed with CM1. (G) Quantitative PCR of LINC00176 in HCT116-/- cells transfected with si Δ40p53 (30nM) and non-specific si (Nsp si). (H) Western blot analysis of cell extracts from HCT116-/- cells transfected with siΔ40p53 (30nM) and non-specific si (Nsp si), probed with CM1. (I) Quantitative PCR of LINC00176 in HCT116-/- cells transfected with control and p53 only. (J) Western blot analysis of cell extracts from HCT116-/- cells transfected with control and p53 only, probed with CM1.

Figure 2. Regulation of LINC00176 under different stress conditions

(A) Schematic of different stress induction in HCT116-/- cells. (B) Quantitative PCR of LINC00176 in HCT116-/- treated with Doxorubicin (2μM) for 16 h. (C) Western blot analysis of cell extracts from HCT116-/- treated with Doxorubicin (2μM) for 16 h, probed with CM1. (D) Quantitative PCR of LINC00176 in HCT116-/- treated with Thapsigargin (0.1μM and 1μM) for 16 h. (E) Western blot analysis of cell extracts from HCT116-/- treated with Thapsigargin (0.1μM and 1μM) for 16 h, probed with CM1. (F) Quantitative PCR of LINC00176 in HCT116-/- deprived of Glucose in medium for 8 h. (G) Western blot analysis of cell extracts from HCT116-/- deprived of Glucose in medium for 8 h, probed with CM1. (H) Schematic of experiment in HCT116-/- transfected with siΔ40p53 followed by Thapsigargin (0.1μM) treatment. (I) Quantitative PCR of LINC00176 in HCT116-/- transfected with siΔ40p53 followed by Thapsigargin (0.1μM) treatment. (J) Western blot analysis of cell extracts from HCT116-/- transfected with siΔ40p53 followed by Thapsigargin (0.1μM) treatment, probed with CM1. (K) Western blot analysis of cell extracts from HCT116-/- transfected with siΔ40p53 followed by Thapsigargin (0.1μM) treatment, probed with Bip Antibody.

Figure 3. Mechanistic basis of LINC00176 regulation by Δ40p53/p53

(A) Schematic of possible routes of LINC00176 regulation. (B) Quantitative PCR of
LINC00176 promoters in HCT116-/- cells processed for ChIP Assay 48 h post seeding.
Δ40p53 was pulled down with 1801 Ab (Santa Cruz) and associated LINC00176 promoters
were probed using qPCR. (C) Quantitative PCR of LINC00176 promoters in HCT116+/+
cells processed for ChIP Assay 48 h post seeding. p53 was pulled down with 1801 Ab (Santa
Cruz) and associated LINC00176 promoters were probed using qPCR. (D) Quantitative PCR
of LINC00176 promoters in HCT116-/- cells transfected with p53 only and control, which
has been processed for ChIP Assay 48 h post transfection. Δ40p53/p53 was pulled down with
1801 Ab (Santa Cruz) and associated LINC00176 promoters were probed using qPCR. (E)
Western blot analysis of cell extracts from HCT116-/- transfected with p53 only and control,
probed with CM1. (F) Schematic of experiment to determine Δ40p53 mediated regulation on
LINC00176 RNA stability. (G) Quantitative PCR of LINC00176 in HCT116-/- transfected
with siΔ40p53 followed by Actinomycin D treatment for 0 h, 4 h and 8 h. (H) Western blot
analysis of cell extracts from HCT116-/- transfected with siΔ40p53 followed by Actinomycin
D treatment, probed with CM1.

Figure 4. Deciphering the mechanism of action of LINC00176

(A) Schematic for the probable downstream regulation by LINC00176. (B) Table for the
selected miRNA targets of LINC00176 obtained from Starbase V2.0, as mentioned in
methods section. (C) Quantitative PCR of selected miRNAs in HCT116+/+ and HCT116-/-
cells. (D) Quantitative PCR of LINC00176 in HCT116-/- transfected with siRNA (30nM)
directed to LINC00176 and non-specific si (Nsp si). (E) Quantitative PCR of selected
miRNAs in HCT116-/-cells transfected with siRNA (30nM) directed to LINC00176 and non-
specific si (Nsp si). (F) Quantitative PCR of selected miRNAs in GST pulldown lysates of
HCT116-/-cells (co-transfected with LINC00176 overexpression construct and pMS2-GST
construct) normalised to Input values. (G) Western blot analysis of cell extracts from Input
and Pulldown fraction of HCT116-/- cells (co-transfected with LINC00176 overexpression
construct and pMS2-GST construct), probed with CM1 and GST Antibody.

Figure 5: The impact of Δ40p53-LINC00176 axis on cellular processes

(A)Schematic of the different processes that has been checked for any probable effect by
LINC00176 over-expression or partial silencing. (B) Quantitative PCR of LINC00176 in
HCT116-/- cells transfected with LINC00176 overexpression construct. (C) Quantitative

PCR of LINC00176 in HCT116-/- cells transfected with LINC00176 shRNA construct. (D)

Quantitative PCR of EMT markers (E-cadherin, Vimentin, Slug) in HCT116-/- cells

transfected with LINC00176 overexpression construct. (E) Western blot analysis of cell

extracts from HCT116-/-cells transfected with LINC00176 overexpression construct, probed

with E-cadherin Antibody, Slug Antibody and CM1. (F) Quantitative PCR of EMT markers

(E-cadherin, Vimentin, Slug) in HCT116-/- cells transfected with LINC00176 shRNA

construct. (G) HCT116-/- cells were transfected with LINC00176 overexpression construct,

at 36 h post transfection cells were reseeded on 96 well plate for MTT Assay. MTT

incorporation was calculated at Day 0, Day 2, Day 4 and Day 6 post seeding (Data

represented for Set 1). (H) HCT116-/- cells were transfected with LINC00176 shRNA

construct, at 36 h post transfection cells were reseeded on 96 well plate for MTT Assay. MTT

incorporation was calculated at Day 0, Day 2, Day 4 and Day 6 post seeding (Data

represented for Set 1). (I) HCT116-/- cells were transfected with 30nM si-LINC00176, 36 h

post transfection cells were reseeded for BrdU assay. After 12 h of seeding BrdU was added

to the cells and incubated for 12 h. Following which, percentage of cell proliferation was

checked by calculating BrdU incorporation.

Graphical Abstract: Model depicting the regulation of Δ40p53-LINC00176-miRNA

axis

(A) In absence of Δ40p53, LINC00176 levels do not increase which leads to increased pool

of free miRNAs which probably binds to several mRNA targets preventing their function,

ultimately resulting in increased cell proliferation. (B) However, when Δ40p53 is present it

upregulates LINC00176 levels both at the level of transcription and post-transcription, which

sequesters oncogenic miRNAs preventing it to bind to their respective mRNA targets, which

ultimately results in decreased cell proliferation.

Figure: 1

Figure 1. Selection and validation of LINC00176 for further studies

Figure: 2

Figure 2. Regulation of LINC00176 under different stress conditions

Figure: 3

Figure 3. Mechanistic basis of LINC00176 regulation by Δ40p53/p53

Figure: 4

Figure 4. Deciphering the mechanism of action of LINC00176

Figure: 5

Figure 5: The impact of Δ40p53-LINC00176 axis on cellular processes

Graphical Abstract

A. Δ40p53 Absent

Transcription

LINC00176 RNA

Processing

Mature LINC00176 RNA

Stability

miRNA pool

Tumor Suppressor mRNAs

Translation

Tumor Suppressor Proteins

Increased Cell Proliferation

B. Δ40p53 Present

Transcription

LINC00176 RNA

Processing

Mature LINC00176 RNA

Stability

Stable LINC00176 RNA

miRNA pool

Tumor Suppressor mRNAs

Translation

Tumor Suppressor Proteins

Decreased Cell Proliferation

Graphical Abstract: Model depicting the regulation of Δ40p53-LINC00176-miRNA axis