Local field potentials and single unit dynamics in motor cortex of unconstrained macaques during different behavioral states

Richy Yun[1,4], Irene Rembado[2,6], Steve I. Perlmutter[2,4,5], Rajesh P. N. Rao[3,4], Eberhard E. Fetz[1,2,4,5]

Departments of [1]Bioengineering, [2]Physiology and Biophysics, [3]Allen School for Computer Science and Engineering, University of Washington
[4]Center for Neurotechnology, University of Washington
[5]Washington National Primate Research Center, University of Washington
[6]currently at the Allen Institute for Brain Science

Abstract

Different sleep states have been shown to be vital for a variety of brain function, including learning, memory, and skill consolidation. However, our understanding of neural dynamics during sleep and the role of prominent LFP frequency bands remain incomplete. To elucidate such dynamics and changes between different behavioral states we collected multichannel LFP and spike data in primary motor cortex of unconstrained macaques for up to 24 hours using the Neurochip3. Each 8 second bin of time was classified into awake and moving (Move), awake and at rest (Rest), REM sleep (REM), or non-REM sleep (NREM) by using dimensionality reduction and clustering on the average spectral density and the acceleration of the head. LFP power showed high delta during NREM, high theta during REM, and high beta when the animal was awake. Cross-frequency phase-amplitude coupling typically showed higher coupling for deeper sleep between all pairs of frequency bands. Two notable exceptions were high delta-high gamma and theta-high gamma coupling during Move, and high theta-beta coupling during REM. Sorted single units showed decreased firing rate with deeper sleep, though with higher "bursty" patterns during NREM compared to other states. Spike-LFP synchrony showed high delta synchrony during Move, and higher coupling with all other frequency bands with deeper sleep.

26   These results altogether are consistent with previous findings showing reactivation of cortical

27   circuitry during sleep, which may be moderated by delta band LFP.

Introduction

The discovery of rapid-eye movement (REM) sleep by Aserinski and Kleitman showed the presence of active brain processes during sleep, which catalyzed studies on different sleep stages (Aserinsky & Kleitman, 1953). Findings have shown REM sleep to play a role in regulating the autonomic nervous system activity and to be linked with spatial and emotional memory consolidation (Boyce et al., 2016; Rechtschaffen, 1998; Siegel, 2005). Similarly, non-REM (NREM) sleep has been shown to be tied to skill consolidation and memory formation (Ramanathan et al., 2015; Siclari & Tononi, 2017).

Studies have furthered our understanding of different sleep stages by exploring the dynamics of local field potential (LFP) frequency band power and single-unit spiking characteristics in both the cortex and various deep brain structures during sleep. Slow waves and delta frequency band are typically present across the brain during NREM sleep, and high theta power in the hippocampus is present during REM sleep (Rechtschaffen & Kales, 1968; Silber et al., 2007). Single units show changes in firing rate as well as firing patterns during sleep, due to changes in connectivity and excitability, potentially reflecting reactivations of relevant cortical circuit patterns to solidify learning and memory (Arbune et al., 2020; Ramanathan et al., 2015; Tononi & Cirelli, 2014; Xu et al., 2019). Several other features of cortical activity, such as k-complexes and sleep spindles during NREM sleep, have also been identified to be associated with different stages of sleep (Fernandez & Lüthi, 2020; Huber et al., 2004; Ulrich, 2016).

Additional findings have shown that certain plasticity mechanisms, such as long-term potentiation/depression and synaptic tagging and capture, occur exclusively during sleep (Kim et al., 2005; Vecsey et al., 2018). Subsequent research has applied this knowledge by enhancing slow waves prevalent during NREM sleep (Bellesi et al., 2014; Marshall et al., 2006) or

delivering stimulation locked to delta to augment learning and memory (Rembado et al., 2017).

However, despite our expanding tools and techniques to take advantage of sleep mechanisms, the

neural dynamics underlying sleep remain unclear.

Various measures of local field potential (LFP) coupling and spike-LFP synchrony have been

commonly used to elucidate mechanisms of neural dynamics. Cross-frequency phase-amplitude

coupling of different LFP frequency bands, thought to reflect communication between brain

circuitry, has been shown to be modulated by task performance, cognitive engagement, and

memory formation in various brain regions (Canolty & Knight, 2010; Jensen & Colgin, 2007).

Single units are strongly synchronized to specific phases of LFP frequency bands, notably to beta

and gamma cycles in the motor cortex, suggesting beta to be a resting rhythm during movement

execution and gamma to reflect local population activity (Buzsáki et al., 2012; Buzsáki & Wang,

2012; Engel & Fries, 2010; Murthy & Fetz, 1996a, 1996b). LFP and unit analyses have given us

significant insight into neural signaling and the role of both spikes and LFPs in brain function,

but there has yet to be a comprehensive analysis of cross-frequency coupling and spike-LFP

synchrony for specific cortical sites during sleep and wake.

This study analyzes LFP and single unit dynamics in the macaque motor cortex during different

behavioral states to better understand the mechanisms operating during different sleep stages as

well as the roles of various LFP frequency bands. We first used the power spectral density of

LFPs to distinguish between four major behavioral states shown to be relevant to plasticity and

learning in the motor cortex: 1. awake and moving (Move), 2. awake and at rest (Rest), 3. rapid-

eye movement (REM) sleep, and 4. non-REM (NREM) sleep. We tracked single-unit activity

concurrently with LFP recordings and focused on 6 different frequency bands commonly

observed in the cortex – delta, theta, alpha, beta, low gamma, and high gamma. Finally, we

assessed state-dependent changes in cross-frequency coupling between pairs of the frequency

bands as well as spike-LFP synchrony in each frequency band.

Materials and Methods

Experimental design

Implants and surgical procedures

The experiments were conducted on two male *Macaca nemestrina* monkeys (Monkeys J and K). Surgeries were performed under isoflurane anesthesia and aseptic conditions to implant multi-electrode Utah arrays. All procedures conformed to the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* and were approved by the University of Washington Institutional Animal Care and Use Committee.

Implantation of the Utah array was guided by stereotaxic coordinates. A 1.5 cm wide square craniotomy was performed over the hand region of the primary motor cortex to expose the dura. Three sides of the exposed dura were cut to expose the cortex; a Utah array was implanted and the dura was sutured over the array. Two reference wires were inserted below the dura and two were inserted between the dura and the skull. The bone flap from the craniotomy was replaced and held in place by a titanium strap screwed onto the skull with 2.5 mm x 6 mm titanium skull screws. A second smaller titanium strap was fastened to the skull to secure the wire bundle. The connector pedestal for the array was attached to the skull with eight titanium skull screws and the incision closed around the pedestal base. Additional skull screws were placed around the base of the connector pedestal and a thin coat of dental acrylic (methyl methacrylate) was applied to the skull between the screws and the connector base for additional stability.

The animals also received a "halo" implant made with 3/8" aluminum forming an egg-shaped oval 17 cm long and 15.3 cm wide. Four titanium straps were affixed to the skull by titanium bone screws. Two were implanted bilaterally over the occipital ridge, and two were placed

temporally bilaterally. After the plates integrated with the skull for 6 weeks, an aluminum halo

was mounted on four pins seated in each plate. A titanium can with a plexiglass top to house the

Neurochip3 was attached to the halo during all Neurochip3 recording sessions.

Monkey J additionally received EOG implants on the lateral wall and superior margin of both

orbits. EOG electrodes consisted of a titanium washer ((#4, 0.25" OD, 0.032" thick) with a

0.016" hole drilled into it and a 34-gauge silver-plated copper microwire with silicone shielding

(Cooner Wire, AS155-34) threaded through the hole and soldered to the washer. An incision was

made to the dorsal and lateral margins of both orbits to expose the skull. A small hole was drilled

in each incision, a titanium skull screw (2 mm x 6 mm) was used to hold down the EOG

electrodes, and the wires of the electrodes were tunneled beneath the tissue along the skull.

Another incision was made along the top of the skull to secure a pedestal with 8 skull screws and

allow access to the electrode wires. The inside of the pedestal was filled with surgical silicone

adhesive (Kwik-Sil, World Precision Instruments) to prevent infection. Connectors were fastened

to the wires postoperatively.

After each surgery animals received postoperative courses of analgesics and antibiotics. Animals

did not show signs of discomfort or pain related to any implanted devices after recovery, and all

exposed implants were regularly disinfected biweekly with chlorohexidine and treated with

antibiotics to prevent infection.

Electrophysiology

All data was collected with the head-mounted Neurochip3 (Shupe et al., 2021) while the

monkeys were freely behaving in their home cage (Figure 1A). The Neurochip3 is a battery

powered bidirectional brain-computer interface capable of saving data to an SD card, allowing

for wireless recording and stimulation for up to 24 hours. 16 channels of the Utah array, or 14

channels of the array and 2 channels of the EOG, were recorded at 20 kHz sampling rate with a

bandwidth of 0.1 Hz to 5 kHz (Figure 1B). The cortical channels were chosen each day with a

preliminary recording to capture the largest single units present in the array. Most experiments

recorded a different set of channels, depending on best spike recordings.

The Neurochip3 also has an onboard 3-axis accelerometer that was simultaneously recorded at

100 samples per second. Recording sessions lasted between 19 and 24 hours. The lights in the

animal rooms were off during the night for the 12 hours between 6pm and 6am. We collected

data from a total of 16 sessions from Monkey K and 10 sessions from Monkey J.

<u>Data analysis</u>

Classifying sleep states

All analyses were performed using custom MATLAB (Mathworks) and Python code. The power

spectral density (PSD) of cortical local field potentials (LFPs) as well as the onboard

accelerometer data from the Neurochip3 were used to classify different sleep states (Figure 1B).

Data was down sampled to 1 kHz before performing Welch's PSD estimate between 0 and 50 Hz

for each 8 second time-bin. We then converted the PSD into power and found the average power

across all channels for each time-bin. The average power was further normalized by subtracting

the minimum value and dividing by the integral to ensure that relative power at different

frequencies played a larger role than the absolute power.

We then used the normalized average power as inputs to train a stacked sparse autoencoder for

dimensionality reduction, similar to the architecture described in Tsinalis et al. 2016 (Tsinalis et

al., 2016). The encoder was composed of 3 layers of 256, 128, and 64 units each, with batch

normalization and ReLU activation function. The final hidden layer containing the reduced

representation of the data had 32 nodes. The autoencoder was trained with minibatch sizes of 64

for 300 epochs. The loss was calculated with mean squared error with L1 regularization

(regularization weight $\lambda = 1e^{-5}$) using the Adam optimizer (learning rate $\alpha = 1e^{-3}$, decay rate

for first moment $\beta_1 = 0.9$, decay rate for second moment $\beta_2 = 0.999$, constant $\varepsilon = 1e^{-8}$). The

autoencoder was implemented in Python using the PyTorch package (Paszke et al., 2019).

Accelerometer data was included by performing the root sum of squares across all three axes.

We then found the variance of the values within each time bin and applied a logarithmic scale to

better compress the data. Finally, the standard deviation was normalized to that of the encoded

dimension with the largest variance. The processed accelerometer data was included as an

additional dimension in the lower dimensional representation (i.e., as the 33rd dimension).

To classify the data, we used k-means clustering with an assumption of 4 centroids. Data points

of the lower dimensional representation within the 90th percentile of pairwise Euclidean distance

were initially used for finding the centroids of clusters to avoid the influence of outliers. Each

data point was assigned to the centroid with the shortest Euclidean distance.

After clustering, each group of records was assigned to one of four states – 1) awake and moving

(Move) 2) awake and at rest (Rest), 3) rapid-eye movement (REM) sleep, 4) non-REM (NREM)

sleep – by assessing the average accelerometer value and average normalized PSD for each

cluster. First, the cluster with highest average acceleration was assigned to be Move, then the

cluster with highest average delta power (0.5 to 4 Hz) was assigned be NREM, then the cluster

with higher average beta power (15 to 30 Hz) was assigned to be Rest, and the remaining cluster

was assigned to be REM.

To include a temporal aspect and smooth any outliers we performed a majority filter on the

classification. For each time-bin, the majority state across 2 time-bins before to 2 time-bins after

(± 16 seconds) was considered to be the current state. Ties were resolved by keeping the original

classification, or by random choice if the original classification was not part of the tie.

Validation of classification

Two EOG electrodes (one dorsal and one lateral) were simultaneously recorded from with the

Neurochip3 in experiments with Monkey J. EOG signals were extracted by subtracting the lateral

electrode signal from the dorsal electrode signal and then applying a band-pass filter between 1

and 20 Hz with a zero-phase second-order Butterworth filter. We performed in-booth

experiments with flashes of lights guiding the monkey's gaze to ensure we were properly

capturing eye movements.

As the Neurochip3 is mounted to the monkeys' heads, we performed overnight recordings with

the Microsoft Kinect in conjunction to ensure there were no large discrepancies between the on-

board accelerometer and whole-body movements (Libey & Fetz, 2017). Movements with the

Kinect was calculated as the absolute difference between each frame of the infrared depth-

finding camera and the previous frame. We captured frames as quickly as possible with the

processing overhead, around 30 frames per second.

Further validation was performed with k-fold cross-validation with k=20. Each dataset was split

into 20 random groups; classification was performed on 19 of the 20 groups, and the final group

was classified using the centroids from the classification k-means. The training error was

calculated as the difference in classification within the 19 groups used for training the

classification, and the test error as the difference in classification within the final group used to

test the classification. Cross-validation was bootstrapped 50 times for each session to minimize

variability that can potentially be introduced by the random sampling.

Coherence

Magnitude-squared coherence was used to calculate synchrony within the same frequencies:

$$C_{xy} = \frac{|P_{xy}(f)|^2}{P_{xx}(f)P_{yy}(f)} \tag{1}$$

where C_{xy} is the coherence between x and y, $P_{xy}(f)$ is the cross-spectral density between x and

y, and $P_{xx}(f)$ and $P_{yy}(f)$ are the spectral densities of x and y respectively. Coherence was

calculated between all combinations of channel pairs at every 0.1 Hz intervals.

Cross-frequency phase-amplitude coupling

To calculate cross-frequency phase-amplitude coupling we used mean vector length (MVL)

(Canolty et al., 2006) (Figure 1C). The LFP was first filtered into 2 frequency bands of interest

within the 6 frequency bands of interest – 1) delta (0.5-4 Hz), 2) theta (4-8 Hz), 3) alpha (8-12

Hz), 4) beta (15-30 Hz), 5) low gamma (30-70 Hz), and 5) high gamma (70-120 Hz) – using a

zero-phase second-order Butterworth filter. We then calculated the analytic signal, H, of each

band using the Hilbert transform:

$$H(x(t)) = \frac{1}{\pi} P \int_{-\infty}^{+\infty} \frac{x(\tau)}{t - \tau} d\tau \tag{2}$$

200   where P is Cauchy principal value. The phase of the complex valued analytic signal is the

201   instantaneous phase at time t, and the magnitude of the analytic signal is the instantaneous

202   amplitude. The mean vector was then calculated by:

$$MVL = \left| \frac{1}{n} \sum_n r \, exp \, (i\phi) \right| \quad\quad\quad (3)$$

where r is the instantaneous magnitude of the higher frequency band and ϕ is the instantaneous

phase of the lower frequency band. ϕ ranges from 0 to 2π radians where 0 is the peak and π the

trough of oscillations. This calculation creates a vector for each sample in time with the phase

and amplitude of the lower and higher frequency bands, respectively. The magnitude of the

average of those vectors, or MVL, measures the strength of synchrony – zero indicates a uniform

distribution in which the vectors "cancel" each other, and higher values indicate the degree of

synchrony.

However, MVL is highly affected by the amplitude and does not have a normalized maximum

value, which makes interpretation of individual values and comparisons of MVL measurements

across different time points difficult. Thus, we additionally calculated the maximum possible

MVL for each state. Instead of using the phase and amplitude that occurs at the same time

sample in Equation 2, we paired the highest amplitudes with the most commonly occurring

phases. Thus, the largest vectors were all in similar directions providing the maximum possible

length of the mean vector. We then normalized the MVL by dividing by the maximum possible

MVL to assess differences in synchrony between behavioral states.

Spike sorting

Spikes were sorted offline using two-window discrimination. The cortical recording was

bandpass filtered between 1000 and 2000 Hz with a first-order Butterworth filter. Then a

negative threshold and two time-delayed windows were manually chosen to capture the trough

and the peak of the spike waveform. All traces crossing the threshold and passing through the

two windows was denoted as a spike.

As the fidelity of spikes may change over such a long period of recording, we additionally

compared the shapes of the first 1000 detected spikes with the last 1000 detected spikes using the

coefficient of determination (CoD):

$$CoD = 1 - \frac{\sum \left(s_{first}(t) - s_{last}(t)\right)^2}{\sum \left(s_{first}(t) - \overline{s_{first}}\right)^2} \tag{4}$$

where s_{first} is a waveform of one of the first spikes, s_{last} is a waveform of one of the last spikes,

and $\overline{s_{first}}$ is the average of the waveform.

We performed pairwise CoD on all first and last 1000 instances of each spike and compared

them to the pairwise CoD between the first 1000 instances of the spike and the last 1000

instances of a different, randomly chosen spike. If the distribution of the CoD comparing the

same spike was significantly higher than the distribution between the spike and another spike it

was considered to be consistent overnight. Additionally, to ensure we were not capturing

multiple neurons with a similar waveform, we manually assessed the autocorrelograms for each

spike to ensure the presence of a refractory period and proper distribution of inter-spike intervals.

Phase-locking value

We calculated the phase-locking value (PLV) to assess the strength of synchronization of spike

timing to phases of oscillations in specific frequency bands from LFPs recorded on the same

channel (Figure 1D). First, the LFP was filtered into a frequency band of interest. The

instantaneous phase of the lower frequency band was calculated as described above. We then

calculated the PLV:

$$PLV = \frac{1}{n}\left|\sum_{n} exp\ (i\phi)\right| \tag{5}$$

where ϕ is the instantaneous phase at spike times. The PLV effectively converts each phase into

a unitary vector and finds the average of all the vectors. The magnitude of the resulting vector

determines the synchrony of the phases. A value of zero, similar to MVL, indicates a uniform

distribution, or no synchrony. A value of one indicates that all phases are equal, or perfect

synchrony.

The average phase of spike timing was found using the circular mean:

$$average\ phase = arg\left(\sum_{n} exp\ (i\phi)\right) \tag{6}$$

where arg is the argument of a complex number, or the angle between the positive real axis and

the complex number in the complex plane.

Statistical analysis

We used Friedman's test to compare values between the four states due to the non-parametric

nature of the distribution and because we were sampling the same spikes and LFPs across each

state. Tukey's honest significance test was used post hoc to determine significant pairwise

differences.

Results

<u>Classification of sleep states</u>

Although there have been various approaches automating sleep state classification by taking

advantage of the sequential nature of the states as suggested by the American Academy of Sleep

Medicine (AASM) classification (Craik et al., 2019; Silber et al., 2007; Supratak et al., 2017),

studies have shown that including information from neighboring epochs does not necessarily

improve classification (Sekkal et al., 2022; Tsinalis et al., 2016). Most deep learning methods

also rely on supervised learning, but, due to the high inter-scorer variability in manual

classification that these models rely on (Himanen & Hasan, 2000; Younes et al., 2016), we chose

an unsupervised method instead. As a result, we used dimensionality reduction for feature

extraction and subsequent clustering. Autoencoders were chosen as the method for

dimensionality reduction due to their nonlinearity potentially extracting more salient features

compared to linear methods. Denoising autoencoders are often used for feature extraction to

ensure the network does not learn to replicate consistent noise, but our input data inherently

contained random noise due to the recording device being mounted on a freely behaving animal

as well as the short time window of 8 seconds for our power spectral density calculations. As a

result, we used a stacked sparse autoencoder for dimensionality reduction.

Sleep states were classified using the power spectral density of local field potentials (LFPs) of

successive 8-second time-bins as the input to train an autoencoder. We then extracted the values

from the hidden units as the low-dimensional representation of the data. Figure 2A shows

examples of the original spectra and reconstructed outputs after training the autoencoder. The

fluctuations in the spectra have been smoothed out but the salient features, such as peaks at alpha

or beta, were maintained. Figure 2B shows the low-dimensional representation visualized via t-

distributed stochastic neighbor embedding (t-SNE) (van der Maaten & Hinton, 2008) after full

classification. There was some overlap between the states in the representation, which is to be

expected as brain states during sleep are on a continuous spectrum. Finally, we wanted to gain

insight into which feature each of the hidden units of the autoencoder represent. Figure 2C shows

the average of 100 spectra that gave the largest values in each hidden unit and Figure 2D shows

the average of 100 spectra that gave the smallest values in each hidden unit subtracted from

spectra in Figure 2C. This analysis can be interpreted as the features within the spectra that each

hidden unit encodes, and clear peaks in each frequency band can be observed.

We then applied k-means clustering on the lower dimensional representation as well as the

accelerometer data that was collected concurrently through the Neurochip3. We assumed 4

clusters – awake and moving (Move), awake and at rest (Rest), REM sleep (REM) and non-REM

sleep (NREM). Each cluster was assigned to the state depending on the average accelerometer

value and features in the average power spectral density (PSD). Although we have a

physiological basis as to why we chose 4 clusters, we wanted to ensure that 4 clusters was a

reasonable number for our dataset. To that end, we tracked the within-clusters sum-of-squares as

well as the average silhouette coefficient for 1 to 10 clusters (Figure 3A). Both values decrease

with more clusters, but the "elbow" point, or where there is a large reduction in the decrease, was

around 4 or 5 clusters. We additionally calculated the individual silhouette coefficients when

using 4 clusters to show that each cluster had a significant number of data points with

coefficients above the average value (Figure 3B).

Classified states show distinct features

The full classification procedure is outline in Figure 4A and an example of the final classification over a 21-hour period and the corresponding spectral power is shown in Figure 4B. During lights-off we found consistent REM cycles in between NREM epochs occurring every 30 minutes to 2 hours, which became longer and more frequent closer to the morning, consistent with previous findings (Aserinsky & Kleitman, 1953; Hsieh et al., 2008; Kripke et al., 1968). NREM was absent when the lights were on in the animal room, but we often found brief periods of REM sleep concurrent with changes in the PSD, which we attributed to naps (Figure 4B, arrow) (Dijk et al., 1987; Moses et al., 1975). We also often found short periods of Rest or sometimes even Move during the night which were attributed to brief awakenings, typical for a normal night of sleep.

Averaging the spectral power during each sleep state shows high delta power during NREM sleep and high beta power when the animal was awake, as we used these criteria to assign the classified clusters to the appropriate states, but also shows high theta power during REM sleep which was not explicitly a part of the classification scheme (Figure 4C). Additionally, raw LFP traces during the states show high-frequency activity during Move, faster oscillatory activity during Rest in the beta range, slower oscillatory activity during REM sleep in the theta and alpha range, and clear K-complexes and sleep spindles along with low frequency activity around delta during NREM sleep (Figure 4D). T

Validating sleep state classification

As our classification scheme was unsupervised, we simultaneously recorded electrooculography (EOG) signals to validate the identified sleep states (Figure 5A). Eye movements detected with the EOG were clearly associated with identified sleep states; EOG variance was very high when

the animal was awake, very low during NREM sleep, and elevated during REM sleep (Figure

5B). This relationship was present even during very brief windows of detected awakening during

the night and naps during the day, indicating high accuracy of state classification.

The accelerometer used for classification was on the Neurochip3 which was mounted on the

animals' head. To confirm the accuracy of detected movements, we additionally recorded the

monkeys' movements overnight with a Microsoft Kinect. Compared to the movement values

extracted with the Kinect, the accelerometer values were smaller when the animal was awake but

larger when the animal was asleep (Figure 5C). This is likely due to body movements that are

independent of the head (i.e., isolated limb movements) and are not as accurately recorded in the

accelerometer. On the other hand, when the animal is laying down at night most movements

involve the head. However, these differences were minor, and the two signals were comparable

throughout the recording duration.

Finally, we also performed k-fold cross-validation to verify that our classification method was

consistent and robust. We used k=20 and carried out 50 repetitions per session and found the

average test and training error to be less than 4%, or around 40 total minutes (Figure 5D). The

similar but low test and training errors suggest the classification method had low variance (i.e.

not overfit) and high consistency. As a result, the classification was deemed appropriate for the

aims of this study.

Changes in LFP dynamics

The average spectral power per state for each animal is shown in Figure 6A. Delta power was

high during NREM sleep due to the presence of slow waves. There was a beta peak during the

wake states, though the specific frequency range differed between the animals: 25-30 Hz for

Monkey K and 15-20 Hz for Monkey J. REM sleep showed slightly higher theta and alpha

power compared to the wake states.

Average pairwise coherence per state for each animal is shown in Figure 6B. The features were

very similar to those show in the power, but the differences were magnified, likely due to

coherence showing synchrony between pairs of channels and amplifying active oscillatory

signaling over baseline spectral density. In addition to the large delta peak during NREM, there

was a much clearer difference between REM and the wake states, including a more distinct beta

peak.

Cross-frequency phase-amplitude coupling

The phase of lower frequency bands has been observed to be coupled with the amplitude of

higher frequency bands, thought to reflect coordination between brain networks (Canolty &

Knight, 2010; Jensen & Colgin, 2007). We explored the coupling between every pair of low

frequency band phase to high frequency band amplitude to determine whether there were state-

dependent changes. We first plotted the average z-scored spectral power of higher frequencies at

different phases of each frequency band (Figure 7). There was clear coupling between different

phases and frequencies, the strength and specific phase of which changed, depending on the

state.

To quantify the degree of coupling we calculated the normalized mean vector length (nMVL).

Figure 8 shows the distribution of nMVL for each pair of frequency bands during each state. The

distribution of nMVL did have state-dependent changes, notably 1) high delta phase to alpha

amplitude (delta-alpha) coupling during Rest, likely due to increased alpha oscillatory power, 2)

low delta-beta coupling during NREM, perhaps for disinhibition of the cortex leading to

increased plasticity, 3) high delta-high gamma coupling during Move and NREM, potentially

reflecting synchrony of spike activity with delta oscillations, 4) high theta-beta coupling during

REM, showing a possible modulation of beta by deeper brain structures, 5) high theta-high

gamma coupling during Move, which may reflect hippocampal place cells synchronizing cortical

activity during behavior, and 6) high alpha-high gamma coupling during Move, which may be

due to the relevance of alpha to both attention and movement preparation.

To further characterize the coupling, we explored the distribution of the mean vectors to

determine whether the phases aligned with the highest amplitudes were consistent. Figure 9

shows the distribution of mean vector phases for each pair of frequency bands during each state.

The mean phases are consistent across most frequency pairs across all states with stronger

consistency typically in aforementioned significant nMVL values (e.g. delta-high gamma during

NREM or theta-beta during REM).

<u>Spike sorting</u>

Spikes were manually sorted using two-window discrimination and confirmed using the

coefficient of determination (Figures 10A and 10B, see *Materials and Methods – Spike sorting*).

We tracked a total of 193 spikes (121 in Monkey K, 92 in Monkey J) that satisfied our conditions

across all sessions. The firing rates of spikes overnight were strongly and consistently modulated

by different identified sleep states, especially during REM cycles during the night, further

demonstrating the accuracy of the classification method (Figure 10C). Firing rates were

consistently lower with deeper sleep.

Additionally, spikes were designated to be regular firing or fast firing. Although spike type

classification was traditionally performed with the spike width (McCormick et al., 1985),

subsequent studies have suggested that spike width, at least when used alone, may not be the

most indicative of the spike type (Insel & Barnes, 2015; Jung et al., 1998; Vigneswaran et al.,

2011). Assessment of various features of spikes showed that the peak of the inter-spike interval

distribution to be the most distinguishable feature within our dataset (Figure 11). If the peak

occurred before 10 ms the spike was designated to be fast spiking (FS), otherwise it was

designated to be regular spiking (RS) (84 FS spikes, 109 RS spikes). Assessing the differences in

firing rates in individual states showed that both RS and FS spike rates significantly decrease

with deeper sleep (Figure 10E).

Changes in spiking dynamics

To determine how spiking patterns changed between different sleep states we analyzed changes

in the average ISI distributions of RS and FS spikes. The raw ISI distributions showed an overall

decrease on most ISIs with deeper sleep, reflecting the diminished firing rate, but also a slight

change in the timing of the distribution, especially for RS spikes (Figure 12A).

To quantify the differences, we calculated the difference in ISI distributions from Move (Figure

12B, top). FS spikes had a consistent decrease in shorter latency ISIs with deeper sleep, but RS

spikes showed a large increase in short latency ISIs at around 5 ms (arrow). To determine if there

were any changes in larger ISIs, we plotted the distributions against frequency (Figure 12B,

bottom). Both FS and RS spikes showed a peak between 7 and 10 Hz during NREM and a peak

between 20 and 25 Hz during Rest and REM.

We additionally sought to determine how the brain state could affect the relationship between

neurons by calculating the cross correlation between firing rates of recorded spikes. Firing rates

were calculated by convolving the spike train with a gaussian kernel with an approximate width

of 10 ms. Figure 12C shows the cross correlations between smoothed firing rates of pairs of FS

neurons, between FS and RS neurons, and between pairs of RS neurons. There was not a clear

difference in the cross correlation between the four states suggesting relative circuitry is

maintained. The correlation between pairs of FS neurons was stronger than between FS and RS

or pairs of RS neurons, as inhibitory neurons are often more interconnected within the cortex.

Spike-field relationships

Spikes have been shown to be synchronized to various frequency bands, particularly in the motor

cortex (Buzsáki et al., 2012; Murthy & Fetz, 1996b). We first analyzed the phase-amplitude

distribution normalized by amplitude of each LFP band during spike timings (Figure 13). There

was clear synchrony with spikes during NREM sleep with all LFP bands, though at different

phases. During NREM spikes typically occurred at the negative phase of delta, beta, and gamma

activity, and at intermediate phases for theta ($\pi/2$) and alpha ($3\pi/4$). The synchrony was

particularly apparent during high amplitudes, suggesting spikes are synchronized to active

frequency band activity rather than epiphenomena arising from periodicity of spike firing

patterns. During both awake states and REM sleep spikes were synchronized to the trough of

beta and gamma frequencies, though less than during NREM sleep. Spikes during Move were

also potentially synchronized to the delta band. All other combinations did not show apparent

coordination between LFP band phase and spike timing.

To quantify the strength of these relationships, we calculated the phase-locking value (PLV) for

each spike during each state for all LFP bands. Figure 14 shows the RS spikes' distribution of

locked phases as well as the PLV for each frequency band during each state. Spikes were more

synchronized to delta during both Move and NREM, validated by the consistency in the locked

phases (around $\pi/2$ and $\pm\pi$, respectively). For every other frequency band, there was stronger

synchrony during deeper sleep states, though the differences were often not statistically

significant. The phases were extremely consistent for beta and higher frequency bands, but

variable for lower frequency bands during states with lower PLVs.

Figure 15 shows the FS spikes' distribution of locked phases and the PLV for each frequency

band during each state. One large difference from RS spikes was the synchronization with delta

during Move, which was not significantly larger compared to Rest or REM. Otherwise, there was

a much stronger relationship between increase in PLV and deeper sleep states, being statistically

significant for each frequency band. In addition, the phase of synchrony to the low and high

gamma bands was earlier in the phase for FS spikes compared to RS spikes.

Discussion

State classification

Although a guideline to standardize sleep staging was published after the discovery of sleep stages (Rechtschaffen & Kales, 1968), staging has remained an inconsistent and often arduous task. Criticisms of and modifications to the guideline (Himanen & Hasan, 2000; Silber et al., 2007) as well as various automated methods (Hamida & Ahmed, 2013; Penzel & Conradt, 2000; Sun et al., 2020) have been developed, but each approach differs in the set of defined states and the type and amount of data required. In addition, studies on sleep typically use electroencephalogram (EEG) LFP recordings, which are distinctly different from intracortical LFPs due to their lower spatial specificity and the complex frequency and phase filtering of bone and tissue, especially at higher frequencies (Buzsáki et al., 2012; Michel et al., 2004; Sejnowski et al., 2014). Beyond these difficulties, manual scoring is also compromised by high inter-scorer variability (Himanen & Hasan, 2000; Younes et al., 2016).

As a result, we developed our own classification methodology to distinguish between different sleep states. The process was relatively simple, utilizing basic dimensionality reduction and clustering on processed LFP PSDs and accelerometer values, but was remarkably consistent. We used several metrics to confirm the method – EOG recordings, concurrent recordings with the Kinect to track movement, k-fold cross validation, and changes in spike rates; they all indicated accurate and robust classification. One explanation may be that sleep states can vary between different regions of the brain (Mascetti, 2016; Siclari & Tononi, 2017). Our focus on a small region (4x4 mm) of the motor cortex and the high spatial specificity of intracortical LFPs may

have produced more robust results compared to those based on EEG recordings from across the

brain.

Our method could potentially be improved by using a dimensionality reduction method with

higher complexity, such as a deep neural network (Malafeev et al., 2018), different clustering

methods, such as hierarchical clustering (Gerla et al., 2019), and by including the EOG

recordings or other biophysical measurements as additional dimensions of data (Khalighi et al.,

2013). However, developing a flawless classification method was not within scope of the study

and the method presented was deemed to be sufficient.

LFP and spike dynamics during different sleep states

Delta band LFP, or slow waves, is typically elevated across the brain during all stages of NREM

(Rembado et al., 2017; Silber et al., 2007; Xu et al., 2019). Beta power is high when the animal

is awake, particularly in the primary motor cortex as beta oscillations are highly relevant in

motor control (Engel & Fries, 2010; Khanna & Carmena, 2017). These two metrics were used as

criteria for assigning the classified clusters to the appropriate states. However, the state

corresponding to REM sleep, as further confirmed through EOG recordings, presented high theta

power, a criterion we did not impose on the classification scheme. The role of theta band in the

cortex is still obscure, but theta in the hippocampus is also consistently high during REM (Boyce

et al., 2016). We found similar features in the pairwise magnitude-squared coherence; as

coherence shows synchrony between two signals, it effectively emphasizes that the increased

power is reflective of changes in coordinating activity.

We also observed decreased spiking activity with deeper sleep, similar to previous studies

(Steriade et al., 2001; Xu et al., 2019). In addition, the firing patterns of spikes changed due to

sleep state – RS neurons had increased high frequency activity during NREM compared to

Move, and all neurons had an increase low frequency activity during Rest, REM, and NREM

compared to Move. The increase in beta frequency activity of neurons during REM and Rest can

be explained by the fact that beta is seen as a "resting" rhythm in the primary motor cortex

(Engel & Fries, 2010). RS neurons are likely to be excitatory pyramidal cells and FS neurons are

likely to be inhibitory interneurons (McCormick et al., 1985), which may explain the higher

"bursty" firing pattern of RS neurons during NREM sleep, potentially due to reactivation of

cortical circuitry activated during the day.

Cross-frequency coupling and spike-LFP synchrony

Cross frequency coupling shows that the higher frequency band modulates with the lower

frequency band. Such synchrony reflects coordination of local networks operating on shorter

time scales to distributed circuits synchronized at longer time scales; this could potentially play a

role in neural computation of attention, learning, and memory (Canolty & Knight, 2010; Jensen

& Colgin, 2007). Various measures have been proposed to quantify phase-amplitude coupling;

compared to other proposed measures the MVL introduced by Canolty et al. (2006) has been

shown to be accurate, the most sensitive to modulations in coupling strengths, and ideal for high

quality signals over a long recording epochs (Canolty et al., 2006; Hülsemann et al., 2019;

Onslow et al., 2011; Tort et al., 2010).

Spike-LFP synchrony provides another measure of synchrony between the single neurons and

the composite synaptic activity of the local population (Buzsáki et al., 2012; Murthy & Fetz,

1996b; Okun et al., 2010). The magnitude indicates the strength of synchrony whereas the phase

can provide insights into the timing of the spikes relative to the local population.

Most instances of cross-frequency phase-amplitude coupling and spike-LFP synchrony were

stronger for deeper sleep states. This may be due to asynchronous activity during Move being

driven by functional local circuitry (i.e. generating movement) whereas activity during resting

and sleep states are more attuned to baseline macroscopic rhythmic activity, potentially related to

homeostatic plasticity (Tononi & Cirelli, 2014).

Delta and high gamma

One noticeable deviation from deeper sleep resulting in stronger synchrony was delta phase to

high gamma amplitude (delta-high gamma) coupling, which was significantly higher during

Move compared to Rest or REM. Delta oscillations may arise in the thalamus or the cortex and

are elevated during NREM. Delta has also been linked to various perceptual, sensorimotor, and

cognitive operations, potentially suppressing activity not necessary to the task at hand (Güntekin

& Başar, 2016; Harmony, 2013). In contrast, high gamma band LFP is representative of local

spiking activity, often strongly correlated with action potentials, and can also reflect neuronal

synchrony (Ray et al., 2008; Ray & Maunsell, 2011).

Evidence suggests the strength of delta-high gamma coupling is modulated by dopamine in both

the prefrontal cortex and the cortico-basal ganglia network (Andino-Pavlovsky et al., 2017;

Güntekin & Başar, 2016). Delta-high gamma coupling has also previously been shown to be

present during sleep in both the hippocampus and the neocortex (Clemens et al., 2009; Takeuchi

et al., 2015), but the underlying mechanisms are still unclear. Andino-Pavlovsky et al. (2017)

showed high delta-high gamma coupling during slow wave sleep in the rodent prefrontal cortex,

and Takeuchi et al. (2015) showed strong delta-high gamma coupling during REM and several

NREM substages in the primate hippocampus (Andino-Pavlovsky et al., 2017; Takeuchi et al.,

2015). However, neither study pursued the state-dependent comparisons, and their broad

definition of gamma (>25 Hz) makes independent interpretation difficult. Specific comparisons

of delta-high gamma coupling during different sleep states have not been reported.

Spike synchrony to delta band sheds additional light on the delta-high gamma relationship. In

our study, only RS neurons showed higher delta-high gamma coupling during Move, and the

phase of synchrony was consistently located at the falling edge of the wave (Figure 14). During

NREM both RS and FS spikes were synchronized to delta, but the phase of synchrony was

immediately after the trough of the wave. These results suggest delta as a coordinating signal that

reactivates cortical circuitry during sleep for consolidation (Gulati et al., 2017; Ramanathan et

al., 2015; Xu et al., 2019). Spiking activity during the day potentially drives delta activity, as it

occurs right before delta is the most depolarized, and delta during NREM may reactivate the

neural circuitry as it has the highest depolarization right before spiking activity. Similar phase

specificity is also reflected in delta-high gamma coupling during NREM.

Theta in the motor cortex and REM sleep

Theta oscillations have often been observed during REM sleep, and have also been shown to

coordinate hippocampal place cell activity during active exploration (Buzsáki & Moser, 2013;

Cantero et al., 2003). In the cortex, modulation of theta has been most prominently seen with

theta burst stimulation (TBS) – bursts of gamma frequency stimulation delivered at theta

intervals, typically with TMS over the motor region. TBS has been shown to increase the

excitability of the motor system, observed through increased motor evoked potentials, potentially

by mimicking the coupling of theta and gamma oscillations (Cárdenas-Morales et al., 2010;

Huang et al., 2005; Suppa et al., 2016). Additional evidence shows that strengthening theta-high

gamma coupling with transcranial alternating current stimulation (tACS) increases motor

learning (Akkad et al., 2021). In agreement with these previous findings, our results show much

higher theta-high gamma coupling during Move.

Theta has also been shown to be coupled to low gamma. A wealth of recent cross-frequency

coupling literature has reported theta-low gamma coupling in both hippocampus and the

neocortex, showing modulations dependent on task performance, cognitive engagement, and

memory formation (Buzsáki & Wang, 2012; Canolty et al., 2006; Lisman & Jensen, 2013).

Although stronger theta-low gamma coupling has also been observed during REM sleep, it was

absent in human studies, potentially due to hippocampal theta oscillations in humans being

slower than those in rodents (Cantero et al., 2003; Clemens et al., 2009; Jacobs, 2014). Similar to

the human studies, we did not find significant theta-low gamma coupling during REM, but we

also did not find significant delta-low gamma coupling.

Instead, we found very high theta-beta coupling during REM sleep. Theta-beta coupling is not a

common topic of study; there is some evidence that suggests it plays a role in working memory

and decision making, but the literature is sparse and focuses on the frontal lobe (Axmacher et al.,

2010; Cohen et al., 2009; Liang et al., 2021). In our results, the maximum beta amplitude occurs

right before the peak of theta (Figure 9), which means minimum beta occurs right after the

trough of theta. If beta is a regulating, inhibitory rhythm in the motor cortex (Engel & Fries,

2010; Khanna & Carmena, 2017; Kilavik et al., 2013), perhaps theta is increasing the excitability

of the cortex by minimizing beta, thus inducing disinhibition. This may lead to more effective

memory consolidation often seen during REM sleep, relevant to motor skill learning. However,

more research is needed to confirm these speculations.

Concluding Comments

Our study provides the first comprehensive analysis of cross-frequency phase amplitude

coupling and spike-field synchrony across all frequency bands within the cortex during different

behavioral states. We observed an increase in burst firing and high coordination between spikes

during NREM, consistent with previous findings suggesting reactivations of cortical circuitry

during NREM. High cross-frequency phase-amplitude coupling between delta and high gamma

when the animal is awake and moving and during non-REM sleep, as well as spike-field

synchrony with delta band LFP, suggest that delta may encode and subsequently drive these

reactivations. Modulations in theta to low gamma phase-amplitude coupling, commonly

observed in rodents, was not observed, similar to previous human studies. Instead, we observed

high theta to beta coupling during REM, potentially driving motor learning by increasing the

excitability of the cortex. These results support previous findings and may serve as the basis for

future studies into the roles of LFP frequency bands and different sleep states.

Acknowledgements

We thank Larry Shupe for programming and software support and Rebekah Schaefer and Becky

Burch for assistance with animal care, handling, training, and surgery. This work was supported

by the National Institutes of Health (NS012542, RR00166, and NS118781), the National Science

Foundation (EEC-1028725), and by the National Institutes of Health Office of the Director,

Office of Research Infrastructure Programs (ORIP) under award number P51OD010425 and

U42OD011123. The authors declare no competing financial interests.

References

Akkad, H., Dupont-Hadwen, J., Frese, A., Tetkovic, I., Barrett, L., Bestmann, S., & Stagg, C. J. (2021). Increasing motor skill acquisition by driving theta-gamma coupling. *ELife*, 1–15. https://doi.org/10.7554/eLife.67355

Andino-Pavlovsky, V., Souza, A. C., Scheffer-Teixeira, R., Tort, A. B. L., Etchenique, R., & Ribeiro, S. (2017). Dopamine modulates delta-gamma phase-amplitude coupling in the prefrontal cortex of behaving rats. *Frontiers in Neural Circuits*, *11*(May). https://doi.org/10.3389/fncir.2017.00029

Arbune, A. A., Popa, I., Mindruta, I., Beniczky, S., Donos, C., Daneasa, A., Mălîia, M. D., Băjenaru, O. A., Ciurea, J., & Barborica, A. (2020). Sleep modulates effective connectivity: A study using intracranial stimulation and recording. *Clinical Neurophysiology*, *131*(2), 529–541. https://doi.org/10.1016/j.clinph.2019.09.010

Aserinsky, E., & Kleitman, N. (1953). Regularly Occurring Periods of Eye Motility, and Concomitant Phenomena, during Sleep. *Science*, *118*(3062), 273–274.

Axmacher, N., Henseler, M. M., Jensen, O., Weinreich, I., Elger, C. E., & Fell, J. (2010). Cross-frequency coupling supports multi-item working memory in the human hippocampus. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(7), 3228–3233. https://doi.org/10.1073/pnas.0911531107

Bellesi, M., Riedner, B. A., Garcia-Molina, G. N., Cirelli, C., & Tononi, G. (2014). Enhancement of sleep slow waves: Underlying mechanisms and practical consequences. *Frontiers in Systems Neuroscience*, *8*(October), 1–17. https://doi.org/10.3389/fnsys.2014.00208

Boyce, R., Glasgow, S. D., Williams, S., & Adamantidis, A. (2016). Sleep research: Causal evidence for the role of REM sleep theta rhythm in contextual memory consolidation. *Science*, *352*(6287), 812–816. https://doi.org/10.1126/science.aad5252

Buzsáki, G., Anastassiou, C. A., & Koch, C. (2012). The origin of extracellular fields and currents-EEG, ECoG, LFP and spikes. *Nature Reviews Neuroscience*, *13*(6), 407–420. https://doi.org/10.1038/nrn3241

Buzsáki, G., & Moser, E. I. (2013). Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nature Neuroscience*, *16*(2), 130–138. https://doi.org/10.1038/nn.3304

Buzsáki, G., & Wang, X.-J. (2012). Mechanisms of Gamma Oscillations. *Annual Review of Neuroscience*, *6*(9), 2166–2171. https://doi.org/10.1021/nl061786n.Core-Shell

Canolty, R. T., Edwards, E., Dalal, S. S., Soltani, M., Nagarajan, S. S., Kirsch, H. E., Berger, M. S., Barbare, N. M., & Knight, R. T. (2006). High gamma power is phase-locked to theta oscillations in human neocortex. *Science*, *313*(5793), 1626–1628. https://doi.org/10.1126/science.1128115

Canolty, R. T., & Knight, R. T. (2010). The functional role of cross-frequency coupling. *Trends in Cognitive Sciences*, *14*(11), 506–515. https://doi.org/10.1016/j.tics.2010.09.001

Cantero, J. L., Atienza, M., Stickgold, R., Kahana, M. J., Madsen, J. R., & Kocsis, B. (2003). Sleep-Dependent θ Oscillations in the Human Hippocampus and Neocortex. *Journal of Neuroscience*, *23*(34), 10897–10903. https://doi.org/10.1523/jneurosci.23-34-10897.2003

Cárdenas-Morales, L., Nowak, D. A., Kammer, T., Wolf, R. C., & Schönfeldt-Lecuona, C. (2010). Mechanisms and applications of theta-burst rTMS on the human motor cortex. *Brain Topography*, *22*(4), 294–306. https://doi.org/10.1007/s10548-009-0084-7

Clemens, Z., Weiss, B., Szucs, A., Eross, L., Rásonyi, G., & Halász, P. (2009). Phase coupling between rhythmic slow activity and gamma characterizes mesiotemporal rapid-eye-movement sleep in humans. *Neuroscience*, *163*(1), 388–396. https://doi.org/10.1016/j.neuroscience.2009.06.044

Cohen, M. X., Elger, C. E., & Fell, J. (2009). Oscillatory activity and phase-amplitude coupling in the human medial frontal cortex during decision making. *Journal of Cognitive Neuroscience*, *21*(2), 390–402. https://doi.org/10.1162/jocn.2008.21020

Craik, A., He, Y., & Contreras-Vidal, J. L. (2019). Deep learning for electroencephalogram (EEG) classification tasks: A review. *Journal of Neural Engineering*, *16*(3). https://doi.org/10.1088/1741-2552/ab0ab5

Dijk, D. J., Beersma, D. G. M., & Daan, S. (1987). EEG Power Density during Nap Sleep: Reflection of an Hourglass Measuring the Duration of Prior Wakefulness. *Journal of Biological Rhythms*, *2*(3), 207–219. https://doi.org/10.1177/074873048700200304

Engel, A. K., & Fries, P. (2010). Beta-band oscillations-signalling the status quo? *Current Opinion in Neurobiology*, *20*(2), 156–165. https://doi.org/10.1016/j.conb.2010.02.015

Fernandez, L. M. J., & Lüthi, A. (2020). Sleep spindles: Mechanisms and functions. *Physiological Reviews*, *100*(2), 805–868. https://doi.org/10.1152/physrev.00042.2018

Gerla, V., Kremen, V., Macas, M., Dudysova, D., Mladek, A., Sos, P., & Lhotska, L. (2019). Iterative expert-in-the-loop classification of sleep PSG recordings using a hierarchical clustering. *Journal of Neuroscience Methods*, *317*(February), 61–70. https://doi.org/10.1016/j.jneumeth.2019.01.013

Gulati, T., Guo, L., Ramanathan, D. S., Bodepudi, A., Ganguly, K., Service, R., Francisco, S., Affairs, V., Francisco, S., Francisco, S., Francisco, S., Affairs, V., Francisco, S., & Francisco, S. (2017). Neural reactivations during sleep determine network credit assignment. *Nature Neuroscience*, *20*(9), 1277–1284. https://doi.org/10.1038/nn.4601.Neural

Güntekin, B., & Başar, E. (2016). Review of evoked and event-related delta responses in the human brain. *International Journal of Psychophysiology*, *103*, 43–52. https://doi.org/10.1016/j.ijpsycho.2015.02.001

Hamida, S. T. Ben, & Ahmed, B. (2013). Computer based sleep staging: Challenges for the future. *2013 7th IEEE GCC Conference and Exhibition, GCC 2013*, 280–285. https://doi.org/10.1109/IEEEGCC.2013.6705790

Harmony, T. (2013). The functional significance of delta oscillations in cognitive processing. *Frontiers in Integrative Neuroscience*, *7*(DEC), 1–10.

https://doi.org/10.3389/fnint.2013.00083

Himanen, S. L., & Hasan, J. (2000). Limitations of Rechtschaffen and Kales. *Sleep Medicine Reviews*, *4*(2), 149–167. https://doi.org/10.1053/smrv.1999.0086

Hsieh, K. C., Robinson, E. L., & Fuller, C. A. (2008). Sleep architecture in unrestrained rhesus monkeys (Macaca mulatta) synchronized to 24-hour light-dark cycles. *Sleep*, *31*(9), 1239–1250.

Huang, Y. Z., Edwards, M. J., Rounis, E., Bhatia, K. P., & Rothwell, J. C. (2005). Theta burst stimulation of the human motor cortex. *Neuron*, *45*(2), 201–206. https://doi.org/10.1016/j.neuron.2004.12.033

Huber, R., Ghilardi, M. F., Massimini, M., & Tononi, G. (2004). Local sleep and learning. *Nature*, *430*(6995), 78–81. https://doi.org/10.1038/nature02663

Hülsemann, M. J., Naumann, E., & Rasch, B. (2019). Quantification of phase-amplitude coupling in neuronal oscillations:comparison of phase-locking value, mean vector length, modulation index, and generalized-linear-modeling-cross-frequency-coupling. *Frontiers in Neuroscience*, *13*(JUN), 1–15. https://doi.org/10.3389/fnins.2019.00573

Insel, N., & Barnes, C. A. (2015). Differential activation of fast-spiking and regular-firing neuron populations during movement and reward in the dorsal medial frontal cortex. *Cerebral Cortex*, *25*(9), 2631–2647. https://doi.org/10.1093/cercor/bhu062

Jacobs, J. (2014). Hippocampal theta oscillations are slower in humans than in rodents: Implications for models of spatial navigation and memory. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *369*(1635). https://doi.org/10.1098/rstb.2013.0304

Jensen, O., & Colgin, L. L. (2007). Cross-frequency coupling between neuronal oscillations. *Trends in Cognitive Sciences*, *11*(7), 267–269. https://doi.org/10.1016/j.tics.2007.05.003

Jung, M. W., Qin, Y., McNaughton, B. L., & Barnes, C. A. (1998). Firing characteristics of deep layer neurons in prefrontal cortex in rats performing spatial working memory tasks. *Cerebral Cortex*, *8*(5), 437–450. https://doi.org/10.1093/cercor/8.5.437

Khalighi, S., Sousa, T., Pires, G., & Nunes, U. (2013). Automatic sleep staging: A computer assisted approach for optimal combination of features and polysomnographic channels. *Expert Systems with Applications*, *40*(17), 7046–7059. https://doi.org/10.1016/j.eswa.2013.06.023

Khanna, P., & Carmena, J. M. (2017). Beta band oscillations in motor cortex reflect neural population signals that delay movement onset. *ELife*, *6*, 1–31. https://doi.org/10.7554/eLife.24573

Kilavik, B. E., Zaepffel, M., Brovelli, A., MacKay, W. A., & Riehle, A. (2013). The ups and downs of beta oscillations in sensorimotor cortex. *Experimental Neurology*, *245*, 15–26. https://doi.org/10.1016/j.expneurol.2012.09.014

Kim, E. Y., Mahmoud, G. S., & Grover, L. M. (2005). REM sleep deprivation inhibits LTP in vivo in area CA1 of rat hippocampus. *Neuroscience Letters*, *388*(3), 163–167. https://doi.org/10.1016/j.neulet.2005.06.057

Kripke, D. F., Reite, M. L., Pegram, G. V., Stephens, L. M., & Lewis, O. F. (1968). Nocturnal sleep in rhesus monkeys. *Electroencephalography and Clinical Neurophysiology*, *24*(6), 581–586. https://doi.org/10.1016/0013-4694(68)90047-3

Liang, W. K., Tseng, P., Yeh, J. R., Huang, N. E., & Juan, C. H. (2021). Frontoparietal Beta Amplitude Modulation and its Interareal Cross-frequency Coupling in Visual Working Memory. *Neuroscience*, *460*, 69–87. https://doi.org/10.1016/j.neuroscience.2021.02.013

Libey, T., & Fetz, E. E. (2017). Open-source, low cost, free-behavior monitoring, and reward system for neuroscience research in non-human primates. *Frontiers in Neuroscience*, *11*(MAY), 1–12. https://doi.org/10.3389/fnins.2017.00265

Lisman, J. E., & Jensen, O. (2013). The Theta-Gamma Neural Code. *Neuron*, *77*(6), 1002–1016. https://doi.org/10.1016/j.neuron.2013.03.007

Malafeev, A., Laptev, D., Bauer, S., Omlin, X., Wierzbicka, A., Wichniak, A., Jernajczyk, W., Riener, R., Buhmann, J., & Achermann, P. (2018). Automatic human sleep stage scoring using deep neural networks. *Frontiers in Neuroscience*, *12*(NOV), 1–15. https://doi.org/10.3389/fnins.2018.00781

Marshall, L., Helgadóttir, H., Mölle, M., & Born, J. (2006). Boosting slow oscillations during sleep potentiates memory. *Nature*, *444*(7119), 610–613. https://doi.org/10.1038/nature05278

Mascetti, G. G. (2016). Unihemispheric sleep and asymmetrical sleep: Behavioral, neurophysiological, and functional perspectives. *Nature and Science of Sleep*, *8*, 221–228. https://doi.org/10.2147/NSS.S71970

McCormick, D. A., Connors, B. W., Lighthall, J. W., & Prince, D. A. (1985). Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *Journal of Neurophysiology*, *54*(4), 782–806. https://doi.org/10.1152/jn.1985.54.4.782

Michel, C. M., Murray, M. M., Lantz, G., Gonzalez, S., Spinelli, L., & Grave De Peralta, R. (2004). EEG source imaging. *Clinical Neurophysiology*, *115*(10), 2195–2222. https://doi.org/10.1016/j.clinph.2004.06.001

Moses, J. M., Hord, D. J., Lubin, A., Johnson, L. C., & Naitoh, P. (1975). Dynamics of nap sleep during a 40 hour period. *Electroencephalography and Clinical Neurophysiology*, *39*(6), 627–633. https://doi.org/10.1016/0013-4694(75)90075-9

Murthy, V. N., & Fetz, E. E. (1996a). Oscillatory activity in sensorimotor cortex of awake monkeys: synchronization of local field potentials and relation to behavior. *Journal of Neurophysiology*, *76*(6), 3949–3967.

Murthy, V. N., & Fetz, E. E. (1996b). Synchronization of neurons during local field potential oscillations in sensorimotor cortex of awake monkeys. *Journal of Neurophysiology*, *76*(6), 3968–3982. http://www.ncbi.nlm.nih.gov/pubmed/8985893

Okun, M., Naim, A., & Lampl, I. (2010). The subthreshold relation between cortical local field potential and neuronal firing unveiled by intracellular recordings in awake rats. *Journal of Neuroscience*, *30*(12), 4440–4448. https://doi.org/10.1523/JNEUROSCI.5062-09.2010

Onslow, A. C. E., Bogacz, R., & Jones, M. W. (2011). Quantifying phase-amplitude coupling in neuronal network oscillations. *Progress in Biophysics and Molecular Biology*, *105*(1–2), 49–57. https://doi.org/10.1016/j.pbiomolbio.2010.09.007

Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z., Gimelshein, N., Antiga, L., Desmaison, A., Köpf, A., Yang, E., DeVito, Z., Raison, M., Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., … Chintala, S. (2019). PyTorch: An imperative style, high-performance deep learning library. *Advances in Neural Information Processing Systems*, *32*(NeurIPS).

Penzel, T., & Conradt, R. (2000). Computer based sleep recording and analysis. *Sleep Medicine Reviews*, *4*(2), 131–148. https://doi.org/10.1053/smrv.1999.0087

Ramanathan, D. S., Gulati, T., & Ganguly, K. (2015). Sleep-Dependent Reactivation of Ensembles in Motor Cortex Promotes Skill Consolidation. *PLoS Biology*, *13*(9), 1–25. https://doi.org/10.1371/journal.pbio.1002263

Ray, S., Crone, N. E., Niebur, E., Franaszczuk, P. J., & Hsiao, S. S. (2008). Neural correlates of high-gamma oscillations (60-200 Hz) in macaque local field potentials and their potential implications in electrocorticography. *Journal of Neuroscience*, *28*(45), 11526–11536. https://doi.org/10.1523/JNEUROSCI.2848-08.2008

Ray, S., & Maunsell, J. H. R. (2011). Different origins of gamma rhythm and high-gamma activity in macaque visual cortex. *PLoS Biology*, *9*(4). https://doi.org/10.1371/journal.pbio.1000610

Rechtschaffen, A. (1998). Current perspectives on the function of sleep. *Perspectives in Biology and Medicine*, *41*(3), 359–390. https://doi.org/10.1353/pbm.1998.0051

Rechtschaffen, A., & Kales, A. (1968). A manual of standardized terminology, techniques and scoring system for sleep stages of human subjects. *U.S. Public Health Service, U.S. Government Printing Office, Washington D.C.*

Rembado, I., Zanos, S., & Fetz, E. E. (2017). Cycle-triggered cortical stimulation during slow wave sleep facilitates learning a BMI task: A case report in a non-human primate. *Frontiers in Behavioral Neuroscience*, *11*(April), 1–13. https://doi.org/10.3389/fnbeh.2017.00059

Sejnowski, T. J., Churchland, P. S., & Movshon, J. A. (2014). Putting big data to good use in neuroscience. *Nature Neuroscience*, *17*(11), 1440–1441. https://doi.org/10.1038/nn.3839

Sekkal, R. N., Bereksi-Reguig, F., Ruiz-Fernandez, D., Dib, N., & Sekkal, S. (2022). Automatic sleep stage classification: From classical machine learning methods to deep learning. *Biomedical Signal Processing and Control*, *77*(October 2021), 103751. https://doi.org/10.1016/j.bspc.2022.103751

Shupe, L. E., Miles, F. P., Jones, G., Yun, R., Mishler, J., Rembado, I., Murphy, R. L., Perlmutter, S. I., & Fetz, E. E. (2021). Neurochip3: An autonomous multichannel bidirectional brain-computer interface for closed-loop activity-dependent stimulation. *Frontiers in Neuroscience*, *15*(August), 1–15. https://doi.org/10.3389/fnins.2021.718465

Siclari, F., & Tononi, G. (2017). Local aspects of sleep and wakefulness. *Current Opinion in Neurobiology*, *44*, 222–227. https://doi.org/10.1016/j.conb.2017.05.008

Siegel, J. M. (2005). Clues to the functions of mammalian sleep. *Nature*, *437*, 1264–1271.

Silber, M. H., Ancoli-Israel, S., Bonnet, M. H., Chokroverty, S., Grigg-Damberger, M. M.,
Hirshkowitz, M., Kapen, S., Keenan, S. A., Kryger, M. H., Penzel, T., Pressman, M. R., &
Iber, C. (2007). The visual scoring of sleep in adults. *Journal of Clinical Sleep Medicine*,
3(2), 121–131.

Steriade, M., Timofeev, I., & Grenier, F. (2001). Natural waking and sleep states: A view from
inside neocortical neurons. *Journal of Neurophysiology*, *85*(5), 1969–1985.
https://doi.org/10.1152/jn.2001.85.5.1969

Sun, S., Jiang, L. P., Peterson, S. M., Herron, J., Weaver, K., Ko, A., Ojemann, J., & Rao, R. P.
N. (2020). Unsupervised Sleep and Wake State Identification in Long-Term
Electrocorticography Recordings. *Annu Int Conf IEEE Eng Med Biol Soc.*, 629–632.

Suppa, A., Huang, Y. Z., Funke, K., Ridding, M. C., Cheeran, B., Di Lazzaro, V., Ziemann, U.,
& Rothwell, J. C. (2016). Ten Years of Theta Burst Stimulation in Humans: Established
Knowledge, Unknowns and Prospects. *Brain Stimulation*, *9*(3), 323–335.
https://doi.org/10.1016/j.brs.2016.01.006

Supratak, A., Dong, H., Wu, C., & Guo, Y. (2017). DeepSleepNet: A model for automatic sleep
stage scoring based on raw single-channel EEG. *IEEE Transactions on Neural Systems and
Rehabilitation Engineering*, *25*(11), 1998–2008.
https://doi.org/10.1109/TNSRE.2017.2721116

Takeuchi, S., Mima, T., Murai, R., Shimazu, H., Isomura, Y., & Tsujimoto, T. (2015). Gamma
oscillations and their cross-frequency coupling in the primate hippocampus during sleep.
Sleep, *38*(7), 1085-1091G. https://doi.org/10.5665/sleep.4818

Tononi, G., & Cirelli, C. (2014). Sleep and the Price of Plasticity: From Synaptic and Cellular
Homeostasis to Memory Consolidation and Integration. *Neuron*, *81*(1), 12–34.
https://doi.org/10.1016/j.neuron.2013.12.025

Tort, A. B. L., Komorowski, R., Eichenbaum, H., & Kopell, N. (2010). Measuring phase-
amplitude coupling between neuronal oscillations of different frequencies. *Journal of
Neurophysiology*, *104*(2), 1195–1210. https://doi.org/10.1152/jn.00106.2010

Tsinalis, O., Matthews, P. M., & Guo, Y. (2016). Automatic Sleep Stage Scoring Using Time-
Frequency Analysis and Stacked Sparse Autoencoders. *Annals of Biomedical Engineering*,
44(5), 1587–1597. https://doi.org/10.1007/s10439-015-1444-y

Ulrich, D. (2016). Sleep Spindles as Facilitators of Memory Formation and Learning. *Neural
Plasticity*, *2016*. https://doi.org/10.1155/2016/1796715

van der Maaten, L., & Hinton, G. (2008). Visualizing Data using t-SNE. *Journal of Machine
Learning Research*, *9*, 2579–2605.

Vecsey, C. G., Huang, T., & Abel, T. (2018). Sleep deprivation impairs synaptic tagging in
mouse hippocampal slices. *Neurobiology of Learning and Memory*, *154*(March), 136–140.
https://doi.org/10.1016/j.nlm.2018.03.016

Vigneswaran, G., Kraskov, A., & Lemon, R. N. (2011). Large identified pyramidal cells in

macaque motor and premotor cortex exhibit "Thin Spikes": Implications for cell type
classification. *Journal of Neuroscience*, *31*(40), 14235–14242.
https://doi.org/10.1523/JNEUROSCI.3142-11.2011

Xu, W., de Carvalho, F., & Jackson, A. (2019). Sequential Neural Activity in Primary Motor
Cortex during Sleep. *The Journal of Neuroscience*, *39*(19), 3698–3712.
https://doi.org/10.1523/jneurosci.1408-18.2019

Younes, M., Raneri, J., & Hanly, P. (2016). Staging sleep in polysomnograms: Analysis of inter-
scorer variability. *Journal of Clinical Sleep Medicine*, *12*(6), 885–894.
https://doi.org/10.5664/jcsm.5894

Figures

Figure 1. <u>Experimental design</u>

A. All data was gathered using the Neurochip3 on freely behaving macaques for up to 24 hours. **B.** We collected 16 channels of data at 20 kHz and obtained the spectral density of every 8-second bin to classify behavioral states. **C.** We then calculated the cross-frequency phase-amplitude coupling of every frequency band pair. The example shows the instantaneous amplitude (red) of high gamma filtered LFP (black, bottom) increasing at the trough of beta (black, top). **D.** We additionally sorted single units and found the synchrony of spikes to LFP bands. The example shows spikes (red) synchronized with the trough of beta filtered LFP.

Figure 2. <u>Dimensionality reduction</u>

A. Examples of original spectra and the reconstruction at the output of a trained autoencoder for
each classified state. **B.** t-SNE visualization of the low dimensional representation with finalized
states. **C.** An example of the average of 100 spectra that generated the highest values in each
hidden unit. The darkness of the traces is weighted by the variance in the values of all values in
that hidden unit (i.e. more relevant during clustering). **D.** The average of 100 spectra that
generated the lowest values in each hidden unit subtracted from (C).

Figure 3. <u>K-means clustering</u>

A. The average inertia (black), or within-clusters sum-of-squares, and the average silhouette
coefficient (red) of the low dimensional data for different number of clusters. **B.** The silhouette
coefficient of each data point with the final classifications compared to the average (vertical red
line). All clusters have significant portions above the average.

Figure 4. <u>State classification</u>

A. Diagram of the full classification process. Normalized LFP power spectral density was used to train an autoencoder for dimensionality reduction. The lower dimensional representation and normalized accelerometer signals were subsequently clustered with k-means clustering then smoothed with a majority filter to determine different sleep states. **B.** Example of classification and spectra over 22 hours of continuous recording. Brief periods of REM sleep were observed during the day, attributed to naps (arrow). **C.** Averaged spectra across each state for the session shown in (B). There is high delta power during NREM, theta power during REM, and beta power during Move and Rest states. **D.** Example traces of raw LFP during each state. Spindles (black arrow) and k-complexes (red arrow) were observed during NREM sleep.

    Figure 5. <u>Validation of classification</u>

A. An example of filtered differential EOG signals and corresponding eye movements. **B.**
Standard deviation of EOG signals overnight with respect to classified states. Note the increase
in large eye movements during REM and waking. **C.** An example of the normalized variance of
accelerometer data and the normalized variance of the Kinect data with respect to classified
states. **D.** Test and training errors with k-fold cross-validation in percent (left) and total minutes
(right).
881

Figure 6. <u>LFP dynamics</u>

A. Average spectral power in each state for each animal across all experiments. Shaded regions
show standard error. **B.** Average pairwise coherence in each state for each animal across all
experiments. Shaded regions show standard error.

889 **Figure 7. <u>Cross-frequency phase-power distributions</u>**

890 Distribution of lower frequency band phase and higher frequency band power during each state.
891 The power is z-scored for each frequency.
892

Figure 8. <u>Normalized MVL distributions</u>

896 Normalized MVL distributions for each lower frequency band phase and higher frequency band
897 amplitude pair during each state. The black boxes show standard box plots with interquartile
898 range and the white dots show median values. The numbers above each group denote
899 significance compared to another state (Friedman's test, $p < 0.05$).
900

Figure 9. <u>Mean vector phase distributions</u>

903 The distribution of phases of mean vectors for each lower frequency band phase and higher
904 frequency band amplitude pair during each state. The phase of the mean vector shows the phase
905 at which the amplitude of the higher frequency band is the greatest.
906

907
908

Figure 10. <u>Spike sorting</u>

A. An example of a sorted spike. The grey traces show a random sample of 1000 spikes, the
black line is the average, the horizontal dashed red line shows the threshold, and the blue vertical
lines show the two windows. **B.** An example of comparisons of the pairwise coefficient of
determination (CoD) between the first and last 1000 instances of the same spike and the first and
last 1000 instances of two different spikes. **C.** Firing rates binned every 60 seconds of four
neurons overnight with classified sleep states. The changes in firing rate very closely match the
changes in state. **D.** Histogram of the peak of the ISI distribution of each spike. Spikes with
peaks earlier than 10 ms were denoted to be fast spiking (FS) and all others denoted to be regular
spiking (RS). **E.** Percent difference of firing rate in each state from the average overall firing rate
for each spike type. Each state is statistically significantly different from each other state
(Friedman's test, $p < 0.05$).

Figure 11. <u>Spike features</u>

A. Spike width, dV/dt ratio (the fastest rise of the spike divided by the fastest fall of the spike as
defined in McCormick et al. 1985), and the average firing rate with respect to the peak of the
inter-spike interval (ISI) distribution. The spike widths include the addition of a random jitter
sampled from a uniform distribution from -0.02 to 0.02 ms to better visualize the data. The
dashed vertical line denotes the boundary between fast- and regular-spiking neurons (FS and RS
respectively) as defined in this study. **B.** Box plots of the corresponding measures and the
statistical significance between the two spike types (Wilcoxon rank-sum test). The two measures
reflecting the spike shape have significant differences. Although FS neurons have a wider spread
of average firing rates, there is no significant difference between the average firing rates of the
two types of neurons.
934

 Figure 12. <u>Changes in spiking patterns</u>

A. Average ISI distributions of identified fast spiking and regular spiking neurons during each
state. **B.** Difference of ISI distributions from Move plotted against time (left) and frequency
(right). Note the peak around 5 ms during NREM for regular spiking neurons and the peaks
around 7-10 Hz during NREM sleep and the peaks at around 20 and 25 Hz for REM and Rest
respectively (arrows) for both neuron types. All three peaks were significantly different from 0
(Wilcoxon signed-rank test, p<0.05). **C.** Cross correlations of spike firing rates between pairs of
fast spiking (FS) neurons (left), FS and regular spiking (RS) neurons (middle), and pairs of RS
neurons (right) (n = 636, 1328, 1036 respectively).

947 **Figure 13. <u>Phase distributions of LFPs at spike times</u>**

948 Phase distributions normalized for each amplitude during each sleep state for each LFP band.
949

952 **Figure 14. <u>Locked phase and phase locking value distributions for regular</u>**

953 **<u>spiking neurons</u>**

954 Distributions of locked phase (left) and PLVs (right) of regular spiking neurons during each
955 sleep state for each LFP band. The numbers above each group denote significance compared to
956 another state (Friedman's test, $p < 0.05$).
957

Figure 15. <u>Locked phase and phase locking value distributions for fast spiking</u>

<u>neurons</u>

Distributions of locked phase (left) and PLVs (right) of fast spiking neurons during each sleep state for each LFP band. The numbers above each group denote significance compared to another state (Friedman's test, p<0.05).