Local field potentials and single unit dynamics in motor cortex of unconstrained macaques during different behavioral states

Richy Yun[1,4] , Jonathan H. Mishler[1,4] , Steve I. Perlmutter[2,4,5], Rajesh P. N. Rao[3,4], Eberhard E. Fetz[1,2,4,5]

Departments of [1]Bioengineering, [2]Physiology and Biophysics, [3]Allen School for Computer Science and Engineering, University of Washington
[4]Center for Neurotechnology, University of Washington
[5]Washington National Primate Research Center, University of Washington

Abstract

Different sleep states have been shown to be vital for a variety of brain function, including learning, memory, and skill consolidation. However, our understanding of neural dynamics during sleep and the role of prominent LFP frequency bands remain incomplete. To discern changes between different behavioral states we collected multichannel LFP and spike data in primary motor cortex of unconstrained macaques for up to 24 hours using the Neurochip3. Each 8 second bin of time was classified into awake and moving (Move), awake and at rest (Rest), REM sleep (REM), or non-REM sleep (NREM) by using dimensionality reduction and clustering on the average spectral density and the acceleration of the head. LFP power showed high delta during NREM, high theta during REM, and high beta when the animal was awake. Cross-frequency phase-amplitude coupling typically showed higher coupling for deeper sleep between all pairs of frequency bands except for high delta-high gamma and theta-high gamma coupling during Move, and high theta-beta coupling during REM. Sorted single units showed decreased firing rate with deeper sleep, though with higher "bursty" patterns during NREM compared to other states. Spike-LFP synchrony showed high delta synchrony during Move, and higher coupling with all other frequency bands during NREM. These results altogether are consistent with previous findings showing reactivation of cortical circuitry activated during the day, which may be driven by the delta band LFP.

Introduction

The discovery of rapid-eye movement (REM) sleep by Aserinski and Kleitman showed the presence of active brain processes during sleep, which catalyzed studies on different sleep stages (Aserinsky & Kleitman, 1953). Findings have shown REM sleep to play a role in regulating the autonomous nervous system activity and to be linked with spatial and emotional memory consolidation (Boyce et al., 2016; Rechtschaffen, 1998; Siegel, 2005). Similarly, non-REM (NREM) sleep has been shown to be tied to skill consolidation and memory formation (Ramanathan et al., 2015; Siclari & Tononi, 2017).

Studies have furthered our understanding of different sleep stages by exploring the dynamics of local field potential (LFP) frequency band power and single-unit spiking characteristics in both the cortex and various deep brain structures during sleep. Slow waves and delta frequency band are typically present across the brain during NREM sleep, and high theta power in the hippocampus is present during REM sleep (Rechtschaffen & Kales, 1968; Silber et al., 2007). Single units show changes in firing rate as well as firing patterns during sleep, due to changes in connectivity and excitability, potentially reflecting reactivations of relevant cortical circuit patterns to solidify learning and memory (Arbune et al., 2020; Ramanathan et al., 2015; Tononi & Cirelli, 2014; Xu et al., 2019). Several other features of cortical activity, such as k-complexes and sleep spindles during NREM sleep, have also been identified to be associated with different stages of sleep (Fernandez & Lüthi, 2020; Huber et al., 2004; Ulrich, 2016).

Additional findings have shown that certain plasticity mechanisms, such as long-term potentiation/depression and synaptic tagging and capture, occur exclusively during sleep (Kim et al., 2005; Vecsey et al., 2018). Subsequent research has applied this knowledge by enhancing slow waves prevalent during NREM sleep (Bellesi et al., 2014; Marshall et al., 2006) or

delivering stimulation locked to delta (Rembado et al., 2017) to augment learning and memory.

However, despite our expanding tools and techniques to take advantage of sleep mechanisms, the

neural dynamics underlying sleep remain unclear.

Various measures of local field potential (LFP) coupling and spike-LFP synchrony have been

commonly used to elucidate mechanisms of neural dynamics. Cross-frequency phase-amplitude

coupling of different LFP frequency bands, thought to reflect communication between brain

circuitry, has been shown to be modulated by task performance, cognitive engagement, and

memory formation in various brain regions (Canolty & Knight, 2010; Jensen & Colgin, 2007).

Single units are strongly synchronized to specific phases of LFP frequency bands, notably to beta

and gamma cycles in the motor cortex, suggesting beta to be a resting rhythm during movement

execution and gamma to reflect local population activity (Buzsáki et al., 2012; Buzsáki & Wang,

2012; Engel & Fries, 2010; Murthy & Fetz, 1996a, 1996b). These two analyses have given us

significant insight into neural signaling and the role of both spikes and LFPs in brain function,

but there has yet to be a comprehensive analysis of cross-frequency coupling and spike-LFP

synchrony for specific cortical sites during sleep and wake.

This study analyzes LFP and single unit dynamics in the macaque motor cortex during different

behavioral states to better understand the mechanisms operating during different sleep stages as

well as the roles of various LFP frequency bands. We first used the power spectral density of

LFPs to distinguish between four major behavioral states shown to be relevant to plasticity and

learning in the motor cortex: 1. awake and moving (Move), 2. awake and at rest (Rest), 3. rapid-

eye movement (REM) sleep, and 4. non-REM (NREM) sleep. We tracked single-unit activity

concurrently with LFP recordings and focused on 6 different frequency bands commonly

observed in the cortex – delta, theta, alpha, beta, low gamma, and high gamma. Finally, we

assessed state-dependent changes in cross-frequency coupling between pairs of the frequency

bands as well as spike-LFP synchrony with each frequency band.

Materials and Methods

Experimental design

Implants and surgical procedures

The experiments were conducted on two male *Macaca nemestrina* monkeys (Monkeys J and K). Surgeries were performed under isoflurane anesthesia and aseptic conditions to implant multi-electrode Utah arrays. All procedures conformed to the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* and were approved by the University of Washington Institutional Animal Care and Use Committee.

Implantation of the Utah array was guided by stereotaxic coordinates. A 1.5 cm wide square craniotomy was performed over the hand region of the primary motor cortex to expose the dura. Three sides of the exposed dura were cut to expose the cortex; a Utah array was implanted and the dura was sutured over the array. Two reference wires were inserted below the dura and two were inserted between the dura and the skull. The bone flap from the craniotomy was replaced and held in place by a titanium strap screwed onto the skull with 2.5 mm x 6 mm titanium skull screws. A second smaller titanium strap was fastened to the skull to secure the wire bundle. The connector pedestal for the array was attached to the skull with eight titanium skull screws and the incision closed around the pedestal base. Additional skull screws were placed around the base of the connector pedestal and a thin coat of dental acrylic (methyl methacrylate) was applied to the skull between the screws and the connector base for additional stability.

The animals also received a "halo" implant made with 3/8" aluminum into an egg-shaped oval 17 cm long and 15.3 cm wide. Four titanium straps were affixed to the skull by titanium bone screws. Two were implanted bilaterally over the occipital ridge, and two were placed temporally

bilaterally. After the plates integrated with the skull for 6 weeks, an aluminum halo was mounted

on four pins seated in each plate. A titanium can with a plexiglass top to house the Neurochip3

was attached to the halo during all Neurochip3 recording sessions.

Monkey J additionally received EOG implants on the lateral wall and superior margin of both

orbits. EOG electrodes consisted of a titanium washer ((#4, 0.25" OD, 0.032" thick) with a

0.016" hole drilled into it and a 34-gauge silver-plated copper microwire with silicone shielding

(Cooner Wire, AS155-34) threaded through the hole and soldered to the washer. An incision was

made to the dorsal and lateral margins of both orbits to expose the skull. A small hole was drilled

in each incision, a titanium skull screw (2 mm x 6 mm) was used to hold down the EOG

electrodes, and the wires of the electrodes were tunneled beneath the tissue along the skull.

Another incision was made along the top of the skull to secure a pedestal with 8 skull screws and

allow access to the electrode wires. The inside of the pedestal was filled with surgical silicone

adhesive (Kwik-Sil, World Precision Instruments) to prevent infection. Connectors were fastened

to the wires postoperatively.

After each surgery animals received postoperative courses of analgesics and antibiotics. Animals

did not show signs of discomfort or pain related to any implanted devices after recovery, and all

exposed implants were regularly disinfected biweekly with chlorohexidine and treated with

antibiotics to prevent infection.

Electrophysiology

All data was collected with the head-mounted Neurochip3 (Shupe et al., 2021) while the

monkeys were freely behaving in their home cage (Figure 1A). The Neurochip3 is a battery

powered bidirectional brain-computer interface capable of saving data to an SD card, allowing

for wireless recording and stimulation for up to 24 hours. 16 channels of the Utah array, or 14

channels of the array and 2 channels of the EOG, were recorded at 20 kHz sampling rate with a

bandwidth of 0.1 Hz to 5 kHz (Figure 1B). The cortical channels were chosen each day with a

preliminary recording to capture the largest single units present in the array. Most experiments

recorded a different set of channels due to the spikes changing over time.

The Neurochip3 also has an onboard 3-axis accelerometer that was simultaneously recorded at

100 samples per second. Recording sessions lasted between 19 and 24 hours. The animal rooms

had the lights off during the night for 12 hours between 6pm and 6am. We collected data from a

total of 16 sessions from Monkey K and 10 sessions from Monkey J.

Data analysis

Classifying sleep states

All analyses were performed using custom MATLAB (Mathworks) and Python code. The power

spectral density (PSD) of cortical local field potentials (LFPs) as well as the onboard

accelerometer data from the Neurochip3 were used to classify different sleep states (Figure 1B).

Data was down sampled to 1 kHz before performing Welch's PSD estimate between 0 and 50 Hz

for each 8 second time-bin. We then converted the PSD into power and found the average power

across all channels for each time-bin. The average power was further normalized by subtracting

by the minimum value and dividing by the integral to ensure relative power at different

frequencies played a larger role than the absolute power.

We then used the normalized average power as inputs to train a stacked sparse autoencoder for

dimensionality reduction, similar to the architecture described in Tsinalis et al. 2016 (Tsinalis et

al., 2016). The encoder was composed of 3 layers of 256, 128, and 64 units each, with batch

normalization and ReLU activation function. The final hidden layer containing the reduced

representation of the data had 32 nodes. The autoencoder was trained with minibatch sizes of 64

for 300 epochs. The loss was calculated with mean squared error with L1 regularization

(regularization weight $\lambda = 1e^{-5}$) using the Adam optimizer (learning rate $\alpha = 1e^{-3}$, decay rate

for first moment $\beta_1 = 0.9$, decay rate for second moment $\beta_2 = 0.999$, constant $\varepsilon = 1e^{-8}$). The

autoencoder was implemented in Python using the PyTorch package (Paszke et al., 2019).

Accelerometer data was included by performing the root sum of squares across all three axes.

We then found the variance of the values within each time bin and applied a logarithmic scale to

better compress the data. Finally, the standard deviation was normalized to that of the encoded

dimension with the largest variance. The processed accelerometer data was included as an

additional dimension in the lower dimensional representation (i.e. 33$^{\text{rd}}$ dimension).

To classify the data, we used k-means clustering with an assumption of 4 centroids. Data points

of the lower dimensional representation within the 90$^{\text{th}}$ percentile of pairwise Euclidean distance

were initially used for finding the centroids of clusters to avoid the influence of outliers. Each

data point was assigned to the centroid with the shortest Euclidean distance.

After clustering, each group of records was assigned to one of four states – 1) awake and moving

(Move) 2) awake and at rest (Rest), 3) rapid-eye movement (REM) sleep, 4) non-REM (NREM)

sleep – by assessing the average accelerometer value and average normalized PSD for each

cluster. First, the cluster with highest average acceleration was assigned to be Move, then the

cluster with highest average delta power (0.5 to 4 Hz) was assigned be NREM, then the cluster

with higher average beta power (15 to 30 Hz) was assigned to be Rest, and the remaining cluster

was assigned to be REM.

To include a temporal aspect and smooth any outliers we performed a majority filter on the

classification. For each time-bin, the majority state across 2 time-bins before to 2 time-bins after

(±16 seconds) was considered to be the current state. Ties were resolved by keeping the original

classification, or by random choice if the original classification was not part of the tie.

Validation of classification

Two EOG electrodes (one dorsal and one lateral) were simultaneously recorded from with the

Neurochip3 in experiments with Monkey J. EOG signals were extracted by subtracting the lateral

electrode signal from the dorsal electrode signal and then applying a band-pass filter between 1

and 20 Hz with a zero-phase second-order Butterworth filter. We performed in-booth

experiments with flashes of lights guiding the monkey's gaze to ensure we were properly

capturing eye movements.

As the Neurochip3 is mounted to the monkey's heads, we performed overnight recordings with

the Microsoft Kinect in conjunction to ensure there were no large discrepancies between the on-

board accelerometer and whole-body movements (Libey & Fetz, 2017). Movements with the

Kinect was calculated as the absolute difference between each frame of the infrared depth-

finding camera and the previous frame. We captured frames as quickly as possible with the

processing overhead, around 30 frames per second.

Further validation was performed with k-fold cross-validation with k=20. Each dataset was split

into 20 random groups; classification was performed on 19 of the 20 groups, and the final group

was classified using the centroids from the classification k-means. The training error was

calculated as the difference in classification within the 19 groups used for training the

classification, and the test error as the difference in classification within the final group used to

test the classification. Cross-validation was bootstrapped 50 times for each session to minimize

variability that can potentially be introduced by the random sampling.

Coherence

Magnitude-squared coherence was used to calculate synchrony within the same frequencies:

$$C_{xy} = \frac{|P_{xy}(f)|^2}{P_{xx}(f)P_{yy}(f)} \tag{1}$$

where C_{xy} is the coherence between x and y, $P_{xy}(f)$ is the cross-spectral density between x and

y, and $P_{xx}(f)$ and $P_{yy}(f)$ are the spectral densities of x and y respectively. Coherence was

calculated between all combinations of channel pairs at every 0.1 Hz intervals.

Cross-frequency phase-amplitude coupling

To calculate cross-frequency phase-amplitude coupling we used mean vector length (MVL)

(Canolty et al., 2006) (Figure 1C). The LFP was first filtered into 2 frequency bands of interest

within the 6 frequency bands of interest – 1) delta (0.5-4 Hz), 2) theta (4-8 Hz), 3) alpha (8-12

Hz), 4) beta (15-30 Hz), 5) low gamma (30-70 Hz), and 5) high gamma (70-120 Hz) – using a

zero-phase second-order Butterworth filter. We then calculated the analytic signal, H, of each

band using the Hilbert transform:

$$H(x(t)) = \frac{1}{\pi} P \int_{-\infty}^{+\infty} \frac{x(\tau)}{t - \tau} d\tau \tag{2}$$

199   where P is Cauchy principal value. The phase of the complex valued analytic signal is the

200   instantaneous phase at time t, and the magnitude of the analytic signal is the instantaneous

201   amplitude. The mean vector was then calculated by:

$$MVL = \left| \frac{1}{n} \sum_n r \, exp \, (i\phi) \right| \tag{3}$$

where r is the instantaneous magnitude of the higher frequency band and ϕ is the instantaneous

phase of the lower frequency band. ϕ ranges from 0 to 2π radians where 0 is the peak and π the

trough of oscillations. This calculation creates a vector for each sample in time with the phase

and amplitude of the lower and higher frequency bands, respectively. The magnitude of the

average of those vectors, or MVL, measures the strength of synchrony – zero indicates a uniform

distribution in which the vectors "cancel" each other, and higher values indicate the degree of

synchrony.

However, MVL is highly affected by the amplitude and does not have a normalized maximum

value, which makes interpretation of individual values and comparisons of MVL measurements

across different time points difficult. Thus, we additionally calculated the maximum possible

MVL for each state. Instead of using the phase and amplitude that occurs at the same time

sample in Equation 2, we paired the highest amplitudes with the most commonly occurring

phases. Thus, the largest vectors were all in similar directions providing the maximum possible

length of the mean vector. We then normalized the MVL by dividing by the maximum possible

MVL to assess differences in synchrony between behavioral states.

Spike sorting

Spikes were sorted offline using two-window discrimination. The cortical recording was

bandpass filtered between 1000 and 2000 Hz with a first-order Butterworth filter. Then a

negative threshold and two time-delayed windows were manually chosen to capture the trough

and the peak of the spike waveform. All traces crossing the threshold and passing through the

two windows was denoted as a spike.

As the fidelity of spikes may change over such a long period of recording, we additionally

compared the shapes of the first 1000 detected spikes with the last 1000 detected spikes using the

coefficient of determination (CoD):

$$CoD = 1 - \frac{\sum \left(s_{first}(t) - s_{last}(t)\right)^2}{\sum \left(s_{first}(t) - \overline{s_{first}}\right)^2} \tag{4}$$

where s_{first} is a waveform of one of the first spikes, s_{last} is a waveform of one of the last spikes,

and $\overline{s_{first}}$ is the average of the waveform.

We performed pairwise CoD on all first and last 1000 instances of each spike and compared

them to the pairwise CoD between the first 1000 instances of the spike and the last 1000

instances of a different, randomly chosen spike. If the distribution of the CoD comparing the

same spike was significantly higher than the distribution between the spike and another spike it

was considered to be consistent overnight. Additionally, to ensure we were not capturing

multiple neurons with a similar waveform, we manually assessed the autocorrelograms for each

spike to ensure the presence of a refractory period and proper distribution of inter-spike intervals.

Phase-locking value

We calculated the phase-locking value (PLV) to assess the strength of synchronization of spike

timing to phases of oscillations in specific frequency bands from LFPs recorded on the same

channel (Figure 1D). First, the LFP was filtered into a frequency band of interest. The

instantaneous phase of the lower frequency band was calculated as described above. We then

calculated the PLV:

$$PLV = \frac{1}{n}\left|\sum_n exp\ (i\phi)\right| \tag{5}$$

where ϕ is the instantaneous phase at spike times. The PLV effectively converts each phase into

a unitary vector and finds the average of all the vectors. The magnitude of the resulting vector

determines the synchrony of the phases. A value of zero, similar to MVL, indicates a uniform

distribution, or no synchrony. A value of one indicates that all phases are equal, or perfect

synchrony.

The average phase of spike timing was found using the circular mean:

$$average\ phase = arg\left(\sum_n exp\ (i\phi)\right) \tag{6}$$

where arg is the argument of a complex number, or the angle between the positive real axis and

the complex number in the complex plane.

Statistical analysis

We used Friedman's test to compare values between the four states due to the non-parametric

nature of the distribution and because we were sampling the same spikes and LFPs across each

state. Tukey's honest significance test was used post hoc to determine significant pairwise

differences.

Results

Classifying sleep states

Sleep states were classified using the local field potential (LFP) and accelerometer data collected with the Neurochip3 for successive 8-second time-bins using an autoencoder for dimensionality reduction and k-means clustering (Figure 2A, *Classifying sleep states* in *Methods and materials*). We assumed 4 clusters – awake and moving (Move), awake and at rest (Rest), REM sleep (REM) and non-REM sleep (NREM). Each cluster was assigned to the state depending on the average accelerometer value and features in the average power spectral density (PSD).

Figure 2B shows an example of the classification over a 21-hour period and the corresponding spectral power. During lights-off we found consistent REM cycles in between NREM epochs occurring every 30 minutes to 2 hours, which became longer and more frequent closer to the morning, consistent with previous findings (Aserinsky & Kleitman, 1953; Hsieh et al., 2008; Kripke et al., 1968). NREM was absent when the lights were on in the animal room, but we often found brief periods of REM sleep concurrent with changes in the PSD, which we attributed to naps (Figure 2B, arrow) (Dijk et al., 1987; Moses et al., 1975). We also often found short periods of Rest or sometimes even Move during the night which were attributed to brief awakenings, typical in an average night of sleep.

Averaging the spectral power during each sleep state shows high delta power during NREM sleep, high theta power during REM sleep, and high beta power when the animal was awake, consistent with previous findings (Figure 2C) (Xu et al., 2019). Additionally, raw LFP traces during the states show high-frequency activity during Move, faster oscillatory activity during Rest in the beta range, slower oscillatory activity during REM sleep in the theta and alpha range,

276 and clear K-complexes and sleep spindles along with low frequency activity around delta during

277 NREM sleep (Figure 2D).

<u>Confirming sleep state classification</u>

As our classification scheme was unsupervised, we simultaneously recorded electrooculography

(EOG) signals to validate the identified sleep states (Figure 3A). Eye movements detected with

the EOG were clearly associated with identified sleep states; EOG variance was very high when

the animal was awake, very low during NREM sleep, and elevated during REM sleep (Figure

3B). This relationship was present even during very brief windows of detected awakening during

the night and naps during the day, indicating high accuracy of state classification.

The accelerometer used for classification was on the Neurochip3 which was mounted on the

animals' head. To confirm the accuracy of detected movements, we additionally recorded the

monkeys' movements overnight with a Microsoft Kinect. Compared to the movement values

extracted with the Kinect, the accelerometer values were smaller when the animal was awake but

larger when the animal was asleep (Figure 3C). This is likely due to body movements that are

independent of the head (i.e., isolated limb movements) not being as accurately recorded in the

accelerometer. On the other hand, when the animal is laying down at night most movements

involve the head. However, these differences were minor, and the two signals were comparable

throughout the recording duration.

Finally, we also performed k-fold cross-validation to verify that our classification method was

consistent and robust. We used k=20 and carried out 50 repetitions per session and found the

average test and training error to be less than 4%, or around 40 total minutes (Figure 3D). The

similar but low test and training errors suggest the classification method had low variance (i.e.

not overfit) and high consistency. As a result, the classification was deemed sufficient for the

aims of this study.

Changes in LFP dynamics

The average spectral power per state for each animal is shown in Figure 4A. Delta power was

high during NREM sleep due to the presence of slow waves. There was a beta peak during the

wake states, though the specific frequency range differed between the animals: 25-30 Hz for

Monkey K and 15-20 Hz for Monkey J. REM sleep showed slightly higher theta and alpha

power compared to the wake states. However, the differences in power between each state were

not as large as expected, which may be due to different sites being recorded for different

recording sessions resulting in a broader average. Although all sites were within the motor cortex

and within a 4 mm wide square window, there may still be large differences in LFP spectral

power across different days, especially during different behavioral states.

Average pairwise coherence per state for each animal is shown in Figure 4B. The features were

very similar to those show in the power, but the differences were magnified, likely due to

coherence showing synchrony between pairs of channels and amplifying active oscillatory

signaling over baseline spectral density. In addition to the large delta peak during NREM, there

was a much clearer difference between REM and the wake states, including a more distinct beta

peak.

Cross-frequency phase-amplitude coupling

The phase of lower frequency bands has been observed to be coupled with the amplitude of

higher frequency bands, thought to reflect coordination between brain networks (Canolty &

Knight, 2010; Jensen & Colgin, 2007). We explored the coupling between every pair of low

frequency band phase to high frequency band amplitude to determine whether there were state-

dependent changes. We first plotted the average z-scored spectral power of higher frequencies at

different phases of each frequency band (Figure 5). There was clear coupling between different

phases and frequencies, the strength and specific phase of which changed depending on the state.

To quantify the degree of coupling we calculated the normalized mean vector length (nMVL).

Figure 6 shows the distribution of nMVL for each pair of frequency bands during each state.

There were large state-dependent fluctuations in nMVL distributions, most notably during

NREM. There was significantly stronger coupling between delta and theta phases and low and

high gamma amplitudes (delta/theta-gamma) during NREM. Interestingly, delta/theta-high

gamma coupling was also elevated during Move. In contrast, delta/theta-alpha and delta-beta

couplings were significantly lower during NREM.

Other large differences include significantly higher theta-beta coupling during REM and higher

alpha-high gamma coupling during Move. Additionally, beta and low gamma phase to higher

frequency coupling is higher during sleep states than awake states.

To further characterize the coupling, we explored the distribution of the mean vectors to

determine whether the phases aligned with the highest amplitudes were consistent. Figure 7

shows the distribution of mean vector phases for each pair of frequency bands during each state.

There was very high consistency in phases during NREM across most pairs of frequency bands.

Theta-beta coupling was very consistent during REM, further validating the high nMVL

distribution. Beta and low gamma phase to higher frequency coupling was consistent in the

phase of coupling, typically between π and $5\pi/4$.

Spike sorting

Spikes were manually sorted using two-window discrimination (Figure 8A). As spike waveform

shape and magnitude can fluctuate over time due to electrode drift, we calculated the coefficient

of determination (CoD) for all pairs of the first and last 1000 instances of each spike to ensure

we tracked the same spike throughout the recording duration (Figure 8B). A higher distribution

of CoD between the first and last 1000 instances of the same spike compared to the CoD

between different spikes confirmed the consistency of the spike waveform. We tracked a total of

193 spikes (121 in Monkey K, 92 in Monkey J) that satisfied our conditions across all sessions.

The firing rates of spikes overnight were strongly and consistently modulated by different

identified sleep states, especially during REM cycles during the night, further demonstrating the

accuracy of the classification method (Figure 8C). Firing rates were consistently lower with

deeper sleep.

Additionally, spikes were designated to be regular firing or fast firing depending on the location

of the inter-spike interval distribution peak during Move (Figure 8D). If the peak occurred before

10 ms the spike was designated to be fast spiking (FS), otherwise it was designated to be regular

spiking (RS) (84 FS spikes, 109 RS spikes). Assessing the differences in firing rates depending

on state showed that both RS and FS spike rates significantly decrease with deeper sleep (Figure

8E).

Changes in spiking dynamics

To determine how spiking patterns changed between different sleep states we analyzed changes

in the average ISI distributions of RS and FS spikes. The raw ISI distributions showed an overall

decrease on most ISIs with deeper sleep, reflecting the diminished firing rate, but also a slight

change in the timing of the distribution, especially for RS spikes (Figure 9A).

To quantify the differences, we calculated the difference in ISI distributions from Move (Figure

9B, top). FS spikes had a consistent decrease in shorter latency ISIs with deeper sleep, but RS

spikes showed a large increase in short latency ISIs at around 5 ms (arrow). To determine if there

were any changes in larger ISIs, we plotted the distributions against frequency (Figure 9B,

bottom). Both FS and RS spikes showed a peak between 7 and 10 Hz during NREM and a peak

between 20 and 25 Hz during Rest and REM.

We also assessed how spiking activity of one spike affects other spikes by plotting the peri-event

time histogram (PETH) based on changes in individual spike firing rates. For each spike, we

found times when the firing rate was over 3 times the standard deviation from the average and

calculated the PETH for each other spike in the recording from -1 to 1 second around the

increased firing rate. Figure 9C shows the average PETH of each channel pair for each state.

There was consistently high spiking activity immediately following the increase of any spike's

firing rate, as expected from the co-modulation of firing rates seen overnight. However, the peak

of activity was significantly higher during Move and NREM, being almost identical between the

two states, and the lowest during REM. The widths of the peaks were between 150 and 200 ms

and was comparable between all states.

Spike field relationships

Spikes have been shown to be synchronized to various frequency bands, particularly in the motor

cortex (Buzsáki et al., 2012; Murthy & Fetz, 1996b). We first analyzed the phase-amplitude

distribution normalized by amplitude of each LFP band during spike timings (Figure 10). There

was clear synchrony with spikes during NREM sleep with all LFP bands, though at varying

phases. During NREM spikes typically occurred at the negative phase of delta, beta, and gamma

activity, and at intermediate phases for theta ($\pi/2$) and alpha ($3\pi/4$). The synchrony was

particularly apparent during high amplitudes, suggesting spikes are synchronized to active

frequency band activity rather than epiphenomena arising from periodicity of spike firing

patterns. During both awake states and REM sleep spikes were synchronized to the trough of

beta and gamma frequencies, though less than during NREM sleep. Spikes during Move were

also potentially synchronized to the delta band. All other combinations did not show apparent

coordination between LFP band phase and spike timing.

To quantify the strength of these relationships, we calculated the phase-locking value (PLV) for

each spike during each state for all LFP bands. Figure 11 shows the RS spikes' distribution of

locked phases as well as the PLV for each frequency band during each state. Spikes were more

synchronized to delta during both Move and NREM, validated by the consistency in the locked

phases (around $\pi/2$ and $\pm\pi$, respectively). For every other frequency band, there was stronger

synchrony during deeper sleep states, though the differences were often not statistically

significant. The phases were extremely consistent for beta and higher frequency bands, but

variable for lower frequency bands during states with lower PLVs.

Figure 12 shows the FS spikes' distribution of locked phases and the PLV for each frequency

band during each state. One large difference from RS spikes was the synchronization with delta

during Move, which was not significantly larger compared to Rest or REM. Otherwise, there was

a much stronger relationship between increase in PLV and deeper sleep states, being statistically

significant for each frequency band. In addition, the phase of synchrony to the low and high

gamma bands was earlier in the phase for FS spikes compared to RS spikes.

Discussion

State classification

Although a guideline to standardize sleep staging was published after the discovery of sleep stages (Rechtschaffen & Kales, 1968), staging has remained an inconsistent and often arduous task. Criticisms of and modifications to the guideline (Himanen & Hasan, 2000; Silber et al., 2007) as well as various automated methods (Hamida & Ahmed, 2013; Penzel & Conradt, 2000; Sun et al., 2020) have been developed, but each approach differs in the set of defined states and the type and amount of data required. In addition, studies on sleep typically use electroencephalogram (EEG) LFP recordings, which are distinctly different from intracortical LFPs due to their lower spatial specificity and the complex frequency and phase filtering of bone and tissue, especially at higher frequencies (Buzsáki et al., 2012; Michel et al., 2004; Sejnowski et al., 2014). Beyond these difficulties, manual scoring is also compromised by high inter-scorer variability (Himanen & Hasan, 2000; Younes et al., 2016).

As a result, we developed our own classification methodology to distinguish between different sleep states. The process was relatively simple, utilizing basic dimensionality reduction and clustering on processed LFP PSDs and accelerometer values, but was remarkably consistent. We used several metrics to confirm the method – EOG recordings, concurrent recordings with the Kinect to track movement, k-fold cross validation, and changes in spike rates; they all indicated accurate and robust classification. One explanation may be that sleep states can vary between different regions of the brain (Mascetti, 2016; Siclari & Tononi, 2017). Our focus on a small region (4x4 mm) of the motor cortex and the high spatial specificity of intracortical LFPs may

have produced more robust results compared to those based on EEG recordings from across the

brain.

Our method could potentially be improved by using a dimensionality reduction method with

higher complexity, such as a stacked autoencoder (Tsinalis et al., 2016), different clustering

methods, such as hierarchical clustering (Gerla et al., 2019), and by including the EOG

recordings or other biophysical measurements as additional dimensions of data (Khalighi et al.,

2013). However, developing a flawless classification method was not within scope of the study

and thus the method presented was deemed to be sufficiently accurate and of practical use.

LFP and spike dynamics during different sleep states

The average PSD of LFPs during each state displayed features similar to those reported

previously (Rechtschaffen & Kales, 1968; Silber et al., 2007; Xu et al., 2019). We found high

delta power during NREM sleep, as delta is elevated across the brain during all stages of NREM.

Beta power was high during Move and Rest, as we were recording from the primary motor

cortex, and beta oscillations are highly relevant in motor control (Engel & Fries, 2010; Khanna

& Carmena, 2017). Finally, we found high theta power during REM sleep. The role of theta band

in the cortex is still obscure, but theta in the hippocampus is also consistently high during REM

(Boyce et al., 2016). We found similar features in the pairwise magnitude-squared coherence. As

coherence shows synchrony between two signals, it effectively emphasizes coordinating activity

from the baseline PSD.

We also observed decreased spiking activity with deeper sleep, similar to a previous study (Xu et

al., 2019). In addition, the firing patterns of spikes changed due to sleep state – RS neurons had

increased high frequency activity during NREM compared to Move, and all neurons had an

increase low frequency activity during Rest, REM, and NREM compared to Move. The increase

in beta frequency activity of neurons during REM and Rest can be explained by the fact that beta

is seen as a "resting" rhythm in the primary motor cortex (Engel & Fries, 2010). RS neurons are

likely to be excitatory pyramidal cells and FS neurons are likely to be inhibitory interneurons

(McCormick et al., 1985), which may explain the higher "bursty" firing pattern of RS neurons

during NREM sleep, potentially due to reactivation of cortical circuitry activated during the day.

This is further substantiated by the PETH of spikes to increases in firing rate showing similarly

high coordination between spikes during Move and NREM compared to Rest or REM.

Cross-frequency coupling and spike-LFP synchrony

Cross frequency coupling shows that the higher frequency band modulates with the lower

frequency band. Such synchrony reflects coordination of local networks operating on shorter

time scales to distributed circuits synchronized at longer time scales; this could potentially play a

role in neural computation of attention, learning, and memory (Canolty & Knight, 2010; Jensen

& Colgin, 2007). Various measures have been proposed to quantify phase-amplitude coupling;

compared to other proposed measures the MVL introduced by Canolty et al. (2006) has been

shown to be accurate, the most sensitive to modulations in coupling strengths, and ideal for high

quality signals over a long recording epochs (Canolty et al., 2006; Hülsemann et al., 2019;

Onslow et al., 2011; Tort et al., 2010).

Spike-LFP synchrony provides another measure of synchrony between the single neurons and

the composite synaptic activity of the local population (Buzsáki et al., 2012; Murthy & Fetz,

1996b; Okun et al., 2010). The magnitude indicates the strength of synchrony whereas the phase

can provide insights into the timing of the spikes relative to the local population.

Most instances of cross-frequency phase-amplitude coupling and spike-LFP synchrony were

stronger for deeper sleep states. This may be due to asynchronous activity during Move being

driven by functional local circuitry (i.e. generating movement) whereas activity during resting

and sleep states are more attuned to baseline macroscopic rhythmic activity, potentially driven by

homeostatic plasticity (Tononi & Cirelli, 2014).

Delta and high gamma

One noticeable deviation from deeper sleep resulting in stronger synchrony was delta phase to

high gamma amplitude (delta-high gamma) coupling, which was significantly higher during

Move compared to Rest or REM. Delta oscillations may arise in the thalamus or the cortex and

are elevated during NREM. Delta has also been linked to various perceptual, sensorimotor, and

cognitive operations, potentially suppressing activity not necessary to the task at hand (Güntekin

& Başar, 2016; Harmony, 2013). In contrast, high gamma band LFP is representative of local

spiking activity, often strongly correlated with action potentials, and can also reflect neuronal

synchrony (Ray et al., 2008; Ray & Maunsell, 2011).

Evidence suggests the strength of delta-high gamma coupling is modulated by dopamine in both

the prefrontal cortex and the cortico-basal ganglia network (Andino-Pavlovsky et al., 2017;

Güntekin & Başar, 2016). Delta-high gamma coupling has also previously been shown to be

present during sleep in both the hippocampus and the neocortex (Clemens et al., 2009; Takeuchi

et al., 2015), but the underlying mechanisms are still unclear. Andino-Pavlovsky et al. (2017)

showed high delta-high gamma coupling during slow wave sleep in the rodent prefrontal cortex,

and Takeuchi et al. (2015) showed strong delta-high gamma coupling during REM and several

NREM substages in the primate hippocampus (Andino-Pavlovsky et al., 2017; Takeuchi et al.,

2015). However, neither study pursued the state-dependent comparisons, and their broad

definition of gamma (>25 Hz) makes independent interpretation difficult. Specific comparisons

of delta-high gamma coupling during different sleep states have not been reported.

Spike synchrony to delta band sheds additional light on the delta-high gamma relationship. In

our study, only RS neurons showed higher delta-high gamma coupling during Move, and the

phase of synchrony was consistently at the falling edge of the wave (Figure 11). During NREM

both RS and FS spikes were synchronized to delta, but the phase of synchrony was immediately

after the trough of the wave. These results suggest delta as a coordinating signal that reactivates

cortical circuitry during sleep for consolidation (Gulati et al., 2017; Ramanathan et al., 2015; Xu

et al., 2019); spiking activity during the day occurs before delta is the most depolarized,

potentially driving delta activity, and delta during NREM has the highest depolarization right

before spiking activity, which may reactivate the neural circuitry. Such coordination is also

reflected in the cross-frequency coupling observed in our study.

Theta in the motor cortex and REM sleep

Theta oscillations have often been observed during REM sleep, and have also been shown to

coordinate hippocampal place cell activity during active exploration (Buzsáki & Moser, 2013;

Cantero et al., 2003). In the cortex, modulation of theta has been most prominently seen with

theta burst stimulation (TBS) – bursts of gamma frequency stimulation delivered at theta

intervals, typically with TMS over the motor region. TBS has been shown to increase the

excitability of the motor system, observed through increased motor evoked potentials, potentially

by mimicking the coupling of theta and gamma oscillations (Cárdenas-Morales et al., 2010;

Huang et al., 2005; Suppa et al., 2016). Additional evidence shows that strengthening theta-high

gamma coupling with transcranial alternating current stimulation (tACS) increases motor

learning (Akkad et al., 2021). In agreement with these previous findings, our results show much

higher theta-high gamma coupling during Move.

Theta has also been shown to be coupled to low gamma. A wealth of recent cross-frequency

coupling literature has reported theta-low gamma coupling in both hippocampus and the

neocortex, showing modulations dependent on task performance, cognitive engagement, and

memory formation (Buzsáki & Wang, 2012; Canolty et al., 2006; Lisman & Jensen, 2013).

Although stronger theta-low gamma coupling has also been observed during REM sleep, it was

absent in human studies, potentially due to hippocampal theta oscillations in humans being

slower than those in rodents (Cantero et al., 2003; Clemens et al., 2009; Jacobs, 2014). Similar to

the human studies, we did not find significant theta-low gamma coupling during REM, but we

also did not find high delta-low gamma coupling.

Instead, we found very high theta-beta coupling during REM sleep. Theta-beta coupling is not a

common topic of study; there is some evidence that suggests it plays a role in working memory

and decision making, but the literature is sparse and focuses on the frontal lobe (Axmacher et al.,

2010; Cohen et al., 2009; Liang et al., 2021). In our results, the maximum beta amplitude occurs

right before the peak of theta (Figure 7), which leads to minimum beta occurring right after the

trough of theta. If beta is a regulating, inhibitory rhythm in the motor cortex, perhaps theta is

increasing the excitability of the cortex by minimizing beta, and thus inhibition. This may lead to

more effective memory consolidation often seen during REM sleep, relevant to motor skill

learning. However, more research is needed to confirm these conclusions.

Concluding Comments

We analyzed spike and LFP activity in the primate motor cortex during different behavioral

states to better understand the role of sleep and various LFP frequency bands. We observed an

increase in burst firing and high coordination between spikes during NREM, consistent with

previous findings suggesting reactivations of cortical circuity during NREM. High cross-

frequency phase-amplitude coupling between delta and high gamma when the animal is awake

and moving and during non-REM sleep, as well as spike-field synchrony with delta band LFP,

suggest that delta may encode and subsequently drive these reactivations. Modulations in theta to

low gamma phase-amplitude coupling, commonly observed in rodents, was not observed, similar

to previous human studies. Instead, we observed high theta to beta coupling during REM,

potentially driving motor learning by increasing the excitability of the cortex. We also observed

high theta to high gamma coupling during Move, potentially reflecting coordination with

hippocampal place cells.

References

Akkad, H., Dupont-Hadwen, J., Frese, A., Tetkovic, I., Barrett, L., Bestmann, S., & Stagg, C. J. (2021). Increasing motor skill acquisition by driving theta-gamma coupling. *ELife*, 1–15. https://doi.org/10.7554/eLife.67355

Andino-Pavlovsky, V., Souza, A. C., Scheffer-Teixeira, R., Tort, A. B. L., Etchenique, R., & Ribeiro, S. (2017). Dopamine modulates delta-gamma phase-amplitude coupling in the prefrontal cortex of behaving rats. *Frontiers in Neural Circuits*, *11*(May). https://doi.org/10.3389/fncir.2017.00029

Arbune, A. A., Popa, I., Mindruta, I., Beniczky, S., Donos, C., Daneasa, A., Mălîia, M. D., Băjenaru, O. A., Ciurea, J., & Barborica, A. (2020). Sleep modulates effective connectivity: A study using intracranial stimulation and recording. *Clinical Neurophysiology*, *131*(2), 529–541. https://doi.org/10.1016/j.clinph.2019.09.010

Aserinsky, E., & Kleitman, N. (1953). Regularly Occurring Periods of Eye Motility, and Concomitant Phenomena, during Sleep. *Science*, *118*(3062), 273–274.

Axmacher, N., Henseler, M. M., Jensen, O., Weinreich, I., Elger, C. E., & Fell, J. (2010). Cross-frequency coupling supports multi-item working memory in the human hippocampus. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(7), 3228–3233. https://doi.org/10.1073/pnas.0911531107

Bellesi, M., Riedner, B. A., Garcia-Molina, G. N., Cirelli, C., & Tononi, G. (2014). Enhancement of sleep slow waves: Underlying mechanisms and practical consequences. *Frontiers in Systems Neuroscience*, *8*(October), 1–17. https://doi.org/10.3389/fnsys.2014.00208

Boyce, R., Glasgow, S. D., Williams, S., & Adamantidis, A. (2016). Sleep research: Causal evidence for the role of REM sleep theta rhythm in contextual memory consolidation. *Science*, *352*(6287), 812–816. https://doi.org/10.1126/science.aad5252

Buzsáki, G., Anastassiou, C. A., & Koch, C. (2012). The origin of extracellular fields and currents-EEG, ECoG, LFP and spikes. *Nature Reviews Neuroscience*, *13*(6), 407–420. https://doi.org/10.1038/nrn3241

Buzsáki, G., & Moser, E. I. (2013). Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nature Neuroscience*, *16*(2), 130–138. https://doi.org/10.1038/nn.3304

Buzsáki, G., & Wang, X.-J. (2012). Mechanisms of Gamma Oscillations. *Annual Review of Neuroscience*, *6*(9), 2166–2171. https://doi.org/10.1021/nl061786n.Core-Shell

Canolty, R. T., Edwards, E., Dalal, S. S., Soltani, M., Nagarajan, S. S., Kirsch, H. E., Berger, M. S., Barbare, N. M., & Knight, R. T. (2006). High gamma power is phase-locked to theta oscillations in human neocortex. *Science*, *313*(5793), 1626–1628. https://doi.org/10.1126/science.1128115

Canolty, R. T., & Knight, R. T. (2010). The functional role of cross-frequency coupling. *Trends in Cognitive Sciences*, *14*(11), 506–515. https://doi.org/10.1016/j.tics.2010.09.001

Cantero, J. L., Atienza, M., Stickgold, R., Kahana, M. J., Madsen, J. R., & Kocsis, B. (2003). Sleep-Dependent θ Oscillations in the Human Hippocampus and Neocortex. *Journal of Neuroscience*, *23*(34), 10897–10903. https://doi.org/10.1523/jneurosci.23-34-10897.2003

Cárdenas-Morales, L., Nowak, D. A., Kammer, T., Wolf, R. C., & Schönfeldt-Lecuona, C. (2010). Mechanisms and applications of theta-burst rTMS on the human motor cortex. *Brain Topography*, *22*(4), 294–306. https://doi.org/10.1007/s10548-009-0084-7

Clemens, Z., Weiss, B., Szucs, A., Eross, L., Rásonyi, G., & Halász, P. (2009). Phase coupling between rhythmic slow activity and gamma characterizes mesiotemporal rapid-eye-movement sleep in humans. *Neuroscience*, *163*(1), 388–396. https://doi.org/10.1016/j.neuroscience.2009.06.044

Cohen, M. X., Elger, C. E., & Fell, J. (2009). Oscillatory activity and phase-amplitude coupling in the human medial frontal cortex during decision making. *Journal of Cognitive Neuroscience*, *21*(2), 390–402. https://doi.org/10.1162/jocn.2008.21020

Dijk, D. J., Beersma, D. G. M., & Daan, S. (1987). EEG Power Density during Nap Sleep: Reflection of an Hourglass Measuring the Duration of Prior Wakefulness. *Journal of Biological Rhythms*, *2*(3), 207–219. https://doi.org/10.1177/074873048700200304

Engel, A. K., & Fries, P. (2010). Beta-band oscillations-signalling the status quo? *Current Opinion in Neurobiology*, *20*(2), 156–165. https://doi.org/10.1016/j.conb.2010.02.015

Fernandez, L. M. J., & Lüthi, A. (2020). Sleep spindles: Mechanisms and functions. *Physiological Reviews*, *100*(2), 805–868. https://doi.org/10.1152/physrev.00042.2018

Gerla, V., Kremen, V., Macas, M., Dudysova, D., Mladek, A., Sos, P., & Lhotska, L. (2019). Iterative expert-in-the-loop classification of sleep PSG recordings using a hierarchical clustering. *Journal of Neuroscience Methods*, *317*(February), 61–70. https://doi.org/10.1016/j.jneumeth.2019.01.013

Gulati, T., Guo, L., Ramanathan, D. S., Bodepudi, A., Ganguly, K., Service, R., Francisco, S., Affairs, V., Francisco, S., Francisco, S., Francisco, S., Affairs, V., Francisco, S., & Francisco, S. (2017). Neural reactivations during sleep determine network credit assignment. *Nature Neuroscience*, *20*(9), 1277–1284. https://doi.org/10.1038/nn.4601.Neural

Güntekin, B., & Başar, E. (2016). Review of evoked and event-related delta responses in the human brain. *International Journal of Psychophysiology*, *103*, 43–52. https://doi.org/10.1016/j.ijpsycho.2015.02.001

Hamida, S. T. Ben, & Ahmed, B. (2013). Computer based sleep staging: Challenges for the future. *2013 7th IEEE GCC Conference and Exhibition, GCC 2013*, 280–285. https://doi.org/10.1109/IEEEGCC.2013.6705790

Harmony, T. (2013). The functional significance of delta oscillations in cognitive processing. *Frontiers in Integrative Neuroscience*, *7*(DEC), 1–10. https://doi.org/10.3389/fnint.2013.00083

Himanen, S. L., & Hasan, J. (2000). Limitations of Rechtschaffen and Kales. *Sleep Medicine Reviews*, *4*(2), 149–167. https://doi.org/10.1053/smrv.1999.0086

Hsieh, K. C., Robinson, E. L., & Fuller, C. A. (2008). Sleep architecture in unrestrained rhesus monkeys (Macaca mulatta) synchronized to 24-hour light-dark cycles. *Sleep*, *31*(9), 1239–1250.

Huang, Y. Z., Edwards, M. J., Rounis, E., Bhatia, K. P., & Rothwell, J. C. (2005). Theta burst stimulation of the human motor cortex. *Neuron*, *45*(2), 201–206. https://doi.org/10.1016/j.neuron.2004.12.033

Huber, R., Ghilardi, M. F., Massimini, M., & Tononi, G. (2004). Local sleep and learning. *Nature*, *430*(6995), 78–81. https://doi.org/10.1038/nature02663

Hülsemann, M. J., Naumann, E., & Rasch, B. (2019). Quantification of phase-amplitude coupling in neuronal oscillations:comparison of phase-locking value, mean vector length, modulation index, and generalized-linear-modeling-cross-frequency-coupling. *Frontiers in Neuroscience*, *13*(JUN), 1–15. https://doi.org/10.3389/fnins.2019.00573

Jacobs, J. (2014). Hippocampal theta oscillations are slower in humans than in rodents: Implications for models of spatial navigation and memory. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *369*(1635). https://doi.org/10.1098/rstb.2013.0304

Jensen, O., & Colgin, L. L. (2007). Cross-frequency coupling between neuronal oscillations. *Trends in Cognitive Sciences*, *11*(7), 267–269. https://doi.org/10.1016/j.tics.2007.05.003

Khalighi, S., Sousa, T., Pires, G., & Nunes, U. (2013). Automatic sleep staging: A computer assisted approach for optimal combination of features and polysomnographic channels. *Expert Systems with Applications*, *40*(17), 7046–7059. https://doi.org/10.1016/j.eswa.2013.06.023

Khanna, P., & Carmena, J. M. (2017). Beta band oscillations in motor cortex reflect neural population signals that delay movement onset. *ELife*, *6*, 1–31. https://doi.org/10.7554/eLife.24573

Kim, E. Y., Mahmoud, G. S., & Grover, L. M. (2005). REM sleep deprivation inhibits LTP in vivo in area CA1 of rat hippocampus. *Neuroscience Letters*, *388*(3), 163–167. https://doi.org/10.1016/j.neulet.2005.06.057

Kripke, D. F., Reite, M. L., Pegram, G. V., Stephens, L. M., & Lewis, O. F. (1968). Nocturnal sleep in rhesus monkeys. *Electroencephalography and Clinical Neurophysiology*, *24*(6), 581–586. https://doi.org/10.1016/0013-4694(68)90047-3

Liang, W. K., Tseng, P., Yeh, J. R., Huang, N. E., & Juan, C. H. (2021). Frontoparietal Beta Amplitude Modulation and its Interareal Cross-frequency Coupling in Visual Working Memory. *Neuroscience*, *460*, 69–87. https://doi.org/10.1016/j.neuroscience.2021.02.013

Libey, T., & Fetz, E. E. (2017). Open-source, low cost, free-behavior monitoring, and reward system for neuroscience research in non-human primates. *Frontiers in Neuroscience*, *11*(MAY), 1–12. https://doi.org/10.3389/fnins.2017.00265

Lisman, J. E., & Jensen, O. (2013). The Theta-Gamma Neural Code. *Neuron*, *77*(6), 1002–1016. https://doi.org/10.1016/j.neuron.2013.03.007

Marshall, L., Helgadóttir, H., Mölle, M., & Born, J. (2006). Boosting slow oscillations during

sleep potentiates memory. *Nature*, *444*(7119), 610–613.
https://doi.org/10.1038/nature05278

Mascetti, G. G. (2016). Unihemispheric sleep and asymmetrical sleep: Behavioral,
neurophysiological, and functional perspectives. *Nature and Science of Sleep*, *8*, 221–228.
https://doi.org/10.2147/NSS.S71970

McCormick, D. A., Connors, B. W., Lighthall, J. W., & Prince, D. A. (1985). Comparative
electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex.
Journal of Neurophysiology, *54*(4), 782–806. https://doi.org/10.1152/jn.1985.54.4.782

Michel, C. M., Murray, M. M., Lantz, G., Gonzalez, S., Spinelli, L., & Grave De Peralta, R.
(2004). EEG source imaging. *Clinical Neurophysiology*, *115*(10), 2195–2222.
https://doi.org/10.1016/j.clinph.2004.06.001

Moses, J. M., Hord, D. J., Lubin, A., Johnson, L. C., & Naitoh, P. (1975). Dynamics of nap sleep
during a 40 hour period. *Electroencephalography and Clinical Neurophysiology*, *39*(6),
627–633. https://doi.org/10.1016/0013-4694(75)90075-9

Murthy, V. N., & Fetz, E. E. (1996a). Oscillatory activity in sensorimotor cortex of awake
monkeys: synchronization of local field potentials and relation to behavior. *Journal of
Neurophysiology*, *76*(6), 3949–3967.

Murthy, V. N., & Fetz, E. E. (1996b). Synchronization of neurons during local field potential
oscillations in sensorimotor cortex of awake monkeys. *Journal of Neurophysiology*, *76*(6),
3968–3982. http://www.ncbi.nlm.nih.gov/pubmed/8985893

Okun, M., Naim, A., & Lampl, I. (2010). The subthreshold relation between cortical local field
potential and neuronal firing unveiled by intracellular recordings in awake rats. *Journal of
Neuroscience*, *30*(12), 4440–4448. https://doi.org/10.1523/JNEUROSCI.5062-09.2010

Onslow, A. C. E., Bogacz, R., & Jones, M. W. (2011). Quantifying phase-amplitude coupling in
neuronal network oscillations. *Progress in Biophysics and Molecular Biology*, *105*(1–2),
49–57. https://doi.org/10.1016/j.pbiomolbio.2010.09.007

Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z.,
Gimelshein, N., Antiga, L., Desmaison, A., Köpf, A., Yang, E., DeVito, Z., Raison, M.,
Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., … Chintala, S. (2019). PyTorch: An
imperative style, high-performance deep learning library. *Advances in Neural Information
Processing Systems*, *32*(NeurIPS).

Penzel, T., & Conradt, R. (2000). Computer based sleep recording and analysis. *Sleep Medicine
Reviews*, *4*(2), 131–148. https://doi.org/10.1053/smrv.1999.0087

Ramanathan, D. S., Gulati, T., & Ganguly, K. (2015). Sleep-Dependent Reactivation of
Ensembles in Motor Cortex Promotes Skill Consolidation. *PLoS Biology*, *13*(9), 1–25.
https://doi.org/10.1371/journal.pbio.1002263

Ray, S., Crone, N. E., Niebur, E., Franaszczuk, P. J., & Hsiao, S. S. (2008). Neural correlates of
high-gamma oscillations (60-200 Hz) in macaque local field potentials and their potential
implications in electrocorticography. *Journal of Neuroscience*, *28*(45), 11526–11536.
https://doi.org/10.1523/JNEUROSCI.2848-08.2008

Ray, S., & Maunsell, J. H. R. (2011). Different origins of gamma rhythm and high-gamma
activity in macaque visual cortex. *PLoS Biology*, *9*(4).
https://doi.org/10.1371/journal.pbio.1000610

Rechtschaffen, A. (1998). Current perspectives on the function of sleep. *Perspectives in Biology*
and Medicine, *41*(3), 359–390. https://doi.org/10.1353/pbm.1998.0051

Rechtschaffen, A., & Kales, A. (1968). A manual of standardized terminology, techniques and
scoring system for sleep stages of human subjects. *U.S. Public Health Service, U.S.*
Government Printing Office, Washington D.C.

Rembado, I., Zanos, S., & Fetz, E. E. (2017). Cycle-triggered cortical stimulation during slow
wave sleep facilitates learning a BMI task: A case report in a non-human primate. *Frontiers*
in Behavioral Neuroscience, *11*(April), 1–13. https://doi.org/10.3389/fnbeh.2017.00059

Sejnowski, T. J., Churchland, P. S., & Movshon, J. A. (2014). Putting big data to good use in
neuroscience. *Nature Neuroscience*, *17*(11), 1440–1441. https://doi.org/10.1038/nn.3839

Shupe, L. E., Miles, F. P., Jones, G., Yun, R., Mishler, J., Rembado, I., Murphy, R. L.,
Perlmutter, S. I., & Fetz, E. E. (2021). Neurochip3: An autonomous multichannel
bidirectional brain-computer interface for closed-loop activity-dependent stimulation.
Frontiers in Neuroscience, *15*(August), 1–15. https://doi.org/10.3389/fnins.2021.718465

Siclari, F., & Tononi, G. (2017). Local aspects of sleep and wakefulness. *Current Opinion in*
Neurobiology, *44*, 222–227. https://doi.org/10.1016/j.conb.2017.05.008

Siegel, J. M. (2005). Clues to the functions of mammalian sleep. *Nature*, *437*, 1264–1271.

Silber, M. H., Ancoli-Israel, S., Bonnet, M. H., Chokroverty, S., Grigg-Damberger, M. M.,
Hirshkowitz, M., Kapen, S., Keenan, S. A., Kryger, M. H., Penzel, T., Pressman, M. R., &
Iber, C. (2007). The visual scoring of sleep in adults. *Journal of Clinical Sleep Medicine*,
3(2), 121–131.

Sun, S., Jiang, L. P., Peterson, S. M., Herron, J., Weaver, K., Ko, A., Ojemann, J., & Rao, R. P.
N. (2020). Unsupervised Sleep and Wake State Identification in Long-Term
Electrocorticography Recordings. *Annu Int Conf IEEE Eng Med Biol Soc.*, 629–632.

Suppa, A., Huang, Y. Z., Funke, K., Ridding, M. C., Cheeran, B., Di Lazzaro, V., Ziemann, U.,
& Rothwell, J. C. (2016). Ten Years of Theta Burst Stimulation in Humans: Established
Knowledge, Unknowns and Prospects. *Brain Stimulation*, *9*(3), 323–335.
https://doi.org/10.1016/j.brs.2016.01.006

Takeuchi, S., Mima, T., Murai, R., Shimazu, H., Isomura, Y., & Tsujimoto, T. (2015). Gamma
oscillations and their cross-frequency coupling in the primate hippocampus during sleep.
Sleep, *38*(7), 1085-1091G. https://doi.org/10.5665/sleep.4818

Tononi, G., & Cirelli, C. (2014). Sleep and the Price of Plasticity: From Synaptic and Cellular
Homeostasis to Memory Consolidation and Integration. *Neuron*, *81*(1), 12–34.
https://doi.org/10.1016/j.neuron.2013.12.025

Tort, A. B. L., Komorowski, R., Eichenbaum, H., & Kopell, N. (2010). Measuring phase-
amplitude coupling between neuronal oscillations of different frequencies. *Journal of*

Neurophysiology, *104*(2), 1195–1210. https://doi.org/10.1152/jn.00106.2010

Tsinalis, O., Matthews, P. M., & Guo, Y. (2016). Automatic Sleep Stage Scoring Using Time-
Frequency Analysis and Stacked Sparse Autoencoders. *Annals of Biomedical Engineering*,
44(5), 1587–1597. https://doi.org/10.1007/s10439-015-1444-y

Ulrich, D. (2016). Sleep Spindles as Facilitators of Memory Formation and Learning. *Neural*
Plasticity, *2016*. https://doi.org/10.1155/2016/1796715

Vecsey, C. G., Huang, T., & Abel, T. (2018). Sleep deprivation impairs synaptic tagging in
mouse hippocampal slices. *Neurobiology of Learning and Memory*, *154*(March), 136–140.
https://doi.org/10.1016/j.nlm.2018.03.016

Xu, W., de Carvalho, F., & Jackson, A. (2019). Sequential Neural Activity in Primary Motor
Cortex during Sleep. *The Journal of Neuroscience*, *39*(19), 3698–3712.
https://doi.org/10.1523/jneurosci.1408-18.2019

Younes, M., Raneri, J., & Hanly, P. (2016). Staging sleep in polysomnograms: Analysis of inter-
scorer variability. *Journal of Clinical Sleep Medicine*, *12*(6), 885–894.
https://doi.org/10.5664/jcsm.5894

Figures

Figure 1. <u>Experimental design</u>

A. All data was gathered using the Neurochip3 on freely behaving macaques for up to 24 hours.
B. We collected 16 channels of data at 20 kHz and obtained the spectral density of every 8-
second bin to classify behavioral states. **C.** We then calculated the cross-frequency phase-
amplitude coupling of every frequency band pair. The example shows the instantaneous
amplitude (red) of high gamma filtered LFP (black, bottom) increasing at the trough of beta
(black, top). **D.** We additionally sorted single units and found the synchrony of spikes to LFP
bands. The example shows spikes (red) synchronized with the trough of beta filtered LFP.
771

Figure 2. <u>State classification</u>

A. Diagram of classification process. LFP and accelerometer signals are used to train an
autoencoder for dimensionality reduction and subsequent k-means clustering to determine
different sleep states. **B.** Example of classification and spectra over 22 hours of continuous
recording. Brief periods of REM sleep were observed during the day, attributed to naps (arrow).
C. Averaged spectra across each state for the session shown in (B). There is high delta power
during NREM, theta power during REM, and beta power during Move and Rest states. **D.**
Example traces of raw LFP during each state. Spindles (black arrow) and k-complexes (red
arrow) were observed during NREM sleep.

Figure 3. <u>Classification confirmation</u>

A. An example of filtered differential EOG signals and corresponding eye movements. **B.**
Standard deviation of EOG signals overnight with respect to classified states. Note the increase
in large eye movements during REM and waking. **C.** An example of the normalized variance of
accelerometer data and the normalized variance of the Kinect data with respect to classified
states. **D.** Test and training errors with k-fold cross-validation in percent (left) and total minutes
(right).
791

793 **Figure 4. <u>LFP dynamics</u>**

794 **A.** Average spectral power in each state for each animal across all experiments. Shaded regions
795 show standard error. **B.** Average pairwise coherence in each state for each animal across all
796 experiments. Shaded regions show standard error.
797

Figure 5. <u>Cross-frequency phase-power distributions</u>

Distribution of lower frequency band phase and higher frequency band power during each state. The power is z-scored for each frequency.

Figure 6. <u>Normalized MVL distributions</u>

805 Normalized MVL distributions for each lower frequency band phase and higher frequency band
806 amplitude pair during each state. The black boxes show standard box plots with interquartile
807 range and the white dots show median values. The numbers above each group denote
808 significance compared to another state (Friedman's test, $p < 0.05$).
809

 Figure 7. <u>Mean vector phase distributions</u>

812 The distribution of phases of mean vectors for each lower frequency band phase and higher
813 frequency band amplitude pair during each state. The phase of the mean vector shows the phase
814 at which the amplitude of the higher frequency band is the greatest.
815

816
817

Figure 8. Spike sorting

A. An example of a sorted spike. The grey traces show a random sample of 1000 spikes, the black line is the average, the horizontal dashed red line shows the threshold, and the blue vertical lines show the two windows. **B.** An example of comparisons of the pairwise coefficient of determination (CoD) between the first and last 1000 instances of the same spike and the first and last 1000 instances of two different spikes. **C.** Firing rates binned every 60 seconds of four neurons overnight with classified sleep states. The changes in firing rate very closely match the changes in state. **D.** Histogram of the peak of the ISI distribution of each spike. Spikes with peaks earlier than 10 ms were denoted to be fast spiking (FS) and all others denoted to be regular spiking (RS). **E.** Percent difference of firing rate in each state from the average overall firing rate for each spike type. Each state is statistically significantly different from each other state (Friedman's test, $p < 0.05$).

 Figure 9. <u>Changes in spiking patterns</u>

A. Average ISI distributions of identified fast spiking and regular spiking neurons during each
state. **B.** Difference of ISI distributions from Move plotted against time (top) and frequency
(bottom). Note the peak around 5 ms during NREM for regular spiking neurons and the peaks
around 7-10 Hz during NREM sleep and the peaks at around 20 and 25 Hz for REM and Rest
respectively (arrows) for both neuron types. All three peaks were significantly different from 0
(Wilcoxon signed-rank test, p<0.05). **C.** Average normalized PETH triggered from an increase in
firing rate of one spike. Each PETH trace was normalized by subtracting the mean value and
then dividing by the standard deviation.

843   **Figure 10. <u>Phase distributions of LFPs at spike times</u>**

844   Phase distributions normalized for each amplitude during each sleep state for each LFP band.
845

848 **Figure 11. <u>Locked phase and phase locking value distributions for regular</u>**

849 **<u>spiking neurons</u>**

850 Distributions of locked phase (left) and PLVs (right) of regular spiking neurons during each
851 sleep state for each LFP band. The numbers above each group denote significance compared to
852 another state (Friedman's test, $p<0.05$).
853

855   **Figure 12. <u>Locked phase and phase locking value distributions for fast spiking</u>**

856   **<u>neurons</u>**

857   Distributions of locked phase (left) and PLVs (right) of fast spiking neurons during each sleep
858   state for each LFP band. The numbers above each group denote significance compared to
859   another state (Friedman's test, $p < 0.05$).
860