**The reprogrammed mouse heart phosphoproteome in response to the chronic stress of**
**markedly high cardiac-specific adenylyl cyclase type 8 overexpression**

Jia-Hua Qu, Khalid Chakir, Kirill V. Tarasov, Daniel R. Riordon, Maria Grazia Perino, Allwin
Jennifa Silvester, and Edward G. Lakatta *

Laboratory of Cardiovascular Science, Intramural Research Program, National Institute on
Aging, National Institutes of Health, Baltimore, MD 21224, USA.

**Address correspondence to:**

Dr. Edward G. Lakatta*

Laboratory of Cardiovascular Science

Intramural Research Program

National Institute on Aging

National Institutes of Health

251 Bayview Boulevard

Suite 100

Baltimore, MD 21224

USA.

Tel: +1 410-558-8202

lakattae@grc.nia.nih.gov

## Abstract

An appropriate intensity and duration of increased AC-cAMP-PKA-$Ca^{2+}$ signaling is the quintessential component of acute adaptation to cardiac stress. Our previous work demonstrated the existence of an exquisite heart performance and protection profile in response to chronically increased cardiac AC-cAMP-PKA-$Ca^{2+}$ signaling in young adult mice with cardiac-specific overexpression of adenylyl cyclase type 8 (AC8) (TGAC8). Here, we conducted an unbiased phosphoproteomics analysis in order to determine the role of altered protein phosphorylation in the chronic adaptive heart performance and protection profile of left ventricle (LV) of adult TGAC8 at 3-4 months of age. We identified 12,963 phospho-peptides and 3646 phosphoproteins, including 4990 upregulated phosphopeptides and 781 differentially regulated (741 Up and 40 Down) phosphoproteins in TGAC8 vs. WT. Based on differentially regulated phosphoproteins by genotype, we discovered the enrichment of numerous stress response canonical pathways within reprogrammed TGAC8 LV, including protein kinase A (PKA) signaling, PI3K signaling, and AMPK signaling, predicted upstream regulators, e.g., PDPK1, PAK1 and PTK2B, and downstream functions, e.g., cell viability, protein quality control, and metabolism. In addition to PKA, numerous kinases (83) and phosphatases (30) were hyper-phosphorylated in TGAC8 vs. WT. By combining the phosphoproteome with its proteome and with the previously published TGAC8 transcriptome, we determined which cardiac performance and adaptive protection profiles were coordinately regulated at post-translational modification (PTM) (phosphorylation), translational and transcriptional levels. In order to assess the impact of phosphorylation on mRNA transcription, we determined the phosphorylation states of transcriptional factors (TFs) that differed by genotype. Hyper-phosphorylated TFs in TGAC8 were associated with increased mRNA transcription, and immune responses and metabolism pathways. To determine the coordination of proteome and phosphoproteome in reprogramming the TGAC8 LV, we normalized its phosphoproteome to its proteome (by dividing the protein phosphorylation abundance by protein abundance). In functional enrichment analysis of 229 normalized phosphoproteins that were differentially regulated by genotype, enriched IPA canonical pathways in the phosphoproteome of TGAC8 LV before and after normalization showed high consistency. The combination of aforementioned analyses indicated that many stress response signaling pathways, i.e., PI3K/AKT signaling and ubiquitin labeling, were consistently enriched at transcriptional, translational and PTM levels, and that PI3K/AKT signaling was also activated in the TGAC8 LV. Thus, PTM of protein by phosphorylation, per se, contributes immensely to the heart performance and protection profile of TGAC8 LV, and is coordinately regulated with altered transcription and translation in the TGAC8 to activate critical stress response pathways.

**Keywords**

left ventricle, adenylyl cyclase 8, AC-cAMP-PKA-Ca$^{2+}$ signaling, phosphoproteome, proteome, transcriptome, PI3K/AKT signaling pathway.

**Materials and Methods**

**1. Animals**.

The cardiac-specific overexpression of AC8 (TGAC8) mouse, generated by inserting a cDNA coding for human AC8 after the murine α-myosin heavy chain promoter, was a gift from Nicole Defer/Jacques Hanoune, Unite de Recherches, INSERM U-99, Hôpital Henri Mondor, F-94010 Créteil, France. The 12-week old male TGAC mice were used as study cases, and age matched WT littermates, in the C57/BL6 background, were used as controls. All studies were performed in accordance with the Guide for the Care and Use of Laboratory Animals published by the National Institutes of Health (NIH Publication no. 85-23, revised 1996). The experimental protocols were approved by the Animal Care and Use Committee of the National Institutes of Health (protocol #441-LCS-2016).

**2. Isolate and store mouse heart.**

Three-month old male C57 mice (TGAC8 and WT) were euthanized and the heart was isolated. The left ventricle (LV) tissues were extracted and stored in tubes at -80°C.

**3. Mass spectrometry for phosphoproteome and data process.**

Four TGAC8 LVs and five WT LVs were sent to National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda to conduct mass spectrometry using the standard tandem mass tag (TMT) labeling method to obtain the phosphoproteome and total proteome concurrently in the same platform. The raw data were processed in Proteome Discover (ThermoFisher Scientific, software version 2.2). After filtering low quality signal, we remained the medium and high confidence peptides. For total proteins and phosphoproteins, we combined peptides to computed the normalized/standardized expressions with the default parameters. Then we divided the expressions of phosphoproteins to those of total proteins to obtain the normalized phosphoproteome. As the expressions appeared to be near normal distribution, we calculated the fold change by dividing the average value of TGAC8 samples by the average value of WT samples, and calculated p-value using two-sided unpaired Student's t-test.

**4. Analyses of the TGAC8 LV transcriptome.**

The processed transcriptomic data were obtained from our recently published work (ref). The
significantly changed mRNAs (TGAC8 vs. WT, q-value < 0.05) were uploaded into DAVID[2]
(version: 6.8; https://david.ncifcrf.gov/home.jsp) for Gene Ontology (GO) term enrichments, and
were also imported into Ingenuity Pathway Analysis[3] (IPA (QIAGEN, March 2020)) for
upstream kinases and phosphatases prediction. Kinases and phosphatases were selected from the
transcriptome and their changes at mRNA level were displayed to observe their regulation by the
overexpression of AC8.

**5. Analyses of phosphoproteome.**

Both unnormalized and normalized phosphoproteome were analyzed and visualized in PCA,
heatmap, and volcano plots. With the cutoff -log(p-value) > 1.3 and absolute(log2(fold change))
≥ 0.4, we obtained the significantly changed unnormalized and normalized phosphoproteins
(TGAC8 vs. WT) and imported them into IPA for functional analyses respectively, and
compared the canonical pathway enrichments from these two datasets. In IPA, the upstream
kinases were predicted based upon the unnormalized phosphoproteom, and the regulatory
pathways from the predicted upstream kinases to downstream effective signaling pathways were
generated. Molecular and cellular functions were also enriched in IPA. The similar analyses to
those conducted in the TGAC8 LV transcriptome were also done in the TGAC8 LV total
proteome to observe the GO term enrichment, predicted activities of upstream kinases and
phosphatases, and expression of kinases and phosphatases at protein level.

**6. Analyses of phosphopeptides.**

The upregulated phosphopeptides (TGAC8 vs. WT, p-value < 0.05 and log2(fold change) > 0)
were processed in R. After split of peptides with multiple phosphosites into individual peptides
and filtration of ambiguously identified sites, 2264 phosphopeptides were selected. Protein
peptide sequences of full length were downloaded from Uniprot database[1] and matched to the
protein accessions corresponding to processed and selected phosphopeptides. Peptides centered
on the phosphosites and flanked by 6 amino acids on the N and C termini respectively were
extracted from the matched protein peptide sequences, and uploaded into WebLogo[2]
(http://weblogo.threeplusone.com/) and iceLogo[3] (https://iomics.ugent.be/icelogoserver/) for
sequence motif analysis. In WebLogo online analysis, different amino acids were labeled in
different colors representing different chemical properties (refer to figure legend). In iceLogo
online analysis,  a public amphiphilicity index of polar amino acids reported by Mitaku and
colleagues[4] was used as an aid to characterize the amino acid preference at membrane-water
interfaces. In addition, the phosphopeptides of PI3K/Akt substates were summarized and
displayed in stacked bar plots.

## 7. Integrative analyses of transcriptome, proteome and phosphoproteome.

The changes of significantly regulated molecules (TGAC8 vs. WT, q-value or p-value < 0.05) in transcriptome, total proteome, and unnormalized phosphoproteome were displayed in circos plot. In addition, the changes at different level were compared and visualized in correlation scatter plots. The discordant proteins whose changes directions in total proteome and phosphoproteome were opposite were used to analyze the specific regulation at the phosphorylation level by the use of Reactome and GO (Cellular component and biological process) databases.

As to the coordinated regulation, the above mentioned significantly regulated molecules (TGAC8 vs. WT ) in each dataset were imported into IPA for canonical pathway enrichment. PI3K/AKT signaling and protein ubiquitination were two coordinated pathways across all omics. The phosphopeptides of protein substrates of PI3K were extracted from the phosphopeptide dataset and the numbers of changed phosphopeptides (TGAC8 vs. WT) were summarized and visualized in stacked bar plot. The PI3K/AKT signaling pathway were visualized in IPA, and nodes in the pathways were marked in different colors in terms of their changes at different levels.

As kinases and phosphatases are pivotal in phosphorylation, and transcription factors (TFs) are important in transcription regulation, they were extracted from transcriptome, proteome, and phosphoproteome to observe their changes (TGAC8 vs. WT) and deduce their role in the reprogram of phosphoproteome. The upstream transcription regulators including but not only TFs were predicted on the basis of transcriptome and proteome by the utilization of knowledge base in IPA, and their expressions were searched in multiple datasets. Furthermore, TFs that were significantly regulated at phosphorylation level (TGAC8 vs. WT, p-value < 0.05) were defined as significantly changed phosphorylated TFs in this research, and they were uploaded into STRING database[4] (version: 11.0; http://string-db.org/) to generate protein-protein interaction network and enrich KEGG pathways. They were also imported into IPA for canonical pathway, molecular and cellular functions enrichments.

## 8. Data process and graphics production.

Most raw data were precleared and processed using RStudio (version: 1.1.463) in R language (version: 3.5.3), in which the tidyr package and dplyr package were applied. In addition, GraphPad Prism (version: 7), Microsoft Excel (version: 2019) and Adobe Illustrator (version: CC 2019) were also used for statistics and graphics. Venn diagrams were generated in Venny[5] (version: 2.1.0; http://bioinfogp.cnb.csic.es/tools/venny/index.html). Circos plot was generated using Circos software[6] in Perl language.

## Introduction

Cyclic AMP (cAMP) is a pivotal second messenger, which exerts multifaceted effects on most significant physiological processes, such as cardiac contractility, glycogenolysis and lipolysis, hormone release, etc[5]. In mammalians, there are nine transmembrane adenylyl cyclase (AC) (mAC) isoforms (AC1-9) and one soluble AC (sAC, also called AC10) catalyzing the conversion of ATP into cAMP[6], which usually activates the phosphorylation cascade, serving as the central motif of cellular signaling[5,7]. Increased AC-cAMP-PKA-Ca$^{2+}$ signaling is the quintessential component of both acute and chronic cardiac responses.

AC8, the isoform, is highly expressed in brain regions[6,8]. Unlike two dominant cardiac AC isoforms, AC5 and AC6, which are inhibited by intracellular Ca$^{2+}$ and inhibited in heart failure (HF), AC8 is stimulated by Ca$^{2+}$-calmodulin (CM)[9,10] and involved in nervous system functions, like learning and memory[6,8]. Our lab and others have demonstrated that cardiac-specific overexpression of AC8 (TGAC8) markedly increases the heart rate and ejection fraction, similar to acute exercise but chronic and in the absence of corporal exertion, and persists on a 24 hours per day and 7 days per week basis[9,11-13]. Despite this persistent high-level of heart performance driven by markedly elevated level of AC8-cAMP -PKA-Ca$^{2+}$ signaling, which usually proceeds to near-term cardiac dysfunction, the TGAC8 survives for an extended period of time, up to about a year of age without excessive mortality.

It is of utmost importance to unravel how the TGAC8 adapts the heart to this chronic stress, as this knowledge can be applied to the study of cardiac physiology and pathology and even to develop treatments for cardiovascular diseases (CVDs). In this regard, as a first "port-of-call", we had recently compared the LV transcriptome and proteome in TGAC8 and WT littermates. Because PKA signaling pathway was highly enriched in TGAC8 vs. WT, we reasoned that protein phosphorylation will differ between TGAC8 and WT, and the phosphorylation modification will play a crucial role in the remarkable adaptation capacity of the TGAC8 heart in response to markedly increased AC8-cAMP-PKA-Ca$^{2+}$ signaling. We have assessed the concordance of the genotypic differences at the phosphorylation level to some degree in western blot (WB) experiments (Ref to be submitted).

As the body is a quite complex system, regulation at a single level cannot reveal the whole picture in an organism, and multiple levels of regulatory mechanisms are coordinated to defend against stimuli and maintain health. Phosphorylation is one of the most important and most extensively studied post-translational modifications (PTMs) on proteins. Various kinases and phosphatases within cells phosphorylate and dephosphorylate a myriad of proteins respectively to regulate their activities and affect multiple signaling pathways. In the present research, we conducted Tandem Mass Tag (TMT)-labeling MS of left ventricular samples of the three-month old male TGAC8 and WT mice, to unravel the reprogrammed phosphoproteome and proteome of the TGAC8 left ventricle (LV). In addition, by comparing genotypic differences in the proteome and phosphoproteome of the present study with genotypic differences in transcriptome published

212    previously, we sought to determine whether some adaptive mechanisms that emerge in the
213    TGAC8 LV are regulated at transcriptional, translational and PTM (phosphorylation) levels.

## Results

### 1. The TGAC8 reprograms the LV phosphoproteome.

The TMT-labeling quantitative MS identified **12,963 phospho-peptides** (File 1: Sheet 1) and
**3646 phosphoproteins** (File 1: Sheet 2). The principal component analysis (PCA) (Figure 1A)
of the 3646 phosphoproteins in each mouse showed that the TG and WT samples were
distinguished into two groups completely. The heatmap (Figure 1B) showed the intra-group
consistency of upregulated phosphoproteins in TGAC8 compared to WT. Within the cutoff at -
log10(p-value) > 1.3 and absolute value of log2(fold-change(TG/WT)) ≥ 0.04, 781
phosphoproteins were differentially regulated by genotype. Of the 781 phosphoproteins, the
phosphorylation levels of 741 proteins were increased and only 40 were decreased in the TGAC8
vs. WT LV (Figure 1C).

### 2. Functional analyses of the phosphoproteome.

228    We performed the canonical pathway, upstream regulation, and downstream function analyses of
229    the remodeled LV phosphoproteome of TGAC8 using the Ingenuity Pathway Analysis (IPA)
230    software.

### 2.1. Canonical pathway analysis of the remodeled LV phosphoproteome of TGAC8.

To study the mechanism at phosphorylation level, we first performed canonical pathway analysis
using IPA. Top canonical pathways within the phosphoproteome that were statistically
differentially enriched in TGAC8 vs. WT were displayed in the bar plot (Figure 1D and File 2:
Sheet 1). Of these pathways, cardiac hypertrophy signaling (enhanced) is most highly activated
(z-score as high as 4.867) (Figure 1D). The combination of Supplementary Figure 1 and File 2:
Sheet 2 and our recently published work (Ref to be submitted) showed that the phosphorylation
status of numerous molecules, e.g., $Ca^{2+}$ ATPase, RyR2, Ncx, involved in excitation, calcium
cycling, contraction, relaxation and protein synthesis, was elevated in the TGAC8 vs. WT LV.
However, we have demonstrated that at three-month of age, the mouse age in the present study,
the LV mass is not elevated in TGAC8 vs. WT (Ref to be submitted). Of note, transcriptomic,
proteomic and WB analyses indicated that pathologic hypertrophic markers were not increased in
TGAC8 vs. WT (Ref to be submitted). In this regard, the activation of key stress response
signaling pathways, e.g., protein kinase A (PKA) signaling, PI3K signaling, and AMPK
signaling, may protect against pathological changes, that result in an increase in cardiac mass in
TGAC8 despite its chronic increase in performance. For example, activated PKA signaling can
restrict the activity of AC8, acting as a feedback to fine-tune the downstream signaling networks

inside the body[7]. Additionally, nitric oxide (NO) signaling may coordinate cardiac protection in
the increase in cardiac mass (Figure 1D). Increased phosphorylation status of HSP90 and Akt in
TGAC8 synergistically increase eNOS activity[14], and increased NO produced by eNOS may
modulate the function of the ryanodine receptor $Ca^{2+}$ release channel (RyR2), which was
increased on the PKA dependent phosphopeptides in TGAC8, on the cardiac sarcoplasmic
reticulum[15]. The TMT-labeling MS identified significant upregulated phosphorylation in the
peptides of RyR2 (RISQTSQVSIDAAHGYSPR, E9Q401 [2805-2823]) (File 1: Sheet 1) ,
consistent with the PKA induced phosphorylation of RyR2-S2809 mediating the increase in $Ca^{2+}$
spark amplitude[16].

## 2.2. Upstream regulators of the remodeled LV phosphoproteome of TGAC8.

As known, the overexpression of AC8 activates the AC8-cAMP-PKA phosphorylation axis[13].
There were 2264 phosphosites whose names and locations are unambiguous within the 4990
upregulated phosphopeptides (p-value ≤ 0.05 and log2(fold-change(TG/WT)) ≥ 0) in TGAC8 vs.
WT. The majority of these phosphosites were serines (S) and threonines (T), and a few were
tyrosines (Y) (Figure 2A). By using WebLogo, we profiled the phosphorylation motif sequences
(Figure 2B). A typical AKT substate motif, R-X-X-pS/T, was the most abundant substrate motif,
consistent with increased PKA dependent phosphorylation[17,18]. In addition to S and T, the center
phosphosites also included phosphotyrosine (Y). The center phosphosites were mainly flanked
by the P-E-E sequence on the C terminus, endowing a strong hydrophobic and acidic property to
the C terminus of the substrate sequences (Figure 2B), supported by the amphiphilicity index
analysis in iceLogo (Figure 2C). Because phosphorylation substrate motifs were not restricted to
PKA's substrate motif, additional kinases must be also involved in reprogramming the TGAC8
phosphoproteome. Therefore, we were inspired to explore additional kinase-substrate pathways
mediating the effect of the cardiac-overexpression of AC8 here.

IPA predicted that 26 kinases were responsible for the above-mentioned phosphorylation motifs
(File 2: Sheet 3), among which, three had previously been identified to differ by genotype in the
LV transcriptome, 18 were previously shown to differ by genotype in the LV proteome, and 18
were identified in the present phosphoproteome. Of note, most of the identified kinases were
upregulated in TGAC8 vs. WT at transcription, translation and PTM (phosphorylation) levels.
Within the phosphoproteome, eight kinases were predicted to be differentially activated by
genotype, and the phosphorylation level of four of these (PDPK1, PAK1, PTK2B and GSK3B)
was increased in TGAC8 vs. WT. These four kinases each phosphorylates various proteins to
regulate cellular functions. For example, a previous study observed that in human glioblastoma
cell lines, T98G and U87MG cells, the interference of human PDPK1 mRNA by siRNA
decreased the phosphorylation level of human BAD protein (S112, S136 and S155) and activated
cell death to suppress tumorigenesis[19]. Another study reported that in mouse myoblast cell line,
C2C12 cells, PDPK1 protein phosphorylated the mouse ULK1 protein[20]. Increased
phosphorylation states of BAD (upregulated at S112, S136 and S155) and ULK1 (upregulated at
S450, S521 and S637/638) in TGAC8 may be linked to activated PDPK1.

Different kinases can coordinate to reprogram the complex regulation network in the organism.
Consistent with the IPA canonical pathway analysis, the phosphorylation level of 27 proteins
(including the three kinases (PDPK1, PAK1 and PTK2B) auto-phosphorylating themselves)
were elevated by the three kinases to regulate cardiac stress signaling pathways, including PKA
signaling, ERK/MAPK signaling, cardiac hypertrophy signaling (enhanced), and renin-
angiotensin signaling (Figure 1E), to cope with the cardiac stress in the TGAC8. For example,
GSK3B is involved in multiple biological processes. PDPK1 phosphorylated GSK3B at S9[21,22]
and PTK2B phosphorylated GSK3B at S9 and Y216[23,24] to inhibit the kinase activity of GSK3B.
In this phosphoproteome, GSK3B was phosphorylated at multiple sites, including S9, Y216,
S219 and S545, and thus GSK3B inhibition was linked to the cardiac adaptation[25,26].

**2.3. Downstream functions of the remodeled LV phosphoproteome of TGAC8.**

To study the downstream functions of these phosphoproteins, we performed molecular and
cellular functions analysis in IPA. The enriched functions were mainly in three categories, i.e.,
cell viability, protein quality control, and metabolism (Figure 1F and File 2: Sheet5). The
functions of cellular assembly and organization, cellular function and maintenance, and cell
death and survival were ranked as the top three terms, indicating the important role of the
reprogrammed phosphoproteome in improving cell viability in response to the chronic stress of
the overexpression of AC8 in the mouse heart.

**3. Integrative analyses of transcriptome, proteome and phosphoproteome.**

We next integrated changes in PTM (phosphorylation) in TGAC8 vs. WT with genotypic
differences in protein translation and in transcription. The circos plot (Figure 3A) illustrates the
genotypic differences (TGAC8 vs. WT) in **molecules** identified across the transcriptome,
proteome, and phosphoproteome. In addition, the correlation scatter plots (Figure 3B)
demonstrated a moderate level of correlation of genotypic differences of molecular changes
within phosphoproteome and transcriptome, and a somewhat stronger correlation between
genotypic differences of molecular changes within phosphoproteome and proteome (Figure 3C).
Concordant and discordant enrichments of canonical pathways differed by genotype among the
three omics are shown in Figure 3D. A comparison of genotypic differences within the three
omics highlights the importance of protein phosphorylation, and the phosphorylation was
coordinated with and supplemental to transcription and translation.

Because kinases and phosphatases are the key regulators of phosphorylation, and
phosphorylation of transcriptional factors (TFs) are the primary determinant of their activities in
transcription, we compared genotypic differences in kinases, phosphatases and TFs within the
phosphoproteome and proteome with our previously published genotypic differences in the
TGAC8 vs. WT transcriptome (Figure 3E). We identified 196 kinases, 71 phosphatases and 191
TFs in the present phosphoproteome (Figure 3F). The protein phosphorylation status of about

one half of these molecules differed significantly (p-value < 0.05) by genotype (Figure 3G), and
the protein phosphorylation status of majority of these molecules was increased (Figure 3H-3J).
The genotypic differences in the transcription (mRNA expression obtained from RNA-seq) and
translation (protein expression obtained from mass spectrometry) of these molecules, however,
differed from those in the phosphoproteome (Figure 3E). Because there were more molecules
that were unchanged or reduced in the TGAC8 vs. WT transcriptome or proteome than in the
phosphoproteome, it is clear that the changes in protein phosphorylation are of utmost
importance in the coordination of transcription, translation and PTM, that underlie the adaptive
response of the TGAC8 LV to the marked, chronic increase stress of its intrinsically activated
AC-cAMP-PKA-Ca$^{2+}$ signaling. Thus, by utilizing multiple bioinformatic analysis methods, we
not only dissected the reprogrammed phosphoproteome and deduced its functions in the TGAC8
mouse LV, but also profiled the coordination of phosphoproteome, transcriptome and proteome.

**3.1. Implications of coordinated changes in protein phosphorylation and transcription in**
**TGAC8 vs. WT.**

The most likely type of coordination between protein phosphorylation and transcription is related
to phosphorylation states of TFs which is determined by kinases and phosphatases activities.
Analysis of the remodeled LV transcriptome of TGAC8 in Gene Ontology (GO) analysis using
DAVID online software discovered that enrichments of numerous phosphorylation and
dephosphorylation related biological processes and molecular functions significantly differed by
genotype (Figure 4A), and that the coordinated expression of numerous kinases and
phosphatases also differed by genotype (Figure 4B-4C and File 8). Because expression levels of
kinases and phosphatases do not denote their activation states, we utilized IPA to predict the
genotypic differences in activation states of upstream kinase and phosphatase (regulators of the
transcriptome). Intriguingly, 60 kinases were predicted to be upstream regulators of the
remodeled LV transcriptome of TGAC8, and the majority of these kinases were predicted to be
activated in TGAC8 vs. WT (Figure 4D and File 9). In contrast, many fewer phosphatases were
predicted between TGAC8 and WT, and most of these were predicted to be inhibited (Figure 4E
and File 9). Thus, the differential genotypic enrichment of phosphorylation status and predicted
activation of kinases and phosphatases identified in our post hoc analysis of our previously
reported TGAC8 vs. WT transcriptome are linked to the differential genotypic reprogramming of
the phosphoproteome discovered in the present study.

We utilized IPA to predict the activation states of upstream transcription regulators on the basis
of the remodeled LV transcriptomes and proteomes, and then determined their expression levels
in transcriptome, proteome and phosphoproteome (Figure 4F). Because TFs are the main
regulators of transcription, and because phosphorylation of TFs usually regulates their
activities[27-29], we focused our subsequent analysis on functions of those TFs that were
differentially phosphorylated in TGAC8 vs. WT. To confirm the hypothesis that the TGAC8
might exert transcriptional control by TFs phosphorylation, we retrieved all TFs identified in our
previous transcriptome and proteome and mapped them to the current phosphoproteome in order

to determine their genotypic differences in phosphorylation status. Of the 191 TFs also identified in this phosphoproteome, 92 TFs differed significantly by genotype in phosphorylation status (p-value < 0.05). In agreement with the general upregulated transcription in TGAC8 LV (Ref to be submitted), nearly all of the TF that differed by genotype had increased phosphorylation level (Figure 4G and File 3). The only exception was purine rich element binding protein G (Purg), known to be hyper-phosphorylated in solid tumors[30].

We conducted Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis to further probe the predicted facts of TFs that were generally hyper-phosphorylated in TGAC8 vs. WT. Numerous immunity related pathways, cancer related pathways, and metabolism related pathways were enriched (Figure 4H and File 4). We then utilized IPA for in-depth analysis. Numerous metabolic canonical pathways (glucocorticoid receptor signaling and PPAR signaling), a cancer related canonical pathway (regulation of epithelial-mesenchymal transition pathway), and immunity related canonical pathways (role of JAK1, JAK2 and TYK2 in interferon signaling, IL-9 signaling, JAK/Stat signaling, etc.) were enriched (Figure 4I and File 5: Sheet 1). Intriguingly, the sumoylation pathway was markedly inhibited. Because both sumoylation and phosphorylation are important PTMs in transcriptional regulation[31,32], it is possible that the upregulated protein phosphorylation and inhibited protein sumoylation pathway in TFs coordinately activate transcription and improve immunity in the TGAC8 LV. In addition, cell viability and metabolism in molecular and cellular functions were enriched in TGAC8 vs. WT (Figure 4J and File 5: Sheet 2), consistent with enrichments of afore mentioned KEGG pathways and IPA canonical pathways (Figure 4H and 4I).

Taken together, we have discovered the existence of substantial crosstalk between protein phosphorylation and transcription in the TGAC8 LV: phosphorylation of TFs reprograms immunity and metabolism pathways and functions in the TGAC8 LV, likely serving to maintain cell survival in response to the chronic stress of marked AC8 overexpression.

**3.2. Coordinated changes in protein expression and phosphorylation of the remodeled proteome and phosphoproteome of TGAC8.**

Having discovered the coordinated changes in the remodeled TGAC8 transcriptome and phosphoproteome (Figure 4A-4E), we next sought to discover the extent to which changes in the remodeled phosphoproteome of the TGAC8 LV were coordinated with changes in its remodeled proteome. Numerous biological processes and molecular functions that were related to phosphorylation and dephosphorylation were enriched in the remodeled TGAC8 LV proteome (Figure 5A), as were numerous kinases and phosphatases (Figure 5B-5C and File 10). Twenty-nine kinases that differed by genotype were predicted to be upstream regulators of the proteome, and most of these were predicted to be activated in TGAC8 vs. WT. Fewer phosphatases than kinases that differed by genotype were identified as upstream regulators of the remodeled TGAC8 LV proteome (Figure 5D-5E and File 11).

One strategy adopted to probe deeper into the coordinated changes in the proteome and
phosphoproteome of the TGAC8 LV was to normalize the phosphoproteome to the total
proteome (in other words, to divide the phosphorylation expression by protein expression) (File
1: sheet4). Of the 3927 phosphoproteins identified in our mouse LV samples, 2164 were
identified in both the proteome and phosphoproteome, and these were focused for subsequent
analyses. After normalization, WT and TGAC8 samples manifested clear genotypic differences
in PCA analysis (Figure 5G), and phosphorylation status of numerous proteins normalized to the
protein expression level still differed by genotype without cutoff (Figure 5H) and with cutoff (-
log(p-value) > 1.3 and absolute(log2(TG/WT)) ≥ 0.4) (Figure 5I). Even after normalizing the
phosphorylation level to the protein expression level, the enrichments of canonical pathways in
TGAC8 were quite consistent with those of unnormalized phosphoproteome (Figure 5J and File
6). Cardiac hypertrophy signaling (enhanced) and cardiac hypertrophy signaling took the top two
places, emphasizing the important role of phosphorylation in the regulation of cardiac functions
in response to chronic stress when coordinated with changes in the total proteome.

Many important stress-response signaling pathways were enriched from phosphoproteome, e.g.,
PKA signaling, ERK/MAPK signaling, insulin receptor signaling, and PI3K/AKT signaling
(Figure 6A). Two of these pathways, PI3K/AKT signaling pathway and ubiquitin labeling, were
also markedly enriched in our previous analysis on total proteome in TGAC8 LV (Ref to be
submitted). By integrating and comparing the enrichments from transcriptome, proteome and
phosphoproteome, we were able to identify two signaling pathways in the TGAC8 LV,
PI3K/AKT signaling pathway and ubiquitin labeling, that were regulated concordantly in
transcription, translation, and PTM (phosphorylation) (Figure 6B). PI3K/AKT Signaling is not
only enriched but also activated in TGAC8, and this activation is crucial for the increase in
protein synthesis and cell survival in the TGAC8 LV (Ref to be submitted). Protein
ubiquitination is an important process in protein degradation, which is also increased in TGAC8
LV (Ref to be submitted), but the activity of this pathways could not be predicted on the basis of
present knowledge base of IPA.

We next sought to identify which molecules linked to PI3K/AKT signaling differed in TGAC8
vs. WT. Among the significantly changed upstream molecules, the majority were upregulated.
AKT, a central molecule in the pathway, was also increased in proteome and phosphoproteome
(Figure 6C). By searching and mapping the substrates of AKT, we unraveled that the
phosphorylation levels of AKT peptide substrates were mostly elevated in TGAC8 vs. WT
(Figure 6D and Supplementary Figure 3). Therefore, the downstream effects and functions of
AKT signaling were activated in TGAC8 LV: (1) upregulating GSK3 to improve energy
metabolism; (2) activating mTOR pathway to increase protein synthesis and cell growth; (3)
promoting NF-κB mediated transcription; (4) upregulating BAD and eNOS only at
phosphorylational level to enhance cell survival and cardiovascular functions. In summary, to
adapt the heart to stress for improved survival, the TGAC8 upregulates the protein
phosphorylation to reprogram the cardiac LV phosphoproteome.

**4. Discordant regulation of the phosphoproteome and proteome in TGAC8 LV.**

Among the 443 proteins that differed by genotype in both protein phosphorylation and protein expression levels, 33 were changed discordantly at the two levels (in opposite directions, i.e., upregulated in proteome but downregulated in phosphoproteome, or vice versa) (Figure 7A and File 7: Sheet 1). We uploaded the 33 discordant proteins into STRING online software for functional analyses. The GO_CC (Cellular Component) analysis resulted in many **mitochondrial components**, indicating the involvement of phosphorylated proteins in mitochondria (Figure 7B and File 7: Sheet 2). Reactome pathway analysis showed that in addition to mitochondria, metabolic pathways were enriched significantly in TGAC8 (Figure 7C and File 7: Sheet 3). Mitochondria are the hub of metabolism and multiple metabolic processes are executed in mitochondria. Consistently, both metabolism and mitochondria related processes were ranked the top processes in GO_BP (Biological Process) analysis (Figure 7D and File 7: Sheet 4). Therefore, the discordant part between total proteome and phosphoproteome indicated that the phosphorylation plays a critical role in the regulation of mitochondria and metabolism in the TGAC8, distinct from its role in regulation at the translational level.

**5. Deduced schematic of coordination of phosphoproteome with transcriptome and proteome in TGAC8 mouse LV.**

A deduced schematic integration of the remodeled phosphoproteome, proteome and transcriptome of the TGAC8 LV is shown in Figure 8. The TGAC8 mouse first induces a PKA dependent phosphorylation via AC8-cAMP-PKA axis. Many cytoplasmic proteins including TFs are thus phosphorylated and then transported into the nucleus where TFs recognize and bind to the promoters to activate the transcription of more regulatory factors, including kinases, phosphatases, and TFs. The activation states of these factors are changed upon phosphorylated in cytosol, thereby activating additional kinases-controlled phosphorylation cascades and reprogramming the cardiac phosphoproteome comprehensively.

**Discussion**

Our prior study (Ref to be submitted) investigated the adaptive mechanisms that emerge in the TGAC8 heart to marked chronic stress of increased AC8-cAMP-PKA-Ca$^{2+}$ signaling. In that study, a large number of proteins differed in TGAC8 vs. WT. The TGAC8 induced a chronic cardiac exercise phenotype in the heart, i.e., an increase in heart rate and contractility. These functional changes were accompanied by an increased cardiac biomass and energy demand. Multiple signaling pathways, related to nutrient sensing, energy production, cardiac cellular stress, cardiac stress response, growth factor signaling, and cell proliferation, were identified as

downstream mediating the effect of the activated AC8-cAMP-PKA-Ca$^{2+}$ signaling in the
TGAC8 LV.

Our goals here were to determine which proteins in TGAC8 LV differed in expression and
phosphorylation status compared to WT, and which kinases/phosphatases play a role in these
altered phosphorylation statuses in the TGAC8. By the utilization of TMT-labeling MS, we
obtained 12963 phosphopeptides and 3646 phosphoproteins, and observed the general
upregulation of phosphorylation. We observed the enrichments of phosphorylation, upregulated
expression of kinases, and predicted activation of upstream kinases. We made use of
phosphoproteome complementary to transcriptome and proteome to expand our understanding of
the regulatory mechanism how the TGAC8 adapts to chronic stress of activated AC8-cAMP-
PKA-Ca$^{2+}$ signaling. This approach enabled us to appreciate how changes in the transcription,
translation and PTM of proteins in the TGAC8 LV may singly or coordinately enable the
TGAC8 to endure the chronic AC8-induced stress (Figure 8).

First, the TGAC8 induced a PKA dependent phosphorylation via the AC8-cAMP-PKA axis
(typical substrate motif of PKA dependent phosphorylation). Most of previous studies only
focused on PKA phosphorylated targets as downstream effects of activated AC8. For example,
AC8 induced cAMP activates PKA to increase the phosphorylation of many key proteins in
cardiovascular system, e.g., ryanodine-receptor-type 2 (RyR2), L-type Ca$^{2+}$ channels (LTCC),
phospholamban (PLN or PLB), etc[33]. In this research, we identified 29 phosphosites on RyR2, 15
phosphosites on LTCC alpha 1 and beta 2, and 2 phosphosites (S16 and T17) on PLN, and the
majority of them were increased in phosphorylation statuses in the TGAC8 LV (File 1: Sheet 1).
It is worth noting that PLN is phosphorylated on S16 by PKA whereas on T17 by Ca$^{2+}$-
calmodulin-dependent protein kinase (CaMKII)[34].

After the activation of phosphorylation cascade, many cytoplasmic proteins including TFs were
thus phosphorylated and then transported into the nucleus where TFs recognize and bind to the
promoters to activate transcription. For example, Figure 4 F and 4G showed that many
transcriptional regulators were phosphorylated in TGAC8 vs. WT. The phosphorylation of
STAT1[35,36], STAT3[36], HTT[37] and CREB1[38] is associated with the activation of their
transcriptional activities. The elevated TFs phosphorylation and possibly activated transcription
were consistent with the general increase in transcriptome (Ref to be submitted). The hyper-
phosphorylated TFs might contribute to the improved immunity, metabolism and cell viability in
TGAC8 (Figure 4H-4I). For example, STAT family proteins are transcription activators, and
their phosphorylation is involved in inflammation and immune response[39,40]. The present study
identified increased phosphorylation levels in S727 on STAT1 and S726/S727 on STAT3 (File
1: Sheet 1), in line with our previous report which identified hyper-phosphorylation of
components within inflammation and immune response signaling pathways such as JAK/STAT
and JNK pathways (Ref to be submitted).

Upon the activation of RNA transcription, numerous molecules, including kinases, phosphatases,
and TFs, were transcribed more in TGAC8 than WT, thereby, activating other kinases-controlled
phosphorylation cascades in addition to PKA. The additional kinases and phosphorylation
cascades could also be directly activated by the overexpression of AC8 in another manner to be
determined. In this study, with the three datasets, IPA predicted many other kinase-substrate
reactions in addition to PKA.  PDPK1, PAK1 and PTK2B were the three most consistent
kinases. In the context of the overexpression of AC8, these three kinases could be activated to
phosphorylate multiple proteins to activate PKA pathway (positive feedback) and MAPK/ERK
pathway (also known as Ras-Raf-MEK-ERK pathway) to maintain heart health (Figure 1E).
Previously, we showed their western blot results that MAPK/ERK pathway were activated due to
increased protein expression of p21 Ras and the elevated phosphorylation levels of c-Raf,
MEK1/2 and ERK1/2 as well as their increased ratios of phosphorylation to protein expression
(Ref to be submitted). In the present study, S233 and S621/S641 were hyper-phosphorylated in
TGAC8 (File 1: Sheet 1). Importantly, the multifunctional molecule, GSK3B, was the
downstream of PDPK1 and PTK2B in the integrated regulation network. GSK3B is involved in
metabolism and cardiac hypertrophy[41,42]. Interestingly, AC8 is required for glucose homeostasis
and hypothalamic adaptation to high-fat diet[43]. The phosphorylation of GSK3B is also a critical
mediator in convergence of cardioprotection signaling[25,26]. We previously detected the increase
in the protein expression of GSK3B (Ref to be submitted) and here identified the hyper-
phosphorylated S9 on GSK3B (File 1: Sheet 1). The bioinformatic analysis strategy here
revealed a larger kinase-substrate network important in the remodeling of the TGAC8 heart. We
plan to validate these kinase-substrate reactions in follow-up studies and provide the reliable
mechanism in cardiovascular research.

Taken together, the TGAC8 upregulates phosphorylation to reprogram the heart. Based on
results in this study combined with our recently published work on TGAC8 LV (Ref to be
submitted), we generated a comprehensive circuitry (Supplementary Figure 3). Navigating this
figure "by the numbers": marked, chronic, cardiac specific overexpression of AC8 (1)
chronically increases cAMP(2) and PKA-Ca$^{2+}$ signaling (3), resulting, not only in an incessantly
increased cardiac performance, due to elevated heart rate and LV contractility (4), but also
activation of growth factor signaling pathways (5), leading to increases in energy demand due to
increased transcription and increased amino acid, nucleotide, protein, lipid and lipoprotein
synthesis, increased protein quality control, including proteosome activity, autophagy,
mitophagy, cytokine signaling and inflammation (6). Chronic high level of AC8 activation elicits
numerous stress responses (8) including increased nutrient sensing with increase AMPK
signaling (9A), increased ROS scavenging, protection against cell death, and also induction of
signaling to minimize the impact of AC-PKA signaling (9B). Shifts in energy production and
utilization to deal with the marked energy demands of the TGAC8 heart (10, 12) include:
enhancement of glucose metabolism is pronounced (11, 13), whilst utilization of fatty acids to
produce energy is reduced (11) but an increase in glucose catabolism (10, 11) is accompanied by
an increased anabolism (12), supported by increased aerobic glycolysis (11,13); increased

utilization of the pentose phosphate shunt (5, 13) supports increased amino acid, nucleotide,
protein and lipoprotein synthesis induced by growth factor signaling (5), resulting in an increase
in LV biomass (14). This adaptive profile balances energy requirements necessary for an
incessant marked increase in cardiac performance (4), for increased RNA transcription, protein
translation and quality control required for cardiac growth (14), and for other biological
processes within the LV myocardium (6). This pattern of consilient adaptations (6-15) sustains
an incessant increase in cardiac performance while conferring cardio-protection to the TGAC8
heart.

In addition to the fascinating adaptive mechanisms in the TGAC8 LV discussed above, the
present research also raised some interesting open-mind questions, such as cardiac hypertrophy
signaling and sumoylation pathway enriched here.

Although the cardiac β-AR signaling is desensitized in many pathological conditions, like aging,
so as to despair the cardiovascular system functions, cAMP bioavailability could maintain
cardiac functions in the aged heart. Based on this, the overexpression of AC8 in heart was
regarded as a potential therapeutic way to increase the bioavailability of cardiac cAMP
independent of β-AR stimulation to improve cardiac functions[11]. Unexpectedly, the cardiac-
specific chronic activation of the cAMP pathway (overexpressing β1- AR, β2-AR, Gas, PKA, I-
I, etc.) that enhances cardiac functions in young animals induces progressive cardiomyopathy
and premature death during aging[44]. Consistently, in the present research, although the enriched
cardiac hypertrophy signaling (enhanced) was not accompanied with the markable hypertrophy
phenotype in young male TGAC8 mice, it indicated a cardiac remodeling that could lead to
hypertrophy if accompanied with additional stresses, like aging. This assumption was supported
by another research on the TGAC8 mouse which reported early alterations of myocardial
structure, i.e., cardiomyocyte hypertrophy and endocardial fibrosis, in twelve-month old TGAC8
mice compared with their WT littermates[45]. Therefore, our established bioinformatic analysis
strategy is a novel paradigm in **predicting** progressive cardiomyopathy and premature death
using early reprograming of omics in young age. More importantly, the mechanisms discovered
here in the young adult TGAC8 mice could be utilized in maintaining cardiac health in older
TGAC8 mice.

Interestingly, more and more PTMs have been discovered nowadays and they often occur at the
same time or on the same protein to carry important biological processes coordinately[46,47]. For
example, in rat hearts subjected to ischemia, lysine acetylation activates protein kinases to
increase proximal dephosphorylation, which is involved in the regulation of cell survival or
death[48]. In the present research, sumoylation pathway was inhibited along with the upregulated
phosphorylation (Figure 4I), indicating the cross-talk between these two PTMs in the mouse
heart. In human cell-lines, the switch from phosphorylation to sumoylation could repress the
Raf/MEK/ERK signaling to improve the immunity, resulting in the declined infection of
influenza A virus[49]. In human HEK293 cells, the coordinated phosphorylation and sumoylation
can also control MEF2 transcriptional activity[50]. However, the competition of phosphorylation

and sumoylation in cardiomyocytes was unclear. In this research, we observed the upregulated
phosphorylation and inhibited sumoylation pathway enriched from TFs in the TGAC8 mouse
LV. The regulated PTMs was speculated to be involved in the increased transcription and
improved immunity. To validate the effect on sumoylation by elevated phosphorylation in our
next studies, we could perform WB with the antibody against pan-sumoylation to detect the
general sumoylation change in the TGAC8 LV. In addition, we could detect the sumoylation
changes after mimicry upregulation or downregulation of phosphorylation using activators or
inhibitors in primary cardiomyocytes to prove the interplay between phosphorylation and
sumoylation in cardiomyocytes.

In summary, the reprogrammed phosphoproteome activated stress response signaling pathways
to defend against cardiac stress in young mice, improve cell viability and suppress cell death. On
one hand, the reprogramed phosphoproteome enhances transcription to improve the immunity
and metabolism; on the other hand, it coordinates with the proteome to improve mitochondrial
functions and metabolism. Finally, although we revealed the coordination of regulation at
different levels, which provides a roadmap to guide the integration of transcription, translation
and PTM, our results set the stage for additional studies in this remarkable TGAC8 model: What
kinases are specifically involved in the PTM identified here? How are these phosphorylated
proteins specifically distributed in subcellular location? The present study focuses on a
comprehensive in silico analysis of functionally coherent and data-driven transcription,
translation, and post-translation patterns that were well translated in the contribution to AC8-
overexpression cardiac outcomes, while further investigation is warranted to see whether the hub
pathway/function predictions via gene sets, phosphopeptides, and phosphoproteins could be used
as biomarkers or impact the diagnostic specificity for transgenic AC8 cardiac health and
pathology.

**ACKNOWLEDGEMENTS**

We thank all members in Laboratory of Cardiovascular Science, National Institute on Aging, National
Institutes of Health, for technical assistance and discussion. This work utilized the computational
resources of the NIH HPC Biowulf cluster (http://hpc.nih.gov).

**SOURCES OF FUNDING**

Intramural Research Program at National Institute on Aging, National Institutes of Health.

**DISCLOSURES**

None.

## Reference

Consortium, T. U. UniProt: a worldwide hub of protein knowledge. *Nucleic acids research* **47**, D506-D515, doi:10.1093/nar/gky1049 (2018).

Crooks, G. E., Hon, G., Chandonia, J. M. & Brenner, S. E. WebLogo: a sequence logo generator. *Genome research* **14**, 1188-1190, doi:10.1101/gr.849004 (2004).

Colaert, N., Helsens, K., Martens, L., Vandekerckhove, J. & Gevaert, K. Improved visualization of protein consensus sequences by iceLogo. *Nat Methods* **6**, 786-787, doi:10.1038/nmeth1109-786 (2009).

Mitaku, S., Hirokawa, T. & Tsuji, T. Amphiphilicity index of polar amino acids as an aid in the characterization of amino acid preference at membrane-water interfaces. *Bioinformatics* **18**, 608-616, doi:10.1093/bioinformatics/18.4.608 (2002).

Halls, M. L. & Cooper, D. M. F. Adenylyl cyclase signalling complexes - Pharmacological challenges and opportunities. *Pharmacol Ther* **172**, 171-180, doi:10.1016/j.pharmthera.2017.01.001 (2017).

Dessauer, C. W. *et al.* International Union of Basic and Clinical Pharmacology. CI. Structures and Small Molecule Modulators of Mammalian Adenylyl Cyclases. *Pharmacological reviews* **69**, 93-139, doi:10.1124/pr.116.013078 (2017).

Willoughby, D. *et al.* A key phosphorylation site in AC8 mediates regulation of Ca(2+)-dependent cAMP dynamics by an AC8-AKAP79-PKA signalling complex. *J Cell Sci* **125**, 5850-5859, doi:10.1242/jcs.111427 (2012).

Sadana, R. & Dessauer, C. W. Physiological roles for G protein-regulated adenylyl cyclase isoforms: insights from knockout and overexpression studies. *Neuro-Signals* **17**, 5-22, doi:10.1159/000166277 (2009).

Georget, M. *et al.* Augmentation of cardiac contractility with no change in L-type Ca2+ current in transgenic mice with a cardiac-directed expression of the human adenylyl cyclase type 8 (AC8). *Faseb j* **16**, 1636-1638, doi:10.1096/fj.02-0292fje (2002).

Ferguson, G. D. & Storm, D. R. Why calcium-stimulated adenylyl cyclases? *Physiology (Bethesda, Md.)* **19**, 271-276, doi:10.1152/physiol.00010.2004 (2004).

Lipskaia, L. *et al.* Enhanced cardiac function in transgenic mice expressing a Ca(2+)-stimulated adenylyl cyclase. *Circ Res* **86**, 795-801, doi:10.1161/01.res.86.7.795 (2000).

Esposito, G. *et al.* Increased myocardial contractility and enhanced exercise function in transgenic mice overexpressing either adenylyl cyclase 5 or 8. *Basic Res Cardiol* **103**, 22-30, doi:10.1007/s00395-007-0688-6 (2008).

Moen, J. M. *et al.* Overexpression of a Neuronal Type Adenylyl Cyclase (Type 8) in Sinoatrial Node Markedly Impacts Heart Rate and Rhythm. *Front Neurosci* **13**, 615, doi:10.3389/fnins.2019.00615 (2019).

Takahashi, S. & Mendelsohn, M. E. Synergistic activation of endothelial nitric-oxide synthase (eNOS) by HSP90 and Akt: calcium-independent eNOS activation involves formation of an HSP90-Akt-CaM-bound eNOS complex. *The Journal of biological chemistry* **278**, 30821-30827, doi:10.1074/jbc.M304471200 (2003).

15 Lim, G., Venetucci, L., Eisner, D. A. & Casadei, B. Does nitric oxide modulate cardiac ryanodine
receptor function? Implications for excitation-contraction coupling. *Cardiovasc Res* **77**, 256-264,
doi:10.1093/cvr/cvm012 (2008).
16 Meissner, G. in *Structure and Function of Calcium Release Channels  Current Topics in*
*Membranes*   91-113 (2010).
17 Hunzicker-Dunn, M. E. *et al.* PKA and GAB2 play central roles in the FSH signaling pathway to
PI3K and AKT in ovarian granulosa cells. *Proceedings of the National Academy of Sciences of the*
*United States of America* **109**, E2979-2988, doi:10.1073/pnas.1205661109 (2012).
18 Bellis, A. *et al.* Cross-talk between PKA and Akt protects endothelial cells from apoptosis in the
late ischemic preconditioning. *Arterioscler Thromb Vasc Biol* **29**, 1207-1212,
doi:10.1161/atvbaha.109.184135 (2009).
19 Desai, S., Pillai, P., Win-Piazza, H. & Acevedo-Duncan, M. PKC-iota promotes glioblastoma cell
survival by phosphorylating and inhibiting BAD through a phosphatidylinositol 3-kinase pathway.
*Biochim Biophys Acta* **1813**, 1190-1197, doi:10.1016/j.bbamcr.2011.03.007 (2011).
20 Li, Y. *et al.* microRNA-378 promotes autophagy and inhibits apoptosis in skeletal muscle.
*Proceedings of the National Academy of Sciences of the United States of America* **115**, E10849-
e10858, doi:10.1073/pnas.1803377115 (2018).
21 Chen, X. *et al.* PDK1 regulates platelet activation and arterial thrombosis. *Blood* **121**, 3718-3726,
doi:10.1182/blood-2012-10-461897 (2013).
22 Ericson, K. *et al.* Genetic inactivation of AKT1, AKT2, and PDPK1 in human colorectal cancer cells
clarifies their roles in tumor growth regulation. *Proceedings of the National Academy of Sciences*
*of the United States of America* **107**, 2598-2603, doi:10.1073/pnas.0914018107 (2010).
23 Hartigan, J. A., Xiong, W. C. & Johnson, G. V. Glycogen synthase kinase 3beta is tyrosine
phosphorylated by PYK2. *Biochemical and biophysical research communications* **284**, 485-489,
doi:10.1006/bbrc.2001.4986 (2001).
24 Tsai, C. C. *et al.* Glycogen synthase kinase-3beta facilitates IFN-gamma-induced STAT1 activation
by regulating Src homology-2 domain-containing phosphatase 2. *J Immunol* **183**, 856-864,
doi:10.4049/jimmunol.0804033 (2009).
25 Juhaszova, M. *et al.* Glycogen synthase kinase-3beta mediates convergence of protection
signaling to inhibit the mitochondrial permeability transition pore. *The Journal of clinical*
*investigation* **113**, 1535-1549, doi:10.1172/JCI19906 (2004).
26 Juhaszova, M. *et al.* Role of glycogen synthase kinase-3beta in cardioprotection. *Circ Res* **104**,
1240-1252, doi:10.1161/CIRCRESAHA.109.197996 (2009).
27 Bohmann, D. Transcription factor phosphorylation: a link between signal transduction and the
regulation of gene expression. *Cancer cells (Cold Spring Harbor, N.Y. : 1989)* **2**, 337-344 (1990).
28 Weigel, N. L. & Moore, N. L. Steroid receptor phosphorylation: a key modulator of multiple
receptor functions. *Molecular endocrinology (Baltimore, Md.)* **21**, 2311-2319,
doi:10.1210/me.2007-0101 (2007).
29 Mayr, B. & Montminy, M. Transcriptional regulation by the phosphorylation-dependent factor
CREB. *Nature reviews. Molecular cell biology* **2**, 599-609, doi:10.1038/35085068 (2001).
30 Zanivan, S. *et al.* Solid tumor proteome and phosphoproteome analysis by high resolution mass
spectrometry. *J Proteome Res* **7**, 5314-5326, doi:10.1021/pr800599n (2008).

31    Tomanov, K. *et al.* Sumoylation and phosphorylation: hidden and overt links. *Journal of*
*Experimental Botany* **69**, 4583-4590, doi:10.1093/jxb/ery167 (2018).
32    Khan, M. *et al.* Interplay between phosphorylation and SUMOylation events determines CESTA
protein fate in brassinosteroid signalling. *Nat Commun* **5**, 4687, doi:10.1038/ncomms5687
(2014).
33    Guellich, A., Mehel, H. & Fischmeister, R. Cyclic AMP synthesis and hydrolysis in the normal and
failing heart. *Pflugers Arch* **466**, 1163-1175, doi:10.1007/s00424-014-1515-1 (2014).
34    Wegener, A. D., Simmerman, H. K., Lindemann, J. P. & Jones, L. R. Phospholamban
phosphorylation in intact ventricles. Phosphorylation of serine 16 and threonine 17 in response
to beta-adrenergic stimulation. *The Journal of biological chemistry* **264**, 11468-11474 (1989).
35    Pilz, A. *et al.* Phosphorylation of the Stat1 transactivating domain is required for the response to
type I interferons. *EMBO reports* **4**, 368-373, doi:10.1038/sj.embor.embor802 (2003).
36    Wen, Z., Zhong, Z. & Darnell, J. E., Jr. Maximal activation of transcription by Stat1 and Stat3
requires both tyrosine and serine phosphorylation. *Cell* **82**, 241-250, doi:10.1016/0092-
8674(95)90311-9 (1995).
37    Kumar, A., Vaish, M. & Ratan, R. R. Transcriptional dysregulation in Huntington's disease: a
failure of adaptive transcriptional homeostasis. *Drug discovery today* **19**, 956-962,
doi:10.1016/j.drudis.2014.03.016 (2014).
38    Wang, H., Xu, J., Lazarovici, P., Quirion, R. & Zheng, W. cAMP Response Element-Binding Protein
(CREB): A Possible Signaling Molecule Link in the Pathophysiology of Schizophrenia. *Front Mol*
*Neurosci* **11**, 255-255, doi:10.3389/fnmol.2018.00255 (2018).
39    Seif, F. *et al.* The role of JAK-STAT signaling pathway and its regulators in the fate of T helper
cells. *Cell Communication and Signaling* **15**, 23, doi:10.1186/s12964-017-0177-y (2017).
40    Decker, T. & Kovarik, P. Serine phosphorylation of STATs. *Oncogene* **19**, 2628-2637,
doi:10.1038/sj.onc.1203481 (2000).
41    Beurel, E., Grieco, S. F. & Jope, R. S. Glycogen synthase kinase-3 (GSK3): regulation, actions, and
diseases. *Pharmacol Ther* **148**, 114-131, doi:10.1016/j.pharmthera.2014.11.016 (2015).
42    Lal, H., Ahmad, F., Woodgett, J. & Force, T. The GSK-3 family as therapeutic target for myocardial
diseases. *Circ Res* **116**, 138-149, doi:10.1161/circresaha.116.303613 (2015).
43    Raoux, M. *et al.* Multilevel control of glucose homeostasis by adenylyl cyclase 8. *Diabetologia*
**58**, 749-757, doi:10.1007/s00125-014-3445-z (2015).
44    El-Armouche, A. & Eschenhagen, T. Beta-adrenergic stimulation and myocardial function in the
failing heart. *Heart failure reviews* **14**, 225-241, doi:10.1007/s10741-008-9132-8 (2009).
45    Mougenot, N. *et al.* Cardiac adenylyl cyclase overexpression precipitates and aggravates age-
related myocardial dysfunction. *Cardiovascular research* **115**, 1778-1790,
doi:10.1093/cvr/cvy306 (2019).
46    Beltrao, P., Bork, P., Krogan, N. J. & van Noort, V. Evolution and functional cross-talk of protein
post-translational modifications. *Molecular systems biology* **9**, 714, doi:10.1002/msb.201304521
(2013).
47    Venne, A. S., Kollipara, L. & Zahedi, R. P. The next level of complexity: crosstalk of
posttranslational modifications. *Proteomics* **14**, 513-524, doi:10.1002/pmic.201300344 (2014).

48    Parker, B. L. *et al.* Structural basis for phosphorylation and lysine acetylation cross-talk in a
kinase motif associated with myocardial ischemia and cardioprotection. *The Journal of biological*
*chemistry* **289**, 25890-25906, doi:10.1074/jbc.M114.556035 (2014).
49    Wang, C. *et al.* HA Triggers the Switch from MEK1 SUMOylation to Phosphorylation of the ERK
Pathway in Influenza A Virus-Infected Cells and Facilitates Its Infection. *Frontiers in cellular and*
*infection microbiology* **7**, 27, doi:10.3389/fcimb.2017.00027 (2017).
50    Grégoire, S. *et al.* Control of MEF2 transcriptional activity by coordinated phosphorylation and
sumoylation. *The Journal of biological chemistry* **281**, 4423-4433, doi:10.1074/jbc.M509471200
(2006).

**Figure Legends**

**Figure 1. TGAC8 reprograms the LV phosphorylation state. A**. PCA plot of phosphoproteome in
TGAC8 vs. WT LVs. **B**. Heatmap of scaled expressions of phosphorylated proteins. **C**. Volcano plot of
phosphoproteome. Red dots represent upregulated phosphoproteins with -log10(p-value) > 1.3 and
log2(fold change) ≥ 0.4; blue dots represent downregulated phosphoproteins with -log10(p-value) > 1.3
and log2(fold change) ≤ -0.4; grey dots represent nonsignificantly regulated phosphoproteins with -
log10(p-value) ≤ 1.3 or absolute(log2(fold change)) < 0.4. **D**. Top 21 canonical pathways enriched from
phosphoproteome. Orange and blue bars represent activated and inhibited pathways respectively. The
darker the color is, the greater the absolute(z-score) is, and the more significantly enriched the pathway is.
**E**. Regulatory pathway from upstream kinases. Three dots in the center represent the predicted upstream
kinases. Surrounding dots represent identified phosphoproteins differentially regulated in
phosphoproteome. Terms on the bottom represent canonical pathways in which the phosphoproteins are
involved. Meanings of colors refer to the box of prediction legend besides the graph. **F**. Enriched
molecular and cellular functions from the phosphoproteome. They are related to three categories,
including cell viability, protein quality control, and metabolism.

**Figure 2. Characteristics of upregulated phosphopeptides. A**. Composition of phosphosites that reside
on the upregulated phosphopeptides. **B**. Sequence logo diagrams of upregulated phosphopeptides centered
on the phosphosites and flanked by 6 amino acids on the N and C termini respectively using WebLogo
online software. Colors of amino acids represent different chemical properties. **C**. Test the amphiphilicity
feature of the upregulated phosphopeptides using iceLogo online software and referring to a published
work by Mitaku and colleagues in 2002.

**Figure 3. Integrative analysis of transcriptome, proteome and phosphoproteome. A**. Circos plot of
gene expression at different levels from transcription, to protein, and to phosphorylation. Bars in each
layer of circle represent changes of significantly regulated molecules (q-value or p-value < 0.05) in
transcriptome, total proteome, and unnormalized phosphoproteome. Bar height, color and direction
represent log2(fold change). Red and upward indicate upregulation, and blue and downward indicate
downregulation. **B-C**. Scatter plots of correlation between phosphoproteome and transcriptome (**B**) or
total proteome (**C**). **D**. Comparison of canonical pathway enrichments from the transcriptome, total

proteome, and phosphoproteome. Enrichment z-scores were calculated in IPA, and were scaled to
generate heatmap in R language. **E**. Integrative graph of expression of all phosphoproteins and selected
types of molecules. Bar plot on the bottom displays expression of all significantly changed (p-value <
0.05) phosphoproteins in unnormalized phosphoproteome. Y-axis denotes log2(fold change), and x-axis
denotes each molecule ranked by log2(fold change). Bars were colored by -log10(p-value). Stick plots on
the top display kinases, phosphatases and transcription factors (TFs) selected from the bottom bar plot and
their changes in transcriptome (RNA), total proteome (pro), and normalized phosphoproteome (norm).
From top to bottom: each large block denotes the changes of certain types of molecules (from kinases to
phosphatases, to TFs). In each block, the three rows from top to bottom indicate the changes in
transcriptome, total proteome, and normalized phosphoproteome. Each stick represents a molecule and
color represents the change: red: (p-value or q-value < 0.05) and log2FC > 0; blue: (p-value or q-value <
0.05) and log2FC < 0; black: (p-value or q-value ≥ 0.05) or log2FC = 0; grey: unidentified in one of the
omics. Their locations are matched to those in the bottom bar plot. **F-J**. Pie plots of composition of
phosphoproteins in the unnormalized phosphoproteome. Regardless of p-value, numbers of four types of
proteins, including kinases, phosphatases, TFs and others, are displayed in **F**. After filtration with cutoff
p-value < 0.05, numbers of those types of proteins are displayed in **G**. Extracted from G, numbers of
upregulated and downregulated kinases (**H**), phosphatases (**I**) and TFs (**J**) are displayed.

**Figure 4. Coordination of phosphoproteome and transcriptome. A**. GO term enrichments of
biological processes and molecular functions about phosphorylation and dephosphorylation from
transcriptome. **B**-**C**. Changes of kinases (**B**) and phosphatases (**C**) whose q-values are less than 0.05 in
transcriptome. **D-E**. States of upstream kinases (**D**) and phosphatases (**E**) predicted based on
transcriptome. Z-scores were calculated in IPA. Positive and negative z-scores indicate activated and
inhibited states respectively. The larger absolute values of z-score indicate more activated or inhibited
states. **F**. Heatmap of predicted upstream transcription regulators (left) and their changes in different
omics datasets (right). Left: Fill color represent activation z-scores in transcriptome and total proteome.
Red and blue indicate activated and inhibited states respectively. Molecules are ranked by z-scores in
transcriptome. Right: Fill color represent changes in transcriptome, total proteome, unnormalized
phosphoproteome, and normalized phosphoproteome: orange: (p-value or q-value < 0.05) and log2FC >
0; yellow: (p-value or q-value ⩾ 0.05) and log2FC > 0; green: (p-value or q-value < 0.05) and log2FC <
0; light blue: (p-value or q-value ⩾ 0.05) and log2FC < 0; grey: unidentified in one of the omics. **G**.
Changes of differentially regulated TFs with p-value < 0.05 in the unnormalized phosphoproteome. **H**.
KEGG pathway enrichments of the differentially regulated TFs. They are in four primary categories:
cancer, immunity, metabolism and others. **I**. Top 20 canonical pathways enriched from the
phosphoproteome using IPA. The darker the color is, the greater the absolute(z-score) is, and the more
significantly enriched the pathway is. **J**. Enriched molecular and cellular functions from the
phosphoproteome using IPA. They are in two primary categories: cell viability and metabolism.

**Figure 5. Analysis of normalized phosphoproteome. A**. GO term enrichments of biological processes
and molecular functions about phosphorylation and dephosphorylation from the present proteome. **B**-**C**.
Changes of kinases (**B**) and phosphatases (**C**) whose q-values are less than 0.05 in previous proteome. **D-**
**E**. States of upstream kinases (**D**) and phosphatases (**E**) predicted based on the present proteome. Z-scores

were calculated in IPA. Positive and negative z-scores indicate activated and inhibited states respectively.
The larger absolute values of z-score indicate more activated or inhibited states. **F**. Venn plot of new total
proteome and unnormalized phosphoproteome that were conducted concurrently in the same platform. **G**.
PCA plot of phosphoproteome normalized to total proteome. **H**. Heatmap of expressions of
phosphoproteins normalized to total proteins. **I**. Volcano plot of normalized phosphoproteome. Red dots
represent upregulated phosphoproteins with -log10(p-value) > 1.3 and log2(fold change) $\geqslant$ 0.4; blue dots
represent downregulated phosphoproteins with -log10(p-value) > 1.3 and log2(fold change) $\leqslant$ -0.4; grey
dots represent nonsignificantly regulated phosphoproteins with -log10(p-value) $\leqslant$ 1.3 or
absolute(log2(fold change)) < 0.4. **J**. Comparison of canonical pathway enrichments between
unnormalized and normalized phosphoproteome. Orange fill colors indicate that all comparable pathways
on the top are activated in both omics datasets. The darker the color is, the greater the absolute(z-score) is,
and the more significantly enriched the pathway is.

**Figure 6. Coordinated regulation of transcriptome, proteome and phosphoproteome on PI3K/AKT**
**signaling. A**. Canonical pathway enrichment of phosphoproteome only with cutoff p-value < 0.05. The
darker the color is, the greater the absolute(z-score) is, and the more significantly enriched the pathway is.
**B**. Comparison of PI3K/AKT signaling and protein ubiquitination pathways enriched from transcriptome
(red outline), proteome (blue outline) and phosphoproteome (green outline). **C**. Pathway graph of
PI3K/AKT signaling marked with changes in transcriptome (left box), proteome (middle box) and
phosphoproteome (right box). Fill colors indicate changes: red and green indicate upregulation and
downregulation respectively; grey indicates nonsignificant regulation. **D**. Phosphopeptide number of
substrates of PI3K. Y-axis represents protein substates of PI3K; x-axis represents number of
phosphopeptides. Red and blue bars indicate upregulated (p-value < 0.05 and log2(fold change) > 0) and
downregulated (p-value < 0.05 and log2(fold change) < 0) phosphopeptides respectively.

**Figure 7. Analyses of proteins changed discordantly in proteome and phosphoproteome. A**. Venn
diagram of significantly regulated proteins in total proteome and significantly regulated phosphoproteins
in unnormalized phosphoproteome that were conducted concurrently in the same platform. **B-D**.
Functional analyses of proteins whose expressions are changed discordantly at total protein and
phosphorylation levels. Mitochondria and others are enriched in cellular component analysis (**B**).
Functions related to metabolism, mitochondria and others are enriched in Reactome pathway (**C**) and
biological process analyses (**D**).

**Figure 8. Deduced schematic of coordination of phosphoproteome with transcriptome and**
**proteome in TGAC8 mouse LV.** Based on our results, we deduced a mechanism of reprogramed
phosphoproteome in TGAC8 mouse LV: First, TGAC8 mouse induces a PKA dependent phosphorylation
via AC8-cAMP-PKA axis. Many cytoplasmic proteins including TFs are thus phosphorylated and then
transported into the nuclear where TFs recognize and bind to the promotors to activate the transcription of
more regulatory factors, including kinases, phosphatases, and TFs. The activation states of these factors
are changed upon phosphorylated in cytosol, thereby activating additional kinases-controlled

phosphorylation cascades and reprogramming the cardiac phosphoproteome comprehensively. Left
column: AC8-cAMP-PKA signaling; middle column: phosphorylation (phosphoproteome); lower right:
transcription (transcriptome); upper right: translation (proteome).

**Supplementary Figure Legends**

**S-Figure 1. The enriched pathway, Cardiac Hypertrophy Signaling (Enhanced), from the**
**phosphoproteome.** Cardiac Hypertrophy Signaling (Enhanced) is visualized using IPA. Dots with purple
outline are identified phosphoproteins or protein complexes. Red and green fill colors indicate
upregulation and downregulation respectively.

**S-Figure 2. The regulatory pathway under the control of a predicted upstream kinase, PDPK1.**
PDPK1 in the center is one representative of predicted upstream kinases. Surrounding dots represent
identified phosphoproteins differentially regulated in phosphoproteome. Meanings of colors refer to the
box of prediction legend besides the graph. Blue glows represent that the increased phosphorylation levels
inhibit the activity of corresponding proteins.

**S-Figure 3. Phosphopeptide number of substrates of PI3K.** Y-axis represents protein substates of
PI3K; x-axis represents number of phosphopeptides. Red and blue bars indicate upregulated and
downregulated phosphopeptides respectively; green bars indicate nonsignificantly regulated ones. A-B.
Red (up): $p$-value $< 0.1$ and $\log 2$(fold change) $> 0$; blue (down): $p$-value $< 0.1$ and $\log 2$(fold change) $> 0$;
green (ns): $p$-value $\geq 0.1$ or $\log 2$(fold change) $= 0$. C. Red (up): $p$-value $< 0.05$ and $\log 2$(fold change) $> 0$;
blue (down): $p$-value $< 0.05$ and $\log 2$(fold change) $> 0$; green (ns): $p$-value $\geq 0.05$ or $\log 2$(fold change) $=$
$0$.

**S-Figure 4.  Label findings from phosphoproteome in the exquisite mechanism graph.** Based on the
present phosphoproteome, we optimized our previously deduced mechanism graph and labeled findings
from the phosphoproteome using red stars. TGAC8 mouse induces a chronic cardiac exercise on heart,
manifested as increased heart rate and contractility accompanied with increased energy demand and
cardiac biomass. Multiple signaling pathways, related to nutrient sensing, energy production, cardiac
cellular stress, cardiac stress response, growth factor signaling, and cell proliferation, are potential
downstream mediating the effect of the activated AC8-cAMP-PKA-Ca2+ signaling in mouse LV. The
altered cardiac functions were also observed associated with protein phosphorylation in our previous
study and labeled in blue star here.

# Figure 1. TGAC8 reprograms the LV phosphorylation state.

**Figure 2. Characteristics of upregulated phosphopeptides.**

**Figure 3. Integrative analysis of transcriptome, proteome and phosphoproteome.**

# Figure 4. Coordination of phosphoproteome and transcriptome.

# Figure 5. Analysis of normalized phosphoproteome.

# Figure 6. PI3K/AKT signaling.

**Figure 7. Analyses of proteins changed discordantly in proteome and phosphoproteome.**

**Figure 8. Deduced schematic of coordination of phosphoproteome with transcriptome and proteome in TGAC8 mouse LV.**