Explaining and Predicting the Effects of Neurostimulation via Neuronal

Excitation/Inhibition on Learning

Nienke E. R. van Bueren[1,2,3*], Sanne H. G. van der Ven[2], Shachar Hochman[3], Francesco

Sella[4], Roi Cohen Kadosh[1,3*]

[1]Wellcome Centre for Integrative Neuroimaging, Department of Experimental Psychology,

University of Oxford, Oxford OX2 6GG, United Kingdom

[2]Behavioural Science Institute, Radboud University Nijmegen, 6525 GD Nijmegen, the

Netherlands

[3]School of Psychology, University of Surrey, Guildford GU2 7XH, United Kingdom

[4]Centre for Mathematical Cognition, Loughborough University, Loughborough LE11 3TU,

United Kingdom

[*]Correspondence to:

E-mail:

r.cohenkadosh@surrey.ac.uk

nienke.vanbueren@ru.nl

Abstract

Previous research has highlighted the role of the excitation/inhibition ratio (E/I) for typical and atypical development, mental health, cognition, and learning. Other research has highlighted the benefits of high-frequency random noise stimulation (tRNS)—an excitatory form of neurostimulation—on learning. We examined the E/I as a potential mechanism and studied whether tRNS effect on learning depends on E/I as measured by the aperiodic exponent. In addition to manipulating E/I using tRNS, we also manipulated the level of learning (learning/overlearning) that has been shown to influence E/I. Participants (n=102) received either sham stimulation or 20 min tRNS over the dorsolateral prefrontal cortex (DLPFC) during a mathematical learning task. We showed that tRNS increased E/I, as reflected by the aperiodic exponent, and that lower E/I predicted greater benefit from tRNS specifically for the learning task. In contrast to previous Magnetic Resonance Spectroscopy-based E/I studies, we found no effect of the level of learning on E/I. A further analysis using a different data set suggest that both measures of E/I (EEG vs MRS) may reflect, at least partly, different biological mechanisms. Our results highlight the role of E/I as a marker for neurostimulation efficacy and learning. This mechanistic understanding provides better opportunities for augmented learning and personalized interventions.

Keywords: mathematics; skill acquisition, learning; overlearning; aperiodic activity; brain stimulation

Introduction

Previous human and animal studies have indicated the importance of the ratio of neuronal

excitation to inhibition (E/I) for learning, i.e., skill acquisition (Ahmad et al., 2022; Dorrn et

al., 2010; Nishiyama et al., 2010; Shibata et al., 2017). Previous magnetic resonance

spectroscopy (MRS) studies highlighted the neurotransmitters glutamate and gamma-

aminobutyric acid (GABA) as the underlying building blocks of E/I, and suggested their

important role in memory and learning, including predicting educational levels later in life

(Barron et al., 2016; Shibata et al., 2017; Wijtenburg et al., 2017; Zacharopoulos et al., 2021a;

Zacharopoulos et al., 2021b).

Recent findings show that an excitatory form of noninvasive neurostimulation—high-

frequency transcranial random noise stimulation (tRNS, Terney et al., 2008)—influences E/I in

mice by reducing GABAergic activity (Sanchez-Leon et al., 2020). During tRNS, a small

current with randomized frequency and current intensity is applied over targeted brain areas. It

is assumed that tRNS amplifies subthreshold neuronal activity, that by itself does not reach the

necessary threshold to yield an action potential (i.e., stochastic resonance; Fertonani &

Miniussi, 2017). Thus, this amplification of the signal has been linked to increases in signal-to-

noise ratio, which is assumed to relate to successful enhancements in learning, perception, and

cognitive performance (Antal & Herrmann, 2016; Cappelletti et al., 2013; Fertonani et al., 2011;

Herpich et al., 2019; Snowball et al., 2013; Van der Groen & Wenderoth, 2016). Despite the

growing interest in applying tRNS in cognitive studies, there is little understanding of the

neurophysiological changes induced (Potok et al., 2022). Mostly, tRNS studies that use an

electroencephalogram (EEG) focus on periodic brain activity, such as theta/beta ratio, for the

predicted efficacy of electrical stimulation on skill acquisition (Harty & Cohen Kadosh, 2019;

Rufener et al., 2017). Considerable research has been done to investigate oscillatory rhythms

as potential electrophysiological predictors for cognitive or behavioral processing in healthy

and clinical populations (Harty & Cohen Kadosh, 2019, Uhlhaas & Singer, 2010; Voytek &

Knight, 2015). However, explaining tRNS, as well as other neurostimulation, efficacy by

investigating the E/I has been overlooked, despite the emerging theoretical motivation (Krause

et al., 2013).

Recently, the interest in electrophysiology has been expanded from this oscillatory (i.e.,

periodic or spectral power) perspective to include an aperiodic perspective, e.g., aperiodic

activity (Ahmad et al., 2022; Donoghue et al., 2020). Aperiodic activity is shown in the EEG

spectrum as a 1/f-like structure and is dominant in the spectrum even when there is no periodic

or oscillatory activity. The power of aperiodic activity decreases exponentially with increasing

frequency (see **Figure 1A**), which is reflected as a negative slope in log-log space (see **Figure**

1B). In contrast to the previous assumption that aperiodic activity reflects background noise in

the EEG spectrum, accumulating evidence points to the importance of aperiodic activity in

understanding brain functions and behavior. Also, periodic activity has been shown to be

confounded due to misestimating spectral power since participants vary in center frequencies if

a predefined spectral range is applied (Lansbergen et al., 2011). Therefore, Donoghue et al.

(2020) recommend to parameterize neural power spectra by also analyzing the aperiodic

activity in the spectrum. Aperiodic activity consists of an aperiodic exponent that can be defined

as x in a 1/f^x function, which reflects the previously mentioned negative slope in log-log space

and, thus, the pattern of power across frequencies. The exponent of aperiodic activity is thought

to underlie the integration of underlying synaptic currents (Buzsáki et al., 2013), and a likely

mechanism of changes in the aperiodic exponent has been linked to the E/I of field potentials

shown by EEG recordings (Gao et al., 2017). A higher E/I relates to a lower aperiodic exponent

and vice versa. The power of inhibitory GABA currents leads to a rapid decay in the power

spectrum at higher frequencies, and, thus a steeper (negatively sloped) exponent (see **Figure**

1C). The opposite happens for excitatory currents, where power is stable for lower frequencies

and declines more slowly for higher frequencies, which is shown in a flatter (closer to zero)

exponent (see **Figure 1C**). Shortly, <u>the higher the</u> E/I, the lower the exponent value (see **Figure**

1D).

Figure 1| E/I and the aperiodic exponent. A) A simplified overview of the difference between
periodic and aperiodic activity in the EEG power-frequency spectrum. **B)** The aperiodic
exponent in log-log space as shown in the EEG spectrum. **C)** and **D)** are adapted with
permission from Gao et al. (2017) which show that high E/I is related to a flatter (closer to zero)
aperiodic exponent and low E/I (i.e., high inhibition) to a negatively steep exponent, compared
to the local field potential (LFP).

Donoghue and colleagues (2020) showed that decreased aperiodic activity (i.e., lower

exponent) in the EEG spectrum relates to flattening the power spectrum as seen in ageing, and

that it is also related to behavioral performance. Other developmental and clinical studies also

indicated that inter-individual differences are important in aperiodic activity in health and

disease (He et al., 2019; Molina et al., 2020; Robertson et al., 2019; Voytek et al., 2015).

Recent MRS findings have linked better mathematical skills to higher E/I in young adults

and the reverse in younger participants (Zacharopoulos et al., 2021b). Moreover, MRS-based

E/I can predict future mathematical reasoning (Zacharopoulos et al., 2021c). It is unknown

whether the E/I can be influenced by tRNS, and if this relates to better mathematical

achievement. In the present study we address this question applying tRNS while participants

solved arithmetic multiplications during a mathematical learning paradigm. Furthermore, we

manipulated the difficulty of the to-be acquired skill to induce different degrees of skill

acquisition: learning or overlearning. Based on previous studies, we defined learning as

practicing a skill during performance improvement (i.e., before learning plateaus) and

overlearning as the point after performance improvement when a plateau has been reached

(Shibata et al., 2017). Learning and overlearning have been linked to an increase and a decrease

in E/I, respectively, in an MRS study (Shibata et al., 2017).

We aimed to impact E/I directly using tRNS, as well as indirectly by manipulating the level

of skill acquisition (learning/overlearning) to examine whether: 1) tRNS will increase E/I as

measured by aperiodic exponent; 2) The direction of change in aperiodic exponent between

pre- and posttest depends on the learning condition: decreasing in the learning condition and

increasing in the overlearning condition; 3) tRNS efficacy on a learning/overlearning task

depends on the individual baseline aperiodic exponent. That is, the tRNS-induced reduction of

the aperiodic exponent differs across participants, depending on their baseline aperiodic

exponent (i.e., E/I levels) (Krause et al., 2013).

<div align="center">

Results

</div>

Phase of Skill Acquisition Efficacy

Both groups were matched on baseline performance and aperiodic exponent (see Methods 'Baseline Matching'). To determine the efficacy of our learning and overlearning task manipulation, the average learning slope (based on response times (RTs); Newell & Rosenbloom, 1981) was computed for all participants who received sham stimulation (see **Figure 2** and **Figure S1** for individual data). This allowed us to prevent confounding the effect of learning/overlearning with the effect of active tRNS. For participants in the learning task, the slope showed a negative linear gradient, whereas participants in the overlearning task showed a clear plateau of performance improvement due to the repetition of presented stimuli. This indicated the efficacy of our task design in manipulating learning and overlearning.

Figure 2| Averaged learning curves of the median RTs of the learning or overlearning task of the sham stimulation. The mean learning curve of the participants (n=22) during learning shows a linear gradient as shown in purple. The mean learning curve of the participants (n=21) during overlearning (in green) shows a clear plateau of performance improvement after approximately block 10, and faster RTs overall. Shading indicates 95% confidence intervals.

The Impact of tRNS and Learning on the Aperiodic Exponent

Then we investigated the effects of tRNS and type of mathematical task

(learning/overlearning) on the aperiodic exponent. The aperiodic exponent change was

calculated by subtracting the pre from the post aperiodic exponent, with positive values

indicating an increase in the exponent from pre to post learning/overlearning. We ran an

ANCOVA with the factors task (learning/overlearning) and stimulation (tRNS/sham), while

controlling for the individual plateau (as it may impact E/I; Shibata et al., 2017) (See the

calculation of the amount of learning in the Methods section). The main effect of stimulation

was significant ($F(1,67)$=6.63, p=.012, MSE=1.20, $\eta^2_{partial}$=.09). No significant main effects

were found for task ($F(1,67)$=0.26, p=.611, MSE=0.04, $\eta^2_{partial}$ =.005), and individual plateau

($F(1,67)$=0.37, p=.540, MSE=0.06, $\eta^2_{partial}$ =.01). Also, no interaction effect of stimulation X

task was found ($F(1,67)$=3.59, p=.062, MSE=0.65, $\eta^2_{partial}$ =.05). We repeated the same analysis

without controlling for the individual plateau. This did not affect the results.

To further explain the significant effect of stimulation, we plotted the aperiodic exponent

change for each stimulation group separately (see **Figure 3** and **Figure S2** for individual data).

The aperiodic exponent change of participants who received tRNS (i.e., more excitation

induced) was lower (i.e., flatter) (M=-0.17, SEM=0.07) after stimulation compared to those

who received sham stimulation (M=0.08, SEM=0.06). This corroborates with the excitatory

effects of tRNS, leading to a lower aperiodic exponent, which reflects a higher E/I. However,

it should be noted that in contrast to our expectations, type of task (learning/overlearning) did

not influence the aperiodic exponent change from pre to post.

Figure 3| ANCOVA showing differences in aperiodic exponent for the stimulation groups (tRNS/sham) on the change in aperiodic exponent. Participants who received tRNS show an increase in E/I as indicated by the mean (±S.E.M.) decreased aperiodic exponent (change: post-pre exponent in $\mu V^2 Hz^{-1}$). Participants who received sham stimulation showed a mean (±S.E.M.) decrease in E/I as indicated by an increased exponent from pre to post. *$p < .05$.

Because a frequentist approach makes it difficult to distinguish a true lack of effect (of task on aperiodic exponent) from a lack of power to detect an effect, we reran the same ANCOVA on the aperiodic exponent change using a Bayesian approach. Our results, as presented in the Supplementary Information, strengthens the conclusion that tRNS impacted the aperiodic exponent, while task had no effect. These findings match the idea that tRNS leads to higher excitation and therefore a lower (i.e., flatter) aperiodic exponent and that this effect is independent of learning/overlearning.

The Aperiodic Exponent Moderates Response Times on a Learning and Overlearning Task

As shown in the previous paragraph, the aperiodic exponent was not influenced by the type of task. To investigate if tRNS efficacy on a learning/overlearning task depends on the individual baseline aperiodic exponent, we ran a Bayesian mixed effects model with the *brms* package to predict RTs for each trial during the learning and overlearning task. Note that we also evaluated the models for accuracy instead of RTs as dependent variable, but due to the

emphasis on RTs in cognitive skill acquisition (Newell & Rosenbloom, 1981; Snowball et al.,

2013; Tzelgov et al., 2000) and participant instructions to avoid errors inducing a high accuracy

(see Methods 'Baseline Ability Task'), we only reported the accuracy results in the

Supplementary Information.

Fixed effects entailed the aperiodic exponent at baseline, trial (1-180), task

(learning/overlearning) and stimulation (tRNS/sham). The model included a random intercept

for trial for all participants. Our effect of interest was the threeway tRNS X baseline aperiodic

exponent X task interaction over trials, and therefore we compared different models varying in

number of included interaction effects. Model comparisons were made by means of leave-one-

out (LOO) cross-validation, including a basic learning model that contained RTs as dependent

variable, and trial and task as fixed effects. As can be seen in **Table S1,** the differences in

predictive performance between the different models are negligible (see also Sivula et al.

2020)). In other words, the predictive value of the different interaction models on the data is

very similar. Nevertheless, we investigated the best predicting model further, that is, the model

with the three-way interaction of tRNS X baseline aperiodic exponent X task with trial as a

covariate. Inference regarding the effects was conducted by inspecting the 95% highest

posterior density (HPD) CrI of the posterior distribution of the effect of interest. If the null value

(i.e., zero) was not included in the interval, it means there is a 95% chance or more that the

effect exists, and there is more support for the alternative hypothesis. We dissected the

interactions concerning the influence of stimulation and task based on the exponent using the

emmeans package with 95% HPD CrI (Lenth, 2018). See the Supplementary Information for

all model comparisons, caterpillar plots, and the posterior predictive check for the most

complex model, Rhat=1).

Table 1 shows the slope of the exponent between the two tasks (learning and overlearning)

and stimulation groups (tRNS and sham) for the best predicted model. It shows that participants

with a high exponent (i.e., low excitation levels) who received tRNS, improved significantly in

terms of lower RTs during the learning task (pd=98.96%). There was no evidence that tRNS

improved performance in the overlearning task.

Table 1| Output of the interaction contrasts between stimulation, task and aperiodic exponent

at baseline using the *emtrends* function (n=75)

Task	Stimulation	Trend Aperiodic exponent	lower and upper HPD
Learning	Sham	-0.36	[-1.30, 0.65]
Overlearning	Sham	0.06	[-0.43, 0.56]
Learning	**tRNS**	**-1.25**	**[-2.64, -0.11]**
Overlearning	tRNS	$-0.50e^{-3}$	[-0.30, 0.27]

Note. The categorical variables task (learning=0 & overlearning=1) and stimulation (sham=0

& tRNS=1) with orthonormal coding. HPD=Highest Posterior Density.

Additionally, we checked the posterior distributions that captures the uncertainty

surrounding the magnitude of an effect. Typically, a posterior distribution higher or equal to

75% (below or above zero) is chosen as a threshold to indicate that an effect is present. The

choice for a certain cut-off criterion depends on the potential risks and benefits of the

intervention (i.e., Ahn et al., 2018), and in this context it means that there is a 75% chance that

the alternative hypothesis (i.e., the presence of an effect) is true. **Figure 4A** shows that there is

a 90% probability that tRNS lowers median RTs on average during both tasks and thus

improves performance (see **Figure S3** for all main effects). The most important effect is the

three-way interaction between tRNS X task X aperiodic exponent at baseline (see **Figure 4B**

and **Figure S3**). Notably, the posterior probability of the presence of a three-way interaction

between tRNS X task X baseline aperiodic exponent is 82% (see **Figure 4B**).

To understand the source of this 3-way interaction we dissected it by running the model for

learning and overlearning separately (see **Figure 4C**). For the learning task the posterior

distribution for the interaction between stimulation and the baseline aperiodic exponent was

91%. We therefore further dissected the model for sham and tRNS separately for the main effect

of the baseline aperiodic exponent in the learning task. We did not find a difference in

performance between those with low and high baseline aperiodic exponent in the sham

condition (posterior distribution=58%). However, when tRNS was applied, those with a high

baseline aperiodic exponent performed better, and those with a low exponent performed worse

(posterior distribution=94%). In contrast, for the overlearning task, the posterior distribution

was 56% indicating no support for an interaction between stimulation and baseline aperiodic

exponent in this task.

Figure 4| Posterior density of tRNS and the three-way interaction between tRNS, task, and baseline aperiodic exponent from the best fitted model. A) The posterior density of stimulation (tRNS) shows that 90% of the posterior distribution is below zero. Indicating that there is a 90% probability that tRNS lowers RTs during the learning and overlearning task. **B)** Posterior distribution of the three-way interaction between tRNS, task, and baseline aperiodic exponent that shows that there is a 82% probability of presence of this interaction. **C)** The left panel indicates the marginal effects of the learning task for low baseline aperiodic exponent values (mean -1 standard deviation (SD)) and high baseline aperiodic exponent values (mean +1 SD). Sham stimulation is indicated in red and tRNS in blue. The right panel indicates the same marginal effects of the overlearning task. This plot shows that tRNS improved performance, but this was restricted to participants with a high baseline aperiodic exponent in the learning task. No effect was found for participants with a low baseline periodic exponent in the learning task, and neither were there any beneficial effects of tRNS in the overlearning task. 95% CrI are indicated.

Sensations

No significant differences arose in terms of felt sensations between the tRNS and sham stimulation group (for statistical details see **Table S3**). Also, no difference was found between the groups in the impact of these sensations on their subjective performance.

Discussion

The aim of the present study was to impact E/I (measured by means of the aperiodic exponent), both directly using tRNS, and indirectly by manipulating the level of skill acquisition (learning/overlearning). We found that the aperiodic exponent decreased after tRNS, indicating an increased E/I. However, we found no effect of task manipulation on the aperiodic exponent, which indicates that the degree of learning a skill did not affect the aperiodic exponent. Finally, we found a three-way interaction: tRNS improved performance of participants with a low baseline aperiodic exponent in the learning task, but there were no effects of tRNS or baseline aperiodic exponent in the overlearning task.

Our finding that tRNS lowered the aperiodic exponent is in line with tRNS experiments in animals showing a reduction of GABAergic activity due to the tRNS excitatory effects (Sanchez-Leon et al., 2020). Our results, which were found using both frequentist and Bayesian approaches, show a decreased aperiodic exponent after applying tRNS related to an increased E/I. This strengthens the notion that delivering electrical random noise to the brain influences the electrophysiological signal. It is thought that tRNS works by enhancing a signal with a near critical signal-to-noise ratio due to introducing noise in the system, described as the phenomenon of stochastic resonance. Previous studies related this increased signal-to-noise ratio from tRNS to enhanced learning, perception, and cognitive performance, which are related to stochastic resonance (Antal & Herrmann, 2016; Cappelletti et al., 2013; Fertonani et al., 2011; Herpich et al., 2019; Snowball et al., 2013; Van der Groen & Wenderoth, 2016). Our findings suggest a working mechanism of tRNS efficacy related to E/I. Whether this mechanism is dependent or independent from stochastic resonance is a question for further research.

Contrary to our expectations, we did not find an effect of task manipulation (i.e., mathematical learning/overlearning) on the aperiodic exponent. It is likely that the effect of our task manipulation was not strong enough. Manipulation on the electrophysiological level by

means of tRNS is a more direct approach to target the aperiodic exponent and is likely to yield

a stronger neuronal effect than cognitive manipulation. This interpretation is in line with the

view that brain stimulation can amplify the cognitive and neural effects of otherwise purely

behavioral approaches (Cappelletti et al., 2013; Cohen Kadosh et al., 2012; Snowball et al.,

2013). While another potential explanation can be attributed to the efficacy of our task

manipulation paradigm, this is unlikely; participants in the learning task did not reach a plateau

of performance improvement, while participants who completed the overlearning task clearly

did show this plateau (see **Figure 2**). Another potential explanation is that both mathematical

learning and overlearning increase the aperiodic exponent (i.e., lower E/I). This explanation is

in line with our finding that the exponent increased for participants in the sham stimulation

group (see **Figure 3**). But this is in contrast to Shibata et al. (2017), who found that perceptual

overlearning led to a reduction in E/I, possibly to protect a newly formed memory trace from

subsequent new information. An alternative explanation, which we elaborate upon later, is that

E/I measures based on EEG and MRS reflect different aspects of E/I.

Previous studies have shown that the efficacy of tRNS depends on the individual baseline

cognitive ability or neural activity (Evans et al., 2018; Frank et al., 2018; Harty & Cohen

Kadosh, 2019). We have extended these findings by showing that enhanced skill acquisition by

tRNS is based on the participants' baseline E/I. First, our results show that the effect of tRNS

is best explained when considering the moderating effects of baseline aperiodic exponent and

task manipulation. To illustrate, the posterior distribution of the three-way-interaction for the

learning and overlearning data indicates the presence of such an interaction effect. Stimulation

improved performance for those with lower E/I (as reflected by a higher aperiodic exponent).

This effect was present only in the learning task. In the overlearning task, tRNS had no effect.

A possible explanation is that participants with low E/I levels benefit more from tRNS

compared to participants with high E/I but only when the task is difficult, indicating an optimum

level depending on task difficulty (Krause et al., 2013; Shibata et al., 2017). This explanation

fits also with the stochastic resonance framework, which predicts non-beneficial or even

detrimental effects when random noise is introduced to an already optimal system (McDonnell

& Ward, 2011). This reveals that the baseline aperiodic exponent is an important predictor of

tRNS efficacy. Thus, we found that tRNS, an excitatory form of neurostimulation, as supported

also by our results, is more beneficial during the learning task (i.e., for low E/I participants)

than during the overlearning task.

As mentioned previously, Shibata et al. (2017) showed that overlearning relates to a shift

from increased E/I, which occurs in learning, to a reduced E/I. While we did not find such a

reduction, our results suggest that in the learning task those with higher E/I at baseline will

perform better than those with lower E/I, unless intervening with tRNS, and in the overlearning

task stimulation is not dependent on E/I. These results suggest, similar to Shibata et al's work,

that E/I is involved in learning. However, our findings indicate that this involvement may not

only be due to E/I alterations related to skill acquisition, but also to the participants' E/I level

at baseline.

While at the beginning of our project some of our predictions were rooted in studies that

used MRS-based E/I measures, our results together with recent findings in the literature, suggest

that there are some discrepancies between EEG-based E/I and MRS-based E/I that must be

acknowledged and addressed in the future. First, some studies found that EEG-based E/I

increases with age (Cellier et al., 2021; Van Bueren et al., 2022), rather than decrease as was

found using MRS (Cohen Kadosh et al., 2015; Zacharopoulos et al., 2021b). Second, our lack

of replication of the effect of learning (increased E/I change) and overlearning (decreased E/I

change) on E/I between before and after task manipulation (Shibata et al., 2017) could be rooted

in the different methodologies used to assess E/I. These discrepancies raise the question of

whether E/I measures based on EEG and MRS reflect different aspects of E/I. Indeed, MRS-

based E/I measure is likely to reflect intra- or extracellular activity, while EEG-based E/I is

based on extracellular concentrations (Buzsáki et al., 2012; Dyke et al., 2017). To tentatively

examine whether MRS-based E/I and EEG-based E/I reflect different aspects of E/I, the results

are incidental, or if the two measures reflect a shared variance of E/I, we used an independent

dataset that would allow us to assess the link between E/I measures derived by MRS and EEG

in 20 young adults (see Supplementary Information). We found that higher glutamate/GABA

measured with MRS (i.e., higher E/I) was significantly associated with an increased aperiodic

exponent (i.e., lower E/I) over the left IPS. No relation was found for the left MFG (see **Figure**

S4). This shows that MRS and EEG may measure different aspects of E/I. These preliminary

results highlight the need to further examine the origin of E/I in MRS and EEG to progress our

basic understanding as well as utilizing these measures for clinical applications.

Our results elucidate one of the underlying mechanisms of tRNS in cognitive and

electrophysiological studies highlighting the role of aperiodic activity, and thus E/I balance.

This study indicates the important role of baseline E/I in skill acquisition and tRNS efficacy.

More specifically, participants with a higher aperiodic exponent (i.e., lower E/I) benefit more

from tRNS during a learning task compared to participants with a lower aperiodic exponent

(i.e., higher E/I). The beneficial effect of tRNS was only found during learning and not during

overlearning. This new understanding has important implications when considering to whom,

when, and in the future what dose of tRNS should be delivered, and increase the importance of

a personalized neurostimulation approach.

Materials and Methods

Participants and Ethical Permission

One hundred and two right-handed participants participated in the study. None of the

participants reported a history of psychiatric, neurological, or skin conditions and all met the

safety criteria for tES participants. All volunteers were naïve to the aim of the study, and written

informed consent was obtained. We ensured that all participants had no more than one cup of

coffee or other sources of caffeine within 1 hour before the start of the study. We excluded

seven participants who displayed an overall accuracy below 70% in the learning (Sham=4 and

tRNS=3) or the overlearning task (Sham=1 and tRNS=1) as they were non-compliant with the

task. Similarly, we excluded four participants because they showed no arithmetic facts learning

in the learning or overlearning (Sham=2 and tRNS=2) tasks, as indexed by the lack of a

significant negative linear regression coefficient of response times as a function of block. Five

participants were excluded due to malfunction of the software (Sham=3 and tRNS=2). For the

electrophysiological analysis, 11 participants were excluded due to artifacts during the pre or

post resting state (rs)-EEG recording (more than 25% of their data were rejected) (see **Table 1**

for demographic data). The final sample (n=75) was composed of 22 participants in the sham-

learning condition, 21 in the sham-overlearning condition, 16 in the tRNS-learning condition,

and 16 in the tRNS-overlearning condition. We excluded three participants (Sham=3) from the

frequentist and Bayesian ANCOVA analyses as they were outliers on Cooks distance with RTs

as outcome variable. For the brms models, we removed outliers on trial level grouped by task

using the median absolute value (mad) with a threshold of 3 as indicated by Leys et al. (2013)

(9.00% trials were removed). The study complied with the standards set by the Declaration of

Helsinki and approved by the ethical advisory committee of the Faculty of Experimental

Psychology at Oxford University (Protocol Number: IDREC, C2-2014-033). The methods of

this study are on Open Science Framework (see https://osf.io/y4xar). However, the analyses

(i.e., neuronal avalanches) presented in this preregistration did not yield significant results (see

Figure S6). We later came across the work on the aperiodic exponent as a measure of E/I,

which we used in this study.

Table 1| Demographic data of the stimulation conditions

Final Sample (N=75)		Stimulation Conditions				
		Sham-Learning (n=22)	tRNS-Learning (n=16)	Sham-Overlearning (n=21)	tRNS-Overlearning (n=16)	
	M(SD)	M(SD)	M(SD)	M(SD)	M(SD)	p*
Age (years)	23.40(4.27)	23.00(3.87)	25.00(5.54)	23.00(2.39)	24.17(5.33)	.45
Aperiodic Baseline	1.43(0.45)	1.38(.53)	1.63(.49)	1.38(.29)	1.37(0.46)	.29
Aperiodic Post	1.38(0.41)	1.39(.45)	1.32(0.21)	1.46(.50)	1.32(.41)	.88
Sex	44 females : 31 males	16 females : 6 males	7 females : 9 males	15 females : 6 males	6 females : 10 males	

401 *p-value of a between-groups ANOVA to compare stimulation groups

Baseline Matching

We investigated whether the participants in the four conditions (i.e., sham-learning, tRNS-
learning, sham-overlearning, tRNS-overlearning) differed in median reaction times (RTs) and
accuracy in the baseline task. A univariate ANOVA with condition as between-participants
factors showed no significant differences for accuracy ($F(3,98)=1.73$, $p=.166$). Subsequently,
all incorrect responses were removed from the baseline task (16% of all trials), and median RTs
for each subject were calculated. Another univariate ANOVA showed no significant differences
for median RTs ($F(3,98)=.11$, $p=.951$) at baseline between the different groups. Furthermore,
there were no differences between the groups regarding subjective levels of sleepiness before
the start of the experiment ($F(3,98)=.58$, $p=.629$). In addition, we ran an exploratory univariate
ANOVA to determine if there were any differences regarding baseline aperiodic activity values
before applying tRNS. We found no significant differences between the groups at baseline for

aperiodic activity ($F(3,71) = 1.25$, $p = .297$) after outlier removal for the ANCOVA. These results

suggest similar baseline values across active and sham stimulation groups.

Tasks

Baseline Ability Task

At the beginning of the session, participants solved four multiplication problems to become

familiar with the testing procedure. Afterwards, participants solved 10 new multiplication

problems to evaluate their baseline ability (see **Table S12**). Every multiplication problem

presented consisted of two-digit times one-digit operands with a two-digit answer (e.g., 16 x

3=48). None of the one-digit operands involved the digits 0 or 1 to minimize variations in

difficulty. Furthermore, the two-digit operand was larger than 15 and not a multiple of 10.

At the beginning of the task, participants pressed the spacebar when ready to solve an

arithmetic problem. Then, each trial started with a fixation screen (**Figure 6A**) after which a

symbol of a headphone set appeared in the middle of the screen, and the arithmetic problem

was presented auditorily. A symbol of a microphone appeared immediately after the arithmetic

problem, and participants could say aloud the response. A noise-sensitive microphone captured

the participant's responses through a Chronos box (Science Plus Group). Lastly, the words

"Retrieve" and "Calculate" appeared on the left and right side of the screen; Participants

indicated whether they had used a retrieval or calculation strategy when solving the arithmetic

problem by pressing the left or right mouse button, respectively.

Participants were instructed to wear a headphone to cancel out any surrounding noise, and

to speak clearly and loudly in the microphone without mumbling or clearing the throat (e.g.,

saying "eh-em", which would be registered as a response). Lastly, participants were informed

that there was no time limit for answering, and they were urged to avoid errors.

Figure 5| A schematic overview of the task structure and experimental protocol. A) First, a fixation screen was shown. Subsequently, a multiplication was presented by voice recording through a headphone. Hereafter, participants were shown a microphone symbol to indicate that they could say the answer into the microphone. This was followed by a 200 ms delay period. Lastly, participants indicated by clicking the left or right mousepad on the keyboard whether they retrieved or calculated the answer. **B)** First, a pre rs-EEG was measured of 8 minutes. Subsequently, a training was presented that contained 4 different multiplication problems. Based on baseline performance, participants either completed the learning or the overlearning task. One block in both the learning and overlearning task consisted of 10 multiplications with 18 blocks, and 180 trials in total. Participants received 20 minutes, 1 mA tRNS during either the learning or overlearning task or sham stimulation. Next, the transfer task was presented with new arithmetic problems containing 10 multiplication repeated three times. The recall task contained the identical multiplications as either the learning or the overlearning task and was repeated three times. Lastly, another post rs-EEG measurement of 8 minutes was assessed. **C)** Placement of the stimulation electrodes over F3 and F4.

Learning and Overlearning Condition

In total, 180 multiplication problems were administered in both the overlearning and the

learning condition (see **Table S13**). The structure of the tasks was identical to the baseline task

(see **Figure 5A**). Both conditions consisted of 18 blocks, comprised of the same number of

trials and were presented in a fixed order. After three blocks, participants had a one-minute

break. Therefore, in the learning condition, a subset of ten problems was presented once in each

block. In the overlearning condition, a subset of five problems was selected, which was

presented twice in each block (i.e., less information to learn in comparison to the learning

condition). This manipulation allowed us to use the same task and duration yet influencing the

stage of skill acquisition that the participants reached. The design was based on a small pilot

study (n=4) in which six multiplication problems were repeated four times. A plateau in

performance improvement was visible after block four.

Transcranial Random Noise Stimulation

TRNS was applied over the bilateral dorsolateral prefrontal (dlPFC; F3 and F4) cortices, as

defined by the international 10-20 system for EEG recording (see **Figure 5C**). Two Pistim

Ag/AgCl electrodes were used with a 1 cm radius and a surface area of 3.14 cm^2 each. A current

was delivered through these electrodes in the form of high frequency noise (100-500 Hz) by a

multichannel transcranial current stimulator (Starstim 8 device, Neuroelectrics, Barcelona). The

impedances of the Pistim electrodes were held at < 10 kΩ and intensity of the current was 1

mA peak-to-peak which has been shown to be safe and painless (Ambrus, Paulus, & Antal,

2010; Fertonani, Pirulli, & Miniussi, 2011). Duration of the stimulation was set to 20 minutes

after onset of the task for the stimulation condition and 30 s for the sham condition with a 15 s

ramp-up and a 15 s ramp-down. This provided the initial skin sensations experienced during

stimulation. Both the participants and the experimenter were blinded to the stimulation

condition. After completing the experiment, all participants filled out a questionnaire in which

they were asked whether they felt any sensations during stimulation (i.e., itchiness, pain,

burning, warmth/heat, pinching, iron taste, and fatigue) and if these sensations affected their

performance. Unfortunately, due to an experimental error, we did not ask them whether they

think they received sham or active stimulation. A follow up study that used the same parameters

and a similar paradigm to this one (Sheffield et al., 2020), on a similar population, participants

in both groups reported being in the stimulation condition at approximately the same rate (sham:

78%; active tRNS: 79%; $\chi(1)$=0.03, p=1). This independent data is supported by our data, which

did not find differences in sensations between both groups (see Results section).

Electrophysiological Data

Rs-EEG recordings were made before baseline allocation and at the end of the experiment

as stated in our preregistration (see **Figure 5B**). Electrophysiological data were obtained with

32 Ag/AgCl electrodes according to the international 10/20 EEG system using the wireless

ENOBIO 32 sensor system (Neuroelectrics, Barcelona) at 500 Hz with no online filters. Note

that we also used recorded rs-EEG from the two stimulation NG Pistim Ag/AgCl electrodes

(F3 and F4). The impedances of the electrodes were held below 5 kΩ. The ground consisted of

the active common mode sense (CMS) and passive driven right leg (DRL) electrode which were

positioned on the right mastoid and connected by adhesive electrodes. Both the pre rs-EEG and

the post rs-EEG had a duration of eight minutes, in which the participants had their eyes open

while watching a fixation point in the middle of the screen in order to avoid mental and muscular

activity.

Data Pre-processing

EEG data were pre-processed using EEGlab toolbox (v14.1.0) (Delorme & Makeig, 2004)

in Matlab software (R2020b). A high-pass filter of 0.1 Hz was applied to minimize slow drifts

and a notch filter at 50 Hz was applied to minimize line noise interference with the signal, and

any data recorded before the presentation of the fixation point was removed. Every data file

was manually checked and high-amplitude artefacts due to muscle movement, sweating or

electrode malfunction were rejected. After pre-processing, Independent Component Analysis

(ICA) was performed to remove stereotyped artefacts such as eye movements (e.g., blinks),

heart rate activity and muscular activity. A maximum of six components per data file were

removed, and a maximum of five bad channels were interpolated. EEG segments that contained

artefacts that could not be removed by ICA were visually inspected and rejected from the

analysis (Delorme et al., 2007). If more than 25 percent of the rs-EEG data were rejected after

pre-processing and ICA, data of the subject were discarded from analysis.

Aperiodic Activity Computation

The rs-EEG data of the remaining participants were separated in 2-second segments with an overlap of 1 second and windowed with a Hann window. Subsequently, data were transformed into the frequency domain via Fast Fourier Transformation (FFT). The FFT was exported from Matlab and important in Python (v.3.7.0; Van Rossum & Drake, 1995), and subsequently analyzed with the FOOOF package (v 1.0.0; Donoghue et al., 2020) over the 1-40 Hz range. This package allows for the decomposition between periodic and aperiodic components of the FFT. The aperiodic activity was calculated for the midline frontal electrode Fz which is the closest electrode to our stimulation electrodes F3 and F4.

Experimental Design and Statistical Analyses

First, participants completed the Stanford sleepiness scale (SSS), which is an introspective measure of subjective sleepiness and the Multidimensional Mood Questionnaire (MDBF), to assess alertness, good-bad mood, tiredness, calmness, and restlessness.

Then, participants completed an rs-EEG pre-measurement of eight minutes where they were informed to sit as still as possible (see **Figure 5B**). Then, a training was presented that consisted of four different arithmetic multiplications. Hereafter, the baseline task was started and a variance minimization procedure (based on response times) followed the baseline task in a double-blind fashion to determine which participants would be allocated to which group (see https://osf.io/y4xar for a detailed explanation of this procedure). This procedure is superior to random assignment for assigning participants to groups before an intervention (Sella et al., 2021), as it results in better matching. The stimulation started together with the learning or the overlearning task. Subsequently, the participants completed a transfer task and lastly a recall task. The effect of stimulation on transfer and recall is of different theoretical interest, but the effect of E/I is. However, E/I did not show an effect on transfer and recall (see **Tables S4-S11**), but there was an effect of tRNS. These results will therefore be reported elsewhere (van Bueren

539   et al., in preparation). More information about the procedure of these tasks, which is beyond the

540   scope of the present manuscript, can be found on https://osf.io/y4xar. At the end of the

541   behavioral tasks, an 8 minute rs-EEG measurement was recorded.

Behavioral Data Cleaning

We excluded responses below 200 ms due to possible noises picked up by the microphone

or mumbling of the subject (0.89% for the baseline task, 2.74% for the learning task, and 7.07%

for the overlearning task). We also excluded wrong responses from the baseline ability task

(16.32%), the learning task (10.65%), and the overlearning task (7.7%). We calculated the

median RTs for each participant in the baseline task and in each block of the learning and

overlearning tasks.

Calculation of the Amount of Learning: Plateau of Performance Improvement

A plateau in performance improvement was computed following the next procedure: the

distribution of RTs from the first block was compared to the distribution of RTs from the next

block using the Wilcoxon test. When no significant difference was found in RTs between the

actual block and the remaining blocks, the actual block was considered as the plateau point.

Therefore, the plateau point is a number between 2 and 18 and an earlier plateau point indicates

a higher amount of overlearning.

Statistical Analyses

All numerical independent variables were standardized to avoid multicollinearity issues. All

inferential statistics reported in the present study were obtained with RStudio version 4.1.1

using the the *wilcox.test* function for the Mann-Whitney U test, the chisq.test for the Chi-Square

Test, the *aov* function for ANCOVA and Univariate ANOVA, the *lm* function for the

exploratory regression, and the *brms* package for the Bayesian mixed effect models (Bürkner,

2017) which are robust to normality violation and can deal with complex models. We used the

open-source project JASP to run the Bayesian ANCOVA (Version 0.14.1.0; JASP Team, 2021).

We originally used *glmer* for our analysis but due to model complexity we revert to *brms*. All

Bayesian models were ran with 4000 iterations (2000 for warmup), 4 chains each, 16 cores.

Additionally, orthonormal contrast coding was used to reliably dissect interactions with the

emmeans package. Due the right skewness of the response times, the shifted lognormal family

was used. Also, all continuous independent variables were centered.

Acknowledgements

570 This research was funded in whole, or in part, by the Wellcome Trust (203139/Z/16/Z). For

571 the purpose of Open Access, the author has applied a CC BY public copyright licence to any

572 Author Accepted Manuscript version arising from this submission. We would like to thank dr.

573 James Sheffield with his help concerning the electrophysiological analyses and design, and dr.

574 George Zacharopoulos with providing MRS-based E/I data.

Competing Interests

576 All authors declare no competing interests.

References

Ahmad, J., Ellis, C., Leech, R., Voytek, B., Garces, P., Jones, E., Buitelaar, J., Loth, E., dos

Santos, F. P., Amil, A. F., Verschure, P. F. M. J., Murphy, D., & McAlonan, G. (2022).

From mechanisms to markers: novel noninvasive EEG proxy markers of the neural

excitation and inhibition system in humans. *Translational Psychiatry*, *12*(1), 1–12.

https://doi.org/10.1038/s41398-022-02218-z

Ahn, H., Suchting, R., Woods, A. J., Miao, H., Green, C., Cho, R. Y., Choi, E., & Fillingim,

R. B. (2018). Bayesian analysis of the effect of transcranial direct current stimulation on

experimental pain sensitivity in older adults with knee osteoarthritis: randomized sham-

controlled pilot clinical study. *Journal of Pain Research*, *11*, 2071.

https://doi.org/10.2147/JPR.S173080

Ambrus, G. G., Paulus, W., & Antal, A. (2010). Cutaneous perception thresholds of electrical

stimulation methods: Comparison of tDCS and tRNS. *Clinical Neurophysiology*,

121(11), 1908–1914. https://doi.org/10.1016/j.clinph.2010.04.020

Anderson, J. R. (2013). Cognitive skills and their acquisition. *Psychology Press*.

Antal, A., & Herrmann, C. S. (2016). Transcranial alternating current and random noise

stimulation: Possible mechanisms. *Neural Plasticity*, *2016*, 1–12.

https://doi.org/10.1155/2016/3616807

Barron, H. C., Vogels, T. P., Emir, U. E., Makin, T. R., O'shea, J., Clare, S., ... & Behrens, T.

E. J. (2016). Unmasking latent inhibitory connections in human cortex to reveal dormant

cortical memories. *Neuron*, *90*(1), 191-203. https://doi.org/10.1016/j.neuron.2016.02.031

Bürkner, P. C. (2017). brms: An R package for Bayesian multilevel models using

Stan. *Journal of statistical software*, *80*, 1-28.

Buzsáki, G., Logothetis, N., & Singer, W. (2013). Scaling brain size, keeping timing:

evolutionary preservation of brain rhythms. *Neuron*, *80*(3), 751–764.

https://doi.org/10.1016/J.NEURON.2013.10.002

Cappelletti, M., Gessaroli, E., Hithersay, R., Mitolo, M., Didino, D., Kanai, R., Cohen

Kadosh, R., & Walsh, V. (2013). Transfer of cognitive training across magnitude

dimensions achieved with concurrent brain stimulation of the parietal lobe. *Journal of*

Neuroscience, *33*(37). https://doi.org/10.1523/JNEUROSCI.1692-13.2013

Cellier, D., Riddle, J., Petersen, I., & Hwang, K. (2021). The development of theta and alpha

neural oscillations from ages 3 to 24 years. *Developmental Cognitive Neuroscience*, *50*,

100969. https://doi.org/10.1016/J.DCN.2021.100969

Cohen Kadosh, R., Levy, N., O'Shea, J., Shea, N., & Savulescu, J. (2012). The neuroethics of

non-invasive brain stimulation. *Current Biology, 22*(4), R108–R111.

https://doi.org/10.1016/j.cub.2012.01.013

Cohen Kadosh, K., Krause, B., King, A. J., Near, J., & Cohen Kadosh, R. (2015). Linking

GABA and glutamate levels to cognitive skill acquisition during development. *Human*

Brain Mapping, *36*(11), 4334–4345. https://doi.org/10.1002/HBM.22921

Deelchand, D. K., Adanyeguh, I. M., Emir, U. E., Nguyen, T. M., Valabregue, R., Henry, P.

G., ... & Öz, G. (2015). Two-site reproducibility of cerebellar and brainstem

neurochemical profiles with short-echo, single-voxel MRS at 3T. *Magnetic resonance in*

medicine, *73*(5), 1718-1725. https://doi.org/10.1002/mrm.25295

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. *Journal of Neuroscience*

Methods, *134*, 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Delorme, A., Sejnowski, T., & Makeig, S. (2007). Enhanced detection of artifacts in EEG

data using higher-order statistics and independent component analysis. *NeuroImage*,

34(4), 1443–1449. https://doi.org/10.1016/j.neuroimage.2006.11.004

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A.

H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing

neural power spectra into periodic and aperiodic components. *Nature Neuroscience 2020*

23:12, *23*(12), 1655–1665. https://doi.org/10.1038/s41593-020-00744-x

Dorrn, A. L., Yuan, K., Barker, A. J., Schreiner, C. E., & Froemke, R. C. (2010).

Developmental sensory experience balances cortical excitation and inhibition. *Nature*

2010 465:7300, *465*(7300), 932–936. https://doi.org/10.1038/nature09119

Emir, U. E., Tuite, P. J., & Öz, G. (2012). Elevated pontine and putamenal GABA levels in

mild-moderate Parkinson disease detected by 7 tesla proton MRS. *PloS one*, *7*(1),

e30918. https://doi.org/10.1371/journal.pone.0030918

Evans, C., Banissy, M. J., & Charlton, R. A. (2018). The efficacy of transcranial random

noise stimulation (tRNS) on mood may depend on individual differences including age

and trait mood. *Clinical Neurophysiology*, *129*(6), 1201-1208.

https://doi.org/10.1016/j.clinph.2018.03.012

Fertonani, A., & Miniussi, C. (2017). Transcranial electrical stimulation: what we know and

do not know about mechanisms. *The Neuroscientist*, *23*(2), 109–123.

https://doi.org/10.1177/1073858416631966

Fertonani, A., Pirulli, C., & Miniussi, C. (2011). Random noise stimulation improves

neuroplasticity in perceptual learning. *Journal of Neuroscience*, *31*(43), 15416-15423.

https://doi.org/10.1523/JNEUROSCI.2002-11.2011

Frank, B., Harty, S., Kluge, A., & Cohen Kadosh, R. (2018). Learning while multitasking:

short and long-term benefits of brain stimulation. *Ergonomics*, *61*(11), 1454-1463.

https://doi.org/10.1080/00140139.2018.1563722

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance

from field potentials. *NeuroImage*, *158*, 70–78.

https://doi.org/10.1016/J.NEUROIMAGE.2017.06.078

Harty, S., & Cohen Kadosh, R. (2019). Suboptimal engagement of high-level cortical regions

predicts random-noise-related gains in sustained attention. *Psychological Science*, *30*(9),

1318-1332. 095679761985665. https://doi.org/10.1177/0956797619856658

He, W., Donoghue, T., Sowman, P. F., Seymour, R. A., Brock, J., Crain, S., Voytek, B., &

Hillebrand, A. (2019). Co-increasing neuronal noise and beta power in the developing

brain. *BioRxiv*, 839258. https://doi.org/10.1101/839258

Herpich, F., Melnick, M. D., Agosta, S., Huxlin, K. R., Tadin, D., & Battelli, L. (2019).

Boosting learning efficacy with noninvasive brain stimulation in intact and brain-

damaged humans. *Journal of Neuroscience*, *39*(28), 5551-5561.

https://doi.org/10.1523/JNEUROSCI.3248-18.2019

Krause, B., & Cohen Kadosh, R. (2014). Not all brains are created equal: the relevance of

individual differences in responsiveness to transcranial electrical stimulation. *Front Syst*

Neurosci, *8*, 25. https://doi.org/10.3389/fnsys.2014.00025

Krause, B., Márquez-Ruiz, J., & Cohen Kadosh, R. (2013). The effect of transcranial direct

current stimulation: a role for cortical excitation/inhibition balance? *Frontiers in Human*

Neuroscience, *7*. https://doi.org/10.3389/fnhum.2013.00602

Lansbergen, M. M., Arns, M., Van Dongen-Boomsma Martine, M., Spronk, D., & Buitelaar,

J. K. (2011). The increase in theta/beta ratio on resting-state EEG in boys with attention-

deficit/hyperactivity disorder is mediated by slow alpha peak frequency. *Progress in*

Neuro-Psychopharmacology and Biological Psychiatry, *35*(1), 47–52.

https://doi.org/10.1016/J.PNPBP.2010.08.004

Lenth, R. (2018) Emmeans: estimated marginal means, aka least-squares means. R package

version 1.2.2. https://CRAN.R-project.org/package=emmeans

Leys, C., Ley, C., Klein, O., Bernard, P., & Licata, L. (2013). Detecting outliers: Do not use

standard deviation around the mean, use absolute deviation around the median. *Journal*

of experimental social psychology, *49*(4), 764-766.

https://doi.org/10.1016/j.jesp.2013.03.013

McDonnell, M. D., & Ward, L. M. (2011). The benefits of noise in neural systems: bridging

theory and experiment. *Nature Reviews Neuroscience*, *12*(7), 415-425.

https://doi.org/10.1038/nrn3061

Molina, J. L., Voytek, B., Thomas, M. L., Joshi, Y. B., Bhakta, S. G., Talledo, J. A.,

Swerdlow, N. R., & Light, G. A. (2020). Memantine effects on electroencephalographic

measures of putative excitatory/inhibitory balance in schizophrenia. *Biological*

Psychiatry: Cognitive Neuroscience and Neuroimaging, *5*(6), 562–568.

https://doi.org/10.1016/J.BPSC.2020.02.004

Newell, A., & Rosenbloom, P. (1981). Mechanisms of skill acquisition. *Cognitive skills and*

their acquisition.

Nishiyama, M., Togashi, K., Aihara, T., & Hong, K. (2010). GABAergic activities control

spike timing- and frequency-dependent long-term depression at hippocampal excitatory

synapses. *Frontiers in Synaptic Neuroscience*, *2*, 22.

https://doi.org/10.3389/FNSYN.2010.00022/BIBTEX

Öz, G., & Tkáč, I. (2011). Short-echo, single-shot, full-intensity 1H MRS for neurochemical

profiling at 4T: validation in the cerebellum and brainstem. *Magnetic resonance in*

medicine, 65(4), 901-910. https://doi.org/10.1002/mrm.22708

Palva, J. M., Zhigalov, A., Hirvonen, J., Korhonen, O., Linkenkaer-Hansen, K., & Palva, S.

(2013). Neuronal long-range temporal correlations and avalanche dynamics are

correlated with behavioral scaling laws. *Proceedings of the National Academy of*

Sciences of the United States of America, *110*(9), 3585–3590.

https://doi.org/10.1073/pnas.1216855110

Potok, W., Van der Groen, O., Bächinger, M., Edwards, D., & Wenderoth, N. (2022).

Transcranial random noise stimulation modulates neural processing of sensory and motor

circuits, from potential cellular Mechanisms to behavior: A scoping review. *ENeuro*,

9(1). https://doi.org/10.1523/ENEURO.0248-21.2021

Robertson, M. M., Furlong, S., Voytek, B., Donoghue, T., Boettiger, C. A., & Sheridan, M.

A. (2019). EEG power spectral slope differs by ADHD status and stimulant medication

exposure in early childhood. *Journal of Neurophysiology, 122(6),* 2427-2437.

https://doi.org/10.1152/JN.00388.2019

Rufener, K. S., Ruhnau, P., Heinze, H.-J., & Zaehle, T. (2017). Transcranial random noise

stimulation (tRNS) shapes the processing of rapidly changing auditory information.

Frontiers in Cellular Neuroscience, *0*, 162. https://doi.org/10.3389/FNCEL.2017.00162

Sánchez-León, C. A., Sánchez-López, Á., Gómez-Climent, M. A., Cordones, I., Cohen

Kadosh, R., & Márquez-Ruiz, J. (2021). Impact of chronic transcranial random noise

stimulation (tRNS) on GABAergic and glutamatergic activity markers in the prefrontal

cortex of juvenile mice. In *Progress in brain research* (Vol. 264, pp. 323-341). Elsevier.

Sarkar, A., Dowker, A., & Cohen Kadosh, R. (2014). Cognitive enhancement or cognitive

cost: trait-specific outcomes of brain stimulation in the case of mathematics anxiety. *The*

Journal of Neuroscience : The Official Journal of the Society for Neuroscience, *34*(50),

16605–16610. https://doi.org/10.1523/JNEUROSCI.3129-14.2014

Shew, W. L., Yang, H., Petermann, T., Roy, R., & Plenz, D. (2009). Neuronal avalanches

imply maximum dynamic range in cortical networks at criticality. *Journal of*

Neuroscience, *29*(49). https://doi.org/10.1523/JNEUROSCI.3864-09.2009

Shibata, K., Sasaki, Y., Bang, J. W., Walsh, E. G., Machizawa, M. G., Tamaki, M., Chang,

L.-H., & Watanabe, T. (2017). Overlearning hyperstabilizes a skill by rapidly making

neurochemical processing inhibitory-dominant. *Nature Neuroscience*, 20(3), 470-475.

https://doi.org/10.1038/nn.4490

Shriki, O., Alstott, J., Carver, F., Holroyd, T., Henson, R. N. A., Smith, M. L., Coppola, R.,

Bullmore, E., & Plenz, D. (2013). Neuronal avalanches in the resting MEG of the human

brain. *Journal of Neuroscience*, *33*(16). https://doi.org/10.1523/JNEUROSCI.4286-

12.2013

Snowball, A., Tachtsidis, I., Popescu, T., Thompson, J., Delazer, M., Zamarian, L., Zhu, T., &

Cohen Kadosh, R. (2013). Long-term enhancement of brain function and cognition using

cognitive training and brain stimulation. *Current Biology, 23*(11), 987-992.

https://doi.org/10.1016/j.cub.2013.04.045

Soher, B. J., Semanchuk, P., Todd, D., Steinberg, J., & Young, K. (2011, May). VeSPA:

integrated applications for RF pulse design, spectral simulation and MRS data analysis.

In *Proc Int Soc Magn Reson Med* (Vol. 19, No. 19, p. 1410).

Terney, D., Chaieb, L., Moliadze, V., Antal, A., & Paulus, W. (2008). Increasing human brain

excitability by transcranial high-frequency random noise stimulation. *Journal of*

Neuroscience, *28*(52), 14147–14155. https://doi.org/10.1523/JNEUROSCI.4248-08.2008

Tzelgov, J., Yehene, V., Kotler, L., & Alon, A. (2000). Automatic comparisons of artificial

digits never compared: learning linear ordering relations. *Journal of Experimental*

Psychology: Learning, Memory, and Cognition, *26*(1), 103.

https://doi.org/10.1037/0278-7393.26.1.103

Uhlhaas, P. J., & Singer, W. (2010). Abnormal neural oscillations and synchrony in

schizophrenia. *Nature Reviews Neuroscience 2010 11:2*, *11*(2), 100–113.

https://doi.org/10.1038/nrn2774

Van Bueren, N. E. R., van der Ven, S. H. G., Roelofs, K., Cohen Kadosh, R., & Kroesbergen,

E. H. (2022). Predicting math ability using working memory, number sense, and

neurophysiology in children and adults. *Brain Sciences*, *12*(5), 550.

https://doi.org/10.3390/brainsci12050550

Van der Groen, O., & Wenderoth, N. (2016). Transcranial random noise stimulation of visual

cortex: Stochastic resonance enhances central mechanisms of perception. *Journal of*

Neuroscience, *36*(19), 5289–5298. https://doi.org/10.1523/JNEUROSCI.4519-15.2016

Van Rossum, G., & Drake Jr, F. L. (1995). Python tutorial: Centrum voor Wiskunde en

Informatica Amsterdam. *The Netherlands*.

Voytek, B., & Knight, R. T. (2015). Dynamic network communication as a unifying neural

basis for cognition, development, aging, and disease. *Biological Psychiatry*, *77*(12),

1089–1097. https://doi.org/10.1016/J.BIOPSYCH.2015.04.016

Voytek, B., Kramer, M. A., Case, J., Lepage, K. Q., Tempesta, Z. R., Knight, R. T., &

Gazzaley, A. (2015). Age-related changes in 1/f neural electrophysiological noise.

Journal of Neuroscience, *35*(38), 13257–13265.

https://doi.org/10.1523/JNEUROSCI.2332-14.2015

Wijtenburg, S. A., West, J., Korenic, S. A., Kuhney, F., Gaston, F. E., Chen, H., Roberts, M.,

Kochunov, P., Hong, L. E., & Rowland, L. M. (2017). Glutamatergic metabolites are

associated with visual plasticity in humans. *Neuroscience Letters*, *644*, 30–36.

https://doi.org/10.1016/J.NEULET.2017.02.020

Zacharopoulos, G., Emir, U., & Cohen Kadosh, R. (2021a). The cross-sectional interplay

between neurochemical profile and brain connectivity. *Human Brain Mapping*, *42*(9),

2722–2733. https://doi.org/10.1002/HBM.25396

Zacharopoulos, G., Sella, F., Cohen Kadosh, K., Hartwright, C., Emir, U., & Cohen Kadosh,

R. (2021b). Predicting learning and achievement using GABA and glutamate

concentrations in human development. *PLOS Biology*, *19*(7), e3001325.

https://doi.org/10.1371/JOURNAL.PBIO.3001325

Zacharopoulos, G., Sella, F., & Cohen Kadosh, R. (2021c). The impact of a lack of

mathematical education on brain development and future attainment. *Proceedings of the*

National Academy of Sciences of the United States of America, *118*(24).

https://doi.org/10.1073/PNAS.2013155118/-/DCSUPPLEMENTAL