Quantum pBac: An effective, high-capacity *piggyBac*-based gene

integration vector system for unlocking gene therapy potential

Wei-Kai Hua[1,*], Jeff C. Hsu[1,*], Yi-Chun Chen[1,*], Peter S. Chang[1], Kuo-Lan Karen

Wen[1], Po-Nan Wang[2], Yi-Shan Yu[1], Ying-Chun Chen[1], I-Cheng Cheng[1], Sareina

Chiung-Yuan Wu[1,†]

[1]GenomeFrontier Therapeutics, Inc. Taipei City, Taiwan (R.O.C.)

[2]Division of Hematology, Chang Gung Medical Foundation, Linkou Branch,

Taipei City, Taiwan (R.O.C.)

[*] These authors contributed equally to this work and should be considered joint

first author. Their authorship order is interchangeable

[†]Correspondence should be addressed to:

S.C.-Y.W. (sareina@genomefrontier.com) 23F, Bldg. C, No. 95, Sec. 1, Xintai

5th Rd., Xizhi Dist., New Taipei City 221416; +886-2-26558766;

This work was performed in Taipei City, Taiwan (R.O.C.)

Short title: *Quantum pBac* for unlocking gene therapy potential

Abstract

Recent advances in gene therapy have brought novel treatment options for

cancer. However, the full potential of this approach has yet to be unlocked due to

the safety concerns and limited payload capacity of commonly utilized viral vectors.

Virus-free DNA transposons, including *piggyBac*, have potential to obviate these

shortcomings. In this study, we improved a previously developed modified

piggyBac system with superior transposition efficiency. We demonstrated that the

internal domain sequences (IDS) within the 3' terminal repeat domain of

hyperactive *piggyBac* (*hyPB*) donor vector contain dominant enhancer elements.

Plasmid-free donor vector devoid of IDS was used in conjunction with a helper

plasmid expressing *Quantum PBase™* v2 to generate an optimal *piggyBac*

system, *Quantum pBac™* (*qPB*), for use in T cells. Cells transfected with *qPB*

expressing CD20/CD19 CAR outperformed those transfected with the same donor

vector and plasmid expressing *hyPB* transposase in terms of CAR-T cell

production. Importantly, *qPB* yielded mainly $CD8^+$ CAR-T_{SCM} cells, and the *qPB*-

induced CAR-T cells effectively eliminated CD20/CD19-expressing tumor cells

both *in vitro* and *in vivo*. Our findings confirm *qPB* as a promising virus-free vector

system with a payload capacity to incorporate multiple genes. This system is highly

efficient and potentially safe for mediating transgene integration.

Introduction

Gene therapy often requires stable, long-term expression of therapeutic transgene(s) in cells. Such prolonged expression can be accomplished by engineering cells either *ex vivo* or *in vivo* using viral or non-viral vector systems. Viral vectors are most commonly used for gene therapy due to their high efficiencies of gene delivery and integration, which confer stable and long-term gene expression. However, viral vectors have several intrinsic limitations, including: (1) limited payload capacity that severely restricts the repertoire of genes that can be integrated[1]; (2) genotoxicity arising from preferential integration into sites near or within active gene loci that may negatively impact the expression and/or function(s) of genes[2–7]; (3) a cellular proclivity for silencing genes introduced by viral vectors, presumably due to cellular immunity[8,9]; and (4) safety concerns related to immunogenicity of viral vectors[10]. Additionally, production of viral vectors for clinical trials is costly, time-consuming (> 6 months), and supply-constrained, presenting significant obstacles to the routine use of viral vectors in medical practice[11].

In recent years, many advances have been made in non-viral gene delivery technologies, such as virus-free DNA transposons. These advancements may be capable of bypassing the shortcomings of viral vectors, and transposon technology has emerged as an especially promising vector system for gene therapy[12,13]. The great promise of transposon technology is largely due to its effective gene integration capability[14]. DNA transposons, also known as mobile elements or jumping genes, are genetic elements with the ability to transverse the genome

through a "cut-and-paste" mechanism. In nature, a simple DNA transposon

contains a transposase gene flanked by terminal repeat sequences. During the

transposition process, the ability of transposase to act on virtually any DNA

sequence flanked by the terminal repeat sequences makes DNA transposons

particularly attractive as vectors for gene therapy. To turn DNA transposons into

tools for genetic engineering, researchers have developed controllable bi-

component vector systems, consisting of (1) a helper plasmid expressing the

transposase and (2) a donor plasmid with exogenous DNA of interest flanked by

the transposon terminal repeat sequences. Currently, *Sleeping Beauty* and

piggyBac are thought to be the most promising DNA transposons for human gene

therapy and have been clinically explored as vectors for several CAR-T cell

therapies. Unlike *Sleeping Beauty*, which was reconstructed from the salmon

genome[15,16], *piggyBac* was derived from the cabbage looper moth *Trichoplusia ni*

and is naturally active in humans[17–19]. By introducing amino acid mutations to the

transposase, a hyperactive *piggyBac* (*hyPB*) transposase, (*hyPBase*) and two

hyperactive transposases of *Sleeping Beauty* (SB100X and hyperactive SB100X,

which is 30% more active than SB100X) were developed[20–24]. When an exogenous

gene is delivered to primary human T cells *ex vivo*, *hyPBase* can increase

transposition efficiency by 2- to 3-fold compared with *piggyBac* or SB100X

transposases. Previously, we shortened the *piggyBac* terminal repeat domain

(TRD) sequences and observed a 2.6-fold increase in transposition activity

mediated by *piggyBac* in HEK293 cells[25]. We also demonstrated that the activity

of *hyPBase* can be increased by another 2- to 3-fold by fusing the transposase

with various peptides[26]. In this study, we address whether the *piggyBac* system

can be further developed for therapeutic application, including increasing its cargo

capacity relative to those of currently available *piggyBac* systems. To do so, we

generated *Quantum pBac™* (*qPB*), a binary *piggyBac* system comprising a

plasmid-free donor vector and a helper plasmid expressing the molecular-

engineered *hyPBase*, *Quantum PBase™* (*qPBase*) *v2*. This simple, robust and

potentially safe vector system can be used to generate potent CD19/CD20 dual-

targeting CAR-T cells for treatment of B cell malignances.

Results

Micro-*piggyBac* possesses significantly lower enhancer activity compared

to mini-*piggyBac*

In the initial clinical trial of retrovirus-based gene therapy for SCID-X1, there

was evidence of malignancies caused by vector-mediated insertional activation of

proto-oncogenes. The currently available minimal *piggyBac* transposon vector,

designated as mini-*piggyBac* here, contains 5' (244 bp) and 3' (313 bp) TRD *cis*

elements, which are transposed along with gene of interest into the genome. Each

TRD contains a Terminal Inverted Repeat (TIR; also known as inverted minimal

terminal repeats) sequence and an internal domain sequence (IDS)[27]. To minimize

the potential risk of insertional mutagenesis caused by gene delivery vectors in

gene therapy, we had previously generated micro-*piggyBac*. This vector contains

the 5' (67 bp) and 3' (40 bp) TIR sequences of mini-*piggyBac*, but the respective

IDS are absent (Figure 1)[25]. Nevertheless, it remained unclear whether the TRDs

of mini-*piggyBac* and/or TIRs of micro-*piggyBac* harbor enhancer and/or silencer

activities. To address this issue, a panel of luciferase reporter constructs

containing individual TRD or TIR sequences was generated and examined in

insect Sf9 cells, human HEK293 cells, and human Jurkat T cells (Figure 1).

Figure 1. Enhancer/silencer activity of mini-*piggyBac* TRDs and micro-*piggyBac* TIRs

(A) Schematic depicts the panel of luciferase reporter constructs containing a CMV minimal

promoter (MP) and firefly luciferase (*Fluc*) gene, with or without the 5' or 3' TRDs from mini-

piggyBac or 5' or 3' TIRs from micro-*piggyBac* inserted upstream of MP. Luciferase activities

exhibited by the reporter constructs of (A) in (B) Sf9, (C) HEK293, and (D) Jurkat cells. Results are

shown as mean fold changes in luciferase activity ± standard deviation (SD; normalized first to

Renilla luciferase activity and then to the luciferase activity obtained from cells transfected with

construct a). Statistical analysis comparing luciferase activities obtained from cells transfected with

construct **a** and those from cells transfected with constructs **b-e** are summarized in the lower panels

(boxed regions). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. N = 3 (triplicate).

Compared to the control (Figure 1A, construct **a**), the 3' TRD of mini-

piggyBac (construct **c**) but not 3' TIR of micro-*piggyBac* (construct **e**) produced

significantly higher luciferase activity across all three cell types (Figure 1B-1D),

suggesting that enhancer activity is present in the 3' IDS of mini-*piggyBac*. On the

other hand, a slightly yet significantly enhanced luciferase activity was detected in

5' TIR of micro-*piggyBac* but not 5' TRD of mini-*piggyBac* in both HEK293 and

Jurkat cells. This result suggests the presence of a minimal level of enhanced

activity in the 5' TIR and/or silencer activity in the 5' IDS of TRD (Figure 1B-1D).

Taken together, the data suggest that micro-*piggyBac* possesses significantly

lower enhancer activity compared to mini-*piggyBac*.

Shortening of donor vector backbone in combination with *Quantum PBase*™

(*qPBase*) *v2* enhances the transposition activity of micro-*piggyBac* in T cells

The significantly reduced enhancer activity of TIRs suggested that micro-

piggyBac may be safer than mini-*piggyBac* for gene therapy applications. We

therefore focused on micro-*piggyBac* and determined whether its transposition

activity may be further enhanced by shortening the donor vector backbone (i.e.,

the sequences outside the TIR-spanning region). We constructed two donor

vectors named micro-*piggyBac*-Short and micro-*piggyBac*-Long. Both of these

vectors contain TIRs (micro) and either retains (Long) or is devoid of (Short)

replication components in its backbone, namely the replication origin and antibiotic

genes. A third donor vector, mini-*piggyBac* Long, which contains TRDs (mini) and

retains the replication components in its backbone (Long), was also constructed

for comparison; its combination with helper plasmid expressing *hyPBase*

(collectively called "hyperactive *piggyBac*" or "*hyPB*") is currently the most

advanced *piggyBac* system available (Figure 2A). We co-electroporated HEK293,

Jurkat or human primary T cells with different combinations of these donor vectors

and helper plasmids expressing wild-type *PBase* (pCMV-Myc-PBase), *hyPBase*

(pCMV-HA-hyPBase), and *qPBase v1* or *v2* (pCMV-qPBase_v1 and pCMV-

qPBase_v2, respectively). We then determined and compared the transposition

efficiencies of these donor vector and helper plasmid combinations (Figure 2A).

Figure 2. Transposition activities of various combinations of donor vectors and helper

plasmids in human cells

(A)   Schematic depicts the *piggyBac* system with various donor vectors and helper plasmids.

Transposition activity of the indicated combination of donor vector and helper plasmid in (B)

HEK293, (C) Jurkat, and (D) primary T cells. (E) Transposition activity of the indicated combination

of linearized donor vector and helper plasmid in HEK293 cells. Results are shown as mean ± SD

(B-E). n.s. not statistically significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. N = 3 (triplicate).

As shown, *hyPBase*, either in combination with mini-*piggyBac*-Long or

micro-*piggyBac*-Short or -Long, markedly enhanced transposition compared to

PBase in T cells but not in HEK293 cells. On the other hand, *qPBase* v1 markedly

enhanced transposition compared to *hyPBase* in HEK293 cells but not in T cells

(Figure 2B-2D). These results suggest that the transposition activity of *hyPBase*

and *qPBase v1* are likely cell-type-dependent. We also found that *qPBase v2*

mediated the highest transposition activity in nearly all of the tested combinations

and cell types. Importantly, when *qPBase v2* is accompanied by micro-*piggyBac*-

Short donor vector, its transposition activity is by far the highest in both Jurkat and

primary T cells, and this combination is clearly superior to the hyperactive

piggyBac system, *hyPBase* in combination with mini-*piggyBac*-Long (Figures 2C

and 2D).

Micro-*piggyBac* is superior to mini-*piggyBac* for advancing adeno-*piggyBac*

hybrid vector for gene therapy

Even though *piggyBac* is capable of integrating sizable DNA (> 100 kb), its

genome engineering efficiency is restricted by the effectiveness of gene delivery

methods. Electroporation is a particularly effective virus-free gene delivery method

commonly used in gene therapy. However, its transfection efficiency is inversely

correlated with the size of DNA delivered, as cell damage may be caused by

introduction of larger transgenes. Thus, to alleviate electroporation-associated

restrictions imposed on the *piggyBac* system, an adenovirus-*piggyBac* hybrid

system (Ad-iPB7) was developed[28]. Since the adenovirus genome exists in a

linear form, we examined the transposition activity of linearized forms of mini-

piggyBac-Long, micro-*piggyBac*-Short and micro-*piggyBac*-Long along with either

qPBase v1 or *qPBase v2*. As shown in Figure 2E, both micro-*piggyBac*-Short and

micro-*piggyBac*-Long were superior to mini-*piggyBac*-Long. Moreover, when

linearized micro-*piggyBac*-Long donor vector was combined with *qPBase v2*, a

significantly greater transposition efficiency was observed relative to that when

qPBase v1 transposase was used (Figure 2E).

Minicircle micro-*piggyBac* is significantly more active than its parental

counterpart in both Jurkat and primary T cells

A minicircle DNA is a plasmid-free nucleic acid sequence that is normally

generated by an *in vivo* recombination process, which excises prokaryotic plasmid

components such as the antibiotic resistance gene and bacterial origin of

replication. Minicircle DNA constructs offer several advantages, including

enhancements in gene delivery efficiency and stable transgene expression. These

advantages may be attributed to the small vector size of minicircle DNA, the

absence of antibiotic resistance genes, and the lack of gene silencing induced by

bacterial backbone sequences required for plasmid replication. We demonstrated

that removal of backbone sequences outside the TIR-spanning regions via a

cloning procedure (which shortens the vector length) enhanced transposition

efficiency (Figure 2). Therefore, we next used the Mark Kay minicircle system to

generate a minicircle form of the donor vector (Q-tdTomato-IRES-hygro) from its

parental plasmid counterpart (pQ-tdTomato-IRES-hygro) (Figure 3A)[29]. We carried

out the same transposition assay as described for Figure 2 and determined

whether transposition of donor vector in the minicircle form into the T cell genome

would be more efficient than that of its parental plasmid counterpart.

Figure 3. Transposition activity of various transposases with donor vector in parental

plasmid or minicircle counterpart forms

(A) Schematic depicts the generation of the *Quantum pBac™* (*qPB*) donor vector. A gene cassette

carries *tdTomato* and *hygromycin* genes linked by IRES (Q-tdTomato-IRES-hygro) from the

parental plasmid (pQ-tdTomato-IRES-hygro). Transposition activities of the indicated combinations

of pQ-tdTomato-IRES-hygro or Q-tdTomato-IRES-hygro donor vectors and helper plasmids in (B)

Jurkat and (C) primary T cells. (D) Schematic depicts the two-component *qPB* system. Results are

shown as mean ± SD. n.s. not statistically significant, ** p < 0.01, *** p < 0.001. N = 3 (triplicate).

When donor vectors of parental plasmid (pQ-tdTomato-IRES-hygro) and

minicircle (Q-tdTomato-IRES-hygro) forms were compared, it was clear that

minicircle donor vector mediated higher transposition efficiencies than the parental

plasmid form, irrespective of which helper plasmid was co-electroporated.

Moreover, cells co-electroporated with minicircle donor vector and helper plasmid

expressing *qPBase v2* consistently exhibited significantly higher transposition

activity than those receiving all other combinations, including those with the helper

plasmid expressing *hyPBase*. Based on these data in both Jurkat (Figure 3B) and

primary T cells (Figure 3C), we selected micro-*piggyBac* vector in the minicircle

DNA form as the donor vector and a series of recombinant *qPBase* (including

qPBase v1 and *v2*) as the helper plasmids to form the *Quantum pBac™* (*qPB*)

system (Figure 3D).

When combined with CD20/CD19 CAR *Quantum pBac™* (*qPB*) donor vector,

***Quantum PBase™* (*qPBase v2*) outperforms *hyPBase* in CAR-T production**

We next evaluated the performance of CAR-T cells produced by

electroporating activated human primary T cells with vectors of the *qPB* system.

To do so, we used anti-CD20/CD19 CAR as the transgene cassette of *qPB* donor

vector and helper plasmids that encode either *hyPBase* or *qPBase v2*. (Figure 4A).

Figure 4. CAR-T cell production using *Quantum pBac™* (*qPB*) donor vector and helper

plasmid

(A) Schematic depicts the CD20/CD19 CAR *qPB* donor vector and helper plasmid pCMV-hyPBase

or pCMV-qPBase_v2. Characterization of (B) cell viabilities one day after electroporation (Day 1).

Percentages of CAR$^+$ cells on (C) day 1 and (D) day 10 following electroporation of activated

primary T cells. (E) The percentages of CD4 and CD8 CAR$^+$ T$_{SCM}$ cells on day 10. (F) Fold

expansion of cells after 10 days of culture, (G) cytotoxicity of day 10 CAR-T cells on Raji target

cells at the indicated E:T ratios, and (H) CAR copy number after 10 days of culture. Results are

shown as mean ± SD (B-H). n.s., p=0.3 not statistically significant, * p < 0.05, ** p < 0.01, *** p <

0.001. N = 3 (triplicate) except in (H) where testing of each sample was repeated three times per

sample.

One day after electroporation, no significant differences in viability or %

CAR$^+$ were found among cells electroporated with CD20/CD19 CAR *qPB* donor

vector and control helper plasmid (pCDNA3.1, Mock control) and those with helper

plasmid expressing either *hyPBase* or *qPBase v2* (Figure 4B, 4C). Notably, after

10 days of culture, significantly more CAR$^+$ cells were observed in cultures

electroporated with CD20/CD19 CAR *qPB* donor vector and helper plasmid

expressing *qPBase v2* compared to *hyPBase* (Figure 4D). Moreover, the

expansion and cytotoxic effects of cells electroporated with CD20/CD19 CAR *qPB*

donor vector and helper plasmid expressing *qPBase v2* were also significantly

greater than those seen in the other two groups of cells (Figure 4F and 4G). The

CAR copy numbers between *hyPBase*- and *qPBase v2*-transposed cells were not

significantly different (Figure 4H), even when CAR copy number is calculated to

reflect only CAR$^+$ cells (data not shown). These observations suggest that cells

electroporated with helper plasmid expressing *qPBase v2* were superior with

regards to their transposition efficiency, expansion capacity and anti-tumor

potency than cells electroporated with *hyPBase*-expressing plasmids.

Quantum PBase™ produces CAR$^+$ T cells that are more highly enriched with

T$_{SCM}$ subset than those produced by *hyPBase*

Since T$_{SCM}$ enrichment is recognized as an important potential indicator of

clinical efficacy, we also compared the percentage of T$_{SCM}$ in cells electroporated

with CD20/CD19 CAR *qPB* donor vector and helper plasmid expressing either

hyPBase or *qPBase v2* (Figure 4E). We found that in both CD4 and CD8

populations, the percentages of T$_{SCM}$ cells were higher in cells electroporated with

helper plasmid expressing *qPBase v2* than *hyPBase*. Next, we focused on the

qPB system (*qPB* donor vector with *qPBase v2* helper plasmid) and sought to

determine whether there may be donor-dependent variations in CAR$^+$ cell

production using this system. We analyzed the performance of CAR-T cells

derived from peripheral blood mononuclear cells (PBMCs) of six healthy donors

(Figure 5).

Figure 5. Characterization of healthy donor CAR-T cells produced using the *Quantum*

pBac™ (qPB) system

Percentages of (A) CAR$^+$ and (B) transposase$^+$ (*qPBase$^+$*) cells on days 1 and 14 following

electroporation. (C) Fold expansion of cells and (D) distributions of CD4$^+$ and CD8$^+$ T cell subtypes

in CAR$^+$ T cells after 14 days of culture. Distributions of five T cell differentiation subsets T$_N$, T$_{SCM}$,

T$_{CM}$, T$_{EM}$ and T$_{EFF}$ among (E) CD4$^+$ CAR$^+$ and (F) CD8$^+$ CAR$^+$ T cells on day 14. Results were

derived from cells taken from six healthy donors. Horizontal lines in (A-F) represent mean ± SD. *

p < 0.05, *** p < 0.001. N = 6 PMBC donors.

As shown in Figure 5A, the average percentage of CAR$^+$ T cells significantly

increased from day 1 to 14 following electroporation, with only one of six donors

(donor 6) exhibiting a decrease in percentage of CAR$^+$ T cells. Moreover, fourteen

days after electroporation, the percentage of *qPBase*$^+$ cells decreased to minimal

levels (< 0.4%) in all of the donor groups (Figure 5B), suggesting successful

clearance of unwanted helper plasmids from T cells following completion of "cut-

and-paste" gene integration. There was high variability among the PBMC donor

groups in terms of the extent of CAR$^+$ T cell expansion during the 14-day culture

period (27- to 248-fold; Figure 5C), indicating a donor-dependent effect on

expansion capacity of the cells. We also profiled the T cell subtypes (CD4$^+$ and

CD8$^+$; Figure 5D) and T cell subsets based on differentiation stages (Figure 5E,

5F) among the CAR$^+$ T cells derived from the PBMC donors. CD8$^+$ T cells

comprised the major T cell subtype population (Figure 5D), despite the known

higher prevalence of CD4$^+$ T cells in human PBMCs. To analyze this

phenomenon, we carried out a time-course experiment to determine CD8/CD4

ratios before and up to 14 days following electroporation (Supplemental Figure 1).

We categorized the time course into three stages: transient transfection (up to 5

days post-electroporation), stable transgene integration (after 5-7 days post-

electroporation with undetectable transient CAR expression, Figure 4D), and cell

expansion (10 and 14 days post-electroporation). As shown, the CD8 enrichment

appeared to occur at the transfection (Supplemental Figure 1A and 1C) and the

cell expansion stages (after seven days in culture), but not at the stable transgene

integration stage (Supplemental Figure 1B, 1C and 1D). Furthermore, the major

CAR$^+$ T cell subset was T$_{SCM}$ for both CD4 and CD8 populations (Figure 5E, 5F).

Quantum pBac™ (_qPB_) system produces functional CAR$^+$ T cells that kill

target cells _in vitro_

We next sought to determine whether CAR-T cells generated using the _qPB_

system (_qPB_ donor vector with _qPBase v2_ helper plasmid) are functional _in vitro_.

We conducted _in vitro_ cytotoxicity assays by co-incubating antigen-expressing

target cells with either non-transfected control Pan-T cells or CD20/CD19 dual-

targeting CAR-T cells (Figure 6).

D

Figure 6. *In vitro* functional characterization of CAR-T cells produced using the *Quantum*

pBac™ (*qPB*) system

In vitro cytotoxicity of Pan-T and/or CAR-T cells derived from healthy donor(s) (A) against the

indicated target cells, and (B) against Raji cells at the indicated Effector: Target (E: T). (C)

Representative flow cytometry data show the distributions of T cell differentiation subsets T_N/T_{SCM},

T_{CM}, T_{EM} and T_{EFF} in $CD4^+$ and $CD8^+$ subtypes for donor 1- and donor 2-derived CAR^+ T cells. (D)

IFN-g secretion by donor 1 and donor 2 CAR-T cells following antigen stimulation. Pan-T cells (non-

gene modified cells) served as a control. Data shown in (B and D) represent mean ± SD. N = 3

(triplicate). Bar in (A) represents 500 mm.

As shown in Figure 6A, compared to pan-T cells, CD20/CD19 dual-targeting

CAR-T cells eradicated more $CD19^+CD20^+$ Raji cells (third row) and K562 target

cells engineered to express either CD19 (first row) or CD20 (second row). On the

other hand, CD20/CD19 dual-targeting CAR-T cells failed to eradicate BCMA

(irrelevant antigen)-expressing control K562 cells (fourth row). These results

demonstrated that the CAR-T cells specifically target and kill both CD20- and

CD19-expressing cells. The cytotoxic functions of CAR-T cells from two donors

(donors 1 and 2) were further assessed at different E:T ratios against Raji cells.

We observed dose- and time-dependent killing by CAR-T cells derived from both

donors (Figure 6B). Compared with donor 1-derived CAR-T cells, donor 2 CAR-T

cells were much more potent in killing Raji cells, which was consistent with the

relatively higher level of IFN-γ detected in the 48-hour culture medium of donor 2

CAR-T cells (Figure 6D). Notably, 74% of Raji cells were killed by donor 2 CAR-T

cells after a 96-h co-culture period even at a E:T ratio of 1:10 (Figure 6B). This

finding is in contrast to the 15.5% killing of Raji cells by donor 1 CAR-T cells at the

same E:T ratio. Together, these results suggest a higher level of persistence for

CAR-T cells derived from donor 2 as compared with those from donor 1.

Supporting this notion, the percentages of CAR$^+$ T$_{SCM}$ cells were higher among

CAR-T cells from donor 2 (74.4% and 81.7% for CD4$^+$ and CD8$^+$ cells,

respectively) than those from donor 1 (52.1% and 48.3% for CD4$^+$ and CD8$^+$ cells,

respectively; Figure 6C).

Effective tumor clearance by *Quantum pBac™* (*qPB*)-generated CAR-T cells

in Raji-bearing immunodeficient mice

Next, we tested the *in vivo* anti-tumor potency of donor 1 and donor 2 CAR-

T cells in Raji-bearing immunodeficient mice (Figure 7A). Similar to the *in vitro*

cytotoxicity results, injections of low, medium and high doses of donor 1 CAR-T

cells for five days killed Raji tumor cells in a dose-dependent fashion, and the

tumors were completely eradicated by Days 5 and 9 in mice injected respectively

with high and medium doses of donor 1 CAR-T cells (Figure 7B).

A

Figure 7. *In vivo* functional characterization of CAR-T cells produced using the *Quantum*

pBac™ (*qPB*) system in Raji-bearing immunodeficient mice

(A) Schematic depicts the design of the *in vivo* functional characterization experiment. (B)

Bioluminescence imaging (IVIS) revealed the extent of tumor cell persistence 1 day prior to, and 5

and 9 days after CAR-T or Pan-T cell injection. N = 3 or 4 mice/group.

Consistent with these findings, a medium dose of donor 2 CAR-T cells also

eradicated Raji tumor cells from the Raji-bearing mice. Notably, the Raji tumor cell

killing by donor 2 CAR-T cells appeared to have occurred at an earlier time point

(at Day 5 following CAR-T cell injection) than that induced by donor 1 CAR-T cells

(at Day 9 following CAR-T cell injection) (Figure 7B), in agreement with the high *in*

vitro cytotoxicity induced by donor 2 CAR-T cells.

Discussion

Sleeping Beauty and *piggyBac* are two DNA transposons that have been

explored in recent clinical studies on gene and cell therapy[14,30]. In addition to the

multiple advantages over viral vectors mentioned earlier, the *piggyBac* transposon

system has at least four other major benefits. (1) It has a large cargo capacity (>

100 kb)[31]. (2) There is a low frequency of footprint-induced mutations caused by

integrant remobilization[32–34]. (3) *piggyBac* is perhaps the most active transposon

system in human cells[35]. (4) It is the most flexible transposon system in terms of

retained activity upon molecular engineering of the transposase; this aspect may

be crucial for applications that require modifications of the transposase to direct

site-specific *piggyBac* genomic integration. These unique features also make

piggyBac superior to *Sleeping Beauty* as a gene therapy vector. Nevertheless,

Sleeping Beauty is thought to be less genotoxic than *piggyBac* for two main

reasons: First, *piggyBac*-like terminal repeat elements are prevalent in the human

genome[36]. Second, unlike the far more random genome integration profile of

Sleeping Beauty, the genome integration profile of *piggyBac* is associated with

euchromatin and largely excluded from heterochromatin[37,38]. The relatively high

likelihood of *piggyBac* integration proximate to or within genes raises safety

concerns, since enhancer elements of retroviral vector long terminal repeats (LTR)

have previously been implicated as triggers of insertional mutagenesis in SCID-X1

patients[39].

The first above mentioned safety concern of *piggyBac* system was

thoroughly evaluated in a previous study,[40] which demonstrated that the

expression of transposase did not result in mobilization of endogenous *piggyBac-*

like sequences from the human genome[40]. The authors also did not find a

significant increase in double-stranded DNA breaks or a selective growth

advantage in *piggyBac*-harboring cells within long-term cultures of primary human

cells that were modified with eGFP-transposons[40]. To address the second above

mentioned concern (related to the potential risk of insertional mutagenesis

resulting from integrant enhancer activity), we determined whether enhancer

activity was present in the TRDs and/or the TIRs. We discovered that the IDS

region of 3' TRD contained significant enhancer activity (Figure 1). By removing

the IDS, our micro-*piggyBac* is expected to exhibit significantly lower levels of

enhancer activity, which in turn may improve the safety profile of the donor vector.

However, it has been well documented that both TIRs and other sequences

contained within the TRDs are crucial for efficient integration of *piggyBac*

transposon into the host genome. Several attempts to decrease the potential

genotoxicity by reducing the size of the required TRDs to approximately 100 base

pairs of TIRs have resulted in significant losses in transposition efficiency[27,41–43].

In contrast to these findings, our previous study[25] and the present study both

demonstrated that by deleting IDS from both ends and leaving only TIRs (107 bp

total in size), the transposition efficiency of *piggyBac* was in fact increased in

several instances (Figure 2 and Figure 3).[25]

Importantly, a significant increase in transposition efficiency was observed

in all cell types in which cells were co-transfected with a helper plasmid expressing

qPBase v2 when compared to those transfected with plasmid expressing wild type

PBase. This finding suggests that *qPBase v2* is a robust and cell type-independent

piggyBac transposase that can exhibit high transposition activity irrespective of the

type and configuration of the *piggyBac* donor vector. We also demonstrated that

qPBase v2 mediated significantly higher transposition activities than *hyPBase*

(1.8-fold increase in percentage of CAR$^+$ cells on day 10 over *hyPBase*; Figure

4D), while their CAR gene copy numbers did not significantly differ. Our previous

study has shown in a HEK293 genome-wide analysis of *qPBase v2*'s integration

sites that *qPBase v2* displayed a much more random integration profile with no

detectable hot spots, and has a lower preference for CpG islands and cancer-

related genes than *qPBase v1* or wild-type *PBase*.[26] Therefore, *qPBase v2*'s

enhanced transposition activities over *hyPBase* may be due to an increased

preference of *qPBase v2* for regions of the chromosome which promotes

transgene expression. With regards to the potential reason why cells produced

using *qPBase v2* expanded more than those produced using *hyPBase*, both the

above mentioned significantly higher transposition activities of *qPBase v2* as well

as the higher percentages of both CD4 and CD8 CAR$^+$ T$_{SCM}$ (Figure 4D) likely play

a role. It is noteworthy that while the CAR copy numbers in our comparison

experiments seemed relatively high, we have determined in a separate study

related to manufacturing CAR-T cells for clinical applications that the total DNA

electroporated is positively-associated with CAR copy number in CAR-T cells. We

have subsequently optimized the donor:helper ratios to maximize generation of

CAR⁺ T cells while maintaining acceptable CAR copy numbers.[44] Collectively, the

observations of this study suggest that micro-*piggyBac* in conjunction with *qPBase*

v2 comprises a superior *piggyBac* system that outperforms other *piggyBac*

systems comprising *hyPBase*.

In addition to the impacts of TRDs/TIRs and transposase on the

performance of the *piggyBac* system, the method of introducing these components

into host cells can also greatly influence the transposition efficiency, potential

safety profile, and transgene stability of the *piggyBac* system. To ensure co-

transfection of both the transgene delivery cassette and the transposase gene, a

single plasmid transposon system was previously created wherein TRDs were

placed within the delivery cassette, and the transposase gene was located in the

helper part of the same plasmid[27]. However, such an arrangement had a few

disadvantages, including: (1) a reduced rate of gene delivery due to an increase in

size of DNA, a phenomenon particularly seen in electroporation-based gene

delivery, (2) a high degree of plasmid backbone DNA integration when using

transposon plasmids[27], and (3) a high rate of transposase gene integration into the

host genome[45]. For these reasons, we opted to utilize a two-component system,

consisting of donor vector and helper plasmid.

467   The use of a donor vector in plasmid form may have several undesirable

468 qualities for clinical translation, including the presence of bacterial genetic

469 elements and antibiotic-resistance genes. These features are undesirable since

470 they pose a significant safety risk if the plasmid backbone becomes transposed

471 into the host's genome. Additionally, unmethylated CpG motifs that are highly

472 enriched in the bacterial backbone of plasmids have been shown to trigger strong

473 inflammatory responses in host cells through toll-like receptor-9 associated

474 mechanisms. The sequences may also induce transgene silencing, presumably

475 via cellular immune responses[46] and/or interferon secretion[47].

476   To reduce the chance that such events may occur, a new *piggyBac* system

477 was established by incorporating *piggyBac* transposon into a doggybone™ DNA

478 (dbDNA) vector, which is a linear, covalently closed, minimal DNA vector produced

479 enzymatically *in vitro*[48]. A recent report has demonstrated that this *piggyBac*-

480 incorporated dbDNA can be used to generate stable CD19-targeting CAR-T cells

481 with an efficiency similar to that of its plasmid counterpart; the efficiencies were

482 comparable when a minimum of approximately 470 bp of additional randomly

483 selected DNA flanking the transposon was included.   However, due to its linear

484 configuration, dbDNA is expected to be less efficient for electroporation-based

485 gene delivery as compared to its supercoiled circular counterpart. In this study,

486 we set out to increase *piggyBac*'s cargo capacity by using Mark Kay minicircle

487 technology to generate the smallest known *piggyBac* transposon donor vector with

488 TIRs (107 bp) and only 88 bp of the backbone sequence flanking the TIRs. The

489 small backbone sequence in combination with the truncated TRDs (i.e., TIRs) also

resulted in a marked decrease (more than 2 kb) in the size of the donor vector.

The smaller size donor vector in turn reduced DNA toxicity (i.e., electroporation-

induced cell damage) and enhanced the transfection rate, which respectively

improved cell expansion and transposition efficiency.   This increase in

transposition efficiency was further amplified when the donor vector carried a

larger transgene (4.0 kb vs 5.34 kb). The end result of our modifications is that we

now have a *piggyBac* system that is capable of successfully delivering and

transposing larger size transgenes compared to *hyPB*. We demonstrated that use

of this transposon donor vector in conjunction with a highly active *qPBase v2*,

collectively designated as *Quantum pBac™* (*qPB*), represents a superior *piggyBac*

system that is minimalistic, highly efficient, and potentially safe (Figure 3).

Currently, the failure rate for producing sufficient number of CAR$^+$ T cells

for therapy is quite high, and this high rate may be traced back to the scarcity of T

cells in some patients. Moreover, the T cells that remain in these patients are often

exhausted and senesced, making the cells more difficult to expand to reach clinical

quantity and quality criteria. In this study, we have addressed this obstacle with

our *qPB* system utilizing *qPBase v2*. This system can be utilized to effectively

generate CD19/CD20 dual-targeting CAR-T cells with a 1.8-fold increase in the

percentage of CAR$^+$ cells on day 10, and a 2-fold increase in the 10-day expansion

capacity, compared to *hyPBase* (Figure 4). Together, these increases in CAR-T

cell performance result in a 3.6-fold increase in the final yield of CAR-T cells. We

showed that this difference in yield is likely due to an enhanced transposition

efficiency achievable by the *qPB* system (Figure 3), and the increased yield should

facilitate the rapid generation of CAR$^+$ cells at a clinical scale. Likewise, the

significantly higher cytolytic activities of CAR-T cells produced using *qPBase v2*

compared with those produced using *hyPBase* (Figure 4G) likely reflect the

significantly higher percentage of CAR$^+$ cells in the cells produced by *qPBase v2*

than by *hyPBase* (Figure 4D) and greater proportion of T$_{SCM}$ CAR$^+$ T cells (Figure

4E). This advance in technology may prevent the need to extend CAR-T cell

production time; such extensions may be required when using other CAR-T

production systems and may increase the risks of T cell exhaustion and

senescence that compromise CAR-T quality. The higher proportion of CAR$^+$ T cells

that are produced by using the *qPB* system should also allow for

injection/administration of fewer total cells, which may reduce the adverse effects

of CAR-T therapy. We have also shown that as the transgene size increased, the

percentage of CAR$^+$ cells decreased.[49] Since the difference in transposition

efficiencies between plasmid backbone-containing and plasmid backbone-lacking

donor vectors became more pronounced as transgene size increased (4.0 kb in

Figure 2D vs 5.34 kb in Figure 3C), we should be able to improve therapy results

by reducing the size of the donor vector to increase its capacity for therapeutic

transgene payload. Therefore, use of the *qPB* system, which is capable of carrying

a larger transgene cargo due to its reduced size, should enable the production of

efficacious CAR-T cells harboring sizable transgene(s) at a clinical scale.

It is noteworthy that there was a high representation of CD8$^+$ T cells in CAR$^+$

T cells produced using the *qPB* system (Figure 5D), despite the known high

prevalence of CD4$^+$ T cells in human PBMCs. As shown in Supplemental Figure

S1A, the CD8/CD4 ratio initially increased within one day post-electroporation,

suggesting that CD4 T cells may be more sensitive to electroporation-associated

damages. Subsequently, the further increase in CD8/CD4 ratio was positively

associated with a marked increase in cell expansion (see Figure S1C and S1B,

respectively), suggesting that the CD8 T cells were preferentially expanded

compared to the CD4 T cells. The fact that there was no change in CD8/CD4 ratio

at 5 and 7 days post-electroporation (when only transposed and not transfected

transgenes could be detected) also suggests that the pronounced enrichment of

CD8 T cells was not due to preferential transposition of CD8 T cells.   As

demonstrated in another study, this preferential enrichment of CD8 T cells can

serve to complement the more difficult-to-enrich patient CD8 T cells and contribute

to a balanced CD8/CD4 ratio (approximately 1.0) in CAR-T cells produced from

patients with B-cell malignancies.[44]

Additionally, CAR-T cell expansion and persistence have emerged as key

efficacy determinants in cancer patients, and both are positively correlated with the

proportion of T_{SCM} cells in the final CAR-T cell product[50,51]. In line with these

findings, we have demonstrated that our *qPB*-derived CAR-T cells, which are

enriched in T_{SCM} cells (especially among CD8$^+$ CAR$^+$ T cells), exhibited potent anti-

tumor effects both *in vitro* and *in vivo*.

Another strategy to improve CAR-T therapy is to minimize genotoxicity caused

by transposase-induced integrant remobilization. To evaluate this potential

approach, transposase mRNA has been examined in transposon-based CAR-T

clinical trials[52]. However, compared to transposase DNA, the mRNA is generally

less efficient, more costly (for GMP production), and less stable under storage

conditions. Given the low frequency of footprint-induced mutation by *piggyBac*

(<5%) and the fact that only a minimal proportion (<0.4%) of CAR$^+$ T cells

expressed detectable levels of *qPBase* (Figure 5B), transposase in a plasmid form

should be sufficiently safe when applied to highly proliferative *ex vivo*-engineered

cells. Additionally, our recent genome-wide integration profiling of *qPB* in CAR-T

products derived from two distinct donors further support the notion that it is safe

to use in T cell engineering[49].

In summary, given its larger payload capacity compared to currently available

piggyBac systems and high level of efficiency, *qPB* is likely the most suitable

system for development of next generation virus-free gene and cell therapies,

especially for multiplex CAR-T therapy. As we have demonstrated in this study,

the *qPB* system exhibits greater transposition activity, causes less damage to cells,

induces a higher percentage of T$_{SCM}$ and CAR$^+$ populations, and expands more of

the engineered cells than previous systems. These attributes lead to an overall

enhanced ability of the *qPB* system to produce highly CAR$^+$ T$_{SCM}$ cells harboring

larger payloads, which should contribute to the success of *piggyBac*-based

multiplex CAR-T gene therapy. The success of *qPB* in CAR-T cell therapy

suggests that the technology may also be suitable for use in other gene therapy

applications as well.

Materials and Methods

Human T cell samples from healthy donors

Blood samples from adult healthy donors were obtained from Chang Gung

Memorial Hospital (Linkou, Taiwan). The acquisition of these samples was

approved by the Institution Review Board at Chang Gung Medical Foundation (IRB

No. 201900578A3).

Vector constructs

Plasmids were constructed following protocols described previously[53].

Minicircle DNAs were purchased from Aldevron (Fargo, ND). All of the PCR

products or junctions of constructs (wherever sequences were ligated) were

confirmed by sequencing.

Vector construction

pGL3-miniP (Construct a; Figure 1A)

pGL3-basic was digested with SacI and HindIII. The CMV mini-promoter

with SacI and HindIII sequences on its 5' and 3' ends, respectively, was

synthesized, double-digested with SacI and HindIII, and ligated to the SacI-HindIII

vector fragment of pGL3-basic to make construct **a** as shown in Figure 1A.

Constructs b-e (Figure 1A)

pGL3-miniP was digested with KpnI and XhoI. The 5'TIR or 3'TIR of micro-

piggyBac or 5'TRD or 3'TRD of mini-*piggyBac* with KpnI and XhoI sequences

added on either end was synthesized, double-digested with KpnI and XhoI, and

ligated to the KpnI-XhoI vector fragment of pGL3-miniP, respectively, to make the

set of constructs (**b-e**) for evaluation of enhancer activity of mini-*piggyBac* and

micro-*piggyBac* TRDs and TIRs, respectively.

Donor vectors (Figure 2A)

"Mini-*piggyBac*-Long" was the same construct as the donor of *piggyBac* system published previously [19]. "Micro-*piggyBac*-Short" was the plasmid named "*pPB*-cassette short" in a previous publication[25]. "Micro-*piggyBac*-Long" was constructed by replacing the backbone of "micro-*piggyBac*-Short" with that of "mini-*piggyBac*-Long" via PCR-based cloning.

Helper plasmid (Figure 2A)

The construction of helper plasmid was described previously[26].

Parental *qPB* donor vector (Figure 3A)

The parental plasmid contained the following components (obtained by DNA synthesis) arranged in a 5' to 3' order: Kanamycin resistance gene, origin of replication (pMB1), 32 I-SceI sites, attB site (PhiC31), 5'TIR of micro-*piggyBac*, multiple cloning site (MSC), 3'TIR of micro-*piggyBac*, and attP site (PhiC31).

pQ-tdTomato-IRES-hygro (Figure 3A)

The DNA for a cassette with the CAG promoter driving a bi-cistronic transcript containing tdTomato gene, internal ribosome entry site (IRES) of Foot-and-mouth disease virus (FMDV), and hygromycin resistance gene was synthesized and cloned into the AseI and EcoRV sites of the parental *qPB* vector to make pQ-tdTomato-IRES-hygro.

pQ-CD20CD19CAR (Figure 4A)

The dual-targeting tandem CD20/CD19 CAR-containing pQ-CD20CD19CAR vector encoded a second-generation CAR composed of an elongation factor 1-alpha (EF1α) promoter that drives an extracellular domain

derived from the single chain variable fragments (scFv) of monoclonal antibodies

directed against the CD19 and CD20 antigens, respectively. The sequence is

further linked to the CD3z chain of the TCR complex by means of a CD8 hinge and

transmembrane domains, together with the 4-1BB co-stimulatory domain.

Following synthesis, the expression cassette was cloned into the MCS of parental

qPB vector to make pQ-CD20CD19CAR.

Minicircle DNA

The minicircle DNAs, Q-tdTomato-IRES-hygro (Figure 3A) and Q-

CD20CD19CAR (Figure 4A), were manufactured by Aldevron from parental

plasmids, pQ-tdTomato-IRES-hygro and pQ-CD2019CAR, respectively.

qPBase

To comply with the FDA regulations, the ampicillin gene in the helper

plasmid expressing *qPBase v2* was replaced by a kanamycin resistance gene

sequence to make *qPBase*. The *qPBase* component was combined with *qPB*

donor vector (minicircle DNA) to form the *qPB* system.

Enhancer assay

The control, pPL-TK (Renilla Luciferase), was co-transfected with the

specified firefly luciferase constructs (i.e., pGL3-miniP, pGL3-miniP-microL, pGL3-

miniP-microR, pGL3-miniP-miniR, pGL3-miniP-miniL, mini-*piggyBac* Long, micro-

piggyBac Short, or micro-*piggyBac* Long) by either FuGENE (HEK293) or

nucleofection (Sf9 cells, and Jurkat T cells). In some experiments, constructs were

first linearized using XmnI or BglII. Forty-eight hours after transfection, cells were

harvested and subjected to the Dual-Luciferase assay (Promega) according to the

manufacturer's instructions.

Transposition assay

HEK293 cells (1 x 10^5) were transfected with 200-334 ng of donor vector

carrying a hygromycin resistance gene and 200-282 ng of helper plasmid in MEM

medium (GeneDireX), 10% FBS (Corning) utilizing the X-tremeGENE™ HP DNA

transfection reagent (Merck). Transfected cells were transferred to 100-mm plates

and cultured under hygromycin (100 μg/ml) selection pressure for 14 days. Cells

were then harvested and fixed with 4% paraformaldehyde. Fixed cells were stained

with 0.2% methylene blue and cell colonies were counted.

Jurkat or primary T cells (2 x 10^5) were electroporated with 400-668 ng of

donor vector carrying a hygromycin resistance gene, which in some experiments

was also linked to a tdTomato gene, and 400-517 ng of helper plasmid in OpTmizer

medium   supplemented   with   *Quantum   Booster*™   (GenomeFrontier).

Electroporation was carried out using a 4D-Nucleofector™ (Lonza) in combination

with the *Quantum Nufect*™ Kit (GenomeFrontier) according to the manufacturer's

instructions. Electroporated cells were transferred to 96-well plates and cultured

under hygromycin (1 mg/ml) selection pressure for 14 days. Cells were then

harvested and stained with AO/PI. Live cell numbers were determined using Celigo

image cytometry (Nexcelom). Transposition efficiency was expressed as number

of either hygromycin-resistant colonies or live cells.

Generation and expansion of CAR-T cells

Peripheral blood mononuclear cells (PBMCs) were isolated from blood samples of healthy donors by Ficoll-Hypaque gradient separation. CD3$^+$ T cells were isolated from PBMCs using EasySep™ Human T Cell Isolation Kit (StemCell Technologies) according to the manufacturer's instructions. T cells were activated by co-incubation with anti-CD3/anti-CD28 antibody conjugated Dynabeads™ (Gibco, Catalogue No. 11141D) in X-VIVO 15 medium (Lonza) for two days at a bead to cells ratio of 3:1. Following the removal of Dynabeads™, activated T cells were harvested and frozen or utilized in experiments. Electroporation of activated T cells was carried out using a Nucleofector™ 2b Device (Lonza) in combination with the *Quantum Nufect*™ Kit (GenomeFrontier) according to the manufacturer's instructions. Cells were electroporated with the combination of CD20/CD19 CAR donor vector, and pCMV-hyPBase or pCMV-qPBase v2 helper vector. Cells were cultured and expanded for 10 or 14 days in OpTmizer medium (Thermo Fisher Scientific) supplemented with 50 IU of IL-2 (PeproTech) and 10% FBS. Thereafter, the cells were harvested for experiments. γ-irradiated aAPCs were added on Day 3 to the T cell expansion cultures at an aAPC:T cell ratio of 1:1.

Evaluation of CAR-T cells performance

CAR expression on T cells was detected by flow cytometry following staining of cells at 4°C for 30 min with F(ab')$_2$ fragment-specific, biotin-conjugated goat anti-mouse antibodies (Jackson ImmunoResearch Laboratories) and R-phycoerythrin (PE)-conjugated streptavidin (Jackson ImmunoResearch Laboratories). Similarly, cells were stained with the following antigen-specific

fluorophore-conjugated antibodies: CD3-Pacific Blue, CD4-Alexa Flour 532

(Thermo Fisher Scientific), CD8-PE-Cy7, CD45RA-BV421, CD62L-PE-Cy5, or

CD95-BV711 (Biolegend). In some experiments, cells were incubated with

propidium iodide (PI, Thermo Fisher Scientific) and/or Acridine orange (AO,

Nexcelom). PI$^-$ cells and T cell differentiation subsets were determined by flow

cytometry based on CD45RA, CD62L and CD95 expression, as follows: T_N

(CD45RA$^+$CD62L$^+$CD95$^-$), T_{SCM} (CD45RA$^+$CD62L$^+$CD95$^+$), T_{CM} (CD45RA$^-$

CD62L$^+$), T_{EM} (CD45RA$^-$CD62L$^-$), and T_{EFF} (CD45RA$^+$CD62L$^-$). *qPBase* expresses

a GFP-transposase fusion protein, which was detected by flow cytometry for GFP$^+$

cells. In some experiments, a hygromycin resistance gene was expressed with a

tdTomato gene and hygromycin-resistant cells were identified by flow cytometry

for tdTomato$^+$ cells. Flow cytometric measurements and analyses were performed

on a SA3800 Spectral Analyzer (Sony). Histograms and dot-plots were generated

using GraphPad Prism software (GraphPad). Live cells were detected using a

Celigo imaging cytometer (Nexcelom) as AO$^+$, PI$^-$ cells.

***In vitro* cytotoxicity assay**

Target-antigen-expressing cells were engineered according to a method

described elsewhere[49]. CD19$^+$ (K562 CD19-tdTomato), CD20$^+$ (K562 CD20-GFP),

CD19$^+$CD20$^+$ (Raji-GFP) or non-relevant K562 BCMA-GFP target cells were

seeded at 5 x 10^3 cells per well in 96-well culture plates (Corning), and control non-

transfected Pan-T or CAR-T cells were added at an E:T ratio of 5:1, 1:1, 1:3 or

1:10. Cytotoxicity of CAR-T cells on target cells was then assessed as previously

described[54]. Briefly, a Celigo imaging cytometer was used to measure the

percentage of live target cells at 0, 24, 48, 72 and 96 h of co-culture. Cell

aggregates were separated by pipetting prior to Celigo imaging. The percent of

specific lysis for each sample was calculated using the formula: [1-(live fluorescent

cell count in the presence of target cells and CAR-T cells / live fluorescent cell

count in the presence of target cells only)] x 100.

In vitro cytokine release assay

Pan-T control cells or CAR-T cells produced from two healthy donors were

thawed and added to cultures containing $CD19^+$ (K562 CD19-tdTomato), $CD20^+$

(K562 CD20-GFP), $CD19^+CD20^+$ (Raji-GFP) or non-relevant K562 BCMA-GFP

tumor target cells. The cells were added at an effector:target (E:T) ratio of 10:1 in

OpTmizer medium supplemented with 50 IU of IL-2 and 10% FBS. After 48 h of

co-culture, supernatant was collected and IFN-γ level in the culture supernatant

was measured by an enzyme-linked immunosorbent assay (Thermo Fisher),

according to the manufacturer's instructions.

Mouse xenograft model

In vivo studies on a mouse xenograft model were conducted at the

Development Center for Biotechnology, Taiwan, using animal protocols and

procedures ethically reviewed and approved by the Taiwan Mouse Clinic IACUC

(2020-R501-035). Briefly, 8-week-old female ASID (NOD.Cg-

Prkdc^{scid}Il2rg^{tm1Wjl}/YckNarl) mice (National Laboratory Animal Center, Taiwan)

were intravenously (i.v.) injected with 1.5×10^5 Raji-Luc/GFP tumor cells. One

week after Raji-Luc/GFP tumor cell injection, mice were injected with Low (7.5 x

10^5), Med (3×10^6), or High (1.0×10^7) doses of CAR-T cells or control Pan-T cells.

Luminescence signals from Raji-Luc/GFP tumor cells were monitored using the

Xenogen-IVIS Imaging System (Caliper Life Sciences).

Statistical analysis

Statistical analyses of differences between two groups were carried out

using Student's t-test (two-tail), and differences among three or more groups were

analyzed by one-way ANOVA with Tukey's multiple comparison test. The analyses

were performed using GraphPad Prism software (GraphPad Software), and

statistical significance was reported as * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

Differences were considered to be statistically significant when $p < 0.05$.

Acknowledgments

The authors thank Ms. Lu-Chun Chen for her assistance throughout the IRB

preparation and approval process. The authors also thank Dr. Pei-Yi Tsai for

assistance with the animal experiments. This study was funded by

GenomeFrontier Therapeutics, Inc.

Author Contributions:

S.C.-Y.W. designed research. Y.-C.C. (Yi-Chun Chen), P.-N.W., Y.-S.Y.,

Y.-C.C. (Ying-Chun Chen), I.-C.C. performed research. W.-K.H., Y.-C.C. (Yi-Chun

Chen), J.C.H., K.-L.K.W., and S.C.-Y.W. analyzed data. J.C.H., P.S.C. and S.C.-

Y.W. wrote the paper.

Data Availability Statement

The authors confirm that the data supporting the findings of this study are available within the article and/or its supplementary materials.

Keywords

gene therapy | transposon | *piggyBac* | CAR-T

References

1. Ghosh, S., Brown, A.M., Jenkins, C., and Campbell, K. (2020). Viral Vector Systems for Gene Therapy: A Comprehensive Literature Review of Progress and Biosafety Challenges. Applied Biosafety *25*, 7–18. 10.1177/1535676019899502.

2. Nowrouzi, A., Glimm, H., von Kalle, C., and Schmidt, M. (2011). Retroviral vectors: Post entry events and genomic alterations. Viruses *3*, 429–455. 10.3390/v3050429.

3. Schröder, A.R.W., Shinn, P., Chen, H., Berry, C., Ecker, J.R., and Bushman, F. (2002). HIV-1 integration in the human genome favors active genes and local hotspots. Cell *110*, 521–529. 10.1016/S0092-8674(02)00864-4.

4. Cesana, D., Sgualdino, J., Rudilosso, L., Merella, S., Naldini, L., and Montini, E. (2012). Whole transcriptome characterization of aberrant splicing events induced by lentiviral vector integrations. Journal of Clinical Investigation *122*, 1667–1676. 10.1172/JCI62189.

5. Modlich, U., Navarro, S., Zychlinski, D., Maetzig, T., Knoess, S., Brugman, M.H., Schambach, A., Charrier, S., Galy, A., Thrasher, A.J., et al. (2009). Insertional transformation of hematopoietic cells by self-inactivating lentiviral and gammaretroviral vectors. Molecular Therapy *17*, 1919–1928. 10.1038/mt.2009.179.

6. Hacein-Bey-Abina, S., Garrigue, A., Wang, G.P., Soulier, J., Lim, A., Morillon, E., Clappier, E., Caccavelli, L., Delabesse, E., Beldjord, K., et al. (2008). Insertional oncogenesis in 4 patients after retrovirus-mediated gene therapy of SCID-X1. Journal of Clinical Investigation *118*, 3132–3142. 10.1172/JCI35700.

7. Cesana, D., Ranzani, M., Volpin, M., Bartholomae, C., Duros, C., Artus, A., Merella, S., Benedicenti, F., Sergi Sergi, L., Sanvito, F., et al. (2014). Uncovering and dissecting the genotoxicity of self-inactivating lentiviral vectors in vivo. Molecular Therapy *22*, 774–785. 10.1038/mt.2014.3.

8. Jähner, D., and Jaenisch, R. (1985). Retrovirus-induced de novo methylation of flanking host sequences correlates with gene inactivity. Nature *315*, 594–597. 10.1038/315594a0.

9. Jähner, D., Stuhlmann, H., Stewart, C.L., Harbers, K., Löhler, J., Simon, I., and Jaenisch, R. (1982). De novo methylation and expression of retroviral genomes during mouse embryogenesis. Nature *298*, 623–628. 10.1038/298623a0.

10. Raper, S.E., Chirmule, N., Lee, F.S., Wivel, N.A., Bagg, A., Gao, G.P., Wilson, J.M., and Batshaw, M.L. (2003). Fatal systemic inflammatory response syndrome in a ornithine transcarbamylase deficient patient following adenoviral gene transfer. Mol Genet Metab *80*, 148–158. 10.1016/j.ymgme.2003.08.016.

11. van der Loo, J.C.M., and Wright, J.F. (2016). Progress and challenges in viral vector manufacturing. Hum Mol Genet *25*, R42–R52. 10.1093/hmg/ddv451.

812   12.   Ramamoorth, M., and Narvekar, A. (2015). Non viral vectors in gene therapy - An
813   overview. Journal of Clinical and Diagnostic Research *9*, GE01–GE06.
814   10.7860/JCDR/2015/10443.5394.

815   13.   Meir, Y.J.J., and Wu, S.C.Y. (2011). Transposon-based vector systems for gene therapy
816   clinical Trials: Challenges and considerations. Chang Gung Med J *34*, 565–579.

817   14.   Tipanee, J., Chai, Y.C., Driessche, T. vanden, and Chuah, M.K. (2017). Preclinical and
818   clinical advances in transposon-based gene therapy. Biosci Rep *37*, BSR20160614.
819   10.1042/BSR20160614.

820   15.   Ivics, Z., Hackett, P.B., Plasterk, R.H., and Izsvák, Z. (1997). Molecular reconstruction of
821   sleeping beauty, a Tc1-like transposon from fish, and its transposition in human cells. Cell
822   *91*, 501–510. 10.1016/S0092-8674(00)80436-5.

823   16.   Ivics, Z., and Izsvák, Z. (2005). A whole lotta jumpin' goin' on: New transposon tools for
824   vertebrate functional genomics. Trends in Genetics *21*, 8–11. 10.1016/j.tig.2004.11.008.

825   17.   Fraser, M.J., Smith, G.E., and Summers, M.D. (1983). Acquisition of Host Cell DNA
826   Sequences by Baculoviruses: Relationship Between Host DNA Insertions and FP Mutants
827   of Autographa californica and Galleria mellonella Nuclear Polyhedrosis Viruses. J Virol *47*,
828   287–300. 10.1128/jvi.47.2.287-300.1983.

829   18.   Ding, S., Wu, X., Li, G., Han, M., Zhuang, Y., and Xu, T. (2005). Efficient transposition of
830   the piggyBac (PB) transposon in mammalian cells and mice. Cell *122*, 473–483.
831   10.1016/j.cell.2005.07.013.

832   19.   Wu, S.C.Y., Meir, Y.J.J., Coates, C.J., Handler, A.M., Pelczar, P., Moisyadi, S., and Kaminski,
833   J.M. (2006). piggyBac is a flexible and highly active transposon as compared to Sleeping
834   Beauty, Tol2, and Mos1 in mammalian cells. Proc Natl Acad Sci U S A *103*, 15008–15013.
835   10.1073/pnas.0606979103.

836   20.   Mátés, L., Chuah, M.K.L., Belay, E., Jerchow, B., Manoj, N., Acosta-Sanchez, A., Grzela,
837   D.P., Schmitt, A., Becker, K., Matrai, J., et al. (2009). Molecular evolution of a novel
838   hyperactive Sleeping Beauty transposase enables robust stable gene transfer in
839   vertebrates. Nat Genet *41*, 753–761. 10.1038/ng.343.

840   21.   Voigt, F., Wiedemann, L., Zuliani, C., Querques, I., Sebe, A., Mátés, L., Izsvák, Z., Ivics, Z.,
841   and Barabas, O. (2016). Sleeping Beauty transposase structure allows rational design of
842   hyperactive variants for genetic engineering. Nat Commun *7*, 11126.
843   10.1038/ncomms11126.

844   22.   Cadiñanos, J., and Bradley, A. (2007). Generation of an inducible and optimized piggyBac
845   transposon system. Nucleic Acids Res *35*, e87. 10.1093/nar/gkm446.

846   23.   Yusa, K., Zhou, L., Li, M.A., Bradley, A., and Craig, N.L. (2011). A hyperactive piggyBac
847   transposase for mammalian applications. Proc Natl Acad Sci U S A *108*, 1531–1536.
848   10.1073/pnas.1008322108.

849   24.   Yusa, K. (2015). piggyBac transposon. In Mobile DNA III (ASM Press), pp. 873–890.
850   10.1128/9781555819217.ch39.

851   25.   Meir, Y.J.J., Weirauch, M.T., Yang, H.S., Chung, P.C., Yu, R.K., and Wu, S.C.Y. (2011).
852   Genome-wide target profiling of piggyBac and Tol2 in HEK 293: Pros and cons for gene
853   discovery and gene therapy. BMC Biotechnol *11*, 28. 10.1186/1472-6750-11-28.

854   26.   Meir, Y.J.J., Lin, A., Huang, M.F., Lin, J.R., Weirauch, M.T., Chou, H.C., Lin, S.J.A., and Wu,
855   S.C.Y. (2013). A versatile, highly efficient, and potentially safer piggyBac transposon
856   system for mammalian genome manipulations. FASEB Journal *27*, 4429–4443.
857   10.1096/fj.12-223586.

858   27.   Solodushko, V., Bitko, V., and Fouty, B. (2014). Minimal piggyBac vectors for chromatin
859   integration. Gene Ther *21*, 1–9. 10.1038/gt.2013.52.

860   28.   Cooney, A.L., Singh, B.K., and Sinn, P.L. (2015). Hybrid nonviral/viral vector systems for
861   improved piggyBac DNA transposon in vivo delivery. Molecular Therapy *23*, 667–674.
862   10.1038/mt.2014.254.

863   29.   Kay, M.A., He, C.Y., and Chen, Z.Y. (2010). A robust system for production of minicircle
864   DNA vectors. Nat Biotechnol *28*, 1287–1289. 10.1038/nbt.1708.

865   30.   Magnani, C.F., Tettamanti, S., Alberti, G., Pisani, I., Biondi, A., Serafini, M., and Gaipa, G.
866   (2020). Transposon-Based CAR T Cells in Acute Leukemias: Where are We Going? Cells *9*,
867   1337. 10.3390/cells9061337.

868   31.   Li, R., Zhuang, Y., Han, M., Xu, T., and Wu, X. (2013). PiggyBac as a high-capacity
869   transgenesis and gene-therapy vector in human cells and mice. DMM Disease Models
870   and Mechanisms *6*, 828–833. 10.1242/dmm.010827.

871   32.   Elick, T.A. (1996). Excision of the piggy Bac transposable element in vitro is a precise
872   event that is enhanced by the expression of its encoded transposase. Genetica *98*, 33–41.
873   10.1007/BF00120216.

874   33.   Li, M.A., Pettitt, S.J., Eckert, S., Ning, Z., Rice, S., Cadiñanos, J., Yusa, K., Conte, N., and
875   Bradley, A. (2013). The piggyBac Transposon Displays Local and Distant Reintegration
876   Preferences and Can Cause Mutations at Noncanonical Integration Sites. Mol Cell Biol *33*,
877   1317–1330. 10.1128/mcb.00670-12.

878   34.   Chen, Q., Luo, W., Veach, R.A., Hickman, A.B., Wilson, M.H., and Dyda, F. (2020).
879   Structural basis of seamless excision and specific targeting by piggyBac transposase. Nat
880   Commun *11*, 3446. 10.1038/s41467-020-17128-1.

881   35.   Doherty, J.E., Huye, L.E., Yusa, K., Zhou, L., Craig, N.L., and Wilson, M.H. (2012).
882   Hyperactive piggybac gene transfer in human cells and in vivo. Hum Gene Ther *23*, 311–
883   320. 10.1089/hum.2011.138.

884   36.   Feschotte, C. (2006). The piggyBac transposon holds promise for human gene therapy.
885   Proc Natl Acad Sci U S A *103*, 14981–14982. 10.1073/pnas.0607282103.

37. Galvan, D.L., Nakazawa, Y., Kaja, A., Kettlun, C., Cooper, L.J.N., Rooney, C.M., and Wilson,
M.H. (2009). Genome-wide mapping of piggybac transposon integrations in primary
human T cells. Journal of Immunotherapy *32*, 837–844. 10.1097/CJI.0b013e3181b2914c.

38. Hamada, M., Nishio, N., Okuno, Y., Suzuki, S., Kawashima, N., Muramatsu, H., Tsubota, S.,
Wilson, M.H., Morita, D., Kataoka, S., et al. (2018). Integration Mapping of piggyBac-
Mediated CD19 Chimeric Antigen Receptor T Cells Analyzed by Novel Tagmentation-
Assisted PCR. EBioMedicine *34*, 18–26. 10.1016/j.ebiom.2018.07.008.

39. Hacein-Bey-Abina, S., von Kalle, C., Schmidt, M., McCormack, M.P., Wulffraat, N.,
Leboulch, P., Lim, A., Osborne, C.S., Pawliuk, R., Morillon, E., et al. (2003). LMO2-
Associated Clonal T Cell Proliferation in Two Patients after Gene Therapy for SCID-X1.
Science (1979) *302*, 415–419. 10.1126/science.1088547.

40. Saha, S., Woodard, L.E., Charron, E.M., Welch, R.C., Rooney, C.M., and Wilson, M.H.
(2015). Evaluating the potential for undesired genomic effects of the piggyBac
transposon system in human cells. Nucleic Acids Res *43*, 1770–1782.
10.1093/nar/gkv017.

41. Li, X., Harrell, R.A., Handler, A.M., Beam, T., Hennessy, K., and Fraser, M.J. (2005).
piggyBac internal sequences are necessary for efficient transformation of target
genomes. Insect Mol Biol *14*, 17–30. 10.1111/j.1365-2583.2004.00525.x.

42. Elick, T.A., Lobo, N., and Fraser, M.J. (1997). Analysis of the cis-acting DNA elements
required for piggyBac transposable element excision. Molecular and General Genetics
255, 605–610. 10.1007/s004380050534.

43. Li, X., Lobo, N., Bauser, C.A., and Fraser, M.J. (2001). The minimum internal and external
sequence requirements for transposition of the eukaryotic transformation vector
piggyBac. Molecular Genetics and Genomics *266*, 190–198. 10.1007/s004380100525.

44. Chang, P.S., Chen, Y.-C., Hua, W.-K., Hsu, J.C., Tsai, J.-C., Huang, Y.-W., Kao, Y.-H., Wu, P.-
H., Chang, Y.-F., Chang, M.-C., et al. (2022). Manufacturing highly potent CD20/CD19-
targeted iCasp9 regulatable CAR-T cells using the Quantum pBac-based CAR-T (qCART)
system for clinical application. bioRxiv. https://doi.org/10.1101/2022.05.03.490475.

45. Urschitz, J., Kawasumi, M., Owens, J., Morozumi, K., Yamashiro, H., Stoytchev, I., Marh, J.,
Dee, J.A., Kawamoto, K., Coates, C.J., et al. (2010). Helper-independent piggyBac plasmids
for gene delivery approaches: Strategies for avoiding potential genotoxic effects. Proc
Natl Acad Sci U S A *107*, 8117–8122. 10.1073/pnas.1003674107.

46. Chen, Z.Y., Yant, S.R., He, C.Y., Meuse, L., Shen, S., and Kay, M.A. (2001). Linear DNAs
concatemerize in vivo and result in sustained transgene expression in mouse liver.
Molecular Therapy *3*, 403–410. 10.1006/mthe.2001.0278.

47. Huerfano, S., Ryabchenko, B., and Forstová, J. (2013). Nucleofection of expression
vectors induces a robust interferon response and inhibition of cell proliferation. DNA Cell
Biol *32*, 467–479. 10.1089/dna.2012.1950.

48.   Bishop, D.C., Caproni, L., Gowrishankar, K., Legiewicz, M., Karbowniczek, K., Tite, J.,
Gottlieb, D.J., and Micklethwaite, K.P. (2020). CAR T Cell Generation by piggyBac
Transposition from Linear Doggybone DNA Vectors Requires Transposon DNA-Flanking
Regions. Mol Ther Methods Clin Dev *17*, 359–368. 10.1016/j.omtm.2019.12.020.

49.   Chen, Y.-C., Hua, W.-K., Hsu, J.C., Chang, P.S., Wen, K.K.-L., Huang, Y.-W., Tsai, J.-C., Kao,
Y.-H., Wu, P.-H., Wang, P.-N., et al. (2022). Quantum CART (qCART), a piggyBac-based
system for development and production of virus-free multiplex CAR-T cell therapy.
bioRxiv. 10.1101/2022.05.03.490469.

50.   Gattinoni, L., Speiser, D.E., Lichterfeld, M., and Bonini, C. (2017). T memory stem cells in
health and disease. Nat Med *23*, 18–27. 10.1038/nm.4241.

51.   Biasco, L., Izotova, N., Rivat, C., Ghorashian, S., Richardson, R., Guvenel, A., Hough, R.,
Wynn, R., Popova, B., Lopes, A., et al. (2021). Clonal expansion of T memory stem cells
determines early anti-leukemic responses and long-term CAR T cell persistence in
patients. Nat Cancer *2*, 629–642. 10.1038/s43018-021-00207-7.

52.   Prommersberger, S., Reiser, M., Beckmann, J., Danhof, S., Amberger, M., Quade-Lyssy, P.,
Einsele, H., Hudecek, M., Bonig, H., and Ivics, Z. (2021). CARAMBA: a first-in-human
clinical trial with SLAMF7 CAR-T cells prepared by virus-free Sleeping Beauty gene
transfer to treat multiple myeloma. Gene Ther *28*, 560–571. 10.1038/s41434-021-00254-
w.

53.   Sambrook, J., and Russell, D.W. (2001). Molecular Cloning: A Laboratory Manual, Third
Edition. In Molecular Cloning: a laboratory a manual, pp. 1.32-1.34.

54.   Sun, Y.-J., Chen, Y.-C., Hua, W.-K., Wu, S.C.-Y., and Chan, L. (2022). Comparison of
chimeric antigen receptor-T cell-mediated cytotoxicity assays with suspension tumor
cells using plate-based image cytometry method. Cytometry A. 10.1002/cyto.a.24673.