1

Aggression and multimodal signaling in noise in a common urban songbird

Çağla Önsal[1], Alper Yelimlieş [1], Çağlar Akçay [1,2]

[1] Department of Psychology, Koç University, İstanbul, Turkey

[2] School of Biological Sciences, Anglia Ruskin University, Cambridge, UK

ORCID

Çağla Önsal: **0000-0003-3131-6756**

Alper Yelimlieş: **0000-0003-2319-6141**

Çağlar Akçay: **0000-0003-0635-9586,**

Corresponding author:

Çağla Önsal

Department of Psychology

Koç University, İstanbul, Turkey

consal20@ku.edu.tr

Abstract

Anthropogenic noise may disrupt signals used to mediate aggressive interactions, leading

to more physical aggression between opponents. One solution to this problem is to switch

signaling effort to a less noisy modality (e.g., the visual modality). In the present study we

investigate aggressive behaviors and signaling in urban and rural male European robins

(*Erithacus rubecula*) in response to simulated intrusions with or without experimental noise.

First, we predicted that urban birds, living in noisier habitats, would be generally more

aggressive than rural birds. We also predicted that during simulated intrusions with experimental

noise, robins would increase their physical aggression and show a multi-modal shift, i.e., respond

with more visual threat displays and sing fewer songs. Finally, we expected the multi-modal shift

in response to noise to be stronger in urban birds compared to rural birds. The results showed

that urban birds were more aggressive than rural robins, but an increase in aggression with

experimental noise was seen only in the rural birds. Urban but not rural birds decreased their

song rate in response to noise. Contrary to the multi-modal shift hypothesis, however, there was

no evidence of a concurrent increase in visual signals. These results point to a complex role of

immediate plasticity and longer-term processes in affecting communication during aggressive

interactions under anthropogenic noise.

Keywords:

Multi-modal signaling, territoriality, anthropogenic noise, multi-modal shift, European robin

Significance Statement

Human activity has an enormous effect on wildlife, including on their social behavior.

Animals living in urban areas often tend to be more aggressive than those living in rural areas,

which may be due to urban acoustic noise making communication between individuals more

difficult. In a study with a common songbird, the European robin, we investigated the role of

urban acoustic noise in aggression and territorial communication. Urban robins were more

aggressive than rural robins, and additional noise in the territory increased aggression in rural but

not urban robins. While urban robins decreased their singing effort with additional noise, they

did not increase visual signals concurrently. These results suggest that noise can indeed make

animals behave more aggressively although the effect may depend on how noisy it is already.

These results further our understanding of how human-made noise changes animal

communication and social behavior.

Introduction

Urban habitats are polluted with anthropogenic noise, often in multiple modalities, which

creates challenges for urban-living wildlife (Brumm and Slabbekoorn 2005). Many species rely

heavily on signals for communication in contexts such as mate attraction and territorial defense,

and noise from vehicles, buildings and other human activities often interferes with these signals

(Francis et al. 2009; Halfwerk and Slabbekoorn 2015; Lee and Thornton 2021). A well-studied

example of the effect of anthropogenic noise on communication is vocal signaling in urban birds:

in response to anthropogenic noise commonly found in cities, many species of birds may

increase repetition rates, amplitude, or frequency characteristics of their acoustic signals (Brumm

2004; Gil and Brumm 2014; Slabbekoorn and den Boer-Visser 2006; Wood and Yezerinac

2006).

Urban living also leads to increased aggressiveness of individuals in urban habitats

compared to the rural habitats (Davies and Sewall 2016; Evans et al. 2010; Hardman and

Dalesman 2018; Phillips and Derryberry 2018; Scales et al 2011). The reasons for increased

aggression in urban habitat are not yet fully understood. It may result from several factors,

including selection for bolder individuals (Evans et al. 2010), increased food resources (Foltz et

al. 2015), increased exposure to harmful chemicals such as lead (McClelland et al. 2019) and less

stable social environment due to high rates of territory turnover in urban habitats (Davis et al.

2013). Anthropogenic noise in urban habitats may also be responsible for increased aggression.

Animals often use signals in aggressive interactions (e.g., during territory defense) to

resolve conflicts with opponents. Use of signals is often beneficial for both parties if they can

avoid costly physical fights in this way (Maynard Smith and Price 1973). Consequently, if

signaling is prevented or the signals are rendered ineffective, individuals may need to resort to

higher levels of physical aggression (Logue et al. 2010). Applied to urban habitats, this

hypothesis suggests that the high levels of urban noise may render long-distance aggressive

signals less effective, which in turn may lead to higher levels of aggression (Phillips and

Derryberry 2018). Consistent with this hypothesis, some studies reported a positive correlation

between ambient noise levels and aggressive behaviors (Akçay et al. 2020; Phillips and

Derryberry 2018; but see Kleist et al. 2016).

Signalers employ various strategies to overcome interference from anthropogenic noise.

We focus here on the flexibility afforded by having signals in more than one modality (Bro-

Jørgensen 2010; Halfwerk and Slabbekoorn 2015; Partan and Marler 1999). When animals have

signals in more than one modality, they may shift their signaling effort from the noisy modality

to a less noisy modality to increase the likelihood that the message of the signals gets through to

the receivers (Partan et al. 2010; Partan 2017). This hypothesis is termed the multi-modal shift

hypothesis.

Few studies tested the multi-modal shift hypothesis in signals used in territorial

interactions. In one study, Ríos-Chelén et al. (2015) found that male red-winged blackbirds

(*Agelaius phoeniceus*) did not use more intense visual signaling in noisier territories although

they modified their acoustic signals (e.g., decreased their song rate). Another study on song

sparrows (*Melospiza melodia*) found that males in noisier urban habitats were both more

aggressive and used proportionally more visual threat signals (wing waves) during territory

defense compared to the males in rural habitats, consistent with a multi-modal shift (Akçay et al.

2020). In a further experiment, however, individual song sparrows did not increase their visual

signaling effort when experimentally presented with noise, suggesting that multi-modal shift

seen in urban song sparrows was not due to immediate plasticity (Akçay and Beecher 2019).

Here we investigate the responses of European robins (*Erithacus rubecula*) living in urban

and rural habitats in Istanbul, Turkey, to simulated territorial intrusions with or without

experimental noise playback. European robins have both visual and acoustic signals that are used

in agonistic interactions (Lack 1943). Previous studies have found that robins respond both to

song and visual signals during territorial intrusions (Chantrey and Workman 1984). Territory

holders sing in response to the song of an intruder, while visual signals are used when the

intruding male is within range of vision. Their most prominent visual signal is the neck display,

which has been observed in response to the sight of a rival male's red neck, or indeed even a ball

of red feathers (Lack 1943). Other visual threat signals include wing flutters, pricking the tail up,

and swaying, where the resident male moves his head from one side to the other (Lack 1943).

Signaling and aggressive behaviors of robins have been investigated in multiple studies. A

study by Mclaughlin and Kunc (2013) found that robins, after being lured by playback of a robin

song from a speaker, tended to move away from the speaker when the speaker switched to

playback of low-frequency noise mimicking typical traffic noise, particularly when the amplitude

of noise was high (90 dB at 1m). In response increasing amplitude of noise, the robins sang

shorter songs with fewer notes and increased the minimum frequency of their songs. In another

study, experimental presentation of wind turbine noise during simulated territorial intrusions led

to a decrease in low frequency elements in the songs, at the same time leading to an increase in

flight rates (Zwart et al. 2016). Song rates did not significantly differ between the noise and no-

noise treatments. Interestingly, fewer robins used visual threat postures under experimental

presentation of noise compared to no noise, although the difference was not significant (Zwart et

al. 2016).

120   The presence of both acoustic and visual signals in territorial defense makes the European

121 robin a suitable candidate for testing the multimodal shift hypothesis but to our knowledge no

122 previous study compared visual signaling between urban and rural robins. We predict that robins

123 in urban habitats will exhibit higher levels of visual signaling. Additionally, if such a multi-

124 modal shift is due to phenotypic plasticity, robins should increase their visual signaling under

125 experimental noise. We also predicted that urban birds would exhibit higher levels of plasticity

126 in signaling than rural birds, because they more frequently experience significant anthropogenic

127 noise levels (Gentry et al. 2017; Lazerte et al. 2016). Finally, in accordance with earlier studies,

128 we also expect to see a greater level of aggression from urban robins compared to rural robins. If

129 increased aggression is due to individual plasticity in response to noise, we also expect higher

130 levels of aggression from robins in response to experimental noise, particularly in urban birds

131 who have more experience with noise.

132   Methods

133   *Study site*s and species

134   We carried out playback experiments with male European robins that held territories in

135 rural areas (forests around Sarıyer, Istanbul, 41° 9' 50.73971"N, 29° 0' 32.25243"E) and urban

136 parks and green areas in Sarıyer, Istanbul, Turkey in April and May 2021 (urban: n=9; rural:

137 n=12). Robin territories were detected by the presence of an already-singing male robin before

138 the first playback or during recording sessions prior to playback. We determined a central

139 location by observing the robin's flights for about 5 minutes, although we did not attempt to map

140 the entire territory. In all trials reported below, only a single bird responded to the playback (for

141 one subject, not included in the final sample, we aborted the trial when a second male came to

142 within 10 m of the speaker). It was not possible to record data blind as our study involved

observing focal individuals in urban or rural habitats and noise manipulation was audible to all

observers.

Stimuli

Playback stimuli were generated on the software Syrinx (John Burt, Portland, OR) from

male European robin songs recorded in March 2021 in four of the nine study sites. We generated

stimulus files by extracting high quality songs from each recording and filtering out low

frequency noise below 1000 Hz. We added a silent period after each song so that stimuli were

presented at a rate of one song per seven seconds. The songs lasted on average (±SD) 2.36

(±0.49). We created one-minute stimuli (consisting of nine different songs) which were repeated

three times to make up three-minute stimuli to be played during the trials. In total, we used 17

files created from the songs of 17 different robins. The stimuli played for each subject came from

a robin whose territory was separated by at least one km from that of the subject's territory.

Subjects received the same song stimulus in both trials. As a visual stimulus, we used a 3-D

printed bird model (dimensions, height: 8 cm, length: 12 cm, width: 4.5 cm) which was hand-

painted to resemble an adult robin (Figure S6).

We generated the experimental acoustic noise stimuli by filtering white noise (created with

Audacity) with the average amplitude spectrum from a 1-minute recording (made with a Marantz

PMD660 and ME66/K6 microphone) of traffic noise in Sarıyer, Istanbul using the package

seewave in R (Sueur et al. 2008). The power spectrum of the noise stimulus can be found in the

supplementary materials (Figure S1).

Experimental procedure and design

We started each trial by placing the robin model attached to a speaker (Anker Soundcore

Bluetooth Speaker, Anker, Inc.) on a natural perch at the estimated center of the resident male's

territory, approximately 1.5 m above the ground. A second Bluetooth speaker (same model as

above) was placed on the ground, face-up below the first, for noise playback. In the control

treatment, the second speaker was placed but not turned on, so the resident male received only

song playback. In the noise treatment, in addition to the song playback, traffic noise was played

at 75 dB SPL at 1 m. The noise playback lasted for the entire duration of the song playback. This

stimulus simulates an acute, if transient, source of noise (e.g., a vehicle idling, or a lawnmower

operating).

Each subject received two 3-minute trials, one with experimental noise and one without

noise. The order of the treatments was counterbalanced, and the two trials were separated by

approximately an hour. Two observers, about 10 m away from the experimental setup, recorded

the songs and calls of the resident robin. The observers also narrated the trials, verbally noting

flights (any airborne movement by the bird), distance with each flight, and visual displays

described in Table 1 onto the recording. We continued recording songs for 3 minutes after the

end of each trial. Recordings were made on a Marantz PMD660 with a Sennheiser ME66/K6

microphone, or on a Zoom H5 handheld recorder with a Zoom SGH6 shotgun microphone.

In 31 of the trials, the bird was already singing when we started the trial. For these birds,

the 3-minute playback period started with the first song played. In 11 of the trials, where the

subject was quiet when the playback started, the 3-minute trial period began with their first

response (song or approach). The average duration of pretrial playback for these 11 trials was

64.9 seconds (SD = 40.8).

After each trial, we measured the ambient noise with a VLIKE VL6708 sound-level meter

with the method described in (Brumm 2004). We took eight measurements (two in each cardinal

direction) within a minute period, which were then averaged. For three subjects, we only had

noise measurements from a single trial.

Response variables

We scanned and annotated the narrated trial recordings using the Syrinx software. The

number of songs and visual displays (number of neck displays and wing flutters as well as the

start and end times for swaying in seconds) were extracted from the verbal notes made on the

recordings. We only analyzed song rates and durations, as overlapping stimulus and subject

songs made it impossible to determine with certainty the note compositions for most songs,

precluding frequency measurements. Only 8 subjects used any visual displays during the

experimental period, we therefore coded visual displays as a binomial variable (visual signal

present vs. absent during a trial). From the recordings, we also extracted the number of flights,

closest approach to the model/speaker and proportion of time spent within 5m of the speaker.

These three spatial variables were taken as aggressive behaviors.

Because the spatial variables of aggression were significantly correlated with each other

(all $p < 0.05$), we carried out a principal component analysis (PCA) using the *principal* function

in package *psych* (Revelle 2021). The first component of PCA (PCA1) explained 61% of

variance and was taken as our primary measure of aggression (see Table 2 for loading

coefficients). The aggression scores thus calculated have been shown to be valid measures of

territorial aggression in songbirds (Akçay et al. 2013). We report the analyses using raw spatial

measures in the supplementary materials.

Data analysis

All analyses were carried out in R (R Core Team 2021). We first checked whether urban

territories had higher ambient noise with a linear mixed model (LMM) using habitat type (urban

vs. rural) as the predictor variable and territory ID as the random variable. We also assessed

whether noise levels were repeatable using the *rptR* package (Stoffel et al. 2017).

We then checked whether the order of trials had a significant effect on aggression scores,

song rates and visual signaling. The order of trials did not have a significant effect on song rate

(LMM, coefficient = -0.68, SE = 0.60, p = 0.28) or aggression score (LMM, coefficient = -0.22,

SE = 0.15, p = 0.16). However, there was a significant order effect on the incidence of visual

displays (GLMM, estimate = -15.32, SE = 6.22, p = 0.01). Eight subjects used visual displays in

the first trial, compared to two in the second trial (both subjects also used visual displays in the

first trial). Because of this order effect, we only used the first trial for each subject in models

including visual displays.

We analyzed song rates and aggression scores with linear mixed models (LMM), using the

lme function in the package *nlme* (Pinheiro et al. 2022). We took habitat type (urban vs. rural)

and experimental treatment (noise vs. control), and their interaction as the predictor variables,

and male ID as the random variable. We applied a generalized linear model (with log-link, using

the "*glm*" function in base R) with visual displays as binomial response variable, and habitat and

treatment as predictor variables, using only the first trials for each subject. Since only two rural

subjects used visual displays, we also carried out this analysis with the subset of only urban birds

(see Supplementary Materials).

Results

Urban habitats had significantly higher levels of ambient noise than rural habitats (urban:

M = 49.0, SD = 7.1; rural: M = 39.9, SD = 3.6) and noise measurements were highly repeatable

between the two trials (intra-class correlation coefficient; r=0.96, standard error: 0.02; p <

0.0001).

Urban birds were significantly more aggressive than rural birds. There was no main effect

of noise treatment but there was a significant interaction effect of habitat and noise treatment

(Table 3, Figure 1a). To understand this interaction effect, we carried out paired t-tests on rural

and urban birds with noise treatment as predictor variable. Rural birds were more aggressive

under the noise compared to no noise treatment (paired t-test; t(11)=2.44, p= 0.033), whereas

there was no effect of experimental noise on aggression in urban birds (paired t-test: t(8)=-1.23,

p= 0.25; Figure 1a). Looking at the spatial variables separately, this interaction effect seems to be

driven mostly by the closest approach measure (see Supplementary Materials).

Song rates did not differ significantly between urban and rural birds and the noise

treatment had no main effect. The interaction effect of habitat and noise treatment however

approached significance (Table 3, Figure 1b). When we analyzed song rates for urban and rural

birds separately there was a significant effect of noise treatment in urban birds with lower song

rates under experimental noise compared to without noise (paired t-test; t(9)=3.15, p= 0.014);

while there was no effect of noise treatment for rural birds (paired t-test; t(11)=1.01, p= 0.33;

Figure 1b). There was no significant main or interaction effect of habitat or noise treatment on

song length (Table 3).

Urban birds used visual threat displays in the first trials significantly more than their rural

counterparts (GLM; χ^2= 9.75, p=0.0018; Figure 2). While there was no main effect of noise

treatment (χ^2= 0.00, p=1.0), the interaction effect of treatment and habitat approached

significance (χ^2= 3.72, p=0.053). This interaction effect was driven by a tendency in urban birds

to use more visual signals in the no noise treatment compared to noise treatment, although the

effect was not significant (GLM within the subset of urban birds; χ^2= 3.22, p=0.07). Out of the

five urban birds that received the no noise treatment in the first trial, all five used visual signals,

compared to only one out of the 4 birds which received the noise treatment first (Figure 2).

Discussion

In the present study, we examined the role of acute anthropogenic noise in determining

aggressiveness and aggressive signaling. We predicted that urban robins, living in noisier

territories, would be more aggressive compared to rural robins in simulated territorial intrusions

and experimental noise during simulated intrusions will change both their aggressive behaviors

and signaling behaviors. Particularly, we expected that experimental acoustic noise should

increase aggression during intrusions and lead to an increase in using signals in the visual

modality, particularly in urban birds who would have more experience dealing with fluctuating

levels of noise.

In line with our first prediction, we found that urban robins responded with significantly

more aggressive behaviors (particularly close approach) to simulated intrusions than rural robins.

The effect of experimental noise treatment on aggressive approach was dependent on the habitat:

contrary to our hypothesis, experimental noise led to increased aggression in the (comparatively

quiet) rural habitats, but it had no effect in the noisy urban habitats. Experimental noise led to a

decrease in song rates in urban and not rural birds. No change was observed in song duration (see

Table 3). Finally, visual signals were more common in urban habitats (consistent with the fact

that urban birds are more aggressive) and tended to be less common under experimental noise in

the urban habitats, albeit not significantly.

Noise and aggressive behaviors in territory defense

Our results on the effect of habitat on aggression replicates earlier findings that urban-

living birds are more aggressive than rural birds (Davies and Sewall 2016; Evans et al. 2010;

Hardman and Dalesman 2018; Phillips and Derryberry 2018). We also extend previous findings

by showing that the effect of experimental noise on aggression was dependent on habitat: urban

males showed no further increases in aggression with experimental noise while rural males

showed a significant increase in aggression. This is opposite of our expectation that urban birds

would show higher levels of phenotypic plasticity in response to experimental noise treatment

(Gentry et al. 2017; LaZerte et al. 2016). The lack of an effect of noise on aggressive behaviors

in urban habitats may be due to several reasons: First, for urban males that are already living in

noisy territories, additional noise may not have as much as an effect as in rural habitats. Urban

birds may also be more habituated to acute increases of noise than rural birds, although they did

show a plastic response in their singing rate as discussed below. Finally, urban birds may not be

able to increase their already high levels of aggression in response to noise playback.

The increased aggression with experimental noise in rural habitats is consistent with the

idea that urban noise has a causal role in increasing aggression. Only a small number of studies

experimentally manipulated noise levels to examine a causal role of noise in increased

aggression. These studies yielded mixed results. Grabarcyzk and Gill (2019) found that house

wrens (*Troglodytes aedon*) males attacked the simulated intruder more frequently when playback

was accompanied with experimental noise than when it wasn't, consistent with the hypothesis

that noise induces higher levels of aggression. Another study in song sparrows however, found

no effect of experimental noise on aggression levels, measured as time spent within one meter of

the speaker, or attacks (Akçay and Beecher 2019). In the latter study, the noise playback started

only when subjects approached to within five meters, which all subjects did within a short period

of time (< 1 minute). Thus, lack of an effect in physical proximity may be due the fact that

subjects already were close to the speaker when the noise playback started.

In another study, Zwart et al. (2016) found that European robins did not show a statistically

significant increase in aggressive behaviors in response to experimental wind turbine noise

during simulated intrusions, although some variables like flights did show a trend consistent with

higher aggression with experimental noise. A more recent study by Reed et al (2021) in lazuli

buntings (*Passerina amoena*) and spotted towhees (*Pipilo maculatus*) found that experimental

presentation of natural noise (such as noise from a river, ocean surf or cicadas) at the landscape

level led to slower detection of a simulated intruder and consequently weaker approach

responses (see also Kleist et al. 2016). Finally, a study conducted with saffron finches (*Sicalis*

flaveola) found birds displayed lower agonistic behaviors under experimental traffic noise,

although this study is harder to interpret in the present context as the experiment was done with

captive birds that lived in small cages (Passos et al. 2020).

Together these studies point to two apparently contradictory effects of noise on territorial

aggression. On one hand, noise may make localization of the simulated intruder and perception

of stimulus features more difficult, leading to slower or weaker approach behaviors (Kleist et al.

2016; Reed et al. 2021; Templeton et al. 2016). On the other hand, assuming the simulated

intruder is located, noise may interfere with the signaling behaviors of subjects which may

induce them to resort to higher physical aggression such as closer approach (e.g., Grabarcyzk

and Gill 2019). Thus, the differences in the findings may be due in part to differences in

experimental designs, particularly with respect to the presentation of the noise stimulus (e.g.,

type of noise, location of noise relative to the conspecific stimulus etc.).

The experimental noise in our study represents a transient increase in noise to a high

amplitude that coincides with the need to confront a territorial intruder. Thus, the effects we see

are responses to acute noises, while urban birds would experience varying noise levels due to

cars passing as well as daily patterns of human activity (Gill et al. 2017). Our study was

explicitly designed to study plastic responses to acute increases in noise and therefore the results

may not apply to chronic but varying amounts of noise. Nevertheless, the situation we simulated

is a realistic one: urban wildlife has to deal with transient increases in noise such as this regularly

(e.g. when a park worker uses a leafblower to clean trails or when a lawnmower works nearby).

Change in multi-modal signals with noise

We found that European robins changed their signaling behaviors in response to noise. In

the acoustic modality, urban but not rural robins decreased their song rates, while birds in neither

habitat changed their song length. From the perspective of the multi-modal shift hypothesis, a

decrease in signals in auditory modality was expected to coincide with an increase in visual

signals. We did not find this second effect. Two caveats are worth mentioning here: First, we

were not able to carry out the experiment blindly with respect to habitat or noise treatment,

which may have biased our observations of visual displays. Clearly, we could not blind observers

regarding which type of habitat they were in, and "blinding" observers with respect to noise

treatment (e.g., by using noise-canceling headphones) would have made keeping track of the

vocal behaviors almost impossible. In any case, given the pattern of results that tend to the

opposite direction of our expectations however, we believe observer bias is unlikely to be an

issue here.

Second, we could not examine individual level plasticity in visual signaling, because of a

significant order effect in visual signals: visual displays were mostly used in the first trials only.

We do not have a good hypothesis as to why we found such an order effect. It is possible that our

stationary 3D model ceased to be a good visual stimulus by the time of the second trial, leading

to a decrease in visual signaling. It is also possible that the second trials may have represented a

lesser threat to the territory owner (given that it simulates the return of a previously retreating

individual) thus eliciting lower threat signals. Lack (1939) noted that repeated presentations of a

taxidermic mount quickly leads the lowered responses which he interpreted as the lack of

realistic response of the immobile taxidermic mount. While something similar may be happening

in our experiment, we note that that there was no order effect in other aggressive or vocal

behaviors.

Whatever the reason, the order effect meant that we could only analyze the visual signals

in the first trials for each subject, halving our sample size and precluding a within-subject

comparison. The between individual comparison of visual signals among urban birds in the first

trials yielded evidence in the opposite direction of what was expected: there was more visual

signaling in trials without experimental noise than with experimental noise, although the

difference was not significant, likely due to the small sample size. Nevertheless, this finding

suggests that while urban birds decrease their acoustic signaling effort, they do not necessarily

depend on visual signals as a back-up as predicted by the multi-modal shift hypothesis.

Our finding is similar to that of a study on European robins by Zwart et al (2016) which

reported slightly lower rates of visual threat signals with experimental noise than without noise.

Another study on song sparrows found no effect of experimental noise during territorial defense

on overall rates of wing waves, a visual threat signal (Akçay and Beecher, 2019), even though

urban song sparrows show higher rates of wing waves than rural birds when controlling for the

total number of acoustic and visual threat signals (Akçay et al. 2020). These studies suggest that

if urban noise causes a multi-modal shift in visual threat signals, it is unlikely to be due to

immediate phenotypic plasticity.

Note that the fact that urban robins used visual displays more frequently than rural robins is

consistent with a multimodal shift due to noise. This finding, however, is also consistent with the

hypothesis that urban birds are simply more aggressive and therefore use visual threat signals

more than rural birds (cf. Akçay et al. 2020). A valid comparison of the use of visual signals

between urban and rural birds would need to somehow correct for aggressiveness. Thus,

currently the evidence for a multi-modal shift in this species is relatively weak.

Behavioral plasticity of urban vs. rural birds in response to noise.

We had expected that urban birds would show a higher level of plasticity in their responses

to experimentally presented noise compared to rural birds. This prediction was based on studies

that showed prior experience with noise (as urban birds would have) leads to more directional

plasticity in acoustic parameters of their song, by e.g. increasing the minimum frequency of their

song in noise (Gentry et al. 2017; LaZerte et al. 2016). Instead, we found contrasting patterns of

response to noise depending on the behavior measured: while in aggression scores, urban birds

showed lower directional plasticity than rural birds (specifically, rural birds increased their

approach distance in noise, while urban birds showed no difference), the opposite was true in

song rates (urban birds decreased song rates in noise while rural birds did not show a difference

between treatments).

These findings suggest that if there is a role of learning in determining responses, it may

take different forms depending on the behavior. It is possible for instance that urban birds in

general have learned to "sit out" transient increases in noise (such as the situation we simulated

here) by reducing song rates and not increasing their approach towards opponents and visual

signaling. Such a strategy may be adaptive, given that closely approaching an opponent and

increasing visual threat signals likely would escalate an aggressive interaction to a fight (Searcy

et al. 2006). In contrast, rural birds may show less plasticity in signing in noise and instead

approach the opponent more closely during noise, because they haven't had the opportunity to

learn how to deal with transient increases in noise. Thus, the contrasting patterns found in urban

and rural birds may still indicate the role of prior experience with noise, although clearly more

controlled experiments are needed to further test this hypothesis.

Conclusion

In summary, our results showed important differences in how urban and rural robins

respond to noise during aggressive interactions. These results suggest that the ambient noise

levels experienced by animals is an important factor in determining their responses to transient

increases in noise.

Disclosure of potential conflicts of interest

The authors declare that they have no conflicts of interest.

Research involving Human Participants and/or Animals

All procedures used in this study follow the ASAB/ABS guidelines for the treatment of

animals in behavioral research and teaching. Subjects were not captured or handled before,

during or after any of the trials. Time spent within a territory did not exceed 15 minutes per trial,

and 30 minutes per day.

Informed consent

The research did not involve human participants.

Data availability statement

The raw data and the R-code to reproduce the analyses reported in the manuscript is

available as a supplementary material and will be deposited at a public repository before the

manuscript is published.

Acknowledgements

This study was funded by the British Ornithological Union Small Ornithological Research

Grant to ÇÖ.

References

Akçay Ç, Beck ML, Sewall KB (2020) Are signals of aggressive intent less honest in urban

habitats? Beh Eco arz179. https://doi.org/10.1093/beheco/arz179

Akçay Ç, Beecher MD (2019) Multi-modal communication: song sparrows increase signal

redundancy in noise. Biol Lett 15:20190513. https://doi.org/10.1098/rsbl.2019.0513

Akçay Ç, Porsuk YK, Avşar A, et al (2020) Song overlapping, noise, and territorial aggression in

great tits. Beh Eco 31:807–814. https://doi.org/10.1093/beheco/araa030

Akçay Ç, Tom ME, Campbell SE, Beecher MD (2013) Song type matching is an honest early

threat signal in a hierarchical animal communication system. Proc R Soc B

280:20122517. https://doi.org/10.1098/rspb.2012.2517

Bro-Jørgensen J (2010) Dynamics of multiple signalling systems: animal communication in a

world in flux. Trends in Ecology and Evolution 25:292–300.

https://doi.org/10.1016/j.tree.2009.11.003

Brumm H (2004) The impact of environmental noise on song amplitude in a territorial bird. J

Anim Ecology 73:434–440. https://doi.org/10.1111/j.0021-8790.2004.00814.x

Brumm H, Slabbekoorn H (2005) Acoustic Communication in Noise. In: Advances in the Study

of Behavior. Elsevier, pp 151–209

Chantrey DF, Workman L (1984) Song and plumage effects on aggressive display by the

European Robin Erithacus rubecula. Ibis 126:366–371. https://doi.org/10.1111/j.1474-

919X.1984.tb00257.x

Davies S, Sewall KB (2016) Agonistic urban birds: elevated territorial aggression of urban song

sparrows is individually consistent within a breeding period. Biol Lett 12:20160315.

https://doi.org/10.1098/rsbl.2016.0315

Davis A, Major RE, Taylor CE (2013) Housing Shortages in Urban Regions: Aggressive

Interactions at Tree Hollows in Forest Remnants. PLoS ONE 8:e59332.

https://doi.org/10.1371/journal.pone.0059332

Evans J, Boudreau K, Hyman J (2010) Behavioural Syndromes in Urban and Rural Populations

of Song Sparrows. Ethology. https://doi.org/10.1111/j.1439-0310.2010.01771.x

Foltz SL, Ross AE, Laing BT, et al (2015) Get off my lawn: increased aggression in urban song

sparrows is related to resource availability. BEHECO 26:1548–1557.

https://doi.org/10.1093/beheco/arv111

Francis CD, Ortega CP, Cruz A (2009) Noise Pollution Changes Avian Communities and

Species Interactions. Current Biology 19:1415–1419.

https://doi.org/10.1016/j.cub.2009.06.052

Gentry KE, Derryberry EP, Danner RM, et al (2017) Immediate signaling flexibility in response

to experimental noise in urban, but not rural, white☐crowned sparrows. Ecosphere 8:.

https://doi.org/10.1002/ecs2.1916

Gil D, Brumm H (2013) Acoustic communication in the urban environment. In: Gil D, Brumm H

(eds) Avian Urban Ecology. Oxford University Press, pp 69–83

Gill SA, Grabarczyk EE, Baker KM, Naghshineh K, Vonhof MJ (2017) Decomposing an urban

soundscape to reveal patterns and drivers of variation in anthropogenic noise. Science of

the Total Environment 599:1191-1201 https://doi.org/10.1016/j.scitotenv.2017.04.229

Grabarczyk EE, Gill SA (2019) Anthropogenic noise affects male house wren response to but

not detection of territorial intruders. PLoS ONE 14:e0220576.

https://doi.org/10.1371/journal.pone.0220576

Halfwerk W, Slabbekoorn H (2015) Pollution going multimodal: the complex impact of the

human-altered sensory environment on animal perception and performance. Biol Lett

11:20141051. https://doi.org/10.1098/rsbl.2014.1051

Hardman SI, Dalesman S (2018) Repeatability and degree of territorial aggression differs among

urban and rural great tits (Parus major). Sci Rep 8:5042. https://doi.org/10.1038/s41598-

018-23463-7

Kleist NJ, Guralnick RP, Cruz A, Francis CD (2016) Anthropogenic noise weakens territorial

response to intruder's songs. Ecosphere 7:. https://doi.org/10.1002/ecs2.1259

LaZerte SE, Slabbekoorn H, Otter KA (2016) Learning to cope: vocal adjustment to urban noise

is correlated with prior experience in black-capped chickadees. Proc R Soc B

283:20161058. https://doi.org/10.1098/rspb.2016.1058

Lack D (1939) The Behaviour of the Robin.—Part I. The Life☐history, with special reference to

Aggressive Behaviour, Sexual Behaviour, and Territory. Part II. A Partial Analysis of

Aggressive and Recognitional Behaviour. Proceedings of the Zoological Society of

London A109:169–219. https://doi.org/10.1111/j.1096-3642.1939.tb03362.x

Lack, D (1943) The life of the Robin, 1st edn. London, Witherby.

Lee VE, Thornton A (2021) Animal Cognition in an Urbanised World. Front Ecol Evol

9:633947. https://doi.org/10.3389/fevo.2021.633947

Logue DM, Abiola IO, Rains D, et al (2010) Does signalling mitigate the cost of agonistic

interactions? A test in a cricket that has lost its song. Proc R Soc B 277:2571–2575.

https://doi.org/10.1098/rspb.2010.0421

Maynard Smith J, Price GR (1973) The Logic of Animal Conflict. Nature 246:15–18.

https://doi.org/10.1038/246015a0

McClelland SC, Durães Ribeiro R, Mielke HW, et al (2019) Sub-lethal exposure to lead is

associated with heightened aggression in an urban songbird. Science of The Total

Environment 654:593–603. https://doi.org/10.1016/j.scitotenv.2018.11.145

McLaughlin KE, Kunc HP (2013) Experimentally increased noise levels change spatial and

singing behaviour. Biol Lett 9:20120771. https://doi.org/10.1098/rsbl.2012.0771

Partan SR, Marler P (1999) Communication Goes Multimodal. Science 283:1272–1273.

https://doi.org/10.1126/science.283.5406.1272

Partan SR (2017) Multimodal shifts in noise: switching channels to communicate through rapid

environmental change. Animal Behaviour 124:325–337.

https://doi.org/10.1016/j.anbehav.2016.08.003

Partan SR, Fulmer AG, Gounard MAM, Redmond JE (2010) Multimodal alarm behavior in

urban and rural gray squirrels studied by means of observation and a mechanical robot.

Current Zoology 56:313–326. https://doi.org/10.1093/czoolo/56.3.313

Passos MFO, Beirão MV, Midamegbe A, et al (2020) Impacts of noise pollution on the agonistic

interactions of the saffron finch (Sicalis flaveola Linnaeus, 1766). Behavioural Processes

180:104222. https://doi.org/10.1016/j.beproc.2020.104222

Phillips JN, Derryberry EP (2018) Urban sparrows respond to a sexually selected trait with

increased aggression in noise. Sci Rep 8:7505. https://doi.org/10.1038/s41598-018-

25834-6

Pinheiro J, Bates D, R Core Team (2022) nlme: Linear and Nonlinear Mixed Effects Models. R

package version 3.1-157, https://CRAN.R-project.org/package=nlme

R Core Team (2021) R: a language and environment for statistical computing. Vienna, Austria:

R Foundation for Statistical Computing http://www.R-project.org/.

Revelle, W (2021) psych: Procedures for Psychological, Psychometric, and Personality Research

(2.1.9). https://CRAN.R-project.org/package=psych

Reed, VA, Toth CA, Wardle RN, Gomes DGE, Barber JR, Francis CD (2021) Natural noise

affects conspecific signal detection and territorial defense behaviors in songbirds.

Behavioral Ecology 32:993–1003

Ríos-Chelén AA, Lee GC, Patricelli GL (2015) Anthropogenic noise is associated with changes

in acoustic but not visual signals in red-winged blackbirds. Behav Ecol Sociobiol

69:1139–1151. https://doi.org/10.1007/s00265-015-1928-7

Scales J, Hyman J, Hughes M (2011) Behavioral Syndromes Break Down in Urban Song

Sparrow Populations: Behavioral Syndromes Break Down in Urban Song Sparrow

Populations. Ethology 117:887–895. https://doi.org/10.1111/j.1439-0310.2011.01943.x

Searcy WA, Anderson RC, Nowicki S (2006) Bird song as a signal of aggressive intent. Behav

Ecol Sociobiol 60:234–241. https://doi.org/10.1007/s00265-006-0161-9

Slabbekoorn H, den Boer-Visser A (2006) Cities Change the Songs of Birds. Current Biology

16:2326–2331. https://doi.org/10.1016/j.cub.2006.10.008

Stoffel MA, Nakagawa S, Schielzeth H (2017) rptR: repeatability estimation and variance

decomposition by generalized linear mixed☐effects models. Methods Ecol Evol 8:1639–

1644. https://doi.org/10.1111/2041-210X.12797

Sueur J, Aubin T, Simonis C (2008) Seewave, a Free Modular Tool for Sound Analysis and

Synthesis. Bioacoustics 18:213–226. https://doi.org/10.1080/09524622.2008.9753600

Templeton CN, Zollinger SA, Brumm H (2016) Traffic noise drowns out great tit alarm calls.

Current Biology 26:R1173–R1174. https://doi.org/10.1016/j.cub.2016.09.058

Wolfenden AD, Slabbekoorn H, Kluk K, Kort SR (2019) Aircraft sound exposure leads to song

frequency decline and elevated aggression in wild chiffchaffs. J Anim Ecol 88:1720–

1731. https://doi.org/10.1111/1365-2656.13059

Wood WE, Yezerinac SM (2006) Song Sparrow (Melospiza Melodia) Song Varies with Urban

Noise. The Auk 123:650–659. https://doi.org/10.1093/auk/123.3.650

Zwart MC, Dunn JC, McGowan PJK, Whittingham MJ (2016) Wind farm noise suppresses

territorial defense behavior in a songbird. Beh Eco 27:101–108.

https://doi.org/10.1093/beheco/arv128

Figure 1. The relationship of aggression scores (A) and song rate (B) with habitat and noise treatment. The boxes indicate interquartile ranges, the middle line indicates median, and whiskers indicate 95% confidence intervals. Dots connected by dotted lines represent data from individual subjects.

Figure 2. Proportion of first trials where the resident male used visual signals, grouped by habitat and noise treatment. The numbers at the bottom of each bar indicates the total number of subjects for each combination.

Table 1. Visual displays of European robins during territorial interactions

Behavior	Description
Neck display	The robin raises his head, displaying his neck.
Wing flutter	The robin flutters his wings.
Swaying	The robin rhythmically sways his body from one side to the other.
Tail up	The robin perks his tail up.

Table 2. Loading coefficients of the Principal Component Analysis.

Factor	Loading coefficient to PC1
Flight rate	0.65
Proportion of time spent within 5 meters	0.83
Closest approach distance	-0.85
SS loadings	1.83
%Variance	61%

Table 3. Coefficients (SE) from the linear mixed models and the p-values from Wald t tests, examining the effect of habitat and experimental noise treatment. Statistically significant values are shown in bold type

	Aggression Score		Song Rate		Song Length	
Predictors	*Estimates (SE)*	*p*	*Estimates (SE)*	*p*	*Estimates (SE)*	*p*
(Intercept)	-0.58 (0.26)	**0.035**	8.44 (0.59)	**<0.001**	1.86(0.14)	**<0.001**
Treatment	1.32 (0.39)	**0.003**	0.61 (0.65)	0.36	0.04(0.21)	0.8454
Habitat	0.32 (0.19)	0.105	0.81 (0.90)	0.38	0.03(0.11)	0.7890
Treatment*Habitat	-0.68 (0.29)	**0.031**	-2.02 (1.00)	0.059	0.11(0.17)	0.5376