Colour polymorphism associated with a gene duplication in male wood tiger moths

Melanie N. Brien[1][*][†], Anna Orteu[2][*], Eugenie C. Yen[2], Juan A. Galarza[1,3], Jimi Kirvesoja[3],

Hannu Pakkanen[4], Kazumasa Wakamatsu[5], Chris D. Jiggins[2][**], Johanna Mappes[1,3][**].

[1]Organismal and Evolutionary Biology Research Program, Faculty of Biological and
Environmental Sciences, University of Helsinki, Finland.

[2]Department of Zoology, University of Cambridge, United Kingdom.

[3]Department of Biological and Environmental Science, University of Jyväskylä, Finland.

[4]Department of Chemistry, University of Jyväskylä, Finland.

[5]Institue for Melanin Chemistry, Fujita Health University, Toyoake, Japan.

*These authors contributed equally.

**Co-senior authors.

[†]Correspondence: mnbrien1@gmail.com

Abstract

Colour is often used as an aposematic warning signal, with predator learning expected to lead

to a single colour pattern within a population. However, there are many puzzling cases where

aposematic signals are also polymorphic. The wood tiger moth, *Arctia plantaginis*, uses

bright hindwing colours as a signal of unpalatability, and males have discrete colour morphs

which vary in frequency geographically. In Finland, both white and yellow morphs can be

found, and these colour morphs also differ in behavioural and life-history traits. Complex

polymorphisms such as these are often explained by supergenes. Here, we show that male

colour is linked to an extra copy of a *yellow* family gene that is only present in the white

morphs. This white-specific duplication, which we name *valkea,* is highly upregulated during

wing development, and could act to reduce recombination, thus potentially representing a

supergene. We also characterise the pigments responsible for yellow, white and black

colouration, showing that yellow is partly produced by pheomelanins, while black is

dopamine-derived eumelanin. The *yellow* family genes have been linked to melanin synthesis

and behavioural traits in other insect species. Our results add to only a few examples of

seemingly paradoxical and complex polymorphisms which are associated with single genes.

Introduction

Colour polymorphisms, defined as the presence of multiple discrete colour phenotypes within a population (Huxley, 1955), provide an ideal trait to study natural and sexual selection. Colour phenotypes are linked to fitness in many contexts including camouflage, mimicry and mating success. Colour is often associated with aposematism, warning predators of unpalatability (Cott, 1940; Cuthill *et al.*, 2017), and in such cases, predator learning should favour the most common colour pattern, leading to positive frequency dependent selection (Endler, 1988). Despite this, aposematic polymorphisms can be stable when selection is context-dependent (Briolat *et al.*, 2019), especially where genetic correlations between colour phenotypes and other traits lead to complex fitness landscapes (reviewed by McKinnon and Pierotti, 2010).

A variety of genetic mechanisms can underpin these types of complex polymorphisms involving multiple associated traits (Orteu and Jiggins, 2020). In many cases, such complex polymorphisms are controlled by 'supergenes' in which divergent alleles at several linked genes are maintained in strong linkage disequilibrium by reduced recombination. The most common mechanism for locally reduced recombination are inversions, which range from single inversions involving a small number of genes, to multiple nested inversions covering large genomic regions (Wang *et al.*, 2013; Küpper *et al.*, 2016; Tuttle *et al.*, 2016). Nonetheless, other mechanisms for reducing recombination, such as centromeres or large genomic deletions, may also play a role. An alternative mechanism is that a single regulatory gene could act in the same way as a supergene, by controlling variation via multiple downstream effects (Thompson and Jiggins, 2014). While there are fewer instances in which multiple phenotypes seem to be controlled by a single gene, one potential example is the common wall lizard, where colour genes have pleiotropic effects on behavioural and reproductive traits (Andrade *et al.*, 2019). Multiple mutations within a single gene can also lead to variation in multiple traits (Linnen *et al.*, 2013).

The wood tiger moth, *Arctia plantaginis,* has a complex polymorphism that has been well studied in an ecological context. Males show polymorphic aposematic hindwing colouration with discrete yellow, white or red hindwing colour morphs found at varying frequencies in different geographic locations. In Finland, for example, both yellow and white morphs can be found, with white morphs varying in frequency from 40 to 75% (Galarza *et al.*, 2014). In Estonia, white morphs make up 97% of the population, while yellows morphs form a

completely monomorphic population in Scotland (Hegna, Galarza and Mappes, 2015) (Figure 1). Long term breeding studies of these moths have shown that male hindwing colour is a Mendelian trait controlled by a single locus with two alleles (Suomalainen, 1938; Nokelainen *et al.*, 2022). White alleles (W) are dominant over the yellow (y). These colour genotypes also covary with behavioural and life-history traits, contributing to the maintenance of this polymorphism. Yellow males are subject to lower levels of predation in the wild (Nokelainen *et al.*, 2012, 2014), while white males have a positive frequency-dependent mating advantage (Gordon *et al.*, 2015). Yellow morphs have stronger chemical defences (Rojas *et al.*, 2017), but show reduced flight activity compared to white males, although yellows may fly at more selective times, i.e. at peak female calling periods (Rojas, Gordon and Mappes, 2015). In summary, there is a trade-off between natural selection through predation and reproductive success, which contributes to the maintenance of this polymorphism (Rönkä *et al.*, 2020).

Despite the large body of research on *A. plantaginis* colour morphs, the genetic basis of this polymorphism is unknown. Here, we explore the genetic basis of male hindwing colour using wild populations and lab crosses to test whether the polymorphism is associated with large structural rearrangements controlling multiple phenotypic elements, or the result of a single gene mutation. We then narrow down the candidate genes using a differential gene expression analysis and identify the pigments producing yellow, white and black colouration on the wings of male *A. plantaginis*.

Figure 1: (A) Sampling locations and frequencies of yellow and white *Arctia plantaginis* in Finland, Scotland and Estonia. (B) Males of the white and yellow colour morphs (Credit: Samuel Waldron).

86 **Results**

A narrow genomic region is associated with hindwing colour

To investigate the genetic basis of male hindwing colouration in *A. plantaginis*, we carried

out a quantitative trait locus (QTL) mapping analysis using crosses between heterozygous

Wy males and homozygous yy females. We used RADseq data aligned to the yellow *A.*

plantaginis reference genome from 172 male offspring (90 white and 82 yellow) from four

families. The QTL analysis identified a single marker associated with male hindwing colour

(Figure 2). This marker was found on scaffold YY_tarseq_206_arrow at position 9,887,968bp

(95% confidence intervals 9,349,978-9,888,009bp) and had a LOD score of 32.8 (p<0.001).

The significant marker explains around 75% of the phenotypic variation and, with one

exception, yellow individuals all had a homozygous yy genotype at this marker. This is

consistent with tight linkage of this marker to the causal SNP.

To further narrow down this region, we ran a genome wide association study (GWAS) using

whole genome sequences of males from four populations: Southern Finland (5 white, 5

yellow) and Central Finland (10 w, 10 y), where male morphs are either white or yellow,

Estonia (4 w), where males are mostly white, and Scotland (12 y), where all males are

yellow. This identified a region of associated SNPs also on scaffold 206 (Figure 2). Two

SNPs, 137bp apart (at positions 9,885,384 and 9,885,521), were significant above a strict

Bonferroni corrected threshold. 162 SNPs were over the threshold of p<0.0001 and, of these,

155 are within a 99kb region on scaffold 206 (9,833,387-9,932,264bp). The top SNPs are

within 2.5kb from the top QTL marker, and the SNP at this marker has a p-value <0.0001.

The 538Kb QTL interval contains 21 genes (Table S1) which were annotated with reference

to *D. melanogaster*. Of these genes, four are part of the *yellow* gene family. The top two

SNPs from the GWAS, and the top marker from the QTL, fall in a non-coding region

upstream of the gene, *yellow-e*, and are also close to an additional *yellow* gene, *yellow-g*.

Identifying structural variation in this region

The trio binning method used by Yen et al. (2020) to assemble the *A. plantaginis* reference

genome produced two reference sequences, one for a white allele and one for a yellow allele.

We extracted the region containing the QTL interval from the yellow reference and aligned it

against the white reference. The alignment showed a duplicated region approximately 117kb

long on scaffold 419 of the white reference from around 6,941,000-7,058,000bp (Figure S1).

The *yellow-e* gene and its flanking regions are within this sequence and are therefore

duplicated in the white reference (Figure 2). One copy of the gene (named jg1310 in the W

annotation) has 7 exons and is similar to the *yellow-e* gene in the yellow reference (99.7%

identity in coding sequences). The second copy unique to the white scaffold (jg1308) has

only the first 5 exons (81.8% identical to the gene in the yellow reference), possibly due to a

stop codon mutation in the 5th exon. For clarity, we named this duplicated white-specific

copy *valkea*, in reference to a Finnish term for 'white'. While all white samples had

consistent coverage of reads across the duplicated region, coverage was patchy in yellow

samples, with many regions having no coverage in yellow samples (Figure 2). Those reads

that map in the *valkea* region in yellow samples are likely to be mapping errors.

To confirm that both of these gene copies are related to *yellow-e*, we compared them to

yellow-e orthologs found in *Bombyx mori*, *Heliconius melpomene* and *Drosophila*

melanogaster, along with other *yellow* genes from *A. plantaginis* and *B. mori*. Both of the

tiger moth genes were most closely related to the *H. melpomene yellow-e* (Figure S2).

Between *valkea* and *yellow-e* there is an additional gene which showed highest similarity to

Drosophila yellow-g2 (when extracted from both the white and yellow references). This gene

is not part of the duplicated sequence and is present as a single copy in both morphs.

Coverage across *yellow-g* and *yellow-e* genome regions in wild samples is similar in both

morphs (Figure 2).

Figure 2: (A) QTL analysis reveals a 500kb region significant on scaffold 206, part of linkage group
9. (B) A GWAS of wild samples showed SNPs associated with hindwing colour along the same
scaffold. The dotted line shows the Bonferroni corrected significance threshold. (C) An insertion in
the white reference sequence contains a copy of the *yellow-e* gene, which we named *valkea*, in
addition to the *yellow-e* present in both white and yellow morphs. (D) Read depth across the candidate
region in all Finnish white and yellow samples.

 Valkea is differentially expressed between morphs

To pinpoint which of these candidate genes is associated with male wing polymorphism in *A.*

plantaginis, we next performed gene expression analyses across several developmental

stages. Based on knowledge of the expression patterns of *yellow* genes and other colour

pattern genes such as *optix* (Reed *et al.*, 2011) and *cortex* (Nadeau *et al.*, 2016) in

Lepidoptera, we hypothesised that changes in gene regulation that control the development of

wing colour morphs in the tiger moth most likely occur during pupal development. Pupal

development in the wood tiger moth lasts for approximately 8 days, and no colour is present

in the wings until day 7, when the yellow pigment appears. A few hours later, black melanin

pigmentation is deposited. We sampled two stages early in development when no colouration

is present in the wings (72 hours post-pupation, and 5 day old pupae), and two stages later in

development: the point when yellow appears in yy morphs (Pre-mel, 7 day old pupae) and the

other after black melanin has also been deposited (Mel, 7-8 days old). Five individuals per

genotype and stage were sampled.

We analysed differential gene expression using the W reference genome, which contains the

duplicated region that includes *valkea*. Developmental stage was the main factor explaining

most of the genome-wide variation in gene expression between samples (Figure S3). Such a

pattern would be expected as many genes are involved in development and thus are likely to

be differentially expressed (DE) between developmental stages. No apparent clustering can

be observed among samples of the same colour morph.

We next compared gene expression between yy and WW individuals at each of the

developmental stages. From the 22 genes identified in the GWAS and QTL analysis, only

two, *yellow-e* and *valkea*, were differentially expressed in any of the comparisons. *Valkea*

was overexpressed in white individuals in the pre-melanin stage with a Log Fold Change of

10.32 and a p-value of 2.18e-06. As *valkea* is only fully present in the W genome, it is not

expected to be expressed at all in the Y genome. *Yellow-e* was also overexpressed in white

individuals with a Log Fold Change of 3.86 and adjusted p-value of 5.62e-06. In other

developmental stages, neither *valkea* nor *yellow-e* showed differences in expression between

morphs (Figure 3).

The GWAS and QTL peaks of association are situated in scaffold 419 of the WW reference

assembly, which in a WW linkage map forms a linkage group along with 6 more scaffolds

(472, 487, 515, 531, 540 and 609). We found that 12 genes present in this linkage group were

differentially expressed, including *valkea* and *yellow-e* (Table S2), and identified their

orthologues in *Drosophila melanogaster*.

We next conducted a genome-wide analysis of the RNA-seq data. Lowly expressed genes

were filtered out with 10,920 genes retained. Overall, 99 genes were differentially expressed

(FDR < 0.05) between the two morphs. Of these, 49 were upregulated in the yy morph, while

the remaining 50 were upregulated in WW individuals. The earliest developmental stage, 72-

hours, was the stage with the highest number of DE genes (n = 48), while the 5-day old stage

had the fewest (n=7). One gene which encodes a C2H2 zinc finger transcription factor in *D.*

melanogaster, *'jg15945',* was over-expressed in yy in the first three stages (Figure S4).

Figure 3: (A) RNA-seq of developing wings shows 99 genes that are differentially expressed between
morphs. (B) *Valkea* is highly expressed in the white morph at the pre-melanin stage. Expression of
valkea in yellow morphs is around 0.

Yellow-e is not associated with colour

We next used a tree topology approach to explore the relationships between alleles around the

duplicated sites among the populations. For four populations (Finnish white and yellow,

Estonian and Scottish populations), three topologies are possible; grouping populations by

colour, by geography and one grouping neither by colour nor geography. At *yellow-e*, the

highest weighting is towards the geography grouping, suggesting that this gene is not

associated with the colour switch (Figure S5).

	Pigment analysis

Since the *yellow* gene family, to which *valkea* is related, is known to be responsible for the

production of melanin pigments, we further investigated the identity of the wing

pigmentation. First, we ruled out the presence of several non-melanin pigment types in the

hindwings, including pterins and carotenoids. Pterins are commonly found in insects and,

along with purine derivatives, papiliochromes and flavonoids, are soluble in strong acids and

bases or in organic solvents (Umebachi, 1975; Kayser, 1985; Shamim *et al.*, 2014). We

placed wing samples from each morph in NaOH overnight, then measured the absorbance of

the supernatant using a spectrophotometer. We also left wings in methanol overnight before

measuring the supernatant. The spectra did not show any peaks indicative of any pigment

dissolved in the sample. Similarly, we found no evidence for carotenoid pigments after

dissolving in a hexane:tert-butyl methyl ether solution (Figure S6). Wings did not fluoresce

under UV light, providing further evidence for the lack of fluorescent pigments including

pterins, flavonoids, flavins and papiliochromes (Umebachi, 1975; Kayser, 1985).

Ommochromes are red and yellow pigments; high performance liquid chromatography

(HPLC) ruled out the presence of these pigments on the moth wings, which we compared to

data from ommochrome-containing *Heliconius* wings and a Xanthurenic acid standard

(Figure S7).

HPLC analysis showed peaks characteristic of pheomelanin (Figure S8). Pheomelanins

produce red-brown colour in grasshoppers and wasps (Galván *et al.*, 2015; Jorge García,

Polidori and Nieves-Aldrey, 2016), and orange-red colours in ants and bumblebees (Hines *et*

al., 2017; Polidori, Jorge and Ornosa, 2017). Insects generally have dopamine-derived

pheomelanin and a breakdown product of this is 4-Amino-3-hydroxyphenylethylamine (4-

AHPEA) (Barek *et al.*, 2018). Yellow wings showed large peaks for 4-AHPEA. White wings

had around 27% of the 4-AHPEA levels seen in yellow wings, and black sections of the

wings had 16%. Hydrogen iodide hydrolysis of wings produced the isomer 3-AHPEA, which

may come from 3-nitrotyramine originating from the decarboxylation of 3-nitrotyrosine.

Reduction of 3-nitrotyrosine produces 3-AHP, another marker of pheomelanin (Wakamatsu,

Ito and Rees, 2002).

Analysis of the black portions of the wing found pyrrole-2,3-dicarboxylic acid (PDCA) and

pyrrole-2,3,5-tricarboxylic acid (PTCA) (Figure S9). Both are components of eumelanin

(Barek *et al.*, 2018), suggesting that the black colouration seen in the wood tiger moth is

predominantly eumelanin derived from dopamine. This is common in producing black

colouration and providing structural components of the exoskeleton. In addition, dopamine is

acylated to both *N*-β-alanyldopamine (NBAD) and *N*-acetyldopamine (NADA) sclerotins.

NADA sclerotins are colourless and likely to be present on the white wings. This analysis of

pigmentation is therefore consistent with a role for *yellow* family genes in regulating the

colour polymorphism.

Discussion

Hindwing colouration of male *Arctia plantaginis* is polymorphic and these colour morphs

vary in multiple behavioural and life-history traits, providing an example of a complex

polymorphism. Here, we have shown that variation in male hindwing colour is associated

with a narrow region in the genome, characterised by a duplicated sequence found only in

white morphs and containing a gene from the *yellow* gene family. The white-specific copy,

valkea, is highly expressed during pupal development, consistent with genetic dominance of

the white allele. The presence of only one gene is in contrast to 'classical' supergenes which

often explain the genetic architecture of complex polymorphisms (Thompson and Jiggins,

2014) and involve chromosomal rearrangements containing multiple genes that allow the

maintenance of co-adapted haplotypes (Joron *et al.*, 2006; Lowry and Willis, 2010; Küpper *et*

al., 2016; Funk *et al.*, 2021).

Although we did not find evidence for a supergene involving multiple loci, this morph-

specific duplication still provides a region of reduced recombination between morphs. This is

similar to the genetic architecture of the *Primula* supergene controlling heterostyly, which

involves a large duplication containing five genes (Huu *et al.*, 2020). The duplicated region is

effectively hemizygous and cannot recombine except in homozygote genotypes, which could

contribute to the maintenance of the complex polymorphism and the linkage of multiple

traits. In polymorphic *Papilio dardanus*, one colour pattern morph is associated with a

duplicated region, again providing physical constraints on recombination (Timmermans *et*

al., 2014). Nonetheless, in the case of the wood tiger moth, it remains unclear how a single

gene, such as *valkea*, can control the development of a broad array of phenotypic traits.

One possible mechanism is that there is a regulatory element along the scaffold which is

controlling colour via the *valkea* gene, but also regulating other genes to control different

phenotypic traits. We found the most significant markers and SNPs located in a non-coding region close to the *yellow* genes, which likely contains a cis-regulatory element (CRE) controlling transcription of *valkea*. In cichlids, for example, a CRE at the gene encoding agouti-related peptide 2 controls variation in strip patterning in two closely related species (Kratochwil *et al.*, 2018).

Differential expression of other genes on the same chromosome controlled by the CRE could explain variation in covarying traits. The over-expression of another gene, possibly encoding a zinc transcription factor, in yellow individuals in the early pupal stages suggests that there is differential expression of unlinked genes as a result of the polymorphism, although since this gene is on a different chromosome to *valkea* it is unlikely to be directly controlled by the CRE. Another hypothesis is that somehow *valkea* itself regulates other genes. However, *yellow* family genes are not known to regulate transcription of other genes, unlike for example *doublesex*, a transcription factor that undergoes alternative splicing and female mimetic wing pattern polymorphism in *Papilio polytes* (Kunte *et al.*, 2014; Nishikawa *et al.*, 2015).

The *yellow* family genes are highly conserved throughout insects (Ferguson *et al.*, 2011). They have been widely linked to colouration (Wittkopp, Vaccaro and Carroll, 2002; Miyazaki *et al.*, 2014; Zhang *et al.*, 2017; Zhang, Mazo-Vargas and Reed, 2017), as well as behaviour, sex-specific phenotypes and reproductive maturation (Wilson *et al.*, 1976). These genes share a common origin with the major royal jelly protein (MRJP) genes (Drapeau *et al.*, 2006) which are crucial in caste development in honeybees. Like the MRJP genes, *yellow* genes in honeybees have diverse spatial and temporal expression patterns. As our focus in *A. plantaginis* has been on wing tissue, we are missing expression of genes in other tissues that could be linked to other traits. Thus, it is not impossible to imagine that a *yellow* gene could have a similar function to a MRJP in regulating the development of a complex phenotype. Recent work with *Bicyclus anynana* showed that *yellow* functions as a repressor of male courtship (Connahs *et al.*, 2022). On the other hand, sex specific behavioural phenotypes of *yellow* mutants in *Drosophila* were found to be due to pigmentation effects (Massey *et al.*, 2019), so more evidence is needed to suggest a functional role for *yellow* genes outside of pigmentation.

The duplication of *yellow-e* and surrounding regions in the white morphs suggests that the yellow morph is the ancestral form. *Valkea* could have evolved in a stepwise fashion, first as

a tandem duplication then with a stop codon mutation altering the gene structure. Gene

duplications can facilitate adaptation and, in some examples, lead to polymorphism. Although

both *valkea* and *yellow-e* are overexpressed in the white morph, the topology weighting

suggests that *yellow-e* is not closely associated with the colour change. *Valkea* likely

represents an example of neofunctionalization, where the duplicated gene gains a different

function to the original gene copy.

Contrary to previous work that attributed red and yellow colours to pterins in another tiger

moth species (Gawne and Nijhout, 2019), we found high levels of 4-AHPEA in the yellow

wings confirming the presence of pheomelanins. These pigments have been widely associated

with red and yellow colours in mammals (e.g. Mcgraw and Wakamatsu, 2004), but only

relatively recently described in insects and likely to be more widespread than previously

thought. Yellow colours can also be produced by NBAD sclerotins which are sclerotizing

precursor molecules made from dopamine and these have an important role in the

sclerotization pathway for hardening the insect cuticle (Andersen, 2007; Barek, Evans and

Sugumaran, 2017) before becoming involved in melanisation (Barek *et al.*, 2018). Thus, we

suggest that the yellow colour arises partly from the NBAD sclerotins and partly from the

presence of pheomelanin pigments, which has been proposed in other Lepidoptera (Matsuoka

and Monteiro, 2018). While some 4-AHPEA also occurred in white wings, this may be due to

its role in production of cross-linking cuticular proteins and chitin during sclerotization

(Sugumaran, 2010). Upregulation of genes on the white allele could be acting as a repressor

of the generation of yellow colour. We suspect that *yellow* family genes play multiple roles

within the melanin production pathway. In the wood tiger moth, *yellow* affects the conversion

of DOPA into black dopamine melanin (Galarza, 2021). *Yellow-e* in particular has been

linked to larval colouration in *Bombyx mori* (Ito *et al.*, 2010) and adult colour in beetles

(Wang *et al.*, 2022), while another gene, *yellow-f*, has a role in eumelanin production (Barek

et al., 2018).

In summary, we identified a structural variant containing a previously undescribed gene,

valkea, which is only present in white morphs of *A. plantaginis*. While functional studies are

needed to determine the exact function of this locus, the presence of a regulatory element

controlling wing colour and other traits via multiple downstream effects could explain how

multiple traits are linked to wing colouration. This complex polymorphism allows multiple

beneficial phenotypes to be inherited together, whereas recombination would separate

multiple loci leading to maladapted individuals. Our results provide the basis for further

exploration of the genetic basis of covarying behavioural and life-history traits, and offer an

intriguing example of how complex polymorphisms are maintained.

Methods

Sampling

Homozygous lines of white (WW) and yellow (yy) *Arctia plantaginis* moths were created

from Finnish populations at the University of Jyväskylä, Finland. Larvae were fed with wild

dandelion (*Taraxacum sp.*) and reared under natural light conditions, with an average day

temperature of 25°C and night temperature between 15-20°C. For the crosses, a heterozygous

male, created from crossing a heterozygous male with a homozygous yy female, was

backcrossed with a yy female. This was repeated to obtain four families totalling 172

offspring and 8 parents (Table S3). Samples from wild populations were caught in Southern

Finland (n=10) and Central Finland (n=20), where male morphs are either white or yellow,

Estonia (n=4), where males are mostly white, and Scotland (n=4), where males are yellow

(Table S4). In addition, we included 8 samples which are F1 offspring of wild Scottish

samples. Forty pupae with known genotypes from lab populations (20 WW and 20 yy) were

used for the RNA extractions.

DNA extraction and sequencing

For the lab crosses, DNA was extracted from two legs crushed with sterilised PVC pestles

using a Qiagen DNeasy Blood & Tissue kit, following the manufacturer's instructions.

Library preparation and GBS sequencing were performed by BGI Genomics on an Illumina

HiSeq X Ten. For the wild samples, DNA was extracted from the thoraces also with a Qiagen

kit. Library preparation and sequencing were performed by Novogene (Hong Kong, China).

150bp paired-end reads were sequenced on an Illumina NovaSeq 6000 platform.

Linkage mapping analysis

FASTQ reads were mapped using bowtie v2.3.2 (Langmead and Salzberg, 2012) to the

yellow *Arctia plantaginis* scaffold-level genome assembly (Yen *et al.*, 2020). BAM files

were sorted and indexed using SAMtools v.1.9 (Li *et al.*, 2009) and duplicates removed using

PicardTools MarkDuplicates (broadinstitute.github.io/picard). Twelve samples which had

aligned <30% were removed. Reads of the remaining samples had an average alignment of

94%. SNPs were called using SAMtools mpileup with minimum mapping quality set to 20

and bcftools call function. Lep-MAP3 (Rastas, 2017) was used for linkage map construction

and we ran the following modules: ParentCall2 which called 105,622 markers, Filtering2,

SeparateChromosomes2 with lodLimit=5 and sizeLimit=100, JoinSingles2All and

OrderMarkers2 with recombination2=0 to denote the lack of female recombination.

Genotypes were phased using the map2genotypes.awk script included with Lep-MAP3.

Markers were named based on the genomic positions of the SNPs in the reference genome

and the map.awk script, and this was used to further order the markers within the linkage

groups. This resolved 30 linkage groups. Although we expect that there are 31 chromosomes

in the moth genome, we suspect that the sex chromosome is missing in this assembly as the

yy individual used in the genome assembly was female (Yen *et al.*, 2020). A small number of

markers which caused long gaps at the beginning or end of linkage groups were manually

removed, leaving the final map 948.7cM long with 19,803 markers. Markers were well

distributed so we began the first analyses with this map. A linkage map was also assembled

using sequences aligned to the white reference and this separated into 31 linkage groups.

The QTL analysis was carried out in R/qtl (Broman *et al.*, 2003). Genotype probabilities were

calculated before running a genome scan using the *scanone* function with the Haley-Knott

method and binary model parameters, and including family as an additive covariate. The

phenotype was labelled as either 0 (Wy) or 1 (yy). We ran 5000 permutations to determine

the significance level for the QTL LOD scores. The *bayesint* function calculated the 95%

Bayesian confidence intervals around the significant marker.

Analysis of whole genome sequences

FASTQ reads were mapped to the yellow *A. plantaginis* genome assembly (Yen *et al.*, 2020)

using BWA-MEM v7.17 (Li, 2013). As before, BAM files were sorted and indexed, and

duplicates were removed. Genotyping and variant calling was carried out with the Genome

Analysis Toolkit (GATK) (McKenna *et al.*, 2010). Variants were called using

HaplotypeCaller (v.3.7) in GVCF mode, then gVCFs combined with GenomicsDBImport

(v.4.0). Joint genotyping was run with GenotypeGVCFs, set with a heterozygosity of 0.01,

and SNPs were called using SelectVariants. Finally, the set of 20,787,772 raw SNPs were

filtered using VariantFiltration and thresholds: quality by depth (QD > 2.0), root mean square

mapping quality (MQ > 50.0), mapping quality rank sum test (MQRankSum > −12.5), read

position rank sum test (ReadPosRankSum > −8.0), Fisher strand bias (FS < 60.0), and strand

odds ratio (SOR < 3.0). A set of 5,227,288 SNPs passed the filtering.

We carried out a Genome Wide Association Study (GWAS) using the R package GenABEL

v.1.8 (Aulchenko *et al.*, 2007). The set of filtered SNPs were converted to BED format with

PLINK2, keeping only biallelic SNPs (cog-genomics.org/plink/2.0/). Sites which were not in

Hardy-Weinberg equilibrium (p<0.01), or had a call rate of <0.5, were excluded. Following

this, 381,266 sites were retained across 40 individuals (out of 57). To account for population

stratification, we performed multidimensional scaling on kinship and identity-by-state (IBS)

information estimated from the data, and included this as a covariate in the association test.

Significance levels were calculated using Bonferroni corrected thresholds to account for

multiple testing. Central and Southern Finnish populations were pooled for this analysis,

based on a previous principal component analysis of these samples (Yen *et al.*, 2020).

In Yen et al. (2020), many of these samples were processed in the same way but aligned to

the white genome assembly (based on a white individual). We used the VCF file from this

study for a Twisst analysis (Martin and Van Belleghem, 2017) which included samples from

the Finnish, Estonian and Scottish populations. A minor allele frequency of 3 was applied to

the VCF with VCFtools, and this was phased using Beagle v5.1 (Browning, Zhou and

Browning, 2018). We generated neighbour joining trees with a window size of 50 using

PhyML and ran Twisst using scripts from github.com/simonhmartin/twisst. Read depth of the

W-mapped samples was calculated in 1kb windows across the candidate region using

BEDtools (v.2.20.1) multicov (Quinlan and Hall, 2010).

For analysis of structural variants, sequences from the white and yellow genome assemblies

were aligned using MAFFT v7.450 (Katoh and Standley, 2013) and viewed with Geneious.

Our focal sequence, scaffold 419 in the white genome, is the reverse complement of scaffold

206 in the yellow genome.

Identification of candidate genes and tree construction

To identify candidate genes in the QTL interval and GWAS region, we ran a protein

BLASTP v.2.4.0 search to identify *Heliconius melpomene* (Hmel2.5) proteins homologous to

predicted *A. plantaginis* proteins in the region from the genome annotation. Informative gene

names were obtained by performing a BLASTP search with the *H. melpomene* proteins

against all *Drosophila melanogaster* proteins in FlyBase v.FB2020_01 (flybase.org/blast).

For the *yellow* gene tree, we used Lepbase (Challi *et al.*, 2016) to search for *yellow* genes in *Bombyx mori* (ASM15162v1). We identified *yellow-e* in *H. melpomene* by searching for major royal jelly proteins, then comparing protein sequences of these against *Drosophila* proteins in Flybase. The sequence for *Dmel yellow-e* was downloaded from Flybase. To make the tree, coding sequences of all genes were aligned in Geneious using MAFFT v7.450 (Katoh and Standley, 2013), then the tree was constructed with PhyML using 10 bootstraps (Guindon *et al.*, 2010).

Differential gene expression

We dissected the wings out of the pupae in Cambridge, UK. Pupae and larvae were sent to Cambridge from Jyväskylä and were kept between 22 and 30˚C. Pupae were sexed and only males were used. Dissections were made at 4 different stages: 72 hours after pupation (72h), 5 days after pupation (5d; counting 0-24 first hours after pupation as day 1), pre melanin deposition (Pre-mel) and post melanin deposition (Mel). We sampled 5 individuals per stage and genotype. Hindwings and forewings were stored separately in RNA-later (Sigma-Aldrich) at 4 ˚C for 2 weeks and later transferred to -20 ˚C, while the rest of the body was stored in pure ethanol.

Total RNA was extracted from hindwing tissue using a standard hybrid protocol. First, we transferred the wing tissue into Trizol Reagent (Invitrogen) and homogenised it using dounce tissue grinders (Sigma-Aldrich). Then, we performed a chloroform phase extraction, followed by DNase treatment (Ambion) for 30 mins at 37˚C. We measured the concentration of total RNA using Qubit Fluorometric Quantitation (Thermofisher) and performed a quality check using an Agilent 4200 TapeStation (Agilent). The extracted total RNA was stored at -20˚C before being sent to Novogene UK for sequencing.

We performed quality control and low-quality base and adapter trimming of the sequence data using *TrimGalore!* We then mapped the trimmed reads to the two *A. plantaginis* genomes using STAR. We performed a second round of mapping (2pass) including as input the output splice junctions from the first round. The *A. plantaginis* genome annotations WW and YY were included in each round of mapping respectively. We then used *FeatureCounts* to count the mapped reads. Finally, we used DESeq2 to analyse the counts and perform the DE analysis.

To identify the gene or genes controlling the development of wing colour in *A. plantaginis*, we performed a genome wide differential expression analysis using limma-voom (Ritchie *et al.*, 2015). First, we defined a categorical variable, 'GenStage', with 8 levels containing the genotype and stage information of every individual sample (e.g. YY72h, WW72h, YY5days, etc.). Then, we built the design matrix fitting a model with GenStage as the only fixed effect factor contributing to the variance in gene expression and included family as a random effect factor (gene expression ~ 0 + GenStage + (1|Family)). We then filtered lowly expressed genes, which resulted in a reduction of the number of tested genes from 17,615 to 11,330 genes in the Y-mapped analysis and 17,930 to 10,920 in the W-mapped one. Finally, the list of genes that are differentially expressed in each stage were extracted using the Benjamini-Hochberg procedure to correct for multiple testing.

Orthology assignment

To infer genome wide orthology between *A. plantaginis* and *D. melanogaster*, we used OrthoFinder (v2.5.4) (Emms and Kelly, 2019). We used proteomes from 6 Lepidoptera species, *Plutela xylostella* (GCA_905116875_2)*, Bombyx mori* (GCF_014905235_1)*, Spodoptera frugiperda* (GCF_011064685_1)*, Parnassius apollo* (GCA_907164705_1)*, Pieris macdunnoughi* (GCA_905332375_1) *amd Pararge aegeria* (GCF_905163445_1) and *Drosophila melanogaster* (GCF_000001215_4)*. We ran the primary_transcript.py utility from OrthoFinder to extract only one transcript per protein, and then ran OrthoFinder with default settings.

Pigment analysis

Solubility and fluorescence tests

Five hindwings from each morph were placed in two separate solvents (0.1M NaOH and 90% MeOH) and left overnight. The supernatant was analysed with an Agilent Cary 8454 UV-Visible spectrophotometer and the spectra compared to known spectra for various pigments. A UV lamp (Philips TL8W/08 F8T5/BLB) used to test for fluorescence on the wings. The presence of carotenoids was tested by placing wings into 1ml of pyridine and leaving at 95°C for 4 hours (McGraw *et al.*, 2002). To these we added 1ml of 1:1 hexane:tert-butyl methyl ether and 2ml of water before shaking and leaving overnight. Again, the supernatant was measured with the spectrometer.

High-Performance Liquid Chromatography (HPLC) to test to eumelanin and ommochrome pigments

To determine the type of melanin producing the black colour on the wings, we cut out approximately 5mg of the black sections of the wings, from both females and males. Eumelanin analysis was carried out according to (Borges *et al.*, 2001). Each sample was added to a tube containing 820µl 0.5 M NaOH, 80µl 3% H_2O_2 and an internal standard (48nmol phthalic acid) and heated in a boiling water bath for 20 minutes. Once cool, 20µl of 10% Na_2SO_3 and 250µl of 6M HCl was added. Samples were then extracted twice with 7ml of ethyl acetate. Ethyl acetate was dried at 45 °C under a stream of nitrogen. The residue was dissolved into 0.5ml of 0.1% formic acid.

We carried out high-performance liquid chromatography on an Agilent 1100 HPLC. 20µl of the sample was injected into a Waters Atlantis T3, 100 x 3.0mm i.d. analytical column (Waters, Milford, MA, USA). The column was set to 25°C and analytes were detected at wavelength 280nm. The HPLC mobile phase consisted of two eluents: UHQ-water/MeOH (98:2; v/v) with 0.1% formic acid and UHQ-water/MeOH (40/60; v/v) with 0.1% formic acid. Flow rate was 0.4 ml/min and the used gradient started with 100% of eluent A and ramped evenly from time 0 to 15 minutes to 40:60 (A:B; v/v), held at 40:60 for 6 minutes and ramped evenly back to initial eluent composition (100% A) over 5 minutes. We compared chromatograms obtained from the samples to the chromatograms obtained from synthetic melanin, ink from sepia officinalis and black human hair.

HPLC was also applied to observe the possible presence of ommochrome pigments. Injection volume was 10µl and for the separation we used the same Waters Atlantis T3 column (100 x 3.0 mm i.d.) set to 30°C. Solvent A was UHQ-water and B was acetonitrile (ACN), both containing 0.1% formic acid. Flow rate was 0.4ml/min and the used gradient was as follows: Initial flow ratio was 98/2 water/ACN (v/v) ramping then evenly from time 1 to 15 min to 30:70 water:ACN (v/v), held for 1.5 min and then ramped evenly back to initial eluent composition over 0.5 minutes. The column was stabilised for 7 minutes before a new run.

Pheomelanin analysis

Samples were analysed for pheomelanin content according to the method of Kolb *et al.* (1997) with modifications. A 2mg sample was placed in a screw-capped tube with 100µl water, 500µl ~55-58% hydrogen iodide (HI), and 20µl 50% hypophosphorous acid (H_3PO_2).

Samples were capped tightly and hydrolysed for 20 hours at 130°. After cooling, samples

were evaporated under nitrogen flow, then dissolved in 1ml of 0.1M HCl and purified with

solid phase extraction. Strata SCX cartridges were preconditioned with 2ml of methanol, 3ml

of water and 1ml of 0.1M HCl. Sample was then applied to the cartridge, washed with 1ml of

0.1M HCl, and finally eluted with 1ml of methanol (MeOH): 0.5M ammonium acetate

($NH_4CH_3CO_2$) (20:80 v/v).

Hydrogen Iodide hydrolysis products were determined by a Dionex high-performance liquid

chromatograph equipped with pulsed amperometric detection (HPLC/PAD). A Phenomenex

Kinetex C18 column (150 x 4.6 mm i.d.; 5μm particle size) with a gradient elution (Table S5)

at a flow rate of 0.9 ml min^{-1} with the eluents: (A) sodium citrate buffer (Hines *et al.*, 2017)

in ultra-high-quality water (internal resistance $\geq$ 18.2 MΩ cm; Milli-Q Plus; Millipore,

Bedford, MA, USA) and (B) methanol were used for the separation. Dionex ED-50 pulsed

amperometric detector (Dionex, Sunnyvale, CA, USA) equipped with a disposable working

electrode by using a Dionex waveform A with potentials presented in Table S6 was used for

detection. 4-AHP, 3-AHPEA and 4-AHPEA standards used in calibration were kindly offered

from the laboratory of Emeritus Professor Wakamatsu and the preparation method is

described in (Wakamatsu *et al.*, 2014).

Acknowledgements

We thank Kaisa Suisto and the greenhouse staff for insect rearing, and Elisa Salmivirta and

Sari Viinikainen for lab assistance. Thanks to Emeritus Prof. Shosuke Ito for providing

pheomelanin standards and to Bodo Wilts for advice on the pigment analyses.

This work was supported by Academy of Finland grants to MB (#343356) and JM (projects

345091 and 328474).

Data availability

All the scripts used for mapping and DE analysis will be uploaded to an online repository.

Scripts and data for the QTL and GWAS analyses can currently be found at

github.com/mnbrien/Aplantaginis-male-colour. Raw sequencing data of wild samples has

previously been deposited in ENA, study accession No. PRJEB36595.

References

Andersen, S. O. (2007) 'Involvement of tyrosine residues, N-terminal amino acids, and β-alanine in insect cuticular sclerotization', *Insect Biochemistry and Molecular Biology*, 37(9), pp. 969–974. doi: https://doi.org/10.1016/j.ibmb.2007.05.002.

Andrade, P. *et al.* (2019) 'Regulatory changes in pterin and carotenoid genes underlie balanced color polymorphisms in the wall lizard', *Proceedings of the National Academy of Sciences of the United States of America*, 116(12), pp. 5633–5642. doi: 10.1073/pnas.1820320116.

Aulchenko, Y. S. *et al.* (2007) 'GenABEL: an R library for genome-wide association analysis', *Bioinformatics*, 23(10), pp. 1294–1296. doi: 10.1093/bioinformatics/btm108.

Barek, H. *et al.* (2018) 'Insect cuticular melanins are distinctly different from those of mammalian epidermal melanins', *Pigment Cell and Melanoma Research*, 31(3), pp. 384–392. doi: 10.1111/pcmr.12672.

Barek, H., Evans, J. and Sugumaran, M. (2017) 'Unraveling complex molecular transformations of N-β-alanyldopamine that account for brown coloration of insect cuticle', *Rapid Communications in Mass Spectrometry*, 31(16), pp. 1363–1373. doi: https://doi.org/10.1002/rcm.7914.

Borges, C. R. *et al.* (2001) 'Relationship of Melanin Degradation Products to Actual Melanin Content: Application to Human Hair', *Analytical Biochemistry*, 290(1), pp. 116–125. doi: https://doi.org/10.1006/abio.2000.4976.

Briolat, E. S. *et al.* (2019) 'Diversity in warning coloration: selective paradox or the norm?', *Biological Reviews*, 94(2), pp. 388–414. doi: 10.1111/brv.12460.

Broman, K. W. *et al.* (2003) 'R/qtl: QTL mapping in experimental crosses', *Bioinformatics*, 19(7), pp. 889–890. doi: 10.1093/bioinformatics/btg112.

Browning, B. L., Zhou, Y. and Browning, S. R. (2018) 'A One-Penny Imputed Genome from Next-Generation Reference Panels', *The American Journal of Human Genetics*, 103(3), pp. 338–348. doi: https://doi.org/10.1016/j.ajhg.2018.07.015.

Challi, R. J. *et al.* (2016) 'Lepbase: the Lepidopteran genome database', *bioRxiv*, p. 56994. doi: 10.1101/056994.

Connahs, H. *et al.* (2022) 'The yellow gene regulates behavioural plasticity by repressing male courtship in Bicyclus anynana butterflies', *Proceedings of the Royal Society B: Biological Sciences*, 289(1972), p. 20212665. doi: 10.1098/rspb.2021.2665.

Cott, H. B. (1940) *Adaptive coloration in animals*. London: Methuen.

Cuthill, I. C. *et al.* (2017) 'The biology of color', *Science*, 357(6350). doi: 10.1126/science.aan0221.

Drapeau, M. D. *et al.* (2006) 'Evolution of the Yellow/Major Royal Jelly Protein family and the emergence of social behavior in honey bees', *Genome Research*, 16(11), pp. 1385–1394.

doi: 10.1101/gr.5012006.

Emms, D. M. and Kelly, S. (2019) 'OrthoFinder: phylogenetic orthology inference for
comparative genomics', *Genome Biology*, 20(1), p. 238. doi: 10.1186/s13059-019-1832-y.

Endler, J. A. (1988) 'Frequency-dependent predation, crypsis and aposematic coloration',
Philosophical Transactions of the Royal Society B: Biological Sciences, 319, pp. 505–523.

Ferguson, L. C. *et al.* (2011) 'Evolution of the insect yellow gene family', *Molecular Biology*
and Evolution, 28(1), pp. 257–272. doi: 10.1093/molbev/msq192.

Funk, E. R. *et al.* (2021) 'A supergene underlies linked variation in color and morphology in
a Holarctic songbird', *Nature Communications*, 12(1), p. 6833. doi: 10.1038/s41467-021-
27173-z.

Galarza, J. A. *et al.* (2014) 'Temporal relationship between genetic and warning signal
variation in the aposematic wood tiger moth (Parasemia plantaginis)', *Molecular Ecology*,
23(20), pp. 4939–4957. doi: 10.1111/mec.12913.

Galarza, J. A. (2021) 'Comparative transcriptomics of albino and warningly-coloured
caterpillars', *Ecology and Evolution*, 11(12), pp. 7507–7517. doi:
https://doi.org/10.1002/ece3.7581.

Galván, I. *et al.* (2015) 'Insects synthesize pheomelanin', *Pigment Cell and Melanoma*
Research, 28(5), pp. 599–602. doi: 10.1111/pcmr.12397.

Gawne, R. and Nijhout, H. F. (2019) 'Expanding the nymphalid groundplan's domain of
applicability: pattern homologies in an arctiid moth (Utetheisa ornatrix)', *Biological Journal*
of the Linnean Society, 126(4), pp. 912–924. doi: 10.1093/biolinnean/bly193.

Gordon, S. P. *et al.* (2015) 'Colour polymorphism torn apart by opposing positive frequency-
dependent selection, yet maintained in space', *Journal of Animal Ecology*, 84(6), pp. 1555–
1564. doi: 10.1111/1365-2656.12416.

Guindon, S. *et al.* (2010) 'New Algorithms and Methods to Estimate Maximum-Likelihood
Phylogenies: Assessing the Performance of PhyML 3.0', *Systematic Biology*, 59(3), pp. 307–
321. doi: 10.1093/sysbio/syq010.

Hegna, R. H., Galarza, J. A. and Mappes, J. (2015) 'Global phylogeography and geographical
variation in warning coloration of the wood tiger moth (Parasemia plantaginis)', *Journal of*
Biogeography, 42(8), pp. 1469–1481. doi: 10.1111/jbi.12513.

Hines, H. M. *et al.* (2017) 'Melanic variation underlies aposematic color variation in two
hymenopteran mimicry systems', *PLoS ONE*, 12(7), pp. 1–17. doi:
10.1371/journal.pone.0182135.

Huu, C. N. *et al.* (2020) 'Supergene evolution via stepwise duplications and
neofunctionalization of a floral-organ identity gene', *Proceedings of the National Academy of*
Sciences of the United States of America, 117(37), pp. 23148–23157. doi:
10.1073/pnas.2006296117.

Huxley, J. (1955) 'Morphism and evolution', *Heredity*, 9(1), pp. 1–52. doi:

10.1038/hdy.1955.1.

Ito, K. *et al.* (2010) 'Yellow-e determines the color pattern of larval head and tail spots of the
silkworm Bombyx mori', *Journal of Biological Chemistry*, 285(8), pp. 5624–5629. doi:
10.1074/jbc.M109.035741.

Jorge García, A., Polidori, C. and Nieves-Aldrey, J. L. (2016) 'Pheomelanin in the secondary
sexual characters of male parasitoid wasps (Hymenoptera: Pteromalidae)', *Arthropod*
Structure & Development, 45(4), pp. 311–319. doi: https://doi.org/10.1016/j.asd.2016.05.001.

Joron, M. *et al.* (2006) 'A Conserved Supergene Locus Controls Colour Pattern Diversity in
Heliconius Butterflies', *PLoS Biology*, 4(10), pp. 1831–1840. doi:
10.1371/journal.pbio.0040303.

Katoh, K. and Standley, D. M. (2013) 'MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability', *Molecular Biology and Evolution*,
30(4), pp. 772–780. doi: 10.1093/molbev/mst010.

Kayser, H. (1985) 'Pigments', in Kerkut, G. A. and Gilbert, L. I. (eds) *Comparative Insect*
Physiology, Biochemistry, and Pharmacology. Oxford: Pergamon Press, pp. 367–415.

Kolb, A. M. *et al.* (1997) 'Determination of Pheomelanin by Measurement of
Aminohydroxyphenylalanine Isomers with High-Performance Liquid Chromatography',
Analytical Biochemistry, 252(2), pp. 293–298. doi: https://doi.org/10.1006/abio.1997.2342.

Kratochwil, C. *et al.* (2018) 'Agouti-related peptide 2 facilitates convergent evolution of
stripe patterns across cichlid fish radiations', *Science*, 362(6413), pp. 457–460. doi:
10.1126/science.aao6809.

Kunte, K. *et al.* (2014) 'doublesex is a mimicry supergene', *Nature*, 507(7491), pp. 229–232.
doi: 10.1038/nature13112.

Küpper, C. *et al.* (2016) 'A supergene determines highly divergent male reproductive morphs
in the ruff', *Nature Genetics*, 48(1), pp. 79–83. doi: 10.1038/ng.3443.

Langmead, B. and Salzberg, S. L. (2012) 'Fast gapped-read alignment with Bowtie 2',
Nature Methods, 9, p. 357. Available at: http://dx.doi.org/10.1038/nmeth.1923.

Li, H. *et al.* (2009) 'The Sequence Alignment/Map format and SAMtools', *Bioinformatics*,
25(16), pp. 2078–2079. doi: 10.1093/bioinformatics/btp352.

Li, H. (2013) 'Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM', *arXiv*, p. 1303.3997.

Linnen, C. R. *et al.* (2013) 'Adaptive Evolution of Multiple Traits Through Multiple
Mutations at a Single Gene', *Science*, 339(6125), pp. 1312–1316. doi:
10.1126/science.1233213.

Lowry, D. B. and Willis, J. H. (2010) 'A Widespread Chromosomal Inversion Polymorphism
Contributes to a Major Life-History Transition, Local Adaptation, and Reproductive
Isolation', *PLOS Biology*, 8(9), p. e1000500. Available at:
https://doi.org/10.1371/journal.pbio.1000500.

Martin, S. H. and Van Belleghem, S. M. (2017) 'Exploring Evolutionary Relationships
Across the Genome Using Topology Weighting', *Genetics*, 206(1), pp. 429 LP – 438. doi:
10.1534/genetics.116.194720.

Massey, J. H. *et al.* (2019) 'The yellow gene influences Drosophila male mating success
through sex comb melanization', *eLife*. Edited by K. VijayRaghavan and C. L. Peichel, 8, p.
e49388. doi: 10.7554/eLife.49388.

Matsuoka, Y. and Monteiro, A. (2018) 'Melanin Pathway Genes Regulate Color and
Morphology of Butterfly Wing Scales', *Cell Reports*, 24(1), pp. 56–65. doi:
10.1016/j.celrep.2018.05.092.

McGraw, K. J. *et al.* (2002) 'The effect of dietary carotenoid access on sexual dichromatism
and plumage pigment composition in the American goldfinch', *Comparative Biochemistry
and Physiology Part B: Biochemistry and Molecular Biology*, 131(2), pp. 261–269. doi:
https://doi.org/10.1016/S1096-4959(01)00500-0.

Mcgraw, K. J. and Wakamatsu, K. (2004) 'Melanin Basis of Ornamental Feather Colors in
Male Zebra Finches', *The Condor*, 106(3), pp. 686–690. doi: 10.1093/condor/106.3.686.

McKenna, A. *et al.* (2010) 'The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data', *Genome Research*, 20, pp. 1297–1303.
doi: 10.1101/gr.107524.110.

McKinnon, J. S. and Pierotti, M. E. R. (2010) 'Colour polymorphism and correlated
characters: Genetic mechanisms and evolution', *Molecular Ecology*, 19(23), pp. 5101–5125.
doi: 10.1111/j.1365-294X.2010.04846.x.

Miyazaki, S. *et al.* (2014) 'Sexually Dimorphic Body Color Is Regulated by Sex-Specific
Expression of Yellow Gene in Ponerine Ant, Diacamma Sp', *PLOS ONE*, 9(3), p. e92875.
Available at: https://doi.org/10.1371/journal.pone.0092875.

Nadeau, N. J. *et al.* (2016) 'The gene *cortex* controls mimicry and crypsis in butterflies and
moths', *Nature*, 534(7605), pp. 106–110. doi: 10.1038/nature17961.

Nishikawa, H. *et al.* (2015) 'A genetic mechanism for female-limited Batesian mimicry in
Papilio butterfly', *Nature Genetics*, 47(4), pp. 405–409. doi: 10.1038/ng.3241.

Nokelainen, O. *et al.* (2012) 'Trade-off between warning signal efficacy and mating success
in the wood tiger moth', *Proceedings of the Royal Society B: Biological Sciences*, 279(1727),
pp. 257–265. doi: 10.1098/rspb.2011.0880.

Nokelainen, O. *et al.* (2014) 'Changes in predator community structure shifts the efficacy of
two warning signals in Arctiid moths', *Journal of Animal Ecology*, 83(3), pp. 598–605. doi:
10.1111/1365-2656.12169.

Nokelainen, O. *et al.* (2022) 'Genetic colour variation visible for predators and conspecifics
is concealed from humans in a polymorphic moth', *Journal of Evolutionary Biology*,
(February), pp. 467–478. doi: 10.1111/jeb.13994.

Orteu, A. and Jiggins, C. D. (2020) 'The genomics of coloration provides insights into
adaptive evolution', *Nature Reviews Genetics*, 21(8), pp. 461–475. doi: 10.1038/s41576-020-

0234-z.

Polidori, C., Jorge, A. and Ornosa, C. (2017) 'Eumelanin and pheomelanin are predominant
pigments in bumblebee (Apidae: Bombus) pubescence', *PeerJ*, 2017(5), pp. 1–21. doi:
10.7717/peerj.3300.

Quinlan, A. R. and Hall, I. M. (2010) 'BEDTools: a flexible suite of utilities for comparing
genomic features', *Bioinformatics*, 26(6), pp. 841–842. doi: 10.1093/bioinformatics/btq033.

Rastas, P. (2017) 'Lep-MAP3: robust linkage mapping even for low-coverage whole genome
sequencing data', *Bioinformatics*, 33, pp. 3726–3732. doi: 10.1093/bioinformatics/btx494.

Reed, R. D. *et al.* (2011) '*optix* drives the repeated convergent evolution of butterfly wing
pattern mimicry', *Science*, 333, pp. 1137–1141. doi: 10.1126/science.1208227.

Ritchie, M. E. *et al.* (2015) 'limma powers differential expression analyses for RNA-
sequencing and microarray studies', *Nucleic Acids Research*, 43(7), pp. e47–e47. doi:
10.1093/nar/gkv007.

Rojas, B. *et al.* (2017) 'How to fight multiple enemies: Target-specific chemical defences in
an aposematic moth', *Proceedings of the Royal Society B: Biological Sciences*, 284(1863).
doi: 10.1098/rspb.2017.1424.

Rojas, B., Gordon, S. P. and Mappes, J. (2015) 'Frequency-dependent flight activity in the
colour polymorphic wood tiger moth', *Current Zoology*, 61(4), pp. 765–772. doi:
10.1093/czoolo/61.4.765.

Rönkä, K. *et al.* (2020) 'Geographic mosaic of selection by avian predators on hindwing
warning colour in a polymorphic aposematic moth', *Ecology Letters*, 23(11), pp. 1654–1663.
doi: https://doi.org/10.1111/ele.13597.

Shamim, G. *et al.* (2014) 'Biochemistry and biosynthesis of insect pigments', *European
Journal of Entomology*, 111(2), pp. 149–164. doi: 10.14411/eje.2014.021.

Sugumaran, M. (2010) 'Chapter 5 - Chemistry of Cuticular Sclerotization', in Simpson, S. J.
B. T.-A. in I. P. (ed.). Academic Press, pp. 151–209. doi: https://doi.org/10.1016/B978-0-12-
381387-9.00005-1.

Suomalainen, E. (1938) 'Die erblichkeitsverhältnisse des männlichen dimorphismus bei
parasemia plantaginis', *Hereditas*, 24, pp. 386–390.

Thompson, M. J. and Jiggins, C. D. (2014) 'Supergenes and their role in evolution', *Heredity*,
113(1), pp. 1–8. doi: 10.1038/hdy.2014.20.

Timmermans, M. J. T. N. *et al.* (2014) 'Comparative genomics of the mimicry switch in
Papilio dardanus', *Proceedings of the Royal Society B: Biological Sciences*, 281(1787). doi:
10.1098/rspb.2014.0465.

Tuttle, E. M. *et al.* (2016) 'Divergence and Functional Degradation of a Sex Chromosome-
like Supergene', *Current Biology*, 26(3), pp. 344–350. doi:
https://doi.org/10.1016/j.cub.2015.11.069.

Umebachi, Y. (1975) 'Further studies on the dopamine derivative, SN-1 derived from the
yellow pigments of Papilio xuthus', *Insect Biochemistry*, 5, pp. 73–92.

Wakamatsu, K. *et al.* (2014) 'Reduction of the Nitro Group to Amine by Hydroiodic Acid to
Synthesize o-Aminophenol Derivatives as Putative Degradative Markers of Neuromelanin',
Molecules . doi: 10.3390/molecules19068039.

Wakamatsu, K., Ito, S. and Rees, J. L. (2002) 'The Usefulness of 4-Amino-3-
hydroxyphenylalanine as a Specific Marker of Pheomelanin', *Pigment Cell Research*, 15(3),
pp. 225–232. doi: https://doi.org/10.1034/j.1600-0749.2002.02009.x.

Wang, J. *et al.* (2013) 'A Y-like social chromosome causes alternative colony organization in
fire ants', *Nature*, 493(7434), pp. 664–668. doi: 10.1038/nature11832.

Wang, P. *et al.* (2022) 'Yellow-b, -c, -e and -h are required for normal body coloration in
Henosepilachna vigintioctopunctata', *Archives of Insect Biochemistry and Physiology*,
109(2), p. e21856. doi: https://doi.org/10.1002/arch.21856.

Wilson, R. *et al.* (1976) 'Behavioural pleiotropy of the yellow gene in Drosophila
melanogaster', *Genetical Research*. 2009/04/14, 28(1), pp. 75–88. doi: DOI:
10.1017/S0016672300016748.

Wittkopp, P. J., Vaccaro, K. and Carroll, S. B. (2002) 'Evolution of yellow gene regulation
and pigmentation in Drosophila', *Current Biology*, 12(18), pp. 1547–1556. doi:
10.1016/S0960-9822(02)01113-2.

Yen, E. C. *et al.* (2020) 'A haplotype-resolved, de novo genome assembly for the wood tiger
moth (Arctia plantaginis) through trio binning', *GigaScience*, 9(8), pp. 1–12. doi:
10.1093/gigascience/giaa088.

Zhang, L. *et al.* (2017) 'Genetic basis of melanin pigmentation in butterfly wings', *Genetics*,
205(4), pp. 1537–1550. doi: 10.1534/genetics.116.196451.

Zhang, L., Mazo-Vargas, A. and Reed, R. D. (2017) 'Single master regulatory gene
coordinates the evolution and development of butterfly color and iridescence', *Proceedings of
the National Academy of Sciences*, 114, pp. 10707–10712. doi: 10.1073/pnas.1709058114.