Colour polymorphism associated with a gene duplication in male wood tiger moths

Melanie N. Brien[1][*][†], Anna Orteu[2][*], Eugenie C. Yen[2], Juan A. Galarza[1,3], Jimi Kirvesoja[3],

Hannu Pakkanen[4], Kazumasa Wakamatsu[5], Chris D. Jiggins[2][**], Johanna Mappes[1,3][**].

[1]*Organismal and Evolutionary Biology Research Program, Faculty of Biological and*
Environmental Sciences, University of Helsinki, Finland.

[2]*Department of Zoology, University of Cambridge, United Kingdom.*

[3]*Department of Biological and Environmental Science, University of Jyväskylä, Finland.*

[4]*Department of Chemistry, University of Jyväskylä, Finland.*

[5]*Institute for Melanin Chemistry, Fujita Health University, Toyoake, Japan.*

*Co-first authors.

**Co-senior authors.

[†]Correspondence: mnbrien1@gmail.com

Abstract

Colour is often used as an aposematic warning signal, with predator learning expected to lead

to a single colour pattern within a population. However, there are many puzzling cases where

aposematic signals are also polymorphic. The wood tiger moth, *Arctia plantaginis*, displays

bright hindwing colours associated with unpalatability, and males have discrete colour

morphs which vary in frequency between localities. In Finland, both white and yellow

morphs can be found, and these colour morphs also differ in behavioural and life-history

traits. Here, we show that male colour is linked to an extra copy of a *yellow* family gene that

is only present in the white morphs. This white-specific duplication, which we name *valkea,*

is highly upregulated during wing development, and CRISPR knockouts validate the role of

valkea in producing white wing colour. We also characterise the pigments responsible for

yellow, white and black colouration, showing that yellow is partly produced by

pheomelanins, while black is dopamine-derived eumelanin. Our results add to a growing

number of studies on the genetic architecture of complex and seemingly paradoxical

polymorphisms, and the role of gene duplications and structural variation in adaptive

evolution.

Introduction

Colour polymorphisms, defined as the presence of multiple discrete colour phenotypes within a population (Huxley, 1955), provide an ideal trait to study natural and sexual selection. Colour phenotypes can have an effect on fitness in many contexts including camouflage, mimicry and mating success. Colour is often associated with aposematism, where it acts as a signal, warning predators of unpalatability (Cott, 1940; Cuthill *et al.*, 2017). In such cases, predator learning should favour the most common colour pattern, leading to positive frequency dependent selection (Endler, 1988). Despite this, aposematic polymorphisms can be stable when selection is context-dependent (Briolat *et al.*, 2019), especially where genetic correlations between colour phenotypes and other traits lead to complex fitness landscapes (reviewed by McKinnon and Pierotti, 2010).

A variety of genetic mechanisms can underpin these types of complex polymorphisms involving multiple associated traits (Orteu and Jiggins, 2020). In many cases, such complex polymorphisms are controlled by 'supergenes' in which divergent alleles at several linked genes are maintained in strong linkage disequilibrium by reduced recombination. The most common mechanism for locally reduced recombination are inversions, which range from single inversions involving a small number of genes, to multiple nested inversions covering large genomic regions (Joron *et al.*, 2011; Wang *et al.*, 2013; Küpper *et al.*, 2016; Funk *et al.*, 2021). Nonetheless, other mechanisms for reducing recombination, such as centromeres or large genomic deletions, may also play a role. An alternative mechanism is that a single regulatory gene controls variation via multiple downstream effects (Thompson and Jiggins, 2014). While there are fewer instances in which multiple phenotypes seem to be controlled by a single gene, one potential example is the common wall lizard, where colour genes have pleiotropic effects on behavioural and reproductive traits (Andrade *et al.*, 2019). Multiple mutations within a single gene can also lead to variation in multiple traits (Linnen *et al.*, 2013).

The wood tiger moth, *Arctia plantaginis,* has a complex polymorphism that has been well studied in an ecological context. Males show polymorphic aposematic hindwing colouration with discrete yellow, white or red hindwing colour morphs found at varying frequencies in different geographic locations. In Finland, for example, both yellow and white morphs can be found, with white morphs varying in frequency from 40 to 75% (Galarza *et al.*, 2014). In Estonia, white morphs make up 97% of the population, while yellows morphs form a

completely monomorphic population in Scotland (Hegna, Galarza and Mappes, 2015) (Figure

1). Long term breeding studies of these moths have shown that male hindwing colour is a

Mendelian trait controlled by a single locus with two alleles (Suomalainen, 1938; Nokelainen

et al., 2022). White alleles (W) are dominant over the yellow (y). These colour genotypes

also covary with behavioural and life-history traits, contributing to the maintenance of this

polymorphism. Yellow males are subject to lower levels of predation in the wild (Nokelainen

et al., 2012, 2014), while white males have a positive frequency-dependent mating advantage

(Gordon *et al.*, 2015). There are differences in chemical defences, and bird reactions to these

defences, between the genotypes (Rojas *et al.*, 2017; Winters *et al.*, 2021). Yellow morphs

show reduced flight activity compared to white males, although yellows may fly at more

selective times, i.e. at peak female calling periods (Rojas, Gordon and Mappes, 2015).

Increased reproductive success in Wy genotype females points towards strong heterozygote

advantage (De Pasqual *et al.*, 2022). In summary, there is a trade-off between natural

selection through predation and reproductive success, which contributes to the maintenance

of this polymorphism (Rönkä *et al.*, 2020).

Despite the large body of research on *A. plantaginis* colour morphs, the genetic basis of this

polymorphism is unknown. Here, we explore male hindwing colour variation using linkage

mapping, whole genome data and gene expression analyses, with wild populations and lab

crosses, to identify the locus controlling the colour polymorphism. We use CRISPR/Cas9

gene knockouts to determine the function of the identified gene, and then characterise the

pigments producing yellow, white, and black colouration on the wings of male *A. plantaginis*.

Our findings aim to provide an example of the genetic architecture controlling a trait that is

part of a complex polymorphism.

Figure 1: (A) Sampling locations and frequencies of yellow and white *Arctia plantaginis* males in Finland, Scotland and Estonia. (B) Males of the white and yellow colour morphs (Credit: Samuel Waldron).

Results

A narrow genomic region is associated with hindwing colour

To investigate the genetic basis of male hindwing colouration in *A. plantaginis*, we carried out a quantitative trait locus (QTL) mapping analysis using crosses between heterozygous Wy males and homozygous yy females. We used RADseq data aligned to the yellow *A. plantaginis* reference genome from 172 male offspring (90 white and 82 yellow) from four families. The QTL analysis identified a single marker associated with male hindwing colour (Figure 2A). This marker was found on scaffold YY_tarseq_206_arrow at position 9,887,968bp (95% confidence intervals 9,349,978-9,888,009bp) and had a LOD score of 32.8 (p<0.001). The significant marker explains around 75% of the phenotypic variation and, with one exception, yellow individuals all had a homozygous yy genotype at this marker.

To further narrow down this region, we ran a genome wide association study (GWAS) using whole genome sequences of males from four populations: polymorphic Southern Finland (5 white, 5 yellow) and Central Finland (10 white, 10 yellow), Estonia (4 white), where males are mostly white, and monomorphic Scotland (12 yellow), where all males are yellow. This identified a region of associated SNPs also on scaffold 206 (Figure 2B). Two SNPs, 137bp apart (at positions 9,885,384 and 9,885,521), were significant above a strict Bonferroni

corrected threshold. 162 SNPs were over the threshold of p<0.0001 and, of these, 155 are

within a 99kb region on scaffold 206 (9,833,387-9,932,264bp). The top SNPs are within

2.5kb from the top QTL marker, and the SNP at this marker has a p-value <0.0001.

The 538Kb QTL interval contains 21 genes (Table S1) which were annotated with reference

to *D. melanogaster*. Of these genes, four are part of the *yellow* gene family. The top two

SNPs from the GWAS, and the top marker from the QTL, fall in a non-coding region

upstream of the gene, *yellow-e*, and are also close to an additional *yellow* gene, *yellow-g*.

Identifying structural variation in this region

The trio binning method used by Yen et al. (2020) to assemble the *A. plantaginis* reference

genome produced two reference sequences, one for a white allele and one for a yellow allele.

We extracted the region containing the QTL interval from the yellow reference and aligned it

against the white reference. The alignment showed a duplicated region approximately 117kb

long on scaffold 419 of the white reference from around 6,941,000-7,058,000bp (Figure 2 -

figure supplement 1). The *yellow-e* gene and its flanking regions are within this sequence and

are therefore duplicated in the white reference (Figure 2C). One copy of the gene (named

jg1310 in the W annotation) has 7 exons and is similar to the *yellow-e* gene in the yellow

reference (99.7% identity in coding sequences). The second copy unique to the white scaffold

(jg1308) has only the first 5 exons (81.8% identical to the gene in the yellow reference),

possibly due to a stop codon mutation in the 5th exon. For clarity, we named this duplicated

white-specific copy *valkea*, in reference to a Finnish word for 'white'. While all white

samples had consistent coverage of reads across the duplicated region, coverage was patchy

in yellow samples, with many regions having no and very low coverage in yellow samples

(Figure 2D). Those reads that map in the *valkea* region in yellow samples are likely to be

mapping errors, because the sequence similarity is high and mapping quality is reduced

within the duplication (Figure 2 – figure supplement 2). When increasing the mapping quality

filtering, read depth decreases more in yellow samples compared to white samples in this

region (Figure 2 – figure supplement 3). We confirmed the absence of this region in yellow

individuals by designing primers within the duplication (Table S2, Appendix Figure 1),

which only amplified in WW and Wy samples, including Finnish, Estonian and lab

populations.

To confirm that both of these gene copies are related to *yellow-e*, we compared them to

yellow-e orthologs found in *Bombyx mori*, *Heliconius melpomene* and *Drosophila*

melanogaster, along with other *yellow* genes from *A. plantaginis* and *B. mori*. Both of the

tiger moth genes were most closely related to the *H. melpomene yellow-e* (Figure 2 – figure

supplement 4). Between *valkea* and *yellow-e* there is an additional gene which showed

highest similarity to *Drosophila yellow-g2* (when extracted from both the white and yellow

references). This gene is not part of the duplicated sequence and is present as a single copy in

both morphs. Coverage across *yellow-g* and *yellow-e* genome regions in wild samples is

similar in both morphs (Figure 2). Upstream of the duplication is an unnamed gene (listed as

jg6744 in the yellow annotation and jg1307 in the white). This is the same orthologous gene

in both reference genomes, having 99.3% identity. Similarly, if we look at the 150kb

upstream region, sequence identity is 99.98%. There are no non-synonymous mutations

between coding sequences of *yellow-e* when comparing the white and yellow references,

although there are differences in the first exon of *yellow-g*. The absence of this duplicated

region in the yellow morphs means we cannot determine if there is a change in linkage

disequilibrium across white and yellow morphs.

Figure 2: A duplicated region in white morphs is associated with male hindwing colour. (A) QTL
analysis of white and yellow F1 males (n=172) reveals a 500kb region significant on scaffold 206,
part of linkage group 9. The dotted line indicates the significance threshold determined by
permutation tests (p=0.05). (B) A GWAS of wild samples (n=46) showed SNPs associated with
hindwing colour along the same scaffold. The dotted line shows the Bonferroni corrected significance
threshold. (C) An insertion in the white reference sequence contains a copy of the *yellow-e* gene,
which we named *valkea*, in addition to the *yellow-e* present in both white and yellow morphs. (D)
Mean read depth across the candidate region in all Finnish white (WW and Wy) and yellow (yy)
samples.

 Valkea is differentially expressed between morphs

To pinpoint which of these candidate genes is associated with male wing polymorphism in *A.*

plantaginis, we next performed gene expression analyses across several developmental

stages. Based on knowledge of the expression patterns of *yellow* genes (Ferguson *et al.*,

2011) and other melanin pathway genes such as *pale, ebony and ddc* in Lepidoptera (Zhang

et al., 2017), we hypothesised that changes in gene regulation that control the development of

wing colour morphs in the tiger moth most likely occur during pupal development. Pupal

development in the wood tiger moth lasts for approximately 8 days, and no colour is present

in the wings until day 7, when the yellow pigment appears. A few hours later, black melanin

pigmentation is deposited. We sampled two stages early in development when no colouration

is present in the wings (72 hours post-pupation, and 5 day old pupae), and two stages later in

development: the point when yellow appears in yy morphs (Pre-mel, 7 day old pupae) and the

other after black melanin has also been deposited (Mel, 7-8 days old). Forty individuals were

sampled in total - five per genotype and stage.

First, we explored the general patterns of expression by mapping RNAseq reads to the white

reference genome, which contains the duplicated region that includes *valkea*. We filtered out

lowly expressed genes, retaining 10,920 genes and used Multidimensional Scaling (MDS), a

dimensionality reduction technique, to explore which factors explain genome-wide variation

in gene expression between samples. We observed that samples clustered based on their

developmental stage, suggesting it is an important factor driving differences in genome-wide

gene expression between samples (Figure 3 – figure supplement 1). Such a pattern would be

expected as many genes are involved in development and thus are likely to be differentially

expressed (DE) between developmental stages. No apparent clustering can be observed

among samples of the same colour morph.

We next compared gene expression between yy and WW individuals at each of the

developmental stages. Overall, 99 genes were differentially expressed (FDR < 0.05) between

the two morphs (Figure 3A). Two of these DE genes, *yellow-e* and *valkea*, are two of the 22

genes identified in the GWAS and QTL analysis. *Valkea* was overexpressed in white

individuals in the pre-melanin stage with a Log Fold Change of 10.32 and a p-value of 2.18e-

06. As *valkea* is only fully present in the W genome, it is not expected to be expressed at all

in the Y genome. *Yellow-e* was also overexpressed in white individuals during the pre-

melanin stage with a Log Fold Change of 3.86 and adjusted p-value of 5.62e-06. In other

developmental stages, neither *valkea* nor *yellow-e* showed differences in expression between

morphs (Figure 3B).

Of the 99 genes differentially expressed between white and yellow individuals across

development, 49 were upregulated in the yy morph, while the remaining 50 were upregulated

in WW individuals. The earliest developmental stage, 72-hours, was the stage with the

highest number of DE genes (n=48), while the 5-day old stage had the fewest (n=7). One

gene which encodes a C2H2 zinc finger transcription factor in *D. melanogaster*, '*jg15945*',

was over-expressed in yy in the first three stages (Figure 3 – figure supplement 2).

Finally, the GWAS and QTL peaks of association are situated in scaffold 419 of the WW

reference assembly, which in a WW linkage map forms a linkage group along with 6 more

scaffolds (472, 487, 515, 531, 540 and 609). We found that 12 genes present in this linkage

group were differentially expressed, including *valkea* and *yellow-e* (Table S3), and identified

their orthologues in *Drosophila melanogaster*.

Figure 3: *Valkea* is overexpressed in white males in the pre-melanin stage. (A) In pink are genes that
are significantly differentially expressed between yellow and white morphs at the pre-melanin stage.
Valkea is the most DE gene (i.e. gene with the highest Log Fold Change). (B) Expression of *valkea*
across developmental timepoints shows that it has higher expression measured in Log2 CPM (Counts
Per Million) in white individuals compared to yellow ones. Expression of *valkea* in yellow morphs is
around 0.

To confirm the function of *valkea* in wing colouration, we used CRISPR/Cas9 to knock out

the gene in white morphs. We tested 5 different guides to target the first 3 exons of *valkea*

and injected Cas9/sgRNA duplexes into a total of 1223 eggs. Of 143 larvae that hatched only

6 developed to adults (Table S4). However, of the five males that did eclose, four had a

visible change in phenotype. Males produced yellow scales instead of white on the dorsal

side of both the forewings and the hindwings (Figure 4). Forewings were more yellow than in

the wildtype yellow males, which usually have lighter forewings compared to hindwings.

White scales on the ventral side of the wings also became yellow, similar to wildtype

yellows. Black melanin patterning did not seem to be affected. Variation in the amount of

melanin can be attributed to the populations from which the individuals originated, with the

darker samples coming from the Finnish population (Figure 4 – figure supplement 1).

Wildtype white morphs also reflect UV, particularly on the hindwings, but this is not seen in

the CRISPR males (Figure 4 – figure supplement 2). This could suggest a change in scale

structure, or that the yellow pigment covers the UV-reflecting structures.

Four out of the five guides tested produced a mutant phenotype, with no differences in the

male phenotype between guides. The similarity of *valkea* to *yellow-e* means that it is likely

that *yellow-e* could also have been knocked-out in these individuals. We used whole genome

sequences of the mutants to confirm that the correct sites in *valkea* had been targeted (Figure

4 – figure supplement 3). All samples also showed evidence of editing at the corresponding

yellow-e exons, which mainly involved insertions. The change in forewing colour could be

attributed to a *yellow-e* mutation. Only one female survived to adulthood, and this had a

mosaic phenotype. This individual had one yellow forewing, similar to the male mutants,

with the rest of the body and wings being wildtype (Figure 4 – figure supplement 1). Since

we did not expect a *valkea* knockout to affect female phenotypes as they always have red

hindwings, this could be further evidence for the effect of *yellow-e* on forewing colour.

Survival of the eggs varied between the guides, although this was largely affected by the

female, as hatching rate between females ranged from 0 to 70%. Females often lay

unfertilised eggs so we expect that hatching rate will be low in some crosses. Using two

guides in combination did not produce any pupae or adults.

White wt Yellow wt

CRISPR dorsal CRISPR ventral

Figure 4: CRISPR/Cas9 knockouts of *valkea* transforms white scales into yellow scales across both
hindwings and forewings. Wildtype WW and yy morphs (top), and the dorsal and ventral sides of one
of the CRISPR knockout males (bottom).

Pigment analysis

Since the *yellow* gene family, to which *valkea* is related, is known to be responsible for the

production of melanin pigments, we further investigated the identity of the wing

pigmentation. First, we ruled out the presence of several non-melanin pigment types in the

hindwings, including pterins and carotenoids. Pterins are commonly found in insects and,

along with purine derivatives, papiliochromes and flavonoids, are soluble in strong acids and

bases or in organic solvents (Umebachi, 1975; Kayser, 1985; Shamim *et al.*, 2014). We

placed wing samples from each morph in NaOH overnight, then measured the absorbance of

the supernatant using a spectrophotometer. We also left wings in methanol overnight before

measuring the supernatant. The spectra did not show any peaks indicative of any pigment

dissolved in the sample. Similarly, we found no evidence for carotenoid pigments after

dissolving in a hexane:tert-butyl methyl ether solution (Appendix Figure 2). Wings did not

fluoresce under UV light, providing further evidence for the lack of fluorescent pigments

including pterins, flavonoids, flavins and papiliochromes (Umebachi, 1975; Kayser, 1985).

Ommochromes are red and yellow pigments; high performance liquid chromatography

(HPLC) ruled out the presence of these pigments on the moth wings, which we compared to

data from ommochrome-containing *Heliconius* wings and a Xanthurenic acid standard

(Appendix Figure 3).

HPLC analysis showed peaks characteristic of pheomelanin (Appendix Figure 4).

Pheomelanins produce red-brown colour in grasshoppers and wasps (Galván *et al.*, 2015;

Jorge García, Polidori and Nieves-Aldrey, 2016), and orange-red colours in ants and

bumblebees (Hines *et al.*, 2017; Polidori, Jorge and Ornosa, 2017). Insects generally have

dopamine-derived pheomelanin and a breakdown product of this is 4-Amino-3-

hydroxyphenylethylamine (4-AHPEA) (Barek *et al.*, 2018). Yellow wings showed large

peaks for 4-AHPEA. White wings had around 27% of the 4-AHPEA levels seen in yellow

wings, and black sections of the wings had 16%. Hydrogen iodide hydrolysis of wings

produced the isomer 3-AHPEA, which may come from 3-nitrotyramine originating from the

decarboxylation of 3-nitrotyrosine. Reduction of 3-nitrotyrosine produces 3-AHP, another

marker of pheomelanin (Wakamatsu, Ito and Rees, 2002).

Analysis of the black portions of the wing found pyrrole-2,3-dicarboxylic acid (PDCA) and

pyrrole-2,3,5-tricarboxylic acid (PTCA) (Appendix Figure 5). Both are components of

eumelanin (Barek *et al.*, 2018), suggesting that the black colouration seen in the wood tiger

moth is predominantly eumelanin derived from dopamine. This is common in producing

black colouration and providing structural components of the exoskeleton. In addition,

dopamine is acylated to both *N*-β-alanyldopamine (NBAD) and *N*-acetyldopamine (NADA)

sclerotins. NADA sclerotins are colourless and likely to be present on the white wings. This

analysis of pigmentation is therefore consistent with a role for *yellow* family genes in

regulating the colour polymorphism.

Discussion

Hindwing colouration of male *Arctia plantaginis* is polymorphic and these colour morphs

vary in multiple behavioural and life-history traits, providing an example of a complex

polymorphism. Here, we have shown that variation in male hindwing colour is associated

with a duplicated sequence found only in white morphs and containing a gene from the

yellow gene family. The white-specific copy, *valkea,* is highly expressed during pupal

development, consistent with genetic dominance of the white allele. When *valkea* is knocked

out in the white morphs, yellow pigment is produced, providing functional evidence that

valkea is responsible for the white phenotype rather than other genes in this region.

These results add to the increasing evidence for the role of gene duplications in the evolution

of adaptive genetic variation. Genes for the metabolisms of proteins in *Heliconius* butterflies

underwent several duplications, facilitating changes in diet and adaptation to pollen feeding

(Smith, Macias-Muñoz and Briscoe, 2016). In *Zerene cesoina* butterflies, recent partial

duplications of the transcription factor *doublesex,* resulting from multiple duplication events,

are associated with sex-specific wing patterning. The duplicated paralog acts as a repressor of

genes producing UV-reflecting wing scales in females (Rodriguez-Caro *et al.*, 2021).

We hypothesise that the morph-specific duplication that we see in *A. plantaginis* provides a

region of reduced recombination between morphs, as the duplicated region is effectively

hemizygous and cannot recombine except in homozygote genotypes, which could contribute

to the maintenance of the complex polymorphism and the linkage of multiple traits. This is

similar to the genetic architecture of the *Primula* supergene controlling heterostyly, which

involves a large duplication containing five genes (Huu *et al.*, 2020). In polymorphic *Papilio*

dardanus, one colour pattern morph is associated with a duplicated region, again providing

physical constraints on recombination (Timmermans *et al.*, 2014). Nonetheless, in the case of

the wood tiger moth, it remains unclear how a single gene, such as *valkea*, can control the

development of a broad array of phenotypic traits.

One possible mechanism is that there is a regulatory element along the scaffold which is

controlling colour via the *valkea* gene, but also regulating other genes to control different

phenotypic traits. We found the most significant markers and SNPs located in a non-coding

region close to the *yellow* genes, which likely contains a cis-regulatory element (CRE)

controlling transcription of *valkea.* In cichlids, for example, a CRE at the gene encoding

agouti-related peptide 2 controls variation in strip patterning in two closely related species

(Kratochwil *et al.*, 2018). Conserved CREs were shown to have wide ranging effects on wing

patterning in multiple Nymphalidae butterflies (Mazo-Vargas *et al.*, 2022).

Differential expression of other genes on the same chromosome controlled by the CRE could

explain variation in covarying traits. The over-expression of another gene, possibly encoding

a zinc transcription factor, in yellow individuals in the early pupal stages suggests that there

is differential expression of unlinked genes as a result of the polymorphism, although since

this gene is on a different chromosome to *valkea* it is unlikely to be directly controlled by the

CRE. Another hypothesis is that somehow *valkea* itself regulates other genes. However,

yellow family genes are not known to regulate transcription of other genes, unlike for

example *doublesex*, which undergoes alternative splicing and female mimetic wing pattern

polymorphism in *Papilio polytes* (Kunte *et al.*, 2014; Nishikawa *et al.*, 2015).

The *yellow* family genes are highly conserved throughout insects (Ferguson *et al.*, 2011).

They have been widely linked to colouration (Wittkopp, Vaccaro and Carroll, 2002;

Miyazaki *et al.*, 2014; Zhang *et al.*, 2017; Zhang, Mazo-Vargas and Reed, 2017), as well as
behaviour, sex-specific phenotypes and reproductive maturation (Wilson *et al.*, 1976). These
genes share a common origin with the major royal jelly protein (MRJP) genes (Drapeau *et*
al., 2006) which are crucial in caste development in honeybees. Like the MRJP genes, *yellow*
genes in honeybees have diverse spatial and temporal expression patterns. As our focus in *A.*
plantaginis has been on wing tissue, we are missing expression of genes in other tissues that
could be linked to other traits. Thus, it is not impossible to imagine that a *yellow* gene could
have a similar function to a MRJP in regulating the development of a complex phenotype.
Recent work with *Bicyclus anynana* showed that *yellow* functions as a repressor of male
courtship (Connahs *et al.*, 2022). On the other hand, sex-specific behavioural phenotypes of
yellow mutants in *Drosophila* were found to be due to pigmentation effects (Massey *et al.*,
2019), so more evidence is needed to suggest a functional role for *yellow* genes outside of
pigmentation.

The duplication of *yellow-e* and surrounding regions in the white morphs suggests that the
yellow morph is the ancestral form. *Valkea* could have evolved in a stepwise fashion, first as
a tandem duplication then with a stop codon mutation altering the gene structure. Gene
duplications can facilitate adaptation and, in some examples, lead to polymorphism. The fact
that the white allele is dominant also supports the hypothesis that yellow is ancestral. Such
invasions of new adaptive alleles are facilitated when the new allele is dominant, as it is then
also expressed when heterozygous, i.e. the Haldane's sieve effect. Melanism, for example,
has repeatedly evolved in mammals due to dominant and semidominant mutations in the
Mc1r locus which have become fixed (Hoekstra, 2006).

Valkea could represent an example of neofunctionalization, where the duplicated gene gains a
different function to the original gene copy. In the CRISPR mutants, both forewings and
hindwings became yellow, and thus we suspect that *valkea* is controlling hindwing colour
while *yellow-e*, which was likely also knocked out due to the similarity of the sequences,
controls forewing colour. Those with the mutant phenotype showed only small deletions or
insertions around the target site. By combining multiple guides we may expect to see larger
deletions (Mazo-Vargas *et al.*, 2022), however none of the eggs that were injected with more
than one guide survived past the larval stage, suggesting that large deletions in *yellow* genes
reduce fitness.

Contrary to previous work that attributed red and yellow colours to pterins in another tiger
moth species (Gawne and Nijhout, 2019), we found high levels of 4-AHPEA in the yellow
wings confirming the presence of pheomelanins. These pigments have been widely associated
with red and yellow colours in mammals (e.g. Mcgraw and Wakamatsu, 2004), but only
relatively recently described in insects and likely to be more widespread than previously
thought. Yellow colours can also be produced by NBAD sclerotins which are sclerotizing
precursor molecules made from dopamine and these have an important role in the
sclerotization pathway for hardening the insect cuticle (Andersen, 2007; Barek, Evans and
Sugumaran, 2017) before becoming involved in melanisation (Barek *et al.*, 2018). Thus, we
suggest that the yellow colour arises partly from the NBAD sclerotins and partly from the
presence of pheomelanin pigments, which has been proposed in other Lepidoptera (Matsuoka
and Monteiro, 2018). While some 4-AHPEA also occurred in white wings, this may be due to
its role in production of cross-linking cuticular proteins and chitin during sclerotization
(Sugumaran, 2010). Upregulation of genes on the white allele could be acting as a repressor
of the generation of yellow colour. If we may speculate, *valkea* could impact the catalysis of
dopamine, having cascading effects down the pathway resulting in the lack of yellow
pigmentation. We suspect that *yellow* family genes play multiple roles within the melanin
production pathway. In the wood tiger moth, *yellow* affects the conversion of DOPA into
black dopamine melanin (Galarza, 2021). *Yellow-e* in particular has been linked to larval
colouration in *Bombyx mori* (Ito *et al.*, 2010) and adult colour in beetles (Wang *et al.*, 2022),
while another gene, *yellow-f*, has a role in eumelanin production (Barek *et al.*, 2018).

In summary, we identified a structural variant which is only present in white morphs of *A.*
plantaginis. This region contains a previously undescribed gene, *valkea,* which when
knocked out results in yellow wings. The presence of a regulatory element controlling wing
colour and other traits via multiple downstream effects could explain how multiple traits are
linked to wing colouration. This complex polymorphism allows multiple beneficial
phenotypes to be inherited together, whereas recombination would separate multiple loci
leading to maladapted individuals. Our results provide the basis for further exploration of the
genetic basis of covarying behavioural and life-history traits, and offer an intriguing example
for the role of gene duplications in adaptive variation.

Methods

Sampling

Homozygous lines of white (WW) and yellow (yy) *Arctia plantaginis* moths were created
from Finnish populations at the University of Jyväskylä, Finland. Larvae were fed with wild
dandelion (*Taraxacum sp.*) and reared under natural light conditions, with an average day
temperature of 25°C and night temperature between 15-20°C. For the crosses, a heterozygous
male, created from crossing a heterozygous male with a homozygous yy female, was
backcrossed with a yy female. This was repeated to obtain four families totalling 172
offspring and 8 parents (Table S5). Samples from wild populations were caught in Southern
Finland (n=10) and Central Finland (n=20), where male morphs are either white or yellow,
Estonia (n=4), where males are mostly white, and Scotland (n=4), where males are yellow
(Table S6). In addition, we included 8 samples which are F1 offspring of wild Scottish
samples. Forty pupae with known genotypes from lab populations (20 WW and 20 yy) were
used for the RNA extractions.

DNA extraction and sequencing

For the lab crosses, DNA was extracted from two legs crushed with sterilised PVC pestles
using a Qiagen DNeasy Blood & Tissue kit, following the manufacturer's instructions.
Library preparation and GBS sequencing were performed by BGI Genomics on an Illumina
HiSeq X Ten. For the wild samples, DNA was extracted from the thoraces also with a Qiagen
kit. Library preparation and sequencing were performed by Novogene (Hong Kong, China).
150bp paired-end reads were sequenced on an Illumina NovaSeq 6000 platform.

Linkage mapping analysis

FASTQ reads were mapped using bowtie v2.3.2 (Langmead and Salzberg, 2012) to the
yellow *Arctia plantaginis* scaffold-level genome assembly (Yen *et al.*, 2020). BAM files
were sorted and indexed using SAMtools v.1.9 (Li *et al.*, 2009) and duplicates removed using
PicardTools MarkDuplicates (broadinstitute.github.io/picard). Twelve samples which had
aligned <30% were removed. Reads of the remaining samples had an average alignment of
94%. SNPs were called using SAMtools mpileup with minimum mapping quality set to 20
and bcftools call function. Lep-MAP3 (Rastas, 2017) was used for linkage map construction
and we ran the following modules: ParentCall2 which called 105,622 markers, Filtering2,

SeparateChromosomes2 with lodLimit=5 and sizeLimit=100, JoinSingles2All and

OrderMarkers2 with recombination2=0 to denote the lack of female recombination.

Genotypes were phased using the map2genotypes.awk script included with Lep-MAP3.

Markers were named based on the genomic positions of the SNPs in the reference genome

and the map.awk script, and this was used to further order the markers within the linkage

groups. This resolved 30 linkage groups. Although we expect that there are 31 chromosomes

in the moth genome, we suspect that the sex chromosome is missing in this assembly as the

yy individual used in the genome assembly was female (Yen *et al.*, 2020). A small number of

markers which caused long gaps at the beginning or end of linkage groups were manually

removed, leaving the final map 948.7cM long with 19,803 markers. Markers were well

distributed so we began the first analyses with this map. A linkage map was also assembled

using sequences aligned to the white reference and this separated into 31 linkage groups.

The QTL analysis was carried out in R/qtl (Broman *et al.*, 2003). Genotype probabilities were

calculated before running a genome scan using the *scanone* function with the Haley-Knott

method and binary model parameters, and including family as an additive covariate. The

phenotype was labelled as either 0 (Wy) or 1 (yy). We ran 5000 permutations to determine

the significance level for the QTL LOD scores. The *bayesint* function calculated the 95%

Bayesian confidence intervals around the significant marker.

Analysis of whole genome sequences

FASTQ reads were mapped to the yellow *A. plantaginis* genome assembly (Yen *et al.*, 2020)

using BWA-MEM v7.17 (Li, 2013). As before, BAM files were sorted and indexed, and

duplicates were removed. Genotyping and variant calling was carried out with the Genome

Analysis Toolkit (GATK) (McKenna *et al.*, 2010). Variants were called using

HaplotypeCaller (v.3.7) in GVCF mode, then gVCFs combined with GenomicsDBImport

(v.4.0). Joint genotyping was run with GenotypeGVCFs, set with a heterozygosity of 0.01,

and SNPs were called using SelectVariants. Finally, the set of 20,787,772 raw SNPs were

filtered using VariantFiltration and thresholds: quality by depth (QD > 2.0), root mean square

mapping quality (MQ > 50.0), mapping quality rank sum test (MQRankSum > -12.5), read

position rank sum test (ReadPosRankSum > -8.0), Fisher strand bias (FS < 60.0), and strand

odds ratio (SOR < 3.0). A set of 5,227,288 SNPs passed the filtering.

We carried out a Genome Wide Association Study (GWAS) using the R package GenABEL

v.1.8 (Aulchenko *et al.*, 2007). The set of filtered SNPs were converted to BED format with

PLINK2, keeping only biallelic SNPs (cog-genomics.org/plink/2.0/). Sites which were not in

Hardy-Weinberg equilibrium ($p<0.01$), or had a call rate of <0.5, were excluded. Following

this, 381,266 sites were retained across 40 individuals (out of 57). To account for population

stratification, we performed multidimensional scaling on kinship and identity-by-state (IBS)

information estimated from the data, and included this as a covariate in the association test.

Significance levels were calculated using Bonferroni corrected thresholds to account for

multiple testing. Central and Southern Finnish populations were pooled for this analysis,

based on a previous principal component analysis of these samples (Yen *et al.*, 2020).

In Yen et al. (2020), many of these samples were processed in the same way but aligned to

the white genome assembly (based on a white individual). Read depth of the W-mapped

samples was calculated in 1kb windows across the candidate region using BEDtools

(v.2.20.1) multicov (Quinlan and Hall, 2010). For visualisation, lines were smoothed using

LOESS and span=0.01 within ggplot2.

For analysis of structural variants, sequences from the white and yellow genome assemblies

were aligned using MAFFT v7.450 (Katoh and Standley, 2013) and viewed with Geneious.

Our focal sequence, scaffold 419 in the white genome, is the reverse complement of scaffold

206 in the yellow genome.

Identification of candidate genes and tree construction

To identify candidate genes in the QTL interval and GWAS region, we ran a protein

BLASTP v.2.4.0 search to identify *Heliconius melpomene* (Hmel2.5) proteins homologous to

predicted *A. plantaginis* proteins in the region from the genome annotation. Informative gene

names were obtained by performing a BLASTP search with the *H. melpomene* proteins

against all *Drosophila melanogaster* proteins in FlyBase v.FB2020_01 (flybase.org/blast).

For the *yellow* gene tree, we used Lepbase (Challi *et al.*, 2016) to search for *yellow* genes in

Bombyx mori (ASM15162v1). We identified *yellow-e* in *H. melpomene* by searching for

major royal jelly proteins, then comparing protein sequences of these against *Drosophila*

proteins in Flybase. The sequence for *Dmel yellow-e* was downloaded from Flybase. To make

the tree, coding sequences of all genes were aligned in Geneious using MAFFT v7.450

(Katoh and Standley, 2013), then the tree was constructed with PhyML using 10 bootstraps

(Guindon *et al.*, 2010).

Differential gene expression

We dissected the wings out of the pupae in Cambridge, UK. Pupae and larvae were sent to

Cambridge from Jyväskylä and were kept between 22 and 30˚C. Pupae were sexed and only

males were used. Dissections were made at 4 different stages: 72 hours after pupation (72h),

5 days after pupation (5d; counting 0-24 first hours after pupation as day 1), pre melanin

deposition (Pre-mel) and post melanin deposition (Mel). We sampled 5 individuals per stage

and genotype. Hindwings and forewings were stored separately in RNA-later (Sigma-

Aldrich) at 4 ˚C for 2 weeks and later transferred to -20 ˚C, while the rest of the body was

stored in pure ethanol. Only hindwings were used for RNAseq analysis.

Total RNA was extracted from hindwing tissue using a standard hybrid protocol. First, we

transferred the wing tissue into Trizol Reagent (Invitrogen) and homogenised it using dounce

tissue grinders (Sigma-Aldrich). Then, we performed a chloroform phase extraction, followed

by DNase treatment (Ambion) for 30 mins at 37˚C. We measured the concentration of total

RNA using Qubit Fluorometric Quantitation (Thermofisher) and performed a quality check

using an Agilent 4200 TapeStation (Agilent). The extracted total RNA was stored at -20˚C

before being sent to Novogene UK for sequencing. Each individual was sequenced

separately, with a total of 40 individual samples sequenced (5 individuals per stage and

genotype).

We performed quality control and low-quality base and adapter trimming of the sequence

data using *TrimGalore!* We then mapped the trimmed reads to the two *A. plantaginis*

genomes using STAR (Dobin *et al.*, 2013). We performed a second round of mapping (2pass)

including as input the output splice junctions from the first round. The *A. plantaginis* genome

annotations WW and YY were included in each round of mapping respectively. We then used

FeatureCounts to count the mapped reads. Finally, we used DESeq2 to analyse the counts

and perform the DE analysis.

To identify the gene or genes controlling the development of wing colour in *A. plantaginis*,

we performed a genome wide differential expression analysis using limma-voom (Ritchie *et*

al., 2015). First, we defined a categorical variable, 'GenStage', with 8 levels containing the

genotype and stage information of every individual sample (e.g. YY72h, WW72h, YY5days,

etc.). Then, we built the design matrix fitting a model with GenStage as the only fixed effect

factor contributing to the variance in gene expression and included family as a random effect

factor (gene expression ~ 0 + GenStage + (1|Family)). We then filtered lowly expressed

genes using the filterByExpr function in limma, which resulted in a reduction of the number

of tested genes from 17,615 to 11,330 genes in the Y-mapped analysis and 17,930 to 10,920

in the W-mapped one. Then, we normalised the expression of the genes using the

calcNormFactors function with TMM normalisation in limma and fit the design matrix using

the voom function. We built a contrast matrix including the comparisons of interest, in which

we compared the expression of the genotypes in each stage (i.e. h72=WW72h-YY72h,

d5=WWd5-YYd5, Premel=WWPremel-YYPremel, Mel=WWMel-YYMel), and fit the

contrast matrix to the data using the contrasts.fit function. Finally, we used the eBayer

function on the fit dataset and we extracted the list of genes that are differentially expressed

in each stage using the Benjamini-Hochberg procedure to correct for multiple testing. We

evaluated the genome wide gene expression using multidimensional scaling (MDS) using the

plotMDS function of the limma package.

Orthology assignment

To infer genome wide orthology between *A. plantaginis* and *D. melanogaster*, we used

OrthoFinder (v2.5.4) (Emms and Kelly, 2019). We used proteomes from six Lepidoptera

species, *Plutela xylostella* (GCA_905116875_2)*, Bombyx mori* (GCF_014905235_1)*,*

Spodoptera frugiperda (GCF_011064685_1)*, Parnassius apollo* (GCA_907164705_1)*,*

Pieris macdunnoughi (GCA_905332375_1) and *Pararge aegeria* (GCF_905163445_1) and

Drosophila melanogaster (GCF_000001215_4)*.* We ran the primary_transcript.py utility

from OrthoFinder to extract only one transcript per protein, and then ran OrthoFinder with

default settings.

CRISPR/Cas9 genome editing

Guide RNAs were designed within the first 3 exons of *valkea* in the white genome annotation

using Geneious (v. 2022.1.1). Guides were chosen with minimal off-target effects, high

activity scores and high specificity scores based on the Geneious algorithm (Table S7).

Guides in the first two exons of *valkea* showed off-target sites in *yellow-e*, however those in

exon 3 showed no off-target sites. Guide RNAs were synthesised by Sigma-Aldrich. Moths

from the greenhouse populations, originating from Finnish and Estonian populations, at the

University of Helsinki, Finland, were genotyped using DNA extracted from leg tissue using

the Chemagic DNA tissue kit (Chemagen) and the primers detailed below. They were paired

and left to mate overnight. Females were watched over the next 3-4 days for egg laying, and

the eggs were removed and injected less than 6 hours after laying. Eggs were glued to

microscope slides and injected with a 1:1 sgRNA/Cas9 mix with phenol red dye using pulled

borosilicate glass capillaries. The injection mix contained 1ug/ul Cas9, 500ng/ul sgRNA and

0.5% phenol red. Guides and Cas9 were diluted using low concentration TE buffer. Different

combinations of guides were also injected in some samples, in which case these were mixed

in a 1:1 ratio. Injections were performed using a MPPI-3 pressure injector with back pressure

unit (ASI). In total, we injected 1223 eggs from 18 [WW x WW] or [WW x Wy] crosses.

Larvae were kept individually in petri dishes and fed daily with dandelion leaves. After

eclosion, legs were taken from adults for DNA extraction. Library preparation and whole

genome sequencing (using Illumina NovaSeq 6000) of 5 CRISPR mutants was performed by

CeGaT (Tübingen, Germany).

Genotyping white and yellow alleles

We used Primer3 to design primers within the duplicated region. Primers were expected to

only amplify in WW and Wy individuals. The alignment of the white and yellow sequences

was then used to design primers for genotyping the locus (Table S2). We looked for short

insertions or deletions that were fixed between the WW and yy within the *valkea/yellow-e*

region, and put primers around these structural variants. Primers were tested on DNA

extractions from moths of known genotypes, including both sexes, wild and lab samples. We

used Sanger sequencing of the PCR product to confirm the correct sequences were

amplified. A set of primers successfully amplified a 449bp region downstream of *valkea*

within the duplication. This amplified in WW and Wy samples, but not in yy (Appendix

Figure 1). We found that white alleles have a 35bp deletion within an intron of the *yellow-e*

gene. We amplified a 163bp region around this (YY_tarseq_206_arrow:9,846,212-9,846,375)

using a standard PCR protocol which allowed us to identify the allele based on the size of the

PCR product. Yellow alleles produce the full 163bp sequence, while white alleles produce a

smaller 128bp product (Appendix Figure 1). Heterozygotes have a copy of each and show

both bands on a gel.

Pigment analysis

Solubility and fluorescence tests

Five hindwings from each morph were placed in two separate solvents (0.1M NaOH and 90%

MeOH) and left overnight. The supernatant was analysed with an Agilent Cary 8454 UV-

Visible spectrophotometer and the spectra compared to known spectra for various pigments.

A UV lamp (Philips TL8W/08 F8T5/BLB) was used to test for fluorescence on the wings.

The presence of carotenoids was tested by placing wings into 1ml of pyridine and leaving at

95°C for 4 hours (McGraw *et al.*, 2002). To these we added 1ml of 1:1 hexane:tert-butyl

methyl ether and 2ml of water before shaking and leaving overnight. Again, the supernatant

was measured with the spectrometer.

High-Performance Liquid Chromatography (HPLC) to test to for eumelanin and

ommochrome pigments

To determine the type of melanin producing the black colour on the wings, we cut out

approximately 5mg of the black sections of the wings, from both females and males.

Eumelanin analysis was carried out according to (Borges *et al.*, 2001). Each sample was

added to a tube containing 820μl 0.5 M NaOH, 80μl 3% H_2O_2 and an internal standard

(48nmol phthalic acid) and heated in a boiling water bath for 20 minutes. Once cool, 20μl of

10% Na_2SO_3 and 250μl of 6M HCl were added. Samples were then extracted twice with 7ml

of ethyl acetate. Ethyl acetate was dried at 45 °C under a stream of nitrogen. The residue was

dissolved into 0.5ml of 0.1% formic acid.

We carried out high-performance liquid chromatography on an Agilent 1100 HPLC. 20μl of

the sample was injected into a Waters Atlantis T3, 100 x 3.0mm i.d. analytical column

(Waters, Milford, MA, USA). The column was set to 25°C and analytes were detected at

wavelength 280nm. The HPLC mobile phase consisted of two eluents: UHQ-water/MeOH

(98:2; v/v) with 0.1% formic acid and UHQ-water/MeOH (40:60; v/v) with 0.1% formic acid.

Flow rate was 0.4 ml/min and the used gradient started with 100% of eluent A and ramped

evenly from time 0 to 15 minutes to 40:60 (A:B; v/v), held at 40:60 for 6 minutes and ramped

evenly back to initial eluent composition (100% A) over 5 minutes. We compared

chromatograms obtained from the samples to the chromatograms obtained from synthetic

melanin, ink from sepia officinalis and black human hair.

HPLC was also applied to observe the possible presence of ommochrome pigments. Injection volume was 10μl and for the separation we used the same Waters Atlantis T3 column (100 x 3.0 mm i.d.) set to 30°C. Solvent A was UHQ-water and B was acetonitrile (ACN), both containing 0.1% formic acid. Flow rate was 0.4ml/min and the used gradient was as follows: Initial flow ratio was 98/2 water/ACN (v/v) ramping then evenly from time 1 to 15 min to 30:70 water:ACN (v/v), held for 1.5 min and then ramped evenly back to initial eluent composition over 0.5 minutes. The column was stabilised for 7 minutes before a new run.

Pheomelanin analysis

Samples were analysed for pheomelanin content according to the method of Kolb *et al.* (1997) with modifications. A 2mg sample was placed in a screw-capped tube with 100μl water, 500μl ~55-58% hydrogen iodide (HI), and 20μl 50% hypophosphorous acid (H_3PO_2). Samples were capped tightly and hydrolysed for 20 hours at 130°. After cooling, samples were evaporated under nitrogen flow, then dissolved in 1ml of 0.1M HCl and purified with solid phase extraction. Strata SCX cartridges were preconditioned with 2ml of methanol, 3ml of water and 1ml of 0.1M HCl. Sample was then applied to the cartridge, washed with 1ml of 0.1M HCl, and finally eluted with 1ml of methanol (MeOH): 0.5M ammonium acetate ($NH_4CH_3CO_2$) (20:80 v/v).

Hydrogen Iodide hydrolysis products were determined by a Dionex high-performance liquid chromatograph equipped with pulsed amperometric detection (HPLC/PAD). A Phenomenex Kinetex C18 column (150 x 4.6 mm i.d.; 5μm particle size) with a gradient elution (Table S8) at a flow rate of 0.9 ml min[-1] with the eluents: (A) sodium citrate buffer (Hines *et al.*, 2017) in ultra-high-quality water (internal resistance $\geq$ 18.2 MΩ cm; Milli-Q Plus; Millipore, Bedford, MA, USA) and (B) methanol were used for the separation. Dionex ED-50 pulsed amperometric detector (Dionex, Sunnyvale, CA, USA) equipped with a disposable working electrode by using a Dionex waveform A with potentials presented in Table S9 was used for detection. The preparation method for the 4-AHP, 3-AHPEA and 4-AHPEA standards used in calibration is described in (Wakamatsu *et al.*, 2014).

Acknowledgements

We thank Alma Oksanen, Kaisa Suisto and the greenhouse staff for insect rearing, and Elisa Salmivirta and Sari Viinikainen for lab assistance. Thanks to Emeritus Prof. Shosuke Ito for kindly providing pheomelanin standards, and to Bodo Wilts for advice on the pigment

647 analyses. We thank Muktai Kuwalekar, Claudius Kratochwil, Rachel Blow, Ian Warren, Tom

648 Generalovic and Joe Hanly for providing equipment and advice regarding the CRISPR

649 injections.

Funding

651 This work was supported by Academy of Finland grants to MB (#343356) and JM (projects

652 345091 and 328474).

Data availability

654 Scripts and data for the QTL, GWAS and DE analyses can be found at

655 doi:10.5281/zenodo.6560628. RADseq, RNAseq data, and WGS of CRISPR samples were

656 deposited to SRA under study accession number PRJNA937225. Raw sequencing data of

657 wild samples has previously been deposited in ENA, study accession No. PRJEB36595.

References

Andersen, S. O. (2007) 'Involvement of tyrosine residues, N-terminal amino acids, and β-
alanine in insect cuticular sclerotization', *Insect Biochemistry and Molecular Biology*, 37(9),
pp. 969–974. doi: https://doi.org/10.1016/j.ibmb.2007.05.002.

Andrade, P. *et al.* (2019) 'Regulatory changes in pterin and carotenoid genes underlie
balanced color polymorphisms in the wall lizard', *Proceedings of the National Academy of*
Sciences of the United States of America, 116(12), pp. 5633–5642. doi:
10.1073/pnas.1820320116.

Aulchenko, Y. S. *et al.* (2007) 'GenABEL: an R library for genome-wide association
analysis', *Bioinformatics*, 23(10), pp. 1294–1296. doi: 10.1093/bioinformatics/btm108.

Barek, H. *et al.* (2018) 'Insect cuticular melanins are distinctly different from those of
mammalian epidermal melanins', *Pigment Cell and Melanoma Research*, 31(3), pp. 384–392.
doi: 10.1111/pcmr.12672.

Barek, H., Evans, J. and Sugumaran, M. (2017) 'Unraveling complex molecular
transformations of N-β-alanyldopamine that account for brown coloration of insect cuticle',
Rapid Communications in Mass Spectrometry, 31(16), pp. 1363–1373. doi:
https://doi.org/10.1002/rcm.7914.

Borges, C. R. *et al.* (2001) 'Relationship of Melanin Degradation Products to Actual Melanin
Content: Application to Human Hair', *Analytical Biochemistry*, 290(1), pp. 116–125. doi:
https://doi.org/10.1006/abio.2000.4976.

Briolat, E. S. *et al.* (2019) 'Diversity in warning coloration: selective paradox or the norm?', *Biological Reviews*, 94(2), pp. 388–414. doi: 10.1111/brv.12460.

Broman, K. W. *et al.* (2003) 'R/qtl: QTL mapping in experimental crosses', *Bioinformatics*, 19(7), pp. 889–890. doi: 10.1093/bioinformatics/btg112.

Challi, R. J. *et al.* (2016) 'Lepbase: the Lepidopteran genome database', *bioRxiv*, p. 56994. doi: 10.1101/056994.

Connahs, H. *et al.* (2022) 'The yellow gene regulates behavioural plasticity by repressing male courtship in Bicyclus anynana butterflies', *Proceedings of the Royal Society B: Biological Sciences*, 289(1972), p. 20212665. doi: 10.1098/rspb.2021.2665.

Cott, H. B. (1940) *Adaptive coloration in animals*. London: Methuen.

Cuthill, I. C. *et al.* (2017) 'The biology of color', *Science*, 357(6350). doi: 10.1126/science.aan0221.

Dobin, A. *et al.* (2013) 'STAR: ultrafast universal RNA-seq aligner', *Bioinformatics*, 29(1), pp. 15–21. doi: 10.1093/bioinformatics/bts635.

Drapeau, M. D. *et al.* (2006) 'Evolution of the Yellow/Major Royal Jelly Protein family and the emergence of social behavior in honey bees', *Genome Research*, 16(11), pp. 1385–1394. doi: 10.1101/gr.5012006.

Emms, D. M. and Kelly, S. (2019) 'OrthoFinder: phylogenetic orthology inference for comparative genomics', *Genome Biology*, 20(1), p. 238. doi: 10.1186/s13059-019-1832-y.

Endler, J. A. (1988) 'Frequency-dependent predation, crypsis and aposematic coloration', *Philosophical Transactions of the Royal Society B: Biological Sciences*, 319, pp. 505–523.

Ferguson, L. C. *et al.* (2011) 'Evolution of the insect yellow gene family', *Molecular Biology and Evolution*, 28(1), pp. 257–272. doi: 10.1093/molbev/msq192.

Funk, E. R. *et al.* (2021) 'A supergene underlies linked variation in color and morphology in a Holarctic songbird', *Nature Communications*, 12(1), p. 6833. doi: 10.1038/s41467-021-27173-z.

Galarza, J. A. *et al.* (2014) 'Temporal relationship between genetic and warning signal variation in the aposematic wood tiger moth (Parasemia plantaginis)', *Molecular Ecology*, 23(20), pp. 4939–4957. doi: 10.1111/mec.12913.

Galarza, J. A. (2021) 'Comparative transcriptomics of albino and warningly-coloured caterpillars', *Ecology and Evolution*, 11(12), pp. 7507–7517. doi: https://doi.org/10.1002/ece3.7581.

Galván, I. *et al.* (2015) 'Insects synthesize pheomelanin', *Pigment Cell and Melanoma Research*, 28(5), pp. 599–602. doi: 10.1111/pcmr.12397.

Gawne, R. and Nijhout, H. F. (2019) 'Expanding the nymphalid groundplan's domain of applicability: pattern homologies in an arctiid moth (Utetheisa ornatrix)', *Biological Journal of the Linnean Society*, 126(4), pp. 912–924. doi: 10.1093/biolinnean/bly193.

Gordon, S. P. *et al.* (2015) 'Colour polymorphism torn apart by opposing positive frequency-
dependent selection, yet maintained in space', *Journal of Animal Ecology*, 84(6), pp. 1555–
1564. doi: 10.1111/1365-2656.12416.

Guindon, S. *et al.* (2010) 'New Algorithms and Methods to Estimate Maximum-Likelihood
Phylogenies: Assessing the Performance of PhyML 3.0', *Systematic Biology*, 59(3), pp. 307–
321. doi: 10.1093/sysbio/syq010.

Hegna, R. H., Galarza, J. A. and Mappes, J. (2015) 'Global phylogeography and geographical
variation in warning coloration of the wood tiger moth (Parasemia plantaginis)', *Journal of
Biogeography*, 42(8), pp. 1469–1481. doi: 10.1111/jbi.12513.

Hines, H. M. *et al.* (2017) 'Melanic variation underlies aposematic color variation in two
hymenopteran mimicry systems', *PLoS ONE*, 12(7), pp. 1–17. doi:
10.1371/journal.pone.0182135.

Hoekstra, H. E. (2006) 'Genetics, development and evolution of adaptive pigmentation in
vertebrates', *Heredity*, 97(3), pp. 222–234. doi: 10.1038/sj.hdy.6800861.

Huu, C. N. *et al.* (2020) 'Supergene evolution via stepwise duplications and
neofunctionalization of a floral-organ identity gene', *Proceedings of the National Academy of
Sciences of the United States of America*, 117(37), pp. 23148–23157. doi:
10.1073/pnas.2006296117.

Huxley, J. (1955) 'Morphism and evolution', *Heredity*, 9(1), pp. 1–52. doi:
10.1038/hdy.1955.1.

Ito, K. *et al.* (2010) 'Yellow-e determines the color pattern of larval head and tail spots of the
silkworm Bombyx mori', *Journal of Biological Chemistry*, 285(8), pp. 5624–5629. doi:
10.1074/jbc.M109.035741.

Jorge García, A., Polidori, C. and Nieves-Aldrey, J. L. (2016) 'Pheomelanin in the secondary
sexual characters of male parasitoid wasps (Hymenoptera: Pteromalidae)', *Arthropod
Structure & Development*, 45(4), pp. 311–319. doi: https://doi.org/10.1016/j.asd.2016.05.001.

Joron, M. *et al.* (2011) 'Chromosomal rearrangements maintain a polymorphic supergene
controlling butterfly mimicry', *Nature*, 477(7363), pp. 203–206. doi: 10.1038/nature10341.

Katoh, K. and Standley, D. M. (2013) 'MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability', *Molecular Biology and Evolution*,
30(4), pp. 772–780. doi: 10.1093/molbev/mst010.

Kayser, H. (1985) 'Pigments', in Kerkut, G. A. and Gilbert, L. I. (eds) *Comparative Insect
Physiology, Biochemistry, and Pharmacology*. Oxford: Pergamon Press, pp. 367–415.

Kolb, A. M. *et al.* (1997) 'Determination of Pheomelanin by Measurement of
Aminohydroxyphenylalanine Isomers with High-Performance Liquid Chromatography',
Analytical Biochemistry, 252(2), pp. 293–298. doi: https://doi.org/10.1006/abio.1997.2342.

Kratochwil, C. *et al.* (2018) 'Agouti-related peptide 2 facilitates convergent evolution of
stripe patterns across cichlid fish radiations', *Science*, 362(6413), pp. 457–460. doi:
10.1126/science.aao6809.

Kunte, K. *et al.* (2014) 'doublesex is a mimicry supergene', *Nature*, 507(7491), pp. 229–232.
doi: 10.1038/nature13112.

Küpper, C. *et al.* (2016) 'A supergene determines highly divergent male reproductive morphs
in the ruff', *Nature Genetics*, 48(1), pp. 79–83. doi: 10.1038/ng.3443.

Langmead, B. and Salzberg, S. L. (2012) 'Fast gapped-read alignment with Bowtie 2',
Nature Methods, 9, p. 357. Available at: http://dx.doi.org/10.1038/nmeth.1923.

Li, H. *et al.* (2009) 'The Sequence Alignment/Map format and SAMtools', *Bioinformatics*,
25(16), pp. 2078–2079. doi: 10.1093/bioinformatics/btp352.

Li, H. (2013) 'Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM', *arXiv*, p. 1303.3997.

Linnen, C. R. *et al.* (2013) 'Adaptive Evolution of Multiple Traits Through Multiple
Mutations at a Single Gene', *Science*, 339(6125), pp. 1312–1316. doi:
10.1126/science.1233213.

Massey, J. H. *et al.* (2019) 'The yellow gene influences Drosophila male mating success
through sex comb melanization', *eLife*. Edited by K. VijayRaghavan and C. L. Peichel, 8, p.
e49388. doi: 10.7554/eLife.49388.

Matsuoka, Y. and Monteiro, A. (2018) 'Melanin Pathway Genes Regulate Color and
Morphology of Butterfly Wing Scales', *Cell Reports*, 24(1), pp. 56–65. doi:
10.1016/j.celrep.2018.05.092.

Mazo-Vargas, A. *et al.* (2022) 'Deep cis-regulatory homology of the butterfly wing pattern
ground plan', *Science*, 378(6617), pp. 304–308. doi: 10.1126/science.abi9407.

McGraw, K. J. *et al.* (2002) 'The effect of dietary carotenoid access on sexual dichromatism
and plumage pigment composition in the American goldfinch', *Comparative Biochemistry
and Physiology Part B: Biochemistry and Molecular Biology*, 131(2), pp. 261–269. doi:
https://doi.org/10.1016/S1096-4959(01)00500-0.

Mcgraw, K. J. and Wakamatsu, K. (2004) 'Melanin Basis of Ornamental Feather Colors in
Male Zebra Finches', *The Condor*, 106(3), pp. 686–690. doi: 10.1093/condor/106.3.686.

McKenna, A. *et al.* (2010) 'The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data', *Genome Research*, 20, pp. 1297–1303.
doi: 10.1101/gr.107524.110.

McKinnon, J. S. and Pierotti, M. E. R. (2010) 'Colour polymorphism and correlated
characters: Genetic mechanisms and evolution', *Molecular Ecology*, 19(23), pp. 5101–5125.
doi: 10.1111/j.1365-294X.2010.04846.x.

Miyazaki, S. *et al.* (2014) 'Sexually Dimorphic Body Color Is Regulated by Sex-Specific
Expression of Yellow Gene in Ponerine Ant, Diacamma Sp', *PLOS ONE*, 9(3), p. e92875.
Available at: https://doi.org/10.1371/journal.pone.0092875.

Nishikawa, H. *et al.* (2015) 'A genetic mechanism for female-limited Batesian mimicry in
Papilio butterfly', *Nature Genetics*, 47(4), pp. 405–409. doi: 10.1038/ng.3241.

Nokelainen, O. *et al.* (2012) 'Trade-off between warning signal efficacy and mating success
in the wood tiger moth', *Proceedings of the Royal Society B: Biological Sciences*, 279(1727),
pp. 257–265. doi: 10.1098/rspb.2011.0880.

Nokelainen, O. *et al.* (2014) 'Changes in predator community structure shifts the efficacy of
two warning signals in Arctiid moths', *Journal of Animal Ecology*, 83(3), pp. 598–605. doi:
10.1111/1365-2656.12169.

Nokelainen, O. *et al.* (2022) 'Genetic colour variation visible for predators and conspecifics
is concealed from humans in a polymorphic moth', *Journal of Evolutionary Biology*,
(February), pp. 467–478. doi: 10.1111/jeb.13994.

Orteu, A. and Jiggins, C. D. (2020) 'The genomics of coloration provides insights into
adaptive evolution', *Nature Reviews Genetics*, 21(8), pp. 461–475. doi: 10.1038/s41576-020-
0234-z.

De Pasqual, C. *et al.* (2022) 'Heterozygote advantage and pleiotropy contribute to
intraspecific color trait variability', *Evolution*, pp. 2389–2403. doi: 10.1111/evo.14597.

Polidori, C., Jorge, A. and Ornosa, C. (2017) 'Eumelanin and pheomelanin are predominant
pigments in bumblebee (Apidae: Bombus) pubescence', *PeerJ*, 2017(5), pp. 1–21. doi:
10.7717/peerj.3300.

Quinlan, A. R. and Hall, I. M. (2010) 'BEDTools: a flexible suite of utilities for comparing
genomic features', *Bioinformatics*, 26(6), pp. 841–842. doi: 10.1093/bioinformatics/btq033.

Rastas, P. (2017) 'Lep-MAP3: robust linkage mapping even for low-coverage whole genome
sequencing data', *Bioinformatics*, 33, pp. 3726–3732. doi: 10.1093/bioinformatics/btx494.

Ritchie, M. E. *et al.* (2015) 'limma powers differential expression analyses for RNA-
sequencing and microarray studies', *Nucleic Acids Research*, 43(7), pp. e47–e47. doi:
10.1093/nar/gkv007.

Rodriguez-Caro, F. *et al.* (2021) 'Novel Doublesex Duplication Associated with Sexually
Dimorphic Development of Dogface Butterfly Wings', *Molecular Biology and Evolution*,
38(11), pp. 5021–5033. doi: 10.1093/molbev/msab228.

Rojas, B. *et al.* (2017) 'How to fight multiple enemies: Target-specific chemical defences in
an aposematic moth', *Proceedings of the Royal Society B: Biological Sciences*, 284(1863).
doi: 10.1098/rspb.2017.1424.

Rojas, B., Gordon, S. P. and Mappes, J. (2015) 'Frequency-dependent flight activity in the
colour polymorphic wood tiger moth', *Current Zoology*, 61(4), pp. 765–772. doi:
10.1093/czoolo/61.4.765.

Rönkä, K. *et al.* (2020) 'Geographic mosaic of selection by avian predators on hindwing
warning colour in a polymorphic aposematic moth', *Ecology Letters*, 23(11), pp. 1654–1663.
doi: https://doi.org/10.1111/ele.13597.

Shamim, G. *et al.* (2014) 'Biochemistry and biosynthesis of insect pigments', *European
Journal of Entomology*, 111(2), pp. 149–164. doi: 10.14411/eje.2014.021.

Smith, G., Macias-Muñoz, A. and Briscoe, A. D. (2016) 'Gene Duplication and Gene
Expression Changes Play a Role in the Evolution of Candidate Pollen Feeding Genes in
Heliconius Butterflies', *Genome Biology and Evolution*, 8(8), pp. 2581–2596. doi:
10.1093/gbe/evw180.

Sugumaran, M. (2010) 'Chapter 5 - Chemistry of Cuticular Sclerotization', in Simpson, S. J.
B. T.-A. in I. P. (ed.). Academic Press, pp. 151–209. doi: https://doi.org/10.1016/B978-0-12-
381387-9.00005-1.

Suomalainen, E. (1938) 'Die erblichkeitsverhältnisse des männlichen dimorphismus bei
parasemia plantaginis', *Hereditas*, 24, pp. 386–390.

Thompson, M. J. and Jiggins, C. D. (2014) 'Supergenes and their role in evolution', *Heredity*,
113(1), pp. 1–8. doi: 10.1038/hdy.2014.20.

Timmermans, M. J. T. N. *et al.* (2014) 'Comparative genomics of the mimicry switch in
Papilio dardanus', *Proceedings of the Royal Society B: Biological Sciences*, 281(1787). doi:
10.1098/rspb.2014.0465.

Umebachi, Y. (1975) 'Further studies on the dopamine derivative, SN-1 derived from the
yellow pigments of Papilio xuthus', *Insect Biochemistry*, 5, pp. 73–92.

Wakamatsu, K. *et al.* (2014) 'Reduction of the Nitro Group to Amine by Hydroiodic Acid to
Synthesize o-Aminophenol Derivatives as Putative Degradative Markers of Neuromelanin',
Molecules . doi: 10.3390/molecules19068039.

Wakamatsu, K., Ito, S. and Rees, J. L. (2002) 'The Usefulness of 4-Amino-3-
hydroxyphenylalanine as a Specific Marker of Pheomelanin', *Pigment Cell Research*, 15(3),
pp. 225–232. doi: https://doi.org/10.1034/j.1600-0749.2002.02009.x.

Wang, J. *et al.* (2013) 'A Y-like social chromosome causes alternative colony organization in
fire ants', *Nature*, 493(7434), pp. 664–668. doi: 10.1038/nature11832.

Wang, P. *et al.* (2022) 'Yellow-b, -c, -e and -h are required for normal body coloration in
Henosepilachna vigintioctopunctata', *Archives of Insect Biochemistry and Physiology*,
109(2), p. e21856. doi: https://doi.org/10.1002/arch.21856.

Wilson, R. *et al.* (1976) 'Behavioural pleiotropy of the yellow gene in Drosophila
melanogaster', *Genetical Research*. 2009/04/14, 28(1), pp. 75–88. doi: DOI:
10.1017/S0016672300016748.

Winters, A. E. *et al.* (2021) 'Multimodal Aposematic Defenses Through the Predation
Sequence', *Frontiers in Ecology and Evolution*, 9(August), pp. 1–18. doi:
10.3389/fevo.2021.657740.

Wittkopp, P. J., Vaccaro, K. and Carroll, S. B. (2002) 'Evolution of yellow gene regulation
and pigmentation in Drosophila', *Current Biology*, 12(18), pp. 1547–1556. doi:
10.1016/S0960-9822(02)01113-2.

Yen, E. C. *et al.* (2020) 'A haplotype-resolved, de novo genome assembly for the wood tiger
moth (Arctia plantaginis) through trio binning', *GigaScience*, 9(8), pp. 1–12. doi:
10.1093/gigascience/giaa088.

Zhang, L. *et al.* (2017) 'Genetic basis of melanin pigmentation in butterfly wings', *Genetics*,
205(4), pp. 1537–1550. doi: 10.1534/genetics.116.196451.

Zhang, L., Mazo-Vargas, A. and Reed, R. D. (2017) 'Single master regulatory gene
coordinates the evolution and development of butterfly color and iridescence', *Proceedings of*
the National Academy of Sciences, 114, pp. 10707–10712. doi: 10.1073/pnas.1709058114.