Comparison of R9.4.1/Kit10 and R10/Kit12 Oxford Nanopore flowcells and chemistries in bacterial genome reconstruction

1.1 Author names

Nicholas Sanderson[#*1,2] (ORCiD: 0000-0001-6370-1961)/Natalia Kapel[#1,2] (ORCiD: 0000-0002-5914-9999), Gillian Rodger[1,3], Hermione Webster[1,2], Samuel Lipworth[2,3], Teresa street[1,2], Tim Peto[1,2,3], Derrick Crook[1,2,3] (ORCiD: 0000-0002-0590-2850), Nicole Stoesser[*1,2,3] (ORCiD: 0000-0002-4508-7969)

[#]Contributed equally

*Corresponding/alternative corresponding author

1.2 Affiliation

[1] NIHR Oxford Biomedical Research Centre, University of Oxford, Oxford, UK

[2] Nuffield Department of Medicine, University of Oxford, Oxford, UK

[3] NIHR Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance at University of Oxford in partnership with Public Health England, Oxford, UK

Corresponding authors

Nicholas Sanderson: nicholas.sanderson@ndm.ox.ac.uk

Nicole Stoesser: nicole.stoesser@ndm.ox.ac.uk

1.3 Keyword

Genome sequencing, hybrid assembly, long-read assembly

Assemblies have been made available at:
https://figshare.com/articles/online_resource/q20_comparison_genome_assemblies/196838
67.

Code and analysis outputs are available at:
https://gitlab.com/ModernisingMedicalMicrobiology/assembly_comparison_analysis/-
/tree/main (tagged version v0.5.5).

2. Abstract

Complete, accurate, cost-effective, and high-throughput reconstruction of bacterial genomes
for large-scale genomic epidemiological studies is currently only possible with hybrid
assembly, combining long- (typically using nanopore sequencing) and short-read (Illumina)
datasets. Being able to utilise nanopore-only data would be a significant advance. Oxford
Nanopore Technologies (ONT) have recently released a new flowcell (R10.4) and chemistry
(Kit12), which reportedly generate per-read accuracies rivalling those of Illumina data. To
evaluate this, we sequenced DNA extracts from four commonly studied bacterial pathogens,
namely *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and
Staphylococcus aureus, using Illumina and ONT's R9.4.1/Kit10, R10.3/Kit12, R10.4/Kit12
flowcells/chemistries. We compared raw read accuracy and assembly accuracy for each
modality, considering the impact of different nanopore basecalling models, commonly used
assemblers, sequencing depth, and the use of duplex versus simplex reads. Duplex
nanopore reads combined with "super accuracy" (sup) basecalling have high per-read
accuracies and could be used to robustly reconstruct bacterial genomes without the use of
Illumina data. However, the per-run yield of duplex reads generated in our hands with
standard sequencing protocols was low (typically <10%), with substantial implications for
cost and throughput if relying on nanopore data only to enable bacterial genome
reconstruction. In addition, recovery of small plasmids with the best-performing long-read
assembler (Flye) was inconsistent. R10.4/Kit12 combined with sup basecalling holds
promise as a singular sequencing technology in the reconstruction of commonly studied
bacterial genomes, but hybrid assembly (Illumina+R9.4.1 hac) currently remains the highest
throughput, most robust, and cost-effective approach.

3. Impact statement

Our understanding of microbes has been greatly enhanced by the capacity to evaluate their
genetic make-up using a technology known as whole genome sequencing. Sequencers
represent microbial genomes as stretches of shorter sequence known as 'reads', which are
then assembled using computational algorithms. Different types of sequencing approach
have advantages and disadvantages with respect to the accuracy and length of the reads
they generate; this in turn affects how reliably genomes can be assembled.

Currently, to completely reconstruct bacterial genomes in a high-throughput and cost-
effective manner, researchers tend to use two different types of sequencing data, namely
Illumina (short-read) and nanopore (long-read) data. Illumina data are highly accurate;
nanopore data are much longer, and this combination facilitates accurate and complete
bacterial genomes in a so-called "hybrid assembly". However, new developments in
nanopore sequencing have reportedly greatly improved the accuracy of nanopore data,
hinting at the possibility of requiring only a single sequencing approach for bacterial
genomics.

Here we evaluate these improvements in nanopore sequencing in the reconstruction of four
bacterial reference strains, where the true sequence is already known. We show that
although these improvements are extremely promising, for high-throughput, low-cost
reconstruction of bacterial genomes hybrid assembly currently remains the optimal
approach.

4. Data summary

The authors confirm all supporting data, code and protocols have been provided
within the article, through supplementary data files, or in publicly accessible
repositories.

Nanopore fast5 and fastq data are available in the ENA under project accession:
PRJEB51164.

Assemblies have been made available at:
https://figshare.com/articles/online_resource/q20_comparison_genome_assemblies/196838
67.

Code and analysis outputs are available at:
https://gitlab.com/ModernisingMedicalMicrobiology/assembly_comparison_analysis/-
/tree/main (tagged version v0.5.5).

5. Introduction

Bacterial whole genome sequencing has become a prominent tool in the biological sciences,
with wide-ranging applications from epidemiology to diagnostics(1). Important considerations
include sequencing throughput, read length (which facilitates complete reconstruction of
bacterial chromosomes and plasmids), read accuracy, accessibility and cost. Historically,

short-read Illumina sequencing has been the leading high-throughput, high-accuracy
technology, but is limited in its capacity to completely reconstruct genomes, particularly in
the presence of repetitive sequences. Nanopore sequencing (Oxford Nanopore
Technologies [ONT]) has become one of the most widely adopted long-read sequencing
approaches, enabled by affordable, small-footprint sequencing platforms, but has been
limited to some extent by its accuracy. Combining short and long-read sets from both
technologies in the form of hybrid assembly has facilitated cost-effective, highly accurate and
scalable genome reconstruction for large bacterial isolate collections(2, 3), such as by
multiplexing 96 *E. coli* isolates on a single nanopore flowcell(3). For nanopore sequencing,
developments in multiplexing, rapid library preparation and flow cell reuse after washing
have streamlined this process(4).

ONT have undertaken iterative development of their sequencing flowcells and chemistries,
releasing the R10.3 (FLO-MIN111) flowcells for consumers in January 2020 and the Kit12
(Q20+) chemistry and R10.4 flowcell (FLO-MIN112) in their store in late 2021. The proposed
advantages of the R10.4/Kit12 system include: (i) a new motor to facilitate more controlled
passage of the nucleic acid template through the sequencing pore thereby avoiding template
slippage; (ii) "duplex" read sequencing - where the forward and reverse strand of a single
nucleic acid molecule are sequenced in succession to improve accuracy; and (iii) an
optimized pore with a longer pore head to better resolve homopolymers.

These new developments however come with some potential disadvantages. Sequencing
yields for the R10.3 flowcells were lower than those using R.9.4.1 flowcells (thought to be
due to the slower passage of template through pores)(5). The use of R10 flowcells also
currently requires a ligation-based library preparation, which results in longer sequencing
turnaround times when compared with rapid transposase-based library preparation kits
which can be used with R9.4.1 flowcells. Ligation-based preparations may also miss the
capture and sequencing of small plasmids(6). The reported improvements in per-read
accuracy with R10/Kit12 are also potentially dependent on the use of super accuracy (sup)
basecalling models; however, on the same computing infrastructure sup basecalling takes 2-
8x longer than the previous typical approach using high accuracy (hac) basecalling models,
which may preclude "on-machine" basecalling in real-time(7).

Sequencing accuracy can be characterized using several different metrics, including: (i) raw
read accuracy (the accuracy achieved when reading a single nucleic acid fragment once)
and (ii) assembly accuracy (the capacity to accurately reconstruct complete genomes in
terms of structure, sequence identity and coding sequence content). We therefore set out to
compare data and assemblies generated by R9.4.1/Kit10 and R10/Kit12 nanopore
flowcells/chemistries, comparing these with Illumina-only sequence data and hybrid
assembly, and investigating the impact of sup versus hac basecalling and metrics for duplex
sequencing reads. We undertook this comparison for four reference bacterial strains

reflecting different species, genome sizes, %GC content, plasmid content and plasmid sizes.
We also evaluated the impact of sequencing depth on the capacity to reconstruct the
reference bacterial genomes, and whether flowcell washing would still enable flow cell reuse
with the new flowcells and chemistry.

6. Methods

Bacterial isolates and DNA extraction

Four reference bacterial strains were sequenced for this study, namely: *Escherichia coli*
CFT073 (Genbank accession: NC_004431.1), *Klebsiella pneumoniae* MGH78578
(NC_009648.1-NC_009653.1), *Pseudomonas aeruginosa* PAO1 (NC_002516.2) and
Staphylococcus aureus MRSA252 (NC_002952.2). Stock cultures were stored at -80°C in
nutrient broth supplemented with 10% glycerol. For DNA extraction, stocks were sub-
cultured on Columbia blood agar at 37°C overnight.

Long fragment DNA extraction from sub-cultured strains was performed using the Qiagen
Genomic tip 100/G kit (Qiagen). Quality and fragment length assessments were measured
with the Qubit fluorometer (ThermoFisher Scientific) and TapeStation (Agilent). The same
DNA extract, stored in elution buffer at 4°C was used for all sequencing experiments. DNA
concentration and fragment lengths were evaluated longitudinally to ensure that there was
minimal obvious degradation (Tables S1-4, Figs.S1-3).

Nanopore sequencing

The experimental workflow is shown in Fig.1. For the experiment using the R9.4.1 (FLO-
MIN106) flowcell (denoted as R.9.4 throughout), ONT sequencing libraries were prepared by
multiplexing DNA extracts from all four isolates using the Rapid Barcoding Sequencing
(SQK-RBK004) kit according to the manufacturer's protocol; sequencing was undertaken on
a GridION for 48 hours.

For the experiments using the R10.3 (FLO-MIN111) and R10.4 (FLO-MIN112) flowcells,
ONT sequencing libraries were prepared from DNA extracts using the Q20+ Early Access
Kit (SQK-Q20EA) ligation-based protocol. During adapter ligation and clean-up the long
fragment buffer was used to enrich for DNA fragments >3kb. Each DNA extract was
sequenced on a single flowcell. After sequencing the *S. aureus* MRSA252 library, the R10.4
(FLO-MIN112) flowcell was washed with the flowcell wash kit (EXP-WSH004) according to
the manufacturer's protocol, before reusing the flowcell to sequence the *P. aeruginosa*
PAO1 library. For the R10.3 experiments, sequencing was undertaken on a GridION for 48
hours; for the unplexed R10.4 experiments sequencing times were terminated prematurely.
The flowcell usage strategy and pore counts for each flowcell prior to use are summarised in
Table S5.

Finally, in a separate experiment, the four DNA extracts were also multiplexed on the R10.4
(FLO-MIN112) flowcell using the Native Barcoding Kit (SQK-NBD112.24); sequencing was
undertaken on a GridION for 48 hours.

Illumina sequencing

DNA extracts for all isolates were also sequenced on the Illumina MiSeq, as part of a wider
run plexing 20 bacterial extracts. Libraries were constructed following the Illumina DNA Prep
protocol, according to the manufacturer's instructions (including standard normalization for
libraries ["Protocol A"]). Library DNA concentrations were quantified by Qubit fluorometry and
size distributions of libraries determined using the TapeStation, as above. Sequencing was
performed using the MiSeq Reagent Kit v2, generating 150 bp paired-end reads.

Data processing and bioinformatic methods

R10.4 duplex read pairs were identified and prepared for basecalling using ONT's duplex
tools (https://pypi.org/project/duplex-tools/; v 0.2.9). R9.4, R10.3, and R10.4 raw nanopore
reads were hac basecalled with Guppy (ONT) versions 5.0.12+eb1a981
(dna_r9.4.1_450bps_hac.cfg), 5.0.13+bbad529 (res_dna_r103_q20ea_crf_v034.cfg), and
5.0.16+b9fcd7b (dna_r10.4_e8.1_hac.cfg) respectively, as recommended by ONT. R9.4,
R10.3, R10.4 (all reads) and R10.4 duplex raw nanopore reads were also basecalled using
sup models dna_r9.4.1_e8.1_sup.cfg, dna_r10.3_450bps_sup.cfg, dna_r10.4_e8.1_sup.cfg.
Basecalled read summary statistics were generated with seqkit stats using '-T' and '-all'
flags(8).

Nanopore reads were subsampled using Rasusa(9) to depths of 10, 20, 30, 40, 50, and 100
average coverage. Nanopore reads were assembled with Canu (version 2.2, using
maxInputCoverage=100 and otherwise default parameters)(10), or Flye (using the --meta
and --nano-hq parameters and otherwise defaults, version 2.9-b1768)(11), both of which are
commonly used long-read only assemblers that have been shown to optimize long-read only
assembly quality(12). Subsampled nanopore reads were combined with Illumina reads for
hybrid assembly using Unicycler (version 0.4.8, default parameters)(13). The SPAdes
(version 3.15.3)(14) assemblies generated from Illumina data as part of the Unicycler
pipeline were used as the Illumina-only assemblies for comparative evaluations.

Given the previous discrepancies observed between multiple resequenced assemblies for *E.
coli* CFT073 and *K. pneumoniae* MGH78578(15), and the genetic and phenotypic
differences observed in different laboratory sub-culture stocks of *P. aeruginosa* PAO1(16,
17), we generated an Illumina-corrected reference sequence to use as the "gold standard"

comparator for this evaluation. Reference genomes for *E. coli* CFT073 (Genbank accession:

AE014075.1), *K. pneumoniae* MGH78578 (CP000647.1), *P. aeruginosa* PAO1

(NC_002516.2), *S. aureus* MRSA252 (NC_002952.2) and the respective Illumina datasets

generated for this study were used as inputs for the SNIPPY pipeline (version 4.6.0)

(https://github.com/tseemann/snippy); output consensus fasta files represented the new

Illumina-corrected reference sequences used in this study.

Assembled contigs from nanopore, Illumina, and hybrid assemblies were compared against

the Illumina-corrected reference sequences using DNAdiff version 1.3(18).

Assembled contigs from nanopore, Illumina, and hybrid assemblies as well as the Illumina-

corrected reference sequences were annotated with Prokka (version 1.14.6)(19), using the

corresponding reference GenBank files to ascertain reference proteins using the '--proteins'

flag.

Translated amino acid sequences for Prokka annotations in the different test assemblies

(Canu, Flye [long-read only], Unicycler [hybrid long-/short-read], SPAdes [short-read only])

and Illumina-corrected reference sequences were compared using the script AAcompare.py

in the workflow provided (see below for the repository link). This looked for exact amino acid

sequence matches (i.e. 100% identity along 100% of the translated protein) between the

Illumina-corrected reference and assembled contigs to determine how intact assembled

coding sequences were for each assembly method.

Per read error rates were calculated by mapping the raw reads to the Illumina corrected

references sequences using minimap2 (version 2.22-r1101)(18). The percent identity was

calculated from the query distance (NM tag) divided by the query length, multiplied by 100,

using the bamreadstats.py script provided in the gitlab repository (link below).

A workflow for this analysis has been written using nextflow(18) and is available on gitlab

(https://gitlab.com/ModernisingMedicalMicrobiology/assembly_comparison). Outputs from

the analyses are also available in this repository (tagged version v0.5.5).

Data visualization

Figures and plots for this manuscript were generated using the ggplot2 and patchwork

packages in R, and Biorender (www.biorender.com).

7. Results

Sequencing yield and read length distributions

The total data yield after 48 hours of sequencing from the R9.4 flowcell was 11.0Gb (four
isolate extracts multiplexed on one sequencing run), compared with 4.0Gb for the R10.4
multiplexed run (Table 1, Fig.S4). For the individual R10.3 flowcells a median of
8.2Gb/flowcell (IQR: 7.3-8.8Gb) were generated by 48 hours of sequencing, and
6.7Gb/flowcell (IQR: 6.6-7.4Gb) for the R10.4 flowcells respectively by 20-30 hours of
sequencing (Table 1, Fig.S4). 21.3Mb of data were generated for the extracts from the
Illumina runs (Table 1).

Read length distributions for a subsample of 1000 reads by modality and species are shown
in Fig.2; overall, across species for nanopore data the median read length was 3580bp, the
maximum read length 388620bp and the minimum read length 77bp. Median read lengths
generated using R9.4 were longer (6273bp versus 2930bp for R10.4; two-sample Wilcoxon
test, p<0.001, comparison for hac basecalled data; Fig.2A). N50s are represented in Table
1; median N50 across species was 19496bp for R9.4.1 hac, 16002bp for R10.3, 20976bp for
R10.4 (all) and 16425bp for R10.4 duplex reads.

Duplex reads

The median proportion of duplex reads across the four unplexed, single-extract R10.4 runs
was 4.5% (3.8% for *E. coli*, 6.1% for *K. pneumoniae*, 4.5% for *P. aeruginosa*, and 4.5% for
S. aureus). For the multiplexed R10.4 run for each species these proportions were 2.3%,
5.4%, 6.0% and 4.7%.

Raw read accuracy by sequencing modality and species

Raw read accuracy (% identity when mapped to the reference) for a subsample of 1000
reads by sequencing data type/process (i.e. "sequencing modality") and species was highest
(as expected) for Illumina reads (modal accuracy: 100.0%), followed by R10.4 duplex reads
basecalled with the sup model (modal accuracy: 99.9%); modal accuracies for all the other
approaches were >97.0% (Fig.3). Sup basecalling improved modal accuracy for R10.4
reads, but not R10.3 or R9.4 reads; multiplexing had no impact (Fig.3). Median and modal
accuracies for each sequencing modality by species are detailed in Table S6.

In terms of insertions and deletions with respect to the reference, for long-read modalities
R10.4 sup called duplex data performed best (Fig.4A, 4B). The median number of insertions
observed per read was 0.94, 0.45, 0.37 and 0.0 for R9.4 hac, R10.3 hac and R10.4 sup and
R10.4 sup duplex respectively (two-sample Wilcoxon test for each versus R9.4 hac as the

reference category; all p<0.001), and for deletions 1.31, 0.73, 0.63 and 0.10 respectively
(two-sample Wilcoxon test for each versus R9.4 hac as the reference category; all p<0.001).

Assembly accuracy with respect to number of expected contigs in the reference sequences
and reference sequence size

We evaluated the capacity of each sequencing approach to accurately reconstruct (i) the
number of known contigs present in each reference isolate, and (ii) what percentage of the
Illumina-corrected reference was covered. All isolates contained single chromosomes only,
except the *K. pneumoniae* reference, which contained a chromosome and five plasmids
ranging in size from 3478-175879bp (Table 2).

Approaches using all the data and Unicycler or Flye largely generated single chromosomal
contigs, except those using R10.4 duplex reads only, particularly for multiplexed extracts,
likely because these reads were insufficient to cover the whole genome (Table 2; Fig.S5A).
Illumina-only assemblies generated much larger numbers of contigs as expected (Table 2).
Using all the data, single *K. pneumoniae* plasmid contigs were mostly obtained using any of
the long-read data and Flye, or hybrid assembly with Unicycler (Table 2, Fig.S5B). Using all
the data, Flye long-read only assemblies largely all missed the two smallest plasmids (Table
2, Fig.S5B).

Sub-sampling the data to 10x, 20x, 30x, 40x, 50x or 100x depth had variable effect - for the
most part single chromosomal contigs were assembled using long-reads only with >20x
depth; Unicycler could mostly be used with 10x long-read depth (Fig.S5A). The same effect
was seen for plasmids, except Flye struggled to reliably assemble the two largest plasmids
into single contigs with lower sequencing depths (Fig.S5B). Canu assemblies failed with 10x
sub-sampling, as expected given the default cut-offs.

For chromosomes, Canu long-read only assemblies tended to over-assemble structures (i.e.
reference coverage >100%, Fig.5A) whilst Illumina-only assemblies under-assembled
structures. Reference coverage % for Unicycler hybrid (R9.4+Illumina) was largely
unaffected by sub-sampling the data to 10x, 20x, 30x, 40x, 50x or 100x (Fig.5A). For
plasmids, Canu assembly again largely over-assembled the structures; Unicycler hybrid
(R9.4+Illumina) assembly was the only approach which consistently assembled all plasmids
at near 100% reference coverage across all sub-sampling depths (Fig.5B).

Assembly accuracy with respect to insertions, deletions and nucleotide-level mismatches

For each sequencing and assembly modality the number of indels and nucleotide-level
mismatches (SNPs) were evaluated (Figs.6A, 6B). Overall, across all species, the R10.4 sup
duplex data assembled with Canu had the smallest number of SNPs (median: 13 SNPs),
followed by the Illumina data (35 SNPs); for indels the Illumina data had the smallest number
(median: 1 indel) followed by the R10.4 sup duplex data assembled with Canu (median: 18
indels). Larger numbers of errors were observed when sequencing *E. coli* and *K.*
pneumoniae (Figs.6A, 6B). For the Unicycler hybrid assemblies a median of 216 SNPs and
28 indels were observed across all species. A detailed breakdown of numbers is presented
in Table S7.

Assembly accuracy with respect to coding sequence content

Amongst the long-read only assemblies, coding sequence content was most accurately
reconstructed using R10.4 sup or sup duplex data and then R10.3 data (Fig.8). Notably, the
duplex datasets from the unplexed 10.4 runs were used, as from multiplexed runs the duplex
yields were insufficient to facilitate assembly in most cases (Table 2). Long-read only
assembly with R9.4 data missed up to 10-15% of coding sequences (Fig.8).

8. Discussion

In this pragmatic study evaluating the impact of different nanopore sequencing flowcells and
chemistries on the capacity to fully reconstruct genomes of four commonly studied bacteria,
we have shown that sup basecalled R10.4/Kit12 data and sup called duplex data are
approaching accuracies that would enable these to be effectively used for the reconstruction
of bacterial genomes without requiring Illumina data to generate hybrids. However, hybrid
assembly (Illumina+9.4.1 hac data) remains the most robust approach in terms of contig
(both chromosomes and plasmids) and CDS recovery without over-assembly, and facilitates
the multiplexing of large numbers of isolates per flowcell, given that in this and at least one
other study(3), <10x long-read depth is required for the accurate reconstruction of
chromosomes and plasmids by combining R9.4.1 and Illumina data using Unicycler. Highly
accurate long-read only assembly and genome reconstructions was optimized by generating
duplex reads, which in our hands made up a small proportion of the output (<10%); as such,
it would come at a significant cost per isolate as a result of being able to only generate data
for 1-2 isolates per flowcell. Very approximate costs per genome therefore for hybrid
assembly versus duplex/long-read-only assembly would be £50-70/genome versus £300-
600/genome.

Although barcoding up to 96 isolates has recently been enabled for the R10.4/Kit12
combination, the data yields per flowcell (~4Gb) would likely preclude viable assembly for 96
E. coli isolates with a typical genome size of ~5Mb (would give <8x coverage). There is also
a current requirement to use a ligation-based library preparation, which lengthens the

processing time, and may impact on plasmid recovery(6). We observed issues with
recovering small plasmids (<5kb) using Flye in this study although both of these small
plasmids could be reliably recovered in Canu assemblies; consistent with this a previous
evaluation has shown that 8-15% of small plasmids are not recovered using these long-read-
only assemblers(12). Similarly, as shown in this study and in other work(12), the basic Canu
workflow 'over-assembles' the data, and contigs require trimming of overlaps in order to
recreate accurate, single, circularized structures.

There are currently few other published studies on the performance of R10.4/Kit12 for
bacterial analyses. We found only one preprint investigating its use on a mock microbial
community (7 bacterial species and 1 fungal species) which found similar modal accuracy
scores of 99% using sup basecalling, and a requirement of 40x to be able to reliably
assemble a genome(20). Their hypothesis was that improved read accuracies were due to
an improved ability to call homopolymers, which we did not investigate in this manuscript. It
was unclear what proportion of reads they characterized as duplex reads.

There are several limitations with our study. We have not exhaustively investigated all
possible approaches to genome assembly, but rather taken a pragmatic approach in
assembling the data with several commonly used assemblers, without additional bespoke
management of workflows. We had low duplex read yields compared with those reported by
ONT (up to 30-40% per flowcell); further optimization is needed to see if these can be
achieved. We have investigated only a limited number of isolates and plasmids, but these
represent a range of %GC and sizes, and are likely to reflect genetic content more widely in
other species; we have not generated replicate datasets. Improvements and upgrades to
nanopore flowcells, chemistries and basecallers occur regularly and nanopore will be
releasing the R10.4.1 flowcell and kit14 chemistries later this year which may further
optimise the quality of long-read only outputs.

In summary, the combination of R10.4/Kit12 flowcells/chemistries look promising for highly
accurate, long-read only bacterial genome assembly; however, this is dependent on the
generation of duplex reads, which currently make up only a small proportion of sequencing
yield, and requires superior accuracy basecalling, which will prolong workflow times. In
addition, for large-scale projects to fully reconstruct 100s-1000s of bacterial isolates, hybrid
assembly, multiplexing and the use of flowcells/chemistries that support rapid barcoding are
currently better suited for higher throughput and are more cost-effective per reconstructed
genome.

9. Author statements

9.1 Authors and contributors

NK, NSa, DWC and NS designed the study. NK and GR performed the laboratory experiments and sequencing. NSa performed the bioinformatics analysis. NSt generated the data visualisations. NSt, NSa and NK wrote the first draft. All authors reviewed and approved the final draft.

9.2 Conflicts of interest

Oxford Nanopore Technologies supplied the R10.3 and R10.4 flowcells free of charge for this study. They were also involved in discussions regarding which data processing approaches to use to optimise basecalling and assembly outputs; however, they did not impact on the presentation of any of the results.

9.3 Funding information

This study was funded by the National Institute for Health Research (NIHR) Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance (NIHR200915), a partnership between the UK Health Security Agency (UKHSA) and the University of Oxford, and was supported by the NIHR Oxford Biomedical Research Centre (BRC). The computational aspects of this research were funded from the NIHR Oxford BRC with additional support from the Wellcome Trust Core Award Grant Number 203141/Z/16/Z. The views expressed are those of the author(s) and not necessarily those of the NHS, NIHR, UKHSA or the Department of Health and Social Care. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising. NS is an NIHR Oxford BRC Senior Research Fellow and an Oxford Martin Fellow.

9.4 Ethical approval

Not applicable

9.5 Consent for publication

Not applicable

9.6 Acknowledgements

We are grateful to Dr Celiq Souque and Prof Craig Maclean at the Department of Zoology, University of Oxford, for supplying the *Pseudomonas aeruginosa* PAO1 strain.

10. References

1. Van Goethem N, Descamps T, Devleesschauwer B, Roosens NHC, Boon NAM, Van Oyen H, et al. Status and potential of bacterial genomics for public health practice: a scoping review. Implementation science : IS. 2019;14(1):79.

2. Shaw LP, Chau KK, Kavanagh J, AbuOun M, Stubberfield E, Gweon HS, et al. Niche and local geography shape the pangenome of wastewater- and livestock-associated Enterobacteriaceae. Science advances. 2021;7(15).

3. Arredondo-Alonso S, Pöntinen AK, Cléon F, Gladstone RA, Schürch AC, Johnsen PJ, et al. A high-throughput multiplexing and selection strategy to complete bacterial genomes. GigaScience. 2021;10(12).

4. Lipworth S, Pickford H, Sanderson N, Chau KK, Kavanagh J, Barker L, et al. Optimized use of Oxford Nanopore flowcells for hybrid assemblies. Microb Genom. 2020;6(11).

5. Oxford Nanopore Technologies. https://nanoporetech.com/about-us/news/r103-newest-nanopore-high-accuracy-nanopore-sequencing-now-available-store; last accessed: 07/Apr/2022.

6. Wick RR, Judd LM, Wyres KL, Holt KE. Recovery of small plasmid sequences via Oxford Nanopore sequencing. Microb Genom. 2021;7(8).

7. Benton M. 2021. Nanopore Guppy GPU basecalling on Windows using WSL2https://hackmd.io/@Miles/rkYKDHPsO. Blog post, last accessed: 07/Apr/2022.

8. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation. PLoS One. 2016;11(10):e0163962.

9. Hall MB. Rasusa: Randomly subsample sequencing reads to a specified coverage. The Journal of Open Source Software.

10. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res. 2017;27(5):722-36.

11. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat graphs. Nature biotechnology. 2019;37(5):540-6.

12. Wick RR, Holt KE. Benchmarking of long-read assemblers for prokaryote whole genome sequencing. F1000Research. 2019;8:2138.

13. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. PLoS Comput Biol. 2017;13(6):e1005595.

14.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De
Novo Assembler. Curr Protoc Bioinformatics. 2020;70(1):e102.

15.   De Maio N, Shaw LP, Hubbard A, George S, Sanderson ND, Swann J, et al.
Comparison of long-read sequencing technologies in the hybrid assembly of complex
bacterial genomes. Microb Genom. 2019.

16.   Klockgether J, Munder A, Neugebauer J, Davenport CF, Stanke F, Larbig KD, et al.
Genome diversity of Pseudomonas aeruginosa PAO1 laboratory strains. J Bacteriol.
2010;192(4):1113-21.

17.   Chandler CE, Horspool AM, Hill PJ, Wozniak DJ, Schertzer JW, Rasko DA, et al.
Genomic and Phenotypic Diversity among Ten Laboratory Isolates of Pseudomonas
aeruginosa PAO1. J Bacteriol. 2019;201(5).

18.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al.
Versatile and open software for comparing large genomes. Genome Biol. 2004;5(2):R12.

19.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.
2014;30(14):2068-9.

20.   Sereika MK, R.H.; Karst, S.M.; Michaelsen, T.Y.; Soresnes, E.A.; Wollenberg, R.D.;
Albertsen, M. Oxford Nanopore R10.4 long-read sequencing enables near-perfect bacterial
genomes from pure cultures and metagenomes without short-read or reference polishing.
BioRxiv.

11. Figures and tables

Table 1. Sequencing read statistics by sequencing modality and bacterial species.
Note for R.9.4/Kit10 four isolates were plexed and the total data output is a composite of the individual outputs; for the R10.3/Kit12 and R10.4/Kit12 evaluations each isolate extract was initially run separately. The same flowcell was washed and then re-used for the R10.4 evaluation for the *S. aureus* and then *P. aeruginosa* isolates. Finally, the four DNA extracts were also multiplexed on a single R10.4/Kit12 run.

Species	Sequencing modality/sub-group	Total reads	Total bases	N50	Percentage of reads with a phred score of ≥20
E. coli	Illumina	3801912	574088712	151	97.93
	R9.4 (multiplexed run)	353317	2364469570	11705	67.1
	R9.4 (multiplexed run; sup called)	339077	2242222750	11535	70.03
	R10.3 (single extract/run)	1073327	5964466078	9852	79.05
	R10.3 (single extract/run; sup called)	1072758	5936766616	9827	73.5
	R10.4 (single extract/run ; overall)	1174227	6124985330	10507	66.2
	R10.4 (single extract/run; sup called)	1167782	6131556595	10562	79.09
	R10.4 (single extract/run; sup called and duplex reads)	52171	229801689	7274	98.21
	R10.4 (multiplexed	286239	671853044	5327	72.62

	run)				
	R10.4 (multiplexed run; sup called and duplex reads	6447	10999797	3403	98.06
K. pneumoniae	Illumina	3202356	483555756	151	97.45
	R9.4 (multiplexed run)	377192	3646791131	17396	65.23
	R9.4 (multiplexed run; sup called)	361657	3458646526	17157	68.59
	R10.3 (single extract/run)	789562	7772922913	19228	77.29
	R10.3 (single extract/run; sup called)	774119	7658992847	19124	70.24
	R10.4 (single extract/run ; overall)	869853	7481444246	18612	65.83
	R10.4 (single extract/run; sup called)	865400	7495921601	18697	79.79
	R10.4 (single extract/run; sup called and duplex reads)	54177	452672411	16484	98.62
	R10.4 (multiplexed run)	224555	1667146081	15525	72.1
	R10.4 (multiplexed run; sup called and duplex reads	12114	95832563	15245	98.82
P. aeruginosa	Illumina	5299866	800279766	151	97.25

	R9.4 (multiplexed run)	361977	4302642519	21597	66.49
	R9.4 (multiplexed run; sup called)	351155	4138688286	21342	71.55
	R10.3 (single extract/run)	1024134	8524041501	17666	81.81
	R10.3 (single extract/run; sup called)	1017748	8528041241	17683	76.05
	R10.4 (single extract/run ; overall)	556000	5851279980	24126	67.35
	R10.4 (single extract/run; sup called)	638801	6378501910	23860	82.24
	R10.4 (single extract/run; sup called and duplex reads)	22859	261812617	21432	98.58
	R10.4 (multiplexed run)	208693	1412016443	14627	73.91
	R10.4 (multiplexed run; sup called and duplex reads	12468	93018395	14095	98.83
S. aureus	Illumina	9033160	1364007160	151	98.98
	R9.4 (multiplexed run)	40194	725665757	33599	72.67
	R9.4 (multiplexed run; sup called)	39155	699249807	33066	75.51

R10.3 (single extract/run)	1625258	9724520340	14338	82.06
R10.3 (single extract/run; sup called)	1645001	9819093990	14337	78.84
R10.4 (single extract/run ; overall)	950361	7371346901	23339	74.06
R10.4 (single extract/run; sup called)	945421	7382123466	23446	84.24
R10.4 (single extract/run; sup called and duplex reads)	47087	334258567	16366	98.8
R10.4 (multiplexed run)	80512	287957484	14301	80.04
R10.4 (multiplexed run; sup called and duplex reads	3753	12755562	10232	99.08

Table 2. Number of unique contigs by sequencing modality and bacterial species. Using the complete data available (i.e. no subsampling). The first number in each cell represents the number of contigs assembled and matching to the Illumina-corrected reference using dnadiff, the total number of contigs assembled is shown in curved brackets, and the proportion of the reference chromosomal contig covered in square brackets. For *E. coli*, *P. aeruginosa*, *S. aureus* the total number of expected contigs is 1, for *K. pneumoniae* 1 chromosome + 5 plasmids. Orange shading shows absent contigs, and/or incomplete assembly (n >1 contig matching to reference), and/or extra contigs not matching to reference. Green shaded cells denote complete singular contigs which reflect the reference DNA content at 100+/-1%. "-" denotes no relevant contig assembled.

Sequencing modality	Assembler	Plexed Y/N	*E. coli*, chromosome (1) 5231428bp 51% GC	*P. aeruginosa*, chromosome (1) 6264404bp 66.2%GC	*S. aureus*, chromosome (1) 2902619bp 32.8%GC	*K. pneumoniae*, chromosome (1) 57.0%GC	*K. pneumoniae*, pKPN3 plasmid (1) 175879 51.7%GC	*K. pneumoniae*, pKPN4 plasmid (1) 107576bp 53.4%GC	*K. pneumoniae*, pKPN5 (plasmid (1) 88582bp 53.8%GC	*K. pneumoniae*, pKPN6 (plasmid (1) 4259bp 41.4%GC	*K. pneumoniae*, pKPN7 (plasmid (1) 3478bp 45.7%GC
Illumina	SPAdes	Y	226 (326) [98.64%]	115 (152) [99.72%]	86 (150) [98.42%]	117 (312) [98.9%]	41 [79.7%]	45 [100.72%]	22 [82.49%]	1 [102.82%]	1 [103.45%]
R9.4.1 hac+Illumina	Unicycler	Y	1 (1) [100.09%]	1 (1) [100.49%]	1 (1) [100.69%]	1 (7) [100.1%]	1 [100%]	1 [100%]	1 [100%]	2 [100%]	1 [96.15%]
R9.4.1 hac	Canu	Y	1 (1) [100.59%]	1 (1) [101.29%]	1 (1) [103.18%]	1 (24) [100.72%]	1 [131.55%]	1 [172.51%]	1 [100%]	2 [133.5%]	1 [100%]

R9.4.1 sup	Canu	Y	1 (1) [100.65%]	1 (1) [101.18%]	1 (1) [102.43%]	1 (7) [100.96%]	1 [129.41%]	1 [100%]	1 [125.07%]	1 [100%]	1 [100%]
R10.3 hac	Canu	N	1 (6) [100.49%]	1 (2) [101.19%]	1 (3) [102.78%]	1 (8) [100.72%]	1 [100%]	1 [100%]	1 [100%]	1 [100%]	1 [100%]
R10.3 sup	Canu	N	1 (2) [100.6%]	1 (3) [101.13%]	1 (3) [102.85]	1 (8) [100.74%]	1 [100%]	1 [138.13%]	1 [143.94%]	1 [192.58%]	1 [100%]
R10.4 hac	Canu	N	1 (6) [100.62%]	1 (1) [101.06%]	1 (2) [103.6%]	1 (8) [100.8%]	1 [127.89%]	1 [141.81%]	1 [147.6%]	1 [100%]	1 [112.85%]
R10.4 sup	Canu	N	1 (3) [100.61%]	1 (2) [100.1%]	1 (2) [102.33%]	1 (7) [100%]	1 [100.94%]	1 [100%]	1 [145.43%]	1 [100%]	1 [100%]
R10.4 sup duplex	Canu	N	4 (42) [100.12%]	1 (14) [101.5%]	1 (29) [101.72%]	1 (41) [100.1%]	1 [100%]	3 [130.98%]	1 [149.19%]	1 [100%]	1 [100%]
R10.4 hac	Canu	Y	1 (2) [100.42%]	1 (1) [100.87%]	1 (1) [102.01%]	1 (7) [100.82%]	1 [124.01%]	1 [100%]	1 [142.89%]	1 [100%]	1 [100%]
R10.4 sup	Canu	Y	1 (1) [100.48%]	1 (1) [100.17%]	1 (1) [102.29%]	1 (12) [100.61%]	1 [117.3%]	2 [134.06%]	1 [137.68%]	1 [100%]	1 [100%]
R10.4 sup duplex	Canu	Y	-*	23 (25) [99.2%]	-*	15 (25) [99.2%]	1 [80.39%]	2 [97.73%]	1 [112.86%]	1 [94.01%]	1 [100%]
R9.4.1 hac	Flye	Y	1 (1) [100.09%]	1 (1) [100.14%]	1 (1) [100.7%]	1 (4) [100.11%]	1 [75.7%]	1 [103.68%]	1 [100%]	-	-
R9.4.1 sup	Flye	Y	1 (1) [100.09%]	1 (1) [100.47%]	1 (1) [100%]	1 (5) [100.1%]	1 [100%]	1 [99.99%]	1 [100%]	-	-

R10.3 hac	Flye	N	1 (1) [100.10%]	1 (1) [100.44%]	1 (1) [100.69%]	1 (4) [100.1%]	1 [95.83%]	1 [73.13%]	1 [100%]	-	-
R10.3 sup	Flye	N	1 (1) [100.09%]	1 (1) [100.44%]	1 (1) [100.69%]	1 (4) [100.1%]	1 [100%]	1 [100%]	1 [100%]	-	-
R10.4 hac	Flye	N	1 (1) [100.10%]	1 (1) [100.11%]	1 (1) [100.69%]	1 (4) [100.1%]	1 [100%]	1 [100%]	1 [100%]	-	-
R10.4 sup	Flye	N	1 (1) [100.09%]	1 (1) [100.4%]	1 (1) [100.69%]	1 (5) [100.1%]	1 [100%]	1 [98.92%]	1 [99.99%]	-	-
R10.4 sup duplex	Flye	N	1 (3) [100.10%]	1 (1) [100.8%]	1 (2) [100.69%]	1 (7) [100.21%]	1 [94.27%]	1 [102.93%]	1 [101.54%]	-	1 [100%]
R10.4 hac	Flye	Y	1 (1) [100.10%]	1 (1) [100.16%]	1 (1) [100.69%]	1 (5) [100.1%]	1 [100%]	1 [100%]	1 [100%]	-	-
R10.4 sup	Flye	Y	1 (1) [100.00%]	1 (1) [100.48%]	1 (1) [100.69%]	1 (5) [100.11%]	1 [100%]	1 [100%]	1 [100%]	-	-
R10.4 sup duplex	Flye	Y	37 (38) [8.47%]	1 (5) [100.48%]	25 (25) [83.64%]	1 (4) [100.4%]	1 [84.71%]	1 [105.27%]	1 [100%]	-	-

* Insufficient read depth for canu to assemble

Figure 1. Experimental workflow

Figure 2. Read length distributions by (A) modality and (B) by modality and species. Boxplots
reflect median (central line) and IQR (box hinges) values, whiskers the smallest and largest values
1.5*IQR, and dots the outlying points beyond these ranges. Note the y-axis is a log-scale. Median
differences in read length were significant across the whole dataset (Kruskal-wallis, p<0.001); other
significance values represent comparisons with the average read length for R9.4 hac as the reference
category ("ns" - not significant, "****" - p<0.001).

(A)

(B)

Figure 3. Median and modal raw read accuracy (% identity when reads are mapped to the

Illumina-corrected reference) for each of the major nanopore sequencing sequencing modalities,

flowcells/kit and basecalling combinations. Reads matching to the reference with <75% identity

have been excluded. Complete details summarising all accuracies across all modality, flowcell/kit and

basecalling combinations, and stratified by species are represented in Supplementary Table S6.

531 **Figure 4. Number of insertions (panel A) and deletions (panel B) amongst reads mapped to the**
532 **Illumina-corrected reference for all sequencing modalities.**

533 (A)

535 (B)

  Figure 5. Assembly reference coverage percentage (%) by sequencing modality, assembler and

  species. Panel A represents the data for chromosomes and panel B evaluations for the five plasmids

  known to occur in the *K. pneumoniae* reference strain (labeled by their lengths in bp).

  (A)

  (B)

 Figure 6. Assembly accuracy by sequencing modality, assembler and species - SNPs and indels.

Accuracy evaluated on the basis of contig comparisons to Illumina-corrected references using

dnadiff. Accuracy measured as total SNPs (A) and total indels (B), and compared to the Illumina-

corrected reference. NB - SPAdes was only used on Illumina data, and Unicycler hybrid assembly was

only performed on R9.4.1+Illumina data.

(A).

(B)

 Figure 7. Impact of subsampling of long-read datasets on assembly accuracy. Presented here in

each panel for R9.4.1 (top), R10.4 sup (bottom left) and R10.4 sup duplex (bottom right) data only,

stratified by species. NB SPAdes and Unicycler (nanopore+Illumina) data are presented only for R9.4

outputs; Canu assemblies at <10x were not evaluated because the default setting sets a threshold

above this.

(A) *E. coli*

(B) *K. pneumoniae* (chromosome only)

564   (C) *P. aeruginosa*

566   (D) *S. aureus*

Figure 8. Coding sequence (CDS) recovery on the basis of exact CDS (amino acid sequence)
matches with respect to the Prokka-annotated Illumina-corrected reference (chromosome+all
plasmids for _K. pneumoniae_). For long-read data only Flye assemblies have been shown; for R10.3
and R10.4 datasets these were from non-multiplexed evaluations (i.e. only single extracts per
flowcell). For Unicycler, the assembly using R.9.4 hac+Illumina data is shown. The total number of
genes identified correctly is shown as a number at the top of each bar.