Succinate mediates inflammation-induced adrenocortical dysfunction

Ivona Mateska[1]*, Anke Witt[1], Eman Hagag[1], Anupam Sinha[1], Canelif Yilmaz[1], Evangelia Thanou[2], Na Sun[3], Ourania Kolliniati[4], Maria Patschin[1], Heba Abdelmegeed[1], Holger Henneicke[5,6], Waldemar Kanczkowski[1], Ben Wielockx[1], Christos Tsatsanis[4], Andreas Dahl[7], Axel Walch[3], Ka Wan Li[2], Mirko Peitzsch[1], Triantafyllos Chavakis[1], Vasileia Ismini Alexaki[1]*

[1] Institute of Clinical Chemistry and Laboratory Medicine, University Hospital, Technische Universität Dresden, Germany

[2] Center of Neurogenomics and Cognitive Research (CNCR), Department of Molecular and Cellular Neurobiology, Vrije Universiteit, Amsterdam, Netherlands

[3] Research Unit Analytical Pathology, German Research Center for Environmental Health, Helmholtz Zentrum München, Munich, Germany

[4] Department of Clinical Chemistry, Medical School, University of Crete, Heraklion, Greece

[5] Center for Regenerative Therapies, TU Dresden, Technische Universität Dresden, Germany

[6] Department of Medicine III & Center for Healthy Ageing, Technische Universität Dresden, Germany

[7] DRESDEN-concept Genome Center, Center for Molecular and Cellular Bioengineering, Technische Universität Dresden, Germany

***Correspondence**:

Dr. V.I. Alexaki, Institute for Clinical Chemistry and Laboratory Medicine, University Hospital, Technische Universität Dresden, Fetscherstrasse 74, 01307 Dresden, Germany; +49 351 458 16273; VasileiaIsmini.Alexaki@uniklinikum-dresden.de

Dr. I. Mateska, Institute for Clinical Chemistry and Laboratory Medicine, University Hospital, Technische Universität Dresden, Fetscherstrasse 74, 01307 Dresden, Germany, Ivona.Mateska@uniklinikum-dresden.de

Keywords: adrenal gland / LPS / IL-1β / DNMT1 / Succinate dehydrogenase / succinate / glucocorticoids

Acknowledgements

We thank Christine Mund, Denise Kaden and Catleen Conrad for technical assistance. We acknowledge the technical support from the Core Facility Cellular Imaging of the Medical Faculty Carl Gustav Carus, TU Dresden for confocal imaging, and from the Light Microscopy Facility of the CMCB Technology Platform at TU Dresden, for laser microdissection.

Abstract

The hypothalamus-pituitary-adrenal (HPA) axis is activated in response to inflammation leading to increased production of anti-inflammatory glucocorticoids by the adrenal cortex, thereby representing an endogenous feedback loop. However, severe inflammation reduces the responsiveness of the adrenal gland to adrenocorticotropic hormone (ACTH) although the underlying mechanisms are poorly understood. Here, we show by transcriptomic, proteomic and metabolomic analyses that LPS-induced systemic inflammation triggers profound metabolic changes in steroidogenic adrenocortical cells, including downregulation of the TCA cycle and oxidative phosphorylation. Inflammation disrupts the TCA cycle at the level of succinate dehydrogenase (SDH) leading to succinate accumulation and disturbed steroidogenesis. Mechanistically, IL-1β reduces SDHB expression through upregulation of DNA methyltransferase 1 (DNMT1) and methylation of the SDHB promoter. Consequently, increased succinate levels impair oxidative phosphorylation and ATP synthesis, leading to reduced steroidogenesis. Together, we demonstrate that the IL-1β-DNMT1-SDHB-succinate axis disrupts steroidogenesis. Our findings not only provide a mechanistic explanation for the adrenal dysfunction in severe inflammation but also a potential target for therapeutic intervention.

Introduction

Stress triggers the hypothalamic–pituitary–adrenal (HPA) axis, i.e. the release of corticotropin-releasing hormone (CRH) from the hypothalamus, followed by adrenocorticotropic hormone (ACTH) secretion from the anterior pituitary, which stimulates the synthesis of glucocorticoid hormones in the adrenal cortex, primarily cortisol in humans and corticosterone in rodents [1–3]. Similarly to any other stress stimulus, inflammation activates the HPA axis leading to increased glucocorticoid release, which is required to restrain the inflammatory response [4-8]. Adrenalectomized rodents show increased mortality after induction of systemic inflammation, while glucocorticoid administration increases survival [9,10]. Essentially, severe inflammation in sepsis is associated with impaired adrenal gland function [11–15] but the mechanisms remain poorly understood.

In immune cells, such as macrophages, dendritic cells and T cells, inflammation triggers cellular metabolic reprograming, enabling the cells to meet the increased demands for fast energy supply and anabolic processes [16–18]. How inflammation may affect cellular metabolism in other cell types and how this affects their function is less explored. Here we show that LPS-induced inflammation profoundly changes the cellular metabolism of steroidogenic adrenocortical cells, perturbing the TCA cycle at the level of succinate dehydrogenase B (SDHB). This is coupled to succinate accumulation, which impairs oxidative phosphorylation and leads to reduced steroidogenesis. Mechanistically, IL-1β inhibits *SDHB* expression through DNA methyltransferase 1 (DNMT1)-dependent DNA methylation of the *SDHB* promoter.

Methods

Animal experiments

Eight to twelve-week-old male C57BL/6J were injected i.p. with 1 mg/kg LPS (LPS-EB Ultrapure; InVivoGen) or PBS, and sacrificed after 6 h (for gene expression analyses) or 24 h (for all other analyses). *Irg1$^{-/-}$* and littermate control mice were injected with 3 mg/kg LPS and sacrificed after 16 h.

Laser capture microdissection of adrenal cortex

Adrenal glands frozen in liquid nitrogen were cut in 25 – 30 μm thick sections, mounted on polyethylene naphthalate (PEN) membrane slides (Zeiss), dehydrated in increasing concentrations of ice-cold ethanol (75 %, 95 %, 100 %) for 45 sec each, and air-dried at room temperature (RT). Laser Capture Microdissection was performed with a Zeiss PALM MicroBeam LCM system. The adrenal cortex from 8 to 12 sections was microdissected and the tissue was collected on Adhesive Caps (Zeiss).

Bioinformatic analysis of RNA-Seq data

For transcriptome mapping, strand-specific paired-end sequencing libraries from total RNA were constructed using TruSeq stranded Total RNA kit (Illumina Inc). Sequencing was performed on an Illumina HiSeq3000 (1x75 basepairs). Low quality nucleotides were removed with the Illumina fastq filter (http://cancan.cshl.edu/labmembers/gordon/fastq_illumina_filter/) and reads were further subjected to adaptor trimming using cutadapt [19]. Alignment of the reads to the Mouse genome was done using STAR Aligner [20] using the parameters: "–runMode alignReads –outSAMstrandField intronMotif –outSAMtype BAM SortedByCoordinate --readFilesCommand zcat". Mouse Genome version GRCm38 (release M12 GENCODE) was used for the alignment. The parameters: 'htseq-count -f bam -s reverse -m union -a 20', HTSeq-0.6.1p1 [21] were used to count the reads that map to the genes in the aligned sample files. The GTF file (gencode.vM12.annotation.gtf) used for read quantification was downloaded from Gencode (https://www.gencodegenes.org/mouse/release_M12.html). Gene centric differential

expression analysis was performed using DESeq2_1.8.1 [22]. The raw read counts for the genes across the samples were normalized using 'rlog' command of DESeq2 and subsequently these values were used to render a PCA plot using ggplot2_1.0.1 [23].

Pathway and functional analyses were performed using GSEA [24] and EGSEA [25]. GSEA is a stand-alone software with a GUI. To run GSEA, a ranked list of all the genes from DESeq2 based calculations was created by taking the -log10 of the p-value and multiplying it with the sign the of the fold change. This ranked list was then queried against Molecular Signatures Database (MsigDB), Reactome, KEGG and GO based repositories. EGSEA is an R/Bioconductor based command-line package. For doing functional analyses using EGSEA a differentially expressed list of genes with parameters log2foldchange > 0.3 and padj < 0.05 was used. Same database repositories as above were used for performing the functional analyses. For constructing pathway-specific heatmaps, the "rlog-normalized" expression values of the significantly expressed genes (padj < 0.05) were mapped on to the KEGG and GO pathways. These pathway-specific expression matrices were then scaled using z-transformation. The resulting matrices were visually rendered using MORPHEUS.

Cell sorting

The adrenal cortex was separated from the medulla under a dissecting microscope and was digested in 1.6 mg/ml collagenase I (Sigma-Aldrich) and 1.6 mg/ml BSA in PBS, for 25 min at 37 °C while shaking at 900 rpm. The dissociated tissue was passed through a 22 G needle and 100 μm cell strainer and centrifuged at 300 g for 5 min at 4 °C. The cell suspension was washed in MACS buffer (0.5 % BSA, 2 mM EDTA in PBS) and CD31$^+$ and CD45$^+$ cells were sequentially positively selected using anti-CD31 and anti-CD45 MicroBeads (Miltenyi Biotec), respectively, according to manufacturer's instructions. Briefly, pelleted cells resuspended in 190 μl MACS buffer, were mixed with 10 μl anti-CD31 MicroBeads, incubated for 15 min at 4 °C, washed with 2 ml MACS buffer and centrifuged at 300 g for 10 min at 4 °C. Then, the cell pellet was resuspended in 500 μl MACS buffer, applied onto MS Column placed on MACS Separator, and the flow-through (CD31$^-$ cells) was collected. CD31$^+$ cells were

positively sorted from the MS Columns. The flow-through was centrifuged at 300 g for 5 min at 4 °C, and the pelleted cells were subjected to the same procedure using anti-CD45 MicroBeads, collecting the flow-through containing CD31$^-$CD45$^-$ adrenocortical cells. CD45$^+$ cells were positively sorted from the MS Columns.

MS/MS Proteomic analysis

CD31$^-$CD45$^-$ adrenocortical cells were sorted and snap-frozen. Samples were randomized and a gel-based sample preparation protocol was followed [26]. In brief, cell pellets were resuspended in SDS loading buffer and 30% acrylamide, boiled at 98 °C for 6 min, and 5 μg protein per sample were separated in 10% SDS gels (SurePAGE Bis-Tris gels, GenScript) for approximately 10 min at 120 V. The gels were fixed in 50 % (v/v) ethanol and 3 % (v/v) phosphoric acid and briefly stained with Colloidal Coomassie Blue. Sample containing lanes were sliced and cut into blocks of approximately 1 mm^3, destained in 50 mM NH$_4$HCO$_3$ and 50 % (v/v) acetonitrile, dehydrated using 100 % acetonitrile, and rehydrated in 50 mM NH$_4$HCO$_3$ containing 10 μg / ml trypsin (sequence grade; Promega). After incubation overnight at 37 °C peptides were extracted and collected in a new tube, dried using a SpeedVac (Eppendorf), and stored at - 20 °C until LC-MS analysis. Peptides were dissolved in 0.1 % formic acid, and 75 ng were loaded into EvoTips (EV2003, EvoSep) and washed according to manufacturer's guidelines. The samples were run on a 15 cm x 75 μm, 1.9 μm Performance Column (EV1112, EvoSep) using the Evosep one liquid chromatography system with the 30 samples per day program. Peptides were analyzed by the TimsTof pro2 mass spectrometer (Bruker) with the diaPASEF method [27]. Data were analyzed using DIA-NN. The fasta database used was uniport mouse_UP000000589_10090. Deep learning was used to generate the in silico spectral library. Output was filtered at 0.01 FDR [28]. The Mass Spectrometry Downstream Analysis Pipeline (MS-DAP) (version beta 0.2.5.1) (https://github.com/ftwkoopmans/msdap) was used for quality control and candidate discovery [29]. Differential abundance analysis between groups was performed on log transformed protein abundances. Empirical Bayes

moderated t-statistics with multiple testing correction by FDR, as implemented by the eBayes functions from the limma R package, was used as was previously described [30].

Bioinformatics analysis of proteomics data

From the proteomics data, the missing data was imputed using the "impute" command running of "DEP" [31] package in R/Bioconductor [32] environment. The imputation was performed using "knn" function. The resultant imputed matrix was used for further analyses. Pathway and functional analyses were performed using GSEA [23] and EGSEA [24]. CLI version of GSEA v4.1 was run using the imputed matrix. Different pathway sets from Molecular Signatures Database (MsigDB) v7.2 like, HALLMARK, Biocarta, Reactome, KEGG, GO and WIKIPATHWAYS were queried for functional enrichment. Geneset permutations were performed 1,000 times to calculate the different statistical parameters. For doing functional analyses using EGSEA, imputed matrix was used. Same database repositories as above were used for performing the functional analyses.

Quantitative RT – PCR

Total RNA was isolated from frozen adrenal glands with the TRI Reagent (MRC) after mechanical tissue disruption, extracted with chloroform and the NucleoSpin RNA Mini kit (Macherey-Nagel). Total RNA from sorted cells was isolated with the Rneasy Plus Micro Kit (Qiagen) according to manufacturer's instructions. cDNA was synthesized with the iScript cDNA Synthesis kit (Biorad) and gene expression was determined using the SsoFast Eva Green Supermix (Bio-Rad), with a CFX384 real-time System C1000 Thermal Cycler (Bio-Rad) and the Bio-Rad CFX Manager 3.1 software. The relative gene expression was calculated using the ΔΔCt method, *18S* was used as a reference gene. Primers are listed in Table 5.

Cell culture and in vitro treatments

CD31$^-$CD45$^-$ adrenocortical cells were plated on 0.2 % gelatin-coated wells of 96-well plates in DMEM/F12 medium supplemented with 1 % fetal bovine serum (FBS), 50 U/mL

penicillin and 50 μg/mL streptomycin (all from Gibco), and let to attach for an hour before treatments. Cells from both adrenal glands from each mouse were pooled together and plated in 2 wells of a 96-well plate. Mouse adrenal explants were dissected from surrounding fat and left in DMEM/F12 medium with 1 % FBS, 50 U/mL penicillin and 50 μg/mL streptomycin for an hour before treatments. NCI-H295R cells (obtained from ATCC) were maintained in DMEM/F12 medium supplemented with 2.5 % Nu-Serum type I (Corning), 1 % Insulin Transferrin Selenium (ITS; Gibco), 50 U/mL penicillin and 50 μg/mL streptomycin.

Cells or explants were treated with DMM (20 mM; Sigma-Aldrich), DES (5 mM; Sigma-Aldrich), FCCP (1 μM; Agilent Technologies), OM (500 nM; Agilent Technologies), AG-221 (10 μM; Selleckchem), 4-OI (125 μM; Cayman Chemical), IL-1β (20 ng/ml, PeproTech), IL-6 (20 ng/ml, PeproTech), TNFα (20 ng/ml, PeproTech), LPS (1 μg/ml; InVivoGen), ACTH (100 ng/ml; Sigma-Aldrich) or Forskolin (10 μM; Sigma-Aldrich). siRNA transfections were done with ON-TARGETplus SMARTpool siRNA against *SDHB* (10 nM), *Sdhb* (30 nM), *Idh2* (30 nM) or *Dnmt1* (30 nM) (all from Horizon Discovery), with Lipofectamine RNAiMAX transfection reagent (Invitrogen), using a reverse transfection protocol per manufacturer's instructions.

Steroid hormone measurement

Steroid hormones were analyzed by LC-MS/MS in cell culture or explant supernatants as described previously [33]. Fifty to hundred μL cell culture supernatants were extracted by solid phase extraction using positive pressure, followed by a dry-down under gentle stream of nitrogen. Residues were reconstituted in 100 μL of the initial LC mobile phase and 10 μL were injected for detection by the triple quadrupole mass spectrometer in multiple reaction-monitoring scan mode using positive atmospheric pressure chemical ionization. Quantification of steroid concentrations was done by comparisons of ratios of analyte peak areas to respective peak areas of stable isotope labeled internal standards obtained in samples to those of calibrators.

Measurement of TCA cycle metabolites

TCA cycle metabolites were determined by LC-MS/MS as described before [34]. Itaconate was included in the existing LC-MS/MS method using multi-reaction monitoring (MRM)-derived ion transition of 128.9->85.1. For quantification of itaconate ratios of analyte peak areas to respective peak areas of the stable isotope labeled internal standard (itaconic acid-[13]C5; Bio-Connect B.V., The Netherlands; MRM transition 133.9->89.1) obtained in samples were compared to those of calibrators.

MALDI-FT-ICR-MSI

Tissue preparation steps for MALDI imaging mass spectrometry (MALDI-MSI) analysis was performed as previously described [35,36]. Frozen mouse adrenals were cryosectioned at 12 μm (CM1950, Leica Microsystems, Wetzlar, Germany) and thaw-mounted onto indium-tin-oxide coated conductive slides (Bruker Daltonik, Bremen, Germany). The matrix solution consisted of 10 mg/ml 1,5-Diaminonaphthalene (Sigma-Aldrich, Germany) in water/acetonitrile 30:70 (v/v). SunCollectTM automatic sprayer (Sunchrom, Friedrichsdorf, Germany) was used for matrix application. The MALDI-MSI measurement was performed on a Bruker Solarix 7T FT-ICR-MS (Bruker Daltonik, Bremen, Germany) in negative ion mode using 100 laser shots at a frequency of 1000 Hz. The MALDI-MSI data were acquired over a mass range of m/z 75-250 with 50 μm lateral resolution. Following the MALDI imaging experiments, the tissue sections were stained with hematoxylin and eosin (H&E) and scanned with an AxioScan.Z1 digital slide scanner (Zeiss, Jena, Germany) equipped with a 20x magnification objective. After the MALDI-MSI measurement, the acquired data underwent spectra processing in FlexImaging v. 5.0 (Bruker Daltonik, Bremen, Germany) and SciLS Lab v. 2021 (Bruker Daltonik, Bremen, Germany). The mass spectra were root-mean-square normalized. MS Peak intensity of isocitrate and succinate of adrenal cortex regions were exported and applied for relative quantification analysis.

ATP measurement

Total ATP was measured in adrenal glands using the ATP Assay Kit (ab83355, Abcam). Briefly, adrenal glands were collected, washed with PBS and immediately homogenized in 100 μl ATP assay buffer. Samples were cleared using the Deproteinizing Sample Preparation Kit – TCA (ab204708, Abcam). Samples were incubated for 30 min with the ATP reaction mix and fluorescence (Ex / Em = 535 / 587 nm) was measured using the Synergy HT microplate reader. The recorded measurements were normalized to the weight of the adrenal gland.

ROS measurement

ROS was detected using the DCFDA/H2DCFDA Cellular ROS Detection Assay Kit (ab113851, Abcam). NCI-H295R cells were plated at 80,000 cells / well in 96-well plate with black walls and clear bottom (Corning) and were incubated with 20 μM DCFDA Solution for 45 min at 37 °C in dark. Fluorescence (Ex / Em = 485 / 535 nm) was measured using the Synergy HT microplate reader.

ADP / ATP ratio measurement

Intracellular ATP / ADP ratio was determined with the ADP / ATP Ratio Assay Kit (MAK135, Sigma-Aldrich). NCI-H295R cells were plated at 80,000 cells / well in 96-well plate with white flat bottom wells (Corning). Luminescence was measured using the Synergy HT microplate reader.

NADP/NADPH measurement

Intracellular NADP/NADPH ratio was measured with the NADP/NADPH Assay Kit (Fluorometric) (ab176724, Abcam). NCI-H295R cells were plated at 5×10^6 cells / 10 cm – diameter dish. Fluorescence (Ex / Em = 540 / 590 nm) was measured using the Synergy HT microplate reader. NADPH in adrenal tissue homogenates were analyzed by LC-MS/MS using an adapted method previously described [37].

Enzyme activity measurement

SDH and IDH activities were measured using respective colorimetric assay kits (MAK197, Sigma-Aldrich, ab102528, Abcam). Cortices from both adrenal glands of each mouse were pooled and processed together. Absorbance (at 600 nm for SDH or 450 nm for IDH) was detected using the Synergy HT microplate reader.

Seahorse assay

OCR measurements were performed with a Seahorse XF96 Analyzer (Agilent Technologies). NCI-H295R cells were plated at 80,000 cells / well in 0.2 % gelatin-precoated XF96 cell culture microplate (Agilent). The experimental medium used was XF Base Medium supplemented with glucose (10 mM), pyruvate (1 mM) and glutamine (2 mM).

Measurement of mitochondrial load and membrane potential

The adrenal cortex was digested and dissociated cells were incubated with Mitotracker Green (0.25 μM; Invitrogen), TMRE (2.5 μM; Thermofisher), CD31-PeCy7 (1:100; eBioscience) and CD45-PeCy7 (1:100; eBioscience) for 30 min in FACS buffer (0.5 % BSA, 2 mM EDTA in PBS) at 37°C in dark. Live cells were selected by Hoechst staining. NCI-H295R cells were incubated with MitotrackerGreen (100 nM) and TMRE (100 nM) for 30 min at 37 °C in dark. FACS was performed using LSR Fortessa X20 flow cytometer and data were analyzed with the FlowJo software.

Western blotting

Cells were lysed with 10 mM Tris-HCl, pH7.4 + 1 % SDS + 1 mM sodium vanadate, cell lysates were centrifuged at 16,000 g for 5 min at 4 °C, supernatants were collected and total protein concentration was measured using Pierce BCA Protein Assay Kit (Thermo Scientific). Gel electrophoresis was performed according to standard protocols [38]. Protein samples were prepared with 5x Reducing Laemmli buffer, denatured at 95 °C for 5 min and loaded on a 10 % acrylamide gel (Invitrogen) for sodium dodecyl sulfate polyacrylamide gel electrophoresis

(SDS-PAGE). PageRuler Prestained Protein Ladder (Thermo Fisher Scientific) was used as a protein size ladder. The separated proteins were transferred on Amersham Protran nitrocellulose membrane (GE Healthcare Lifescience). After blocking with 5 % skimmed milk in TBS-T (0.1 % Tween-20 (Sigma-Aldrich) in 1x Tris-buffered saline) for 1 hour at RT, membranes were incubated overnight at 4 °C with anti-SDHB (1:1000; Sigma-Aldrich, #HPA002868), anti-DNMT1 (1:1000; Cell Signaling, #5032) or anti-Tubulin (1:3000; Sigma-Aldrich, T5186), diluted in 5 % BSA in TBS-T. After washing, membranes were incubated for 1 hour at RT with secondary antibodies: goat anti-rabbit IgG HRP- conjugated (1:3000; Jackson ImmunoResearch) or goat anti-mouse IgG HRP-conjugated (1:3000; Jackson ImmunoResearch), diluted in 5 % skimmed milk in TBS-T. The signal was detected using the Western Blot Ultra-Sensitive HRP Substrate (Takara) and imaged using the Fusion FX Imaging system (PeqLab Biotechnologie).

DNA methylation measurement

Genomic DNA from $2x10^6$ NCI-H295R cells was isolated with the Quick-DNA Miniprep Kit (Zymo research). Bisulfite treatment was performed using the EZ DNA Methylathion™ Kit (Zymo Research), following the manufacturer's protocol. For each sample 500 ng genomic DNA was used, bisulfite treated for 14h in the dark and, after a desulphonation and cleaning step, eluted in 10 μL nuclease-free water. The SDHB promoter region was amplified with primers for a methylated and a non-methylated sequence (listed in Table 5), using the Qiagen Multiplex PCR Kit. Equal amount of DNA not treated with bisulfite was amplified as a loading control. The PCR products were then electrophoresed on 3 % agarose gel and visualized under UV illumination using the Fusion FX Imaging system (Vilber). The ratio of methylated to non-methylated DNA was calculated after gel intensity quantification in Fiji / ImageJ.

Immunofluorescent staining

Adrenal glands cleaned from surrounding fat tissue were fixed in 4 % PFA in PBS, washed overnight in PBS, cryopreserved in 30 % sucrose (AppliChem GmbH) in PBS overnight

at 4°C, embedded in O.C.T. Compound (Tissue-Tek) and frozen at −80 °C. Each adrenal gland was cut into 8 µm thick serial sections. Before staining, adrenal sections were pre-warmed at RT for 30 min and antigen retrieval was performed by boiling in citrate buffer (pH 6) for 6 min. Adrenal sections were washed with PBS, permeabilized with 0.1 % Triton X-100 in PBS for 20 min, treated with TrueBlack Lipofuscin Quencher (1:40 in 70 % ethanol; Biotium) for 30 sec to reduce autofluorescence and blocked in Dako Protein Block, serum-free buffer for 1 hour at RT. Then, sections were incubated overnight at 4 °C with primary antibodies, washed with PBS, and incubated for 1 hour at RT with the secondary antibodies together with DAPI (1:5,000; Roche), all diluted in Dako Antibody Diluent. Antibodies and dyes used were: anti-SDHB (1:300; Sigma-Aldrich, #HPA002868), anti-IDH2 (1:50; Sigma-Aldrich, #HPA007831), anti-SF-1 (1:100; TransGenic Inc. #KO610), Lectin Esculentum DyLight488 (1:300; Vector Laboratories, #DL-1174), 4-Hydroxynonenal (4-HNE, 1:200; Abcam, # ab48506), Alexa Fluor 555 donkey anti-rabbit (1:300; Life Technologies, #A-31572), Alexa Fluor 647 chicken anti-rat (1:300; Invitrogen, #A21472) and Alexa Fluor 555 donkey anti-mouse (1:300; Invitrogen, #A31570). After washing with PBS, cryosections were mounted with Fluoromount (Sigma-Aldrich), covered with 0.17 mm cover glass, fixed with nail polish and kept at 4 °C until imaging.

Image acquisition and image analysis

Z-series microscopic images for SDHB and IDH2 staining were acquired on Zeiss LSM 880 inverted confocal microscope (Zeiss, Jena, Germany), illuminated with laser lines at 405 nm, 488 nm, 561 nm and 633 nm, and detected by two photomultiplier tube detectors. EC Plan-Neofluoar objective with 40x magnification, 1.30 numerical aperture and M27 thread, working with an oil immersion medium Immersol 518 F, was used. Microscopic images of SF-1 and 4-HNE stainings were acquired with an Axio Observer Z1/7 inverted microscope with Apotome mode (Zeiss, Jena, Germany), illuminated with LED-Module 385 nm and 567 nm, on a Plan-Apochromat objective with 10x magnification, 0.45 numerical aperture and M27 thread. Laser power, photomultiplier gain and pinhole size were set for each antibody individually and kept constant for all image acquisitions. For each condition, at least 3 view-fields were imaged per

tissue section. Images were acquired with the ZEN 3.2 blue edition software, and processed and quantified with the Fiji / ImageJ software on maximum intensity Z-projection images.

Statistical analysis

The statistical analysis and data plotting were done with the GraphPrism 7 software. The statistical tests used are described in each figure legend, p < 0.05 was set as a significance level.

Graphical design

Figure 7 was created with Biorender.com.

Results

1. Metabolic reprograming of the adrenal cortex in inflammation.

To explore inflammation-induced alterations in the adrenal cortex, we performed RNA-Seq in microdissected adrenal cortices from mice treated for 6 h i.p. with 1 mg / kg LPS or PBS, which revealed 2,609 differentially expressed genes, out of which 1,363 were down- and 1,246 were upregulated (Fig. 1a). Gene set enrichment analysis (GSEA) using the Molecular Signatures Database (MSigDB) hallmark gene set collection [39] showed a significant enrichment of inflammatory response-related gene sets in the adrenal cortex of LPS-treated mice (Fig. 1b). In acute inflammation, leukocytes infiltrate the adrenal cortex [7, 9] and resident macrophages are activated [40,41]. In order to delineate the inflammatory response in the adrenocortical steroidogenic cells, CD31⁻CD45⁻ cells were sorted: enrichment in steroidogenic cells was evidenced by high *Star* expression (Suppl. Fig. 1a), and purity was verified by the absence of *Cd31* and *Cd45* expression (Suppl. Fig. 1b,c). Proteomic analysis in the sorted CD31⁻CD45⁻ adrenocortical cell population and GSEA of GO terms confirmed the enrichments of innate immune response-related proteins in adrenocortical cells of LPS-injected mice (Fig. 1c), suggesting that steroidogenic adrenocortical cells respond to inflammatory stimuli.

LPS treatment leads to increased plasma corticosterone levels [7, 9, 10]. Numerous studies have shown that elevated glucocorticoid levels are primarily driven by activation of the HPA axis, and coincide with increased circulating ACTH levels [7, 9, 10]. This is accompanied by increased expression of genes related to steroid biosynthesis [42]. We confirmed an increased expression of the cholesterol transporter *Star* [43] and the terminal enzyme for glucocorticoids synthesis *Cyp11b1* [2] in adrenocortical cells of LPS mice (Suppl. Fig. 2a,b). However, the expression of other steroidogenic enzymes, such as *3βHSD* and *Cyp21a1* was reduced, while *Cyp11a1* remained unchanged (Suppl. Fig. 2c-e). Similarly, protein levels of the transcription factor SF-1, a key inducer of steroidogenesis [44], were somewhat reduced after LPS injection (Suppl. Fig. 2f). Therefore, the observed changes in plasma glucocorticoid levels which accompany inflammation cannot be solely explained by the transcriptional changes in steroidogenic enzymes.

Next, we explored the cell metabolic changes induced by LPS in the adrenal cortex. By GSEA of the RNA-Seq data we observed negative regulation of gene sets related to carbohydrate metabolism in the adrenal cortex of LPS-injected mice (Fig. 1d). Proteomic analysis was performed in CD31⁻CD45⁻ adrenocortical cells (Suppl. Fig. 1) to examine the effects of inflammation on the metabolism of specifically steroidogenic adrenocortical cells, thus evading the well-described inflammation-induced metabolic changes in immune cells [16-18]. Similarly to the RNA-Seq data, GSEA of the proteomic data showed significant negative enrichment of proteins associated with carbohydrate metabolism in the steroidogenic cells (Fig. 1e). EGSEA pathway analysis of the RNA-Seq and proteomic data revealed that TCA cycle, oxidative phosphorylation, tyrosine metabolism, fatty acid degradation, D-Glutamine and D-glutamate metabolism, glutathione metabolism and other metabolic pathways were significantly enriched among the downregulated genes and proteins in the adrenal cortex and in steroidogenic cells of LPS mice (Table 1, Table 2).

Fig. 1.

LPS-induced inflammation changes the transcriptional and proteomic profile of the adrenal cortex. **a** Volcano plot showing differentially expressed genes in the microdissected adrenal gland cortex of mice treated for 6 h with PBS or LPS. **b** GSEA for immune pathways in the adrenal cortex of LPS- versus PBS mice. **c** GSEA for proteins associated with the innate immune response in CD31⁻CD45⁻ adrenocortical cells of mice treated for 24 h with PBS or LPS. **d** RNA-Seq based GSEA for carbohydrate metabolism in the adrenal cortex of LPS- versus PBS mice. **e** GSEA for proteins associated with carbohydrate metabolism in CD31⁻CD45⁻ adrenocortical cells of LPS versus PBS mice. NES: normalized enrichment score. **a**,**b**,**d** n = 3 mice per group, **c,e** n = 6 mice per group, padj < 0.05 was used as a cut-off for significance.

2. Inflammation disrupts the TCA cycle in adrenocortical cells at the levels of isocitrate dehydrogenase and succinate dehydrogenase.

Inflammation downregulates the TCA cycle and oxidative phosphorylation in inflammatory activated macrophages [16], however little is known about inflammation-induced metabolic changes in other cell types. We show that TCA cycle-related gene expression was downregulated in the adrenal cortex of LPS-treated mice (Fig. 2a,b, Table 1). Expression of genes encoding key TCA cycle enzymes, including succinate dehydrogenase *Sdhb* and *Sdhc*, isocitrate dehydrogenase 2 and 3 (*Idh2* and *Idh3b*) and malate dehydrogenase 1 (*Mdh1*), was reduced in the adrenal cortex of LPS-injected mice (Fig. 2b). Proteomic GSEA confirmed the TCA cycle downregulation in steroidogenic adrenocortical cells of LPS mice (Fig. 2c, Table 2). Accordingly, CD31⁻CD45⁻ adrenocortical cells from LPS mice displayed reduced *Idh1*, *Idh2*, *Sdhb* and *Sdhc* expression (Fig. 2d,e) and LPS treatment attenuated the IDH and SDH enzymatic activities in the adrenal cortex (Fig. 2f,g). Additionally, immunofluorescent staining showed that IDH2 and SDHB proteins are highly expressed in SF-1⁺ (steroidogenic) cells (Fig. 2h,i). In endothelial and immune cells of the adrenal cortex of LPS mice, *Idh1* and *Idh2* gene expression was reduced, *Sdhb* gene expression was increased, while expression of *Sdhc* was unaltered (Suppl. Fig. 3a,b). Collectively, these data indicate that the reduced activity of SDH

and IDH in the adrenal cortex of LPS-treated mice is mainly due to their downregulated expression in steroidogenic adrenocortical cells.

Fig. 2

Systemic inflammation disrupts the TCA cycle in the adrenal cortex. **a,b** Transcriptome analysis in the microdissected adrenal gland cortex of mice treated for 6 h with PBS or LPS (n = 3 mice per group). **a** GSEA for TCA cycle genes. **b** Heatmap of differentially expressed TCA cycle genes (padj < 0.05). **c** GSEA analysis for TCA cycle proteins in CD31⁻CD45⁻ adrenocortical cells of mice treated for 24 h with PBS or LPS (n = 6 mice per group). **d,e** mRNA expression of *Idh1*, *Idh2*, *Sdhb* and *Sdhc* in adrenocortical CD31⁻CD45⁻ cells of mice treated

for 6 h with PBS or LPS (n = 8 mice per group, shown is one from two experiments). **f,g** Quantification of IDH and SDH activities in the adrenal cortex of mice treated for 24 h with LPS or PBS (n = 6 mice per group). Values are normalized to the total protein amount in the adrenal cortex. **h,i** Immunofluorescence images of the adrenal gland, stained for IDH2 (red) or SDHB (red), SF-1 (magenta), Isolectin (staining endothelial cells, green), and DAPI (blue). Scale bar, 30 μm. **j-o** TCA cycle metabolites (isocitrate, α-ketoglutarate, succinate, fumarate) were measured by LC–MS/MS in adrenal glands of mice 24 h after injection with PBS or LPS (n = 4 mice per group, shown is one from two experiments). **p,q** MALDI-MSI for isocitrate and succinate in the adrenal cortex of mice treated for 24 h with PBS or LPS (n = 3 mice per group). Representative images and quantifications are shown. Scale bar, 500 μm. Data are presented as mean ± s.d., and analyzed with Mann-Whitney test. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

In order to confirm that inflammation disrupts the TCA cycle in adrenocortical cells, we profiled the changes in metabolite levels in the adrenal glands of PBS- and LPS-treated mice using liquid chromatography – tandem mass spectrometry (LC-MS/MS). The levels of isocitrate and succinate, as well as the ratios of isocitrate / α-ketoglutarate and succinate / fumarate were increased in the adrenal glands of LPS-treated mice (Fig. 2j-o). Furthermore, MALDI MS–Imaging confirmed the increased levels of isocitrate and succinate in the adrenal cortex of LPS mice (Fig. 2p,q). These data collectively demonstrate that inflammation disrupts IDH and SDH activities and increases the levels of their substrates isocitrate and succinate in adrenocortical cells.

3. **Inflammation reduces oxidative phosphorylation and increases oxidative stress in the adrenal cortex.**

Next, we investigated how inflammation affects mitochondrial oxidative metabolism in adrenocortical cells. GSEA of the RNA-Seq and proteomic data in the adrenal cortex and CD31[-] CD45[-] adrenocortical cells, respectively, revealed that oxidative phosphorylation was

significantly enriched among the downregulated genes (Fig. 3a) and proteins (Fig. 3b), and expression of a large number of oxidative phosphorylation-associated genes was reduced in the adrenal cortex of LPS mice (Fig. 3c). In accordance, ATP levels were reduced in the adrenal gland (Fig. 3d) and the mitochondrial membrane potential of CD31⁻CD45⁻ adrenocortical cells was decreased in mice treated with LPS (Fig. 3e). In pro-inflammatory macrophages, a TCA cycle 'break' at the level of SDH is associated with repurposing of mitochondria from oxidative phosphorylation-mediated ATP synthesis to ROS production [45]. EGSEA pathway analysis showed that upon LPS treatment several pathways involved in the regulation of and the cellular response to oxidative stress in the adrenal cortex were enriched at mRNA (Table 3) and protein level (Table 4). This was confirmed by increased 4-hydroxynonenal (4-HNE) staining, indicating higher oxidative stress-associated damage in the adrenal cortex of LPS-treated mice (Fig. 3f). Antioxidant defense mechanisms are particularly important in the adrenal cortex, since electron leakage through the reactions catalyzed by CYP11A1 and CYP11B1 during glucocorticoid synthesis contributes significantly to mitochondrial ROS production [46]. Cells neutralize ROS to maintain their cellular redox environment by using the reducing equivalents NADPH and glutathione [47]. In addition, NADPH serves as a co-factor for mitochondrial steroidogenic enzymes [48]. NADPH levels and glutathione metabolism-related gene expression were significantly decreased in the adrenal glands of LPS mice (Fig. 3g,h, Table 1, Table 2). These findings collectively suggest that inflammation in the adrenal cortex is associated with increased oxidative stress, perturbed mitochondrial oxidative metabolism, reduced anti-oxidant capacity and increased ROS production.

Fig. 3

Oxidative phosphorylation is reduced and oxidative stress is increased in the adrenal cortex of LPS treated mice. **a** GSEA for oxidative phosphorylation-related genes in the adrenal cortex of mice treated for 6 h with PBS or LPS (n = 3 mice per group). **b** GSEA for oxidative phosphorylation-associated proteins in CD31⁻CD45⁻ adrenocortical cells of mice treated for 24 h with PBS or LPS (n = 6 mice per group). **c** Heatmap of differentially expressed genes related to oxidative phosphorylation (padj < 0.05). **d** Measurement of ATP in adrenal glands of mice

treated for 24 h with PBS or LPS (n = 10-11 mice per group). **e** Measurement of mitochondrial membrane potential by TMRE staining and mitochondrial load by Mitotracker Green FM in CD31$^-$CD45$^-$adrenocortical cells of PBS or LPS mice. Data are presented as ratio of the median fluorescence intensities of TMRE to Mitotracker Green FM (n = 6 mice per group). **f** Representative immunofluorescence images of adrenal gland sections from PBS- and LPS-treated mice (24 h post-injection), stained for 4-HNE (magenta) and DAPI (blue). Scale bar, 300 μm. Quantification of the mean fluorescence intensity of 4-HNE staining in the adrenal cortex of PBS- or LPS-treated mice (n = 6 mice per group). **g** NADPH measurement by LC–MS/MS in adrenal glands of mice treated with PBS or LPS for 24 h (n = 8 mice per group). Data are given as observed peak area intensities of NADPH. **h** GSEA for glutathione metabolism of RNA-Seq data in the adrenal cortex of LPS versus PBS mice. Data present mean ± s.d. and were analyzed with Mann-Whitney test. *p < 0.05, **p < 0.01.

4. Increased succinate levels impair mitochondrial metabolism and steroidogenesis in adrenocortical cells.

SDH is complex II of the electron transport chain (ETC), coupling succinate oxidation with the respiratory chain [49]. Inhibition of SDH function with dimethyl malonate (DMM), which is hydrolyzed to the competitive SDH inhibitor malonate [24/50], or treatment of adrenocortical cells with the cell-permeable succinate analog diethyl succinate (DES) increased the amount of succinate and the succinate / fumarate ratio in adrenal gland explants (Fig. 4a) and human adrenocortical carcinoma cells NCI-H295R (Fig. 4b) and decreased the oxygen consumption rate (OCR) and ATP production in the latter (Fig. 4c,d). This was associated with reduced mitochondrial membrane potential (Fig. 4e), but not mitochondrial load (Fig. 4f). Furthermore, DMM and DES increased ROS (Fig. 4g) and decreased NADPH levels (Fig. 4h), suggesting that in adrenocortical cells, as in macrophages [45], succinate repurposes mitochondrial metabolism from oxidative phosphorylation to ROS production. Such changes in the mitochondrial function were not observed when inhibiting IDH activity with enasidenib (AG221)

[51] (Fig. 4i-k). AG221 increased isocitrate and the isocitrate / α-ketoglutarate ratio (Fig. 4i), but did not affect OCR (Fig. 4j) or the mitochondrial membrane potential (Fig. 4k).

Fig. 4

Increased succinate levels impair mitochondrial function in adrenocortical cells. **a,b** Succinate and fumarate levels were measured by LC–MS/MS in adrenal gland explants (**a**) and NCI-H295R cells (**b**) treated with DMM or DES for 24 h (n = 5 and n = 4 for **a** and **b**, respectively). **c** OCR measurement with Seahorse technology in NCI-H295R cells treated with DMM or DES for 24 h (n = 6). **d** Measurement of ATP/ADP ratio in NCI-H295R cells treated with DMM or DES for 24 h (n = 4-8). **e,f** TMRE and Mitotracker Green FM staining assessed by flow cytometry in NCI-H295R cells treated with DES for 4 h, MFI is shown (n = 7 for **e** and n = 4, one from two experiments for **f**). **g** ROS measurement in NCI-H295R cells treated with DMM or DES for 2 h (n = 10 - 12). **h** Measurement of NADPH/NADP ratio in NCI-H295R cells treated with DMM for 24 h (n = 3 - 4). **i** Isocitrate levels measured by LC–MS/MS in NCI-H295R cells treated for 24 h with AG221 or DMSO (n = 6). **j** OCR measurement in NCI-H295R cells treated

for 24 h with AG221 or DMSO (n = 10). **k** TMRE staining and flow cytometry in NCI-H295R cells treated for 4 h with AG221 or DMSO, MFI is shown (n = 7). Data are presented as mean ± s.d., and analyzed with one-way ANOVA (**a**,**d**,**g**) or Mann-Whitney test (**b**,**e**,**f**,**h**,**i**,**k**). Data in **c** and **j** are shown as mean ± s.e.m. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

Key steps of steroidogenesis take place in the mitochondria [49], thus, we asked whether disruption of SDH activity affects steroidogenic function. We inhibited SDH activity with DMM in human and mouse adrenocortical cells, and induced glucocorticoid production by forskolin or ACTH, respectively. SDH inhibition considerably impaired glucocorticoid and progesterone production in mouse primary adrenocortical cells (Fig. 5a,b), adrenal gland explants (Fig. 5c-e) and human adrenocortical NCI-H295R cells (Suppl. Fig. 4a,b). Similarly, DES diminished glucocorticoid production in mouse (Fig. 5a,b) and human adrenocortical cells (Suppl. Fig. a,b). Confirming these data, *Sdhb* silencing (Suppl. Fig. 5a,b) impaired glucocorticoid synthesis in mouse (Fig. 5f-h) and human adrenocortical cells (Suppl. Fig. 4c), implying that proper adrenocortical steroidogenesis relies on intact SDH activity. Recently it was shown that SDH activity and intracellular succinate are required for CYP11A1-mediated pregnenolone synthesis, the first step of steroidogenesis [52]. Adding to this knowledge, our data demonstrate that increasing succinate concentrations impair steroidogenesis (Suppl. Fig. 4d-f). Moreover, the proton gradient uncoupler FCCP (Fig. 5i) and the ATP synthase inhibitor oligomycin (Fig. 5j-m) both strongly reduced steroidogenesis in mouse (Fig. 5k-m) and human adrenocortical cells (Fig. 5i,j), demonstrating the well-established requirement of intact mitochondrial membrane potential and ATP generation for steroidogenic function [52,53]. In accordance, DMM and DES downregulated the expression of *Cyp11a1* and *Cyp11b1* (Fig. 5n,o), genes encoding for enzymes with critical roles in glucocorticoid synthesis, that catalyze the conversion of cholesterol to pregnenolone and the final step of corticosterone/cortisol production, respectively [2, 49]. Importantly, treatment of adrenal gland explants with LPS reduced corticosterone secretion in response to ACTH, similarly to DMM of DES (Fig. 5p).

In contrast, inhibition of IDH activity with AG221 (Fig. 4i), did not alter glucocorticoid production in mouse adrenocortical cells (Suppl. Fig. 6a,b), adrenal gland explants (Suppl. Fig. 6c,d) or

human adrenocortical cells (Suppl. Fig. 6e,f), nor did *Idh2* silencing in mouse adrenocortical cells (Suppl. Fig. 5c, 6g,h). Taken together, these results imply that SDH but not IDH activity is required for adrenocortical steroidogenesis.

Fig. 5

Disruption of SDH function impairs glucocorticoid production. **a-e** Primary adrenocortical cells (**a**,**b**) and adrenal explants (**c-e**) were treated for 24 h with DMM or DES and for another 45 min with ACTH (10 ng/mL or 100 ng/mL, respectively) (n = 5-6). **f-g** Primary adrenocortical cells were transfected with si*Sdhb* or non-targeting siRNA (siCtrl) and 24 h post-transfection they were treated for 45 min with ACTH (n = 7-8). **i,j** NCI-H295R cells were treated for 24 h with FCCP (**i**) or oligomycin (OM) (**j**) and for another 30 min with Forskolin (Fsk) (n = 6). **k-m** Primary adrenocortical cells were treated for 24 h with oligomycin (OM) and for another 45 min with ACTH (n = 6). **n,o** *Cyp11a1* and *Cyp11b1* expression in primary adrenocortical cells treated for 24 h with DMM or DES and for 45 min with ACTH (n = 5-6). **p** Adrenal gland explants were treated for 24 h with LPS and for 45 min with ACTH (n = 4-5). Measurements of steroid

hormones in **a-m,p** were performed in supernatants of primary adrenocortical cell cultures or adrenal gland explants by LC-MS/MS. Data are shown as mean ± s.d. and analyzed with one-way ANOVA. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. BLD = below level of detection.

5. Itaconate is not responsible for reduced SDH activity and steroidogenesis in adrenocortical cells

In inflammatory macrophages SDH function is inhibited by itaconate [54], a byproduct of the TCA cycle produced from cis-aconitate in a reaction catalyzed by aconitate decarboxylase 1 (ACOD1) [55]. The expression of *Irg1*, the gene encoding for ACOD1, and itaconate levels are strongly upregulated in macrophages upon inflammation [54]. We asked whether itaconate might affect SDH activity in the adrenal cortex. *Irg1* expression was upregulated in the adrenal cortex of LPS-treated mice but this increase derived from CD45$^+$ cells, while *Irg1* was not expressed in CD31$^-$CD45$^-$ adrenocortical cells (Suppl. Fig. 7a). Accordingly, LPS treatment significantly elevated itaconate levels in the CD31$^+$CD45$^+$ fraction, while it did not increase itaconate levels in CD31$^-$CD45$^-$ adrenocortical cells (Suppl. Fig. 7b,c). Itaconate can be secreted from LPS-stimulated macrophages [54], and could thereby affect SDH activity in adrenocortical cells. Therefore, we tested whether exogenously given itaconate may affect steroidogenesis by treating primary adrenocortical cells with the cell permeable itaconate derivative 4-octyl itaconate (4-OI). Adrenocortical cells internalized the added itaconate derivative (Suppl. Fig. 7d), which however did not alter succinate or fumarate levels or the succinate / fumarate ratio (Suppl. Fig. 7e-g), nor did it affect glucocorticoid production (Suppl. Fig. 7h,i). Additionally, SDH activity in the adrenal cortex of *Irg1*-KO mice injected with LPS was not different from that in their wild-type counterparts (Suppl. Fig. 7j). Hence, neither is itaconate produced nor does it affect SDH activity through paracrine routes in adrenocortical cells.

6. IL-1β downregulates SDHB expression and steroidogenesis in a DNMT1-dependent manner.

Systemic inflammation induces substantial leukocyte recruitment in the adrenal gland, accompanied by elevated production of pro-inflammatory cytokines [7, 56]. IL-1β is highly produced by inflammatory monocytes and macrophages [57]. RNA-Seq in the adrenal cortex, including recruited immune cells, showed increased expression of *Il-1β* in LPS- compared to PBS-injected mice (log2fc = 1.46, padj = 0.019). Furthermore, there was significant positive enrichment of genes associated with IL-1β secretion in the adrenal cortex of mice treated with LPS (Fig. 6a). The IL-1β receptor *Il-1r1* is expressed in $CD31^-CD45^-$ adrenocortical cells and its expression was upregulated in adrenocortical cells sorted from LPS treated mice (Fig. 6b). In accordance, proteins related to IL-1β signaling were positively enriched in $CD31^-CD45^-$ adrenocortical cells of LPS mice (Fig. 6c). Essentially, IL-1β, but not IL-6 or TNFα, reduced *SDHB* expression in NCI-H295R cells (Fig. 6d). One way of transcriptional gene repression is covalent attachment of methyl groups on the cytosine 5' position within the gene promoter sequence, a reaction catalyzed by DNA methyltransferases [58]. Proteomics revealed significant upregulation of DNA methyltransferase 1 (DNMT1) in $CD31^-CD45^-$ adrenocortical cells of LPS mice (log2fc = 0.421, padj = 0.015), which we confirmed by western blot analysis (Fig. 6e). In accordance, IL-1β increased DNA methylation of the *SDHB* promoter (Fig. 6f), and *Dnmt1* silencing restored *Sdhb* expression in IL-1β treated adrenocortical cells (Fig. 6g, Suppl. Fig. 5d). Lastly, IL-1β reduced adrenocortical steroidogenesis (Fig. 6h-j), similarly to DMM, DES and LPS (Fig. 5a-e, 5p).

Fig. 6

IL-1β reduces SDHB expression and glucocorticoid synthesis in a DNMT1-dependent manner.

a GSEA for genes related to positive regulation of IL-1β secretion in the adrenal cortex of mice treated for 6 h with PBS or LPS (n = 3 mice per group). **b** *Il-1r1* expression in CD31⁻CD45⁻ adrenocortical cells of mice 6 h post-injection with PBS or LPS (n = 6 mice per group). **c** GSEA for proteins related to IL-1β signaling in CD31⁻CD45⁻ adrenocortical cells of mice treated for 24 h with PBS or LPS (n = 6 mice per group). **d** *SDHB* expression in NCI-H295R cells treated for 2 h with IL-1β, IL-6 or TNFα (n = 6). **e** Western blot analysis for DNMT1 in CD31⁻CD45⁻ adrenocortical cells of mice 24 h after injection of PBS (P) or LPS (L) (n = 4 mice per group), α-TUBULIN was used as loading control. The asterisk (*) depicts an unspecific band.

Quantification of the western blot, shown as relative intensity of DNMT1 to α-TUBULIN. **f** NCI-H295R cells were treated for 2 h with IL-1β (n = 4), and the ratio of methylated to unmethylated DNA in the *SDHB* promoter was assayed after bisulfite conversion. Representative gel electrophoresis images of bisulfite converted and non-treated DNA (M – methylated, U – unmethylated). **g** *Sdhb* expression in primary adrenocortical cells transfected with si*Dnmt1* or siCtrl and 24 h post-transfection treated for 6 h with IL-1β (n = 7-8). **h-j** Primary adrenocortical cells were treated for 6 h with IL-1β and for another 45 min with ACTH (10 ng/mL) (n = 11-12). Steroid hormones were measured in the culture supernatant by LC-MS/MS. Data are shown as mean ± s.d. and analyzed with Mann-Whitney (**b**,**d**,**e**,**f**) or Wilcoxon test (**g-j**). *$p < 0.05$, **$p < 0.01$.

Discussion

Glucocorticoid production in response to inflammation is essential for survival. The adrenal gland shows great resilience to damage induced by inflammation due to its strong regenerative capacity [7, 59,60]. This maintains glucocorticoid release during infection or sterile inflammation, which is vital to restrain and resolve inflammation [1,4]. However, severe sepsis is associated with adrenocortical impairment [11-15,61]. We show here that the inflamed adrenal cortex undergoes cellular metabolic reprograming which involves perturbations in the TCA cycle and oxidative phosphorylation, finally leading to impaired steroidogenesis. Our findings provide a mechanistic explanation of inflammation-related impaired adrenocortical steroidogenesis through cell metabolic reprogramming of steroidogenic adrenocortical cells.

In specific, we demonstrate that IL-1β reduces *SDHB* expression through DNMT1-dependent DNA methylation of the *SDHB* promoter. Several studies have shown that inflammation promotes DNA methylation and thereby regulates gene expression [62-66]. Particularly IL-1β was demonstrated to increase DNA methylation in different genes in a cell type-specific manner [63,67]. In accordance, DNMT1 expression was shown to increase upon acute inflammation in human peripheral blood mononuclear cells or mouse spleens [62,68], as well as in fibroblasts treated with IL-1β [67]. Moreover, reduced *SDH* promoter methylation associates with enhanced SDHB expression and reduced succinate levels in villi from individuals with recurrent spontaneous abortion [69]. These reports stand in accordance with our findings showing regulation of SDHB expression through its promoter methylation by an IL-1β – DNMT axis in steroidogenic adrenocortical cells. In contrast, itaconate, which was shown to reduce SDH activity in macrophages [54], does not regulate SDH in adrenocortical cells.

Consequently, accumulation of succinate leads to impaired oxidative phosphorylation and ATP synthesis, coupled to reduced steroidogenesis. Intact mitochondrial membrane potential and ATP generation are essential requirements for steroidogenic function [52,53]. This we confirmed by treatment of adrenocortical cells with the mitochondrial uncoupler FCCP or the ATP synthase inhibitor oligomycin, which both diminished steroidogenesis. Interestingly, a switch from the canonical towards a non-canonical TCA cycle, involving metabolism of

mitochondrially derived citrate to acetyl-CoA, was recently described and may be activated in inflammation [70,71]. Whether a shift to the non-canonical TCA cycle regulates steroidogenesis remains to be elucidated.

SDH regulates ETC-mediated ROS formation: SDH inhibition or increased succinate levels augment ROS generation in tumors and macrophages [45,72-75]. Similarly, we show in adrenocortical cells that SDH inhibition or high succinate levels lead to increased ROS levels at the expense of mitochondrial oxidative function and ATP production. Adrenocortical disorders such as triple A syndrome and familial glucocorticoid deficiency can be driven by increased oxidative stress in the adrenal cortex [46]. In fact, mutations in genes encoding for proteins conferring antioxidant protection were implicated in the development of adrenocortical deficiencies [46]. Hence, SDH dysfunction leading to oxidative stress may be an important component of the pathophysiology of adrenocortical insufficiency, a notion which merits further investigation.

In conclusion, we demonstrate that tight regulation of succinate levels is essential for normal steroidogenesis, while disruption of SDH expression in inflammation through the IL-1β-DNMT1 axis contributes to adrenocortical dysfunction (Fig. 7). This study expands the current knowledge on regulation of glucocorticoid production and identifies potential targets for therapeutic interventions.

Fig. 7

Illustration of the regulation of adrenocortical steroidogenesis by inflammation. IL-1β reduces SDHB expression through upregulation of DNA methyltransferase 1 (DNMT1) and methylation of the SDHB promoter. Consequently, increased succinate levels impair oxidative phosphorylation, leading to reduced steroidogenesis.

References

1. Chrousos GP (1995) The hipothalamic-pituitary-adrenal axis and immune-mediated inflammation. N Engl J Med 332(20):1351–1362

2. Payne AH, Hales DB (2004) Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. Endocr Rev 25(6):947–70

3. Lightman SL, Birnie MT, Conway-Campbell BL (2021) Dynamics of ACTH and cortisol secretion and implications for disease. Endocr Rev 41(3):470–490

4. Alexaki VI, Henneicke H. The Role of Glucocorticoids in the Management of COVID-19 (2021) Horm Metab Res 53(1):9–15

5. Kanczkowski W, Chatzigeorgiou A, Samus M, Nguyen T, Zacharowski K, Chavakis T, Bornstein SR (2013) Characterization of the LPS-induced inflammation of the adrenal gland in mice. Mol Cell Endocrinol 371(1–2):228–235

6. Kanczkowski W, Alexaki VI, Nguyen T et al (2013) Hypothalamo-pituitary and immune-dependent adrenal regulation during systemic inflammation. Proc Natl Acad Sci USA 110(36):14801–14806

7. Kanczkowski W, Chatzigeorgiou A, Grossklaus S, Sprott D, Bornstein SR, Chavakis T (2013) Role of the endothelial-derived endogenous anti-inflammatory factor del-1 in inflammation-mediated adrenal gland dysfunction. Endocrinol 154(3):1181–1189

8. Alexaki VI (2021) The Impact of Obesity on Microglial Function: Immune, Metabolic and Endocrine Perspectives. Cells 10(7):1584

9. Bertini R, Bianchi M, Ghezzi P. Adrenalectomy sensitizes mice to the lethal effects of interleukin 1 and tumor necrosis factor (1988) J Exp Med 167(5):1708–1712

10. Butler LD, Layman NK, Riedl PE, Cain RL, Shellhaas J, Evans GF, Zuckerman SH (1989) Neuroendocrine regulation of in vivo cytokine production and effects: I. In vivo regulatory networks involving the neuroendocrine system, interleukin-1 and tumor necrosis factor-α. J Neuroimmunol 24(1–2):143–153

11. Annane D, Sébille V, Troché G, Raphaël JC, Gajdos P, Bellissant E (2000) A 3-level prognostic classification in septic shock based on cortisol levels and cortisol response to

corticotropin. JAMA 283(8):1038–1045

12. Boonen E, Bornstein SR, Van den Berghe G (2015) New insights into the controversy of adrenal function during critical illness. Lancet Diabetes Endocrinol 3(10):805–815

13. Boonen E, Meersseman P, Vervenne H et al (2014) Reduced nocturnal ACTH-driven cortisol secretion during critical illness. Am J Physiol - Endocrinol Metab 306(8):2–11

14. Den Brinker M, Joosten KFM, Liem O et al (2005) Adrenal insufficiency in meningococcal sepsis: Bioavailable cortisol levels and impact of interleukin-6 levels and intubation with etomidate on adrenal function and mortality. J Clin Endocrinol Metab 90(9):5110–5117

15. Jennewein C, Nguyen T, Kanczkowski W et al (2016) Mortality of Septic Mice Strongly Correlates with Adrenal Gland Inflammation. Crit Care Med 44(4):e190–e199

16. Ryan DG, O'Neill LAJ (2020) Krebs Cycle Reborn in Macrophage Immunometabolism. Annu Rev Immunol 38: 289–313

17. O'Neill LAJ, Pearce EJ (2016) Immunometabolism governs dendritic cell and macrophage function. J Exp Med 213: 15–23

18. Geltink RIK, Kyle RL Pearce EL (2018) Unraveling the Complex Interplay between T Cell Metabolism and Function. Annu Rev Immunol 36: 461–488

19. Martin M (2011) Cutadept removes adapter sequences from high-throughput sequncing reads. EMBnet.journal 2011;7(10):2803–2809

20. Dobin A, Davis CA, Schlesinger F et al (2013) STAR: Ultrafast universal RNA-seq aligner. Bioinformatics 29(1):15–21

21. Anders S, Pyl PT, Huber W (2015) HTSeq-A Python framework to work with high-throughput sequencing data. Bioinformatics 31(2):166–169

22. Anders S, Huber W (2010) Differential expression analysis for sequence count data. Genome Biol 11(R106):4310–4315

23. Wickham H (2009) ggplot2: elegant graphics for data analysis. Springer New York

24. Subramanian A, Tamayo P, Mootha VK et al (2005) Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci USA 102(43):15545–15550

25. Alhamdoosh M, Ng M, Wilson NJ, Sheridan JM, Huynh H, Wilson MJ, Ritchie ME (2017)

Combining multiple tools outperforms individual methods in gene set enrichment analyses. Bioinformatics 33(3):414–424

26. Chen N, Van Der Schors RC, Smit AB. A 1D-PAGE/LC-ESI Linear Ion Trap Orbitrap MS Approach for the Analysis of Synapse Proteomes and Synaptic Protein Complexes. Humana Press; 2011

27. Meier F, Brunner A-D, Frank M et al (2020) diaPASEF: parallel accumulation–serial fragmentation combined with data-independent acquisition. Nat Methods 17(12):1229–1236

28. Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M (2020) DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. Nat Methods 17(1):41–44

29. Hondius DC, Koopmans F, Leistner C et al (2021) The proteome of granulovacuolar degeneration and neurofibrillary tangles in Alzheimer's disease. Acta Neuropathol141(3):341–358

30. Koopmans F, Pandya NJ, Franke SK, Phillippens IHCMH, Paliukhovich I, Li KW, Smit AB (2018) Comparative Hippocampal Synaptic Proteomes of Rodents and Primates: Differences in Neuroplasticity-Related Proteins. Front Mol Neurosci 11:364

31. Zhang X, Smits AH, van Tilburg GB, Ovaa H, Huber W, Vermeulen M (2018) Proteome-wide identification of ubiquitin interactions using UbIA-MS. Nat Protoc 13(3):530–550

32. Team RC (2021) R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria

33. Peitzsch M, Dekkers T, Haase M et al (2015) An LC-MS/MS method for steroid profiling during adrenal venous sampling for investigation of primary aldosteronism. J Steroid Chem Mol Biol 145:75–84

34. Richter S, Gieldon L, Pang Y et al (2019) Metabolome-guided genomics to identify pathogenic variants in isocitrate dehydrogenase, fumarate hydratase, and succinate dehydrogenase genes in pheochromocytoma and paraganglioma. Genet Med 21(3):705–717

35. Aichler M, Borgmann D, Krumsiek J et al (2017) N-acyl Taurines and Acylcarnitines Cause an Imbalance in Insulin Synthesis and Secretion Provoking β Cell Dysfunction in Type 2 Diabetes. Cell Metab 25(6):1334-1347

36. Sun N, Fernandez IE, Wei M et al (2018) Pharmacometabolic response to pirfenidone in pulmonary fibrosis detected by MALDI-FTICR-MSI. Eur Respir J 52(3) :1702314

37. Yuan M, Breitkopf SB, Yang X, Asara JM (2012) A positive/negative ion-switching, targeted mass spectrometry-based metabolomics platform for bodily fluids, cells, and fresh and fixed tissue. Nat Protoc 7(5):872–881

38. Laemelli UK (1970) Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. Nature 227(5259):680–685

39. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P (2015) The Molecular Signatures Database Hallmark Gene Set Collection. Cell Syst 1(6):417–425

40. González-Hernández JA, Bornstein SR, Ehrhart-Bornstein M, Geschwend JE, Adler G, Scherbaum WA (1994) Macrophages within the human adrenal gland. Cell Tissue Res 278(2):201–205

41. Schober A, Huber K, Fey J, Unsicker K (1998) Distinct populations of macrophages in the adult rat adrenal gland: A subpopulation with neurotrophin-4-like immunoreactivity. Cell Tissue Res 291(3):365–373

42. Chen LS, Singh SP, Schuster M, Grinenko T, Bornstein SR, Kanczkowski W (2019) RNA-seq analysis of LPS-induced transcriptional changes and its possible implications for the adrenal gland dysregulation during sepsis J Steroid Biochem Mol Biol 191:105360

43. Miller WL (2007) Steroidogenic acute regulatory protein (StAR), a novel mitochondrial cholesterol transporter. Biochim Biophys Acta - Mol Cell Biol Lipids 1771(6):663–676

44. Parker KL, Schimmer BP (1997) Steroidogenic factor 1: a key mediator of endocrine development and function. Endocr Rev 18(3):361–377

45. Mills EL, Kelly B, Logan A et al (2016) Succinate Dehydrogenase Supports Metabolic Repurposing of Mitochondria to Drive Inflammatory Macrophages. Cell 167(2):457-470

46. Prasad R, Kowalczyk JC, Meimaridou E, Storr HL, Metherell LA (2014) Oxidative stress and adrenocortical insufficiency. J Endocrinol 221(3):R63-73

47. Xiao W, Loscalzo J (2020) Metabolic Responses to Reductive Stress. Antioxid Redox Sign 32(18):1330–1347

48. Frederiks WM, Kümmerlin IPED, Bosch KS, Vreeling-Sindelárová H, Jonker A, Van

Noorden CJF (2007) NADPH production by the pentose phosphate pathway in the zona fasciculata of rat adrenal gland. J Histochem Cytochem 55(9):975–980

49. Midzak A, Papadopoulos V (2016) Adrenal mitochondria and steroidogenesis: From individual proteins to functional protein assemblies. Front Endocrinol (Lausanne) 7:106

50. Moosavi B, Zhu XL, Yang WC, Yang GF (2020) Genetic, epigenetic and biochemical regulation of succinate dehydrogenase function. Biol Chem 401(3):319–330

51. Yen K, Travins J, Wang F et al (2017) AG-221, a first-in-class therapy targeting acute myeloid leukemia harboring oncogenic IDH2 mutations. Cancer Discov 7(5):478–493

52. Bose HS, Marshall B, Debnath DK, Perry EW, Whittal RM (2020) Electron Transport Chain Complex II Regulates Steroid Metabolism iScience 23(7):101295

53. King SR, Liu Z, Soh J, Eimerl S, Stocco DM (1999) Effects of disruption of the mitochondrial electrochemical gradient on steroidogenesis and the steroidogenic acute regulatory (StAR) protein. J Steroid Biochem Mol Biol 69(1–6):143–154

54. Lampropoulou V, Sergushichev A, Bambouskova M et al (2016) Itaconate Links Inhibition of Succinate Dehydrogenase with Macrophage Metabolic Remodeling and Regulation of Inflammation Cell Metab 24(1):158–166

55. Michelucci A, Cordes T, Ghelfi J et al (2013) Immune-responsive gene 1 protein links metabolism to immunity by catalyzing itaconic acid production. Proc Natl Acad Sci USA 110(19):7820–7825

56. Chen L-S, Singh SP, Müller G, Bornstein SR, Kanczkowski W (2020) Transcriptional Analysis of Sepsis-Induced Activation and Damage of the Adrenal Endothelial Microvascular Cells. Front Endocrinol (Lausanne) 10:944

57. Netea MG, Simon A, van de Veerdonk F, Kullberg B-J, van der Meer JWM, Boosten LAB (2010) IL-1β processing in host defense: Beyond the inflammasomes. PLoS Pathog 6(2):e1000661

58. Suzuki MM, Bird A (2008) DNA methylation landscapes: Provocative insights from epigenomics. Nat Rev Genet 9(6):465–476

59. Mateska I, Nanda K, Dye NA, Alexaki VI, Eaton S (2020) Range of SHH signaling in adrenal gland is limited by membrane contact to cells with primary cilia. J Cell Biol 219(12) :e201910087

60. Lyraki R, Schedl A (2021) Adrenal cortex renewal in health and disease. Nat Rev Endocrinol 17(7):421–434

61. Annane D, Maxime V, Ibrahim F, Alvarez JC, Abe E, Boudou P (2006) Diagnosis of adrenal insufficiency in severe sepsis and septic shock. Am J Respir Crit Care Med 174(12):1319–1326

62. Koos B, Moderegger EL, Rump K et al (2020) LPS-Induced Endotoxemia Evokes Epigenetic Alterations in Mitochondrial DNA That Impacts Inflammatory Response. Cells 9(10):2282

63. Li R, Ong SL, Tran LM et al (2020) Chronic IL-1β-induced inflammation regulates epithelial-to-mesenchymal transition memory phenotypes via epigenetic modifications in non-small cell lung cancer. Sci Rep 10(1):377

64. Rodriguez RM, Suarez-Alvarez B, Lavín JL et al (2019) Signal Integration and Transcriptional Regulation of the Inflammatory Response Mediated by the GM-/M-CSF Signaling Axis in Human Monocytes. Cell Rep 29(4):860-872

65. Weiss E, Vlahos A, Kim B et al (2021) Transcriptomic Remodelling of Fetal Endothelial Cells During Establishment of Inflammatory Memory. Front Immunol 12:757393

66. Morante-Palacios O, Lorente-Sorolla C, Ciudad L et al (2021) JAK2-STAT Epigenetically Regulates Tolerized Genes in Monocytes in the First Encounter With Gram-Negative Bacterial Endotoxins in Sepsis. Front Immunol 12:734652

67. Seutter S, Winfield J, Esbitt A et al (2020) Interleukin 1β and Prostaglandin E2 affect expression of DNA methylating and demethylating enzymes in human gingival fibroblasts. Int Immunopharmacol 78:105920

68. Cao L, Zhu T, Lang X et al (2020) Inhibiting DNA Methylation Improves Survival in Severe Sepsis by Regulating NF-κB Pathway. Front Immunol 11:1360

69. Wang XH, Xu S, Zhou X-Y et al (2021) Low chorionic villous succinate accumulation associates with recurrent spontaneous abortion risk. Nat Commun 12(1):3428

70. Arnold PK, Jackson BT, Paras KI et al (2022) A non-canonical tricarboxylic acid cycle underlies cellular identity. Nature 603(7901):477–481

71. Mateska I, Alexaki VI (2022) Light shed on a non-canonical TCA cycle : cell state regulation beyond mitochondrial energy production. Signal Transduct Target Ther 7(1):201

72. Hadrava Vanova K, Kraus M, Neuzil J, Rohlena J (2020) Mitochondrial complex II and reactive oxygen species in disease and therapy. Redox Rep 25(1):26–32

73. Ralph SJ, Moreno-Sánchez R, Neuzil J, Rodríguez-Enríquez S (2011) Inhibitors of succinate: Quinone reductase/Complex II regulate production of mitochondrial reactive oxygen species and protect normal cells from ischemic damage but induce specific cancer cell death. Pharm Res 28(11):2695–2730

74. Guzy RD, Sharma B, Bell E, Chandel NS, Schumacker PT (2008) Loss of the SdhB, but Not the SdhA, Subunit of Complex II Triggers Reactive Oxygen Species-Dependent Hypoxia-Inducible Factor Activation and Tumorigenesis. Mol Cell Biol 28(2):718–731

75. Selak MA, Armour SM, MacKenzie ED et al (2005) Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF-α prolyl hydroxylase. Cancer Cell 7(1):77–85

Statements & Declarations

Funding

This work was supported by grants from the Deutsche Forschungsgemeinschaft (SFB/TRR 205 to V.I. Alexaki, B. Wielockx and M. Peitzsch) and the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement (No 765704 to V.I. Alexaki).

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author contributions

I.M.: planned and performed experiments, analyzed data, prepared the figures, wrote the manuscript; A.W: planned and performed experiments, analyzed data, edited the manuscript; E.H.: performed experiments, analyzed data; C.Y.: performed experiments; A.S.: analyzed data; E.T.: performed experiments; N.S.: performed experiments, analyzed data; O.K.: performed experiments; M.P.: performed experiments; H.A.: performed experiments; H.H.: edited the manuscript; W.K.: performed experiments; B.W.: edited and discussed the manuscript; C.T.: planned experiments; A.D.: planned experiments; A.W.: planned experiments; K.W.L: planned experiments; M.P.: analyzed data; T.C.: edited and discussed the manuscript; V.I.A.: conceived the study, supervised the project, wrote and edited the manuscript.

Data Availability

The datasets generated during the current study are available from the corresponding author.

RNA-Seq data are available in https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE200220

Ethics approval

The animal experiments were approved by the Landesdirektion Sachsen Germany.

Tables

Table 1. Cellular metabolic pathways transcriptionally regulated by inflammation in the adrenal cortex. The pathway analysis of differentially expressed genes was done with the software package EGSEA and queried against the KEGG Pathways repository. Pathways with p < 0.05 are shown.

ID	Metabolic pathway	Number of expressed genes	p value	p adj	avg.logfc	Direc-tion
mmu00190	Oxidative phosphorylation	132/134	8.32E-15	7.32E-13	0.613034377	Down
mmu00280	Valine, leucine and isoleucine degradation	55/56	1.16E-05	0.001119154	0.725758563	Down
mmu00511	Other glycan degradation	18/18	3.82E-05	0.001119154	0.29921956	Down
mmu00980	Metabolism of xenobiotics by cytochrome P450	65/65	0.000282456	0.006214029	0.650304677	Down
mmu00350	Tyrosine metabolism	38/39	0.001754788	0.030884268	0.305935415	Down
mmu00640	Propanoate metabolism	31/31	0.002151447	0.031554563	0.886410728	Down
mmu00020	Citrate cycle (TCA cycle)	32/32	0.003091828	0.034636464	0.21912487	Down
mmu01200	Carbon metabolism	118/118	0.003685749	0.034636464	0.715139933	Down
mmu00471	D-Glutamine and D-glutamate metabolism	3/3	0.003980826	0.034636464	0.338725016	Up
mmu00300	Lysine biosynthesis	2/2	0.004518807	0.034636464	0.156233777	Down
mmu00630	Glyoxylate and dicarboxylate metabolism	29/29	0.006079126	0.034636464	0.886410728	Down

mmu0 0071	Fatty acid degradation	49/49	0.0061 24673	0.0346 36464	0.4014 41917	Down
mmu0 1210	2-Oxocarboxylic acid metabolism	19/19	0.0062 97539	0.0346 36464	0.9426 61129	Down
mmu0 0920	Sulfur metabolism	11/11	0.0062 97539	0.0346 36464	0.6018 64913	Down
mmu0 0480	Glutathione metabolism	58/59	0.0062 97539	0.0346 36464	0.5502 57753	Down
mmu0 0510	N-Glycan biosynthesis	49/49	0.0062 97539	0.0346 36464	0.2544 26968	Down
mmu0 0450	Selenocompound metabolism	17/17	0.0091 15527	0.0471 86259	0.3291 18527	Up
mmu0 0514	Other types of O-glycan biosynthesis	22/22	0.0106 14958	0.0506 93394	0.2049 24928	Up
mmu0 0440	Phosphonate and phosphinate metabolism	6/6	0.0109 45165	0.0506 93394	0.2915 10253	Up
mmu0 0120	Primary bile acid biosynthesis	16/16	0.0119 3422	0.0525 10566	2.7135 91551	Down
mmu0 0565	Ether lipid metabolism	44/44	0.0167 11812	0.0617 24814	0.4338 37026	Down
mmu0 0520	Amino sugar and nucleotide sugar metabolism	49/49	0.0182 12661	0.0617 24814	0.6526 46241	Down
mmu0 0790	Folate biosynthesis	14/14	0.0182 1569	0.0617 24814	0.4200 00237	Down
mmu0 0230	Purine metabolism	174/178	0.0187 93387	0.0617 24814	0.8301 15541	Down
mmu0 0603	Glycosphingolipid biosynthesis – globo series	16/16	0.0198 32034	0.0617 24814	0.5342 87429	Up
mmu0 0534	Glycosaminoglycan biosynthesis – 42eparan sulfate/heparin	24/24	0.0204 91524	0.0617 24814	0.4713 78176	Up

mmu0 0270	Cysteine and methionine metabolism	46/48	0.0209 59507	0.0617 24814	0.6018 64913	Down
mmu0 1100	Metabolic pathways	1303/1315	0.0224 19002	0.0617 24814	0.6839 47211	Down
mmu0 0531	Glycosaminoglycan degradation	21/21	0.0231 39686	0.0617 24814	0.9117 53864	Down
mmu0 0604	Glycosphingolipid biosynthesis – ganglio series	15/15	0.0232 68716	0.0617 24814	0.4572 85673	Down
mmu0 0250	Alanine, aspartate and glutamate metabolism	36/37	0.0236 85519	0.0617 24814	0.9426 61129	Down
mmu0 0240	Pyrimidine metabolism	101/104	0.0242 35887	0.0617 24814	0.3081 35641	Up
mmu0 0130	Ubiquinone and other terpenoid-quinone biosynthesis	11/11	0.0246 30584	0.0617 24814	0.4099 1964	Down
mmu0 0330	Arginine and proline metabolism	49/50	0.0250 88792	0.0617 24814	0.4795 54867	Down
mmu0 0564	Glycerophospholipid metabolism	94/94	0.0254 48859	0.0617 24814	0.4981 46256	Down
mmu0 0910	Nitrogen metabolism	17/17	0.0257 82601	0.0617 24814	1.7655 82115	Up
mmu0 0982	Drug metabolism – cytochrome P450	67/67	0.0272 01304	0.0617 24814	0.6205 62175	Down
mmu0 0785	Lipoic acid metabolism	3/3	0.0275 07553	0.0617 24814	0.1334 03173	Down
mmu0 0051	Fructose and mannose metabolism	35/35	0.0296 79889	0.0617 24814	0.6526 46241	Down
mmu0 0561	Glycerolipid metabolism	59/59	0.0304 68773	0.0617 24814	0.4155 946	Down
mmu0 0512	Mucin type O-glycan biosynthesis	28/28	0.0305 40327	0.0617 24814	0.4724 58211	Up

mmu00052	Galactose metabolism	32/32	0.031376559	0.061724814	0.42838689	Down
mmu01230	Biosynthesis of amino acids	78/78	0.031800904	0.061724814	0.942661129	Down
mmu00533	Glycosaminoglycan biosynthesis – keratan sulfate	14/14	0.031872109	0.061724814	0.27544313	Down
mmu00900	Terpenoid backbone biosynthesis	22/23	0.032232321	0.061724814	0.320850492	Up
mmu00730	Thiamine metabolism	15/15	0.032265244	0.061724814	0.323024942	Down
mmu00500	Starch and sucrose metabolism	33/33	0.033413545	0.062178073	0.452533726	Up
mmu00770	Pantothenate and CoA biosynthesis	18/18	0.033915312	0.062178073	0.307590054	Down
mmu00062	Fatty acid elongation	27/27	0.035703854	0.062200773	0.536297353	Down
mmu00592	alpha-Linolenic acid metabolism	25/25	0.035861114	0.062200773	0.568488547	Down
mmu00562	Inositol phosphate metabolism	70/70	0.037060597	0.062200773	0.197317479	Up
mmu00760	Nicotinate and nicotinamide metabolism	34/35	0.037777703	0.062200773	0.44361797	Down
mmu00740	Riboflavin metabolism	8/8	0.038531607	0.062200773	0.336638606	Down
mmu00670	One carbon pool by folate	19/19	0.039807001	0.062200773	0.293141874	Up
mmu00310	Lysine degradation	57/59	0.040221284	0.062200773	0.409052131	Up
mmu00010	Glycolysis / Gluconeogenesis	66/66	0.040372513	0.062200773	0.4474754	Down

ID	Metabolic pathway		p.value	p.adj		Down
mmu0 0053	Ascorbate and aldarate metabolism	27/27	0.0405 35303	0.0622 00773	0.3190 47565	Down
mmu0 0030	Pentose phosphate pathway	32/32	0.0409 95964	0.0622 00773	0.3163 47941	Down

Table 2. Cellular metabolic pathways regulated on protein level by inflammation in adrenocortical cells. The pathway analysis of differentially expressed proteins was done with the software package EGSEA and queried against the KEGG Pathways repository. Pathways with $p < 0.05$ are shown.

ID	Metabolic pathway	Number of detected proteins	p.value	p.adj	avg.l ogfc	Direc- tion
mmu0 0534	Glycosaminoglycan biosynthesis – heparan sulfate / heparin	4/24	1.50E-07	1.29E-05	0.13	Up
mmu0 0280	Valine, leucine and isoleucine degradation	37/57	5.20E-07	2.23E-05	0.01	Down
mmu0 0983	Drug metabolism – other enzymes	25/92	3.80E-05	0.0008 56142	0.02	Down
mmu0 0562	Inositol phosphate metabolism	19/72	5.89E-05	0.0008 56142	0.03	Up
mmu0 0260	Glycine, serine and threonine metabolism	19/40	6.89E-05	0.0008 56142	0.03	Down
mmu0 0240	Pyrimidine metabolism	25/58	6.98E-05	0.0008 56142	0.04	Up
mmu0 0982	Drug metabolism – cytochrome P450	16/71	7.76E-05	0.0008 56142	0.02	Down
mmu0 0230	Purine metabolism	44/133	7.96E-05	0.0008 56142	0.04	Up

mmu0 0790	Folate biosynthesis	9/26	9.75E-05	0.0009 31452	0.09	Up
mmu0 1100	Metabolic pathways	593/1608	0.0001 34694	0.0011 51176	0.03	Down
mmu0 0190	Oxidative phosphorylation	70/135	0.0001 60368	0.0011 51176	0.02	Down
mmu0 0980	Metabolism of xenobiotics by cytochrome P450	19/73	0.0001 75997	0.0011 51176	0.02	Down
mmu0 1240	Biosynthesis of cofactors	74/154	0.0001 78113	0.0011 51176	0.04	Down
mmu0 0730	Thiamine metabolism	5/15	0.0001 87401	0.0011 51176	0.02	Down
mmu0 0020	Citrate cycle (TCA cycle)	30/32	0.0002 0177	0.0011 56812	0.01	Down
mmu0 1200	Carbon metabolism	81/121	0.0004 04805	0.0021 75826	0.02	Down
mmu0 0760	Nicotinate and nicotinamide metabolism	11/41	0.0004 59649	0.0023 25283	0.04	Up
mmu0 0860	Porphyrin and chlorophyll metabolism	19/43	0.0005 49279	0.0026 24333	0.04	Up
mmu0 0360	Phenylalanine metabolism	7/23	0.0007 44591	0.0033 70253	0.01	Up
mmu0 0630	Glyoxylate and dicarboxylate metabolism	20/32	0.0007 94408	0.0034 15956	0.02	Down
mmu0 0480	Glutathione metabolism	28/72	0.0010 37994	0.0042 50832	0.02	Down
mmu0 0061	Fatty acid biosynthesis	12/19	0.0011 9934	0.0043 23466	0.04	Up
mmu0 0052	Galactose metabolism	17/32	0.0012 12032	0.0043 23466	0.02	Down

mmu00350	Tyrosine metabolism	12/40	0.001275764	0.004323466	0.02	Down
mmu00900	Terpenoid backbone biosynthesis	8/23	0.001283046	0.004323466	0.01	Down
mmu00140	Steroid hormone biosynthesis	12/92	0.001307094	0.004323466	0.02	Down
mmu00511	Other glycan degradation	11/18	0.001620057	0.005160182	0.03	Down
mmu00520	Amino sugar and nucleotide sugar metabolism	29/51	0.001922509	0.005904848	0.02	Down
mmu00040	Pentose and glucuronate interconversions	9/35	0.002291756	0.006796243	0.02	Down
mmu00620	Pyruvate metabolism	34/44	0.004741112	0.013428366	0.02	Down
mmu00524	Neomycin, kanamycin and gentamicin biosynthesis	3/5	0.004840458	0.013428366	0.02	Down
mmu00053	Ascorbate and aldarate metabolism	9/31	0.005220665	0.013429037	0.01	Down
mmu00830	Retinol metabolism	8/97	0.005279453	0.013429037	0.01	Down
mmu00531	Glycosaminoglycan degradation	10/21	0.005309154	0.013429037	0.02	Down
mmu00450	Selenocompound metabolism	9/17	0.006951355	0.017080471	0.02	Down
mmu00250	Alanine, aspartate and glutamate metabolism	17/39	0.007955263	0.019004239	0.02	Down
mmu01230	Biosynthesis of amino acids	45/79	0.008873164	0.020624111	0.02	Down
mmu01212	Fatty acid metabolism	40/62	0.009336325	0.021129578	0.03	Down

mmu00500	Starch and sucrose metabolism	14/34	0.011170016	0.024631316	0.02	Down
mmu00514	Other types of O-glycan biosynthesis	15/43	0.011876496	0.025349557	0.04	Up
mmu01210	2-Oxocarboxylic acid metabolism	11/20	0.01227639	0.025349557	0.01	Down
mmu00650	Butanoate metabolism	13/28	0.012404185	0.025349557	0.01	Down
mmu00670	One carbon pool by folate	9/19	0.012674778	0.025349557	0.05	Down
mmu00310	Lysine degradation	20/64	0.01401052	0.027384197	0.02	Down
mmu00590	Arachidonic acid metabolism	9/86	0.015253177	0.029150517	0.01	Down
mmu00770	Pantothenate and CoA biosynthesis	8/21	0.016632522	0.031095584	0.02	Down
mmu00592	alpha-Linolenic acid metabolism	3/25	0.017819371	0.032605657	0.03	Up
mmu00780	Biotin metabolism	3/3	0.018650597	0.033415653	0.01	Down
mmu00640	Propanoate metabolism	25/31	0.020182799	0.035173564	0.02	Down
mmu00290	Valine, leucine and isoleucine biosynthesis	2/4	0.02053428	0.035173564	0.02	Down
mmu00920	Sulfur metabolism	7/11	0.021206387	0.035173564	0.02	Down
mmu00062	Fatty acid elongation	15/19	0.021267736	0.035173564	0.02	Down
mmu00604	Glycosphingolipid biosynthesis – ganglio series	5/15	0.022065753	0.035804807	0.02	Down

mmu0 0563	Glycosylphosphatidylinositol (GPI)-anchor biosynthesis	8/26	0.0235 53511	0.0361 86307	0.04	Down
mmu0 0750	Vitamin B6 metabolism	3/9	0.0235 53511	0.0361 86307	0.03	Up
mmu0 0220	Arginine biosynthesis	7/20	0.0242 45413	0.0361 86307	0.02	Down
mmu0 0270	Cysteine and methionine metabolism	29/53	0.0254 10276	0.0361 86307	0.03	Up
mmu0 0051	Fructose and mannose metabolism	17/36	0.0255 1991	0.0361 86307	0.04	Down
mmu0 0071	Fatty acid degradation	30/52	0.0256 67032	0.0361 86307	0.02	Down
mmu0 0330	Arginine and proline metabolism	23/54	0.0256 67032	0.0361 86307	0.02	Down
mmu0 0561	Glycerolipid metabolism	23/62	0.0256 67032	0.0361 86307	0.03	Down
mmu0 0010	Glycolysis / Gluconeogenesis	40/67	0.0507 28869	0.0681 66917	0.02	Down
mmu0 0030	Pentose phosphate pathway	17/33	0.0507 28869	0.0681 66917	0.03	Down
mmu0 0410	beta-Alanine metabolism	19/32	0.0507 28869	0.0681 66917	0.01	Down

Table 3. ROS pathways are transcriptionally upregulated in the adrenal cortex of LPS-treated mice. The pathway analysis of differentially expressed genes was done with the software package EGSEA and queried against the GO Gene Sets repository. Pathways with padj. < 0.05 are shown.

ID	Gene Set	Number of expressed genes	p value	p adj	avg.l ogfc	Direc -tion
M13 446	GO Regulation of reactive oxygen species metabolic process	271/275	3.75E-08	1.26E-06	1.01 00	Up
M13 580	GO Positive regulation of reactive oxygen species metabolic process	182/186	2.33E-07	6.26E-06	1.01 00	Up
M16 953	GO Response to reactive oxygen species	300/317	0.0034 22132	0.0090 35375	0.86 00	Up
M16 581	GO Cellular response to reactive oxygen species	173/177	0.0025 37297	0.0090 35375	0.76 00	Up
M10 618	GO Negative regulation of response to reactive oxygen species	24/24	0.0072 384	0.0109 42115	0.71 00	Up
M15 379	GO Regulation of reactive oxygen species biosynthetic process	145/148	5.90E-05	0.0007 70609	0.70 00	Up
M10 827	GO Positive regulation of reactive oxygen species biosynthetic process	120/123	0.0004 65606	0.0045 4261	0.70 00	Up
M15 990	GO Reactive oxygen species metabolic process	163/167	0.0089 36465	0.0125 68942	0.67 00	Up
M16 764	GO Regulation of response to reactive oxygen species	43/43	0.0067 53207	0.0104 98628	0.62 00	Down
M16 007	GO Negative regulation of reactive oxygen species biosynthetic process	23/23	0.0164 34387	0.0202 74996	0.60 00	Up
M12 185	GO Reactive oxygen species biosynthetic process	32/33	0.0064 83259	0.0102 4232	0.57 00	Down
M10 894	GO Negative regulation of reactive oxygen species metabolic process	59/59	0.0065 38654	0.0102 87661	0.55 00	Up

Table 4. ROS – related protein expression is upregulated in the adrenal cortex of LPS-treated mice. The pathway analysis of differentially expressed proteins was done with the software package EGSEA and queried against the GO Gene Sets repository. Pathways with padj. < 0.05 are shown.

ID	Protein Set	Number of detected proteins	p.value	p.adj	avg.logfc	Direction
M13 446	GO REGULATION OF REACTIVE OXYGEN SPECIES METABOLIC PROCESS	62/275	6.30E-06	0.000 12401	0.03	Up
M15 379	GO REGULATION OF REACTIVE OXYGEN SPECIES BIOSYNTHETIC PROCESS	34/148	9.43E-06	0.000 14372	0.03	Up
M13 580	GO POSITIVE REGULATION OF REACTIVE OXYGEN SPECIES METABOLIC PROCESS	29/186	1.11E-05	0.000 15778	0.03	Up
M10 827	GO POSITIVE REGULATION OF REACTIVE OXYGEN SPECIES BIOSYNTHETIC PROCESS	20/123	1.17E-05	0.000 16203	0.03	Up
M10 894	GO NEGATIVE REGULATION OF REACTIVE OXYGEN SPECIES METABOLIC PROCESS	23/59	5.29E-05	0.000 37856	0.05	Down
M16 007	GO NEGATIVE REGULATION OF REACTIVE OXYGEN SPECIES BIOSYNTHETIC PROCESS	12/23	0.00014 126	0.000 73167	0.07	Up
M16 581	GO CELLULAR RESPONSE TO REACTIVE OXYGEN SPECIES	50/177	0.00015 378	0.000 77551	0.03	Down
M15 990	GO REACTIVE OXYGEN SPECIES METABOLIC PROCESS	35/167	0.00037 436	0.001 50479	0.02	Down

| M16 953 | GO RESPONSE TO REACTIVE OXYGEN SPECIES | 83/317 | 0.00076 076 | 0.002 62274 | 0.03 | Up |
| M12 185 | GO REACTIVE OXYGEN SPECIES BIOSYNTHETIC PROCESS | 9/33 | 0.00192 844 | 0.005 45214 | 0.02 | Down |

Table 5. Primer sequences.

Gene name	Forward Sequence (5' → 3')	Reverse Sequence (5' → 3')
mouse *18S* rRNA	GTTCCGACCATAAACGATGCC	TGGTGGTGCCCTTCCGTCAAT
mouse *Idh1*	GTGGTGGAGATGCAAGGAGAT	TGGTCATTGGTGGCATCACG
mouse *Idh2*	GATGGACGGTGACGAGATGAC	GGTCTGGTCACGGTTTGGA
mouse *Sdhb*	GGACCTCAGCAAAGTCTCCAA	TGCAGATACTGTTGCTTGCC
mouse *Sdhc*	GCTAAGGAGGAGATGGAGCG	AGAGACCCCTCCACTCAAGG
mouse *Star*	CTGTCCACCACATTGACCTG	CAGCTATGCAGTGGGAGACA
mouse *Cyp11b1*	TCACCATGTGCTGAAATCCTTCCA	GGAAGAGAAGAGAGGGCAATGTGT
mouse *3bHsd2*	GCGGCTGCTGCACAGGAATAAAG	TCACCAGGCAGCTCCATCCA
mouse *Cyp21a1*	TGGGGATGCAAGATGTGGTGGT	GGTCGGCCAGCAAAGTCCAC
mouse *Cyp11a1*	GGATGCTGGAGGAGATCGT	GAAGTCTGGAGGCAGGTTGA
mouse *Cd31*	TGCAGGAGTCCTTCTCCACT	ACGGTTTGATTCCACTTTGC
mouse *Cd45*	CCAGTCATGCTACCACAACG	TGGACATCTTTGAGGTCTGCC
mouse *Irg1*	CTCCCACCGACATATGCTGC	GCTTCCGATAGAGCTGTGA
mouse *Il-1r1*	TGGAAGTCTTGTGTGCCCTT	TCCGAAGAAGCTCACGTTGT
mouse *Dnmt1*	CTGGAAGAGGTAACAGCGGG	CGTCCAAGTGAGTTTCCGGT
human *18S*	TGCCCTATCAACTTTCGATG	GATGTGGTAGCCGTTTCTCA
human *SDHB*	CAAGGCTGGAGACAAACCTCA	GGGTGCAAGCTAGAGTGTTG
human methylated *SDHB* promoter	AGTGGGTCCTCAGTGGATGTA	GGCGATAGTTTGGTGGCAGA
human unmethylated *SDHB* promoter	CGCGATGTTCGACGGGATA	CTTCACACCCCGCAAATCTC