RESEARCH ARTICLE
Long-term population decline of a genetically homogenous continental-wide top Arctic
predator

Marianne Gousy-Leblanc[1*], Jean-François Therrien[1,2], Thomas Broquet[3], Delphine Rioux[4], Nadine Curt-
Grand-Gaudin[4], Nathalie Tissot[4], Sophie Tissot[4], Ildiko Szabo[5], Laurie Wilson[6], Jack T. Evans[7], Victoria
Bowes[8], Gilles Gauthier[9], Karen L. Wiebe[10], Glenn Yannic[4#], Nicolas Lecomte[1*#]

[1]Canada Research Chair in Polar and Boreal Ecology, Department of Biology, University of Moncton, Moncton, NB, Canada;
Central d'Études Nordiques and Quebec Central of Biodiversity Science, QC, Canada; [2]Hawk Mountain Sanctuary, Acopian
Center for Conservation Learning, Orwigsburg, PA 17961, USA;[3]UMR 7144, Station Biologique de Roscoff, CNRS &
Sorbonne Université, Roscoff, France; [4]Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LECA, Grenoble,
France;[5]Collections Curator, Cowan Tetrapod Collection, UBC Beaty Biodiversity Museum, 2212 Main Mall, Vancouver,
BC, Canada; [6]Wildlife Biologist, Environment and Climate Change Canada, Pacific Wildlife Research Centre, RR#1 Delta,
BC, Canada;[7]Wildlife Biologist, Ministry of Forests, Lands, Natural Resource Operations and Rural Development, Surrey,
BC, Canada; [8]Avian Pathologist, Animal Health Center, BC Ministry of Agriculture, Abbotsford, BC, Canada;[9]Department de
Biologie and Centre d'Études Nordiques, Université Laval, QC, Canada; [10]University of Saskatchewan, Department of
Biology, 112 Science Place, Saskatoon, SK, Canada; #Co-senior authors: These authors contributed equally to the paper.*
Corresponding authors: Marianne Gousy-Leblanc, marianne.gousyl@gmail.com; Nicolas Lecomte,
nicolas.lecomte@umoncton.ca
19
ORCID ID : Marianne Gousy-Leblanc: 0000-0001-7805-8120-Glenn Yannic: 0000-0002-6477-2312- Jean-François Therrien: 0000-0002-1046-
0962- Nicolas Lecomte: 0000-0002-8473-5375- Thomas Broquet: 0000-0002-2986-2822- Laurie Wilson: 0000-0002-1972-1196- Gilles
Gauthier: 0000-0002-2624-3508

ABSTRACT

Genetic analysis can provide valuable information for conservation programs by unraveling the

demographic trajectory of populations, by estimating effective population size, or by inferring

genetic differentiation between populations. Here, we investigated the genetic differentiation

within the Snowy Owl (*Bubo scandiacus*), a species identified as vulnerable by the IUCN, to (i)

quantify connectivity among wintering areas, (ii) to evaluate current genetic diversity and

effective population size and (iii) to infer changes in the historical effective population size

changes from the last millennia to the recent past. The Snowy Owl, a highly mobile top predator,

breeds across the Arctic tundra which is a region especially sensitive to current climate change.

Using SNP-based analyses on Snowy Owls sampled across the North American nonbreeding

range, we found an absence of genetic differentiation among individuals located up to 4,650 km

apart. Our results suggest high genetic intermixing and effective dispersal at the continental scale

despite documented philopatry to nonbreeding sites in winter. Reconstructing the population

demographic indicated that North American Snowy Owls have been steadily declining since the

Last Glacial Maximum ca 20,000 years ago and concurrently with global increases in

temperature. Conservation programs should now consider North American Snowy Owls as a

single, genetically homogenous continental-wide population which is most likely sensitive to the

long-term global warming occurring since the Last Glacial Maximum.

Keywords: Snowy Owl, genetic structure, *Bubo scandiacus*, conservation, SNPs, effective population

size

INTRODUCTION

Natural ecosystems are currently changing at an unpreceded rate as a consequence of global

climate change (IPCC 2018, Taylor *et al.* 2020). These changes are particularly apparent in the

Arctic, where climate warming is estimated to be three times as fast as the global average (ACIA

2004, Box *et al.* 2019, Meredith *et al.* 2020). Species that live in the Arctic are therefore facing a

variety of changes and challenges (Gilg *et al.* 2012), and may respond by shifting their

geographic ranges, adjusting behaviour or phenotypes, or adapting to the new local conditions

(Parmesan 2006, Brown *et al.* 2016, Kelly 2019). In a scenario where environments are changing

so quickly and in complex ways, genetic information may be an efficient tool to inform

conservation and management programs (Schwartz *et al.* 2007, Hoban *et al.* 2021a, Hohenlohe *et*

al. 2021). First, understanding genetic structure allows one to infer dispersal of individuals and

hence connectivity among populations. The genetic structure of populations is also important for

wildlife conservation because it provides a baseline of the geographic scale at which programs

should be implemented and helps to define populations and management units (Fraser &

Bernatchez 2001, Funk *et al.* 2012, Yannic *et al.* 2016). Furthermore, genetic tools provide key

information on the current effective population size of populations (i.e. an essential parameter for

monitoring vital rate ;Hoban et al. 2013, 2021b), are useful when calculating the loss of diversity

due to inbreeding or intraspecific hybridization (Yannic *et al.* 2017), and help to predict the

ability of populations to persist and adapt to new environmental conditions (Hoffmann & Sgrò

2011). In addition, the long-term demographic trajectory of a species may also be retrieved from

genetic data (Cristofari *et al.* 2018, Cole *et al.* 2019, Cleary *et al.* 2021), which can help to

understand how species have responded to past major climatic and environmental changes. Thus,

understanding, predicting, and managing biodiversity responses to rapid climate change demand

a full consideration of the genetic differentiation within a species, an assessment of the species'

evolutionary potential and knowledge of demographic trends.

The Snowy Owl (*Bubo scandiacus*), declining according to the IUCN (BirdLife International

2020), provides a case study of how conservation actions can be informed and refined using

genetic information. This owl is a highly mobile top predator which breeds exclusively on the

Arctic tundra (Holt et al. 2020). In Canada, Snowy Owls are declining at an estimated -0.03 to -

2.85% rate per year depending on the region (Christmas bird counts; Meehan et al. 2018). Using

very different estimation and extrapolation methods, the global Snowy Owl breeding population

was first estimated at *ca* 290,000 individuals (Rich *et al.* 2004) and then more recently at *ca*

28,000 mature individuals (Potapov & Sale 2012). Both estimates are, however, probably

inaccurate because they are not based on actual continental-wide surveys (Fuller *et al.* 2003,

Therrien *et al.* 2014, Holt *et al.* 2020).

Currently, there is little genetic information on Snowy Owl populations. A previous study based

on mtDNA concluded there was little phylogeographic differentiation across the geographic

range of the species, suggesting one global panmictic population (Marthinsen *et al.* 2009). These

authors estimated the global species effective population size (*Ne*) at *ca* 14,000 individuals. This

estimate, derived from mtDNA, corresponds to the number of breeding females and does not

equate to the whole population size as it is considered by IUCN (BirdLife International 2020).

Indeed, not only is *Ne* not a direct proxy for census population size (e.g., Ferchaud et al. 2016),

but this estimate disregards the number of breeding males or the variance in reproductive success

among individuals (Storz *et al.* 2001, Wang *et al.* 2016). Therefore, further research using

genome-wide nuclear data would provide a better understanding of the genetic structure of the

population and a more accurate estimate of the current effective population size of Snowy Owls.

Snowy Owls exhibit a diversity of movement strategies throughout the year (McCabe *et al.*

2021). During the non-breeding season, individuals regularly overwinter in the Canadian Prairies

and the American Great Plains (Chang & Wiebe 2018), and throughout the entire temperate

regions of North America during irruption years (Kerlinger & Lein 1988, Robillard *et al.* 2016,

Therrien *et al.* 2017, Holt *et al.* 2020). Conversely, some individuals stay in the Arctic

throughout winter (Therrien *et al.* 2011, Robillard *et al.* 2018). Snowy Owls from the central

regions of North America appear to follow similar and constant migration routes from breeding

to winter areas (Curk et al. 2020; K. L. Wiebe personal communication), and other studies

concluded there was some site fidelity to wintering areas by Snowy Owls (Therrien *et al.* 2011,

Robillard *et al.* 2018). In contrast, individuals from the eastern part of the North-American

continent have non-regular and unpredictable movement patterns (Therrien *et al.* 2014, Robillard

et al. 2018). In summer, there is evidence that Snowy Owls use breeding sites far apart between

consecutive years, e.g. 725 km on average and up to 2, 224 km, for individuals in eastern Canada

(Therrien et al. (2014) and 1,088 km for individuals in northern Alaska; Fuller et al. 2003),

suggesting a lack of breeding site fidelity for the species (Therrien *et al.* 2014, Doyle *et al.* 2017,

Holt *et al.* 2020). The large-scale movements of the species among breeding and nonbreeding

seasons suggest there could be widespread connectivity among populations. Thus, because

Snowy Owls in different parts of their geographic range exhibit both regular and predictable

migratory movements and also large-scale unpredictable movements, our primary objective was

to study how such movements affect overall genetic structure of the species at a continental

scale.

Using SNP data, we specifically tested whether the large dispersal capacity of the species

resulted in a high degree of genetic mixing or whether, on a continental scale within North

America, the owls showed sufficient wintering site fidelity and non-random migration patterns to

exhibit population structure. In a conservation context, we also attempted to estimate the current

effective population size (Ne) of the species in North America. Our last objective was to

investigate the long-term population trajectory of Snowy Owls in North America since the Last

Glacial Maximum (~ 21k years ago) and beyond. Understanding how past shifts in climate may

have affected population size might provide insight into the resilience of wildlife resilience to

climate change in the near future. Past climatic fluctuations deeply affect long-term (historical)

effective population size for Arctic terrestrial (Prost *et al.* 2010, Lorenzen *et al.* 2011, Yannic *et*

al. 2014) and marine species (Louis *et al.* 2020, Cleary *et al.* 2021), and this never been explored

for Snowy Owls, an important species which structures the Arctic trophic network (Legagneux *et*

al. 2012, Gauthier *et al.* 2013).

METHODS

Sample Collection

We used feathers (*n*=74), blood (*n*=2) and tissue (liver [*n*=48], muscle [*n*=26]) from Snowy Owls

collected via live-trapping or carcass collection (brought to rehabilitation centers, government

agencies, and veterinary laboratories). Sample collection covered most of the southern

nonbreeding region in North America, including British Columbia (*n*=50), Saskatchewan (*n*=29),

Minnesota (*n=3*), Ontario (*n=1*), Wisconsin (*n=9*), Michigan (*n=16*), Maryland (*n=3*),

Pennsylvania (*n=2*), New York (*n=34*), Québec (Nunavik; *n=1*), Nova Scotia (*n=1*), and Prince

Edward Island (*n=1*; Figure 1, Table 1, Supplementary Material Table S1). Sampling spanned a

4,650-km distance. We collected samples during winter between 2012 and 2018 (except for two

samples that were collected in 2007 and 2008). For each individual, we assessed sex and age

class (first year or adult) according to morphological and molt measurements (Solheim 2012).

The spatial distribution of the samples was slightly clustered into four groups (i.e. wintering

areas) according to longitude (see Figure 1). We called these wintering areas: West (W: British

Columbia); Central (C: Saskatchewan); Great Lakes areas (GL: Michigan, Wisconsin, Ontario,

and Maryland); and East (E), which included all other individuals (including the one from

Nunavik; Figure 1).

DNA Extraction

We used Qiagen DNeasy Blood and Tissue kit (Qiagen, Inc., Valencia, CA, USA) to extract

genomic DNA following the manufacturer's instructions and assessed DNA quality and checked

for DNA degradation on agarose gels. We quantified DNA concentration for each sample using

the QuantiFluor dsDNA System kit (Promega), using samples with a minimum DNA

concentration of 3.20 ng/µl after extraction for subsequent double digested RAD sequencing

(ddRADSeq) library preparation.

ddRADSeq Library Construction and Sequencing

We built four ddRADSeq libraries from the Snowy Owl genomic DNA following a modified

version of the protocol in Peterson et al. (2012) and detailed in Gagnon et al. (2019). We

randomized the individuals for each wintering area in each library. The Genomic DNA (100 ng)

was digested using the enzymes SbfI and MspI and fragments selected between 300 bp and 500

bp using BluePippin size-selection system (Sage Science). To control for library quality and

consistency, we replicated 15.2 % of our samples (*n*=28) following the recommendations of

Mastretta-Yanes et al. (2015). We included 47 samples per library including replicates and a

negative control. The four libraries were then sequenced on two full lanes (two libraries per lane)

of paired-end (2 x 125bp) Illumina Hi-Seq 2500 (Fasteris SA, Switzerland).

ddRADSeq Data Processing

We used Stacks*1.44 (Catchen et al. 2011, 2013) to demultiplex data, build a *de novo* SNP

catalog and call genotypes. Following Mastretta☐Yanes et al. (2015), we tested different sets of

Stacks core parameters (ustack – m (2 to 6), –M (2 to 6) and –max_locus_stacks (2 to 6) and

cstack –n (0 to 5) – Supplementary Material Figure S1 and S2), by varying one parameter at a

time while holding the others at their default values. We then selected the set of parameters that

minimized error rates between replicates (*n* = 28 pairs) and maximized the amount of data

recovered. The optimal values were: –m (5), –M (2), –max_locus_stacks (3) and –n (2; error

rates for each set of parameters are presented in Supplementary Material Figure S1 and S2). To

produce the final data set, we ran Stacks with all parameters set to their optimal values. We

performed the next filtering steps in *R**3.6.2 (R Core Team 2019) from the original VCF file

obtained from stacks and by keeping only loci that were typed for at least 85% of samples and

only samples that were typed for at least 80% of loci. We then used *dartR* (Gruber *et al.* 2018)

and *Radiator* (Gosselin *et al.* 2017) packages in *R* and *PDGspider* (Lischer & Excoffier 2012) to

convert SNPs VCF files to other formats (e.g. genlight; Jombart 2008).

Population Genetic Structure, Genetic Distance, and Isolation-by-distance

We used complementary analyses to quantifying population genetic structure. Unless mentioned,

all analyses were in *R**3.6.2 (R Core Team 2019). We first used Principal Component Analysis

(PCA), considering the a priori defined wintering regions with the function *glPCA* using the

package *Adegenet* *2.1.1 (Jombart 2008). Next, to compute pairwise F_{ST} among regions

according to Weir and Cockerham (1984) we used the *gl.fst.pop* in the *dartR* package (Gruber et

al. 2018; See the Supplementary Material Appendix). Also, we performed a Discriminant

Analysis of Principal Components (DAPC; Jombart et al. 2010; also implemented in the package

Adegenet (Jombart 2008)) to infer the number of genetic clusters that best fit our data. In

addition, we used a Maximum Likelihood approach using ADMIXTURE*1.3 software (Alexander

et al. 2009) to estimate the number of populations underlying the genetic dataset based on

individual relatedness without our *a priori* wintering regions. Finally, we tested for Isolation-by-

distance at the individual level by investigating the between the natural logarithm of Euclidean

distance and relatedness (i.e. *beta*, according to Weir and Goudet 2017) distance matrices

between each pair of individuals. An extended description of the methods is provided in

Supplementary Material Appendix.

Descriptive Genetic Diversity

We calculated the observed heterozygosity (*Ho*) and the expected heterozygosity (*He*) for each

locus using the function *gl.Ho* and *gl.Hs* in the *dartR* (Gruber *et al.* 2018) package in *R**3.6.2 (R

Core Team 2019). We also computed *Ho* and *Hs* for each wintering area. To test whether a

significant difference in *Hs* among wintering areas can exist, we performed a Linear Mixed

Model (LMM) with wintering area as the fixed effect and the loci ID as a random effect. We

verified linearity assumptions of model residuals, and we then computed a post-hoc Tukey test to

contrast wintering areas.

Current Effective Population Size (*Ne*)

We used NEESTIMATOR*2.1 (Do *et al.* 2014) to estimate the current effective population size

(*Ne*) of Snowy Owls in North America. We used the linkage disequilibrium (LD) method with a

random mating model. Following Waples and Do (2010), we excluded rare alleles (i.e., those

SNP with a minor allele frequency ≤ 0.05) to avoid estimation bias.

Population Trajectory Reconstruction

We reconstructed the population history of Snowy Owls using the Stairway Plot 2 method (Liu and Fu 2020). The Stairway Plot infer detailed population demographic history using the site frequency spectrum (SFS: Liu and Fu 2015) from DNA sequence data. This method uses a flexible, multi-epoch model as used in the Skyline Plot method (Strimmer and Pybus 2001, Navascués et al. 2017), and based on the expected composite likelihood of a given SNP frequency spectrum (SFS). It provides information on the history of population effective size (*Ne*) over time until the recent past. We estimated the folded SFS using the VCF file and a Python script, *EasySFS* (https://github.com/isaacovercast/easySFS; Covercast 2017) and included the total number of observed nucleic sites (both polymorphic and monomorphic sites). The "two-epoch" model, with 67% of sites for training and 200 bootstraps was used as recommended. We tested four different numbers of breaking points (i.e. to define the boundaries of each epoch; Liu and Fu 2020) as described in Liu and Fu (2015), i.e. 74, 149, 223, and 298. We assumed a mutation rate of 1.9×10^{-9} which is a recent estimate for birds in general (Zhang et al. 2014) and a generation time of 7.93 years (a generally accepted value; BirdLife International 2020). However, because we have no information on the mutation rate in Snowy owls, and because there is also some uncertainty over generation times in the literature (e.g. 4.7 years, Heggøy et al. 2017; 7.93, BirdLife International 2020), we ran sensitivity analyses using various mutation rates retrieved from the literature on birds (i.e. $\mu = 4.6 \times 10^{-9}$, Smeds et al. 2016); $\mu = 1.9 \times 10^{-9}$; Nam et al. 2010, Zhang et al. 2014), and ranges of generation times. We also added the major past glaciation events to the resulting Stairway plot (Clark *et al.* 2009, 2012, Mann *et al.* 2009).

RESULTS

SNPs Genotyping

After the *de novo* SNP calling and filtering procedure, the data set encompassed 13,793 SNPs

distributed over 5,987 loci. We only kept the SNPs that had less than 15% of missing data, i.e.,

that were typed for at least 85% of birds. Individuals scored on average 5,767 SNPs ± 198 (SD)

[4,826; 5,921], resulting in 3.7% missing data in the genotype matrix. Error rates estimated on

the 13,793 SNPs obtained from replicated samples were 0.004 +/- 0.003 and dropped to 0.003

+/- 0.002 when considering single SNP per loci (*n*=5,987 SNPs). For subsequent analyses, we

only considered a single SNP per locus, for a final dataset consisting of 5,987 SNPs genotyped

on 150 Snowy Owl individuals.

Population Genetic Structure, Genetic Distance, and Isolation-by-distance

Results from the PCA (Supplementary Material Figure S3), the DACP (Supplementary Material

Figure S4), and from ADMIXTURE (Supplementary Material Figure S5-S6) were congruent and

all revealed a single genetic cluster (*K*=1). When ADMIXTURE was conducted separately on each

dataset of yearling owls versus the adult owls, both also revealed a single genetic cluster (*K*=1).

Pairwise F_{ST} between wintering areas averaged $7e^{-5} ± 3e^{-4}$ (Figure 1). However, pairwise F_{ST}

analysis on adults from W and E, the two most distant wintering areas, revealed a small, but

significant genetic differentiation ($F_{ST} = 0.002$; $p = 0.017$; Figure 1 and Supplementary Table

S2), which also held for the adult females between W and E ($F_{ST} = 0.002$; p = 0.006;

Supplementary Table S2) and the adult females between W and GL ($F_{ST} = 0.001$; $p = 0.042$;

Supplementary Table S2). There was also a low genetic differentiation between males from W

and C ($F_{ST} = 0.002$; $p=0.042$; Supplementary Table S2). Global F_{IS} was 0.0332 ± 0.2431. Mean

F_{IS} were low to moderate (but all significantly different from zero; all Wilcoxon signed rank tests

p-values < 0.001) and were similar within wintering areas: W= 0.037 ± 0.19; C= 0.038 ± 0.21;

GL = 0.048 ± 0.22 and E = 0.040 ± 0.20.

On average, genetic relatedness was 0.017 ± 0.014 (Figure 2), and the maximum value between

two individuals was 0.093, one sampled in the E wintering area (New York) and one in the GL

area (Michigan; 2,763 km apart). The minimum value between the two individuals was 1.98 x10^{-}

^{6}, one from the E (New York) and one from the GL areas (Wisconsin; 2,457 km apart). Genetic

distance was similar within and between wintering areas (Figure 2) and between adults and first

year (Figure S7). We did not find any pattern of Isolation-by-distance (at a scale of 4,650 km), i)

at the individual level (i.e. relatedness vs ln-transformed geographic distance): Mantel's $r =$

0.005, $p = 0.39$ and LMM slope=-8.87 e^{-5}, 95% CI= -1.86e^{-4}; 9.30e^{-6}, and ii) at the population

level (i.e. $F_{ST}/(1-F_{ST})$ vs ln-transformed geographic distance): Mantel's $r = -0.258$, $p = 0.70$ and

LMM; slope=-3.69 e^{-4}, 95% CI= -6.8 e^{-4}; 4.75e^{-5}).

Descriptive Genetic Diversity

At the locus scale, the average observed heterozygosity (Ho) per locus was 0.142 ± 0.143 (SD)

and the expected heterozygosity per locus (He) was 0.148 ± 0.140. At the wintering area scale,

we found a significant decrease of gene diversity from West to East (Hs; slope= -1.847e^{-2}, SE=

5.664e^{-3}, 95% CI= -0.0296, -0.00737; Figure 2).

Curent Effective Population Size (Ne)

Using the LD method with a lowest allele frequency set at 0.05, the current effective population

size (Ne) for the North American population of Snowy Owls was estimated to be 15,792

individuals (95% CI 10,850-28,950). Excluding singleton alleles only, the Ne value was 15,401

(95% CI 12,782-19,366).

Population Trajectory Reconstruction

The Stairway plot approach showed that the coalescence Ne of Snowy Owls in North America

declined steadily in recent times (Figure 3). There was an expansion of Ne around 100,000 years

ago, followed by a long period of stability. Coalescence Ne then started to decline around 6,000

years ago, which coincides with an acceleration of deglaciation in North America between 6,000

and 8,000 years ago. This is also approximately timed with the beginning of the Holocene

period, which was characterized by global temperature increases. During the Little Ice Age

period (*ca* 1,400-1,700 AD), the rate of decline in coalescence *Ne* apparently slowed down. The

decline started to speed up again around 200 years ago, approximately at the onset of the

worldwide retreat of glaciers and the acceleration of air temperature warming (Figure 3). We

cannot infer population changes for the last 100 years because of the uncertainties in the final

steps of the Stairway Plot method (Liu and Fu 2015, 2020). The sensitivity analysis showed that

despite using different values for mutation rate or generation time, all demography simulations

resulted in a declining trend (Supplementary Material Figure S8).

DISCUSSION

Continent-scale Genetic Homogeneity

Based on the information retrieved from Snowy Owls on the wintering grounds, our results all

indicate that populations at the continental scale are genetically homogenous in North America.

Two non-mutually exclusive hypotheses could explain the absence of genetic differentiation on a

4,500 km scale: breeding dispersal or juvenile (natal) dispersal. Irrespective of their wintering

areas, if owls disperse widely during the pre-breeding period or between successive reproduction

events (i.e., breeding dispersal, Clobert et al. 2012) individuals from different wintering areas

may come into contact, mate, and breed together in the high Arctic. There is some evidence that

Snowy Owls have low breeding site fidelity and that long distances may separate breeding sites

in consecutive years (Fuller et al. 2003, Therrien et al. 2014, Doyle et al. 2017, Holt et al. 2020).

Thus, mating is likely panmictic and not related to the winter origin of birds.

A second possibility is that high juvenile dispersal from the site of birth to the site of

reproduction is driving the genetic mixing. Natal dispersal distance is typically much larger than

breeding dispersal in many birds (Clobert et al. 2012) and probably plays a role in genetic

intermixing in other owl species such as the Northern Spotted Owl; *Strix occidentalis caurina*;

(Miller *et al.* 2018), but nothing is known about natal dispersal in Snowy Owls. This hypothesis

could explain the absence of genetic differentiation if juveniles are dispersing among wintering

areas during the first years of life, i.e. before their first reproduction. Alternatively, genetic

homogenization could occur as a result of periodic winter irruptions when, after a summer with

high productivity linked to abundant lemming prey in the Arctic, large numbers of juveniles

travel farther south than average during their first winter (Kerlinger & Lein 1988, Potapov &

Sale 2012, Holt *et al.* 2020). High gene flow is probably facilitated by the high mobility and

nomadic behavior of the species, allowing high breeding and natal dispersal or a combination of

both (Yannic et al. 2016).

The pairwise F_{ST} values among winter areas of Snowy Owls (ranging -0.0003 to 0.0004) are low

compared to the range of values based on SNPs from other raptor species. Gousy-Leblanc *et al.*

(2021) recently reviewed genetic differentiation within raptors and reported values for four

species based on data from breeding grounds at smaller scales than our current study and with

different ecology than Snowy Owls (*i.e*., fully sedentary, or partially migratory species): Great

Gray Owl *Strix nebulosa* F_{ST} = 0.03-0.15 (Mendelsohn *et al.* 2020), Prairie Falcon *Falco*

mexicanus F_{ST} = 0.01-0.03 (Doyle *et al.* 2018), Bald Eagle *Haliaeetus leucocephalu; F_{ST} =*

0.037-0.203 (Judkins *et al.* 2019), and Northern Goshawk *Accipiter gentilis; F_{ST} = 0-0.093*

(Geraldes *et al.* 2019). All estimates were at least an order of magnitude higher than those we

found in Snowy Owls, and none of these species had movements as nomadic as those observed

in Snowy Owls.

Conversely, the Gyrfalcon (*Falco rusticolus*), an Arctic raptor which inhabits similar habitats as

the Snowy Owl, shows genetic differentiation between the continental populations of Greenland

and Iceland, despite the high dispersal capability of the species (Johnson *et al.* 2007). For the

Eurasian Arctic Peregrine Falcon (*Falco peregrinus*), four distinct genetic groups were detected

at the continental scale, which supposedly arose from the use of distinct population-specific

migration routes, and fidelity to breeding and non-breeding areas (Gu *et al.* 2021). At such a

large scale and for species with a high dispersal capacity, various factors like fidelity to breeding

or non-breeding areas, low breeding and natal dispersal, the use of distinct migration routes, and

the interplay between those factors could impede gene flow among populations, promoting

genetic differentiation.

Current Genetic Diversity and Effective Population Size (*Ne*)

For Snowy Owls, we found a slight statistically significant genetic diversity between the most

distant regions, i.e., the Western and Eastern regions in North America. This could be explained

by the potential inclusion (i.e. via migration) of individuals in the "Western" category which

originated from Alaska or Russia but whose dispersal was not far enough to reach the Eastern

population (Fuller *et al.* 2003, Holt *et al.* 2020). Such heterogeneous migration could slightly

influence allele frequencies across the circumpolar distribution, while not greatly impacting the

genetic differentiation overall. Indeed, we observed that the highest *Hs* occured in the West. The

global population should nonetheless be considered as one population unit genetically.

The average expected heterozygosity (0.148 ± 0.140 SD) at the locus scale for Snowy Owls is

slightly lower compared to the *He* values computed with SNPs for three other species of raptors.

Two species that were partial migrants displayed genetic differentiation at large scale, *He* =

0.293 in the Bald Eagle, *H. leucocephalus* (Judkins *et al.* 2019) and *He* = 0.340 in the Golden

Eagle *Aquila chrysaetos* (Doyle *et al.* 2016). One obligate migrant species, the Prairie Falcon

Falco mexicanus, showed no population genetic differentiation at large scale *He* = 0.332 (Doyle

et al. 2018). However, *He* in Snowy Owls is quite similar to the Burrowing Owl *Athene*

cunicularia, another nomadic facultative migrant species (*He* = 0.112) that, exhibited population

genetic differentiation at low scale (Mueller et al. 2018).

Although Snowy Owls appear to have lower gene diversity than other raptors, the estimated

current effective population size of the species in North America (*Ne* = 15,792 individuals) is

large enough to preserve the evolutionary potential of the population and allow it to persist and

adapt in a changing environment as suggested by population modelling (see Kamath et al. (2015)

and Hoban et al. 2020). In general, a minimum *Ne* of 100 individuals is recommended to prevent

loss of genetic diversity by genetic drift and a *Ne* of 1,000 individuals to maintain long-term

evolutionary potential (Frankham *et al.* 2014). In a conservation context, *Ne/Nc* where *Nc* is the

number of individuals counted in a census population, is important for disentangling the relative

effects of population size and genetic factors on the persistence of species (Frankham 1995,

Palstra & Ruzzante 2008, Ferchaud *et al.* 2016, Waples 2016). Considering our estimated *Ne* of

15,792 individuals (95% CI between 10,850- 28,950) and a *Nc* of about 30,000 individuals in

North America (Rosenberg *et al.* 2016, BirdLife International 2020), we can calculate a *Ne*/*Nc*

ratio of about 0.57 [range 0.36 - 0.96]. Estimates of *Nc* for Snowy Owls, are extremely uncertain

and may be as low as 7,000-8,000 pairs in North America (Potapov & Sale 2012, BirdLife

International 2020). *Ne/Nc* ratio values derived from these lower *Nc* estimates would then be

around 1.05 [0.77 - 1.81]. All these *Ne/Nc* values are nonetheless higher than those in many other

avian taxa (Frankham 1995) so our results suggest that the North American population as a

whole does not face an imminent risk of inbreeding depression or genetic impoverishment.

Past Population Trajectory

Demographic reconstruction suggests that the population size of Snowy Owls declined over the

last few thousand years, starting around the Last Glacial Maximum (LGM). A recent

reconstruction of demographic history in the Peregrine Falcon (*Falco peregrinus*), another Arctic

raptor, resulted in a similar population size trajectory as the Snowy Owl (i.e. expansion during

the LGM and a steady decline since the beginning of the Holocene; Gu et al. 2021). Perhaps an

increase in the total area of tundra habitat may explain the population expansion during the LGM

and the subsequent decline in the population occurred when the tundra began to contract

northward (Gu *et al.* 2021). Similarly, the population size of Snowy Owls may be closely

matched to the amount of Arctic tundra because the species breeds exclusively in that habitat. An

implication is that large and long-term environmental changes over the last few thousand years,

and in particular to temperature increases since the LGM (Clark et al. 2009) may greatly affect

the population size of Snowy Owls. Indeed, population reconstructions showed that temperature

increases reduced the effective population size of three penguin genera (*Eudyptes*, *Pygoscelis*,

and *Aptenodytes*) in Antarctica (Cole *et al.* 2019), highlighting ecosystem-wide responses to

climate changes in the Antarctic ocean in the past. Similarly, the climate warming that occurred

after the LGM in the Northern Hemisphere reduced effective population sizes, genetic diversity,

and/ or census counts of other Arctic species like the polar bear *Ursus maritimus* (Miller *et al.*

2012) and the Arctic fox *Vulpes lagopus* (Larsson *et al.* 2019). Global warming has also been

implicated as a major factor in the mass extinction of Late Quaternary megafauna in the

Northern Hemisphere (Lorenzen et al. 2011, Lord et al. 2020, Stewart et al. 2021; but see

Sandom et al. 2014).

Globally, the warming period of the Holocene has been an important influence on population

size and/or range distribution of small mammals including lemmings (Prost *et al.* 2010, 2013,

Lanier *et al.* 2015, Fedorov *et al.* 2020) which are the main prey of breeding Snowy Owls. Thus,

climate change has the potential to influence entire ecosystems in the Arctic by disrupting the

main trophic links between species and Snowy Owls may have declined since the mid-Holocene

both as direct result of warmer temperatures (e.g. physiological stress) and/or due to the indirect

effect of the climate-induced population reductions and geographic range contractions of its main

prey, i.e., the small mammals in the Arctic.

Conclusion

Our findings should help further define some research priorities for Snowy Owls identified by

Holt et al. (2020). First, the lack of significant genetic differentiation over the span of a continent

spanning > 4,500 km does not support the hypothesis that well-separated migration routes, i.e.,

north-south corridors to distinct winter grounds are serving to isolate distinct breeding

populations of Snowy Owls. Second, and potentially most concerning, our results showed that

Snowy Owls experienced a continuous decline in numbers since the end of the Last Glacial

Maximum (18-21k years ago) in North America. Although our estimate of current effective

population size does not suggest that Snowy Owls are at risk for imminent genetic problems, the

finding that the owls were highly sensitive to global changes in the past implies that the

population size may decline more precipitously in the future if Arctic warming accelerates, as

predicted by many climate changes models.

ACKNOWLEDGEMENTS

This work was supported by the Canada Research Chair Program, Polar Knowledge Canada,

NSERC, and the Canadian Foundation for Innovation (N.L.). M.G.L. benefited from a NSERC

and a FRQNT graduate grant. We thank C. Miquel as well as the AnaBM (USMB) and AEEM

(UGA) laboratory facilities for their help and support during the laboratory work. We thank

everyone that contributed samples, including D. Brinker, D. Genesky, G. Jacobs, T. McDonald,

S. Weidensaul, A. Robillard as well as Project SNOWstorm members and supporters. In BC, all

of the following organization contributed snowy owl carcasses: BC Ministry of Agriculture,

Food and Fisheries, BC Ministry of Environment and Climate Change staff from multiple

regions, BC Wildlife Park, Hillspring Wildlife Rehabilitation Center, Mars Wildlife Rescue

Centre, North Island Wildlife Recovery Assoc., Northern Lights Wildlife Shelter, Orphaned

Wildlife Rehabilitation Soc., Northern Wildlife Rescue, South Okanagan Rehabilitation Centre

for Owls, University of Northern British Columbia, Vancouver Airport Authority, and Wildlife

Rescue Assoc; in the Maritimes: P-Y Daoust, D. Weeks, F. de Biep. For helping to trap owls in

Saskatchewan, we thank D. Zazelenchuk and M. Stoffel. This work benefited from access to the

Biogenouest genomic platform at Station Biologique de Roscoff and we are grateful to the

Roscoff Bioinformatics platform ABiMS for providing storage and computing resources. We

thank E. Milot and A.-M. Dion-Coté for the Ms review. **Funding statement:** The Canada

Research Program, Polar Knowledge Canada, NSERC, and the Canadian Foundation for

Innovation supported N.L. M.G-L. received NSERC and FRQNT grants for her MSc. **Ethics**

statement: All live captures followed Canadian Animal Care Committees guidelines. **Author**

contributions: Conceived the idea, design, experiment**:** N.L., M.G.-L., G. Y., and J-F. T. Data

collection: M. G-L, J-F. T., I. S., L. W., J. T. E., V. B, G. G., K. L. W., and N.L. Performed the

laboratory work: D. R., N. C-G-G, S. T. and N. T. Analyzed the data: M. G-L., G. Y., T.B., N.L.,

and J-F. T. Wrote first drafts: M.G-L, G.Y., N.L., J-F. Final version of the paper revised and

agreed by all authors. **Data depository:** Data used in the article can be found on Figshare:

https://figshare.com/s/f1f01dc772d2f8c8698c.

LITERATURE CITED

ACIA. 2004. *Impacts of a warming Arctic-Arctic climate impact assessment*. Cambridge
University Press.
Alexander, D.H., Novembre, J. & Lange, K. 2009. Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res.* **19**: 1655–1664.
Bates, D., Maechler, M., Bolker, B., Walker, S. & Haubo Bojesen Christensen, R. 2015.
lme4: Linear mixed-effects models using Eigen and S4.
BirdLife International. 2020. IUCN Red List for birds.
Box, J.E., Colgan, W.T., Christensen, T.R., Schmidt, N.M., Lund, M., Parmentier, F.-J.W.,
Brown, R., Bhatt, U.S., Euskirchen, E.S. & Romanovsky, V.E. 2019. Key indicators of Arctic
climate change: 1971–2017. *Environ. Res. Lett.* **14**: 045010.
Brown, C.J., O'Connor, M.I., Poloczanska, E.S., Schoeman, D.S., Buckley, L.B., Burrows,
M.T., Duarte, C.M., Halpern, B.S., Pandolfi, J.M., Parmesan, C. & Richardson, A.J. 2016.
Ecological and methodological drivers of species' distribution and phenology responses to
climate change. *Glob. Change Biol.* **22**: 1548–1560.
Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. & Cresko, W.A. 2013. Stacks: an
analysis tool set for population genomics. *Mol. Ecol.* **22**: 3124–3140.
Catchen, J.M., Amores, A., Hohenlohe, P., Cresko, W. & Postlethwait, J.H. 2011. Stacks:
building and genotyping loci de novo from short-read sequences. *G3 Genes Genomes Genet.* **1**:
171–182.
Chang, A.M. & Wiebe, K.L. 2018. Habitat selection by wintering male and female Snowy
Owls on the Canadian prairies in relation to prey abundance and a competitor, the Great Horned
Owl. *J. Field Ornithol.* **89**: 64–77.
Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica,
J.X., Hostetler, S.W. & McCabe, A.M. 2009. The Last Glacial Maximum. *Science* **325**: 710–
714.
Clark, P.U., Shakun, J.D., Baker, P.A., Bartlein, P.J., Brewer, S., Brook, E., Carlson, A.E.,
Cheng, H., Kaufman, D.S., Liu, Z., Marchitto, T.M., Mix, A.C., Morrill, C., Otto-Bliesner,
B.L., Pahnke, K., Russell, J.M., Whitlock, C., Adkins, J.F., Blois, J.L., Clark, J., Colman,
S.M., Curry, W.B., Flower, B.P., He, F., Johnson, T.C., Lynch-Stieglitz, J., Markgraf, V.,
McManus, J., Mitrovica, J.X., Moreno, P.I. & Williams, J.W. 2012. Global climate evolution
during the last deglaciation. *Proc. Natl. Acad. Sci.* **109**: E1134–E1142.
Cleary, A.C., Hoffman, J.I., Forcada, J., Lydersen, C., Lowther, A.D. & Kovacs, K.M.
2021. 50,000 years of ice and seals: Impacts of the Last Glacial Maximum on Antarctic fur seals.
Ecol. Evol. **11**: 14003–14011.
Cole, T.L., Dutoit, L., Dussex, N., Hart, T., Alexander, A., Younger, J.L., Clucas, G.V.,
Frugone, M.J., Cherel, Y., Cuthbert, R., Ellenberg, U., Fiddaman, S.R., Hiscock, J.,
Houston, D., Jouventin, P., Mattern, T., Miller, G., Miskelly, C., Nolan, P., Polito, M.J.,
Quillfeldt, P., Ryan, P.G., Smith, A., Tennyson, A.J.D., Thompson, D., Wienecke, B.,
Vianna, J.A. & Waters, J.M. 2019. Receding ice drove parallel expansions in Southern Ocean
penguins. *Proc. Natl. Acad. Sci.* **116**: 26690–26696.

Covercast, I. 2017. *easySFS*. Github repository.
Cristofari, R., Liu, X., Bonadonna, F., Cherel, Y., Pistorius, P., Le Maho, Y., Raybaud, V.,
Stenseth, N.C., Le Bohec, C. & Trucchi, E. 2018. Climate-driven range shifts of the king
penguin in a fragmented ecosystem. *Nat. Clim. Change* **8**: 245–251.
Curk, T., Pokrovsky, I., Lecomte, N., Aarvak, T., Burnham, K., Dietz, A., Franke, A.,
Gauthier, G., Jacobsen, K.-O., Kidd, J., Lewis, S.B., Øien, I.J., Solheim, R., Wiebe, K.,
Wikelski, M., Therrien, J.-F. & Safi, K. 2020. Arctic avian predators synchronise their spring
migration with the northern progression of snowmelt. *Sci. Rep.* **10**: 7220.
Do, C., Waples, R.S., Peel, D., Macbeth, G.M., Tillett, B.J. & Ovenden, J.R. 2014.
NeEstimator v2: re-implementation of software for the estimation of contemporary effective
population size from genetic data. *Mol. Ecol. Resour.* **14**: 209–214.
Doyle, F.I., Therrien, J.-F., Reid, D.G., Gauthier, G. & Krebs, C.J. 2017. Seasonal
Movements of Female Snowy Owls Breeding in the Western North American Arctic. *J. Raptor*
Res. **51**: 428–438.
Doyle, J.M., Bell, D.A., Bloom, P.H., Emmons, G., Fesnock, A., Katzner, T.E., LaPré, L.,
Leonard, K., SanMiguel, P., Westerman, R. & Andrew DeWoody, J. 2018. New insights into
the phylogenetics and population structure of the prairie falcon (*Falco mexicanus*). *BMC*
Genomics **19**: 233.
Doyle, J.M., Katzner, T.E., Roemer, G.W., Cain, J.W., Millsap, B.A., McIntyre, C.L.,
Sonsthagen, S.A., Fernandez, N.B., Wheeler, M., Bulut, Z., Bloom, P.H. & Andrew
DeWoody, J. 2016. Genetic structure and viability selection in the golden eagle (*Aquila*
chrysaetos), a vagile raptor with a Holarctic distribution. *Conserv. Genet.* **17**: 1307–1322.
Fedorov, V.B., Trucchi, E., Goropashnaya, A.V., Waltari, E., Whidden, S.E. & Stenseth,
N.Chr. 2020. Impact of past climate warming on genomic diversity and demographic history of
collared lemmings across the Eurasian Arctic. *Proc. Natl. Acad. Sci.* **117**: 3026–3033.
Ferchaud, A.-L., Perrier, C., April, J., Hernandez, C., Dionne, M. & Bernatchez, L. 2016.
Making sense of the relationships between Ne, Nb and Nc towards defining conservation
thresholds in Atlantic salmon (*Salmo salar*). *Heredity* **117**: 268–278.
Frankham, R. 1995. Effective population size/adult population size ratios in wildlife: a review.
Genet. Res. **66**: 95–107.
Frankham, R., Bradshaw, C.J.A. & Brook, B.W. 2014. Genetics in conservation management:
Revised recommendations for the 50/500 rules, Red List criteria and population viability
analyses. *Biol. Conserv.* **170**: 56–63.
Fraser, D.J. & Bernatchez, L. 2001. Adaptive evolutionary conservation: towards a unified
concept for defining conservation units. *Mol. Ecol.* **10**: 2741–2752.
Fuller, M., Holt, D. & Schueck, L. 2003. Snowy owl movements: variation on the migration
theme. In: *Avian Migration* (P. Berthold, E. Gwinner, & E. Sonnenschein, eds), pp. 359–365.
Springer, Berlin, Germany.
Funk, W.C., McKay, J.K., Hohenlohe, P.A. & Allendorf, F.W. 2012. Harnessing genomics
for delineating conservation units. *Trends Ecol. Evol.* **27**: 489–496.
Gagnon, M., Yannic, G., Perrier, C. & Côté, S.D. 2019. No evidence of inbreeding depression
in fast declining herds of migratory caribou. *J. Evol. Biol.* **32**: 1368–1381.
Gauthier, G., Bêty, J., Cadieux, M.-C., Legagneux, P., Doiron, M., Chevallier, C., Lai, S.,
Tarroux, A. & Berteaux, D. 2013. Long-term monitoring at multiple trophic levels suggests
heterogeneity in responses to climate change in the Canadian Arctic tundra. *Philos. Trans. R.*
Soc. B Biol. Sci. **368**: 20120482.
Geraldes, A., Askelson, K.K., Nikelski, E., Doyle, F.I., Harrower, W.L., Winker, K. &

Irwin, D.E. 2019. Population genomic analyses reveal a highly differentiated and endangered
genetic cluster of northern goshawks (*Accipiter gentilis laingi*) in Haida Gwaii. *Evol. Appl.* **12**:
757–772.

Gilg, O., Kovacs, K.M., Aars, J., Fort, J., Gauthier, G., Grémillet, D., Ims, R.A., Meltofte,
H., Moreau, J., Post, E., Schmidt, N.M., Yannic, G. & Bollache, L. 2012. Climate change and
the ecology and evolution of Arctic vertebrates: Climate change impacts on Arctic vertebrates.
Ann. N. Y. Acad. Sci. **1249**: 166–190.

Gosselin, T., Lamothe, M. & Grewe, P. 2017. *Radiator: RADseq Data Exploration, m*
anipulation and Visualization using R.

Gousy-Leblanc, M., Yannic, G., Therrien, J.-F. & Lecomte, N. 2021. Mapping our
knowledge on birds of prey population genetics. *Conserv. Genet.* **22**: 685–702.

Gousy-Leblanc M., Therrien J.-F., Broquet T., Rioux D., Curt-Grand-Gaudin N., Tissot,
Tissot S., Szabo I., Wilson L., Evans J.T., Bowes V., Gauthier G., Wiebe K.L., Yannic G.,
Lecomte N. 2022. Long-term population decline of a genetically homogenous continental-wide
top Arctic predator. Figshare, Dataset, https://figshare.com/s/f1f01dc772d2f8c8698c.

Gruber, B., Unmack, P.J., Berry, O.F. & Georges, A. 2018. DARTR☐: An R package to
facilitate analysis of SNP data generated from reduced representation genome sequencing. *Mol.*
Ecol. Resour. **18**: 691–699.

Gu, Z., Pan, S., Lin, Z., Hu, L., Dai, X., Chang, J., Xue, Y., Su, H., Long, J., Sun, M.,
Ganusevich, S., Sokolov, V., Sokolov, A., Pokrovsky, I., Ji, F., Bruford, M.W., Dixon, A. &
Zhan, X. 2021. Climate-driven flyway changes and memory-based long-distance migration.
Nature **591**: 259–264.

Heggøy, O., Aarvak, T., Øien, I.J., Jacobsen, K.-O., Solheim, R., Zazelenchuk, D., Stoffel,
M. & Kleven, O. 2017. Effects of satellite transmitters on survival in Snowy Owls *Bubo*
scandiacus. *Ornis Nor.* **40**: 33.

Hoban, S., Bruford, M.W., Funk, W.C., Galbusera, P., Griffith, M.P., Grueber, C.E.,
Heuertz, M., Hunter, M.E., Hvilsom, C., Stroil, B.K., Kershaw, F., Khoury, C.K., Laikre,
L., Lopes-Fernandes, M., MacDonald, A.J., Mergeay, J., Meek, M., Mittan, C., Mukassabi,
T.A., O'Brien, D., Ogden, R., Palma-Silva, C., Ramakrishnan, U., Segelbacher, G., Shaw,
R.E., Sjögren-Gulve, P., Veličković, N. & Vernesi, C. 2021a. Global Commitments to
Conserving and Monitoring Genetic Diversity Are Now Necessary and Feasible. *BioScience* **71**:
964–976.

Hoban, S., Paz-Vinas, I., Aitken, S., Bertola, L., Breed, M.F., Bruford, M., Funk, C.,
Grueber, C., Heuertz, M. & Hohenlohe, P. 2021b. Effective population size remains a
suitable, pragmatic indicator of genetic diversity for all species, including forest trees. *Biol.*
Conserv. **253**: 108906.

Hoban, S.M., Hauffe, H.C., Pérez-Espona, S., Arntzen, J.W., Bertorelle, G., Bryja, J., Frith,
K., Gaggiotti, O.E., Galbusera, P., Godoy, J.A., Hoelzel, A.R., Nichols, R.A., Primmer,
C.R., Russo, I.-R., Segelbacher, G., Siegismund, H.R., Sihvonen, M., Vernesi, C., Vilà, C. &
Bruford, M.W. 2013. Bringing genetic diversity to the forefront of conservation policy and
management. *Conserv. Genet. Resour.* **5**: 593–598.

Hoffmann, A.A. & Sgrò, C.M. 2011. Climate change and evolutionary adaptation. *Nature* **470**:
479–485.

Hohenlohe, P.A., Funk, W.C. & Rajora, O.P. 2021. Population genomics for wildlife
conservation and management. *Mol. Ecol.* **30**: 62–82.

Holt, D.W., Larson, M.D., Smith, N., Evans, D. & Parmelee, D.F. 2020. The snowy owl
(*Bubo scandiacus*). In: *Birds of the World* (S. M. Billerman, ed). Cornell Lab of Ornithology,

Ithaca, New York, USA.

IPCC. 2018. Global warning of 1.5 C.

Johnson, J.A., Burnham, K.K., Burnham, W.A. & Mindell, D.P. 2007. Genetic structure
among continental and island populations of gyrfalcons: Gyrfalcon population structure. *Mol.*
Ecol. **16**: 3145–3160.

Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics **24**: 1403–1405.

Jombart, T., Devillard, S. & Balloux, F. 2010. Discriminant analysis of principal components:
a new method for the analysis of genetically structured populations. *BMC Genet.* **11**: 94.

Judkins, M.E., Couger, B.M., Warren, W.C. & Van Den Bussche, R.A. 2019. A 50K SNP
array reveals genetic structure for bald eagles (*Haliaeetus leucocephalus*). *Conserv. Genet.* **21**:
65–76.

Kelly, M. 2019. Adaptation to climate change through genetic accommodation and assimilation
of plastic phenotypes. *Philos. Trans. R. Soc. B Biol. Sci.* **374**: 20180176.

Kerlinger, P. & Lein, M.R. 1988. Population Ecology of Snowy Owls during Winter on the
Great Plains of North America. *The Condor* **90**: 866–874.

Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A. & Mayrose, I. 2015.
CLUMPAK☐: a program for identifying clustering modes and packaging population structure
inferences across *K. Mol. Ecol. Resour.* **15**: 1179–1191.

Lanier, H.C., Gunderson, A.M., Weksler, M., Fedorov, V.B. & Olson, L.E. 2015.
Comparative Phylogeography Highlights the Double-Edged Sword of Climate Change Faced by
Arctic- and Alpine-Adapted Mammals. *PLOS ONE* **10**: e0118396.

Larsson, P., Götherström, A., Androsov, S., Germonpré, M., Bergfeldt, N., Fedorov, S.,
Eide, N.E., Sokolova, N., Berteaux, D., Angerbjörn, A., Flagstad, Ø., Plotnikov, V., Norén,
K., Díez-del-Molino, D., Dussex, N., Stanton, D.W.G. & Dalén, L. 2019. Consequences of
past climate change and recent human persecution on mitogenomic diversity in the arctic fox.
Philos. Trans. R. Soc. B Biol. Sci.

Legagneux, P., Gauthier, G., Berteaux, D., Bêty, J., Cadieux, M.-C., Bilodeau, F., Bolduc,
E., McKinnon, L., Tarroux, A., Therrien, J.-F., Morissette, L. & Krebs, C.J. 2012.
Disentangling trophic relationships in a High Arctic tundra ecosystem through food web
modeling. *Ecology* **93**: 1707–1716.

Lischer, H.E.L. & Excoffier, L. 2012. PGDSpider: an automated data conversion tool for
connecting population genetics and genomics programs. *Bioinformatics* **28**: 298–299.

Liu, X. & Fu, Y.-X. 2015. Exploring population size changes using SNP frequency spectra. *Nat.*
Genet. **47**: 555–559.

Liu, X. & Fu, Y.-X. 2020. Stairway Plot 2: demographic history inference with folded SNP
frequency spectra. *Genome Biol.* **21**: 280.

Lord, E., Dussex, N., Kierczak, M., Díez-del-Molino, D., Ryder, O.A., Stanton, D.W.,
Gilbert, M.T.P., Sánchez-Barreiro, F., Zhang, G. & Sinding, M.-H.S. 2020. Pre-extinction
demographic stability and genomic signatures of adaptation in the woolly rhinoceros. *Curr. Biol.*
30: 3871–3879.

Lorenzen, E.D., Nogués-Bravo, D., Orlando, L., Weinstock, J., Binladen, J., Marske, K.A.,
Ugan, A., Borregaard, M.K., Gilbert, M.T.P., Nielsen, R., Ho, S.Y.W., Goebel, T., Graf,
K.E., Byers, D., Stenderup, J.T., Rasmussen, M., Campos, P.F., Leonard, J.A., Koepfli, K.-
P., Froese, D., Zazula, G., Stafford, T.W., Aaris-Sørensen, K., Batra, P., Haywood, A.M.,
Singarayer, J.S., Valdes, P.J., Boeskorov, G., Burns, J.A., Davydov, S.P., Haile, J., Jenkins,
D.L., Kosintsev, P., Kuznetsova, T., Lai, X., Martin, L.D., McDonald, H.G., Mol, D.,

Meldgaard, M., Munch, K., Stephan, E., Sablin, M., Sommer, R.S., Sipko, T., Scott, E.,
Suchard, M.A., Tikhonov, A., Willerslev, R., Wayne, R.K., Cooper, A., Hofreiter, M., Sher,
A., Shapiro, B., Rahbek, C. & Willerslev, E. 2011. Species-specific responses of Late
Quaternary megafauna to climate and humans. *Nature* **479**: 359–364.
Louis, M., Skovrind, M., Samaniego Castruita, J.A., Garilao, C., Kaschner, K.,
Gopalakrishnan, S., Haile, J.S., Lydersen, C., Kovacs, K.M., Garde, E., Heide-Jørgensen,
M.P., Postma, L., Ferguson, S.H., Willerslev, E. & Lorenzen, E.D. 2020. Influence of past
climate change on phylogeography and demographic history of narwhals, *Monodon monoceros*.
Proc. R. Soc. B Biol. Sci. **287**: 20192964.
Mann, M.E., Zhang, Z., Rutherford, S., Bradley, R.S., Hughes, M.K., Shindell, D.,
Ammann, C., Faluvegi, G. & Ni, F. 2009. Global Signatures and Dynamical Origins of the
Little Ice Age and Medieval Climate Anomaly. *Science* **326**: 1256–1260.
Mantel, N. 1967. The detection of disease clustering and a generalized regression approach.
Cancer Res. **27**: 209–220.
Marthinsen, G., Wennerberg, L., Solheim, R. & Lifjeld, J.T. 2009. No phylogeographic
structure in the circumpolar snowy owl (*Bubo scandiacus*). *Conserv. Genet.* **10**: 923–933.
Mastretta-Yanes, A., Arrigo, N., Alvarez, N., Jorgensen, T.H., Piñero, D. & Emerson, B.C.
2015. Restriction site-associated DNA sequencing, genotyping error estimation and *de novo*
assembly optimization for population genetic inference. *Mol. Ecol. Resour.* **15**: 28–41.
McCabe, R.A., Therrien, J.-F., Wiebe, K.L., Gauthier, G., Brinker, D., Weidensaul, S. &
Elliott, K. 2021. Landscape cover type, not social dominance, is associated with the winter
movement patterns of Snowy Owls in temperate areas. *The Auk* **138**: ukaa082.
Meehan, T.D., LeBaron, G.S., Dale, K., Michel, N.L., Verutes, G.M. & Langham, G.M.
2018. *Abundance trends of birds wintering in the USA and Canada, from Audubon Christmas*
Bird Counts, 1966-2017, version 2.1. National Audubon Society, New York, New York, USA.
Meirmans, P.G. 2012. The trouble with isolation by distance. *Mol. Ecol.* **21**: 2839–2846.
Mendelsohn, B., Bedrosian, B., Love Stowell, S.M., Gagne, R.B., LaCava, M.E.F., Godwin,
B.L., Hull, J.M. & Ernest, H.B. 2020. Population genomic diversity and structure at the
discontinuous southern range of the Great Gray Owl in North America. *Conserv. Genet.*, doi:
10.1007/s10592-020-01280-8.
Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, C., Ekaykin, A., Hollowed, A.,
Kofinas, G., Mackintosh, A., Melbourne-Thomas, J. & Muelbert, M.M.C. 2020. Polar
Regions. In: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate* (H.-O.
Pörtner, D. C. Roberts, V. Masson-Delmott, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck,
A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, & N. M. Weyer, eds).
Miller, J.M., Cullingham, C.I. & Peery, R.M. 2020. The influence of a priori grouping on
inference of genetic clusters: simulation study and literature review of the DAPC method.
Heredity, doi: 10.1038/s41437-020-0348-2.
Miller, M.P., Davis, R.J., Forsman, E.D., Mullins, T.D. & Haig, S.M. 2018. Isolation by
distance versus landscape resistance: Understanding dominant patterns of genetic structure in
Northern Spotted Owls (*Strix occidentalis caurina*). *PLOS ONE* **13**: e0201720.
Miller, W., Schuster, S.C., Welch, A.J., Ratan, A., Bedoya-Reina, O.C., Zhao, F., Kim,
H.L., Burhans, R.C., Drautz, D.I., Wittekindt, N.E., Tomsho, L.P., Ibarra-Laclette, E.,
Herrera-Estrella, L., Peacock, E., Farley, S., Sage, G.K., Rode, K., Obbard, M., Montiel,
R., Bachmann, L., Ingolfsson, O., Aars, J., Mailund, T., Wiig, O., Talbot, S.L. & Lindqvist,
C. 2012. Polar and brown bear genomes reveal ancient admixture and demographic footprints of
past climate change. *Proc. Natl. Acad. Sci.* **109**: E2382–E2390.

Mueller, J.C., Kuhl, H., Boerno, S., Tella, J.L., Carrete, M. & Kempenaers, B. 2018.
Evolution of genomic variation in the burrowing owl in response to recent colonization of urban
areas. *Proc. R. Soc. B Biol. Sci.* **285**: 20180206.
Nam, K., Mugal, C., Nabholz, B., Schielzeth, H., Wolf, J.B., Backström, N., Künstner, A.,
Balakrishnan, C.N., Heger, A., Ponting, C.P., Clayton, D.F. & Ellegren, H. 2010. Molecular
evolution of genes in avian genomes. *Genome Biol.* **11**: 17.
Navascués, M., Leblois, R. & Burgarella, C. 2017. Demographic inference through
approximate-Bayesian-computation skyline plots. *PeerJ* **5**: e3530.
Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R.B., Simpson,
G.L., Solymos, P., Stevens, M.H.H. & Wagner, H. 2013. Vegan: Community ecology package.
R Package Version 2–0.
Palstra, F.P. & Ruzzante, D.E. 2008. Genetic estimates of contemporary effective population
size: what can they tell us about the importance of genetic stochasticity for wild population
persistence? *Mol. Ecol.* **17**: 3428–3447.
Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. *Annu.*
Rev. Ecol. Evol. Syst. **37**: 637–669.
Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H.S. & Hoekstra, H.E. 2012. Double Digest
RADseq: An Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and
Non-Model Species. *PLoS ONE* **7**: 11.
Potapov, E. & Sale, R. 2012. *The snowy owl*. T&A Poyser, London, United kingdon.
Prost, S., Guralnick, R.P., Waltari, E. & Fedorov, V.B. 2013. Losing ground: past history and
future fate of Arctic small mammals in a changing climate. *Glob. Change Biol.* 11.
Prost, S., Smirnov, N., Fedorov, V.B., Sommer, R.S., Stiller, M., Nagel, D., Knapp, M. &
Hofreiter, M. 2010. Influence of Climate Warming on Arctic Mammals? New Insights from
Ancient DNA Studies of the Collared Lemming *Dicrostonyx torquatus*. *PLoS ONE* **5**: e10447.
QGIS Development team. 2020. *QGIS Geographic Information System. Open Source*
Geospatial Foundation.
R Core Team. 2021. *R: A language and environment for statistical computing.* R Foundation for
Statistical Computing, Vienna, Austria.
Rich, T.D., Beardmore, H., Blancher, P.J., Bradstreet, M.S.W., Butcher, G.S., Demarest,
D.W., Dunn, E.H., Hunter, W.C., Inigo-Elias, E.E., Martell, A.M., Panjabi, A.O., Pashley,
D.N., Rosenberg, K.V., Rustay, C.M., Wendt, J.S. & Will, T.C. 2004. *Partners in Flight*
North American Landbird Conservation Plan. Cornell Lab of Ornithology, Ithaca, New York,
USA.
Robillard, A., Gauthier, G., Therrien, J.-F. & Bêty, J. 2018. Wintering space use and site
fidelity in a nomadic species, the snowy owl. *J. Avian Biol.* **49**: jav-01707.
Robillard, A., Therrien, J.F., Gauthier, G., Clark, K.M. & Bêty, J. 2016. Pulsed resources at
tundra breeding sites affect winter irruptions at temperate latitudes of a top predator, the snowy
owl. *Oecologia* **181**: 423–433.
Rosenberg, K.V., Kennedy, J.A., Dettmers, R., Ford, R.P., Reynolds, D., Alexander, J.D.,
Beardmore, C.J., Blancher, P.J., Bogart, R.E., Butcher, G.S., Camfield, A.F., Couturier, A.,
Demarest, D.W., Easton, W.E., Giocomo, J.J., Keller, R.H., Mini, A.E., Panjabi, A.O.,
Pashley, D.N., Rich, T.D., Ruth, J.M., Stabins, H., Stanton, J.C. & Will, T. 2016. *Partners in*
Flight Landbird Conservation Plan: 2016 Revision for Canada and Continental United States.
Rousset, F. 1997. Genetic differentiation and estimation of gene flow from F-statistics under
isolation by distance. *Genetics* **145**: 1219–1228.
Sandom, C., Faurby, S., Sandel, B. & Svenning, J.-C. 2014. Global late Quaternary

megafauna extinctions linked to humans, not climate change. *Proc. R. Soc. B Biol. Sci.* **281**:
20133254.
Schwartz, M., Luikart, G. & Waples, R. 2007. Genetic monitoring as a promising tool for
conservation and management. *Trends Ecol. Evol.* **22**: 25–33.
Smeds, L., Qvarnström, A. & Ellegren, H. 2016. Direct estimate of the rate of germline
mutation in a bird. *Genome Res.* **26**: 1211–1218.
Solheim, R. 2012. Wing feather moult and age determination of Snowy Owls *Bubo scandiacus*.
Ornis Nor. **35**: 48–67.
Stewart, M., Carleton, W.C. & Groucutt, H.S. 2021. Climate change, not human population
growth, correlates with Late Quaternary megafauna declines in North America. *Nat. Commun.*
12: 1–15.
Storz, J.F., Ramakrishnan, U. & Alberts, S.C. 2001. Determinants of effective population size
for loci with different modes of inheritance. *J. Hered.* **92**: 497–502.
Strimmer, K. & Pybus, O.G. 2001. Exploring the Demographic History of DNA Sequences
Using the Generalized Skyline Plot. *Mol. Biol. Evol.* **18**: 2298–2305.
Taylor, J.J., Lawler, J.P., Aronsson, M., Barry, T., Bjorkman, A.D., Christensen, T.,
Coulson, S.J., Cuyler, C., Ehrich, D. & Falk, K. 2020. Arctic terrestrial biodiversity status and
trends: a synopsis of science supporting the CBMP State of Arctic Terrestrial Biodiversity
Report. *Ambio* **49**: 833–847.
Therrien, J.-F., Gauthier, G. & Bêty, J. 2011. An avian terrestrial predator of the Arctic relies
on the marine ecosystem during winter. *J. Avian Biol.* **42**: 363–369.
Therrien, J.-F., Gauthier, G., Pinaud, D. & Bêty, J. 2014. Irruptive movements and breeding
dispersal of snowy owls: a specialized predator exploiting a pulsed resource. *J. Avian Biol.* **45**:
536–544.
Therrien, J.-F., Weidensaul, S., Brinker, D., Huy, S., Miller, T., Jacobs, E., Weber, D.,
McDonald, T., Lanzone, M., Smith, N. & Lecomte, N. 2017. Winter Use of a Highly Diverse
Suite of Habitats by Irruptive Snowy Owls. *Northeast. Nat.* **24**: B81–B89.
Van Strien, M.J., Keller, D. & Holderegger, R. 2012. A new analytical approach to landscape
genetic modelling: least-cost transect analysis and linear mixed models: New approach in
landscape genetics. *Mol. Ecol.* **21**: 4010–4023.
Wang, I.J., Santiago, E. & Caballero, A. 2016. Prediction and estimation of effective
population size. *Heredity* **117**: 193–206.
Waples, R.S. 2016. Making sense of genetic estimates of effective population size. *Mol. Ecol.*
25: 4689–4691.
Waples, R.S. & Do, C. 2010. Linkage disequilibrium estimates of contemporary N_e using
highly variable genetic markers: a largely untapped resource for applied conservation and
evolution. *Evol. Appl.* **3**: 244–262.
Weir, B.S. & Cockerham, C.C. 1984. Estimating F-statistics for the analysis of population
structure. *evolution* **38**: 1358–1370.
Weir, B.S. & Goudet, J. 2017. A Unified Characterization of Population Structure and
Relatedness. *Genetics* **206**: 2085–2103.
Wright, S. 1943. Isolation by distance. *Genetics* **28**: 114–138.
Yannic, G., Pellissier, L., Ortego, J., Lecomte, N., Couturier, S., Cuyler, C., Dussault, C.,
Hundertmark, K.J., Irvine, R.J., Jenkins, D.A., Kolpashikov, L., Mager, K., Musiani, M.,
Parker, K.L., Røed, K.H., Sipko, T., Þórisson, S.G., Weckworth, B.V., Guisan, A.,
Bernatchez, L. & Côté, S.D. 2014. Genetic diversity in caribou linked to past and future climate
change. *Nat. Clim. Change* **4**: 132–137.

Yannic, G., Statham, M.J., Denoyelle, L., Szor, G., Qulaut, G.Q., Sacks, B.N. & Lecomte,
N. 2017. Investigating the ancestry of putative hybrids: are Arctic fox and red fox hybridizing?
Polar Biol. **40**: 2055–2062.
Yannic, G., St-Laurent, M.-H., Ortego, J., Taillon, J., Beauchemin, A., Bernatchez, L.,
Dussault, C. & Côté, S.D. 2016. Integrating ecological and genetic structure to define
management units for caribou in Eastern Canada. *Conserv. Genet.* **17**: 437–453.
Zhang, G., Li, C., Li, Q., Li, B., Larkin, D.M., Lee, C., Storz, J.F., Antunes, A., Greenwold,
M.J., Meredith, R.W., Odeen, A., Cui, J., Zhou, Q., Xu, L., Pan, H., Wang, Z., Jin, L.,
Zhang, P., Hu, H., Yang, W., Hu, J., Xiao, J., Yang, Z., Liu, Y., Xie, Q., Yu, H., Lian, J.,
Wen, P., Zhang, F., Li, H., Zeng, Y., Xiong, Z., Liu, S., Zhou, L., Huang, Z., An, N., Wang,
J., Zheng, Q., Xiong, Y., Wang, G., Wang, B., Wang, J., Fan, Y., da Fonseca, R.R., Alfaro-
Nunez, A., Schubert, M., Orlando, L., Mourier, T., Howard, J.T., Ganapathy, G., Pfenning,
A., Whitney, O., Rivas, M.V., Hara, E., Smith, J., Farre, M., Narayan, J., Slavov, G.,
Romanov, M.N., Borges, R., Machado, J.P., Khan, I., Springer, M.S., Gatesy, J., Hoffmann,
F.G., Opazo, J.C., Hastad, O., Sawyer, R.H., Kim, H., Kim, K.-W., Kim, H.J., Cho, S., Li,
N., Huang, Y., Bruford, M.W., Zhan, X., Dixon, A., Bertelsen, M.F., Derryberry, E.,
Warren, W., Wilson, R.K., Li, S., Ray, D.A., Green, R.E., O'Brien, S.J., Griffin, D.,
Johnson, W.E., Haussler, D., Ryder, O.A., Willerslev, E., Graves, G.R., Alstrom, P.,
Fjeldsa, J., Mindell, D.P., Edwards, S.V., Braun, E.L., Rahbek, C., Burt, D.W., Houde, P.,
Zhang, Y., Yang, H., Wang, J., Avian Genome Consortium, Jarvis, E.D., Gilbert, M.T.P.,
Wang, J., Ye, C., Liang, S., Yan, Z., Zepeda, M.L., Campos, P.F., Velazquez, A.M.V.,
Samaniego, J.A., Avila-Arcos, M., Martin, M.D., Barnett, R., Ribeiro, A.M., Mello, C.V.,
Lovell, P.V., Almeida, D., Maldonado, E., Pereira, J., Sunagar, K., Philip, S., Dominguez-
Bello, M.G., Bunce, M., Lambert, D., Brumfield, R.T., Sheldon, F.H., Holmes, E.C.,
Gardner, P.P., Steeves, T.E., Stadler, P.F., Burge, S.W., Lyons, E., Smith, J., McCarthy, F.,
Pitel, F., Rhoads, D. & Froman, D.P. 2014. Comparative genomics reveals insights into avian
genome evolution and adaptation. *Science* **346**: 1311–1320.
Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C. & Weir, B.S. 2012. A high-
performance computing toolset for relatedness and principal component analysis of SNP data.
Bioinformatics **28**: 3326–3328.
2012. *Dispersal ecology and evolution* (J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock,
eds). Oxford University Press, Oxford, United Kingdom.
2019. *Handbook of the Birds of the word alive* (J. Del Hoyo, A. Elliott, J. Sargatal, D. A.
Christie, & G. Kirwan, eds). Lynx Edicions, Barcelona.
797

 TABLES

799   **Table 1.** Sampling distribution of Snowy Owls per sex or age for each wintering area.

		Wintering areas				
		West	Central	Great Lakes	East	Total
	Male	24	14	13	18	69
Sex	Female	25	14	9	24	72
	Unkown	1	1	5	0	7
	Adult	28	5	13	25	71
Age	First year	20	24	12	15	71
	Unknown	2	0	4	2	8
	Total	**50**	**29**	**29**	**42**	**150**

Figure 1. Sampling sites of wintering Snowy Owls (n=150) collected during November-March

between 2007 and 2018. Green areas represent the geographic year-round range and yellow areas

represent nonbreeding range of the species (BirdLife International 2020). Pairwise *Fst* values

among wintering areas (W-West, C-Central, GL- Great Lakes, E-East) and the associated *p*-

values are displayed in the legend section.

Figure 2. Violin plot of genetic distance [beta estimation of relatedness between every pair of

individuals] for all individual Snowy Owls pooled (Total) and for each wintering area (W-West,

C-Central, GL- Great Lakes, E-East).

Figure 3. Reconstruction of the effective population size of the snowy owl population (*Ne*) from

600 Ka until 100 years ago in North America, assuming mutation rate of 1.9×10^{-9} and a

generation time of 7.93 based on a Stairway Plot 2 (Liu and Fu 2020). The method precludes

reliable estimates for the most recent 100 years. Grey lines indicate the 95% confidence interval.

Light grey areas represent glaciation events according to Clark et al. (2009, 2012); Mann et al.

(2009). Events in red are characterized by higher temperature and events in blue are characterized

by lower temperatures.

SUPPLEMENTARY MATERIAL

Population genetic structure, genetic distance and Isolation-by-distance

Unless mentioned, all analyses were in *R* (3.6.2.; R Core Team 2019). We first computed a Principal Component Analysis (PCA) with the function *glPCA* using the package *Adegenet**2.1.1 (Jombart 2008) using the four wintering regions (W, C, GL and E), retaining 100 PCA axes according to the eigenvalues plot. To infer the number of clusters of genetically related individuals to best fit our data, we used a Discriminant Analysis of Principal Components (DAPC; Jombart et al. 2010) also implemented in the package *Adegenet* (Jombart 2008). We followed the recommendation from Miller et al. (2020) to report the methods and results. Without prior information (*de novo*, unsupervised DAPC), we computed the function *find.cluster()* to determine the optimal number of clusters. Precisely, we ran successive *K*-means clustering with an increase in the number of clusters (K=1-10). We used the smallest Bayesian information criterion (BIC) value associated with K for defining the number of clusters. We explored grouping individuals by sex or age (adult only), but we did not find any pattern. We also used ADMIXTURE v 1.3 (Alexander *et al.* 2009) to estimate the number of clusters in our dataset. We ran 10 replicates of ADMIXTURE (without any prior) allowing for the number of clusters (*K*) in the model to vary from 1 to 10. We generated random seeds for each run (beta < 0.0001). We chose the *K* value that minimized cross-validation error and hence best fit the data (Alexander *et al.* 2009). The output generated from 10 independent runs across each *K* was summarized and graphically represented using the main pipeline implemented in CLUMPAK with default options (Kopelman *et al.* 2015). We performed 10 independent runs of ADMIXTURE with adults and first years separated. Additionally, we computed pairwise F_{ST} among wintering areas (W vs. C vs. GL vs. E) with the *gl.fst.pop* in the *dartR* package (Gruber *et al.* 2018). We calculated pairwise F_{ST} by gender (male *vs.* female) and age group (adult *vs.* first year) among wintering areas. We computed global and within wintering regions F_{IS} using *gl.basics.stats* from *DartR* and tested if mean F_{IS} were significantly different from zero using Wilcoxon signed rank tests with continuity correction. We computed a *beta* estimator genetic distance matrix (i.e. a relatedness estimation matrix; according to Weir and Goudet 2017) between each pair of individuals with the function *snpgdsIndvBeta* in the package *SNPRelate* (Zheng *et al.* 2012). Generated values were homolog to the F_{ST} genetic distance (Weir and Cockerham 1984) measured between pairs of individuals instead of pairs of populations (Weir and Goudet 2017). A value of 0.5 is a first degree of relatedness (expected e.g., for full-sibs) and 0.25 a second degree (e.g. between half-sibs). We also performed a genetic distance matrix between all individuals located in the same wintering areas and a genetic distance matrix per age group (adult *vs.* first year). We calculated Geographic Euclidean distances between each pair of individuals using distance matrix in QGIS v 3.8. (QGIS Development team 2020). We assessed Isolation-by-distance (IBD; Wright 1943), the tendency for genetic similarity to reflect geographic proximity (Meirmans 2012), with two analyses. First, we computed a Mantel test (Mantel 1967) between the *beta* (i.e. genetic distance matrix) and the geographic matrices using the mantel function in the *Vegan* package (Oksanen *et al.* 2013) with 10,000 permutations to assess significance. Second, we also used a maximum-likelihood population-effects model (MLPE) in a linear mixed model (LMM), in which the covariate structure is specifically designed to account for the specific non-independency among column and row values in an n x n distance matrices (Van Strien *et al.* 2012). We fitted LMMs with *lme4* package (Bates *et al.* 2015). We verified linearity assumptions. We also ran the same analysis (Mantel and LMM) at the population scale, i.e. testing the relationship between the logarithm of Euclidian geographic [km] and the pairwise genetic distance $F_{ST}/(1-F_{ST})$ among all pairs of wintering regions following (Rousset 1997).

TABLES

Table S1: Sample information. **ID:** unique sample identification; **Lab ID:** identification used during laboratory work (UBC Beaty Biodiversity; UM Université de Moncton; USMB Université Savoie Mont-Blanc); **Sample type**: liver, tissue, feather or blood; **Wintering area:** West (British Columbia); Central (Saskatchewan); GL, Great lakes areas (Michigan, Wisconsin, Ontario, and Maryland); and East, i.e. all other individuals from the east coast of North America including the single Nunavut sample; **Year:** year of collection; **Sex:** sex of the individual; Age: Adult or first year (First);**Stored location:** actual location and long-term storage of the sample.

ID	Lab ID	Sample type	Wintering area	Year	Sex	Age	Storage location	Organisation	Country	Accession No.
L12771	Bs_256	liver	WEST	2012	M	Adult	BC: Delta	UBC	Canada	B016769
L12772	Bs_257a	liver	WEST	2012	F	Adult	BC: Port Coquitlam	UBC	Canada	B016770
L12773	Bs_258	liver	WEST	2012	F	Adult	BC: Hope	UBC	Canada	B016771
L12774	Bs_259a	liver	WEST	2012	M	Adult	BC: Powell River	UBC	Canada	B016772
L12954	Bs_260	liver	WEST	2012	F	Adult	BC: Squamish	UBC	Canada	B016773
L13573	Bs_261	liver	WEST	2012	M	Adult	BC: Port Hardy	UBC	Canada	B016776
L13574	Bs_262	liver	WEST	2012	M	Adult	BC: Parksville	UBC	Canada	B016777
L13576	Bs_263	liver	WEST	2012	M	Adult	BC: Courtenay	UBC	Canada	B016779
L13577	Bs_264	liver	WEST	2012	M	Adult	BC: Qualicum Beach	UBC	Canada	B016780
L13578	Bs_265	feather	WEST	2012	F	Adult	BC: Richmond	UBC	Canada	B016781
L13585	Bs_266	liver	WEST	2013	F	Adult	BC: Smithers	UBC	Canada	B016871
L13586	Bs_267	liver	WEST	2013	M	Adult	BC: Richmond	UBC	Canada	B016872
L13587	Bs_268a	liver	WEST	2013	F	First	BC: Richmond	UBC	Canada	B016873
L13588	Bs_269b	liver	WEST	2012	F	Adult	BC: Prince George	UBC	Canada	B016874
L13589	Bs_270	liver	WEST	2012	F	Adult	BC: Fort St James	UBC	Canada	B016875
L13590	Bs_271	liver	WEST	2012	M	First	BC: Vanderhoof	UBC	Canada	B016876
L13591	Bs_272	liver	WEST	2012	M	Adult	BC: Vanderhoof	UBC	Canada	B016877
L13592	Bs_273	liver	WEST	2012	F	First	BC: Prince George	UBC	Canada	B016878

L13594	Bs_275	liver	WEST	2012	F	First	BC: Prince George	UBC	Canada	B016880
L13595	Bs_276	liver	WEST	2012	F	Adult	BC: Prince George	UBC	Canada	B016881
L13596	Bs_277	liver	WEST	2012	F	Adult	BC: Fort St James	UBC	Canada	B016882
L13597	Bs_278	liver	WEST	2013	M	First	BC: Kelowna	UBC	Canada	B016883
L13598	Bs_279b	liver	WEST	2012	M	Adult	BC: Cranbrook	UBC	Canada	B016884
L13599	Bs_280	liver	WEST	NA	M	First	BC: Dawson Creek	UBC	Canada	B016885
L13601	Bs_282	liver	WEST	NA	F	First	BC: Taylor	UBC	Canada	B016887
L13602	Bs_283	liver	WEST	2013	F	First	BC: Dawson Creek	UBC	Canada	B016888
L13603	Bs_284b	liver	WEST	2008	F	First	BC: Cecil Lake	UBC	Canada	B016889
L13604	Bs_285	liver	WEST	NA	M	First	BC: Fort St John	UBC	Canada	B016890
L13605	Bs_286	liver	WEST	2008	Unk	First	BC: Fort St John	UBC	Canada	B016891
L13606	Bs_287	liver	WEST	2012	M	Adult	BC: Clinton	UBC	Canada	B016892
L13607	Bs_288	liver	WEST	2012	F	First	BC: Kamloops	UBC	Canada	B016893
L13608	Bs_289	liver	WEST	2013	M	First	BC: Ladner	UBC	Canada	B016894
L13609	Bs_290	feather	WEST	2013	M	First	BC: Ladner	UBC	Canada	B016895
L13610	Bs_291	liver	WEST	2012	F	First	BC: Richmond	UBC	Canada	B017061
L13611	Bs_292	liver	WEST	2012	F	Adult	BC: Richmond	UBC	Canada	B017062
L13612	Bs_293	liver	WEST	2012	M	Unk	BC: Richmond	UBC	Canada	B017063
L13614	Bs_294	liver	WEST	2013	F	First	BC: Ladner	UBC	Canada	B017064
L13616	Bs_295	liver	WEST	2013	F	First	BC: Penticton	UBC	Canada	B017069
L13617	Bs_296	liver	WEST	2013	F	First	BC: Delta	UBC	Canada	B017066
L13625	Bs_297a	liver	WEST	2012	M	Adult	BC: Penticton	UBC	Canada	B016782
L13626	Bs_298b	liver	WEST	2012	F	First	BC: Penticton	UBC	Canada	B016783
L13776	Bs_299	liver	WEST	2012	M	Adult	BC: Delta	UBC	Canada	B016784
L13777	Bs_300	liver	WEST	2012	F	Adult	BC: Williams_Lake	UBC	Canada	B016785
L13778	Bs_301	liver	WEST	2012	F	Adult	BC: Pemberton	UBC	Canada	B016786
L13779	Bs_302	liver	WEST	2012	M	Adult	BC: Williams_Lake	UBC	Canada	B016787

L13780	Bs_303	liver	WEST	2012	M	Adult	BC: Williams_Lake	UBC	Canada	B016788
L13781	Bs_304	liver	WEST	2012	M	Adult	BC: Richmond	UBC	Canada	B016789
L13782	Bs_305	liver	WEST	2013	M	First	BC: Abbotsford	UBC	Canada	B016896
L13783	Bs_306b	liver	WEST	2013	M	Adult	BC: Chilliwack	UBC	Canada	B016897
L13955	Bs_307	liver	WEST	2012	M	Adult	BC: Coquitlam	UBC	Canada	B016791
65900455	Bs_047	feather	EAST	2015	M	Adult	NB: Moncton	UM	Canada	-
67903623	Bs_049	feather	EAST	2018	F	First	NB: Moncton	UM	Canada	-
92808790	Bs_055b	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
69900560	Bs_056	feather	EAST	2014	M	Adult	NB: Moncton	UM	Canada	-
78851220	Bs_057	feather	EAST	2018	M	Adult	NB: Moncton	UM	Canada	-
109801605	Bs_061	feather	EAST	2018	M	Adult	NB: Moncton	UM	Canada	-
108805542	Bs_062	feather	EAST	2014	M	Adult	NB: Moncton	UM	Canada	-
65900454	Bs_064	feather	EAST	2015	M	Adult	NB: Moncton	UM	Canada	-
78836716	Bs_070	feather	EAST	2017	M	Adult	NB: Moncton	UM	Canada	-
92808795	Bs_075	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
92808796	Bs_076	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
109801896	Bs_083	feather	EAST	2018	M	First	NB: Moncton	UM	Canada	-
111854551	Bs_086	feather	EAST	2017	M	Adult	NB: Moncton	UM	Canada	-
92808793	Bs_095	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
69900559	Bs_100a	feather	EAST	2014	F	Adult	NB: Moncton	UM	Canada	-
65900451	Bs_113	feather	EAST	2015	M	Adult	NB: Moncton	UM	Canada	-
67903626	Bs_114	feather	EAST	2018	F	First	NB: Moncton	UM	Canada	-
69900561	Bs_115	feather	EAST	2014	F	Adult	NB: Moncton	UM	Canada	-
78836713	Bs_118	feather	EAST	2015	F	Adult	NB: Moncton	UM	Canada	-
65900458	Bs_125	feather	EAST	2017	F	Adult	NB: Moncton	UM	Canada	-
69900557	Bs_126	feather	EAST	2014	F	First	NB: Moncton	UM	Canada	-
69900563	Bs_127	feather	EAST	2014	F	Adult	NB: Moncton	UM	Canada	-
92808800	Bs_128	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
92808798	Bs_129	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-

69900562	Bs_130	feather	EAST	2014	F	Adult	NB: Moncton	UM	Canada	-
Bubo5	Bs_136	feather	EAST	2007	F	Adult	NB: Moncton	UM	Canada	-
A1570718	Bs_188	feather	EAST	2018	M	Adult	NB: Moncton	UM	Canada	-
A952818	Bs_189	feather	EAST	2018	F	Adult	NB: Moncton	UM	Canada	-
60868075	Bs_219	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
60868076	Bs_220	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
60868077	Bs_221	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
60868097	Bs_223a	feather	EAST	2014	M	Adult	NB: Moncton	UM	Canada	-
60868093	Bs_227	feather	EAST	2014	F	Adult	NB: Moncton	UM	Canada	-
60880995	Bs_228	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
59946963	Bs_238a	feather	EAST	2014	F	Adult	NB: Moncton	UM	Canada	-
59929600	Bs_242	feather	EAST EAST	2014	F	First	NB: Moncton	UM	Canada	-
65903413	Bs_243a	feather	EAST	2015	F	Adult	NB: Moncton	UM	Canada	-
59929599	Bs_245	feather	EAST	2014	F	First	NB: Moncton	UM	Canada	-
59950750	Bs_246	feather	EAST	2014	F	First	NB: Moncton	UM	Canada	-
59950748	Bs_247	feather	EAST	2014	M	First	NB: Moncton	UM	Canada	-
60868085	Bs_251	feather	EAST	2014	M	Adult	NB: Moncton	UM	Canada	-
62923832	Bs_254	feather	EAST	2014	F	Adult	NB: Moncton	UM	Canada	-
WDL16056	Bs_033	feather	GL	2016	F	Adult	NB: Moncton	UM	Canada	-
WDL16083	Bs_034	feather	GL	2016	F	Adult	NB: Moncton	UM	Canada	-
WDL16118	Bs_037	feather	GL	2016	M	First	NB: Moncton	UM	Canada	-
WDL16119	Bs_038a	feather	GL	2016	M	Adult	NB: Moncton	UM	Canada	-
WDL16172	Bs_040	feather	GL	2016	M	Adult	NB: Moncton	UM	Canada	-
WDL16547	Bs_042	feather	GL	2016	F	Adult	NB: Moncton	UM	Canada	-
70902651	Bs_058	dry_blood	GL	2014	F	Adult	NB: Moncton	UM	Canada	-
70902652	Bs_081	dry_blood	GL	2014	F	First	NB: Moncton	UM	Canada	-
65900450	Bs_101	feather	GL	2015	F	Adult	NB: Moncton	UM	Canada	-
59932503	Bs_116b	feather	GL	2014	F	First	NB: Moncton	UM	Canada	-
59932502	Bs_117	feather	GL	2013	F	First	NB: Moncton	UM	Canada	-

78820173	Bs_132	feather	GL	2014	M	Adult	NB: Moncton	UM	Canada	-
WDL15976	Bs_159	feather	GL	2015	M	First	NB: Moncton	UM	Canada	-
WDL151203	Bs_161	feather	GL	2015	F	Adult	NB: Moncton	UM	Canada	-
WDL151059	Bs_171	feather	GL	2015	M	First	NB: Moncton	UM	Canada	-
WDL151045	Bs_172b	feather	GL	2015	M	First	NB: Moncton	UM	Canada	-
WDL15180	Bs_177	feather	GL	2015	F	First	NB: Moncton	UM	Canada	-
WDL15027	Bs_178a	feather	GL	2015	F	Adult	NB: Moncton	UM	Canada	-
WDL15015	Bs_183	feather	GL	2015	F	Adult	NB: Moncton	UM	Canada	-
WDL15213	Bs_184	feather	GL	2015	M	Adult	NB: Moncton	UM	Canada	-
WDL15072	Bs_185a	feather	GL	2015	F	Adult	NB: Moncton	UM	Canada	-
26903001	Bs_207	feather	GL	2016	Unk	Unk	NB: Moncton	UM	Canada	-
26919001	Bs_209	feather	GL	2016	Unk	Unk	NB: Moncton	UM	Canada	-
26919001	Bs_211a	feather	GL	2016	Unk	Unk	NB: Moncton	UM	Canada	-
27219001	Bs_212	feather	GL	2016	Unk	Unk	NB: Moncton	UM	Canada	-
26864001	Bs_215	feather	GL	2016	Unk	Unk	NB: Moncton	UM	Canada	-
26863001	Bs_216	feather	GL	2016	Unk	Unk	NB: Moncton	UM	Canada	-
26879001	Bs_217	feather	GL	2016	Unk	Unk	NB: Moncton	UM	Canada	-
WDL151095	Bs_170	feather	GL	2015	M	First	NB: Moncton	UM	Canada	-
200421	Bs_001	tissu	CENTRAL	2016	M	Adult	Le-Bourget-du-Lac	USMB	France	-
207465	Bs_002	tissu	CENTRAL	2017	F	First	Le-Bourget-du-Lac	USMB	France	-
500000	Bs_003	tissu	CENTRAL	2018	M	First	Le-Bourget-du-Lac	USMB	France	-
202636	Bs_004	tissu	CENTRAL	2017	F	Adult	Le-Bourget-du-Lac	USMB	France	-
185184	Bs_005	tissu	CENTRAL	2014	M	Adult	Le-Bourget-du-Lac	USMB	France	-
207951	Bs_006a	tissu	CENTRAL	2017	F	First	Le-Bourget-du-Lac	USMB	France	-
208485	Bs_007a	tissu	CENTRAL	2018	M	Adult	Le-Bourget-du-Lac	USMB	France	-
199380	Bs_008	tissu	CENTRAL	2016	M	First	Le-Bourget-du-Lac	USMB	France	-
207829	Bs_009	tissu	CENTRAL	2017	F	First	Le-Bourget-du-	USMB	France	-

							Lac			
207131	Bs_010	tissu	CENTRAL	2017	M	First	Le-Bourget-du-Lac	USMB	France	-
500001	Bs_011a	tissu	CENTRAL	2018	M	First	Le-Bourget-du-Lac	USMB	France	-
216308	Bs_012	tissu	CENTRAL	2018	Unk	First	Le-Bourget-du-Lac	USMB	France	-
62951004	Bs_013	tissu	CENTRAL	2015	F	Adult	Le-Bourget-du-Lac	USMB	France	-
500003	Bs_014	tissu	CENTRAL	2015	F	First	Le-Bourget-du-Lac	USMB	France	-
500002	Bs_015	tissu	CENTRAL	2017	M	First	Le-Bourget-du-Lac	USMB	France	-
208151	Bs_016	tissu	CENTRAL	2015	M	First	Le-Bourget-du-Lac	USMB	France	-
500006	Bs_017a	tissu	CENTRAL	2015	M	First	Le-Bourget-du-Lac	USMB	France	-
500005	Bs_018	tissu	CENTRAL	2015	M	First	Le-Bourget-du-Lac	USMB	France	-
187393	Bs_019	tissu	CENTRAL	2015	F	First	Le-Bourget-du-Lac	USMB	France	-
191937	Bs_020b	tissu	CENTRAL	2015	F	First	Le-Bourget-du-Lac	USMB	France	-
500004	Bs_021	tissu	CENTRAL	2015	F	First	Le-Bourget-du-Lac	USMB	France	-
192027	Bs_022b	tissu	CENTRAL	2015	F	First	Le-Bourget-du-Lac	USMB	France	-
195064	Bs_023	tissu	CENTRAL	2016	F	First	Le-Bourget-du-Lac	USMB	France	-
18	Bs_024	tissu	CENTRAL	2015	M	First	Le-Bourget-du-Lac	USMB	France	-
191707	Bs_025b	tissu	CENTRAL	2016	F	First	Le-Bourget-du-Lac	USMB	France	-
194340	Bs_026	tissu	CENTRAL	2016	F	First	Le-Bourget-du-Lac	USMB	France	-
153613	Bs_027a	feather	CENTRAL	2008	F	First	NB: Moncton	UM	Canada	-
PDS1535117	Bs_150	feather	CENTRAL	2016	M	First	NB: Moncton	UM	Canada	-
PDS1535118	Bs_151	feather	CENTRAL	2016	M	First	NB: Moncton	UM	Canada	-

Table S2: Pairwise *Fst* value (below diagonal) and p-value (above diagonal) between wintering areas (West (W), Central (C), Great lakes (GL), East (E) for A) adult individuals only B) first-year individuals only C) female only D) male only. Significant values are represented by a star and sample size per area is in bold. Negative values are equivalent to $F_{ST} = 0$.

A)

	W (**28**)	C(**5**)	GL (**12**)	E (**14**)
W	-	0.070	0.059	0.017*
C	0.002738	-	0.659	0.720
GL	0.001420	-0.000773	-	0.590
E	**0.001756**	-0.001077	-0.000211	-

B)

	W(**20**)	C (**24**)	GL (**12**)	E (**15**)
W	-	0.229	0.110	0.653
C	0.000414	-	0.171	0.892
GL	0.001152	0.000853	-	0.398
E	0.000272	0.000962	0.000252	-

C)

	W(**28**)	C(**5**)	GL (**13**)	E (**24**)
W	-	0.055	0.042*	0.006*
C	0.00274	-	0.546	0.356
GL	**0.00144**	-0.000226	-	0.200
E	**0.00154**	0.000542	0.000707	-

D)

	W (**24**)	C (**14**)	GL (**13**)	E (**18**)
W	-	0.610	0.903	0.230
C	-0.000185	-	0.848	0.076
GL	-0.000964	-0.00103	-	0.814
E	0.000493	0.00145	0.000826	-

FIGURES

Figure S1. Effect of the variation of different *Stacks* core parameters locus error rated (proportion of missing loci in only one of the two replicates of a pair). Each box represents the result of a single *Stacks* run in which all parameters were set to their default values (m = 3, M = 2, N = M+2, n = 1, max_locus_stacks (mx.lcs) = 3, model = SNP) except for one that varied (indicated on the x-axis). See Mastretta-Yanes et al. (2015) for further discussion on parameters.

918   **Figure S2**. Variation of different *Stacks* core parameters on the SNP error rate (proportion of SNP
919   mismatch within a replicate pair). Each box represents the result of a single *Stacks* run in which all
920   parameters were set to their default values (m = 3, M = 2, N = M+2, n = 1, max_locus_stacks (mx.lcs) = 3,
921   model = SNP) except for one that varied (indicated on the x-axis). See Mastretta-Yanes et al. (2015) for
922   further discussion on parameters.

923
924
925   **Figure S3**. PCA of all the individuals according to their wintering areas. The number of retained PCs was
926   100. The number of dimensions (d) of the PCA is 2. Color codes: Orange=West (W, British Columbia,
927   n=50), Purple=Central (SK, Saskatchewan, n=29), Red= Great lake (GL, n=29) and Green=East (E,
928   n=42). The boxes with the letter are the centroid of each group.
929

930
931   **Figure S4**. Relationship between the Bayesian information criterion (BIC) value and the number of
932   clusters in the DAPC model.

933
934   **Figure S5.** Cross-validation results of 10 runs (K=1-10) from the ADMIXTURE software. Selected value of
935   the number of clusters (K) should be the lowest value (Alexander et al., 2009)

936
Figure S6. CLUMPAK results for *K*=2 from 10 runs of ADMIXTURE, without any prior allowing for the
number of clusters (*K*) in the model to vary from 1 to 10. We generated random seeds for each run (beta <
0.0001).

Figure S7. Boxplot of genetic distance (relatedness, beta index) of adults (dark grey) and first year (light
grey) for each wintering area.

Figure S8. Stairway plot results of runs considering different mutation rates (μ: 2×10^{-8} and 1.9×10^{-9}) or
generation time (gt: 4.7, 6, 7.93, 10, and 12.5 years). For the sake of clarity, we did not provide the 95%
confidence intervals in the figure (all very similar to the one presented in Fig. 3). The orange slope is the
one that was selected in Figure 3.