Hybridization drives mitochondrial DNA degeneration and metabolic shift in a species

with biparental mitochondrial inheritance

Mathieu Hénault[1245*], Souhir Marsit[12345#a], Guillaume Charron[1345#b] and Christian R. Landry[12345]

[1]Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, QC, Canada

[2]Département de biochimie, microbiologie et bioinformatique, Université Laval, Québec, QC,

Canada

[3]Département de biologie, Université Laval, Québec, QC, Canada

[4]Quebec Network for Research on Protein Function, Engineering, and Applications (PROTEO),

Université Laval, Québec, QC, Canada

[5]Université Laval Big Data Research Center (BDRC_UL), Québec, QC, Canada

[#a]Current address: Département de biologie, chimie et géographie, Université du Québec à

Rimouski, Rimouski, QC, Canada

[#b]Current address: Centre de foresterie des Laurentides, Ressources naturelles Canada,

Québec, QC, Canada

*Corresponding author

E-mail: mathieu.henault.1@ulaval.ca

ABSTRACT

Mitochondrial DNA (mtDNA) is a cytoplasmic genome that is essential for respiratory metabolism. While uniparental mtDNA inheritance is most common in animals and plants, distinct mtDNA haplotypes can coexist in a state of heteroplasmy, either because of paternal leakage or de novo mutations. MtDNA integrity and the resolution of heteroplasmy have important implications, notably for mitochondrial genetic disorders, genome evolution and speciation. However, knowledge on the factors driving heteroplasmy resolution is sparse. Here, we use *Saccharomyces* yeasts, fungi with constitutive biparental mtDNA inheritance, to investigate the resolution of mtDNA heteroplasmy in a variety of hybrid genotypes. We previously designed 11 crosses along a gradient of parental evolutionary divergence using undomesticated isolates of *Saccharomyces paradoxus* and *Saccharomyces cerevisiae*. Each cross was independently replicated 48 to 96 times, and the resulting 864 hybrids were evolved under relaxed selection for mitochondrial function. Genome sequencing of 446 MA lines revealed extensive mtDNA recombination, but recombination rates were not predicted by parental divergence. We found a strong positive relationship between parental divergence and the rate of large-scale mtDNA deletions, which lead to the loss of respiratory metabolism. We also uncovered associations between mtDNA recombination, deletion, and genome instability that were genotype-specific. Our results show that hybridization in yeast induces mtDNA degeneration, with deep consequences for mtDNA evolution, metabolism and the emergence of reproductive isolation.

INTRODUCTION

Mitochondrial DNAs (mtDNAs) are genomes that are maintained and expressed in mitochondria, which are ATP-producing organelles shared by virtually all Eukaryotes (Müller et al. 2012). Although mtDNAs vary greatly in terms of size and contents across Eukaryotes, all encode genes involved in respiration and protein synthesis (Adams and Palmer 2003). Most animals and plants have uniparental maternal mtDNA inheritance. Many exceptions exist to this rule, either because of constitutive uniparental or biparental inheritance (Breton et al. 2015; Wilson and Xu 2012; Wang et al. 2015), or leakage of paternal mtDNA (Parakatselaki and Ladoukakis 2021). Leakage and biparental mtDNA inheritance lead to a state of heteroplasmy, which is the intracellular coexistence of multiple mtDNA haplotypes. Evidence for paternal leakage is growing, even in species long considered to have strict uniparental maternal inheritance (Ladoukakis and Zouros 2017; Kondo et al. 1990; Payne et al. 2013; Gyllensten et al. 1991). Heteroplasmy may have important implications for mtDNA evolution, notably by enabling recombination between otherwise clonal mtDNAs (Rokas et al. 2003; Tsaousis et al. 2005; Ladoukakis et al. 2011). In humans, heteroplasmy is determinant in the onset of mitochondrial genetic disease, which are linked to inherited or de novo mtDNA variants (Stewart and Chinnery 2015; Wallace 2015; Stewart and Chinnery 2021). Additionally, when diverged populations or species hybridize, heteroplasmy can lead to the coexistence of more or less diverged mtDNAs which can then interact.

Genes encoded on mtDNAs are necessary, but not sufficient for carrying their essential metabolic function. The essential interactions between gene products encoded by mtDNAs and nuclear genomes require tight coevolution (Burton and Barreto 2012; Piccinini et al. 2021). Mitonuclear coevolution has important implications in the context of heteroplasmy in hybrids. Because mtDNA segregation is non-mendelian, heteroplasmy is generally transient and a single mtDNA haplotype is eventually fixed by vegetative segregation (Birky 2001). The return to homoplasmy can fix either parental haplotype, or a recombinant mtDNA containing alleles from both parents (Rokas et al. 2003; Tsaousis et al. 2005). Heteroplasmy resolution can lead to the fixation of alleles involved in mitonuclear incompatibilities, thus compromising mitochondrial function and hybrid fitness. In addition, mitochondrial alleles can interact epistatically within recombinant mtDNAs (mito-mito epistasis) (Wolters et al. 2018).

The genotype of a hybrid may contribute to determine the outcome of heteroplasmy resolution. One general prediction is that the potential for epistatic interactions scales with the level of

evolutionary divergence between the parents (Orr 1995). As such, mitonuclear incompatibilities

are expected to be more frequent between distantly related genomes compared to closely

related ones (Burton 2022). Under the same rationale, mito-mito incompatibilities are expected

to show increased prevalence between more diverged mtDNAs. Nucleotide sequence

divergence between mtDNAs is also expected to reduce the opportunities for homologous

recombination. Little is known about the path to heteroplasmy resolution in hybrids and its

consequences for hybrid evolution. Although the study of genomes sampled from natural

populations yielded important insights into mtDNA introgression, recombination and horizontal

transfer (Wu et al. 2015; Rice et al. 2013; Peris et al. 2017; Leducq et al. 2017), how parental

divergence shapes the neutral landscape of mtDNA recombination is largely unknown.

Additionally, whether parental divergence impacts mitochondrial respiratory function in hybrids

remains unexplored. Answering these questions requires a model system in which

heteroplasmy is dominant and that can easily generate a large diversity of F1 hybrid genotypes.

Like many other fungi, yeasts of the *Saccharomyces* genus have biparental mtDNA inheritance

(Solieri 2010; Wilson and Xu 2012). This enables the inheritance of either parental mtDNA in F1

hybrids, as well as recombination between parental haplotypes (Dujon et al. 1974). Because

yeast F1 hybrids are initially heteroplasmic and quickly transition to homoplasmy by mitotic

vegetative segregation (Solieri 2010), they are a powerful system to investigate the factors

driving heteroplasmy resolution. In addition, respiratory metabolism in yeast is facultative,

enabling the study of neutral mtDNA evolution without the selective pressure to maintain

respiration. The undomesticated species *Saccharomyces paradoxus* is found in natural lineages

with wide variation in genetic diversity and evolutionary divergence (Kuehne et al. 2007; Xia et

al. 2017; Leducq et al. 2016, 2014), allowing laboratory crosses that span various levels of

parental divergence.

Mutation accumulation (MA) is a type of experimental evolution that minimizes the power of

natural selection. Combined with genome sequencing, MA experiments produce near-unbiased

estimates of the spectrum of genomic changes (Lynch et al. 2016). We previously performed a

large-scale MA experiment on yeast hybrids (Charron et al. 2019; Hénault et al. 2020). Parental

strains comprised undomesticated isolates of *S. paradoxus* (11 strains from three lineages) and

Saccharomyces cerevisiae (two strains, Fig. 1A). Using these parental backgrounds, we

designed 11 crosses spanning a range of evolutionary divergence from intra-lineage to

interspecific. Each cross was replicated 48 to 96 times with independent matings, totalling 864

hybrid MA lines. We submitted each line to periodical extreme bottlenecks by streaking for

single colonies on solid medium (Joseph and Hall 2004; Lynch et al. 2008) (Fig. 1B) for ~770
mitotic generations (Charron et al. 2019; Hénault et al. 2020). Here, we use this collection and
the associated whole-genome sequencing datasets (Hénault et al. 2020; Marsit et al. 2021) to
investigate the resolution of heteroplasmy in a variety of hybrid genotypes.

RESULTS

Parental strains of the MA experiment display extensive variation in mtDNA content

We sequenced the genome of the 13 MA parental strains (Supplemental Table S1) using
Oxford Nanopore long reads and produced high-quality mtDNA assemblies (Supplemental Fig.
S1). Our long-read assemblies were consistent with a subset of five mtDNAs previously
assembled from short reads (Leducq et al. 2017) (Supplemental Fig. S2). Assembly annotation
revealed variation in mtDNA content, mostly in the presence of introns in the *COB* and *COX1*
genes (Fig. 1C). MtDNA content was consistent within lineages, although important diversity
was found in *SpB*. To enable comparative analyses based on a common reference sequence,
the mtDNA with the largest feature set (LL2012_028) was used as a template to build an
exhaustive artificial reference sequence by complementing it with introns found in other
assemblies, but absent from LL2012_028.

Figure 1. Design of the MA experiment on *Saccharomyces* hybrids and mtDNA contents of the
parental strains. (*A*) Design of the MA crosses. 864 independent MA lines were subdivided into
11 crosses spanning intra-lineage (*SpC*×*SpC*, CC; *SpB*×*SpB*, BB), intraspecific (*SpB*×*SpC*, BC;
SpB×*SpA*, BA) and interspecific (*SpB*×*S. cerevisiae*, BSc) parental divergence. Each cross was
replicated to initiate 48 to 96 independent evolution lines, all arising from distinct mating events.
The phylogenetic tree summarizes the evolutionary divergence between lineages and species
(substitutions per site, based on nuclear genome-wide variants). (*B*) Microbial MA experiments
minimize the efficiency of natural selection. Periodic extreme bottlenecks are achieved through
streaking for single colonies on solid medium, which amplifies genetic drift. (*C*) Annotation
summary of high-quality reference mtDNA assemblies for each parental strain. Squares denote
the presence of each feature in the corresponding genome. Light blue, protein-coding genes;
dark blue, protein-coding exons; white, introns; grey, RNA-coding genes; black, RNA-coding
exons; red, intronic or free-standing ORFs; green, tRNAs.

The landscape of mtDNA recombination is not predicted by parental divergence

For each cross, 36 to 47 hybrid MA lines were previously selected at random for short-read

whole-genome sequencing, both at the initial timepoint (~60 generations post-mating) and final

timepoint (after ~770 mitotic generations) of the MA experiment (Hénault et al. 2020; Marsit et

al. 2021) (Supplemental Table S2). We used this short-read data along with our mtDNA

assemblies to identify a set of confident marker variants that discriminate the parental mtDNA

haplotypes. We then used genotypes at marker positions to identify the recombination tracts for

individual MA lines, yielding 482 independent mtDNA haplotypes. All samples were

homoplasmic at both timepoints, with the vast majority of minor allele frequencies below 0.5%

(Supplemental Fig. S3). However, for many lines, the mtDNA haplotypes sampled at the initial

and final timepoints had incompatible genotypes, suggesting persistent segregation of distinct

haplotypes in some MA lines. The frequency at which distinct haplotypes were sampled varied

among crosses, reaching up to 33% of the lines in BA2 (Supplemental Table S3). All MA

crosses experienced some extent of recombination, except BA1 which exhibited no recombinant

mtDNA (Fig. 2A, Supplemental Fig. S4-14). Some crosses exhibited recombination hotspots,

yielding high local similarity among haplotypes (Fig. 2B).

149
Figure 2. Recombination is pervasive between parental mtDNA haplotypes of hybrid MA lines.
(*A*) Stacked concentric lines show recombination tracts of independent mtDNA haplotypes,
colored by parental ancestry. Outer circles show annotation summary as in Fig 1C, with
coordinates in kb. (*B*) Density of markers (red) and recombination junctions (black) in 250 bp
windows. Recombination junction density is normalized by marker density.

With the exception of BA1, MA crosses exhibited frequencies of recombinant mtDNAs ranging
from 28.9% to 57.5% (Fig. 3A). CC2 and BC1 crosses harbored the mtDNA haplotypes with the
highest count of recombination tracts. Recombination rate displayed no significant trend with
evolutionary divergence between the parents of a cross (based on nuclear genome-wide
variants; Fig. 3B, Supplemental Fig. S15A). Notably, the interspecific BSc crosses had
recombination rates similar to intraspecific crosses. Excluding BSc crosses did not yield better
correlations, suggesting an absence of relationship with evolutionary divergence, even at the
intraspecific level. The average length of recombination tracts was higher for BSc crosses and
had a significant positive correlation with parental divergence (Fig. 3C). Neither recombination
rate nor tract length correlated with marker counts (Fig. 3B,C), suggesting that these estimates
are largely unaffected by parental marker density. This indicates that mtDNA recombination rate
is independent of the global evolutionary divergence level between the parents of a hybrid,
contrary to the expectation that divergence should hamper recombination.

168
Figure 3. MtDNA protein-coding exons are enriched in recombination junctions . (*A*) Left: counts
of recombination tracts per mtDNA haplotype. Right: percentages of recombinant haplotypes.
(*B*) Left: recombination rate distributions for individual lines. Hollow circles show average
recombination rates. Right: Pearson's correlations between average recombination rate and
number of markers or parental evolutionary divergence (substitutions per site, based on nuclear
genome-wide variants). The correlation in red excludes BSc crosses. (*C*) Left: length
distributions for individual recombination tracts. Hollow circles show average recombination tract
lengths. Right: Pearson's correlations between average recombination tract length and number
of markers or parental evolutionary divergence (substitutions per site, based on nuclear
genome-wide variants). The correlation in red excludes BSc crosses. (*D*) Distributions of
annotation feature types for markers (top) and recombination breakpoints (bottom). (*E*)
Overrepresentation of each annotation feature type in recombination junctions, calculated as the
difference in frequency between breakpoints and markers for each feature type. FDR-corrected
p-values of chi-squared tests for frequency deviations within each cross are shown (***:
p≤0.001). (*F*) Percentage of nucleotide identity between parental mtDNAs for each annotation

feature type. (*G*) Overrepresentation of each annotation feature type in recombination junctions
against the percentage of nucleotide identity. Values for each cross are represented as
separate points.

We classified recombination breakpoints by annotation feature type and contrasted the resulting
distributions with parental markers (Fig. 3D). We found a strong overrepresentation of protein-
coding exons in recombination breakpoints in most crosses (Fig. 3E). We next quantified the
nucleotide identity between parental alleles of each cross for all mtDNA features, and asked
whether it correlated with overrepresentation in recombination junctions. High sequence identity
for some features in BSc crosses was consistent with inter-specific introgressions
(Supplemental Fig. S16), many of which were previously described (Leducq et al. 2017; Peris et
al. 2017). Despite these introgressions, a phylogenetic tree built on mtDNA-encoded protein-
coding DNA sequences showed the same topology as the nuclear-encoded genes, with all main
lineages having 100% branch support (Supplemental Figure S15B). While protein-coding exons
retain high nucleotide sequence identity in all crosses (Fig. 3F, Supplemental Figure S15C,D),
this does not solely explain their overrepresentation, as other annotation types displayed similar
identity levels (Fig. 3G, Supplemental Fig. S17). In summary, mtDNA recombination occurs
preferentially near protein-coding exon sequences and at a rate that is largely independent of
parental evolutionary divergence.

We asked whether the overrepresentation of exons in recombination junctions could be
explained by adjacent intron mobilization. Although intron mobility cannot be assessed strictly
from genomic data, we looked for genomic signatures of putative mobilization. We focused on
introns which show presence/absence polymorphism between the parents of each cross,
making the conservative assumption that all polymorphic introns can mobilize. We examined the
depth of coverage at intron-exon junctions, which shows distinctive profiles in the presence or
absence of an intron. We identified eight putative mobility events not involving recombination,
seven corresponding to *cox1*-I1 in BB1 and one corresponding to *cob*-I3 in BC1 (Supplemental
Fig. S18). A total of 15 recombination junctions were consistent with mobilization-mediated
recombination (Supplemental Fig. S19). Notably, *cox1*-I1 accounted for six of the eight putative
recombination-associated mobilization events in BB crosses. Although many intron
presence/absence polymorphisms were consistent with mobilization-associated recombination,
mobilization potentially explained only a minor fraction of the total recombination junctions (7/68
in BB1, 1/34 in BB2, 3/80 in BC1, 3/56 in BSc1 and 1/32 in BSc2).

Recombination may yield biases in the inheritance of mitochondrial alleles. MtDNA-wide
inheritance ratios did not correlate with parental mtDNA copy number ratios estimated from
depth of coverage of long-read sequencing libraries (Supplemental Fig. S20). However, copy
number estimates may have low accuracy since mtDNA abundance depends on the growth
phase (Galeota-Sprung et al. 2021) and we did not optimize the DNA extraction cultures for
synchronicity. They nevertheless suggest that parental mtDNA copy numbers do not determine
inheritance probability. We found significant deviations from mtDNA-wide inheritance ratios for
some genes (Supplemental Fig. S21). Similar patterns were found in the two BSc crosses, with
segments centered on *COX2* being mostly inherited from the *SpB* parent.

Parental divergence predicts the frequency of loss of respiratory function and large-scale
mtDNA deletions

We next characterized the loss of respiratory function in MA lines. The loss of respiration in
yeast is spontaneous and most often explained by the loss of mtDNA integrity through large-
scale deletions (Bernardi 1979). We measured the growth of all MA lines on media containing
fermentable (glucose, YPD) or non-fermentable (glycerol and ethanol, YPEG) carbon sources
using high-density colony array imaging. Since respiratory effects of mtDNA haplotypes or
mitonuclear interactions are often revealed with exposure to high temperatures (Li et al. 2019;
Paliwal et al. 2014), growth was measured at both 25°C and 37°C. We classified lines as non-
respiring if no growth was scored on YPEG at both temperatures. The frequency of respiration
loss was higher for more divergent crosses (Fig. 4A). We quantified the depth of sequencing
coverage along the mtDNA sequence and classified lines as harboring complete mtDNAs or
mtDNAs with deletions (Supplemental Fig. S22-32). All parental strains had complete mtDNAs.
The frequencies of mtDNAs with deletions closely matched the frequencies of respiration loss
(Fig. 4B), suggesting that the loss of mtDNA integrity is the main cause underlying changes in
respiratory function. Frequencies of deleted mtDNAs and had a significant positive correlation
with parental divergence (Pearson's correlation, r=0.88, p-value=3.5×10^{-4}, Fig. 4C). For most
crosses, deletions in mtDNAs had significantly negative effects on growth in YPD (Fig. 4D),
similar to the effect of respiration loss in *S. cerevisiae* (Ephrussi et al. 1949; Vowinckel et al.
2021). In contrast, mtDNA deletions were associated with complete loss of growth in YPEG
(Fig. 4D).

We asked whether the frequency of respiration loss in each cross could be explained by
additive parental effects. Spontaneous rates of respiration loss in parental strains were

estimated by measuring the frequency of non-respiring colonies, which display a characteristic
small size (petite) on medium with limited glucose. Parental petite frequencies ranged from
0.2% to 4.9% (Supplemental Fig. S33A). However, the average parental petite frequency for
each cross did not correlate with the percentage of lines that lost respiratory function (Pearson's
correlation, r=0.14, p-value=0.68; Supplemental Fig. S33B), indicating no additive parental
effects. In addition, we observed that stocks of the haploid parental strains contained variable
standing proportions of petite-forming cells (Supplemental Fig. S33C). The frequency of petite-
forming cells did not correlate with the fraction of MA lines from the initial timepoint harboring
non-recombinant, deleted mtDNAs of the corresponding parent (Pearson's correlation, r=-0.29,
p-value=0.36; Supplemental Fig. S33D). Thus, the frequency of respiration loss in MA crosses
appears to be the product of hybridization, rather than additive parental effects or direct
inheritance of defective mtDNAs.

A total of 25 MA lines had complete mtDNA haplotypes at the initial timepoint, but harbored
mtDNAs with large deletions at the final timepoint. From these, 12 were consistent with partial or
complete deletion of mtDNAs occurring during evolution (Supplemental Fig. S34). The
remaining 13 lines showed incompatible genotypes between timepoints, indicating that distinct
mtDNA haplotypes were still segregating in the lines at the initial timepoint. Likewise, ten lines
gained complete mtDNAs after evolution, which is only consistent with the segregation of
distinct haplotypes.

We asked if the genotype of non-deleted mtDNA haplotypes impacted the growth phenotypes of
respiring lines. The only significant difference was found between non-recombinant *SpB* and
SpA mtDNAs in the BA1 cross on YPEG at 37°C (Supplemental Fig. S35,36), with *SpB* having a
significant growth advantage (Mann-Whitney U test p-value=1.56×10^{-6}, FDR-corrected). Lines
with *SpB* mtDNAs also tended to grow better on YPEG at 37°C in the BC2 cross, which had
many of the largest effect sizes (Supplemental Fig. S36), hinting towards a similar
thermotolerance phenotype against *SpC* and recombinant mtDNAs. These results suggest that
mitochondrial genetic variation is linked to thermal and respiratory adaptations in wild *S.*
paradoxus.

Figure 4. Frequent deletions in mtDNAs lead to the loss of respiratory function and occur preferentially in crosses with higher parental divergence. (*A*) Percentage of all MA lines with loss of respiration. Counts of lines (at initial and final timepoints combined) are shown. (*B*) Percentage of sequenced MA lines per cross with mtDNA deletions. Counts of lines (initial and final timepoints combined) are shown. (*C*) Pearson's correlation between percentage of mtDNAs with deletions and parental divergence for each cross (substitutions per site, based on nuclear genome-wide variants). The correlation shown in red excludes BSc crosses. (*D*) Area under growth curves (AUC) for lines without and with mtDNA deletions in the four tested conditions. FDR-corrected p-values for Mann-Whitney U tests between mtDNAs without and with deletions, *: p≤0.05, **: p≤0.01, ***: p≤0.001. Images show colonies of representative lines (i.e. the closest to the median) after four days.

We next investigated which mtDNA segments were mostly involved in deletions. We compared

the depth of coverage profiles along mtDNAs and found recurring patterns among mtDNAs of

non-respiring lines (Fig. 5). Notably, many BSc haplotypes comprised only *COX2* and *COX3*.

This recurrent pattern emerged at two levels: among independently evolved lines, and in the

independent crosses BSc1 and BSc2. These results suggest that mtDNA deletions may be

predictable and repeatable in hybrids, and that high mtDNA deletion rates may be explained by

deletion hotspots.

Depth of coverage profiles from short read mappings are highly heterogeneous along the

mtDNA sequence, with some features displaying very low depth even in respiring lines (Fig. 5,

Supplemental Fig. S22-32). To circumvent the difficulty of identifying deletion breakpoints from

short reads, we randomly selected 122 lines from the final timepoint for long-read sequencing

and assembled their mtDNAs. 15 haplotypes harbored large-scale deletions, and most deletion

breakpoints were in unannotated regions (Supplemental Fig. S37, Supplemental Table S4). This

result indicates that the types of sequences involved in mtDNA deletions are different from

mtDNA recombination, suggesting distinct mechanisms.

303
Figure 5. Recurrent mtDNA deletion patterns emerge in non-respiring lines. The left part of
each heatmap shows the normalized depth of coverage of short-read sequencing libraries on
the reference mtDNA sequence, grouped by mtDNA features. The right part of each heatmap
shows the growth AUC in the four conditions tested. Rows represent individual MA lines
(including both initial and final timepoints). Lines are hierarchically clustered by depth of
coverage profile similarity. The top lines represent the parental strains.

Aspects of mtDNA evolution show contrasted associations between MA crosses

We next characterized the associations between mtDNA recombination, mtDNA deletion and

loss of respiration. All three were frequently observed at the initial timepoint, indicating that they

can occur shortly after F1 zygotes are formed. We also observed distinct haplotypes sampled

from the same line between timepoints, suggesting persistent heteroplasmy in some lines. We

thus analyzed the initial and final timepoints separately. As expected, there was a strong

negative relationship between mtDNA deletion and respiration (Fig. 6A,C). There were also

significant associations between mtDNA deletion and recombination in BA2 and BSc1, although

with opposite signs. While recombinant mtDNA haplotypes were more likely to harbor deletions

in BA2, only one BSc1 haplotype showed both recombination and deletion (Fig. 6B,D).

Figure 6. Associations between aspects of mtDNA evolution in MA lines. (*A*) Associations for lines at the initial timepoint. The heatmap shows the odds ratio for each pair of binary variables among mtDNA deletion, mtDNA recombination, respiration, mtDNA instability and aneuploidy instability. FDR-corrected p-values of Fisher's exact test, *: p≤0.05, **: p≤0.01, ***: p≤0.001. (*B*) Counts of MA lines for each comparison with a statistically significant association, classified as true (T) or false (F) for the given variable. The shaded background comprises associations between mtDNA deletion and respiration that were not statistically significant. (*C,D*) Same as A and B, but for lines at the final timepoint.

While most MA lines had the expected negative association between mtDNA deletions and respiration, we observed many exceptions. Many non-respiring lines had non-deleted mtDNAs, which could be explained by other types of mutations (mitochondrial or nuclear) leading to respiration loss. However, mtDNAs with large deletions (which are incompatible with respiration) were found in six and three respiring lines at the initial and final timepoints, respectively (Fig. 6B,D). Moreover, loss of respiration without deletion was found in six lines at the initial timepoint, all in BA2, which are unlikely explained by de novo mutations given the short evolution time. These inconsistencies may be explained by differences between the cryopreserved archive stocks used for genome sequencing, and the copies of those stocks used for phenotyping. Spot assays in the four conditions revealed that both stocks exhibited the same phenotypes (Supplemental Fig. S38), thus ruling out an effect of stock propagation.

We looked for genomic changes other than mtDNA deletions that could have occurred rapidly in MA line evolution and explain respiration loss. We searched for de novo nucleotide variants in mtDNAs of the MA lines by looking for confident variants that are private to single lines. We found a handful of candidate de novo mutations at the initial timepoint but none were in coding regions (Supplemental Table S5), making them poor candidates to explain respiration loss of function. We next focused on aneuploidies in the nuclear genome, which were previously shown to arise early in MA (Fijarczyk et al. 2021; Marsit et al. 2021). Several inconsistent lines had aneuploidies, including five BA2 lines with monosomies for chromosomes I, III, VI and IX and/or trisomies for chromosomes IV and XII (Supplemental Fig. S39). However, parental allele frequencies on these chromosomes closely matched the diploid expectation of 50%. We quantified the deviation in depth of coverage from the value expected given the copy number call for each chromosome. Many lines with aneuploidy calls showed high deviations from the expectation of true aneuploidies (Supplemental Fig. S39A). We validated that these deviations were not artifacts from variation in library size (Supplemental Fig. S40). Continuous variation in depth of coverage and symmetrical parental allele frequencies are consistent with unbiased, segregating aneuploidies rather than fixed aneuploidies. These aneuploidies arose from single

euploid clones in the population expansion leading to DNA extraction. We thus term this
phenotype aneuploidy instability.

Focusing on BA2 at the initial timepoint, we assessed the presence of two mtDNA loci (*ATP6*
and 21S rRNA) by PCR for all lines (Supplemental Table S6). While globally consistent with the
sequencing libraries, both PCR amplicons were detected in the two respiring lines genotyped as
having mtDNA deletions (Supplemental Fig. S39). Furthermore, at least one amplicon was
missing for five out of six non-respiring lines with complete mtDNAs. This suggests that minor
mtDNA variants segregating in the archive stocks were selected by chance when single clones
were isolated for genome sequencing. We term the inconsistency between mitochondrial
deletion and respiration mtDNA instability. BA2 had the highest proportion of lines with mtDNA
instability, and those were primarily found at the initial timepoint (Supplemental Fig. S41A). We
found a statistically significant association of mtDNA instability with aneuploidy instability in BA2
at the initial timepoint (Fig. 6B). Out of eight lines displaying mtDNA instability, five also were
among the lines with the highest aneuploidy instability. There was a similar trend in other
crosses, although only BA2 was statistically significant (Supplemental Fig. S41B). This suggests
that a proportion of hybrid lines, specifically in the BA2 cross, suffered from high genomic
instability, manifested by both frequent aneuploidy and mtDNA deletion.

DISCUSSION

How diverged mtDNA haplotypes interact to shape the resolution of heteroplasmy and the
evolution of mitochondrial functions remains vastly unexplored. In this study, we used
Saccharomyces yeasts as a model to probe the neutral evolution of mtDNAs in heteroplasmic
hybrids through experimental evolution by MA. We uncovered that recombination rates are
independent of parental divergence, and recombination occurs preferentially near protein-
coding exons. On the other hand, parental divergence is a strong predictor of the rate of large-
scale mtDNA deletions, which is the primary mechanism for loss of respiratory metabolism. We
also uncovered that some crosses have contrasted associations between mtDNA recombination
and deletion, as well as shared genomic instability at the level of mtDNAs and nuclear
aneuploidies.

Recombination between mtDNAs in heteroplasmic *S. cerevisiae* has been known for decades
(Dujon et al. 1974). Recombination is likely central in the replication of mtDNAs, making it an
inherent part of mitochondrial genetics rather than an accidental byproduct of heteroplasmy

(Dujon 2020). Genomic studies brought further support for the idea that mtDNA recombination is
pervasive (Fritsch et al. 2014; Bágeľová Poláková et al. 2021). Fritsch and collaborators
characterized recombination in pools of diploid F1 hybrids from intraspecific crosses of *S.*
cerevisiae, and found that recombination hotspots were concentrated in tRNA clusters and
rRNA genes (Fritsch et al. 2014). Bágeľová Poláková and collaborators investigated three
recombinant mtDNA haplotypes in F1 hybrids between *S. cerevisiae* and *S. paradoxus*, and
attributed most of the observed recombination events to the mobilization of introns and free-
standing ORFs (Bágeľová Poláková et al. 2021). Thus, our study is the first large-scale report of
mtDNA recombination at the intra- and inter-specific levels, allowing the resolution of 482
independent mtDNA haplotypes. Our data confirms that mtDNA recombination is pervasive in a
variety of hybrid genotypes. In contrast to previous studies, we found an association of
recombination junctions with protein-coding exons and only a minor potential contribution for
putative intron mobilization.

We observed associations between mtDNA recombination and deletion that suggest a
mechanistic interaction, although this remains to be investigated. The most contrasting patterns
were found between BA2 and BSc1, two crosses with frequent mtDNA deletions. In BA2,
haplotypes harboring deletions were almost always recombinant, and non-deleted haplotypes
rarely showed evidence for recombination. In contrast, in BSc1, mtDNA recombination and
deletion were almost mutually exclusive. This is consistent with differences in mtDNA
architecture between *S. paradoxus* lineages *SpB* and *SpA*. While *SpB* mtDNAs are collinear
with other North American lineages and with *S. cerevisiae*, *SpA* harbors rearranged mtDNAs
(Leducq et al. 2017; Yue et al. 2017). As such, homologous recombination between rearranged
mtDNAs could lead to frequent deletions. However, deletion breakpoints were often located in
unannotated regions, contrasting with recombination breakpoints and suggesting a distinct
mechanism. The association between mtDNA instability and aneuploidy instability in BA2 is
more challenging to interpret. BA2 was also the cross with the most frequent segregation of
distinct mtDNA haplotypes between timepoints, suggesting rampant, persistent heteroplasmy.
Nevertheless, it raises the possibility that both mtDNA and aneuploidy instability could be linked
or share a common underlying cause. These types of genomic instability in BA2 add to many
others previously characterized in our MA crosses (Marsit et al. 2021).

Experimental evolution by MA minimizes the efficiency of natural selection, enabling a
selectively unbiased characterization of mtDNA evolution. The characterization of de novo
mutation in nuclear genomes of our MA lines revealed signatures that were entirely consistent

with neutral evolution (Fijarczyk et al. 2021). The frequent loss of respiratory function also
suggests the absence of purifying selection. Although large-scale deletions leading to
respiration mostly happened before the initial timepoint, a dozen MA lines had mtDNA
haplotypes consistent with deletion occurring between the initial and final timepoints. While the
MA design minimized the efficiency of natural selection at the level of individual cells, it did not
alleviate it at the intracellular level. MtDNA haplotypes can outcompete others if they have a
replication advantage, regardless of their effect on organismal fitness (Ma and O'Farrell 2016;
Taylor et al. 2002). As such, some mtDNA haplotypes could have reached homoplasmy
because of replication advantages.

Our results show that respiratory function is primarily lost through large-scale deletions within
mtDNAs. The level of replication of our MA experiment yielded a high number of recombinant
haplotypes. Nevertheless, cases of respiration loss in our experiment are likely entirely
explained by mtDNA deletions. Additionally, we found no significant negative effect of non-
deleted, recombinant mtDNA haplotypes on respiration. In BSc1, the non-significant growth
difference between lines with *SpB* versus recombinant haplotypes on YPEG at 25°C was the
second largest effect size, suggesting that some recombinant haplotypes might be
disadvantaged. Despite this hypothetical case, we found no support for negative mito-mito
epistasis, contrary to previous reports (Wolters et al. 2018; Leducq et al. 2017).

Mitonuclear incompatibilities are known to contribute to reproductive isolation between species
of the *Saccharomyces* genus (Chou et al. 2010; Lee et al. 2008; Jhuang et al. 2017), along with
other types of reproductive barriers (Hunter et al. 1996). Even at the intraspecific level in *S.*
cerevisiae and *S. paradoxus*, crosses between diverged isolates often have low fertility (Greig et
al. 2003; Charron et al. 2014; Leducq et al. 2016; Eberlein et al. 2019), including most of our MA
crosses (Charron et al. 2019; Marsit et al. 2021). Our results show that loss of respiration adds
another reproductive barrier, since meiosis (sporulation) is dependent on functioning respiratory
metabolism (Jambhekar and Amon 2008). Yeast can divide strictly mitotically in permissive
laboratory conditions, and mitosis is dominant in the wild (Tsai et al. 2008). However,
sporulation is triggered in starvation conditions (Bautz Freese et al. 1982) and is likely to be
essential in nature, thus selecting for mtDNA integrity. Surprisingly, natural isolates of *S.*
cerevisiae were reported to completely lack mtDNA (De Chiara et al. 2020), suggesting that loss
of sporulation ability can arise in nature. As such, mtDNA degeneration in hybrids acts as an
additional mechanism of reproductive isolation, both within and between species.

In conclusion, our study provides the first large-scale investigation of neutral mtDNA evolution in
heteroplasmic hybrids. Our results show that distinct aspects of neutral mtDNA evolution can be
either genotype-specific or, in contrast, highly predictable given the genomic composition of
hybrids. We show that hybridization can trigger mtDNA degeneration with profound metabolic
consequences and the emergence of reproductive isolation.

METHODS

Long read library preparation and sequencing

Stocks of the 13 haploid parental strains (Supplemental Table S1) and of 122 randomly selected
MA lines from the final timepoint (Charron et al. 2019; Hénault et al. 2020) were thawed,
streaked on YPD agar medium (1% yeast extract, 2% bio-tryptone, 2% glucose, 2% agar) and
incubated at room temperature for three days. Cultures in YPD medium (1% yeast extract, 2%
bio-tryptone, 2% glucose) were inoculated with single colonies and incubated overnight at room
temperature without shaking. DNA was extracted from the cultures following a standard phenol-
chloroform protocol. Oxford Nanopore native (PCR-free) genomic DNA libraries were prepared
in multiplex with kits SQK-LSK109 and EXP-NBD104 (Oxford Nanopore Technologies, Oxford,
UK). Libraries were sequenced with FLO-MIN106 (revC) flowcells on a MinION sequencer
(MIN-101B). The sequencing and basecalling were run on a MinIT computer (MNT-001) with
MinKNOW v3.3.2 and guppy v3.0.3. Reads were demultiplexed using guppy_basecaller v3.1.5.

Long read mtDNA assemblies

Long reads matching mtDNA sequences were extracted from whole-genome libraries. For each
strain, we made a compound reference genome comprising the nuclear genome of the
corresponding *S. paradoxus* lineage(s) (*SpA*: LL2012_001, *SpB*: MSH-604, *SpC*: LL2011_012)
(Eberlein et al. 2019), or *S. cerevisiae* (YPS128) (Yue et al. 2017), supplemented with a
concatenated mitochondrial contig comprising *S. paradoxus SpB* (YPS138), *SpA* (CBS432) and
(YPS138) and *S. cerevisiae* (S288c) (Yue et al. 2017) mtDNAs joined by 100 Ns. Nanopore
libraries were mapped on the compound references using minimap2 v2.20-r1061 (Li 2018a)
with preset map-ont. Reads mapping to the mtDNA contig were extracted using samtools v1.15
(Li et al. 2009) and seqtk v1.3-r106 (Li 2018b). Assembly of the extracted reads was performed
using wtdbg2 v2.5 (Ruan and Li 2019) with options –AS 2 –g 72k –X 50 and a grid search of the
following parameters: –k {23, 21, 19, 17, 15, 13, 11}, –p {8, 5, 2, 1, 0}, –l {4096 2048 1024}.

For parental strains, draft assemblies were aligned against the reference mtDNA assembly (Yue
et al. 2017) of the closest species/lineage (*SpB* and *SpC*: YPS138, *SpA*: CBS432, *S. cerevisiae*:
S288c) using MUMmer v3.23 (Kurtz et al. 2004). For each strain, the best assembly was
selected on the basis of a quality score computed from the alignment of each query assembly
using custom Python v3.9.10 (Van Rossum and Drake 2009) scripts. A genome coverage
metric was first computed as the sum of absolute deviations from 1 of both query and reference
coverage. A contiguity metric was computed as the inverse of the largest contig size in the
assembly. Both coverage and continuity metrics were normalized to z-scores and summed to
yield the final score. Assemblies with the lowest score were circularized at the start of the *ATP6*
gene and resulting duplications were manually trimmed. Circularized assemblies were polished
by first aligning the basecalled long reads to draft assemblies using minimap2 v2.20-r1061 with
preset map-ont. Alignments were polished using the raw nanopore signal using nanopolish
v0.13.3 (Loman et al. 2015) with parameter ––min-candidate-frequency 0.1. Short read libraries
for each parental strain (Hénault et al. 2020; Marsit et al. 2021) were aligned on draft
assemblies using bwa-mem v0.7.17 and polishing was performed using Pilon v1.22 (Walker et
al. 2014). Assemblies were aligned against the reference mtDNA sequence of *S. cerevisiae*
S288c (Yue et al. 2017) using MUMmer v3.23. For strains YPS744, LL2011_012, MSH-587-1,
MSH-604 and UWOPS-91-202, assemblies were compared to the corresponding mtDNA short-
read assemblies from (Leducq et al. 2017) using the dnadiff tool from the MUMmer v3.23 suite.

For MA lines, draft assemblies were polished with basecalled long reads using medaka v1.4.4
(https://github.com/nanoporetech/medaka) with model r941_min_high_g303. Assemblies were
circularized at the start of the *ATP6* gene (if present) and aligned to the artificial mtDNA
reference (see below) using MUMmer v3.23.

MtDNA assembly annotation and generation of mtDNA artificial reference

MtDNA assemblies of the 13 parental strains were annotated using the MFannot server
(https://megasun.bch.umontreal.ca/cgi-bin/mfannot/mfannotInterface.pl) and annotations were
manually curated. MFannot master files were converted to the GFF3 format using the
mfannot2gff.pl script from the MITONOTATE pipeline (Seah 2016). A reference library of intron
sequences was assembled from the mtDNA assemblies and annotations of *S. paradoxus*
strains CBS432 (GenBank accession JQ862335.1) (Procházka et al. 2012) and CBS7400
(GenBank accession KX657749.1) (Sulo et al. 2017), and *S. cerevisiae* S288c (GenBank
accession AJ011856.1) (Foury et al. 1998). Introns annotated from our assemblies were

renamed after their best hit from a search against the reference library using BLASTN v2.11.0
(Camacho et al. 2009), following the nomenclature of (Lambowitz and Belfort 1993). From the
comparative analysis of mtDNA annotations, we built an artificial mtDNA reference comprising
the largest set of features. We used the LL2012_028 genome as a template, and added introns
cob-I3, *cob*-I4, *cob*-I5 and *cox1*-I3γ from YPS744 or LL2013_054 at the orthologous splice sites
using custom Python v3.9.10 scripts. GC clusters on the artificial mtDNA reference were
annotated using the four reference sequences presented in Table 1 of (Wu and Hao 2015) as
queries for BLASTN v.2.11.0 with parameter word_size 7. The resulting hits were collapsed
using bedtools merge v2.30.0 (Quinlan and Hall 2010) and manually inspected to add or merge
adjacent GC-rich segments. Parental mtDNAs were aligned to the artificial mtDNA contig using
mugsy v1r2.3 (Angiuoli and Salzberg 2011), and liftover tables were built using a custom Python
v3.9.10 script.

Phylogenetic analysis of protein-coding genes

Maximum likelihood phylogenetic trees were constructed using protein-coding genes of the
nuclear and mitochondrial genomes. For mtDNAs, spliced protein-coding sequences of the eight
canonical genes were aligned using MUSCLE v3.8.31 (Edgar 2004) and concatenated into a
multiple sequence alignment of 6699 positions. For nuclear genomes, 81 orthogroups (each
comprising a total of 26 *S. paradoxus* and *S. cerevisiae* sequences) were randomly chosen from
(Eberlein et al. 2017), and protein-coding DNA sequences were concatenated to yield a multiple
sequence alignment of 124,041 positions. Maximum likelihood phylogenetic trees were
computed using RAxML_NG v1.1.0 (Kozlov et al. 2019) with the GTR+G model initiated with 10
parsimony trees and 200 bootstraps. For mitochondrial genes, the evolutionary divergence
between the parents of each cross were extracted. For nuclear genes, the evolutionary
divergence value of each cross was computed as the average of all pairwise distances within or
between the corresponding parental lineage(s) or species. Phylogenetic trees were built for
individual spliced protein-coding genes, introns and ORFs). DNA sequences were aligned using
MUSCLE v3.8.31 and trees were inferred using the neighbor-joining algorithm implemented in
the ape v5.6-2 library (Paradis and Schliep 2019) in R v4.2.1 (R Core Team 2016).

Short read alignment and variant calling

Short read libraries corresponding to the 13 parental haploid strains and 447 hybrid yeast MA
lines, sampled at both the initial (~60 mitotic generations after zygote formation) and final (after

~770 mitotic generations) timepoints, were downloaded from NCBI (BioProject PRJNA515073).
Short reads were trimmed using Trimmomatic v0.36 (Bolger et al. 2014) with parameters
ILLUMINACLIP:{custom adapters file}:2:30:10 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36
(Hénault et al. 2020). We built a concatenated reference genome comprising nuclear contigs of
S. paradoxus SpB strain MSH-604 (Eberlein et al. 2019) and *S. cerevisiae* strain YPS128 (Yue
et al. 2017), to which we appended the artificial mtDNA contig. Reads were aligned to this
reference using bwa mem v0.7.17 and mappings to the mtDNA contig were extracted using
samtools v1.8. Variant calling was performed separately for each cross using freebayes v1.3.5
(Garrison and Marth 2012). Multiallelic variants were decomposed using vcfbreakmulti from
vcflib v1.0.2 (Garrison 2018).

Definition of mtDNA recombination tracts

The characterization of recombination in mtDNA haplotypes was performed using custom
Python v3.9.10 scripts. For each cross, marker variants were identified for the parental
genomes based on variant calls from short-read alignments on the artificial mtDNA reference.
Candidate markers were filtered as follows. Each variant had to be supported by at least two
read alignments. Parental alleles at each marker locus were validated using whole-genome
alignments of the high-quality genome assemblies of each parental mtDNA and the associated
liftover tables. Markers were required to have nucleotide calls and assembly alignments for both
parents, thus excluding insertion and deletions.

For MA lines of each cross, variants corresponding to parental markers were filtered to exclude
loci with low depth of coverage. For each line we defined a threshold as the highest value
between the 10th percentile of depth of coverage values for the nuclear genome of each line, or
20 reads. Variants passed the filter if supporting reads for the major allele were higher than the
threshold. Since each MA line was sampled for sequencing at two timepoints, mtDNA
haplotypes were compared across timepoints of individual lines to determine whether they were
clonal or independent haplotypes that segregated from initially heteroplasmic lines. For each
line, we computed the pairwise identity between mtDNA haplotypes as the proportion of
identical calls at marker variants (excluding missing positions in either mtDNA). If the identity
was 95% or higher, we kept only one mtDNA (preferentially the latest timepoint) as
representative of the line. Independent haplotypes within each cross were clustered by identity
using the nearest point algorithm from scipy v1.8.0 (Virtanen et al. 2020).

Recombination tracts were defined by scanning and grouping adjacent marker variants of the
same genotype along the genome, with a tolerance of one marker of the opposite genotype (i.e.
two marker variants of the opposite genotype were required to break a tract). Tracts were
circularized at the ends of the mtDNA contig. Recombination breakpoints were defined as the
marker variants at the ends of each tract. Recombination breakpoint and marker variant
densities were computed for 100 bp windows along the genome. Recombination rates were
computed as the number of recombination tracts per haplotype divided by the combined length
of parental mtDNAs. Average recombination rate and tract length for each cross were correlated
with marker variant counts and parental evolutionary divergence values (see section
Phylogenetic analysis of protein-coding genes).

MtDNA inheritance biases

Recombination tracts were used to compute the proportion of each mtDNA haplotype that was
inherited from each parent. The inheritance ratios were averaged over all independent
haplotypes of a cross to yield genome-wide average inheritance ratios. The depth of coverage
of long reads on each parental mtDNA assembly compared to nuclear chromosomes was used
to compute a relative mtDNA copy number. For individual genes within haplotypes, discrete
inheritance calls were generated by attributing either parental ancestry to annotations that had
ratios of 10% or less, or 90% or more. Genes with ratios between those values were considered
as recombinant and excluded. Binomial tests were used to test the deviation of non-
recombinant genes from genome-wide inheritance ratios.

Growth measurements of the MA lines collection

During the MA experiment, MA lines were sampled at regular intervals and archived in
cryopreserved stocks in 96 well microplates. Archives stocks sampled at the initial and final
timepoints were replicated on YPD medium and cryopreserved as stock copies. For high-
throughput phenotyping, stock copies were thawed and spotted on YPD agar medium in
OmniTray plates (Thermo Fisher Scientific, Waltham, USA) using a BM5-SC1 colony
processing robot (S&P Robotics Inc., Toronto, Canada) and incubated at room temperature for
four days. Colonies were replicated on YPD agar by condensing them into six 384-position
source arrays and incubated at room temperature for three days. Source arrays were replicated
twice on YPD agar medium and twice on YPEG agar medium (1% yeast extract, 2% bio-
tryptone, 3% glycerol, 3% ethanol, 2% agar). Plates were incubated at 25°C or 37°C for the first

round of selection during two days. Plates were replicated for the second round of selection on
the same media and photographed after 24, 48 and 96 hours of incubation with the robotic
platform. Colony sizes were quantified from the plate photos using gitter v1.1 (Wagih and Parts
2014) in R v3.3.2 (R Core Team 2016). Colony sizes in pixels were transformed in log and
summed across imaging timepoints to estimate the area under growth curves (AUC). One
colony (BSc2, line F53, initial timepoint, YPD 37°C) was removed due to visible contamination.
Calls of growth absence based on pixel quantifications were manually curated with raw images.

Measurement of parental petite frequencies

Parental rates of spontaneous petite formation were measured by streaking haploid stocks on
YPD agar + NAT (100 µg ml^{-1} nourseothricin) or YPD agar + G418 (200 µg ml^{-1} geneticin),
depending on the selection marker at the *HO* locus, and incubated at 25°C for six days. Two
single large colonies per strain were resuspended in sterile water and diluted to an optical
density at 600 nm (OD$_{600}$) of 2.5×10^{-4}, and 200 µl were plated on YPdG agar medium (1% yeast
extract, 2% bio-tryptone, 3% glycerol, 0.1% glucose, 2% agar). Plates were incubated at 25°C
for four days and photographed. The standing frequency of non-respiring cells within parental
stock cultures was measured by spotting 20 µl of stocks on YPD agar and incubating for three
days. Cells were resuspended in sterile water and diluted to an OD$_{600}$ of 5×10^{-4}, and 100 µl were
plated on YPdG agar. Plates were incubated at 25°C for five days and photographed.

Characterization of mtDNA deletions

Depth of coverage values along mtDNAs were extracted from short read alignments using
samtools v1.15. Median depth values were computed for 100 bp windows and normalized by
the depth of coverage on nuclear chromosomes. Depth values were grouped by annotation type
and the fraction of positions with normalized depth values higher than 0.1 was extracted. For
each cross, these fractions were normalized into z-scores, and an mtDNA completion score was
computed as the sum of z-scores across annotation types for individual mtDNA haplotypes.
Haplotypes were sorted by mtDNA completion score to classify them as complete or deleted,
and the classification was manually curated by examination of the depth of coverage profiles.
Haplotypes were clustered by Pearson's correlation coefficient between normalized depth
profiles using the nearest point algorithm from scipy v1.8.0.

Detection of intron presence/absence polymorphisms

Presence or absence of introns in mtDNA haplotypes of MA lines was called by analyzing depth of coverage profiles at intron-exon junctions. For each polymorphic intron, the marker variants closest to both junctions (upstream or downstream) were identified. 100 bp windows centered around each junction were used to analyze depth of coverage profiles. Depth of coverage values were normalized by the 80^{th} percentile of depth values within a window. Euclidean distance and Pearson's correlation coefficient were used to compare profiles of each line with both parental profiles. Evidence for intron presence was called at a junction if the euclidean distance and Pearson's correlation coefficient were respectively lower and higher for comparisons to the parent harboring the intron. An intron was called present if its two junctions showed consistent evidence. Putative intron mobilization events not involving recombination were defined as intron presence with two flanking markers of the opposite ancestry. Putative recombination-associated mobilization events were defined as intron presence with one flanking marker of the opposite ancestry, consistent with neighboring recombination tracts.

Confirmation of growth phenotypes by spot assays

A subset of lines classified as having mtDNA instability at the initial timepoint were selected for growth phenotypes validation by spot assays. Original archive stocks and copies were thawed, spotted on YPD agar medium and incubated at room temperature. Spots were used to inoculate precultures in YPD medium that were incubated overnight at 30°C. Precultures were diluted at 1 OD ml^{-1} in sterile water and five 5X serial dilutions were prepared. Aliquots of 5 µl of each dilution were spotted twice on YPD agar and YPEG agar. Plates were incubated at 25°C or 37°C for three days and photographed.

De novo mutations identification

Candidate de novo single nucleotide mutations were identified from variant calls based on short read alignments. First, variant calls supported by at least five reads and which had an alternative allele ratio of at least 0.8 were extracted. Variants which were only observed in a single line were further checked for the absence of matching alleles in the parental variant calls from all crosses, and from lines of all the other crosses. For each remaining candidate, we extracted the proportion of lines in the cross with missing data at that locus. We defined a more stringent alignment criteria for supporting reads by requiring the length of the aligned reference segment to be between 141 and 161 bp (for 151 bp reads), and counted those high confidence

alignments for each candidate variant. Finally, we counted the number of supporting reads for

matching alleles in the whole dataset, excluding the line of interest. The final set of candidate de

novo mutations consisted in variants for which the total number of supporting reads was at least

twice the number of supporting reads in the whole dataset excluding the line, and had at least

two supporting reads with high confidence alignments.

Depth of coverage deviations on nuclear chromosomes

Variant calls from short-read alignments and aneuploidy calls were previously done by Fijarczyk

and collaborators (Fijarczyk et al. 2021). BSc crosses were excluded from this analysis as the

mapping strategy was different (using concatenated *SpB* and *S. cerevisiae* nuclear reference

genomes). For each cross, 10000 nuclear variants were chosen at random. Allele ratios were

examined along nuclear chromosomes. For individual lines, the median depth of coverage of

each chromosome was normalized by the genome-wide median to yield the depth ratio. The

copy number ratio was defined by dividing chromosome copy number calls by the global ploidy

level of each line. Depth of coverage deviation scores were computed as (depth ratio – copy

number ratio). Absolute deviation scores were averaged over all chromosomes and lines with

an average absolute deviation score of at least 0.08 were classified as having the aneuploidy

instability phenotype.

MtDNA genotyping by PCR

Two loci were genotyped for presence or absence by PCR for all lines of the BA2 cross at the

initial and final timepoints. DNA was extracted by resuspending colonies in 40 µl NaOH 20 mM

and incubating at 95°C for 20 min. Two mtDNA loci in *ATP6* and 21S rRNA were amplified with

primers and cycles detailed in Supplemental Table S6 (Leducq et al. 2017). Amplicons of the

nuclear mating type loci were amplified as positive controls with primers and cycles detailed in

Supplemental Table S6 (Huxley et al. 1990). Presence and absence were scored by agarose

gel electrophoresis.

DATA ACCESS

Long read datasets are available at NCBI under accession PRJNA828354. Scripts, genome

assemblies and annotations are available at https://github.com/Landrylab/mito_ma.

COMPETING INTEREST STATEMENT

We declare no competing interest.

ACKNOWLEDGEMENTS

We thank Anna Fijarczyk for comments on the manuscript and contributions to sequencing data
preprocessing and analysis. We thank Hélène Martin for contributions to sequencing data
preprocessing. We thank the members of the Landry laboratory for discussions on the project.
This project was supported by funding to C.R.L. from a NSERC discovery grant and FRQNT
team grant, and a NSERC Alexander Graham Bell doctoral scholarship to M.H. CRL holds the
Canada Research Chair in Cellular Systems and Synthetic Biology.

AUTHOR CONTRIBUTIONS

CRL and MH designed the project. MH performed the analyses and experiments. MH, SM and
GC performed the MA evolution experiment and short read sequencing as cited. MH wrote the
manuscript with inputs from CRL, SM and GC.

REFERENCES

Adams KL, Palmer JD. 2003. Evolution of mitochondrial gene content: Gene loss and transfer to
the nucleus. *Mol Phylogenet Evol* **29**: 380–395.

Angiuoli SV, Salzberg SL. 2011. Mugsy: Fast multiple alignment of closely related whole
genomes. *Bioinformatics* **27**: 334–342.

Bágeľová Poláková S, Lichtner Ž, Szemes T, Smolejová M, Sulo P. 2021. Mitochondrial DNA
duplication, recombination, and introgression during interspecific hybridization. *Sci Rep* **11**:
12726.

Bautz Freese E, Chu MI, Freese E. 1982. Initiation of Yeast Sporulation by Partial Carbon,
Nitrogen, or Phosphate Deprivation. *J Bacteriol* **149**: 840–851.

Bernardi G. 1979. The petite mutation in yeast. *Trends Biochem Sci* **4**: 197–201.

Birky CW Jr. 2001. The inheritance of genes in mitochondria and chloroplasts: laws,
mechanisms, and models. *Annu Rev Genet* **35**: 125–148.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**: 2114–2120.

Breton S, Stewart DT, Bonen L. 2015. Atypical mitochondrial inheritance patterns in eukaryotes.
Genome **58**: 423–431.

Burton RS. 2022. The role of mitonuclear incompatibilities in allopatric speciation. *Cell Mol Life*
726   *Sci* **79**: 103.

727 Burton RS, Barreto FS. 2012. A disproportionate role for mtDNA in Dobzhansky-Muller
728   incompatibilities? *Mol Ecol* **21**: 4942–4957.

729 Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
730   BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 421.

731 Charron G, Leducq JB, Landry CR. 2014. Chromosomal variation segregates within incipient
732   species and correlates with reproductive isolation. *Mol Ecol* **23**: 4362–4372.

733 Charron G, Marsit S, Hénault M, Martin H, Landry CR. 2019. Spontaneous whole-genome
734   duplication restores fertility in interspecific hybrids. *Nat Commun* **10**.
735   http://dx.doi.org/10.1038/s41467-019-12041-8.

736 Chou J-Y, Hung Y-S, Lin K-H, Lee H-Y, Leu J-Y. 2010. Multiple Molecular Mechanisms Cause
737   Reproductive Isolation between Three Yeast Species ed. M.A.F. Noor. *PLoS Biol* **8**:
738   e1000432.

739 De Chiara M, Friedrich A, Barré B, Breitenbach M, Schacherer J, Liti G. 2020. Discordant
740   evolution of mitochondrial and nuclear yeast genomes at population level. *BMC Biol* **18**: 49.

741 Dujon B. 2020. Mitochondrial genetics revisited. *Yeast* **37**: 191–205.

742 Dujon B, Slonimski PP, Weill L. 1974. Mitochondrial genetics IX: A model for recombination and
743   segregation of mitochondrial genomes in saccharomyces cerevisiae. *Genetics* **78**: 415–
744   437.

745 Eberlein C, Hénault M, Fijarczyk A, Charron G, Bouvier M, Kohn LM, Anderson JB, Landry CR.
746   2019. Hybridization is a recurrent evolutionary stimulus in wild yeast speciation. *Nat*
747   *Commun* **10**: 923.

748 Eberlein C, Nielly-Thibault L, Maaroufi H, Dubé AK, Leducq J-B, Charron G, Landry CR. 2017.
749   The Rapid Evolution of an Ohnolog Contributes to the Ecological Specialization of Incipient
750   Yeast Species. *Mol Biol Evol* **34**: 2173–2186.

751 Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
752   throughput. *Nucleic Acids Res* **32**: 1792–1797.

753 Ephrussi B, Hottinguer H, Tavlitzki J. 1949. Action de l'acriflavine sur les levures. I. La mutation"
754   petite colonie." *Ann Inst Pasteur* **76**: 351–367.

755 Fijarczyk A, Hénault M, Marsit S, Charron G, Landry CR. 2021. Heterogeneous Mutation Rates
756   and Spectra in Yeast Hybrids. *Genome Biol Evol* **13**.
757   http://dx.doi.org/10.1093/gbe/evab282.

758 Foury F, Roganti T, Lecrenier N, Purnelle B. 1998. The complete sequence of the mitochondrial
759   genome of Saccharomyces cerevisiae. *FEBS Lett* **440**: 325–331.

760 Fritsch ES, Chabbert CD, Klaus B, Steinmetz LM. 2014. A genome-wide map of mitochondrial
761   DNA recombination in yeast. *Genetics* **198**: 755–771.

Galeota-Sprung B, Fernandez A, Sniegowski P. 2021. Changes to the mtDNA copy number
during yeast culture growth. *bioRxiv* 2021.09.02.458779.
https://www.biorxiv.org/content/10.1101/2021.09.02.458779v4.full (Accessed March 1,
2022).

Garrison E. 2018. *vcflib*. https://github.com/vcflib/vcflib.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing.
arXiv preprint arXiv:12073907 [q-bioGN]. http://arxiv.org/abs/1207.3907.

Greig D, Travisano M, Louis EJ, Borts RH. 2003. A role for the mismatch repair system during
incipient speciation in Saccharomyces. *J Evol Biol* **16**: 429–437.

Gyllensten U, Wharton D, Josefsson A, Wilson AC. 1991. Paternal inheritance of mitochondrial
DNA in mice. *Nature* **352**: 255–257. http://dx.doi.org/10.1038/352255a0.

Hénault M, Marsit S, Charron G, Landry CR. 2020. The effect of hybridization on transposable
element accumulation in an undomesticated fungal species. *Elife* **9**.
http://dx.doi.org/10.7554/eLife.60474.

Hunter N, Chambers SR, Louis EJ, Borts RH. 1996. The mismatch repair system contributes to
meiotic sterility in an interspecific yeast hybrid. *EMBO J* **15**: 1726–1733.

Huxley C, Green ED, Dunham I. 1990. Rapid assessment of S. cerevisiae mating type by PCR.
Trends Genet **6**: 236.

Jambhekar A, Amon A. 2008. Control of meiosis by respiration. *Curr Biol* **18**: 969–975.

Jhuang HY, Lee HY, Leu JY. 2017. Mitochondrial-nuclear co-evolution leads to hybrid
incompatibility through pentatricopeptide repeat proteins. *EMBO Rep* **18**: 87–101.

Joseph SB, Hall DW. 2004. Spontaneous mutations in diploid Saccharomyces cerevisiae: more
beneficial than expected. *Genetics* **168**: 1817–1825.

Kondo R, Satta Y, Matsuura ET, Ishiwa H, Takahata N, Chigusa SI. 1990. Incomplete maternal
transmission of mitochondrial DNA in Drosophila. *Genetics* **126**: 657–663.

Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAxML-NG: a fast, scalable and
user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**: 4453–
4455.

Kuehne HA, Murphy HA, Francis CAA, Sniegowski PD. 2007. Allopatric Divergence, Secondary
Contact, and Genetic Isolation in Wild Yeast Populations. *Curr Biol* **17**: 407–411.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004.
Versatile and open software for comparing large genomes. *Genome Biol* **5**: R12.

Ladoukakis ED, Theologidis I, Rodakis GC, Zouros E. 2011. Homologous recombination
between highly diverged mitochondrial sequences: examples from maternally and
paternally transmitted genomes. *Mol Biol Evol* **28**: 1847–1859.

Ladoukakis ED, Zouros E. 2017. Evolution and inheritance of animal mitochondrial DNA: rules
and exceptions. *J Biol Res* **24**: 2.

Lambowitz AM, Belfort M. 1993. Introns as mobile genetic elements. *Annu Rev Biochem* **62**:
587–622.

Leducq J-B, Charron G, Samani P, Dubé AK, Sylvester K, James B, Almeida P, Sampaio JP,
Hittinger CT, Bell G, et al. 2014. Local climatic adaptation in a widespread microorganism.
Proc Biol Sci **281**: 20132472.

Leducq J-B, Henault M, Charron G, Nielly-Thibault L, Terrat Y, Fiumera HL, Shapiro BJ, Landry
CR. 2017. Mitochondrial Recombination and Introgression during Speciation by
Hybridization. *Mol Biol Evol* **34**: 1947–1959.

Leducq JB, Nielly-Thibault L, Charron G, Eberlein C, Verta JP, Samani P, Sylvester K, Hittinger
CT, Bell G, Landry CR. 2016. Speciation driven by hybridization and chromosomal plasticity
in a wild yeast. *Nature Microbiology* **1**: 1–58.

Lee H-Y, Chou J-Y, Cheong L, Chang N-H, Yang S-Y, Leu J-Y. 2008. Incompatibility of Nuclear
and Mitochondrial Genomes Causes Hybrid Sterility between Two Yeast Species. *Cell* **135**:
1065–1073.

Li H. 2018a. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**: 3094–
3100.

Li H. 2018b. Seqtk: a fast and lightweight tool for processing FASTA or FASTQ sequences.
https://github.com/lh3/seqtk.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R.
2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Li XC, Peris D, Hittinger CT, Sia EA, Fay JC. 2019. Mitochondria-encoded genes contribute to
evolution of heat and cold tolerance in yeast. *Sci Adv* **5**: eaav1848.

Loman NJ, Quick J, Simpson JT. 2015. A complete bacterial genome assembled de novo using
only nanopore sequencing data. *Nat Methods* **12**: 733–735.

Lynch M, Ackerman MS, Gout J-F, Long H, Sung W, Thomas WK, Foster PL. 2016. Genetic
drift, selection and the evolution of the mutation rate. *Nat Rev Genet* **17**: 704–714.

Lynch M, Sung W, Morris K, Coffey N, Landry CR, Dopman EB, Dickinson WJ, Okamoto K,
Kulkarni S, Hartl DL, et al. 2008. A genome-wide view of the spectrum of spontaneous
mutations in yeast. *Proc Natl Acad Sci U S A* **105**: 9272–9277.

Ma H, O'Farrell PH. 2016. Selfish drive can trump function when animal mitochondrial genomes
compete. *Nat Genet* **48**: 798–802.

Marsit S, Hénault M, Charron G, Fijarczyk A, Landry CR. 2021. The neutral rate of whole-
genome duplication varies among yeast species and their hybrids. *Nat Commun* **12**: 3126.

Müller M, Mentel M, van Hellemond JJ, Henze K, Woehle C, Gould SB, Yu R-Y, van der Giezen
M, Tielens AGM, Martin WF. 2012. Biochemistry and evolution of anaerobic energy
metabolism in eukaryotes. *Microbiol Mol Biol Rev* **76**: 444–495.

Orr HA. 1995. The population genetics of speciation: The evolution of hybrid incompatibilities.
Genetics **139**: 1805–1813.

Paliwal S, Fiumera AC, Fiumera HL. 2014. Mitochondrial-Nuclear Epistasis Contributes to
Phenotypic Variation and Coadaptation in Natural Isolates of Saccharomyces cerevisiae.
Genetics **198**: 1251–1265.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics* **35**: 526–528.

Parakatselaki M-E, Ladoukakis ED. 2021. mtDNA Heteroplasmy: Origin, Detection,
Significance, and Evolutionary Consequences. *Life* **11**.
http://dx.doi.org/10.3390/life11070633.

Payne BAI, Wilson IJ, Yu-Wai-Man P, Coxhead J, Deehan D, Horvath R, Taylor RW, Samuels
DC, Santibanez-Koref M, Chinnery PF. 2013. Universal heteroplasmy of human
mitochondrial DNA. *Hum Mol Genet* **22**: 384–390.

Peris D, Arias A, Orlić S, Belloch C, Pérez-Través L, Querol A, Barrio E. 2017. Mitochondrial
introgression suggests extensive ancestral hybridization events among Saccharomyces
species. *Mol Phylogenet Evol* **108**: 49–60.

Piccinini G, Iannello M, Puccio G, Plazzi F, Havird JC, Ghiselli F. 2021. Mitonuclear
Coevolution, but not Nuclear Compensation, Drives Evolution of OXPHOS Complexes in
Bivalves. *Mol Biol Evol* **38**: 2597–2614.

Procházka E, Franko F, Poláková S, Sulo P. 2012. A complete sequence of Saccharomyces
paradoxus mitochondrial genome that restores the respiration in S. cerevisiae. *FEMS Yeast
Res* **12**: 819–830.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841–842.

R Core Team. 2016. R: A Language and Environment for Statistical Computing. https://www.R-
project.org/.

Rice DW, Alverson AJ, Richardson AO, Young GJ, Sanchez-Puerta MV, Munzinger J, Barry K,
Boore JL, Zhang Y, dePamphilis CW, et al. 2013. Horizontal transfer of entire genomes via
mitochondrial fusion in the angiosperm Amborella. *Science* **342**: 1468–1473.

Rokas A, Ladoukakis E, Zouros E. 2003. Animal mitochondrial DNA recombination revisited.
Trends Ecol Evol **18**: 411–417.

Ruan J, Li H. 2019. Fast and accurate long-read assembly with wtdbg2. *Nat Methods*.
http://dx.doi.org/10.1038/s41592-019-0669-3.

Seah B. 2016. *kbseah/mitonotate: Mitonotate v1.0*. https://zenodo.org/record/159532.

Solieri L. 2010. Mitochondrial inheritance in budding yeasts: towards an integrated
understanding. *Trends Microbiol* **18**: 521–530.

Stewart JB, Chinnery PF. 2021. Extreme heterogeneity of human mitochondrial DNA from
organelles to populations. *Nat Rev Genet* **22**: 106–118.

Stewart JB, Chinnery PF. 2015. The dynamics of mitochondrial DNA heteroplasmy: implications
for human health and disease. *Nat Rev Genet* **16**: 530–542.

Sulo P, Szabóová D, Bielik P, Poláková S, Šoltys K, Jatzová K, Szemes T. 2017. The
evolutionary history of Saccharomyces species inferred from completed mitochondrial
genomes and revision in the "yeast mitochondrial genetic code." *DNA Res* **24**: 571–583.

Taylor DR, Zeyl C, Cooke E. 2002. Conflicting levels of selection in the accumulation of
mitochondrial defects in Saccharomyces cerevisiae. *Proc Natl Acad Sci U S A* **99**: 3690–
3694.

Tsai IJ, Bensasson D, Burt A, Koufopanou V. 2008. Population genomics of the wild yeast
Saccharomyces paradoxus: Quantifying the life cycle. *Proc Natl Acad Sci U S A* **105**: 4957–
4962.

Tsaousis AD, Martin DP, Ladoukakis ED, Posada D, Zouros E. 2005. Widespread
recombination in published animal mtDNA sequences. *Mol Biol Evol* **22**: 925–933.

Van Rossum G, Drake FL. 2009. *Python 3 Reference Manual: (Python Documentation Manual
Part 2)*. CreateSpace Independent Publishing Platform.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E,
Peterson P, Weckesser W, Bright J, et al. 2020. SciPy 1.0: fundamental algorithms for
scientific computing in Python. *Nat Methods* **17**: 261–272.

Vowinckel J, Hartl J, Marx H, Kerick M, Runggatscher K, Keller MA, Mülleder M, Day J, Weber
M, Rinnerthaler M, et al. 2021. The metabolic growth limitations of petite cells lacking the
mitochondrial genome. *Nat Metab* **3**: 1521–1535.

Wagih O, Parts L. 2014. Gitter: A robust and accurate method for quantification of colony sizes
from plate images. *G3: Genes, Genomes, Genetics* **4**: 547–552.

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,
Wortman J, Young SK, et al. 2014. Pilon: An Integrated Tool for Comprehensive Microbial
Variant Detection and Genome Assembly Improvement ed. J. Wang. *PLoS One* **9**:
e112963.

Wallace DC. 2015. Mitochondrial DNA variation in human radiation and disease. *Cell* **163**: 33–
38.

Wang Z, Wilson A, Xu J. 2015. Mitochondrial DNA inheritance in the human fungal pathogen
Cryptococcus gattii. *Fungal Genet Biol* **75**: 1–10.

Wilson AJ, Xu J. 2012. Mitochondrial inheritance: Diverse patterns and mechanisms with an
emphasis on fungi. *Mycology* **3**: 158–166.

Wolters JF, Charron G, Gaspary A, Landry CR, Fiumera AC, Fiumera HL. 2018. Mitochondrial
recombination reveals mito–mito epistasis in yeast. *Genetics* **209**: 307–319.

Wu B, Buljic A, Hao W. 2015. Extensive horizontal transfer and homologous recombination
generate highly chimeric mitochondrial genomes in yeast. *Mol Biol Evol* **32**: 2559–2570.

Wu B, Hao W. 2015. A Dynamic Mobile DNA Family in the Yeast Mitochondrial Genome. *G3* **5**:
1273–1282.

Xia W, Nielly-Thibault L, Charron G, Landry CR, Kasimer D, Anderson JB, Kohn LM. 2017.

Population genomics reveals structure at the individual, host-tree scale and persistence of
genotypic variants of the undomesticated yeast Saccharomyces paradoxus in a natural
woodland. *Mol Ecol* **26**: 995–1007.

Yue JX, Li J, Aigrain L, Hallin J, Persson K, Oliver K, Bergström A, Coupland P, Warringer J,
Lagomarsino MC, et al. 2017. Contrasting evolutionary genome dynamics between
domesticated and wild yeasts. *Nat Genet* **49**: 913–924.