Transcription-induced active forces suppress chromatin motion

Sucheol Shin[a], **Guang Shi**[a,b], **Hyun Woo Cho**[c], **and D. Thirumalai**[a,d]

[a]Department of Chemistry, The University of Texas at Austin, Texas 78712, USA; [b]Department of Materials Science, University of Illinois, Urbana, Illinois 61801, USA; [c]Department of Fine Chemistry and Center for Functional Biomaterials, Seoul National University of Science and Technology, Seoul 01811, Republic of Korea; [d]Department of physics, The University of Texas at Austin, Texas 78712, USA

This manuscript was compiled on August 23, 2023

The organization of interphase chromosomes in a number of species is starting to emerge thanks to advances in a variety of experimental techniques. However, much less is known about the dynamics, especially in the functional states of chromatin. Recent experiments have shown that the mobility of individual loci in human interphase chromosome decreases during transcription, and increases upon inhibiting transcription. This is a counter-intuitive finding because it is thought that the active mechanical force (F) on the order of ten pico-newtons, generated by RNA polymerase II (RNAPII) that is presumably transmitted to the gene-rich region of the chromatin, would render it more open, thus enhancing the mobility. Inspired by these observations, we developed a minimal active copolymer model for interphase chromosomes to investigate how F affects the dynamical properties of chromatin. The movements of the loci in the gene-rich region are suppressed in an intermediate range of F, and are enhanced at small and large F values. In the intermediate F, the bond length between consecutive loci increases, becoming commensurate with the distance at the minimum of the attractive interaction between non-bonded loci. This results in a transient disorder-to-order transition, leading to the decreased mobility during transcription. Strikingly, the F-dependent change in the locus dynamics preserves the organization of the chromosome at $F = 0$. Transient ordering of the loci, which is not found in the polymers with random epigenetic profiles, in the gene-rich region might be a plausible mechanism for nucleating a dynamic network involving transcription factors, RNAPII, and chromatin.

Advances in experimental techniques (1, 2) have elucidated the organizational details of chromosomes, thus deepening our understandings of how gene regulation is connected to chromatin structure (3). Relatively less is known about the dynamics of the densely packed interphase chromosomes in the cell nucleus. Experimental and theoretical studies have shown that the locus dynamics is heterogeneous, exhibiting sub-diffusive behavior (4–8), which is consistent with models that predict glass-like dynamics (9–11). However, it is challenging to understand the dynamic nature of chromosomes that governs the complex subnuclear processes, such as gene transcription.

The link between transcriptional activity and changes in chromosome dynamics is important in understanding the dynamics of chromosomes in distinct cell types and states (12, 13). It is reasonable to expect that transcription of an active gene-rich region could make it more expanded and dynamic (14, 15). However, active RNA polymerase (RNAP) II suppressed the movement of individual loci in chromatin (13, 16). Let us first summarize the key experimental results of Ref. 13, which inspired our study: (1) By imaging the motion of individual nucleosomes (monomers in the chromosomes) in live human cells, it was shown that the mean square displacements (MSDs) of the nucleosomes during active transcription are constrained (see Fig. S1 in SI Appendix). (2) When the cells were treated with α-amanitin (α-AM), a transcription inhibitor which selectively blocks the translocation of RNAPII (17, 18), the mobilities of the nucleosomes were enhanced. This finding is counter-intuitive because the translocation generates mechanical forces (19, 20) that should render the chromatin more dynamic. (3) The enhanced motion was restricted only to gene-rich loci (euchromatin) that are predominantly localized in the cell interior (21, 22) whereas the dynamics of gene-poor heterochromatin, found in the periphery (21–23), is unaffected by transcription. Based on these observations, it was hypothesized that RNAPs and other protein complexes, facilitating transcription, transiently stabilize chromatin by forming dynamic clusters (24–28), whose structural characteristics are unknown. This hypothesis has been challenged because the inhibition mainly leads to stalling of RNAPs bound to chromatin (17, 18). Moreover, transcriptional inhibition does not significantly alter the higher-order structures of chromosomes (28, 29). These observations raise the question: Is there a physical explanation for the increased chromatin dynamics upon inhibition of transcription and a decrease during transcription? We provide a plausible answer to this question by extensive simulations of a simple minimal active copolymer model, which captures the organization of chromosomes in the absence of active forces.

Using the experimental results as a backdrop, we theorize that RNAPII exerts active force (19, 20) in a vectorial manner on the active loci. We then examine the effects of active force on the organization and dynamics of chromosome using the

Significance Statement

In order to explain a physically counter-intuitive experimental finding that chromatin mobility is reduced during transcription, we introduced a polymer model for interphase chromosome that accounts for RNA polymerase (RNAP) induced active force. The simulations show that, over a range of active forces, the mobility of the gene-rich loci is suppressed. Outside this range, chromosomes are compact and exhibit glass-like dynamics. Our study, which accounts for the experimental observations, leads to a novel and testable mechanism of how transcription could shape the coexistence of fluid- and solid-like properties within chromosomes.

S.S. and D.T. designed research; S.S., G.S., and D.T. performed research; S.S., G.S, H.W.C, and D.T. analyzed data, and S.S., G.S, and D.T. wrote the paper

The authors declare no conflict of interest.

Fig. 1. Schematic depiction of the model. (A) Chromosome is modeled as a copolymer chain consisting of active/inactive (green/purple-colored) loci. Specific locus pairs are connected by loop mediators (orange color). We apply active forces to all the gene-rich A-type loci in the form of an extensile force dipole (red arrows) along each bond vector between the A-type loci. The green and purple boxes illustrate the microscopic origin of the dipolar extensile active forces. Compared to the inactive B-type loci (bottom), the A-type loci include the nucleosomes in a more dynamic state along with the actively transcribing RNAPII, such that their effective excluded volumes increase modestly, as represented by the light-red shade. The enhanced volume exclusion gives rise to repulsion between the consecutive A-type loci and the increase of the bond distance. (B) Illustration for how active forces are imposed on the i^{th} locus and its bonded loci, where the solid arrow is the net force from the force dipoles (dashed arrows).

Model

Chromosome Copolymer Model. We model an interphase chromosome as a flexible self-avoiding copolymer (Fig. 1), whose potential energy function is given in Methods. Non-adjacent pairs of loci are subject to favorable interactions, modeled by the Lennard-Jones (LJ) potential (Eq. 6), depending on the locus type. The relative strengths of interactions between the loci is constrained by the Flory-Huggins theory (31, 32), which ensures microphase separation between the A and B loci—an important organization principle of interphase chromosomes. Certain locus pairs, separated by varying genomic distances, are linked to each other, representing the chromatin loops mediated by CCCTC-binding factors (CTCFs) (33). In this study, we simulate a 4.8-Mb segment of human chromosome 5 (Chr5: 145.87–150.67 Mb) using $N = 4,000$ loci, that is, 1.2 kb ~ 6 nucleosomes per locus. The A- to B-type ratio is $N_{\text{A}}/N_{\text{B}} = 982/3018 \approx 1/3$. See Methods for the details about

the CCM polymer. In the absence of active forces, the CCM faithfully captures the experimental results, showing microphase separation between euchromatin (A-type loci) and heterochromatin (B-type loci) on large length scale and formation of substructures on a smaller length scale in interphase chromosomes. The CCM leads to a condensed polymer, exhibiting glass-like dynamics in the absence of activity, in agreement with experiments (4, 5, 10, 30). Brownian dynamics simulations of a single chromosome in the presence of active forces, produces the following key results: (i) In accord with experiments (13), the dynamics of the active loci, measured using the MSD, is suppressed upon application of the active force. Despite the use of minimal model, our simulations qualitatively capture the increase in the MSD for the transcriptionally inactive case relative to the active case. (ii) The decrease in the mobility of the A loci occurs only over a finite range of activity level. Surprisingly, in this range the segregated A loci undergo a transient disorder-to-order transition whereas the B loci remain fluid-like. (iii) Chromosome in which the epigenetic (A/B) profile is random cannot capture the experimental observations, implying that the sequence inherent in the chromosomes plays a vital role.

assignment of the locus type and the loop anchors in the CCM polymer.

In our model, the loop topology does not change with time, unlike polymer models, which examined the consequences of dynamic extrusion of loops on interphase and mitotic structures (34–36). We simulate a given chromosome region up to ~10 seconds (see Methods for details) that is much shorter than the lifetime of the CTCF loops (15–30 mins) (37, 38). Furthermore, the loop extrusion rate is ~0.5–2 kb/s (39, 40), which also would not significantly alter the topology of loops on the simulation time scales. Therefore, the assumption that loops are static is reasonable. We discuss, in a later section, the potential effects of the removal of loops along with the implications (see Discussion)

Active forces in transcriptional activity. Previous theoretical studies (41–49) have considered different forms of active forces on homopolymers and melts of polymer rings. Potential connection to chromatin has been proposed using polymer models in the presence of active forces (6, 50–55). However, these have not accounted for the counter-intuitive finding from the experiment of transcription inhibition (13). Here we applied active forces on the CCM polymer chain to mimic the force generated by the actively transcribing RNAP during transcription elongation (19, 20).

In our model, active forces act along each bond vector between the A-type loci in an extensile manner, ensuring momentum conservation (see Fig. 1). The force, associated with the translocation of actively transcribing RNAP (19, 20), should result in the increased volume exclusion between the bonded A loci, as shown in the green box in Fig. 1A. We reason that the dipolar extensile active force, with local momentum conservation, models the effective repulsion between the active loci in the coarse-grained level (see SI Appendix, Sec. 1 for more details about the biological relevance of the force). Along the A-A bond vector, $\mathbf{b}_i = \mathbf{r}_{i+1} - \mathbf{r}_i$ ($\mathbf{r}_i$ is the position of the i^{th} locus), force, $\delta\mathbf{f}_{i+1}^{(\text{f})} = f_0\hat{\mathbf{b}}_i$, acts on the $(i + 1)^{\text{th}}$ locus in the forward direction (f_0 is the force magnitude and $\hat{\mathbf{b}}_i = \mathbf{b}_i/|\mathbf{b}_i|$), and $\delta\mathbf{f}_i^{(\text{b})} = -\delta\mathbf{f}_{i+1}^{(\text{f})}$ is exerted on the i^{th} locus in the opposite direction (Fig. 1B). If the $(i − 1)^{\text{th}}$ locus is also

A-type, another pair of active force is exerted along the bond vector, $\mathbf{b}_{i-1} = \mathbf{r}_i - \mathbf{r}_{i-1}$, and thus the net active force on the i^{th} locus is given by $\mathbf{f}_i = \delta\mathbf{f}_i^{(f)} + \delta\mathbf{f}_i^{(b)} = f_0(\hat{\mathbf{b}}_{i-1} - \hat{\mathbf{b}}_i)$ (solid arrow in Fig. 1B). We simulate the CCM chain dynamics by numerically solving the overdamped Langevin equation, given in Eq. 10, which includes the active forces (see Methods for simulation details). We use the dimensionless parameter, $F \equiv f_0\sigma/k_BT$, as a measure of the force magnitude, where σ is the diameter of a single locus, k_B is the Boltzmann constant, and T is the temperature.

In the simulations, we apply active forces to all the A-type loci at each time step, which can be justified as follows. First, consider the relevant length scale. Since the average human gene size is ~27 kb (56), our choice of the resolution is sufficiently small to distinguish between the gene and the intergenic regions which are assigned A- and B-types, respectively. On each gene body, there are multiple active RNAPII. The experimentally inferred density of RNAPII is 1.0–1.3 RNAPII per kb (57). Hence, it is a reasonable approximation that each individual A-type locus is occupied by a single active RNAPII. The tension force locally generated by the RNAPII would propagate beyond the gene-transcribing region on a time scale that is faster than our simulation time unit (see SI Appendix, Sec. 1 for details). As a result, the force is felt throughout the A-type loci. Single-molecule measurements (20) showed that the tension force on DNA was maintained even if the RNAP pauses the elongation step intermittently (~1–10 s). This observation provides the basis for applying the active forces on the A-type loci throughout the simulations.

Results

MSD comparison. We calculated the MSDs separately for euchromatin and heterochromatin loci at lag time t,

$$\langle\Delta r_\mu^2(t)\rangle = \frac{1}{N_\mu}\left\langle\sum_{i=1}^N \delta_{\nu(i)\mu}|\mathbf{r}_i(t) - \mathbf{r}_i(0)|^2\right\rangle, \qquad [1]$$

where $\langle\cdots\rangle$ is the ensemble average, $\nu(i)$ is the type of the i^{th} locus, and $\delta_{\nu(i)\mu}$ is the Kronecker delta ($\delta_{xy} = 1$ if $x = y$, or 0 else) that picks out the loci of specific type μ (either A or B). Fig. 2A shows that $\langle\Delta r_A^2(t)\rangle$ is smaller at $F = 80$ compared to $F = 0$. This result is qualitatively similar to the nucleosome MSDs measured from the interior section of the cell nucleus treated with the transcription inhibitor α-AM (13). $\langle\Delta r_B^2(t)\rangle$ at $F = 80$ is also smaller than at $F = 0$, but the difference is marginal compared to $\langle\Delta r_A^2(t)\rangle$, which is consistent with the experimental results for the nuclear periphery (Fig. S2A). In Fig. 2A, the magnitude of $\langle\Delta r_A^2(t)\rangle$ is smaller in simulations than in experiment, where the difference in magnitude depends on the conversion of simulation length/time to real units (see Methods for the details). The level of agreement with experiments is striking, considering that we used a minimal model with no parameters to describe a very complicated phenomenon. We could probably have obtained better agreement with experiments by tweaking the parameters in the model, which we think is not necessary for the purposes of this study.

In Fig. 2A, each MSD plot as a function of time is shown with scaling behavior, $\langle\Delta r_A^2(t)\rangle \sim t^\alpha$. The scaling exponent, α, provides the link between chromatin structure and the mobility. It was established previously (6) that, $\alpha = \frac{2\nu}{(2\nu+1)}$

Fig. 2. Transcription-induced active forces reduce the mobilities of euchromatin loci. (A) $\langle\Delta r_A^2(t)\rangle$ from simulations with $F = 0$ and $F = 80$ (solid lines), compared with the euchromatin MSD from the experiment that inhibits transcription using α-AM (13) (circles). The dotted lines are the fits to the experimental data. The error bars for the simulation data are obtained from five independent trajectories. (B) Bar graphs comparing the increase in $\langle\Delta r_A^2(t)\rangle$ ($\langle\Delta r_B^2(t)\rangle$) shown in panel A (Fig. S2A) between experiment and simulation results. (C) MSDs for the A and B loci at $t = 0.5$ s as a function of F. The dotted lines are a guide to the eye. (D) Radial distributions of the A-locus displacement at $t = 10\tau_B$ (dashed) and $t = 1,000\tau_B$ (solid), compared between $F = 0$ (gray) and $F = 80$ (dark-red). The inset shows the 2-D projection of the trajectory of an active locus for $10^4\tau_B$ at $F = 0$ and $F = 80$.

where ν, the Flory exponent related to the radius of gyration ($R_g \sim N^\nu$) of the polymer chain. For an ideal Rouse polymer, $\nu = 0.5$, and for a polymer in a good solvent, which takes the excluded volume between the monomers into account, $\nu \approx 0.6$ (58). The fit to the experimental data (gray dotted line in Fig. 2A) when transcription is inhibited yields $\alpha \approx 0.57$, which would be consistent with the prediction for the SAW behavior. In all other cases, α is less than 0.5 (the value for the Rouse polymer) both in experiments and simulations, which is indicative of restricted motility. The scaling exponent in the experimental MSD increases from $\alpha = 0.48\,(\pm0.05)$ to $\alpha = 0.57\,(\pm0.07)$ upon inhibiting transcription. We estimated the errors using 10,000 samples reconstructed from the normal distribution for each data point, following the procedure described elsewhere (59). Our simulations capture the change in α such that $\Delta\alpha$(active $\to$ passive) $= 0.05$ (versus 0.09 from the experiments). In contrast, recent experiments for chromatin dynamics in different contexts, such as loop formation and response to mechanical perturbation, reported that the MSD exponents are similar to the Rouse (37, 60) or SAW (38) polymer, i.e., $\alpha \gtrsim 0.5$. These results imply fundamentally different physical nature in transcriptionally active regions, which constrains the polymer dynamics to $\alpha < 0.5$.

In Fig. 2B, we compare the transcription-inhibited increase ($\Delta\overline{\text{MSD}}$) in the MSD, between experiment and simulations (see Eqs. S1–S2 in SI Appendix). We use $F = 80$, which shows the smallest MSD (see Fig. 2C), as the control. The value of f_0 for $F = 80$ is in the range, $f_0 \approx 3$–16 pN (see Methods), which accords well with the typical forces exerted by RNAP (19). Comparison between $\Delta\overline{\text{MSD}}$ for the A loci (simulation) and the interior measurements (experiment) is less precise

than between the B loci and the periphery. The difference may arise because the interior measurements could include the heterochromatin contribution to some extent, whereas the periphery measurements exclude the euchromatin. Nevertheless, $\Delta\overline{\text{MSD}}$ for all the loci is in better agreement with experiment (Fig. S2C). The reasonable agreement between simulations and experiment is surprising because it is obtained *without adjusting any parameter to fit the data*. Although comparisons in Fig. 2B are made with $F = 80$, we obtain qualitatively similar results for F in the range, $60 \leq F \leq 90$ (Fig. S3).

The simulated MSD, at a given lag time, changes non-monotonically as F changes. Remarkably, the change is confined to the A loci (Figs. 2C and S2D–S2E); $\langle \Delta r_{\text{A}}^2(0.5\,\text{s}) \rangle$ increases modestly as F increases from zero to $F \lesssim 30$, and decreases when F exceeds thirty. There is an abrupt reduction at $F \approx 50$. In the range, $50 \lesssim F \lesssim 80$, $\langle \Delta r_{\text{A}}^2(0.5\,\text{s}) \rangle$ continues to decrease before an increase at higher F values. In contrast, the MSD of the B loci does not change significantly with F (Fig. 2C). The non-monotonic trend in $\langle \Delta r_{\text{A}}^2(0.5\,\text{s}) \rangle$ upon changing F is reminiscent of a *re-entrant* phase behavior in which a system transitions from an initial phase to a different phase and back to the initial phase as a parameter (F in our case) is varied. Such a behavior occurs in a broad range of different systems (61–64). We present additional analyses in the next section to confirm the *dynamic re-entrance*.

To obtain microscopic insights into the simulated MSDs, we calculated the van Hove function, which is simply the distribution of locus displacement at a given lag time,

$$G_s^\mu(\mathbf{r}, t) = \frac{1}{N_\mu} \left\langle \sum_{i=1}^N \delta_{\nu(i)\mu} \delta(\mathbf{r} + \mathbf{r}_i(0) - \mathbf{r}_i(t)) \right\rangle \quad . \quad [2]$$

Figure 2D compares the radial distribution for the displacement of A-type loci, $4\pi r^2 G_s^{\text{A}}(r, t)$, between $F = 0$ and $F = 80$ at $t = 10\tau_B \approx 0.007\,\text{s}$ and $1{,}000\tau_B \approx 0.7\,\text{s}$ (see Methods for the unit conversion). At short lag times, both the distributions for $F = 0$ and $F = 80$ show a peak at $r < 0.5\sigma$, which means that a given locus is unlikely to diffuse far from the neighboring loci in a condensed microenvironment. At long lag times, the distribution for $F = 0$ becomes more populated at $r \geq \sigma$, as the loci escape from the regime caged by the neighbors. The remarkably broad and flat curve for $r^2 G_s^{\text{A}}(r, t = 10^3\tau_B)$ at $F = 0$, resembling a uniform distribution, signifies the heterogeneous dynamics of the A-type loci, which differs from the Gaussian distribution expected for the Rouse polymer. The A-type loci at $F = 80$ do not diffuse as much at $F = 0$ even at long lag times (Fig. 2D), and their displacements are largely within the length scale of σ. In contrast, there is no significant difference in $G_s^{\text{B}}(r, t)$ between $F = 0$ and $F = 80$ (Fig. S2F). Notably, the second peak of $G_s^{\text{A}}(r, t)$ at $F = 80$ hints at the solid-like lattice (65).

Dynamic re-entrant behavior. Here, we further characterize the dynamics of the CCM polymer chain in the presence of RNAPII-induced active forces. The purpose is to investigate how the re-entrant behavior emerges in the CCM polymer whose dynamics is glass-like in the absence of the activity (10). To probe how the glass-like behavior of the polymer chain (6, 9, 10) is affected by the active forces, we calculated the

Fig. 3. Re-entrance in the relaxation time and dynamic heterogeneity is observed upon increasing the activity. (A) Plot of $\langle F_s(k_{\max}, t) \rangle$ for different F. The dashed line is a stretched exponential fit for $F = 0$. (B) τ_α (blue) and β (orange) of $\langle F_s(k_{\max}, t) \rangle$ as a function of F. The dotted lines are a guide to the eye. (C) Plot of $\chi_4(t)$ [Eq. (4)] for different F. (D) The maximum value of $\chi_4(t)$ as a function of F.

self-intermediate scattering function,

$$F_s(|\mathbf{k}|, t) = \frac{1}{N} \sum_{j=1}^N e^{i\mathbf{k} \cdot [\mathbf{r}_j(t) - \mathbf{r}_j(0)]} \quad , \quad [3]$$

where $\mathbf{k}$ is the wave vector. The scattering function, the Fourier transform of the van Hove function (Eq. 2), quantifies the extent of the loci displacements over time t relative to the length scale set by the wave vector ($\sim 2\pi/|\mathbf{k}|$). We computed the ensemble-averaged $\langle F_s(k_{\max}, t) \rangle$ with $k_{\max} = 2\pi/r_{\text{LJ}}^* \approx 5.60\sigma^{-1}$, where $r_{\text{LJ}}^* = 2^{1/6}\sigma \approx 1.12\sigma$ is the distance at the minimum of the non-bonding (LJ) potential (see Methods), which the nearest neighbors are likely to be located at (see the radial distribution function in Fig. 4 below). The decay of $\langle F_s(k_{\max}, t) \rangle$ indicates the structural relaxation on the length scale $\gtrsim \sigma$.

Time-dependent variations in $\langle F_s(k_{\max}, t) \rangle$ (Fig. 3A) show stretched exponential behavior ($e^{-(t/\tau_a)^\beta}$; $\beta < 1/3$ at all F values), which is one signature of glass-like dynamics. Note that the decay is even slower if F is increased, which is in agreement with the results shown in Figs. 2 and S2. The relaxation time, τ_α, calculated using $\langle F_s(k_{\max}, \tau_\alpha) \rangle = 0.2$, shows that the relaxation is slowest at $F \approx 80$ (Fig. 3B), which occurs after the dynamical transition in $\langle \Delta r_{\text{A}}^2(0.5\,\text{s}) \rangle$ at $F \approx 50$ and before $\langle \Delta r_{\text{A}}^2(0.5\,\text{s}) \rangle$ increases beyond $F = 100$ (Fig. 2C). Similarly, when the tails of $\langle F_s(k_{\max}, t) \rangle$ were fit with $e^{-(t/\tau_a)^\beta}$, the exponent β also exhibits the analogous trend (Fig. 3B); that is, as τ_α increases, β decreases, indicating the enhancement in the extent of the glass-like behavior.

Dynamic heterogeneity, another hallmark of glass-like dynamics (66, 67), was calculated using the fourth-order susceptibility (68), which is defined as the mean squared fluctuations in the scattering function,

$$\chi_4(t) = N \left[\langle F_s(k_{\max}, t)^2 \rangle - \langle F_s(k_{\max}, t) \rangle^2 \right] \quad . \quad [4]$$

We find that $\chi_4(t)$, at all F values, has a broad peak spanning a wide range of times, reflecting the heterogeneous motion of

the loci (Fig. 3C). The peak height, χ_4^*, increases till $F \approx 50$ and subsequently decreases (Fig. 3D). When F exceeds 100, χ_4^* decreases precipitously. Our results suggest that there are two transitions: one at $F \approx 50$ where the dynamics slows down and the other, which is a reentrant transition beyond $F = 100$, signaled by an enhancement in the mobilities of the A-type loci. Although the system is finite, these transitions are discernible.

Like the MSD, when $\langle F_s(k_{\max}, t) \rangle$ and $\chi_4(t)$ was decomposed into the contributions from A and B loci (Eq. S3 and S4), we find that the decrease in the dynamics and the enhanced heterogeneity are driven by the active loci (Fig. S4). These observations, including the non-monotonicity in τ_α and β that exhibit a dynamic reentrant behavior, prompted us to examine if the dynamical changes in the A-type loci are accompanied by any structural alterations.

Fig. 4. F-induced structural transition. (A) RDF for A-A pairs at different F (solid lines; see the legend in panel B), where $g(r)$ for a FCC crystal is shown with the dotted line (scaled arbitrarily). The inset shows $g_{AA}(r)$ for the quenched polymer with $F = 80$. (B) Distributions of the BOO parameter, q_{12}, for A loci as a function of F. The dashed line is for the quenched A loci at $F = 80$. (C) Height of the dominant peak in $g_{AA}(r)$ (blue) and $\langle q_{12} \rangle_A$ (orange) as a function of F. Simulation snapshots for $F = 0$ and $F = 80$ are in the gray and yellow boxes, respectively.

Transient disorder-to-order transition induced by active forces. The radial distribution function (RDF) for A-A locus pairs, $g_{AA}(r)$, with signature of a dense fluid, shows no visible change for $F \lesssim 30$ (Fig. 4A). In contrast, the height of the primary peak, g_{AA}^*, increases sharply beyond $F = 30$ (Fig. 4C). Remarkably, $g_{AA}(r)$ for $F = 80$ exhibits secondary peaks that are characteristics of the FCC (face-centered cubic) crystal phase (Fig. 4A, arrows). Upon further increase in F, these peaks disappear (Fig. S5A) and g_{AA}^* reverts to the level of the passive case (Fig. 4C). In other words, the active forces, which preserve fluid-like disorder in the A loci at low F values, induce a structural transition to solid-like order in the intermediate range of F values, which is followed by reentrance to a fluid-like behavior at higher F values. In contrast, $g_{BB}(r)$ exhibits dense fluid-like behavior at all F values (Fig. S5B). We confirm that the FCC lattice is the minimum energy configuration by determining the *inherent structure* (69, 70) for the A loci at $F = 80$ by quenching the active polymer to a low temperature (see Methods) (Fig. 4A, inset). In Fig. S5C,

$g_{AA}(r)$ at $F = 0$ reflects the structure of a dense fluid upon the quench whereas $g_{AA}(r)$ at $F = 80$ is quantitatively close to that for a perfect FCC crystal. Quenching does not alter the structure of the B loci at $F = 80$ or $g_{AA}(r)$ at $F = 0$ (Figs. S5C-S5D).

To assess local order, we calculated the bond-orientational order (BOO) parameter for 12-fold rotational symmetry, q_{12} (Eqs. 12–13) (71, 72). For a perfect FCC crystal, $q_{12} \approx 0.6$ (73). The distribution for A loci, $P_A(q_{12})$ (Eq. 14), is centered around $q_{12} = 0.3$ at $F = 0$ (Fig. 4B), representing a disordered liquid state (Fig. 4C, gray box). As F increases, the distribution shifts towards the right, especially in the $50 \leq F \leq 80$ range. The increase of $\langle q_{12} \rangle_A$ (Eq. 15) indicates a transition to a FCC-like ordered state that is visible in the simulations (Fig. 4C, yellow box). $P_A(q_{12})$ at $F = 80$ is broad, whereas the inherent structure gives a narrower distribution peaked near $q_{12} = 0.6$ (Fig. 4B, dashed line), which shows that the ordered solid-like state coexists with the disordered fluid-like state within thermal fluctuations. The maximum in $P_A(q_{12})$ shifts to the left for $F > 80$ (Fig. S5E) and $\langle q_{12} \rangle_A$ decreases, suggestive of F-induced reentrant transition. The distribution $P_B(q_{12})$ for the B-type loci is independent of F (Fig. S5F). These results show that FCC-like ordering emerges in $50 \lesssim F \lesssim 100$ range. Outside this range, the RDFs display the characteristics of a dense fluid. In addition, the F-dependent bending angle distributions for three consecutive A loci (Eq. S5) reflect the FCC-like ordering (Fig. S6). The transitions in the A-type loci may be summarized as fluid $\rightarrow$ FCC/fluid $\rightarrow$ fluid, as F changes from 0 to 120.

Since the overall RDF at $F = 80$ does not show the FCC peaks (Fig. S7A), the F-induced order may not be seen directly in the conventional experiments. Moreover, the ordering is transient, as implied in the coexistence of order and disorder. By transient we mean that solid-like order persists only when the polymerase induces active forces during transcription. The time correlation function of the BOO parameter, $C_{q_{12}}^A(t)$ [Eq. (S6)], decays rapidly, at most on the order, $\approx 0.7\,\mathrm{s}$ (Fig. S7B), much shorter than the transcription time scale ($\sim$ minute). More details and the visualization of the transient ordering are given in SI Appendix, Sec. 5 and Movie S1. Remarkably, the transient ordering preserves the large-scale structure of chromosome (see SI Appendix, Sec. 6; Fig. S8), in agreement with the chromosome conformation capture experiments with transcription inhibition (28, 29). Here we emphasize that the solid-like nature coexisting with the liquid is sufficient to explain the experimentally observed suppression of euchromatin motions. The predicted transient ordering could be detected by future high resolution imaging experiments.

Origin of F-induced order. The emergence of solid-like order in the A-type loci is explained using the effective A-A interaction generated by F. We calculated the *effective pair potential* for an A-A bond,

$$u_b^{\mathrm{eff}}(r) = u_b^0(r) - f_0(r - b_0) , \qquad [5]$$

where $u_b^0(r)$ and b_0 are the F-independent bonding potential, and the corresponding equilibrium bond length, respectively. The $f_0(r - b_0)$ term is the work done by the active force to stretch the bond from b_0. The equation of motion for $F \neq 0$ represents dynamics in the *effective equilibrium* under

the potential energy involving $u_b^{eff}(r)$ (Eqs. 16–17). Such effective equilibrium concept was useful for characterizing various physical quantities in active systems (47, 54, 74–77).

Plots of $u_b^{eff}(r)$ in Fig. 5A show that the effective equilibrium bond length, r_{min}, increases as F increases. This prediction is confirmed by the direct measurement of A-A bond distance from the simulations (Figs. S9A–S9B). Note that $r_{min}(F = 80) \approx r_{LJ}^* \approx 1.12\sigma$, where r_{LJ}^* is the distance at the minimum of the LJ potential (Fig. 5B). The F-induced extension of A-A bonds makes r_{min} commensurate with r_{LJ}^*, which is conducive to FCC ordering (78) in the active loci. At $F = 0$, since the bond distance ($b_0 = 0.96\sigma$; see Methods) is smaller than the preferred non-bonding interaction distance, r_{LJ}^*, the lattice-matching configurations cannot be formed given the chain connectivity (Fig. 5C, left). In contrast, at $F = 80$, the condition that $r_{min} \approx r_{LJ}^*$ allows the A loci to be arranged into the FCC lattice, leading to the transition to the ordered state (Fig. 5C, center). Upon increasing the activity further (e.g., $F = 120$), the bond distance becomes too large to be fit with the lattice structure, so the loci re-enter the disordered fluid state (Fig. 5C, right).

Fig. 5. Effective potential energy accounts for the structural transition of the A-type loci. (A) Effective pair potential of a single A-A bond, $u_b^{eff}(r)$ (Eq. 5), as a function of F. The LJ potential, $u_{LJ}(r)$, which is vertically shifted, is shown in blue dashed line for comparison. (B) Distance at the minimum of $u_b^{eff}(r)$ as a function of F, where the dashed line indicates r_{LJ}^* (see the main text). (C) Schematic illustration for how the F-induced bond extension enables FCC ordering in the A-type loci (green circles). The blue shade shows r_{LJ}^* at which non-bonded pairs have the most favorable interactions. The double red arrows indicate the bond extension upon increasing F. (D) Mean of the effective potential energy of A loci (left) and mean fluctuations (right) with respect to F. The arrows indicate the two structural transitions.

We can also describe the ordering behavior using thermodynamic properties based on $u_b^{eff}(r)$. We calculated the mean and variance of the effective potential energy of the A loci, $E_{eff,A}$ (Eq. 18). Fig. 5D shows that $\langle E_{eff,A} \rangle$ decreases smoothly as F changes, without pronounced change in the slope, as might

be expected for a structural transition (78). Nevertheless, $\langle (\delta E_{eff,A})^2 \rangle$ indicates signatures of a transition more dramatically, with peaks at $F = 50$ and $F = 100$ (Fig. 5D, arrows I and II). Thus, both ordering and reentrant fluid behavior coincide with the boundaries of the dynamic transitions noted in Figs. 2B, 3B, and 3D.

Discussion

We introduced a minimal active copolymer model for Chr5 in order to explain the non-intuitive experimental observation that during transcription the mobilities of the loci are suppressed. Despite the simplicity of the polymer model, we reproduce semi-quantitatively the experimental observations. In particular, the MSD exponent α ($\langle \Delta r_\mu^2(t) \rangle \sim t^\alpha$) decreases by 0.05 in the simulations whereas the decrease is 0.09 between the transcription inhibited and active states in the experiments (Fig. 2A). We find this result and the results in Fig. 2 surprising because no parameter in the very simple model was adjusted to obtain the semi-quantitative agreement with experiments (13).

We hasten to add that the model indeed is oversimplified, and is the major limitation of our study. The actual machinery of transcription active (or inhibited) state is extremely complicated involving the interplay of transcription factors, RNAPII, chromatin, and other cofactors. These elements are believed to drive the formation of a hub of connected modules resulting in condensates (24–27). Although the structural features of chromatin in the condensates are unknown, our simulations suggest that for a very short period ($\approx$ 1s) chromatin could adopt solid-like properties when RNAPII exerts an active force ($\approx$ 5–10 pN) during transcription.

Robustness of the conclusions. We performed four tests in order to ensure that the results are robust. (1) Simulations of a segment of chromosome 10, with a larger fraction of active loci, show qualitatively similar behavior (See SI Appendix, Sec. 7; Fig. S10). (2) For a copolymer chain whose A/B sequence is random (See SI Appendix, Sec. 8; Fig. S11), we find that at all F, MSD for the A-type loci increases monotonically (Fig. 6A) in contrast to the non-monotonic behavior found in Chr5 with a given epigenetic (A/B) profile (Fig. 2C). The B-type MSD decreases at high F (Fig. 6A), which is also different from the behavior in Chr5. In addition, in the random sequences the loci exhibit fluid-like behavior at all values of F, and hence cannot account for the experimental observation (Fig. 6B). Thus, F-induced decrease in the motility of the A loci, accompanied by transient ordering, occurs only in copolymer chains exhibiting microphase separation between A and B loci—an intrinsic property of interphase chromosomes (79). In other words, the epigenetic profile determines the motilities of the individual loci (or nucleosomes) during transcription. (3) We performed simulations by removing all the loop anchors (analog of topological constraints (47); see SI Appendix, Sec. 9). Fig. S12 shows that the F-induced dynamical slowdown and ordering is not significantly affected by the loop anchors. The modest increase in the MSD upon removal of loops, at $F = 0$, is qualitatively consistent with the experimental measurements of increased chromatin mobility when cohesin loading factors were deleted (12). The experimental results suggest that a reduction in transcriptional activity may be induced by loss of loops, which is assumed to be a consequence of depletion

Fig. 6. A copolymer chain with the same length and the fraction of active loci as Chr5 but randomly shuffled epigenetic profile does not capture the experimental observations. (A) Ratio of the MSD at $t_0 = 100\tau_B$ for the active random copolymer to the passive case as a function of F, shown in black color. The data for Chr5 is shown in blue color. The same quantities computed for the A and B loci of the random copolymer are also plotted in green and purple color, respectively. The dotted lines are a guide to the eye. The simulation snapshot shows that the copolymer chain is in a mixed state without discernible microphase separation. (B) Probability distributions of q_{12} for the A loci at different activity levels, where the dashed and dotted lines are the distributions for Chr5 with $F = 0$ and $F = 80$, respectively.

of cohesin (see "Connection to other experiments" below for more discussion). (4) We applied the active forces in a non-continuous manner over time (see SI Appendix, Sec. 10), since the transcription elongation process shows intermittent pauses (80–82). Although the tension force applied on DNA is maintained even during transcriptional pauses (20), how the force is generated and transmitted by the RNAP elongation complex onto DNA is unknown. To provide insights into this issue, we explore theoretically the potential effects of fluctuations in the applied forces on chromatin dynamics. The similar extent of transient ordering and dynamical suppression is induced by the stochastic active forces by the same mechanism (Fig. S14). These tests show that our conclusions are robust, and the model might serve as a minimal description of the effect of transcription-induced active forces on real chromosomes.

Emergence of solid-like ordering induced by active forces. In the single copolymer chain, which reflects the compact nature of interphase chromosomes (1, 83), transient solid-like ordering would occur only if the distance between two bonded loci is roughly equal to the distance at which the pair potential between the non-bonded loci is a minimum (for the LJ potential, $r_{\text{LJ}}^* = 2^{1/6}\sigma$) (78). This argument is generic because it only depends on the presence of attractive interactions between non-bonded loci, which is needed to ensure that interphase chromosomes are compact. Application of the active forces increases r_{min}, and over a small range of forces r_{min} becomes commensurate with r_{LJ}^* (Figs. 5A–5C). It should be noted that the criterion for solid-like ordering is *independent* of the interaction parameters characterizing the chromosome, and depends only on the magnitude of the force (see SI Appendix, Sec. 11; Fig. S15). If the bonds are rigid, chain segments in the polymer chain, under confinement and active extensile forces, would align exhibiting a nematic order (51). Thus, the mechanisms of our FCC-like ordering and the nematic order in Ref. 51 are qualitatively different. However, both the studies show that active forces induce ordering in chromatin. Additional comparisons between the two models are given in SI Appendix, Sec. 12.

Studies of activity-induced changes in polymers. The roles of active processes and associated forces in chromosome organization and dynamics have been probed in theoretical studies based on polymer models (6, 50–55). It was shown that athermal random noise, when applied selectively on the active loci, affects the nuclear positioning of the active loci (6, 50, 54). Interestingly, using a combination of analytical theory and simulations (55), it has been shown that a modest difference in activity between active and inactive loci leads to microphase separation (55), a key feature of organization in interphase chromosomes. Using a model that is similar to that used here (54), it has been shown that active forces are important in driving heterochromatin to the periphery of the nucleus, which attests to the dynamic nature of chromatin structures. Typically applying active forces resulted in the increased MSD of the polymer chain (44–46, 55), which would not capture the suppressed chromatin mobility in human cells with active transcription (8, 13). Our minimal model, in the absence of active forces, leads to compact globule that exhibits glass-like dynamics over a long time (10), which may be needed to reproduce the experimental findings. Furthermore, it is likely that explicit inclusion of the precise epigenetic states, as done here for chromosomes 5 and 10, is required for observing the unexpected role of active forces on locus mobilities.

Connection to other experiments. Gu *et al.* (14) reported that transcription enhances the mobility of specific gene promoters/enhancers, which seems to contradict the results in a related experimental study (13). Gu *et al.* tracked the motion of only a few loci whereas Nagashima *et al.* (13) measured the motility of many loci across the nucleus. In addition, the active genes chosen by Gu *et al.* seem to be outliers in terms of changes in the gene expression level upon cell differentiation (14). Those genes show about 50–100 times difference in their expression levels upon activation or suppression following cell differentiation, although expression most of the genes change by a factor of 4–5. We surmise that the remarkably high level of gene transcription activity results in the mobility changes that are opposite to the global trend for all other genes (8, 13, 16). Using our results, we interpret, tentatively, that the enhanced motility observed for the activated genes (14) likely corresponds to $0 \leq F \leq 30$ or $90 \leq F \leq 120$, or possibly to the case where these genes are situated with no other co-activated genes in their proximity (see Discussion below). On the other hand, a majority of the active loci tracked in Nagashima *et al.* (13) likely corresponds to $60 \leq F \leq 90$.

In an another experiment (12), it was shown that acute depletion of cohesin, which mediates the formation of chromatin loops along with CTCFs (33, 39, 84, 85), results in increased motility of nucleosomes. Our polymer simulations capture the enhanced dynamics upon loop loss at $F = 0$. Indeed, when we compare the MSDs between $F = 80$ with loops and $F = 0$ without loops, a near quantitative agreement with the experiment (~20–25% increase) is obtained. This result implies that the loop loss may lead to the decreased transcription activity of some genes, possibly due to the disruption of interactions between the cognate promoter and enhancer. Although the downregulation of genes was observed upon loop loss (84–86), the overall effect on gene expression is not clearly understood since the loop formation is dynamic (37, 38, 87), and is likely coupled to transcription (88, 89). More precise theory that incorporates the effect of dynamic loops on transcription needs

to be constructed using the experiments that measure locus motion and real-time gene expression simultaneously (90).

The physical picture suggested here needs additional validation by future experiments. Direct observation of transient ordering could be challenging in the resolution limit of currently available in imaging methods. Possibly, the live-cell magnetic tweezer technique (60) may be utilized to test our finding. Recent experiments support our conclusions indirectly (30, 91, 92). In particular, Bohrer and Larson (92) highlighted the significant correlation between the spatial proximity and the co-expression of genes by analyzing the DNA and RNA FISH data (93). The role of gene co-activation is implicitly considered in our study because we apply active forces on all the gene-transcribing loci in a given chromosome region. We do not observe the activity-induced decrease in chromatin mobility when only one gene or a few loci are active (see SI Appendix, Sec. 13; Fig. S16). Based on these simulations, we conclude tentatively that motility of loci would increase (14) when only a single gene is transcribed whereas RNAPII-induced forces would suppress chromatin loci mobility when a few genes are co-expressed.

Limitations. Our model is not without limitations. For instance, explicit effect of transcription factors may bring the theory closer to experiments (94) but will come at the expense of complicated simulations. In addition, our model does not include hydrodynamic effects (51, 95, 96) that contribute significantly to the correlated motion of chromatin observed in the experiments (16). On the experimental side (13), the mobility change might arise due to DNA damage that could be induced by transcriptional inhibition (16, 97). The DNA repair process, which is not considered here, could be an alternate mechanism that could explain the increase in the MSD (16, 98).

For a more realistic picture of gene expression dynamics, our model should be extended to incorporate RNAP-induced force transmission in temporally and/or spatially dependent manner. This extension may also include the implementations of the features at higher resolution, such as the dynamic loop extrusion (35, 37, 38, 89) and the RNAP translocation with nascent RNAs elongated (64). Such a biophysical model would provide clearer pictures on how the gene expression is regulated in conjunction with the dynamically-evolving chromatin topology.

The prediction of transient solid-like ordering might be criticized because currently there is no experimental validation of the proposed mechanism. However, whether chromatin is solid-like (30) or liquid-like (12) continues to be controversial (99). It could adopt both these characteristics depending on length and time scales (100). In light of the unresolved nature of the dynamical state of chromatin, our proposal of the emergence of transient solid-like order is not unreasonable, especially considering that the simulations confirm the non-intuitive experimental observations. Additional experiments are needed to clarify the nature of the chromatin state, which is likely to change depending on the chromatin activity.

Methods

Assignment of the monomer type and loop anchors in CCM. We model chromosomes as a copolymer that embodies the epigenetic information. The locus type and loop anchor locations (see Fig. 1) in the CCM are assigned following the procedures given in our previous study (10). The locus type (euchromatin or heterochromatin) is assigned using the epigenetic information from the Broad ChromHMM track (101–103). The first 1–11 states are labeled as active (euchromatin, A) and states 12–15 as inactive (heterochromatin, B). To identify the loop anchors, we use the locations of unique CTCF loops identified by Rao et al. (33). We assign loop anchors to the loci whose genomic regions are closest to the centroids of the experimentally identified anchors. For the results shown above, we considered chromosome 5 (Chr5) in the region from 145.87 to 150.67 Mb from the GM12878 cell line with the resolution of 1.2 kb per locus. The numbers of active and inactive loci are $N_A = 982$ and $N_B = 3018$, respectively. In this Chr5 region, there are 13 CTCF loops. The simulations were performed using a single copolymer model of Chr5.

Potential energy function of CCM. The CCM energy function for a chromosome is given by $U_0 = U_{NB} + U_B + U_L$, where U_{NB}, U_B, and U_L represent the energetic contributions of non-bonding interactions, bond stretch, and CTCF-mediated loops, respectively. We use the Lennard-Jones potential for the non-bonding pair interactions, given by $U_{NB} = \sum_{i=1}^{N-2} \sum_{j=i+2}^{N} u_{LJ}(r_{i,j}|\epsilon_{\nu(i)\nu(j)}, \sigma)$, where

$$u_{LJ}(r|\epsilon, \sigma) = 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^{6} \right] , \qquad [6]$$

$r_{i,j} = |\mathbf{r}_i - \mathbf{r}_j|$ is the distance between the i^{th} and j^{th} loci, and σ is the diameter of a locus. The diameter of A and B type loci are the same. Here, $\nu(i)$ is the locus type, A or B, so there are three interaction parameters, ϵ_{AA}, ϵ_{BB}, and ϵ_{AB}. The bond stretch energy is written as $U_B = \sum_{i=1}^{N-1} u_b^0(|\mathbf{b}_i|)$, where $u_b^0(r) = u_{FENE}(r) + u_{WCA}(r)$ and $\mathbf{b}_i = \mathbf{r}_{i+1} - \mathbf{r}_i$. $u_{FENE}(r)$ is the finite extensible nonlinear elastic (FENE) potential (104, 105),

$$u_{FENE}(r) = -\frac{1}{2} K_S r_{max}^2 \ln \left(1 - \frac{r^2}{r_{max}^2} \right) , \qquad [7]$$

where r_{max} is the maximum bond length and K_S is the FENE spring constant. $u_{WCA}(r)$ is the Weeks-Chandler-Anderson (WCA) potential (106), given by,

$$u_{WCA}(r) = \left[u_{nb}(r|\epsilon_b, \sigma) + \epsilon_b \right] \Theta(r_{LJ}^* - r) , \qquad [8]$$

where $\Theta(x)$ is the Heaviside step function ($\Theta(x) = 1$ if $x > 0$ and $\Theta(x) = 0$ if $x \leq 0$). The WCA potential is the repulsive tail of the LJ potential, which is shifted and truncated at $r_{LJ}^* = 2^{1/6}\sigma$, and keeps a pair of adjacent loci from collapsing. Finally, the constrains imposed by the loop anchors is modeled using a harmonic potential,

$$U_L = \sum_{\{p,q\}} K_L (r_{p,q} - a)^2 , \qquad [9]$$

where $\{p, q\}$ is the set of indices of the loop anchors, a is the equilibrium bond length between the pair of loop anchors, and K_L is the harmonic spring constant.

The values of the parameters in U_B and U_L are $K_S = 30k_B T/\sigma^2$, $r_{max} = 1.5\sigma$, $\epsilon_b = 1.0k_B T$, $K_L = 300k_B T/\sigma^2$, and $a = 1.13\sigma$. With these parameter values, the equilibrium bond length is given by $b_0 = 0.96\sigma$ at which $u_b^0(r)$ is minimized. For the energetic parameters in U_{NB}, we used $\epsilon_{AA} = \epsilon_{BB} = 2.4k_B T$ and $\epsilon_{AB} = \frac{9}{11}\epsilon_{AA}$. The previous study showed that the CCM with this parameter set reproduces the Hi-C inferred contact maps for chromosomes 5 and 10 well (10). In particular, compartment features on ~ 5 Mb scale and topologically associating domains (TADs) on ~ 0.5 Mb are reproduced using the CCM simulations. In addition, the CCM simulations reproduced the dynamic properties that agree with experimental results.

Simulation details with active force. We performed Brownian dynamics simulations by integrating the following equation of motion, which for the i^{th} locus is given by,

$$\zeta \dot{\mathbf{r}}_i = -\frac{\partial}{\partial \mathbf{r}_i} U_0(r^N(t)) + \mathbf{R}_i(t) + \delta_{\nu(i)A}\mathbf{f}_i(t) , \qquad [10]$$

where ζ is the friction coefficient and $U_0(r^N(t))$ is the potential energy of the CCM polymer chain with the configuration,

$r^N(t) = \{\mathbf{r}_1(t), \cdots, \mathbf{r}_N(t)\}$). The Gaussian white noise, $\mathbf{R}_i(t)$, satisfies $\langle \mathbf{R}_i(t) \cdot \mathbf{R}_j(t') \rangle = 6\zeta k_B T \delta_{ij} \delta(t - t')$, where k_B is the Boltzmann constant and T is temperature. The Kronecker delta, $\delta_{\nu(i)A}$, in the last term of Eq. (10) ensures that $\mathbf{f}_i(t)$ acts only on the A-type loci. The exact form of $\mathbf{f}_i(t)$ is,

$$\mathbf{f}_i(t) = f_0 \left[\delta_{\nu(i-1)A} \hat{\mathbf{b}}_{i-1}(t) - \delta_{\nu(i+1)A} \hat{\mathbf{b}}_i(t) \right] . \qquad [11]$$

The time step for integrating Eq. (10) was chosen as $\delta t = 10^{-4} \tau_B$, where τ_B is the Brownian time, defined by $\tau_B = \sigma^2/D_0$. Here, $D_0 = k_B T/\zeta$ is the diffusion coefficient of a single locus. Using the Stokes-Einstein relation, $D_0 = k_B T/6\pi\eta R$, where η is the viscosity of the medium and R is the radius of a locus, we can evaluate the diffusion coefficient and the simulation time step in real units. We choose $\eta = 0.89 \times 10^{-3}$ Pa · s from the viscosity of water at 25°C and $R \approx \sigma/2$. We take $\sigma = 70$ nm from an approximate mean of the lower and upper bounds for the size of 1.2 kb of chromatin including six nucleosomes and linker DNAs, which are estimated as 20 nm and 130 nm, respectively (10). Hence, in real time and length units, $D_0 = \frac{(1.38 \times 10^{-23} \text{ Pa·m}^3/\text{K})(298\text{K})}{6\pi(0.89 \times 10^{-3} \text{ Pa·s})(35 \times 10^{-9} \text{ m})} \approx 7.0 \, \mu\text{m}^2/\text{s}$ and $\tau_B \approx 0.7$ ms.

For the most part, the results of the simulations are reported in the reduced units. The energy parameters are given in units of $k_B T$. Thus, the fundamental energy unit is $\epsilon = k_B T$, which means that the reduced temperature becomes $T^* = T k_B/\epsilon = 1$. The fundamental length and time units are σ and $\tau = (m\sigma^2/\epsilon)^{1/2}$, respectively, where m is the mass of a single locus. With an estimate of 260 kDa per nucleosome with 200 bps of dsDNA, we obtain $\tau \approx 55.5$ ns in real units so the time step is given by $\delta t \approx 1.26\tau$. The magnitude of the active force is $F = f_0 \sigma/\epsilon = f_0 \sigma/k_B T$. For instance, $F = 80$ corresponds to 3 to 16 pN in real units, based on the lower and upper bounds of σ specified above (20 nm to 130 nm).

Each simulation starts from a collapsed globule configuration that is obtained from the equilibration of an extended polymer chain using the low-friction Langevin thermostat at $T^* = 1.0$ in the absence of active force. We propagate the collapsed configuration with active force for $10^6 \delta t$ until the active polymer reaches a steady state where the radius of gyration and potential energy does not increase further. Subsequently, we run the simulation for additional $2 \times 10^8 \delta t$ and generate five independent trajectories to obtain the statistics needed to calculate the quantities of interest. All the simulations were performed using LAMMPS (107).

Inherent structure for the active polymer. Following the concept introduced by Stillinger and Weber (69), we investigated the *inherent structure* for each locus type in the CCM. The inherent structure is the ideal structure in molecular liquid which is preferred by the geometrical packing of particles, so it is the arrangement of particles expected by removing thermal excitations. Practically, the inherent structure is determined by minimizing the energy of the system using the steepest descent method (108). This procedure, however, is not directly applicable to our system because it involves the active force that is not derived from a given potential energy. Instead, we perform the quench of the CCM with the active force to low temperature. In a previous study, it was shown that quenching the monodisperse colloidal liquid generates a stable BCC (body-centered cubic) crystal, which cannot be easily obtained by the standard steepest descent (70).

To quench the system to a low temperature, we took the configuration from $T = 298K$ ($T^* = 1.0$) and ran Brownian dynamics simulation at $T \approx 3$ K ($T^* = 0.01$). The Brownian time and simulation time step are scaled accordingly such that $\tau_B' = 100\tau_B$ and $\delta t' = 10^{-4} \tau_B' = 100\delta t = 126\tau$. The quench simulation was run for $10^6 \delta t'$ during which the system energy was minimized until it reached a plateau value within $\sim 10^5 \delta t'$. We computed the radial distribution functions, $g_{AA}(r)$ and $g_{BB}(r)$, from the quenched configurations.

Bond-orientational order parameter. Following Ref. 71, we computed the bond-orientational order (BOO) parameter for each individual locus. The BOO with l-fold symmetry for the i^{th} locus is defined

as,

$$q_l(i) = \left[\frac{4\pi}{2l+1} \sum_{m=-l}^{l} |q_{lm}(i)|^2 \right]^{1/2} , \qquad [12]$$

where $q_{lm}(i)$ is the average of the spherical harmonics, Y_{lm}, for the bond angles formed between the i^{th} locus and its nearest neighbors,

$$q_{lm}(i) = \frac{\sum_{j \neq i}^{N} Y_{lm}(\mathbf{r}_j - \mathbf{r}_i)\Theta(r_c - r_{i,j})}{\sum_{j \neq i}^{N} \Theta(r_c - r_{i,j})} . \qquad [13]$$

We use $r_c = 1.4\sigma$, where the radial distribution function has the local minimum after the peak corresponding to the first coordination shell, as the cutoff pair distance for the nearest neighbors. The probability distribution and average of BOO parameter were computed depending on the locus type such that

$$P_\mu(q_{12}) = \frac{1}{N_\mu} \sum_{i=1}^{N} \delta_{\nu(i)\mu} \langle \delta(q_{12}(i) - q_{12}) \rangle , \qquad [14]$$

and

$$\langle q_{12} \rangle_\mu = \frac{1}{N_\mu} \sum_{i=1}^{N} \delta_{\nu(i)\mu} \langle q_{12}(i) \rangle . \qquad [15]$$

The distributions, $P_A(q_{12})$ and $P_B(q_{12})$, for different values of F are shown in Figs. S5E and S5F.

Effective potential energy. Since the active force is not stochastic, it may be treated as a pseudo-conservative force, which contributes to potential energy. In particular, we define the effective potential energy,

$$U_{\text{eff}} = U_0 - f_0 \sum_{i=1}^{N-1} \delta_{\nu(i)A} \delta_{\nu(i+1)A}(|\mathbf{b}_i| - b_0) , \qquad [16]$$

where the second term, denoted by U_a, represents the work due to the active force on the A-A bonds. It can be verified that $(\partial/\partial \mathbf{r}_i)U_a = -\delta_{\nu(i)A}\mathbf{f}_i$, so the equation of motion [Eq. (10)] can be rewritten as,

$$\zeta \dot{\mathbf{r}}_i = -\frac{\partial}{\partial \mathbf{r}_i} U_{\text{eff}}(r^N(t)) + \mathbf{R}_i(t) . \qquad [17]$$

The minus sign in U_a indicates that this potential prefers bond extension. Note that U_a only affects the bond potential energy of the A-A bonds. It is consistent with the effective bond potential, $u_{\text{b}}^{\text{eff}}(r)$ in Eq. 5, with $U_a = \sum_{i=1}^{N-1} \delta_{\nu(i)A} \delta_{\nu(i+1)A}[u_{\text{b}}^{\text{eff}}(|\mathbf{b}_i|) - u_{\text{b}}^0(|\mathbf{b}_i|)]$. In Fig. 5D, we showed the mean and variance of the effective potential energy for the A loci ($E_{\text{eff,A}}$) as a function of F. $E_{\text{eff,A}}$ is given by the sum of all the pairwise interactions involving the A loci, or the total effective potential energy excluding the contributions of B-B pairs,

$$E_{\text{eff,A}} = U_{\text{eff}} - \sum_{i=1}^{N-2} \sum_{j=i+2}^{N} \delta_{\nu(i)B} \delta_{\nu(j)B} u_{\text{LJ}}(r_{i,j}|\epsilon_{\nu(i)\nu(j)}, \sigma)$$
$$- \sum_{i=1}^{N-1} \delta_{\nu(i)B} \delta_{\nu(i+1)B} u_{\text{b}}^0(|\mathbf{b}_i|)$$
$$- \sum_{\{p,q\}} \delta_{\nu(p)B} \delta_{\nu(q)B} K_L(r_{p,q} - a)^2 . \qquad [18]$$

ACKNOWLEDGMENTS. We thank Alexandra Zidovska, Bin Zhang, Debayan Chakraborty, Xin Li, and Davin Jeong for useful discussions. We thank the National Science Foundation (CHE19-00093) and the Welch Foundation through the Collie-Welch Chair (F-0019) for support.

1. Lieberman-Aiden E, et al. (2009) Comprehensive Mapping of Long-Range Interactions Reveals Folding Principles of the Human Genome. *Science* 326(5950):289–293.

2. Bintu B, et al. (2018) Super-resolution chromatin tracing reveals domains and cooperative interactions in single cells. *Science* 362(6413):eaau1783.

3. Finn EH, Misteli T (2019) Molecular basis and biological function of variability in spatial genome organization. *Science* 365(6457):eaaw9498.

4. Lucas JS, Zhang Y, Dudko OK, Murre C (2014) 3D trajectories adopted by coding and regulatory DNA elements: First-passage times for genomic interactions. *Cell* 158(2):339–352.

5. Bronshtein I, et al. (2015) Loss of lamin A function increases chromatin dynamics in the nuclear interior. *Nature Communications* 6(1):8044.

6. Liu L, Shi G, Thirumalai D, Hyeon C (2018) Chain organization of human interphase chromosome determines the spatiotemporal dynamics of chromatin loci. *PLOS Computational Biology* 14(12):e1006617.

7. Ashwin SS, Nozaki T, Maeshima K, Sasai M (2019) Organization of fast and slow chromatin revealed by single-nucleosome dynamics. *Proceedings of the National Academy of Sciences of the United States of America* 116(40):19939–19944.

8. Shaban HA, Barth R, Recoules L, Bystricky K (2020) Hi-D: nanoscale mapping of nuclear dynamics in single living cells. *Genome Biology* 21(1):95.

9. Kang H, Yoon YG, Thirumalai D, Hyeon C (2015) Confinement-Induced Glassy Dynamics in a Model for Chromosome Organization. *Physical Review Letters* 115(19):198102.

10. Shi G, Liu L, Hyeon C, Thirumalai D (2018) Interphase human chromosome exhibits out of equilibrium glassy dynamics. *Nature Communications* 9(1):3161.

11. Di Pierro M, Potoyan DA, Wolynes PG, Onuchic JN (2018) Anomalous diffusion, spatial coherence, and viscoelasticity from the energy landscape of human chromosomes. *Proceedings of the National Academy of Sciences of the United States of America* 115(30):7753–7758.

12. Nozaki T, et al. (2017) Dynamic Organization of Chromatin Domains Revealed by Super-Resolution Live-Cell Imaging. *Molecular Cell* 67(2):282–293.e7.

13. Nagashima R, et al. (2019) Single nucleosome imaging reveals loose genome chromatin networks via active RNA polymerase II. *Journal of Cell Biology* 218(5):1511–1530.

14. Gu B, et al. (2018) Transcription-coupled changes in nuclear mobility of mammalian cis-regulatory elements. *Science* 359(6379):1050–1055.

15. Tortora MM, Salari H, Jost D (2020) Chromosome dynamics during interphase: a biophysical perspective. *Current Opinion in Genetics and Development* 61:37–43.

16. Zidovska A, Weitz DA, Mitchison TJ (2013) Micron-scale coherence in interphase chromatin dynamics. *Proceedings of the National Academy of Sciences of the United States of America* 110(39):15555–15560.

17. Brueckner F, Cramer P (2008) Structural basis of transcription inhibition by α-amanitin and implications for RNA polymerase II translocation. *Nature Structural and Molecular Biology* 15(8):811–818.

18. Bensaude O (2011) Inhibiting eukaryotic transcription. Which compound to choose? How to evaluate its activity? *Transcription* 2(3):103–108.

19. Yin H, et al. (1995) Transcription against an applied force. *Science* 270(5242):1653–1657.

20. Wang MD, et al. (1998) Force and velocity measured for single molecules of RNA polymerase. *Science* 282(5390):902–907.

21. Xing Y, Johnson CV, Moen PT, McNeil JA, Lawrence JB (1995) Nonrandom gene organization: Structural arrangements of specific pre-mRNA transcription and splicing with SC-35 domains. *Journal of Cell Biology* 131(6 II):1635–1647.

22. Croft JA, et al. (1999) Differences in the localization and morphology of chromosomes in the human nucleus. *Journal of Cell Biology* 145(6):1119–1131.

23. Reddy KL, Zullo JM, Bertolino E, Singh H (2008) Transcriptional repression mediated by repositioning of genes to the nuclear lamina. *Nature* 452(7184):243–247.

24. Hnisz D, Shrinivas K, Young RA, Chakraborty AK, Sharp PA (2017) A Phase Separation Model for Transcriptional Control. *Cell* 169(1):13–23.

25. Cho WK, et al. (2018) Mediator and RNA polymerase II clusters associate with transcription-dependent condensates. *Science* 361(6400):412–415.

26. Chong S, et al. (2018) Imaging dynamic and selective low-complexity domain interactions that control gene transcription. *Science* 361(6400):eaar2555.

27. Sabari BR, et al. (2018) Coactivator condensation at super-enhancers links phase separation and gene control. *Science* 361(6400):eaar3958.

28. Hsieh THS, et al. (2020) Resolving the 3D Landscape of Transcription-Linked Mammalian Chromatin Folding. *Molecular Cell* 78(3):539–553.e8.

29. Jiang Y, et al. (2020) Genome-wide analyses of chromatin interactions after the loss of Pol I, Pol II, and Pol III. *Genome Biology* 21(1):158.

30. Strickfaden H, et al. (2020) Condensed Chromatin Behaves like a Solid on the Mesoscale In Vitro and in Living Cells. *Cell* 183(7):1772–1784.

31. Huggins ML (1941) Solutions of long chain compounds. *The Journal of Chemical Physics* 9(5):440.

32. Flory PJ (1941) Thermodynamics of High Polymer Solutions. *The Journal of Chemical Physics* 9(8):660.

33. Rao SS, et al. (2014) A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* 159(7):1665–1680.

34. Fudenberg G, et al. (2016) Formation of Chromosomal Domains by Loop Extrusion. *Cell Reports* 15(9):2038–2049.

35. Nuebler J, Fudenberg G, Imakaev M, Abdennur N, Mirny LA (2018) Chromatin organization by an interplay of loop extrusion and compartmental segregation. *Proceedings of the National Academy of Sciences* 115(29):E6697–E6706.

36. Dey A, Shi G, Takaki R, Thirumalai D (2023) Structural changes in chromosomes driven by multiple condensin motors during mitosis. *Cell Reports* 42(4):112348.

37. Gabriele M, et al. (2022) Dynamics of CTCF- and cohesin-mediated chromatin looping revealed by live-cell imaging. *Science* 376(6592):496–501.

38. Mach P, et al. (2022) Cohesin and CTCF control the dynamics of chromosome folding. *Nature Genetics* 54(12):1907–1918.

39. Kim Y, Shi Z, Zhang H, Finkelstein IJ, Yu H (2019) Human cohesin compacts DNA by loop extrusion. *Science* 366(6471):1345–1349.

40. Golfier S, Quail T, Kimura H, Brugués J (2020) Cohesin and condensin extrude DNA loops in a cell cycle-dependent manner. *eLife* 9:e53885.

41. Ghosh A, Gov NS (2014) Dynamics of active semiflexible polymers. *Biophysical Journal* 107(5):1065–1073.

42. Eisenstecken T, Gompper G, Winkler RG (2016) Conformational properties of active semi-flexible polymers. *Polymers* 8(8):37–42.

43. Sakaue T, Saito T (2017) Active diffusion of model chromosomal loci driven by athermal noise. *Soft Matter* 13(1):81–87.

44. Bianco V, Locatelli E, Malgaretti P (2018) Globule-like conformation and enhanced diffusion of active polymers. *Physical Review Letters* 121(21):217802.

45. Put S, Sakaue T, Vanderzande C (2019) Active dynamics and spatially coherent motion in chromosomes subject to enzymatic force dipoles. *Physical Review E* 99(3):032421.

46. Foglino M, et al. (2019) Non-equilibrium effects of molecular motors on polymers. *Soft Matter* 15(29):5995–6005.

47. Smrek J, Chubak I, Likos CN, Kremer K (2020) Active topological glass. *Nature Communications* 11(1):1–11.

48. Locatelli E, Bianco V, Malgaretti P (2021) Activity-Induced Collapse and Arrest of Active Polymer Rings. *Physical Review Letters* 126(9):97801.

49. Ghosh A, Spakowitz AJ (2022) Active and thermal fluctuations in multi-scale polymer structure and dynamics. *Soft Matter* 18(35):6629–6637.

50. Ganai N, Sengupta S, Menon GI (2014) Chromosome positioning from activity-based segregation. *Nucleic Acids Research* 42(7):4145–4159.

51. Saintillan D, Shelley MJ, Zidovska A (2018) Extensile motor activity drives coherent motions in a model of interphase chromatin. *Proceedings of the National Academy of Sciences of the United States of America* 115(45):11442–11447.

52. Liu K, Patteson AE, Banigan EJ, Schwarz JM (2021) Dynamic Nuclear Structure Emerges from Chromatin Cross-Links and Motors. *Physical Review Letters* 126(15):158101.

53. Mahajan A, Yan W, Zidovska A, Saintillan D, Shelley MJ (2022) Euchromatin Activity Enhances Segregation and Compaction of Heterochromatin in the Cell Nucleus. *Physical Review X* 12(4):41033.

54. Jiang Z, Qi Y, Kamat K, Zhang B (2022) Phase Separation and Correlated Motions in Motorized Genome. *Journal of Physical Chemistry B* 126(30):5619–5628.

55. Goychuk A, Kannan D, Chakraborty AK, Kardar M (2023) Polymer folding through active processes recreates features of genome organization. *Proceedings of the National Academy of Sciences* 120(20):e2221726120.

56. Venter JC, et al. (2001) The sequence of the human genome. *Science* 291(5507):1304–1351.

57. García-Martínez J, Aranda A, Pérez-Ortín JE (2004) Genomic run-on evaluates transcription rates for all yeast genes and identifies gene regulatory mechanisms. *Molecular Cell* 15(2):303–313.

58. Doi M, Edwards SF (1988) *The theory of polymer dynamics.* (Oxford University Press).

59. Backlund MP, Joyner R, Moerner WE (2015) Chromosomal locus tracking with proper accounting of static and dynamic errors. *Physical Review E* 91(6):062716.

60. Keizer VI, et al. (2022) Live-cell micromanipulation of a genomic locus reveals interphase chromatin mechanics. *Science* 377(6605):489–495.

61. Stillinger FH (1976) Phase transitions in the Gaussian core system. *The Journal of Chemical Physics* 65(10):3968–3974.

62. Eckert T, Bartsch E (2002) Re-entrant Glass Transition in a Colloid-Polymer Mixture with Depletion Attractions. *Physical Review Letters* 89(12):20–23.

63. Banerjee PR, Milin AN, Moosa MM, Onuchic PL, Deniz AA (2017) Reentrant Phase Transition Drives Dynamic Substructure Formation in Ribonucleoprotein Droplets. *Angewandte Chemie - International Edition* 56(38):11354–11359.

64. Henninger JE, et al. (2021) RNA-Mediated Feedback Control of Transcriptional Condensates. *Cell* 184(1):207–225.e24.

65. Kim J, Kim C, Sung BJ (2013) Simulation Study of Seemingly Fickian but Heterogeneous Dynamics of Two Dimensional Colloids. *Physical Review Letters* 110(4):047801.

66. Berthier L, Biroli G (2011) Theoretical perspective on the glass transition and amorphous materials. *Reviews of Modern Physics* 83(2):587–645.

67. Kirkpatrick TR, Thirumalai D (2015) Colloquium: Random first order transition theory concepts in biology and physics. *Reviews of Modern Physics* 87(1):183–209.

68. Kirkpatrick TR, Thirumalai D (1988) Comparison between dynamical theories and metastable states in regular and glassy mean-field spin models with underlying first-order-like phase transitions. *Physical Review A* 37(11):4439–4448.

69. Stillinger FH, Weber TA (1984) Packing structures and transitions in liquids and solids. *Science* 225(4666):983–989.

70. Thirumalai D (1989) Liquid and crystalline states of monodisperse charged colloidal particles. *The Journal of Physical Chemistry* 93(15):5637–5644.

71. Steinhardt PJ, Nelson DR, Ronchetti M (1983) Bond-orientational order in liquids and glasses. *Physical Review B* 28(2):784–805.

72. ten Wolde PR, Ruiz-Montero MJ, Frenkel D (1995) Numerical Evidence for bcc Ordering at the Surface of a Critical fcc Nucleus. *Physical Review Letters* 75(14):2714–2717.

73. Omar AK, Klymko K, Grandpre T, Geissler PL (2021) Phase Diagram of Active Brownian Spheres: Crystallization and the Metastability of Motility-Induced Phase Separation. *Physical Review Letters* 126(18):188002.

74. Loi D, Mossa S, Cugliandolo LF (2008) Effective temperature of active matter. *Physical Review E* 77(5):051111.

75. Grosberg AY, Joanny JF (2015) Nonequilibrium statistical mechanics of mixtures of particles in contact with different thermostats. *Physical Review E* 92(3):032118.

76. Mandal R, Klymko K, DeWeese MR (2017) Entropy Production and Fluctuation Theorems for Active Matter. *Physical Review Letters* 119(25):1–6.

77. Omar AK, Klymko K, GrandPre T, Geissler PL, Brady JF (2023) Tuning nonequilibrium phase transitions with inertia. *The Journal of chemical physics* 158(7):074904.

78. Cho HW, Shi G, Kirkpatrick TR, Thirumalai D (2021) Random First Order Transition Theory for Glassy Dynamics in a Single Condensed Polymer. *Physical Review Letters*

126(13):137801.

79. Nagano T, et al. (2017) Cell-cycle dynamics of chromosomal organization at single-cell resolution. *Nature* 547(7661):61–67.

80. Landick R (2006) The regulatory roles and mechanism of transcriptional pausing. *Biochemical Society Transactions* 34(6):1062–1066.

81. Darzacq X, et al. (2007) In vivo dynamics of RNA polymerase II transcription. *Nature Structural and Molecular Biology* 14(9):796–806.

82. Hager GL, McNally JG, Misteli T (2009) Transcription Dynamics. *Molecular Cell* 35(6):741–753.

83. Manuelidis L (1990) A view of Interphase Chromosomes. *Science* 250(4987):1533–1540.

84. Schwarzer W, et al. (2017) Two independent modes of chromatin organization revealed by cohesin removal. *Nature* 551(7678):51–56.

85. Rao SS, et al. (2017) Cohesin Loss Eliminates All Loop Domains. *Cell* 171(2):305–320.e24.

86. Hsieh THS, et al. (2022) Enhancer–promoter interactions and transcription are largely maintained upon acute loss of CTCF, cohesin, WAPL or YY1. *Nature Genetics* 54(12):1919–1932.

87. Hansen AS, Cattoglio C, Darzacq X, Tjian R (2018) Recent evidence that TADs and chromatin loops are dynamic structures. *Nucleus* 9(1):20–32.

88. Banigan EJ, et al. (2023) Transcription shapes 3D chromatin organization by interacting with loop extrusion. *Proceedings of the National Academy of Sciences* 120(11):e2210480120.

89. Zhang S, Übelmesser N, Barbieri M, Papantonis A (2023) Enhancer–promoter contact formation requires RNAPII and antagonizes loop extrusion. *Nature Genetics* 55(5):832–840.

90. Brückner DB, Chen H, Barinov L, Zoller B, Gregor T (2023) Stochastic motion and transcriptional dynamics of pairs of distal DNA loci on a compacted chromosome. *Science* 380(6652):1357–1362.

91. Bignaud A, et al. (2022) Transcriptional units form the elementary constraining building blocks of the bacterial chromosome. *bioRxiv* p. 2022.09.16.507559.

92. Bohrer CH, Larson DR (2023) Synthetic analysis of chromatin tracing and live-cell imaging indicates pervasive spatial coupling between genes. *eLife* 12:e81861.

93. Su JH, Zheng P, Kinrot SS, Bintu B, Zhuang X (2020) Genome-Scale Imaging of the 3D Organization and Transcriptional Activity of Chromatin. *Cell* 182(6):1641–1659.e26.

94. Wagh K, et al. (2023) Dynamic switching of transcriptional regulators between two distinct low-mobility chromatin states. *Science Advances* 9(24):eade1122.

95. Bruinsma R, Grosberg AY, Rabin Y, Zidovska A (2014) Chromatin hydrodynamics. *Biophysical Journal* 106(9):1871–1881.

96. Eshghi I, Zidovska A, Grosberg AY (2022) Symmetry-based classification of forces driving chromatin dynamics. *Soft Matter* 18(42):8134–8146.

97. Lee KB, Wang D, Lippard SJ, Sharp PA (2002) Transcription-coupled and DNA damage-dependent ubiquitination of RNA polymerase II in vitro. *Proceedings of the National Academy of Sciences of the United States of America* 99(7):4239–4244.

98. Eaton JA, Zidovska A (2020) Structural and Dynamical Signatures of Local DNA Damage in Live Cells. *Biophysical Journal* 118(9):2168–2180.

99. Zidovska A (2020) Chromatin: Liquid or Solid? *Cell* 183(7):1737–1739.

100. Nozaki T, et al. (2023) Condensed but liquid-like domain organization of active chromatin regions in living human cells. *Science Advances* 9(14):eadf1488.

101. Rosenbloom KR, et al. (2013) ENCODE Data in the UCSC Genome Browser: Year 5 update. *Nucleic Acids Research* 41(D1):56–63.

102. Ernst J, Kellis M (2010) Discovery and characterization of chromatin states for systematic annotation of the human genome. *Nature Biotechnology* 28(8):817–825.

103. Ernst J, et al. (2011) Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature* 473(7345):43–49.

104. Warner HR (1972) Kinetic Theory and Rheology of Dilute Suspensions of Finitely Extendible Dumbbells. *Industrial and Engineering Chemistry Fundamentals* 11(3):379–387.

105. Kremer K, Grest GS (1990) Dynamics of entangled linear polymer melts: A molecular-dynamics simulation. *The Journal of Chemical Physics* 92(8):5057–5086.

106. Weeks JD, Chandler D, Andersen HC (1971) Role of repulsive forces in determining the equilibrium structure of simple liquids. *The Journal of Chemical Physics* 54(12):5237–5247.

107. Plimpton S (1995) Fast Parallel Algorithms for Short-Range Molecular Dynamics. *Journal of Computational Physics* 117:1–19.

108. Stillinger FH, Weber TA (1985) Inherent structure theory of liquids in the hard-sphere limit. *The Journal of Chemical Physics* 83(9):4767–4775.