# Elevator mechanism dynamics in a sodium-coupled dicarboxylate transporter

Kinz-Thompson, C.D.,[1,†] Lopez-Redondo, M.L.,[2,‡] Mulligan C.,[3] Sauer, D.B.,[2,§] Marden, J.J.,[2] Song, J.,[2] Tajkhorshid, E.,[4] Hunt, J.F.,[5] Stokes, D.L.,[2] Mindell, J.A.,[6*] Wang, D.N,[2*] Gonzalez, Jr., R.L.[1*]

1. Department of Chemistry, Columbia University, New York, New York 10027, US
2. Department of Cell Biology, New York University School of Medicine, New York, New York 10016, US
3. School of Biosciences, University of Kent, Canterbury, Kent, CT2 7NH, UK
4. Theoretical and Computational Biophysics Group, NIH Center for Macromolecular Modeling and Bioinformatics, Beckman Institute for Advanced Science and Technology, Department of Biochemistry, Center for Biophysics and Quantitative Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, US
5. Department of Biological Sciences, Columbia University, New York, New York 10027, US
6. Membrane Transport Biophysics Section, Porter Neuroscience Research Center, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD 20892, US

*Correspondence: rlg2118@columbia.edu, da-neng.wang@med.nyu.edu, mindellj@ninds.nih.gov
†Current address: Department of Chemistry, Rutgers University-Newark, Newark, New Jersey 07012, US
‡Current address: Instituto de Biomedicina de Valencia, Consejo Superior de Investigaciones Científicas (IBV-CSIC), 46010 Valencia, Spain
§Current address: Centre for Medicines Discovery, Nuffield Department of Medicine, University of Oxford, Oxford, OX1 3QU, UK

# Abstract

VcINDY, the sodium-dependent dicarboxylate transporter from *Vibrio cholerae*, is responsible for $C_4$- and $C_5$-carboxylate uptake into cells. The molecular mechanism of how VcINDY physically moves substrates across the membrane, and does so in an energetically efficient manner, is unclear. Here, we use single-molecule fluorescence resonance energy transfer experiments to directly observe the individual mechanistic steps that VcINDY takes to translocate substrates across a lipid bilayer, and then test key predictions of transport cycle mechanistic models. Our data provide the first direct evidence that VcINDY undergoes stochastic, elevator-type conformational motions that enable substrate translocation. Kinetic analysis suggests that the two protomers of the VcINDY homodimer undergo those motions in a non-cooperative manner, and thus catalyze two independent transport reactions. The relative substrate independence of those motions supports the notion that the VcINDY transport cycle maintains strict co-substrate coupling using a cooperative binding mechanism. Finally, thermodynamic modeling provides insight into how such a cooperative binding mechanism provides a generalized approach to optimizing transport for many secondary active transporters.

# Introduction

Secondary active transporters are transmembrane proteins that harvest free energy from electrochemical
gradients to drive the movement of substrates across a membrane (Fig. 1) (Stein and Lieb, 1986). In the absence
of a free energy source, these transporters move substrates back and forth across the membrane without
creating a net change in substrate concentration (Jardetzky, 1966). Net transport occurs because the transporter
can dissipate the electrochemical gradient of a 'driving' substrate to harvest free energy, and then couple that
free energy to the 'uphill' movement of a 'driven' substrate across the membrane (Forrest et al., 2011). The
mechanistic details of this coupling are governed by the transport cycle, which describes how the transporter
binds its substrates, translocates them across the membrane, releases them, and resets itself for the next round
(Fig. 1B). While the specific details of the transport cycle and how it is achieved on a molecular level are unique
to each transporter or transporter family, the active use of free energy suggests that transport cycles must
maintain strict substrate coupling to minimize energetically wasteful 'slippage cycles' (*c.f.*, Fig. 1B) in which
driving substrates are transported without driving the translocation of the driven substrate (Stein and Lieb, 1986).

While all secondary active transporters work by facilitating a general, dynamic equilibrium (*e.g.*,
$mA_\text{outside} + nB_\text{outside} \rightleftharpoons mA_\text{inside} + nB_\text{inside}$, where $A$ and $B$ are co-substrates with stoichiometries $m$ and $n$), the
molecular details of how this chemical reaction maps on to the transport cycle are determined by a transporter's
conformational dynamics. According to the alternating access hypothesis, secondary active transporters must
undergo a reversible conformational transition between at least two distinct conformational states: one with the
substrate binding pocket in a state that is 'outward-facing' with respect to membrane ($C_o$), and one in an 'inward-
facing' state ($C_i$) (Jardetzky, 1966). While there are many possible structural implementations of alternating
access, including the so-called 'rocker-switch', 'rocking-bundle', and 'elevator' transport mechanisms (Drew and
Boudker, 2016), at least two distinct conformational transitions between $C_i$ and $C_o$ must occur in a complete
transport cycle (*c.f.*, Fig. 1B). Here, we investigate the role of such conformational dynamics in the transport
cycle of the prototypical sodium-dependent divalent anion transporter from mesophilic *Vibrio cholerae*, VcINDY,
and elucidate how they are tuned to enable efficient transport and minimize slippage.

VcINDY is a well-characterized member of the divalent anion/sodium symporter (DASS) class of
secondary active transporters, which extends across all kingdoms of life. In mammals, it is represented by the
solute carrier 13 (SLC13) gene family, which includes members that transport di- and trivalent anions, including
sulfate, citrate, and succinate (Bergeron et al., 2013; Markovich, 2012). Within this family, the SLC13A5 citrate
transporter, NaCT, has significant metabolic and neurologic health importance for humans, and mutations in
NaCT can cause epilepsy in children (reviewed in (Sauer et al., 2022b)). Like NaCT, VcINDY is a $Na^+$-driven
carboxylate transporter, but VcINDY is primarily a succinate transporter with an electrogenic transport cycle
stoichiometry of 3 $Na^+$ ions per succinate (Fig. 1A) (Mulligan et al., 2014; Fitzgerald et al., 2017). Over the past
decade, extensive *in vitro* biochemical characterization (Mulligan et al., 2014, 2016; Fitzgerald et al., 2017;
Sampson et al., 2020, 2021), and deep structural insight from X-ray crystallography, cryo-electron microscopy

(cryo-EM), and computational studies (Mancusso et al., 2012; Mulligan et al., 2016; Nie et al., 2017; Sauer et
al., 2021, 2022a), have made VcINDY into the prototypical transporter for understanding the mechanistic details
of transport in the DASS family.

While the biochemical and structural studies published to-date do not yield a description of the molecular
mechanism for the complete VcINDY transport cycle, recent combinations of computational modeling,
biochemical, and structural analysis have provided insight (Mulligan et al., 2016; Sauer et al., 2020). Structural
studies have revealed VcINDY is a homodimeric membrane protein with two-fold rotational symmetry; however,
in all these studies, VcINDY was only observed in a single $C_i$ conformational state in which the substrate binding
pockets of both protomers are accessible only from the cytosolic side (*i.e.*, the global homodimer conformation
is $C_i$-$C_i$) (Mancusso et al., 2012; Nie et al., 2017; Sauer et al., 2020). Computational and crosslinking studies
suggest that VcINDY functions using an elevator mechanism, with a large translocation of the 'transport' domain
that carries substrates from one side of the membrane to the other, and a relatively stationary 'scaffold' domain
that harbors the dimerization interface  (Mulligan et al., 2016). This model is supported by recent structural work
revealing a $C_o$ state for LaINDY, a close homolog from *Lactobacillus acidophilus* (Sauer et al., 2020). While these
lines of evidence support an elevator mechanism, there has been no experimental observation of a transition
between these $C_o$ and $C_i$ states, nor any direct evidence that they are actually responsible for achieving
alternating access during transport. Comprehensive validation of this model requires confirming the proposed
elevator motions, connecting them to the transport cycle, and understanding how those conformational dynamics
enable VcINDY to achieve tight co-substrate coupling between $Na^+$ and succinate to minimize energetically costly
slippage of substrates across the membrane (Fig. 1B).

In this work, using single-molecule fluorescence resonance energy transfer (smFRET) microscopy, we
aimed to directly observe the global conformational dynamics of homodimeric VcINDY in a lipid-bilayer
membrane, allowing us to characterize the structural and kinetic details of its transport cycle. To achieve this,
we designed two complementary VcINDY constructs for smFRET experiments. Each smFRET construct
contained a FRET donor- and acceptor fluorophore pair engineered at residue positions such that the resulting
donor-acceptor distance-dependent FRET efficiency ($E_{FRET}$) would report on the conformational dynamics
responsible for substrate translocation (Fig. 1C). These smFRET constructs can therefore be used to directly
observe the motions that VcINDY undergoes to achieve alternating access (Fig. 1B). In performing smFRET
experiments with these constructs, we obtained direct experimental evidence that (i) global conformational
dynamics of VcINDY during the transport cycle are consistent with the elevator model of transport, (ii) the
individual protomers of homodimeric VcINDY exhibit minimal cooperativity, and (iii) VcINDY minimizes slippage
using a mechanism in which co-substrate binding is highly cooperative. These findings provide the first
comprehensive description of the molecular underpinnings of the VcINDY transport cycle, and, by extension,
deep insight into the transport cycle of all DASS transporters.

## Results

**smFRET constructs designed to report on substrate translocation**

In order to directly monitor the VcINDY conformational dynamics relevant to the transport cycle (Fig. 1B), we designed two smFRET constructs based on the assumption that substrate translocation in VcINDY involves a large-scale conformational change between the $C_o$-$C_o$ and the $C_i$-$C_i$ conformations (*c.f.*, Fig. 1C) (Mancusso et al., 2012; Mulligan et al., 2016; Nie et al., 2017; Sauer et al., 2020). One construct was designed to contain a FRET donor- and acceptor fluorophore pair on the external, periplasmic face of the transport domains of VcINDY whereas the other construct was designed to contain them on the internal, cytosolic face. These complementary designs comprise a 'multi-perspective' approach with an internal self-consistency where any global conformational motions of the VcINDY homodimer occurring during the transport cycle (*e.g.*, an elevator motion) can be detected on both the external and internal sides of the protein.

To design these 'external' and 'internal' smFRET constructs, we identified solvent-exposed residues in the transport domain with optimal distances between the two equivalent positions across the homodimer (Fig. 1C). Specifically, residues were chosen such that the distance would undergo a significant change upon a transition between the structurally defined $C_i$-$C_i$ and the computationally predicted $C_o$-$C_o$ states (Mancusso et al., 2012; Mulligan et al., 2016; Nie et al., 2017; Sauer et al., 2020). In an optimized smFRET experiment, the corresponding change in the distance-dependent $E_{FRET}$ will utilize the full dynamic range of $E_{FRET}$ from zero to one. To achieve this, we sought residues that moved from separations of less than the Förster distance ($R_0$), which corresponds to the half-maximum $E_{FRET}$ of the donor-acceptor pair, to more than the $R_0$, or vice versa. We used Alexa Fluor 555 as our donor and Alexa Fluor 647 as our acceptor with $R_0$ ~51 Å (Lambert, 2019). Given these considerations, we chose Gly211, located on the extracellular face of the transport domain as our external site, with a predicted $E_{FRET}$ change from ~0.20 to ~0.93 upon a transition from the $C_o$-$C_o$ to the $C_i$-$C_i$ state (Fig. 1C). Likewise, we chose Ser436, located on the cytoplasmic face of the transport domain as our internal site, with a predicted $E_{FRET}$ change from ~0.70 and ~0.12 upon transition from the $C_o$-$C_o$ to the $C_i$-$C_i$ state. Single VcINDY homodimer $E_{FRET}$ *versus* time trajectories (*i.e.*, $E_{FRET}$ trajectories) recorded using the external and internal smFRET constructs should then report upon the same set of global VcINDY conformational dynamics involved in substrate translocation but from different perspectives.

Biochemically active variants of the two VcINDY smFRET constructs designed above were created, overexpressed, purified, and labeled with donor and acceptor fluorophores to generate the externally and internally labeled smFRET constructs (VcINDY[ext] and VcINDY[int], respectively) (see Methods and Materials for extended details) (Fig. 1–Figure Supplement 1). Both VcINDY[ext] and VcINDY[int] were able to catalyze substrate translocation in liposomes, and exhibited transport activities that were only three-fold reduced relative to wildtype VcINDY and were similar to the activity of the unlabeled cysteine-containing mutants (Fig. 1–Figure Supplement 2). To investigate the conformational dynamics of VcINDY in a membrane, we reconstituted VcINDY[ext] and VcINDY[int] into lipid bilayer nanodiscs (Fig. 2) (Denisov and Sligar, 2017). Nanodisc-based experimental

conditions are ideal for interrogating the $C_o$ to $C_i$ transition at equilibrium, since both sides of the membrane are

exposed to identical solutions. Furthermore, the nanodiscs used here have a diameter of ~108 Å (Bayburt and

Sligar, 2010), which helped establish single-molecule resolution by ensuring that only a single VcINDY dimer

could occupy each nanodisc. After chromatographic purification, these nanodiscs appeared well dispersed when

imaged using negative stain electron microscopy (Fig. 2—Figure Supplement 1).

smFRET experiments were performed using a widefield, TIRF microscope to record the fluorescence

intensity emitted from individual, immobilized VcINDY-containing nanodiscs (Fig. 2). The corresponding $E_{FRET}$

trajectory for each molecule was then calculated using $E_{FRET} = I_{acceptor} / (I_{donor} + I_{acceptor})$, where $I_x$ is the observed

intensity of either the donor or the acceptor fluorophore. Along with the chromatographic purification used in

nanodisc reconstitution, the selection criteria for $E_{FRET}$ trajectory (see Methods and Materials), ensured that each

$E_{FRET}$ trajectory originated from a single VcINDY dimer (Fig. 1—Figure Supplement 1).

**Multiple $E_{FRET}$ states are stochastically sampled during VcINDY substrate translocation**

To characterize the conformational dynamics of VcINDY during the transport cycle (Fig. 1B), we performed

equilibrium smFRET experiments using VcINDY[ext] or VcINDY[int] in which the buffer contained saturating

concentrations of both $Na^+$ and succinate co-substrates (Buffer HS: High $Na^+$ with Succinate). The

conformational dynamics that occur under these conditions are the same as those involved in vectorial

transport—even in the absence of a chemical free energy driving force. Thus, by saturating the VcINDY smFRET

constructs with both co-substrates, we expected to record $E_{FRET}$ trajectories that report on the equilibrium

conformational dynamics of the substrate translocation reaction between a substrate-bound $C_o$ state and a

substrate-bound $C_i$ state ($C_o$" and $C_i$" in Figure 1B), which we hypothesized resemble the $C_o$-$C_o$ and $C_i$-$C_i$

conformations, respectively (Fig. 1C).

In the presence of both $Na^+$ and succinate, the $E_{FRET}$ trajectories originating from individual VcINDY[ext] or

VcINDY[int] dimers stochastically exhibited instantaneous jumps between transiently stable states characterized

by distinct $E_{FRET}$ values (*i.e.*, $E_{FRET}$ states) (Fig. 3). Qualitatively, the lifetimes of these $E_{FRET}$ states were generally

several seconds, and jumps occurred to $E_{FRET}$ states with both higher and lower average $E_{FRET}$ values. By

inspection, it was not uncommon for individual $E_{FRET}$ trajectories to visit many distinct $E_{FRET}$ states (*e.g.*, Fig. 3,

lower panels), and the same $E_{FRET}$ states were often revisited within a single $E_{FRET}$ trajectory. Most $E_{FRET}$

trajectories, however, exhibited just a few jumps to a low number of distinct $E_{FRET}$ states before relatively rapid

photobleaching of the donor and/or acceptor fluorophores prematurely ended our ability to measure $E_{FRET}$.

Strikingly, the distribution of $E_{FRET}$ states across the entire population of VcINDY[ext] or VcINDY[int] $E_{FRET}$

trajectories spanned nearly the entire range of possible $E_{FRET}$ values, from zero to one, and did not exhibit well-

defined peaks (Fig. 4). Relative to the signal-to-background ratio of individual $E_{FRET}$ trajectories (Fig. 3), the broad

spread of these $E_{FRET}$ state distributions (Fig. 4) suggests that VcINDY dimers exhibit significant conformational

heterogeneity. Regardless, the spread of the $E_{FRET}$ state distributions is consistent with the amplitude of

fluctuations predicted for a large, elevator conformational transition between the $C_o$-$C_o$ and $C_i$-$C_i$ conformations

using the donor-acceptor pairs engineered into both VcINDY[ext] and VcINDY[int] (Fig. 1C).

**VcINDY samples $C_o$-$C_o$, $C_i$-$C_i$, and mixed $C_o$-$C_i$ conformational states during substrate transport**

After observing dynamic $E_{FRET}$ trajectories that exhibited instantaneous jumps between widely distributed $E_{FRET}$

states for both VcINDY[ext] and VcINDY[int] (Figs. 3 and 4), we desired to understand how these $E_{FRET}$ states and

dynamics correspond to the underlying conformational states and dynamics of the VcINDY transport cycle (Fig.

1B). To do so, we sought the simplest, quantitative description of the $E_{FRET}$ trajectories that was globally

consistent with all the $E_{FRET}$ trajectories recorded in an experiment. Because of the instantaneous jumping

behavior of the $E_{FRET}$ trajectories (Fig. 3), we used a global hidden Markov model (HMM) to quantify the common

$E_{FRET}$ states and kinetics present in all the $E_{FRET}$ trajectories (see Methods and Materials for details) (Fig. 4–

Figure Supplements 1-2). In the presence of both co-substrates, we found that the most parsimonious global

HMMs that could account for the observed complexity in the $E_{FRET}$ trajectories had three $E_{FRET}$ states for both

VcINDY[ext] and VcINDY[int] (Fig. 4, Fig. 4–Figure Supplements 3-4). Given that we find three $E_{FRET}$ states in buffer

HS from both smFRET perspectives, and that the $E_{FRET}$ trajectories recorded under different $Na^+$ and succinate

conditions were also best described by three $E_{FRET}$ states (see below) (Fig. 4—Figure Supplement 3), we

concluded that our smFRET data report primarily on transitions between three common, global conformations of

the VcINDY dimer.

According to the alternating access principle, at least some part of these three $E_{FRET}$ states should

correspond to the $C_o$ and $C_i$ states in the transport cycle (Jardetzky, 1966) (Fig. 1B). For VcINDY[ext], the $E_{FRET}$ =

0.25 state is consistent with our estimate of $E_{FRET}$ = 0.20 for the $C_o$-$C_o$ conformation, and the $E_{FRET}$ = 0.85 state

is consistent with our estimate of $E_{FRET}$ = 0.93 for the $C_i$-$C_i$ conformation (Fig. 1C, Fig. 4–Figure Supplement 4).

Likewise, for VcINDY[int], the $E_{FRET}$ = 0.83 state is consistent with our estimate of $E_{FRET}$ = 0.70 for the $C_o$-$C_o$

conformation, and the $E_{FRET}$ = 0.16 state is consistent with our estimate of $E_{FRET}$ = 0.12 for the $C_i$-$C_i$ conformation

(Fig. 1C, Fig. 4–Figure Supplement 4). Thus, the $E_{FRET}$ states we observe in our smFRET experiments are

broadly consistent with the elevator mechanism suggested by previous structural work and illustrate how these

large-scale conformational changes are coupled to substrate transport.

The small differences between the expected and observed $E_{FRET}$ values ($\Delta E_{FRET} \sim 0.1$) might be due to:

(i) differences in the $C_o$-$C_o$ and/or $C_i$-$C_i$ structures which nanodisc-embedded VcINDY adopts in our smFRET

experiments and between those in the computational, X-ray, and cryo-EM studies (Mancusso et al., 2012;

Mulligan et al., 2016; Nie et al., 2017; Sauer et al., 2020), (ii) structural perturbations of VcINDY due to

fluorophore labeling, (iii) linker contributions to the distance between fluorophores, (iv) constraints on fluorophore

mobility due to interactions with protein and/or membrane, or even (iv) photophysical complications, such as

'photoblinking', that effect $E_{FRET}$ estimates. Given the experimental consistency with the structural estimates for

both VcINDY[ext] and VcINDY[int], we assigned the $E_{FRET}$ states discussed above to the $C_o$-$C_o$ and $C_i$-$C_i$

conformational states. Additionally, these assignments are self-consistent with respect to the equilibrium state population distribution of the transport cycle (Fig. 1) in that the most occupied $E_{FRET}$ state in both smFRET experiments was the $C_i$-$C_i$ VcINDY conformational state and was thus independent of the smFRET perspective used (Fig. 4).

The remaining, unassigned $E_{FRET}$ state can be explained as either a 'mixed' $C_o$-$C_i$ conformation of the VcINDY dimer or a new, 'intermediate' conformation distinct from $C_o$-$C_o$ and $C_i$-$C_i$. Based on biochemical and structural considerations, we favor assigning this $E_{FRET}$ state to the more parsimonious, mixed conformation. The existence of such a mixed conformation is only possible if the protomers of VcINDY operate independently, which is suggested by previous Hill coefficient measurements for succinate transport (Mulligan et al., 2014). Additionally, the two-fold rotational symmetry of the VcINDY dimer renders the $C_o$-$C_i$ and $C_i$-$C_o$ conformations structurally degenerate, and thus the $C_o$-$C_i$ and $C_i$-$C_o$ $E_{FRET}$ states would both yield the same $E_{FRET}$ value, which is consistent with our analysis. In contrast, similar symmetry considerations for a VcINDY dimer with a unique intermediate protomer conformation would yield up to six distinct $E_{FRET}$ states in the absence of strict protomer cooperativity or anti-cooperativity. Finally, computational alignment of the VcINDY scaffold domain of the $C_o$-$C_o$ and $C_i$-$C_i$ conformations shows that the donor and acceptor fluorophores in a mixed $C_o$-$C_i$ state would be ~50 Å apart for VcINDY$^{ext}$ and ~59 Å apart for VcINDY$^{int}$—distances that correspond to $E_{FRET}$ values of 0.53 and 0.29, respectively. Given the caveats described above, these estimates are in reasonable agreement with the $E_{FRET}$ values of 0.54 and 0.42 that we obtained from our global HMM analysis of the smFRET data (Fig. 4–Figure Supplement 4).

Despite the limited observation window of any one individual $E_{FRET}$ trajectory, the global HMM analyses we performed provide a quantification of the kinetics present within the collective set of $E_{FRET}$ trajectories (Fig. 4—Figure Supplement 4). Regardless of the smFRET perspective used (VcINDY$^{ext}$ or VcINDY$^{int}$), the elevator motion of a single protomer in the VcINDY dimer across the membrane occurs in these experiments at a rate of ~0.4 s$^{-1}$ (*i.e.*, once every ~2.5 seconds), with no apparent directional dependence. Notably, direct transitions between $C_o$-$C_o$ and $C_i$-$C_i$ where both protomers undergo an apparently simultaneous transition within the temporal resolution of our TIRF microscope were approximately an order of magnitude slower (~0.04 s$^{-1}$), and therefore much less likely to occur. This suggests that the individual protomers of a VcINDY dimer primarily undergo the translocation reaction in an independent and uncoordinated manner. Overall, the elevator motions of VcINDY responsible for substrate translocation that occur in these experiments are approximately an order of magnitude faster than the turn-over rate for transport measured in bulk biochemical transport experiments of ~0.02 s$^{-1}$ (Mulligan et al., 2014). Along with the fact that our smFRET experiments were performed at lower a temperature (21 °C *versus* 30 °C), this suggests that VcINDY undergoes multiple substrate translocation reactions before a productive transport event occurs, and that the rate limiting step of VcINDY, and possibly DASS family, catalyzed transport may involve substrate binding and/or dissociation.

**VcINDY exhibits conformational dynamics that are largely substrate independent**

The basic mechanistic model of a transport cycle for VcINDY makes several fundamental predictions regarding the conformational dynamics that occur during transport (Fig. 1B). Consistent with the alternating access hypothesis (Jardetzky, 1966), VcINDY should be able to switch from a $C_o$ to a $C_i$ conformation in the apo, substrate-free state. Also, to preserve substrate coupling, we expect that the partially bound transporter ($C_o'$ and $C_i'$ in Fig. 5) will not undergo outward to inward transitions (and vice versa) to minimize uncoupled slippage of $Na^+$ or succinate. To test these predictions, we performed additional smFRET experiments using VcINDY[int] in the absence of both co-substrates (*i.e.*, in the apo state) (Buffer LX: Low $Na^+$ with no succinate) as well as in the presence of high concentrations of one co-substrate but the absence of the other (*i.e.*, in the only bound $Na^+$- or only succinate bound states) (Buffer HX: High $Na^+$ with no succinate;  Buffer LS: Low $Na^+$ with Succinate). Because of the consistency between VcINDY[int] and VcINDY[ext] we observed in buffer HS (Figs. 3,4), these substrate-dependence experiments were done with only the VcINDY[int] construct. Where appropriate, we replaced $Na^+$ with choline to maintain ionic strength and osmolarity. We note that all of our buffers likely contain trace (*i.e.*, 'low') amounts of contaminating $Na^+$, but at several orders of magnitude less than the measured Michaelis constant, $K_M = 41.7$ mM, for transport (Mulligan et al., 2014).

As observed for the smFRET experiments recorded in the presence of high concentrations of both co-substrates (Buffer HS), the smFRET experiments recorded under all three additional co-substrate conditions (Buffers LX, HX, and LS) yielded $E_{FRET}$ trajectories that spontaneously exhibited instantaneous jumps between $E_{FRET}$ states spanning a comparably wide range as was seen in buffer HS (Fig. 4). When we analyzed these data using the same global HMM approach described above, the most parsimonious quantification of the sets of $E_{FRET}$ trajectories in each smFRET experiment was again a global HMM with three $E_{FRET}$ states (Fig. 4–Figure Supplement 3). In these global HMMs, both the $E_{FRET}$ states and rate constants for the transitions between them were very similar to those obtained in Buffer HS (Fig. 4–Figure Supplement 4). While there were slight differences in the equilibrium population distribution of the three $E_{FRET}$ states observed in each co-substrate condition, the kinetics of these global HMMs were relatively unchanged with single protomer translocation reactions occurring all at approximately the same rate of ~0.5 $s^{-1}$. Thus, our quantitative kinetic analysis reveals that the conformational motions of VcINDY that are probed along the coordinate between the donor and acceptor fluorophore in VcINDY[int] are relatively independent of the co-substrate composition. Although these results conform to our expectation for the apo state, the observation of frequent transitions in the $E_{FRET}$ trajectories recorded under the single-substrate conditions was unexpected.

**Thermodynamic modeling shows that compensatory tuning of co-substrate binding and dissociation can minimize slippage cycles**

In search of an explanation for the surprising kinetics observed under the single-substrate conditions, we next analyzed the thermodynamic efficiency of the VcINDY transport cycle. We modeled the free-energy transduction

of a secondary active transporter using Hill's method of cycles (Hill, 2005) to determine how the different
molecular processes involved in transport might contribute to slippage. For simplicity, we considered co-
substrate binding as a strictly sequential, two-step process with mirror symmetry (*i.e.*, the same binding/release
order on both faces of the membrane). To be concrete, we considered the most parsimonious molecular
interpretation of this kinetic scheme for secondary active transport where (i) the rate constants are all first-order
or pseudo-first order, (ii) the rate constants are independent of the $C_o$ or $C_i$ state, and (iii) the translocation
reaction is independent of substrate binding status (Fig. 5). The resulting, simplified kinetic scheme has three
Hill cycles—two of which are involved in slippage (cycles *a* and *b*) and one that is responsible for transport (cycle
*c*) (Fig. 5).

We then calculated the thermodynamic efficiency, $\eta$, of a transporter with these assumptions (Fig. 5) (Hill,
2005). For a maximally efficient transporter, $\eta \approx 1$ and the transporter tightly couples free energy from the driving
to the driven co-substrate with very little slippage. In that situation, the thermodynamic efficiency is $\eta \propto 1 -$
$(\eta_{ac} - \eta_{bc}) + \cdots$ with small higher order terms; here we have assumed $A$ is the driving substrate, but the
conclusions discussed below are independent of this. Under our assumptions,

287
$$\eta_{ac} = \frac{J_a}{J_c} = \left( \frac{[A]_o^m - [A]_i^m}{[A]_o^m [B]_o^n - [A]_i^m [B]_i^n} \right) \left( \frac{2k_t + k_d''}{k_a''} \right),$$

288 and

289
$$\eta_{bc} = \frac{J_b}{J_c} = \left( \frac{[B]_o^n - [B]_i^n}{[A]_o^m [B]_o^n - [A]_i^m [B]_i^n} \right) \left( \frac{k_t([A]_o^m + [A]_i^m) + k_a'[A]_o^m [A]_i^m}{k_d'} \right),$$

where $J_\kappa$ is the cycle flux for cycle $\kappa$, concentrations are indicated by square brackets, $A$ and $B$ are the first and
second binding co-substrates with stoichiometries $m$ and $n$, respectively, the translocation rate constant is $k_t$,
association and dissociation rate constants are $k_a$ and $k_d$, respectively, and prime and double-prime denote the
first and second binding reactions, respectively (Fig. 5). The first terms in these equations show that the relative
amount of slippage that a transporter undergoes depends on the instantaneous thermodynamic driving forces
*via* the co-substrate chemical gradients, while the second terms show that it also depends on the translocation
and substrate binding and dissociation reaction kinetics. Thus, the only reliable way nature can optimize this
transporter into a highly efficient free energy transducer over a broad range of environmental conditions is to
have evolved control over slippage *via* evolutionary tuning of these rate constants to reduce the total magnitude
of $\eta_{ac}$ and $\eta_{bc}$. This suggests that optimal transporters have faced evolutionary pressure to decrease $k_t$, $k_a'$
and/or $k_d''$, and to increase $k_a''$ and/or $k_d'$ such that the first binding co-substrate weakly binds until the second
arrives and then both co-substrates can tightly bind. Unfortunately, tuning these rate constants will also affect
the overall transport cycle flux, $J_c$, and this creates a speed-accuracy tradeoff.

The speed-accuracy tradeoff can, however, be navigated. On the molecular level, the clearest way to
reduce slippage is for the binding pocket to stabilize the *apo* and fully bound states (C and C"), by some amount
$\Delta G'$ and $\Delta G''$, respectively, and to source all of that energy from a compensatory destabilization of the partially

bound conformation, C', by an amount $\Delta G' + \Delta G''$, all without affecting the transition state energy barriers (Fig.
6). This is a conceptually similar idea to the use of substrate binding energy to stabilize the transition state for
enzyme-substrate transition state complementarity (Fersht, 2017). In our case, this isoenergetic change
increases $k_a'$ and $k_d''$ while decreasing $k_a''$ and $k_d'$, which reduces slippage and increases the thermodynamic
efficiency of the transporter. It also has the effect of increasing the transport cycle flux *via* ~75% of the terms
defining $J_c$ (see *Methods and Materials* for the expression for $J_c$). Given this substantial ability to speed up
transport in the complicated expression for $J_c$, we suspect that these compensatory, isoenergetic binding and
dissociation reaction tunings optimally navigate the speed-accuracy tradeoff by simultaneously decreasing the
relative fraction of slippage while increasing the total amount of transport. The exact extent of these abilities,
however, depends on the instantaneous internal and external co-substrate concentrations, which can vary
significantly across physiologically relevant conditions (Fig. 6–Figure Supplement 1) (Lo et al., 2006).
Regardless, this thermodynamic modeling of the free energy transduction efficiency of a secondary active
transporter predicts that slippage can be reduced simply by isoenergetic optimization of the energy landscape
of the co-substrate binding reaction (Fig. 6). In other words, increased cooperativity of co-substrate binding can
itself lead to minimized slippage.

Finally, we also used this thermodynamic modeling-based approach to estimate the occupancy of apo
VcINDY in our smFRET experiments performed in presence of only saturating Na⁺ (Buffer HX). Specifically, we
modeled the reported $K_M$ and $V_{max}$ values for the Na⁺ titration and succinate titration experiments that recorded
succinate uptake rates in the biochemical proteoliposome-based transport assay data of Mulligan and coworkers
(see *Methods and Materials* for extended details) (Mulligan et al., 2014). This was an underdetermined problem,
so we assumed that a single, average rate constant defined several molecular processes: $\langle k \rangle \equiv k_t = k_d' = k_d''$.
Markov chain Monte Carlo optimization yielded estimates of $k_a' = 1.5 \times 10^7 M^{-3} s^{-1}$, $k_a'' = 7.0 \times 10^8 M^{-1} s^{-1}$, and
$\langle k \rangle = 1.05 \times 10^3 \ s^{-1}$ (Fig. 5-Figure Supplement 1 and 2), and reasonably matched the original titration data (Fig.
5-Figure Supplement 3). The mismatch is due to the common rate constant assumption, and probably to
differences in the actual binding mechanism assumed here (*e.g.*, perhaps not all three Na⁺ ions bind before
succinate). Regardless, these estimates are consistent with the biochemical transport data in the context of our
kinetic mechanism, so we used them to calculate the steady-state probability of apo VcINDY in Buffer HX to be
6.5% (see *Methods and Materials*). This estimate suggests that even with 100 mM Na⁺ in solution, VcINDY still
spends a non-negligible portion of time with an unoccupied binding pocket. In this light, the majority of the
unexpected elevator motions present in the E_FRET trajectories recorded in the presence of only one co-substrate
might originate from apo, rather than partially bound, VcINDY.

**Dicarboxylate substrate binding pocket is enthalpically stabilized by sodium ions**
To obtain atomic-resolution insight into the Na⁺ and dicarboxylate substrate binding reaction for the VcINDY
transport cycle, we determined how the conformational state of the VcINDY homodimer is influenced by different

dicarboxylate substrates. We therefore separately crystallized detergent-solubilized VcINDY with $Na^+$ and three
different $C_4$-dicarboxylate substrates (succinate, fumarate, and malate) and solved their structures at 3.09 Å,
3.29 Å, and 3.50 Å, respectively. Regardless of the dicarboxylate substrate identity, VcINDY adopted the same
$C_i$-$C_i$ state in all three of these structures. Similarly, as seen in the previous structure of the $C_i$-$C_i$ state (Mancusso
et al., 2012; Nie et al., 2017; Sauer et al., 2020), the binding pocket of the dicarboxylate substrate in all three
structures was partially comprised of two hairpin loops that coordinated distinct $Na^+$ ions, Na1 and Na2 (Fig. 7).
Given this binding geometry, it is possible that tight binding of both co-substrates requires stabilization of these
hairpin loops and that their local conformational rearrangements gates binding and/or release of the co-
substrates.

# Discussion

**350 The VcINDY transport domain undergoes elevator motions to enable alternating access**

The ubiquitous presence in our experiments of stochastic, instantaneous transitions between $E_{FRET}$ states
consistent with the $C_o$-$C_o$ and the $C_i$-$C_i$ conformations is the first direct evidence for the hypothesis that VcINDY
uses an elevator transport mechanism to enable alternating access (Fig. 3) (Mulligan et al., 2016). Because we
directly observed these stochastic, elevator motions in the presence of substrates but absence of an
electrochemical gradient (Buffer HS) and also in the absence of any co-substrates at all (Buffer LX), it is clear
that the VcINDY transport domain can translocate the substrate binding site across the membrane using only
thermal energy—as is expected for the conformational motions of a secondary active transporter that are
responsible for transport. Moreover, the rate constants for that process are on the order of $10^{-1}$ $s^{-1}$ (Fig. 4—Figure
Supplement 4), suggesting that the energy barrier separating the $C_o$ and $C_i$ state along the translocation reaction
coordinate is not prohibitively high. Since the VcINDY transport cycle is much slower and occurs with a rate on
the order of $10^{-2}$ $s^{-1}$ (Mulligan et al., 2014), it seems that multiple back and forth substrate translocation events
occur before a productive transport event is completed. This observation is consistent with a model in which a
molecular process other than transport domain translocation govern the transport rate (*e.g.*, gating dynamics
within the substrate binding pocket itself).

**365 The conformational dynamics of VcINDY protomers are non-cooperative**

The experimental observation of a mixed $C_o$-$C_i$ conformation for the VcINDY dimer constrains the role that the
two protomers can play in the transport cycle. Previously, proteoliposome-based transport experiments
determined that VcINDY transports succinate with a Hill coefficient of 0.88 ± 0.13 (Mulligan et al., 2014). Given
the VcINDY transport stoichiometry of 3:1 $Na^+$:dicarboxylate (Fitzgerald et al., 2017), this Hill coefficient
measurement suggested that the VcINDY transport cycle is non-cooperative and that each VcINDY protomer
can independently transport succinate. We supported this biochemical hypothesis on a molecular level by
interpreting our experimental observation of the $E_{FRET}$ state assigned to the mixed $C_o$-$C_i$ conformation as direct

evidence that the VcINDY protomers undergo non-cooperative substrate translocation reactions during vectorial
transport. Since the VcINDY dimer can significantly occupy the $C_o$-$C_i$ state, the two VcINDY protomers must not
undergo a strictly cooperative translocation reaction (*i.e.*, $C_o$-$C_o \rightleftharpoons C_i$-$C_i$). Furthermore, they must not undergo a
strictly anti-cooperative translocation reaction either (*i.e.*, $C_o$-$C_i \rightleftharpoons C_i$-$C_o$), because this behavior would preclude
the occupation of the $C_o$-$C_o$ and $C_i$-$C_i$ conformations that we observe in our smFRET experiments and in
structural studies. Therefore, our results provide direct evidence that the VcINDY protomers undergo non-
cooperative translocation reactions, and that the VcINDY transport cycle itself can be considered to occur
independently in each protomer.

In support of this idea, a recent structure of CitS, another homodimeric elevator transporter in a different
structural family, reveals a CitS dimer with one inward-facing and one outward-facing protomer (Kim et al., 2017),
and a recent high-speed atomic force microscopy study of the CitS elevator translocation motion reported that
CitS undergoes non-cooperative interprotomer conformational dynamics (Maity et al., 2022). Combined with
evidence that Glt$_{Ph}$, an elevator-type, glutamate transporter that unlike VcINDY is from a thermophilic organism,
also shows independence between the motions of its three protomers (Akyuz et al., 2013; Erkens et al., 2013;
Ruan et al., 2017), these results suggest that such independence may be a general feature of oligomeric elevator
transporters (Garaeva and Slotboom, 2020).

**Slippage cycles are avoided without inhibiting VcINDY transport domain translocation**
To avoid wasting the free energy provided by the $Na^+$ gradient that drives transport of substrates across the
membrane, the conformational motions of VcINDY must be coordinated. In particular, VcINDY must minimize
slippage cycles in which $Na^+$ ions are transported across the membrane independent of dicarboxylate (Fig. 1).
We had hypothesized that VcINDY avoids such slippage cycles by inhibiting the substrate binding pocket
translocation reaction (*i.e.*, $C_o \rightleftharpoons C_i$) when sub-stoichiometric amounts of the co-substates are bound, and by
promoting those motions only when VcINDY is either empty or fully bound. However, the smFRET experiments
we recorded under single substrate conditions, yielded $E_{FRET}$ trajectories that exhibited no discernable
differences in the $E_{FRET}$ states or their rates of transitions, regardless of the presence of relatively high
concentrations of either individual co-substrate in solution (Fig. 4–Figure Supplement 4). Given our determination
that those observed transitions could reasonably correspond to apo VcINDY, it is quite possible that transitions
between $C_o$ and $C_i$ in partially bound states are directly suppressed and that we were not able to observe those
effects, but our smFRET experiments suggest that VcINDY also uses another, additional mechanism to minimize
slippage (Fig. 1B).

There are at least two additional mechanisms that VcINDY can use to minimize slippage. One possibility
is a secondary conformational change—independent of $C_o \rightleftharpoons C_i$ translocation—to 'gate' both binding and release
of co-substrates. For instance, while VcINDY could be able to translocate substrates across the membrane if the
$C_o \rightleftharpoons C_i$ transition is not completely inhibited, a slippage cycle cannot be completed without co-substrate release.

Mechanistically, such gating could be achieved with an additional VcINDY conformational equilibrium that is
relatively independent of the transport domain translocation reaction $C_o \rightleftharpoons C_i$. Without direct structural insight
into such a hypothetical equilibrium, it is attractive to think that it might involve a conformational rearrangement
of the binding pocket and, in particular, the hairpin loops (*c.f.*, Fig. 7). In this scenario, the transitions we observe
in the smFRET data for partially bound VcINDY would reflect transitions between so-called "occluded" states.

Another possible mechanism to minimize slippage is to ensure that co-substrate binding is extremely
cooperative, such that stable binding requires both co-substrates to be present. In our smFRET experiments
performed in Buffers HX and LS, this mechanism would manifest as transitions in the $E_{FRET}$ trajectories that
predominately correspond to transport domain translocation of apo VcINDY. This alternative scenario is
consistent with the non-negligible fraction of apo VcINDY we estimated to be present under these experimental
conditions. Furthermore, it could explain the apparent lack of substrate dependence observed relative to our
smFRET experiments in Buffer LX (Fig. 4-Figure Supplement 4). This cooperative co-substrate binding scenario
agrees with our thermodynamic modeling (Fig. 5), which predicts that an optimized transporter that has evolved
to minimize slippage cycle flux should have extremely stable apo and fully bound states with very weak binding
of the initial co-substrate(s) (Fig. 6). The extent of that optimization reflects how nature has navigated the speed-
accuracy tradeoff between rapid transport and minimal slippage. This raises the possibility that some secondary
active transporters undergo a non-negligible amount of slippage in order to be efficient at transporting the driven
co-substrate as long as they do not significantly waste the free energy of the driving co-substrate.

Experimental evidence is also consistent with a cooperative co-substrate binding mechanism. For
instance, our X-ray crystallographic structures show that Na1, Na2, and dicarboxylate substrates all interact with
common hairpin loop elements (Fig. 7). That the VcINDY conformation is effectively independent of dicarboxylate
identity suggests that this fully bound conformation is very stable—as predicted in our modeling of optimal co-
substrate binding. Moreover, recent experiments demonstrated that, in the presence of high concentrations of
only Na$^+$, residues associated with the substrate binding pocket are protected from PEG modification (Sampson
et al., 2020; Sauer et al., 2022a) indicating some interaction between Na$^+$ and the apo VcINDY binding pocket.
This result is consistent with a Na$^+$-first binding order, and with predictions that slippage is minimized and that
transport is optimized by weak binding of the first co-substrate(s) (Fig. 6). In future work, determination of the
affinities and binding order of co-substrates, together with structure characterization of the sites, will be required
to develop a comprehensive understanding of how co-substrate binding is employed as a mechanism to avoid
slippage cycles during vectorial transport by VcINDY.

# 437 Conclusion

Previous studies of the structure, transport biochemistry, and alternating access mechanism of VcINDY have
established it as a prototypical membrane-bound transporter of the DASS family (Mancusso et al., 2012; Mulligan
et al., 2014, 2016; Nie et al., 2017; Fitzgerald et al., 2017; Sauer et al., 2020; Sampson et al., 2020). Here, we

have advanced this fundamental understanding of the DASS family and secondary active transport by directly
determining how the conformational dynamics of VcINDY enable it to achieve alternating access, and how those
dynamic motions are tuned in order to maintain the efficiency of the transport cycle. Specifically, our study
provides direct evidence that (i) the VcINDY translocation reaction uses a thermally driven, elevator mechanism;
(ii) VcINDY protomer conformational dynamics are non-cooperative; (iii) the VcINDY transport cycle uses
cooperative substrate binding to minimizes slippage; and (iv) cooperative substrate binding is a general approach
to optimize transport efficiency for certain secondary active transporters. Altogether, these findings immediately
suggest future studies aimed at determining the extent to which highly cooperative co-substrate binding and/or
additional conformational dynamics that gate transport (Drew and Boudker, 2016; Lolkema and Slotboom, 2019;
Stein and Lieb, 1986) enable VcINDY to avoid slippage cycles during vectorial transport. Overall, the powerful
smFRET-based experimental modality for interrogating the function and regulation of VcINDY established here
is easily extended to study the transport cycles of other DASS family transporters, and even dimeric transporters
from other families, to further our understanding of secondary active transport.

# Methods and Materials

**Expression and purification of VcINDY**. Expression and purification of wild type VcINDY was carried out
according to our previous protocol (Mancusso et al., 2012). Briefly, *Escherichia coli* (*E. coli*) BL21-AI cells
(Invitrogen) were transformed with a modified pET vector encoding N-terminal 10X His tagged VcINDY. Cells
were grown for 16 hours post-induction at 19°C. After cell breakage, membranes were solubilized in 1.2% N-
decyl-β-maltoside (DM) and the protein was purified on a Ni-NTA affinity column (Qiagen). Following His-tag
removal, VcINDY was further purified by size-exclusion chromatography (SEC) in buffer containing 25 mM Tris
pH 8, 100 mM NaCl, 5% glycerol, 0.1% DM unless otherwise indicated.

**Purification of VcINDY mutants**. Expression and purification of VcINDY mutants G211C and S436C was
performed as with the wildtype protein, but with the following modifications. All buffers contained 10 mM sodium
citrate but not glycerol. Membranes were solubilized in 1.0% N-dodecyl-β-maltoside (DDM). The protein was
eluted from Ni-NTA agarose beads using 300 mM imidazole. Finally, the SEC buffer contained 25 mM Tris pH
8, 100 mM NaCl, 0.1% DDM and 0.5 mM TCEP.

**Expression, purification and biotinylation of membrane scaffolding protein**. Expression and purification of
membrane scaffolding protein (MSP1E3D1) was carried out according to a previously published protocol
(Denisov et al., 2004). In brief, *E. coli* BL21-Gold (DE3) cells (Agilent) were transformed with a modified pET 28a
vector encoding N-terminal 7X His tagged MSP with a mutated cysteine at position 277 for biotinylation. Cells
were grown at 30°C for 4 hrs post 0.1 mM IPTG induction at OD 0.8. After cell breakage, lysate was bound to
Ni-NTA agarose beads (Qiagen) for 4 hours with nutation at 4°C. Protein was eluted with 100 – 250 mM imidazole
and dialyzed into 100 mM Tris pH 7.4, 300 mM NaCl for 1 hr at 4°C. The His-tag was then removed by overnight

TEV digestion at 12°C. Cleaved MSP protein was subsequently biotinylated using EZ-Link Maleimide-PEG11-
Biotin (Promega) according to the manufacturer's instructions. Specifically, a 20-fold molar excess of biotin
reagent dissolved in DMSO was incubated with protein overnight at 4°C. Labeled MSP1E3D1-T277C protein
was further purified by SEC in 100 mM NaCl, 1 mM TCEP, 20 mM Tris pH 7.4.

**Fluorophore Labeling.** Stock solutions of Alexa Fluor 555 C2 maleimide (AF555) and Alexa Fluor 647 C2
maleimide (AF647) were prepared in anhydrous DMSO at 3mM and stored at -20°C. Prior to labeling, a 1 mg/ml
solution of VcINDY (total of 0.5mg to 1mg) was treated with 10mM DTT for 2hrs at 4°C to reduce the cysteines.
DTT was then removed by 3 serial dilutions in SEC buffer (25 mM tris pH 8.0, 100 mM NaCl, 5% glycerol, 0.15%
DDM, 1 mM TCEP) followed by re-concentration with an Amicon concentrator (cutoff 50 kDa). This preparation
was incubated with a 10-fold molar excess of AF555 and AF647 overnight at 4°C, protected from light. The
resulting solution was purified on Superdex 200 10/300 GL size exclusion column to remove the unreacted dye.
Fractions from the elution peak were combined and the labeling stoichiometry was determined with a
spectrophotometer using the following wavelengths and extinction coefficients: VcINDY: $\lambda$=280 nm with $\epsilon$ =
57075 cm$^{-1}$M$^{-1}$; AF555: $\lambda$= 556 nm with $\epsilon$= 155,000 cm$^{-1}$M$^{-1}$; AF647: $\lambda$= 651 nm with $\epsilon$= 270,000 cm$^{-1}$M$^{-1}$. Typical
labeling efficiency was 42% for each dye.

**Nanodisc Reconstitution.** A thin film of *E. coli* polar lipid (EPL) was prepared by adding 10 mg of chloroform
stock solution to a glass test tube. The solvent was evaporated with argon gas and the test tube was placed in
a vacuum chamber at ~1 mTorr for ~2 hr. This lipid film was resuspended by adding 285 µl of nanodisc buffer
(20mM Tris pH7.4; 100mM NaCl, 0.5mM EDTA, 0.5mM TCEP) followed by vortexing and sonication in an
ultrasonic bath (Laboratory Supplies Co. Inc, Model G112SP1T) with 10 sec bursts alternating with 10 sec rests
for a total of 3min. This aqueous lipid solution was solubilized by adding 215 µl of 10% DDM to produce 500 µl
of final lipid stock concentration at 20mg/ml.

To reconstitute VcINDY into nanodiscs, detergent-solubilized EPL was added to VcINDY at a molar ratio
of 280 and incubated on ice for 10 min. Biotinylated MSP was then added to this mixture at an 8-fold molar
excess to produce 500 µl of a solution containing 2.5mM EPL, 72 µM MSP, 9 µM VcINDY and 7.2 mM DDM.
This solution was incubated at 4°C for 1hr followed by the addition of SM-2 Bio-Beads in three steps: 300 mg
initially, 200 mg added 60 min later, and 100 mg added at 90 min later and incubated overnight. During these
incubations, solutions were kept at 4°C and gently stirred.

Nanodiscs containing VcINDY were purified by incubating the solution with 0.5 ml of Ni-NTA beads pre-
equilibrated with nanodisc buffer (20 mM Tris pH=7.4; 100 mM NaCl; 0.5 mM EDTA; 0.5 mM TCEP) for 30 min
at 4°C. The beads were packed into a column and washed with 1 ml of nanodisc buffer. Nanodiscs containing
VcINDY were then eluted with 2 ml nanodisc buffer supplemented with 0.3 M imidazole and the peak elution
fractions were pooled and concentrated with an Amicon concentrator (cutoff 50 kDa). The concentrated sample
was fractionated on Superdex 200 10/300 GL size exclusion column (GE Healthcare) pre-equilibrated and eluted

with nanodisc buffer. The fractions were evaluated by SDS-PAGE and those containing both MSP and VcINDY

were pooled, and concentrated to ~0.2-0.4 mg/ml. Typical yield for 1mg of VcINDY was ~0.4 mg of reconstituted

protein (VcINDY plus MSP). Electron microscopy grids were prepared in 2% (w/v) phosphotungstic acid (pH 7.5)

and were examined in a Talos L120C transmission electron microscope (FEI-Thermo Fisher Scientific).

Nanodisc samples were frozen in aliquots and stored at –80°C.

**Proteoliposome Reconstitution and Transport Assay.** VcINDY was reconstituted into proteoliposomes as

previously described (Mulligan et al., 2016, 2014). These proteoliposomes were then used in succinate transport

assays as previously described (Mulligan et al., 2016, 2014). Briefly, proteoliposomes were exchanged into

external buffer containing tritiated succinate with an inward facing Na⁺ gradient, samples taken at various time

points were quenched and rapidly filtered over a 0.22 $\mu m$ nitrocellulose membrane, and then succinate content

inside the proteoliposomes was measured using scintillation counting. Transport rates were calculated from a

linear fit to at least three data points within the first minute of the transport reaction.

**TIRF Microscopy.** smFRET imaging experiments were performed on a prism-based, wide-field TIRF

microscope at a temperature of 21 ± 1 °C. Briefly, a 532 nm diode-pumped solid-state continuous-wave laser

(Laser Quantum, GEM532) was directed through an acousto-optic modulator (AOM) (Isomet, IMDD-P80L) such

that the +1 order peak is directed toward a prism suspended over a 60x, 1.2 NA water immersion objective

(Nikon). VcINDY embedded nanodiscs were tethered to the surface of quartz slides derivatized with dilute biotin-

poly(ethylene)-glycol (PEG) in PEG *via* a biotin-streptavidin-biotin bridge, and illuminated with the TIR field

generated at the slide/buffer interface. The instantaneous illumination power incident at the prism entry face was

40 mW (estimated density of 4.3 W/cm²), and the AOM was modulated with a 50% duty cycle at 1 MHz, such

that the time averaged power was 20 mW. Fluorescence was collected through the objective, separated with a

multi-wavelength imager (Photometrics, Dual-view), and imaged onto a water-cooled electron-multiplying charge

coupled device (Andor, iXon Ultra 888). Movies were collected for 2000 frames at 25 msec exposures with 2x

binning, 600 EM gain, 1x pre-amp gain, 30.0 MHz readout mode, 0.60 µs vertical speed, and +4 clock voltage

running at –67°C using the open source microscopy software Micro-Manger (Edelstein et al., 2014).

**smFRET Imaging Buffers.** VcINDY homodimers were imaged in the following buffers, which all contained 100

mM KCl, and 20 mM Tris-Cl (pH₂₅C=7.5). **Buffer LX**: 100 mM choline chloride. **Buffer LS**: 100 mM choline

chloride, and 1 mM succinic acid. **Buffer HX**: 100 mM NaCl. **Buffer HS**: 100 mM NaCl, and 1 mM succinic acid.

All buffers also contained 1% β-D-glucose, 1 mM trolox, 0.0025 U glucose oxidase, 0.02 U catalase, 1 mM

1,3,5,7-cyclooctatetraene, and 1 mM 3-nitrobenzyl alcohol as photostabilizing additives.

**smFRET Data Analysis.** Diffraction-limited spots of fluorescence intensity within the first 400 frames (10

seconds) were located in both the donor and acceptor fluorescence channels. Intensity versus time trajectories

were estimated using the iterative, maximum likelihood formulas for intensity estimation $N = \left(\sum_{xy}(d_{xy} - b)\Psi_{xy}\right)/$

$\left(\sum_{xy} \Psi_{xy}^2\right)$ and $b = \left(\sum_{xy}(d_{xy} - N\Psi_{xy})\right)/\left(\sum_{xy} 1\right)$, where (x,y) is a pixel location, N is the fluorescence intensity, b

is the background intensity, and Ψ is a fixed width, Gaussian-shaped point spread function with width defined by

the TIRF microscope optics. Donor and acceptor fluorescence channels were aligned using a 4th order

polynomial transform of ~2800 control points obtained from an image of an array of sub-diffraction limit

nanopatterned features localized in both channels. Trajectories were manually classified to select trajectories

from only donor-acceptor labeled molecules (see Fig. 1—Figure Supplement 1). The photobleaching point for

each $E_{FRET}$ trajectory was inferred using Bayesian inference (see below). A non-linear filter (Haran, 2004) was

applied to fluorescence intensity trajectories for easier visualization in intensity and $E_{FRET}$ trajectory plots, but all

analyses (*e.g.*, histograms, and HMMs) were performed using the unfiltered data.

**Global, variational Bayes hidden Markov modeling (vbHMM)**. Global vbHMMs were estimated in a manner

similar to a previously described single trajectory vbHMM implemented in the software package vbFRET

(Bronson et al., 2009). vbFRET is a Bayesian inference-based 'maximum evidence' method (Kinz-Thompson et

al., 2021), which allowed one to determine the simplest HMM that sufficiently accounts for the complexity of the

smFRET data (Bronson et al., 2009; van de Meent et al., 2014; Hon and Gonzalez, 2019). In this method, one

can find the most parsimonious HMM for smFRET data by estimating several HMMs, each with an increasing

number of hidden states, then calculating the 'evidence' for each HMM and choosing the one with the maximum

evidence value (Bronson et al., 2009). As seen in the directed acyclic graph (Fig. 4—Figure Supplement 1), the

only difference between vbFRET, and the global vbHMM used here is that an ensemble of $E_{FRET}$ versus time

trajectories is assumed to be independent and identically distributed (I.I.D.) from the same HMM parameters in

the global vbHMM model. Thus, all data points then contribute to the expectation and maximization equations

for the hyperparameters governing the posterior probability distribution, as opposed to only those from a single

trajectory is done for vbFRET. This is equivalent to considering one large, concatenated trajectory with vbFRET,

where however there is no conditional dependence for the hidden state on the previous hidden state for

neighboring data points from different trajectories. Instead, the first data point of each trajectory depends on the

initial-state probability distribution denoted here by π. As can be done with vbFRET, model selection can be

performed by calculating the evidence lower bound (ELBO) for each model and choosing that model with the

largest value.

To demonstrate the utility of using model selection with global vbHMMs, we simulated multiple four-state

time series, performed model selection, and compared this to an autocorrelation function (ACF) analysis (Fig.

4—Figure Supplement 2). The two-point autocorrelation function (Berne and Pecora, 2000) of a stationary

stochastic process or signal, y(t), describes how similar the process remains after some time lag, $\tau$, and can be

calculated as

573
$$ACF(\tau) \equiv \langle \delta y(0) \cdot \delta y(\tau) \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} d\delta y(0) \cdot d\delta y(t) \cdot y(0) \cdot y(\tau) \cdot P(\delta y(0), \delta y(\tau))$$

574 where $\delta y(\tau) = y(\tau) - \langle y \rangle$, angular brackets denote an expectation value, P(x) is the probability distribution

575 function of x, and

$$P\big(\delta y(0), \delta y(\tau)\big) = P\big(\delta y(\tau)|\delta y(0)\big) \cdot P\big(\delta y(0)\big).$$

577 For a discrete, zero-mean signal, A(t), composed of N data points, the ACF can be estimated as

$$ACF(\tau) \equiv \langle A(0)A(\tau) \rangle = \lim_{N \to \infty} \frac{1}{N} \sum_{i=1}^{N} A_i A_{i+\tau}$$

Thus, assuming normally distributed values of $A_i A_{i+\tau}$, the ACF and the associated precision is distributed as a

student-T distribution, $T_\nu(x|\mu, \sigma^2)$ with $\nu$ degrees of freedom, as $T_{N-1}(ACF(\tau)|\bar{x}, (\sum_i(x_i - \bar{x})^2)/(N(N-1)))$,

where $x_i = A_i A_{i+\tau}$, and $\bar{x} = \frac{1}{N} \sum_i A_i A_{i+\tau}$. This is derived from the posterior marginal of a normal-distribution

likelihood with a normal-gamma prior with $\kappa$=0, $\alpha$=−1/2, and $\beta$=0 (Murphy, 2007). The same formulas can be

used to calculate an ACF from multiple signal trajectories that are independent and identically distributed; for

instance the ACF at time $\tau$ would then include terms $A_i^0 A_{i+\tau}^0 + A_i^1 A_{i+\tau}^1 + \cdots + A_i^M A_{i+\tau}^M$, where the superscript

denotes an index over $M$ number of signal trajectories. We refer to this as the ensemble ACF.

For some specific types of signals, we can calculate an analytical form of the ACF for comparison to

observed ACFs. First, if a time-dependent signal was stationary at zero and distributed according to

$y(t) \sim \mathcal{N}(0, \sigma^2)$, where $\mathcal{N}$ denotes a normal distribution, and $\sigma^2$ is the variance, the $ACF(t) = \sigma^2 \cdot \delta(t)$, where

$\delta(t)$ is a Dirac delta function, which is one when t=0, and zero otherwise. Second, consider a zero-mean Markov

chain with N states that are normally distributed according to $\mathcal{N}(\mu_i, \sigma_i^2)$ for the i$^{th}$ state, but where $\langle y \rangle = 0$. Also,

consider that the rate constant for the transition from the i$^{th}$ state to the j$^{th}$ state, k$_{ij}$, can be written as a rate

constant matrix, K, with the associated Q matrix (K but where the on-diagonals are the negative sum of the rate

constants out the i$^{th}$ state). Notably, a vector of the probability of being in a particular state, P(t), given an initial

probability vector, P(0), can then be written $P(t) = e^{Qt} \cdot P(0)$, where $e$ denotes a matrix-exponential. If the

Markov chain is in state i, the signal value at a particular time can be written $y(t) \sim \mu_i + N_i$, where $N_i \sim \mathcal{N}(0, \sigma_i^2)$.

Thus, the ACF for this model is

$$\langle y(0)y(t) \rangle = \sum_i \sum_j \int_{N_i} \int_{N_j} (\mu_i + N_i)(\mu_j + N_j) \cdot P(j, t; i, 0; N_t; N_0),$$

598 where

$$P(j, t; i, 0; N_t; N_0) = P(j, t|i, 0) \cdot P_i(t = \infty) \cdot \mathcal{N}(0, \sigma_j^2) \cdot \mathcal{N}(0, \sigma_i^2).$$

599 Therefore

$$\langle y(0)y(t) \rangle = \sum_{i,j} \mu_i \mu_j \cdot d_j \cdot e^{Qt} \cdot d_i \cdot P_i(t = \infty) + \sum_i \sigma_i^2 \cdot P_i(t = \infty) \cdot \delta(t)$$

$$\equiv ACF_{MC}(t) + \langle \sigma^2 \rangle \cdot \delta(t),$$

where $d_i$ is a vector with a 1 in position i, and ACF$_{MC}$ is the contribution from the Markov chain. Additionally, we

note that $P(j, t|i, 0) \cdot P(i) = d_j \cdot e^{Qt} \cdot d_i \cdot P_i(t = \infty)$ can be rewritten with a spectral expansion as $\sum_i C_i e^{-\lambda_i t}$,

where $C_i$ is a weighting factor, and $\lambda_i$ is an eigenvalue of $e^{Qt}$. Thus, for this Markov chain model with normally
distributed noise, the ACF can be written as a weighted sum of *N* exponential decays with a term at t=0
corresponding to the average amount of uncorrelated, normally distributed noise. Therefore, for this model, the
difference between ACF(t=0) and the next data point ACF(t=dt) is mostly $\langle \sigma^2 \rangle$, and the rest of the ACF is
described by various exponential decays to zero.

In order to simulate single-molecule trajectories, we used the stochastic simulation algorithm (Gillespie,
1976) with the parameters in Table 1 with additive, normally distributed noise. From these 400 trajectories, we
estimated global HMMs with one through six states as described above. As expected, the HMM with four states
had the largest ELBO (Fig. 4–Figure Supplement 2). Additionally, we calculated the ACFs of these HMMs as
described above. Notably, of these ACFs, only those for HMMs with four or more states are sufficient to match
the individual and ensemble ACFs of the simulated trajectories (Fig. 4–Figure Supplement 2). Thus, the most
parsimonious global HMM determined by this method also recapitulates the kinetic results of a model-free ACF
method, but has the additional benefit as a model-based method of providing a mechanistic interpretation of
those results.

**Table 1.** $E_{FRET}$ trajectory simulation parameters

| Parameter | Value | Units |
|---|---|---|
| $\mu_1$ | 0.1 | A.U. |
| $\mu_2$ | 0.3 | A.U. |
| $\mu_3$ | 0.7 | A.U. |
| $\mu_4$ | 0.9 | A.U. |
| $k_{12}$ | 0.25 | $sec^{-1}$ |
| $k_{13}$ | 0.50 | $sec^{-1}$ |
| $k_{14}$ | 1.00 | $sec^{-1}$ |
| $k_{21}$ | 0.50 | $sec^{-1}$ |
| $k_{23}$ | 1.00 | $sec^{-1}$ |
| $k_{24}$ | 0.25 | $sec^{-1}$ |
| $k_{31}$ | 1.00 | $sec^{-1}$ |
| $k_{32}$ | 0.25 | $sec^{-1}$ |
| $k_{34}$ | 0.50 | $sec^{-1}$ |
| $k_{41}$ | 0.25 | $sec^{-1}$ |
| $k_{42}$ | 0.50 | $sec^{-1}$ |
| $k_{43}$ | 1.00 | $sec^{-1}$ |
| $\sigma$ | 0.20 | A.U. |
| $\tau$ | 0.01 | $sec$ |
| n | 2000 | frames |
| N | 400 | trajectories |

$\mu$ is an emission mean, $k$ is a rate constant, $\sigma$ is a standard deviation of additive noise, $\tau$ Is the time period of each data point, $n$ is the length of each trajectory, N is the number of trajectories

**Photobleaching detection**. We detect photobleaching by using shape-based Bayesian model selection to
choose the photobleaching time, or lack thereof, in a fluorescence intensity trajectory (Ray et al., 2021). Each
model, $M_T$, that we choose between is similar in that before some time, T, the data points are assumed to be
distributed according to a normal distribution with some mean, $\mu_1$, and some precision, $\lambda_1$ (i.e., inverse variance
$\lambda = \sigma^{-2}$), which we do not care about, while the data points after time T are distributed according to a normal
distribution with mean of zero, and some different precision, $\lambda_2$. We then calculate the probability of each $M_T$,

having marginalized out the means and precisions so that the conclusions are relevant regardless of those

values, and then use Bayesian inference to select the $M_T$ that gives the best estimate of the photobleaching time

as is supported by the available data. Practically, this method assigns all observed fluorescence at the beginning

of a trajectory into the same 'on' state, usually with a large variance to account for different intensity states that

compose the 'on' state, and then finds the last point where the data can be described as dropping to zero. For

an ensemble of intensity trajectories, we perform model selection twice. The first time, we use uniform priors for

the photobleaching time, while the second time, we use the independently calculated photobleaching times to

infer an ensemble photobleaching rate constant, and then repeat the photobleaching time model selection

calculation using this photobleaching rate constant to define the prior probability distribution for T for all of the

independent intensity trajectories.

Mathematically, the likelihood for one of these models $M_T$ is

636
$$P(\{x\}|\mu_1, \lambda_1, \lambda_2, T) = \prod_{t<T} \mathcal{N}(x_t|\mu_1, \lambda_1^{-1}) \prod_{t\geq T} \mathcal{N}(x_t|0, \lambda_2^{-1}),$$

637 where $\mathcal{N}(x|\mu, \sigma^2)$ is a normal distribution with mean μ, and variance σ². Using a prior probability distribution of

638
$$P(\mu_1, \lambda_1, \lambda_2) = NG(\mu_1, \lambda_1|\alpha_1, \beta_1, \kappa_1, m_1) \cdot G(\lambda_2|\alpha_2, \beta_2)$$

639
$$= \left[ (2\pi)^{-\frac{1}{2}} (\kappa_1 \lambda_1)^{\frac{1}{2}} e^{-\frac{\kappa_1 \lambda_1}{2}(\mu_1 - m_1)^2} \cdot \beta_1^{\alpha_1} \Gamma(\alpha_1)^{-1} \lambda_1^{\alpha_1-1} e^{-\beta_1 \lambda_1} \right] \cdot \left[ \beta_2^{\alpha_2} \Gamma(\alpha_2)^{-1} \lambda_2^{\alpha_2-1} e^{-\beta_2 \lambda_2} \right],$$

640 where NG is a joint normal-gamma distribution, and G is a gamma distribution,(Murphy, 2007) we obtain a

641 posterior probability distribution of the same form but with updated hyperparameters (denoted by primes)

642
$$P(\mu_1, \lambda_1, \lambda_2|\{x\}, T) = NG(\mu_1, \lambda_1|\alpha_1', \beta_1', \kappa_1', m_1') \cdot G(\lambda_2|\alpha_2', \beta_2'),$$

643 where

644
$$\alpha_1' = \alpha_1 + T/2$$

645
$$\beta_1' = \beta_1 + \frac{1}{2}\sum_{i=0}^{T}(x_i - \langle x \rangle)^2 + \frac{\kappa_1 T(\langle x \rangle - m_1)^2}{2(\kappa_1 + T)}$$

646
$$\kappa_1' = \kappa_1 + T$$

647
$$m_1' = \frac{\kappa_1 m_1 + T\langle x \rangle}{\kappa_1 + T}$$

648
$$\alpha_2' = \alpha_2 + (N - T)/2$$

649
$$\beta_2' = \beta_2 + \frac{1}{2}\sum_{i=T}^{N} x_i^2 ,$$

650 where N is the total number of data points. However, we do not particularly care about the exact values of $\mu_1$,

651 $\lambda_1$, or $\lambda_2$, so instead, we can marginalize them out to find the probability that of the observed data given the

652 model $M_T$ (i.e., the "evidence"), regardless of these nuisance parameters. In this case, we find

653
$$P(\{x\}|M_T) = P(\{x\}_{t<T}|M_T) \cdot P(\{x\}_{t\geq T}|M_T)$$

654
$$= \left[ \frac{\Gamma(\alpha_1')}{\Gamma(\alpha_1)} \cdot \frac{\beta_1^{\alpha_1}}{\beta_1'^{\alpha_1'}} \cdot \left(\frac{\kappa_1}{\kappa_1'}\right)^{\frac{1}{2}} \cdot (2\pi)^{-\frac{T}{2}} \right] \cdot \left[ \frac{\Gamma(\alpha_2')}{\Gamma(\alpha_2)} \cdot \frac{\beta_2^{\alpha_2}}{\beta_2'^{\alpha_2'}} \cdot (2\pi)^{-\frac{(N-T)}{2}} \right].$$

655  Now, with the evidence for the data given a photobleaching time, $P(\{x\}|M_T)$, we use Bayesian inference to select

656  best photobleaching model (i.e., the model best supported by the data, regardless of the individual parameters)

657  as the largest value of

658
$$P(M_T|\{x\}) = \frac{P(\{x\}|M_T) \cdot P(M_T)}{\sum_{T=0}^{N} P(\{x\}|M_T) \cdot P(M_T)}.$$

As discussed above, $P(M_T)$ is at first taken to be a constant and all trajectories are analyzed independently.

Then, we infer the ensemble photobleaching rate constant using Bayesian inference with an exponential

distribution as a likelihood, and a gamma distribution as the conjugate prior, and therefore also the posterior. As

a result, we estimate $k_{PB} = (1 + n)/(1 + \sum_{i=0}^{n} T_i)$, where $i$ indexes the different trajectories. Then, we use an

exponential distribution as the prior $P(M_T) = k_{PB} e^{-k_{PB} T}$, where k$_{PB}$ is treated as a deterministic value. Finally,

we note that these model selection calculations are performed in log-space, and that we subtract the maximum

of all the log terms (i.e., scale by a constant) from both the numerator and denominator in order the to avoid

computer underflow.

Since the models used for model selection in this algorithm are not comprehensive in the sense that they

do not account for every experimental nuance of the data, we believe it is best to manually examine the selected

model and the underlying fluorescence intensity trajectory to ensure that they are reasonably well matched, and

so that one can manually adjust the photobleaching time when they do not match due to unaccounted for

experimental behaviors (*e.g.*, deviations from Gaussian emissions).

**Calculation of Cycle Fluxes.** For the mechanistic scheme for transport in Fig. 5, the following terms are

necessary to calculate cycle fluxes to be calculated. For shorter notation, we use $t \equiv k_t$, $d' \equiv k_d'$, $d'' = k_d''$, $A_o \equiv$

$k_a'[A]_o^m$, $A_i \equiv k_a'[A]_i^m$, $B_o \equiv k_a''[B]_o^n$, and $B_i \equiv k_a''[B]_i^n$. The cycle flux for any cycle is

675
$$J_\kappa = J_{\kappa+} - J_{\kappa-} = (\Pi_{\kappa+} - \Pi_{\kappa-}) \Sigma_\kappa / \Sigma,$$

676  where

677
$$\Pi_{a+} - \Pi_{a-} = t^2 d'(A_o - A_i),$$

678
$$\Pi_{b+} - \Pi_{b-} = t^2 d''(B_o - B_i),$$

679
$$\Pi_{c+} - \Pi_{c-} = t^2 d' d''(A_o B_o - A_i B_i),$$

680
$$\Sigma_a = 2t d'' + d''^2,$$

681
$$\Sigma_b = t(A_o + A_i) + A_o A_i,$$

682
$$\Sigma_c = 1, \text{ and}$$

$$\Sigma = 2t^3[d''(4d' + 2A_i + A_o) + (2A_iB_i + A_oB_o + A_oB_i + B_oA_i)]$$

$$+ 2t^2[d''^2(2d' + A_i + A_o)$$

$$+ d''((2(A_o + A_i) + (B_o + B_i) + 2d')d' + (A_iB_i + B_oA_i + 2A_oA_i + A_oB_o + A_oB_i))$$

$$+ ((A_oB_o + A_iB_i)d' + (A_oB_oA_i + A_oA_iB_i + A_oB_oB_i + B_oA_iB_i))]$$

$$+ 2t[d''^2((A_o + A_i)d' + A_oA_i + d'^2) + d''((A_oB_o + A_iB_i)d' + (A_oB_oA_i + A_oA_iB_i)) + (A_oB_oA_iB_i)].$$

From these equations, the functional form of $J_c$ for each first order rate constant, $k$, in Fig. 5 is $[ak + b + ck^{-1}]^{-1}$

where the coefficients *a*, *b*, and *c* depend on the other rate constants. This functional form has a maximum at

$k = \sqrt{c/a}$, so there is an optimal value of $k_t$ that maximizes transport cycle flux under for a given set of substrate

concentrations. Finally, we note that $\Sigma$ has 90 terms, and thus so does $J_c$. Due to the changes to the energy

landscape in Fig. 6, 68 of the terms in $J_c$ increase the magnitude of $J_c$, 18 decrease it, and 4 do not change it.

To find the optimal parameters to describe previously published succinate transport data, a target energy

function of $(J^{expt} - (J_b + J_c))^2$ was used for $J^{expt}([X] = K_M) = \frac{V_{max}}{2}$ and $J^{expt}([X] = 1000K_M) = V_{max}$ for the

sodium titration in their Figure 3 ($K_M = 41.7\ mM$, and $V_{max} = 53.5\ nmol\ mg^{-1}min^{-1} = 58.3\ s^{-1}$), and the

succinate titration in their Figure 6 ($K_M = 1.0\ \mu M$, and $V_{max} = 232.6\ nmol\ mg^{-1}min^{-1} = 253.4\ s^{-1}$) (Mulligan et

al., 2014). The collective energy function for both experiments was minimized using an affine invariant MCMC

ensemble sampler (Foreman-Mackey et al., 2013) with 200 walkers, a 1,000 step burn-in period, and a 10,000

step production period from which every 100[th] point was taken as a sample to avoid autocorrelation within the

samples.

The steady-state probability of apo VcINDY in Buffer HX was calculated from the mechanism in Fig. 5 by

setting $[B]_o = [B]_i = 0$, using the estimates for $k_a'$, $\langle k \rangle \equiv k_t$, and $k_d'$ (Fig. 5—Figure Supplement 2) and $[A]_o =$

$[A]_i = [Na^+] = 0.1M$, to calculate the steady state probability $P_{apo} = P_{C_o} + P_{C_i}$ as

$$P_{apo} = \frac{3\langle k \rangle^3 + k_a'[Na^+]^3\langle k \rangle^2}{3\langle k \rangle^3 + 4k_a'[Na^+]^3\langle k \rangle^2 + k_a'^2[Na^+]^6\langle k \rangle}.$$

**Crystallization of Wild Type VcINDY with Substrates**. Crystals of VcINDY were grown with a protein

concentration of 6 mg/ml at 4°C in hanging-drop vapor diffusion. For VcINDY-succinate, crystallization was

carried out with 30% PEG 300, 5 mM sodium succinate, 200 mM sodium chloride, and 100 mM MES pH 5.8.

VcINDY-fumarate crystals were grown with 37.5% PEG 300, 200 μM sodium fumarate, and 100 mM sodium

acetate pH 4.6. For VcINDY-malate, crystals were grown with 25% PEG 400, 20 mM sodium malate, and 100

mM sodium acetate pH 5.0.

**X-ray Data Collection and Structure Determination**. Crystals were frozen in liquid nitrogen with the

crystallization solution serving as the cryoprotectant. X-ray data were collected at the Advanced Photon Source

Beamlines 19-ID and 19-BM. Data processing and scaling were performed using the HKL2000 software

(Otwinowski and Minor, 1997). All crystals were of space group P2₁ with unit cell dimensions around a = 106 Å,

b = 102 Å, c = 171 Å, β = 99°, and contained four molecules per asymmetric unit. The structures were determined
by molecular replacement using the structure of VcINDY (PDB: 5UL9) (Nie et al., 2017) with sodium and citrate
removed as the initial search model, followed by repeated cycles of model building in Coot (Emsley and Cowtan,
2004), and refinement in Phenix (Table 2) (Adams et al., 2010).

**Data and Code Availability.** VcINDY structures with fumarate, malate, and succinate bound were deposited in
the PDB with IDs 6OKZ, 6OL0, and 6OL1, respectively. Code used to process smFRET data is described in a
separate manuscript (in preparation, C.D.K. and R.L.G.).

**Table 2.** X-ray crystallography of VcINDY in complex with substrate

|  | Parameter | VcINDY-succinate | VcINDY-fumarate | VcINDY-malate |
|---|---|---|---|---|
| Data collection | Wavelength (Å) | 0.9791 | 0.9794 | 0.9794 |
|  | Resolution range (Å) | 50.0-3.09 (3.15-3.09) | 50-3.29 (3.35-3.29) | 50.0-3.50 (3.56-3.50) |
|  | Space group | $P2_1$ | $P2_1$ | $P2_1$ |
|  | Cell dimensions: a, b, c (Å) | 105.5, 102.5, 171.1 | 105.8, 102.1, 171.2 | 106.8, 102.4, 171.8 |
|  | Cell angle: $\beta$ (°) | 98.808 | 98.937 | 97.791 |
|  | Total reflections | 128,621 (6,360) | 99,521 (3,598) | 86,871 (4,328) |
|  | Unique reflections | 65,778 (3,229) | 51,510 (1,913) | 44,866 (2,248) |
|  | Multiplicity | 5.0 (5.1) | 6.6 (5.3) | 6.9 (6.4) |
|  | Completeness (%) | 99.9 (100) | 93.9 (71.0) | 96.4 (96.9) |
|  | $I/\sigma(I)$ | 10.3 (0.625) | 10.1 (0.667) | 10.8 (0.571) |
|  | Wilson B (Å$^2$) | 80.97 | 110.64 | 134.20 |
|  | $R_{pim}$ | 8.4 (152) | 4.4 (54.2) | 3.5 (61.5) |
|  | $CC_{1/2}$ | 0.998 (0.443) | 0.102 (0.512) | 0.997 (0.604) |
| Refinement | $R_{work}/R_{free}$ (%) | 25.3% / 29.6% | 24.9% / 28.5% | 26.1% / 30.5% |
|  | No. atoms - Protein/Ligands | 13,325 / 40 | 13,097 / 23 | 12,792 / 42 |
|  | RMSD bond lengths (Å) | 0.003 | 0.003 | 0.001 |
|  | RMSD bond angles (°) | 0.66 | 0.52 | 0.39 |
|  | Ramachandran favored (%) | 88.7 | 91.3 | 93.2 |
|  | Ramachandran allowed (%) | 9.99 | 7.30 | 5.72 |
|  | Ramachandran outliers (%) | 1.35 | 1.44 | 1.12 |
|  | Rotamer outliers (%) | 0.07 | 0.0 | 0.0 |
|  | Clashscore | 34.26 | 29.10 | 23.49 |

Numbers in parentheses are statistics for the highest resolution shell.

# 722 CRediT Authorship Contribution Statement

723 **Colin D. Kinz-Thompson:** Conceptualization, Methodology, Resources, Investigation, Validation, Formal
724 Analysis, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization. **Marisa Lopez-**
725 **Redondo:** Methodology, Resources, Investigation, Validation. **Christopher Mulligan:** Investigation, Validation,
726 Formal Analysis, Visualization. **David B. Sauer:** Investigation, Validation, Formal Analysis, Data Curation,

Visualization. **Jennifer J. Marden:** Resources, Validation. **Jinmei Song:** Resources, Validation. **Emad Tajkhorshid:** Conceptualization, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition. **John F. Hunt:** Conceptualization, Methodology, Resources, Supervision, Funding Acquisition. **David L. Stokes:** Conceptualization, Methodology, Resources, Supervision, Funding Acquisition. **Joseph A. Mindell:** Conceptualization, Methodology, Resources, Data Curation, Writing - Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition. **Da-Neng Wang:** Conceptualization, Methodology, Resources, Data Curation, Writing - Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition. **Ruben L. Gonzalez, Jr.:** Conceptualization, Methodology, Resources, Data Curation, Writing - Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition.

# Acknowledgements

R.L.G. acknowledges support from the National Institutes of Health (NIH) through grants R01-GM084288 and R01-GM137608. D.N.W. acknowledges support from the NIH through grants R01-GM121994, R01-NS108151, and R01-DK099023, the G. Harold and Leila Y. Mathers Foundation, and the TESS Research Foundation. J.M. and C.M. acknowledge support from the National Institute of Neurological Disorders and Stroke (NINDS) Intramural Program, and C.M. acknowledges support from the Biotechnology and Biological Sciences Research Council through grant BB/V007424/1. D.L.S. acknowledges support from NIH through grant R35-GM144109. J.F.H. acknowledges support from NIH through grants R01-GM127883. E.T. acknowledges support from NIH through grants P41-GM104601 and R01-GM123455. D.B.S. acknowledges support, in part, from an American Cancer Society Postdoctoral Fellowship (PF-17-135-01) and from the Office of the Assistant Secretary of Defense for Health Affairs, through the Peer Reviewed Cancer Research Program under Award No. W81XH-16-1-0153. Crystal screening and X-ray data collection were carried out at NSLS-II Brookhaven National Laboratory and at Argonne National Laboratory, Structural Biology Center (SBC) at the Advanced Photon Source. SBC-CAT is operated by Univ. Chicago Argonne, LLC, for the U.S. Department of Energy, Office of Biological and Environmental Research under contract DE-AC02-06CH11357. Electron microscopy was performed at the NYU School of Medicine Microscopy Core.

# References

Adams PD, Afonine PV, Bunkóczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung L-W, Kapral GJ, Grosse-Kunstleve RW, McCoy AJ, Moriarty NW, Oeffner R, Read RJ, Richardson DC, Richardson JS, Terwilliger TC, Zwart PH. 2010. PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**:213–221. doi:10.1107/S0907444909052925

Akyuz N, Altman RB, Blanchard SC, Boudker O. 2013. Transport dynamics in a glutamate transporter homologue. *Nature* **502**:114–118. doi:10.1038/nature12265

Bayburt TH, Sligar SG. 2010. Membrane protein assembly into Nanodiscs. *FEBS Lett* **584**:1721–1727. doi:10.1016/j.febslet.2009.10.024

Bergeron MJ, Clémençon B, Hediger MA, Markovich D. 2013. SLC13 family of $Na^+$-coupled di- and tri-carboxylate/sulfate transporters. *Mol Aspects Med* **34**:299–312. doi:10.1016/j.mam.2012.12.001

Berne BJ, Pecora R. 2000. Dynamic Light Scattering: With Applications to Chemistry, Biology, and Physics,
Unabridged edition. ed. Mineola, N.Y: Dover Publications.
Bronson JE, Fei J, Hofman JM, Gonzalez RL, Wiggins CH. 2009. Learning rates and states from biophysical
time series: a Bayesian approach to model selection and single-molecule FRET data. *Biophys J* **97**:3196–
205. doi:10.1016/j.bpj.2009.09.031
Denisov IG, Grinkova YV, Lazarides AA, Sligar SG. 2004. Directed Self-Assembly of Monodisperse Phospholipid
Bilayer Nanodiscs with Controlled Size. *J Am Chem Soc* **126**:3477–3487. doi:10.1021/ja0393574
Denisov IG, Sligar SG. 2017. Nanodiscs in Membrane Biochemistry and Biophysics. *Chem Rev* **117**:4669–4713.
doi:10.1021/acs.chemrev.6b00690
Drew D, Boudker O. 2016. Shared Molecular Mechanisms of Membrane Transporters. *Annu Rev Biochem*
**85**:543–72. doi:10.1146/annurev-biochem-060815-014520
Edelstein AD, Tsuchida MA, Amodaj N, Pinkard H, Vale RD, Stuurman N. 2014. Advanced methods of
microscope control using µManager software. *J Biol Methods* **1**:10. doi:10.14440/jbm.2014.36
Emsley P, Cowtan K. 2004. Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr*
**60**:2126–2132. doi:10.1107/S0907444904019158
Erkens GB, Hänelt I, Goudsmits JMH, Slotboom DJ, van Oijen AM. 2013. Unsynchronised subunit motion in
single trimeric sodium-coupled aspartate transporters. *Nature* **502**:119–123. doi:10.1038/nature12538
Felle H, Porter JS, Slayman CL, Kaback HR. 1980. Quantitative measurements of membrane potential in
Escherichia coli. *Biochemistry* **19**:3585–3590. doi:10.1021/bi00556a026
Fersht A. 2017. Structure and Mechanism in Protein Science: A Guide to Enzyme Catalysis and Protein Folding.
WORLD SCIENTIFIC. doi:10.1142/10574
Fitzgerald GA, Mulligan C, Mindell JA. 2017. A general method for determining secondary active transporter
substrate stoichiometry. *eLife* **6**:e21016. doi:10.7554/eLife.21016
Foreman-Mackey D, Hogg DW, Lang D, Goodman J. 2013. emcee: The MCMC Hammer. *Publ Astron Soc Pac*
**125**:306–312. doi:10.1086/670067
Forrest LR, Krämer R, Ziegler C. 2011. The structural basis of secondary active transport mechanisms. *Biochim*
*Biophys Acta Bioenerg* **1807**:167–188. doi:10.1016/j.bbabio.2010.10.014
Garaeva AA, Slotboom DJ. 2020. Elevator-type mechanisms of membrane transport. *Biochem Soc Trans*
**48**:1227–1241. doi:10.1042/BST20200290
Gillespie DT. 1976. A general method for numerically simulating the stochastic time evolution of coupled
chemical reactions. *J Comput Phys* **22**:403–434. doi:10.1016/0021-9991(76)90041-3
Haran G. 2004. Noise reduction in single-molecule fluorescence trajectories of folding proteins. *Chem Phys*
**307**:137–145. doi:10.1016/j.chemphys.2004.05.017
Hill TL. 2005. Free energy transduction and biochemical cycle kinetics. Mineola, N.Y: Dover Publications.
Hon J, Gonzalez RL. 2019. Bayesian-Estimated Hierarchical HMMs Enable Robust Analysis of Single-Molecule
Kinetic Heterogeneity. *Biophys J* **116**:1790–1802. doi:10.1016/j.bpj.2019.02.031
Jardetzky O. 1966. Simple allosteric model for membrane pumps. *Nature* **211**:969–970. doi:10.1038/211969a0
Kim JW, Kim Subin, Kim Songwon, Lee H, Lee J, Jin MS. 2017. Structural insights into the elevator-like
mechanism of the sodium / citrate symporter CitS. *Sci Rep* **7**:2548. doi:10.1038/s41598
Kinz-Thompson CD, Ray KK, Gonzalez RL. 2021. Bayesian Inference: The Comprehensive Approach to
Analyzing Single-Molecule Experiments. *Annu Rev Biophys* **50**:191–208. doi:10.1146/annurev-biophys-
082120-103921
Lambert TJ. 2019. FPbase: a community-editable fluorescent protein database. *Nat Methods* **16**:277–278.
doi:10.1038/s41592-019-0352-8
Lo C-J, Leake MC, Berry RM. 2006. Fluorescence Measurement of Intracellular Sodium Concentration in Single
Escherichia coli Cells. *Biophys J* **90**:357–365. doi:10.1529/biophysj.105.071332
Lolkema JS, Slotboom DJ. 2019. Models to determine the kinetic mechanisms of ion-coupled transporters. *J*
*Gen Physiol* **151**:369–380. doi:10.1085/jgp.201812055
Maity S, Trinco G, Buzón P, Anshari ZR, Kodera N, Ngo KX, Ando T, Slotboom DJ, Roos WH. 2022. High-speed
atomic force microscopy reveals a three-state elevator mechanism in the citrate transporter CitS. *Proc*
*Natl Acad Sci* **119**:e2113927119. doi:10.1073/pnas.2113927119

Mancusso R, Gregorio GG, Liu Q, Wang D-N. 2012. Structure and mechanism of a bacterial sodium-dependent
dicarboxylate transporter. *Nature* **491**:1–6. doi:10.1038/nature11542

Markovich D. 2012. Sodium-Sulfate/Carboxylate Cotransporters (SLC13)Current Topics in Membranes.
Elsevier. pp. 239–256. doi:10.1016/B978-0-12-394316-3.00007-7

Mulligan C, Fenollar-Ferrer C, Fitzgerald GA, Vergara-Jaque A, Kaufmann D, Li Y, Forrest LR, Mindell JA. 2016.
The bacterial dicarboxylate transporter VcINDY uses a two-domain elevator-type mechanism. *Nat Struct*
*Mol Biol* **23**:256–263. doi:10.1038/nsmb.3166

Mulligan C, Fitzgerald GA, Wang D-N, Mindell JA. 2014. Functional characterization of a Na$^+$-dependent
dicarboxylate transporter from Vibrio cholerae. *J Gen Physiol* **143**:745–759. doi:10.1085/jgp.201311141

Murphy KP. 2007. Conjugate Bayesian analysis of the Gaussian distribution (Technical Report). University of
British Columbia.

Nie R, Stark S, Symersky J, Kaplan RS, Lu M. 2017. Structure and function of the divalent anion/Na$^+$ symporter
from Vibrio cholerae and a humanized variant. *Nat Commun* **8**:15009. doi:10.1038/ncomms15009

Otwinowski Z, Minor W. 1997. Processing of X-ray diffraction data collected in oscillation mode. *Methods*
*Enzymol* **276**:307–326.

Ray KK, Verma AR, Gonzalez Jr RL, Kinz-Thompson CD. 2021. Inferring the shape of data: A probabilistic
framework for analyzing experiments in the natural sciences. *ArXiv210912462 Phys Q-Bio*.

Ruan Y, Miyagi A, Wang X, Chami M, Boudker O, Scheuring S. 2017. Direct visualization of glutamate
transporter elevator mechanism by high-speed AFM. *Proc Natl Acad Sci* **114**:1584–1588.
doi:10.1073/pnas.1616413114

Sampson CDD, Fàbregas Bellavista C, Stewart MJ, Mulligan C. 2021. Thermostability-based binding assays
reveal complex interplay of cation, substrate and lipid binding in the bacterial DASS transporter, VcINDY.
*Biochem J* **478**:3847–3867. doi:10.1042/BCJ20210061

Sampson CDD, Stewart MJ, Mindell JA, Mulligan C. 2020. Solvent accessibility changes in a Na$^+$-dependent
C4-dicarboxylate transporter suggest differential substrate effects in a multistep mechanism. *J Biol Chem*
**295**:18524–18538. doi:10.1074/jbc.RA120.013894

Sauer DB, Marden JJ, Sudar JC, Song J, Mulligan C, Wang D-N. 2022a. Structural basis of ion – substrate
coupling in the Na$^+$-dependent dicarboxylate transporter VcINDY. *Nat Commun* **13**:2644.
doi:10.1038/s41467-022-30406-4

Sauer DB, Song J, Wang B, Hilton JK, Karpowich NK, Mindell JA, Rice WJ, Wang D-N. 2021. Structure and
inhibition mechanism of the human citrate transporter NaCT. *Nature* **591**:157–161. doi:10.1038/s41586-
021-03230-x

Sauer DB, Trebesch N, Marden JJ, Cocco N, Song J, Koide A, Koide S, Tajkhorshid E, Wang D-N. 2020.
Structural basis for the reaction cycle of DASS dicarboxylate transporters. *eLife* **9**:e61350.
doi:10.7554/eLife.61350

Sauer DB, Wang B, Sudar JC, Song J, Marden J, Rice WJ, Wang D. 2022b. The ups and downs of elevator-
type di-/tricarboxylate membrane transporters. *FEBS J* **289**:1515–1523. doi:10.1111/febs.16158

Stein WD, Lieb WR. 1986. Transport and diffusion across cell membranes. Orlando: Academic Press.

van de Meent J-W, Bronson JE, Wiggins CH, Gonzalez RL. 2014. Empirical Bayes Methods Enable Advanced
Population-Level Analyses of Single-Molecule FRET Experiments. *Biophys J* **106**:1327–1337.
doi:10.1016/j.bpj.2013.12.055

 Figures

**Figure 1.** VcINDY transport cycle. (A) Diagram of VcINDY dimer (green) reversibly transporting sodium ions ($Na^+$) and dicarboxylate substrate ($D^{2-}$) into and out of a cell with a stoichiometry of 3:1; an inward facing $Na^+$ electrochemical gradient provides the free energy for vectorial transport. (B) Minimal mechanistic model of VcINDY transport cycle. Outward- and inward-facing conformations of the VcINDY homodimer ($C_o$, and $C_i$, respectively) bind $Na^+$ and $D^{2-}$ in an unknown order along the horizontal transitions; double-prime signifies both co-substrates are bound. VcINDY protomer conformational motions along the vertical transitions are involved in substrate translocation. At least one closed transport cycle, such as shown here in the clockwise direction in blue, is required for vectorial transport. Slippage reactions occur when an incomplete set of co-substrates is moved across the membrane. (C) Cartoon diagram of smFRET pairs designed to report on the VcINDY translocation reaction for VcINDY$^{ext}$ (left; G211) and VcINDY$^{int}$ (right; S436). The distance between each labeling position, $r$, was determined from the $C_\alpha$-$C_\alpha$ distance, and was measured for a $C_o$, substrate-bound repeat-swap model ($C_o$-$C_o$, PMDB: PM0080216), and a substrate-bound, $C_i$ X-ray crystallography structure ($C_i$-$C_i$, PDB: 5UL9). $E_{FRET}$ was estimated using $R_0$=51 Å. Scaffold domains are shown in gray, and transport domains are shown in green.

**Figure 1–Figure Supplement 1.** Cartoon schematic of approach to yield donor- and acceptor- fluorophore-labeled VcINDY homodimers. Site-directed mutagenesis is used to remove all codons encoding cysteine from a plasmid-based "wild type" VcINDY gene (1) to yield a "cys-less" variant (2), and used further to introduce a single codon encoding cysteine (3). This is over-expressed in *Escherichia coli* (*E. coli*), and purified to yield homodimers with single-cysteine containing protomers (4). The homodimers are labeled with a 1:1 ratio of maleimide-conjugated donor- and acceptor-fluorophores (5). Only those homodimers labeled with both a single donor- and single acceptor-fluorophore are computationally selected after imaging (6).

**Figure 1–Figure Supplement 2.** VcINDY biochemical transport assays. Transport rates of proteoliposome-embedded VcINDY homodimers for the external (left) and internal (right) smFRET pair labeled positions. Wild type (*i.e.*, cysteine-containing) assays were performed as control separately for each labeling position.

**Figure 2.** Schematic diagram of nanodisc-embedded VcINDY homodimer construct. Single cysteine mutant VcINDY labeled with a 1:1 ratio of donor and acceptor fluorophores is mixed with biotinylated membrane scaffold protein (MSP) and *E. coli* polar lipids to make nanodisc-embedded dimers. VcINDY containing nanodiscs are then tethered to a flow cell surface *via* a biotin-streptavidin-biotin bridge, and imaged using TIRF microscopy. The scaffold domains are shown in gray, and transport domains are shown in green.

**Figure 2–Figure Supplement 1.** Negative stain electron microscopy of VcINDY reconstituted into lipid nanodiscs. Both top and side views of nanodiscs were seen. The nanodiscs had a diameter of ~108 Å and each contained one VcINDY dimer. The sample was uniform and no aggregation was observed.

**Figure 3.** Equilibrium smFRET from VcINDY$^{ext}$ (left) and VcINDY$^{int}$ (right) in the presence of sodium and succinate (buffer HS). Example donor (green) and acceptor (red) fluorescence intensity trajectories, with corresponding E$_{FRET}$ trajectories (blue) and idealized trajectories (black) are shown. Several trajectories shown include donor fluorescence intensity after the acceptor fluorophore has photobleached, but this was removed prior to further analysis.

**Figure 4.** Substrate dependence of VcINDY $E_{FRET}$ distributions. Histograms of $E_{FRET}$ values from all external (left) and internal (right) smFRET signals recorded with VcINDY[ext], and VcINDY[int], respectively, in different buffer conditions. The $E_{FRET}$ data in these histograms were not filtered, and $E_{FRET}$ values corresponding to donor photobleaching have been removed (*i.e.*, the last dwell in an $E_{FRET}$ state of zero). The predictive posterior probability distribution for $E_{FRET}$ from the global HMM with the maximum ELBO is shown overtop (solid black) as a heuristic measure of how well the global HMM explains the entirety of each dataset, but this ignores the kinetic aspects of the HMM. The distributions for the individual, consensus $E_{FRET}$ states from the global HMM are shown as black dashed lines.

**Figure 4–Figure Supplement 1.** Directed acyclic graph for a global, variational Bayes HMM. Pink nodes represent hyperparameters, grey nodes represent unobserved variables, and blue nodes represent observed variables. $T_j$ is the number of data points in the $j^{th}$ trajectory, and $D_{i,j}$ corresponds to the $i^{th}$ data point of the $j^{th}$ trajectory. Arrows represent conditional dependencies.

**Figure 4–Figure Supplement 2.** Autocorrelation function analysis of global HMMs with various number of hidden states of simulated, four-state $E_{FRET}$ trajectories. Individual ACFs (black), ensemble ACF (blue), and global HMM ACF (red). The maximum ELBO HMM (n=4; ln(ELBO)= −1.1x10⁴) matches the number of $E_{FRET}$ states in the simulation.

**Figure 4–Figure Supplement 3.** Substrate dependence of evidence lower bound (ELBO) calculations for global HMMs of VcINDY smFRET experiments. The maximum ELBO model is the HMM that best represents the set of $E_{FRET}$ trajectories for those substrate conditions.

**Figure 4–Figure Supplement 4.** Substrate dependence of the HMM with maximum ELBO for global HMMs of VcINDY smFRET experiments. For each kinetic scheme, states are denoted by the mean $E_{FRET}$ value of that state, which was inferred from the HMM. Rate constants for transitions between states are shown above the arrows.

**Figure 5.** Free energy transduction model for a secondary active transporter. First order and pseudo-first order rate constants for the parsimonious kinetic model discussed in the text are given over each transition. For simplicity, co-substrate binding has been reduced to a strictly sequential, two-step process with mirror symmetry (Stein and Lieb, 1986). Substrate A is the first co-substrate to bind with stoichiometry $m$; co-substrate B binds second with stoichiometry $n$. Prime signifies the first co-substrate is bound; double-prime signifies both co-substrates are bound; $C_o$ and $C_i$ denote the outward and inward facing conformation of the transporter. $k_t$ is the translocation rate constant, $k_a$ is a second order association rate constant, and $k_d$ is a dissociation rate constant. Cycles *a* (red) and *b* (green) are responsible for slippage, while cycle *c* (blue) is responsible for transport. The positive direction is defined to be clockwise.

**Figure 5—Figure Supplement 1.** MCMC trajectories for optimal kinetic model parameters of VcINDY proteoliposome-based succinate transport assay. (A) Affine invariant ensemble MCMC sampler trajectories for 200 walkers during burn-in period. These samples were discarded. (B) Production period trajectories from 200 walkers. Only every 100th sample of those shown here was used to minimize autocorrelation.

**Figure 5—Figure Supplement 2.** Corner plot of MCMC samples for optimal kinetic model parameters of VcINDY proteoliposome-based succinate transport assay. Ranges given are the $1\sigma$ level containing 39.3% of the sample volume.

**Figure 5—Figure Supplement 3.** Modeled VcINDY transport rate experiments. Data from Mulligan and coworkers were reproduced from their reported $K_M$, $V_{max}$, and Hill coefficients (Mulligan et al., 2014). Modeled data are the 95% credible intervals generated from MCMC samples of $k_a'$, $k_a''$, and $\langle k \rangle$, and dashed lines are the lowest energy solution.

**Figure 6.** Schematic of optimized energy landscape for binding co-substrates A and B. Original energy levels are denoted with dashed lines for apo- ($C$), partially bound ($C'$), and fully bound ($C''$) transporter. Direction of perturbations to energy levels by amounts $\Delta G'$ and $\Delta G''$ are shown with arrows. Relative positioning of unperturbed $C$, $C'$, and $C''$, and transition states $\ddagger'$ and $\ddagger''$ are arbitrarily chosen, as are the magnitudes of $\Delta G'$ and $\Delta G''$.

**Figure 6—Figure Supplement 1.** Estimated physiological Na$^+$ and succinate concentrations in a bacterial cell in different environments. External (ext) and internal (int) Na$^+$ concentrations were measured for another gammaproteobacteria, *E. coli*, at pH=7 (Lo et al., 2006). For *E. coli* at pH=7, the membrane potential difference ($\Delta\psi$) is approximately -100 mV (Felle et al., 1980). With these numbers, balancing the chemical potential difference and effects of charge for the VcINDY transport reaction at equilibrium provides the internal to external ratio of succinate concentrations for the cell (orange). When $\Delta\psi = 0$, the cell is depolarized (dashed).

**Figure 7.** View of the VcINDY binding pocket from the cytosol. The X-ray crystallography derived structural model is of succinate- and Na$^+$-bound VcINDY (green). Electron density for the substrates is shown contoured at 2.5σ.