Elevator mechanism dynamics in a sodium-coupled dicarboxylate transporter

Kinz-Thompson, C.D.,[1,†] Lopez Redondo, M.,[2] Mulligan C.,[3] Sauer, D.B.,[2,‡] Marden, J.J.,[2] Song, J.,[2] Tajkhorshid, E.,[4] Hunt, J.F.,[5] Stokes, D.L.,[2] Mindell, J.A.,[6*] Wang, D.N,[2*] Gonzalez, Jr., R.L.[1*]

1. Department of Chemistry, Columbia University, New York, New York 10027, US
2. Department of Cell Biology, New York University School of Medicine, New York, New York 10016, US
3. School of Biosciences, University of Kent, Canterbury, Kent, CT2 7NH, UK
4. Theoretical and Computational Biophysics Group, NIH Center for Macromolecular Modeling and Bioinformatics, Beckman Institute for Advanced Science and Technology, Department of Biochemistry, Center for Biophysics and Quantitative Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, US
5. Department of Biological Sciences, Columbia University, New York, New York 10027, US
6. Membrane Transport Biophysics Section, Porter Neuroscience Research Center, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD 20892, US

*Correspondence: rlg2118@columbia.edu, da-neng.wang@med.nyu.edu, mindellj@ninds.nih.gov
†Current address: Department of Chemistry, Rutgers University-Newark, Newark, New Jersey 07012, US
‡Current address: Centre for Medicines Discovery, Nuffield Department of Medicine, University of Oxford, Oxford, UK, OX1 3QU

Abstract

VcINDY, the sodium-dependent dicarboxylate transporter from *Vibrio cholerae*, is responsible for C_4- and C_5-carboxylate uptake into cells. The molecular mechanism of how VcINDY physically moves substrates across the membrane, and does so in an energetically efficient manner, is unclear. Here, we use single-molecule fluorescence resonance energy transfer experiments to directly observe the individual mechanistic steps that VcINDY takes to translocate substrates across a lipid bilayer, and then test key predictions of transport cycle mechanistic models. Our data provide the first direct evidence that VcINDY undergoes stochastic, elevator-type conformational motions that enable substrate translocation. Kinetic analysis suggests that the two protomers of the VcINDY homodimer undergo those motions in a non-cooperative manner, and thus catalyze two independent transport reactions. The relative substrate independence of those motions provides evidence that the VcINDY transport cycle maintains strict co-substrate coupling using a cooperative binding mechanism. Finally, thermodynamic modeling provides insight into how such a cooperative binding mechanism provides a generalized approach to optimizing transport for many secondary active transporters.

Introduction

Secondary active transporters are transmembrane proteins that harvest free energy from electrochemical gradients to drive the movement of substrates across a membrane (Fig. 1) (Stein and Lieb, 1986). In the absence of a free energy source, these transporters move substrates back and forth across the membrane without creating a net change in substrate concentration (Jardetzky, 1966). Net transport occurs because the transporter can dissipate the electrochemical gradient of a 'driving' substrate to harvest free energy, and then couple that free energy to the 'uphill' movement of a 'driven' substrate across the membrane (Forrest et al., 2011). The mechanistic details of this coupling are governed by the transport cycle, which describes how the transporter binds its substrates, translocates them across the membrane, releases them, and resets itself for the next round of transport (Fig. 1B). While the specific details of the transport cycle and how it is achieved on a molecular level are unique to each transporter or transporter family, the active use of free energy suggests that all transport cycles must maintain strict substrate coupling to minimize energetically wasteful 'slippage cycles' (*c.f.*, Fig. 1B) in which driving substrates are transported without driving the translocation of the driven substrate (Stein and Lieb, 1986).

While all secondary active transporters work by facilitating a general, dynamic equilibrium (*e.g.*, $mA_{\text{outside}} + nB_{\text{outside}} \rightleftharpoons mA_{\text{inside}} + nB_{\text{inside}}$, where A and B are co-substrates with stoichiometries m and n), the molecular details of how this chemical reaction maps on to the transport cycle are determined by a transporter's conformational dynamics. According to the alternating access hypothesis, secondary active transporters must undergo a reversible conformational transition between at least two distinct conformational states: one with the substrate binding pocket in a state that is 'outward-facing' with respect to membrane (C_o), and one in an 'inward-facing' state (C_i) (Jardetzky, 1966). While there are many possible structural realizations of alternating access, including the so-called 'rocker-switch', 'rocking-bundle', and 'elevator' transport mechanisms (Drew and Boudker, 2016), at least two distinct conformational transitions between C_i and C_o must occur in a complete transport cycle (*c.f.*, Fig. 1B). Here, we investigate the role of such conformational dynamics in the transport cycle of the prototypical sodium-dependent divalent anion transporter from *Vibrio cholerae*, VcINDY, and elucidate how they are tuned to enable efficient transport and minimize slippage.

VcINDY is a well-characterized member of the divalent anion/sodium symporter (DASS) class of secondary active transporters, which extends across all kingdoms of life. In mammals, it is represented by the solute carrier 13 (SLC13) gene family, which includes members that transport di- and trivalent anions, including sulfate and organic carboxylates such as citrate and succinate (Bergeron et al., 2013; Markovich, 2012). Within this family, the SLC13A5 citrate transporter, NaCT, has notably significant metabolic and neurologic health importance for humans, and mutations in NaCT can cause epilepsy in children (reviewed in (Sauer et al., 2022b)). Like NaCT, VcINDY is a Na$^+$-driven carboxylate transporter, but VcINDY is primarily a succinate transporter with an electrogenic transport cycle stoichiometry of 3 Na$^+$ ions per succinate (Fig. 1A) (Mulligan et al., 2014; Fitzgerald et al., 2017). Over the past decade, extensive *in vitro* biochemical characterization (Mulligan

et al., 2014, 2016; Fitzgerald et al., 2017; Sampson et al., 2020, 2021), and extensive structural insight from X-ray crystallography, cryo-electron microscopy (cryo-EM), and computational studies (Mancusso et al., 2012; Mulligan et al., 2016; Nie et al., 2017; Sauer et al., 2021, 2022a), have made VcINDY into the prototypical transporter for understanding the mechanistic details of transport in the DASS family.

While the biochemical and structural studies published to-date do not yield a description of the molecular mechanism for the complete VcINDY transport cycle, recent combinations of computational modeling, biochemical, and structural analysis have provided insight (Mulligan et al., 2016; Sauer et al., 2020). Structural studies have revealed VcINDY is a homodimeric membrane protein with C_2 symmetry; however, in all these studies, VcINDY was only observed in a single C_i conformational state in which the substrate binding pockets of both protomers are accessible only from the cytosolic face of the protein (*i.e.*, the global homodimer conformation is C_i-C_i) (Mancusso et al., 2012; Nie et al., 2017; Sauer et al., 2020). Computational and crosslinking studies suggest that VcINDY functions using an elevator mechanism, with a large translocation of the 'transport' domain that carries substrates from one side of the membrane to the other, and a relatively stationary 'scaffold' domain that harbors the dimerization interface (Mulligan et al., 2016). This model is supported by recent structural work revealing a C_o state for LaINDY, a close homolog from *Lactobacillus acidophilus* (Sauer et al., 2020). While these lines of evidence support an elevator mechanism, there has been no experimental observation of a transition between these C_o and C_i states, nor any direct evidence that they are responsible for achieving alternating access during transport. Comprehensive validation of this model requires confirming the proposed elevator motions, connecting them to the transport cycle, and understanding how those conformational dynamics enable VcINDY to achieve tight co-substrate coupling between Na^+ and succinate to minimize energetically costly slippage of substrates across the membrane (Fig. 1B).

Using single-molecule fluorescence resonance energy transfer (smFRET) microscopy, we have directly observed the global conformational dynamics of homodimeric VcINDY in a lipid-bilayer membrane, allowing us to characterize the structural and kinetic details of its transport cycle. To achieve this, we designed two complementary VcINDY constructs for smFRET experiments. Each smFRET construct contained a FRET donor- and acceptor fluorophore pair engineered at residue positions such that the resulting, donor-acceptor distance-dependent FRET efficiency (E_{FRET}) would report on the conformational dynamics responsible for substrate translocation (Fig. 2). These smFRET constructs can therefore be used to directly observe the motions that VcINDY undergoes to achieve alternating access (Fig. 1B). In performing smFRET experiments with these constructs, we obtained direct experimental evidence that (i) global conformational dynamics of VcINDY during the transport cycle are consistent with the elevator model of transport, (ii) the individual protomers of homodimeric VcINDY exhibit minimal cooperativity, and (iii) VcINDY minimizes slippage using a mechanism in which co-substrate binding is highly cooperative. These findings provide the first comprehensive description of the molecular underpinnings of the VcINDY transport cycle, and, by extension, deep insight into the transport cycle of all DASS transporters.

Results

smFRET constructs designed to report on substrate translocation.

In order to directly monitor the VcINDY conformational dynamics relevant to the transport cycle (Fig. 1B), we designed two smFRET constructs based on the assumption that substrate translocation in VcINDY involves a large-scale conformational change between the C_o-C_o and the C_i-C_i conformations (*c.f.,* Fig. 2) (Mancusso et al., 2012; Mulligan et al., 2016; Nie et al., 2017; Sauer et al., 2020). One construct was designed to contain a FRET donor- and acceptor fluorophore pair on the external, periplasmic face of the transport domains of VcINDY whereas the other construct was designed to contain them on the internal, cytosolic face. This multi-perspective approach provides a self-consistency such that any global conformational motions of the VcINDY homodimer occurring during the transport cycle (*e.g.*, an elevator motion) can be detected on both the external and internal sides of the protein.

To design these 'external' and 'internal' smFRET constructs, we identified solvent-exposed residues in the transport domain with particular distances between the two equivalent positions across the homodimer (Fig. 2). That distance was chosen such that it would undergo a significant change upon a transition between the structurally defined C_i-C_i and the computationally predicted C_o-C_o states (Mancusso et al., 2012; Mulligan et al., 2016; Nie et al., 2017; Sauer et al., 2020). In an optimized smFRET experiment, the corresponding change in the distance-dependent E_{FRET} will utilize the full dynamic range of E_{FRET} from zero to one. To achieve this, we sought residues that moved from separations of less than the distance corresponding to the half-maximum E_{FRET} (R_0) of the donor-acceptor pair to more than the R_0, or vice versa. For Alexa Fluor 555 as our donor and Alexa Fluor 647 as our acceptor, the corresponding R_0 is ~51 Å (Lambert, 2019). Given these considerations, we chose Gly211, located on the extracellular face of the transport domain as our external site, with a predicted E_{FRET} change from ~0.20 to ~0.93 upon a transition from the C_o-C_o to the C_i-C_i state. Likewise, we chose Ser436, located on the cytoplasmic face of the transport domain as our internal site, with a predicted E_{FRET} change from ~0.70 and ~0.12 upon transition from the C_o-C_o to the C_i-C_i state (Fig. 2). Single VcINDY homodimer E_{FRET} *versus* time trajectories (*i.e.*, E_{FRET} trajectories) recorded using the external and internal smFRET constructs should then report upon the same set of global VcINDY conformational dynamics involved in substrate translocation but from different perspectives.

Biochemically active variants of the two VcINDY smFRET constructs designed above were created, overexpressed, purified, and labeled with donor and acceptor fluorophores to generate the externally and internally labeled smFRET constructs (VcINDY[ext] and VcINDY[int], respectively) (see Methods and Materials for extended details) (Fig. 2–Figure Supplement 1). Both VcINDY[ext] and VcINDY[int] were able to catalyze substrate translocation and exhibited transport activities that were only three-fold reduced relative to wildtype VcINDY similar to the activity of the unlabeled cysteine-containing mutants (Fig. 2–Figure Supplement 2). To investigate the conformational dynamics of VcINDY in a membrane, we reconstituted VcINDY[ext] and VcINDY[int] into lipid bilayer nanodiscs (Fig. 3) (Denisov and Sligar, 2017). Nanodisc-based experimental conditions are ideal for

interrogating the C_o to C_i transition at equilibrium, since both sides of the membrane are exposed to identical solutions. Furthermore, the nanodiscs used here have a diameter of ~110 Å, which helped establish single-molecule resolution by ensuring that only a single VcINDY dimer could occupy each nanodisc.

smFRET experiments were performed using a widefield, TIRF microscope to record the fluorescence intensity emitted from individual, immobilized VcINDY-containing nanodiscs (Fig. 3). The corresponding E_{FRET} trajectory for each molecule was then calculated using $E_{FRET} = I_{acceptor} / (I_{donor} + I_{acceptor})$, where I_x is the observed intensity of the 'x' fluorophore. The VcINDY-containing nanodiscs appeared well dispersed when imaged using negative-stain electron microscopy (Fig. 3–Figure Supplement 1). This observation, along with the chromatographic purification used in nanodisc reconstitution and the selection criteria used to ensure each E_{FRET} trajectory was calculated using emissions from a single donor-acceptor pair (see Methods and Materials), ensured that E_{FRET} trajectories originated from single VcINDY dimers.

Multiple E_{FRET} states are stochastically sampled during VcINDY substrate translocation.

To characterize the conformational dynamics of VcINDY during the transport cycle (Fig. 1B), we performed equilibrium smFRET experiments using VcINDY[ext] or VcINDY[int] in which the buffer contained saturating concentrations of both Na^+ and succinate co-substrates (Buffer HS: High Na^+ with Succinate). The conformational dynamics that occur under these conditions are the same as those involved in vectorial transport—even in the absence of a chemical free energy driving force. Thus, by saturating the VcINDY smFRET constructs with both co-substrates, we expected to record E_{FRET} trajectories that report on the equilibrium conformational dynamics of the substrate translocation reaction between a substrate-bound C_o state and a substrate-bound C_i state (C_o" and C_i" in Figure 1B), which we hypothesized resemble the C_o-C_o and C_i-C_i conformations, respectively (Fig. 2).

In the presence of both Na^+ and succinate, the E_{FRET} trajectories originating from individual VcINDY[ext] or VcINDY[int] dimers stochastically exhibited instantaneous jumps between transiently stable states characterized by distinct E_{FRET} values (*i.e.*, E_{FRET} states) (Fig. 4). Qualitatively, the lifetimes of these E_{FRET} states were generally several seconds, and jumps occurred to E_{FRET} states with both higher and lower average E_{FRET} values. By inspection, it was not uncommon for individual E_{FRET} trajectories to visit many distinct E_{FRET} states (*e.g.*, Fig. 4, lower panels), and the same E_{FRET} states were often revisited within a single E_{FRET} trajectory. Most E_{FRET} trajectories, however, exhibited very few jumps to a low number of distinct E_{FRET} states, because relatively rapid photobleaching of the donor and/or acceptor fluorophores ended our ability the measure more E_{FRET} values before more jumps presumably occurred in those E_{FRET} trajectories.

Strikingly, the distribution of E_{FRET} states across the entire population of VcINDY[ext] or VcINDY[int] E_{FRET} trajectories spanned nearly the entire range of possible E_{FRET} values, from zero to one, and did not exhibit well-defined peaks (Fig. 5). Relative to the signal-to-background ratio of individual E_{FRET} trajectories (Fig. 4), the broad spread of these E_{FRET} state distributions (Fig. 5) suggests that VcINDY dimers exhibit significant conformational heterogeneity. Regardless, the spread of the E_{FRET} state distributions is consistent with the amplitude of

fluctuations predicted for a large, elevator conformational transition between the C_o-C_o and C_i-C_i conformations using the donor-acceptor pairs engineered into both VcINDY[ext] and VcINDY[int] (Fig. 2).

VcINDY samples C_o-C_o, C_i-C_i, and mixed C_o-C_i conformational states during substrate transport.

After observing dynamic E_{FRET} trajectories that exhibited instantaneous jumps between widely distributed E_{FRET} states for both VcINDY[ext] and VcINDY[int] (Figs. 4 and 5), we desired to understand how these E_{FRET} states and dynamics correspond to the underlying conformational states and dynamics of the VcINDY transport cycle (Fig. 1B). To do so, we sought the simplest, quantitative description of the E_{FRET} trajectories that was globally consistent with all the E_{FRET} trajectories recorded in an experiment. Because of the instantaneous jumping behavior of the E_{FRET} trajectories (Fig. 4), we used a global hidden Markov model (HMM) to quantify the common E_{FRET} states and kinetics present in all the E_{FRET} trajectories (see Methods and Materials for extended details) (Fig. 5–Figure Supplements 1-2). In the presence of both co-substrates, we found that the most parsimonious global HMMs that could account for the observed complexity in the E_{FRET} trajectories recorded for VcINDY[ext] or VcINDY[int] both had three E_{FRET} states (Fig. 5, Fig. 5–Figure Supplements 3-4). Given that we find three E_{FRET} states from both perspectives in buffer HS, and that the E_{FRET} trajectories recorded under different Na^+ and succinate conditions were also best described by three E_{FRET} states (see below) (Fig. 5—Figure Supplement 3), we concluded that our smFRET data report primarily on transitions between three common, global conformations of the VcINDY dimer.

According to the alternating access principle, at least some part of these three E_{FRET} states should correspond to the C_o and C_i states in the transport cycle (Jardetzky, 1966) (Fig. 1). For VcINDY[ext], the $E_{FRET} = 0.25$ state is consistent with our estimate of $E_{FRET} = 0.20$ for the C_o-C_o conformation, and the $E_{FRET} = 0.85$ state is consistent with our estimate of $E_{FRET} = 0.93$ for the C_i-C_i conformation (Fig. 2, Fig. 5–Figure Supplement 4). Likewise, for VcINDY[int], the $E_{FRET} = 0.83$ state is consistent with our estimate of $E_{FRET} = 0.70$ for the C_o-C_o conformation, and the $E_{FRET} = 0.16$ state is consistent with our estimate of $E_{FRET} = 0.12$ for the C_i-C_i conformation (Fig. 2, Fig. 5–Figure Supplement 4). Thus, the E_{FRET} states we observe in our smFRET experiments demonstrate that the transport domains of VcINDY undergo large-scale conformational changes during substrate transport and strongly suggest that substrate transport by VcINDY proceeds *via* an elevator mechanism.

The small differences between the expected and observed E_{FRET} values ($\Delta E_{FRET} \sim 0.1$) might be due to: (i) differences in the C_o-C_o and/or C_i-C_i structures which nanodisc-embedded VcINDY adopts in our smFRET experiments and between those in the computational, X-ray, and cryo-EM studies (Mancusso et al., 2012; Mulligan et al., 2016; Nie et al., 2017; Sauer et al., 2020), (ii) structural perturbations of VcINDY due to fluorophore labeling, (iii) linker contributions to the distance between fluorophores, or even (iv) photophysical complications, such as 'photoblinking', that effect E_{FRET} estimates. Given the experimental consistency with the structural estimates for both VcINDY[ext] and VcINDY[int], we assigned the E_{FRET} states discussed above to the C_o-C_o and C_i-C_i conformational states. Additionally, these assignments are self-consistent with respect to the equilibrium state population distribution of the transport cycle (Fig. 1) in that the most occupied E_{FRET} state in

both smFRET experiments was the C_i-C_i VcINDY conformational state, and was thus independent of the smFRET perspective used (Fig. 5).

The remaining, unassigned E_{FRET} state can be explained as either a new, 'intermediate' conformation distinct from C_o-C_o or C_i-C_i, or as a 'mixed' C_o-C_i conformation of the VcINDY dimer. Based on biochemical and structural considerations, we favor assigning this E_{FRET} state to the more parsimonious, mixed conformation. The existence of such a mixed conformation is only possible if the protomers of VcINDY operate independently, which is suggested by previous Hill coefficient measurements for succinate transport (Mulligan et al., 2014). Additionally, the C_2 symmetry of the VcINDY dimer renders the C_o-C_i and C_i-C_o conformations structurally degenerate, and thus the C_o-C_i and C_i-C_o E_{FRET} states would both yield the same E_{FRET} value, which is consistent with our analysis. In contrast, similar symmetry considerations for a VcINDY dimer with a unique intermediate protomer conformation would yield up to six distinct E_{FRET} states in the absence of strict protomer cooperativity or anti-cooperativity. Finally, computational alignment of the VcINDY scaffold domain of the C_o-C_o and C_i-C_i conformations shows that the donor and acceptor fluorophores in a mixed C_o-C_i state would be ~50 Å apart for VcINDY[ext] and ~59 Å apart for VcINDY[int]—distances that correspond to E_{FRET} values of 0.53 and 0.29, respectively. Given the caveats described above, these estimates are in reasonable agreement with the E_{FRET} values of 0.54 and 0.42 that we obtained from our global HMM analysis of the smFRET data (Fig. 5–Figure Supplement 4).

Despite the limited observation window of any one individual E_{FRET} trajectory, the global HMM analyses we performed provide a quantification of the kinetics present within the collective set of E_{FRET} trajectories (Fig. 5—Figure Supplement 4). Regardless of the smFRET perspective used (VcINDY[ext] or VcINDY[int]), the elevator motion of a single protomer in the VcINDY dimer across the membrane occurs in these experiments at a rate of ~0.4 s^{-1} (*i.e.*, once every ~2.5 seconds), with no apparent directional dependence. Notably, direct transitions between C_o-C_o and C_i-C_i where both protomers undergo an apparently simultaneous transition within the temporal resolution of our TIRF microscope were approximately an order of magnitude slower (~0.04 s^{-1}), and therefore much less likely to occur. This suggests that the individual protomers of a VcINDY dimer primarily undergo the translocation reaction in an independent and uncoordinated manner. Overall, the elevator motions of VcINDY responsible for substrate translocation that occur in these experiments are approximately an order of magnitude faster than the turn-over rate for transport measured in bulk biochemical transport experiments of ~0.02 s^{-1} (Mulligan et al., 2014). This suggests that VcINDY undergoes multiple substrate translocation reactions before a productive transport event occurs, and that the rate limiting step of VcINDY, and possibly DASS family, catalyzed transport may involve substrate binding and/or dissociation.

VcINDY exhibits conformational dynamics that are largely substrate independent.

The basic mechanistic model of a transport cycle for VcINDY makes several fundamental predictions regarding the conformational dynamics that occur during transport (Fig. 1B). Consistent with the alternating access hypothesis (Jardetzky, 1966), VcINDY should be able to switch from an outward-facing (C_o) to inward-facing (C_i)

conformation in the apo, substrate-free state. Also, to preserve substrate coupling, we expect that the partially bound transporter (C_o' and C_i' in Fig. 6) will not undergo outward to inward transitions (and vice versa) to minimize uncoupled slippage of Na^+ or succinate. To test these predictions, we performed additional smFRET experiments using VcINDY[int] in the absence of both co-substrates (*i.e.*, in the apo state) (Buffer LX: Low Na^+ with no succinate) as well as in the presence of high concentrations of one co-substrate but the absence of the other (*i.e.*, in the only bound Na^+- or only succinate bound states) (Buffer HX: High Na^+ with no succinate; Buffer LS: Low Na^+ with Succinate). Where appropriate, we replaced Na^+ with choline to maintain ionic strength and osmolarity. We note that all of our buffers likely contain trace (*i.e.*, 'low') amounts of contaminating Na^+, but at several orders of magnitude less than the Michaelis constant, K_M, for transport (Mulligan et al., 2014).

As observed for the smFRET experiments recorded in the presence of high concentrations of both co-substrates (Buffer HS), the smFRET experiments recorded under all three additional co-substrate conditions (Buffers LX, HX, and LS) yielded E_{FRET} trajectories that spontaneously exhibited instantaneous jumps between E_{FRET} states spanning a comparably wide range as was seen in buffer HS (Fig. 5). When we analyzed these data using the same global HMM approach described above (see Methods and Materials for extended details), the most parsimonious quantification of the sets of E_{FRET} trajectories in each smFRET experiment was again a global HMM with three E_{FRET} states (Fig. 5–Figure Supplement 3). In these global HMMs, both the E_{FRET} states and rate constants for the transitions between them were very similar to those obtained in Buffer HS (Fig. 5–Figure Supplement 4). While there were slight differences in the equilibrium population distribution of the three E_{FRET} states observed in each co-substrate condition, the kinetics of these global HMMs were relatively unchanged with single protomer translocation reactions occurring all at approximately the same rate of ~0.5 s^{-1}. Thus, our quantitative kinetic analysis reveals that the conformational motions of VcINDY that are probed along the coordinate between the donor and acceptor fluorophore in VcINDY[int] are relatively independent of the co-substrate composition. Although these results conform to our expectation for the apo state, the observation of frequent transitions in the E_{FRET} trajectories recorded under the single-substrate conditions was unexpected.

Thermodynamic modeling shows that compensatory tuning of co-substrate binding and dissociation can minimize slippage cycles.

In search of an explanation for the surprising kinetics observed under the single-substrate conditions, we next analyzed the thermodynamic efficiency of the VcINDY transport cycle. We modeled the free-energy transduction of a secondary active transporter using Hill's method of cycles (Hill, 2005) to determine how the different molecular processes involved in transport might contribute to slippage. For simplicity, we considered co-substrate binding as a strictly sequential, two-step process with mirror symmetry (*i.e.*, the same binding/release order on both faces of the membrane). To be concrete, we considered the most parsimonious molecular interpretation of this kinetic scheme for secondary active transport where (i) the rate constants are all first-order or pseudo-first order, (ii) the rate constants are independent of the C_o or C_i state, and (iii) the translocation reaction is independent of substrate binding status (Fig. 6). The resulting, simplified kinetic scheme has three

Hill cycles—two of which are involved in slippage (cycles *A* and *B*) and one that is responsible for transport (cycle *C*) (Fig. 6).

We then calculated the thermodynamic efficiency, η, of a transporter with these assumptions (Fig. 6) (Hill, 2005). For a very efficient transporter, $\eta \approx 1$ and the transporter tightly couples free energy from the driving to the driven co-substrate with very little slippage. In that situation, the thermodynamic efficiency is $\eta \propto 1 - (\eta_{ac} - \eta_{bc}) + \cdots$ with small higher order terms; here we have assumed A is the driving substrate, but the conclusions discussed below are independent of this. Under our assumptions,

$$\eta_{ac} = \frac{J_a}{J_c} = \left(\frac{[A]_o^m - [A]_i^m}{[A]_o^m [B]_o^n - [A]_i^m [B]_i^n} \right) \left(\frac{2k_t + k_d''}{k_a''} \right),$$

and

$$\eta_{bc} = \frac{J_b}{J_c} = \left(\frac{[B]_o^n - [B]_i^n}{[A]_o^m [B]_o^n - [A]_i^m [B]_i^n} \right) \left(\frac{k_t([A]_o^m + [A]_i^m) + k_a'[A]_o^m[A]_i^m}{k_d'} \right),$$

where J_κ is the cycle flux for cycle κ, concentrations are indicated by square brackets, A and B are the first and second binding co-substrates with stoichiometries m and n, respectively, the translocation rate constant is k_t, association and dissociation rate constants are k_a and k_d, respectively, and prime and double-prime denote the first and second binding reactions, respectively (Fig. 6). The first terms in these equations show that the relative amount of slippage that a transporter undergoes depends on the instantaneous thermodynamic driving forces *via* the co-substrate chemical gradients, while the second terms show that it also depends on the translocation and substrate binding and dissociation reaction kinetics. Thus, the only reliable way nature can optimize this transporter into a highly efficient free energy transducer over a broad range of environmental conditions is to have evolved control over slippage *via* evolutionary tuning of these rate constants to reduce the total magnitude of η_{ac} and η_{bc}. This suggests that optimal transporters have faced evolutionary pressure to decrease k_t, k_a' and/or k_d'', and to increase k_a'' and/or k_d' such that the first binding co-substrate weakly binds until the second arrives and then both co-substrates can tightly bind. Unfortunately, tuning these rate constants will also affect the overall transport cycle flux, J_c, and this creates a speed-accuracy tradeoff.

The speed-accuracy tradeoff can, however, be navigated. On the molecular level, the clearest way to reduce slippage is for the binding pocket to stabilize the *apo* and fully bound states (C and C"), by some amount $\Delta G'$ and $\Delta G''$, respectively, and to source all of that energy from a compensatory destabilization of the partially bound conformation, C', by an amount $\Delta G' + \Delta G''$, all without affecting the transition state energy barriers (Fig. 7). This is a conceptually similar idea to the use of substrate binding energy to stabilize the transition state for enzyme-substrate transition state complementarity (Fersht, 2017). In our case, this isoenergetic change increases k_a' and k_d'' while decreasing k_a'' and k_d', which reduces slippage and increases the thermodynamic efficiency of the transporter. It also has the effect of increasing the transport cycle flux *via* ~75% of the terms defining J_c (see *Methods and Materials* for the expression for J_c). Given this substantial ability to speed up transport in the complicated expression for J_c, we suspect that these compensatory, isoenergetic binding and dissociation reaction tunings optimally navigate the speed-accuracy tradeoff by simultaneously decreasing the

relative fraction of slippage while increasing the total amount of transport. The exact extent of these abilities, however, depends on the instantaneous internal and external co-substrate concentrations, which can vary significantly across physiologically relevant conditions (Fig. 7–figure supplement 1) (Lo et al., 2006). Regardless, this thermodynamic modeling of the free energy transduction efficiency of a secondary active transporter predicts that slippage can be reduced simply by isoenergetic optimization of the energy landscape of the co-substrate binding reaction (Fig. 7). In other words, increased cooperativity of co-substrate binding can itself lead to minimized slippage.

Finally, we also used this thermodynamic modeling-based approach to estimate the occupancy of apo VcINDY in our smFRET experiments performed in presence of only saturating Na$^+$ (Buffer HX). Specifically, we modeled the reported K_M and V_{max} values for the Na$^+$ titration and succinate titration experiments that recorded succinate uptake rates in the biochemical proteoliposome-based transport assay data of Mulligan and coworkers (see *Methods and Materials* for extended details) (Mulligan et al., 2014). This was an underdetermined problem, so we assumed that a single, average rate constant defined several molecular processes: $\langle k \rangle \equiv k_t = k_d' = k_d''$. Markov chain Monte Carlo optimization yielded estimates of $k_a' = 1.5 \times 10^7 M^{-3} s^{-1}$, $k_a'' = 7.0 \times 10^8 \ M^{-1} s^{-1}$, and $\langle k \rangle = 1.05 \times 10^3 \ s^{-1}$ (Fig. 6-Figure Supplement 1 and 2), and reasonably matched the original titration data (Fig. 6-Figure Supplement 3). The mismatch is due to the common rate constant assumption, and probably to differences in the actual binding mechanism assumed here (*e.g.*, perhaps not all three Na$^+$ ions bind before succinate). Regardless, these estimates are consistent with the biochemical transport data in the context of our kinetic mechanism, so we used them to calculate the steady-state probability of apo VcINDY in Buffer HX to be 6.5% (see *Methods and Materials*). This estimate suggests that even with 100 mM Na$^+$ in solution, VcINDY still spends a non-negligible portion of time with an unoccupied binding pocket. In this light, the majority of the unexpected elevator motions present in the E$_{FRET}$ trajectories recorded in the presence of only one co-substrate might originate from apo, rather than partially bound, VcINDY.

Dicarboxylate substrate binding pocket is enthalpically stabilized by sodium ions.
To obtain atomic-resolution insight into the Na$^+$ and dicarboxylate substrate binding reaction for the VcINDY transport cycle, we determined how the conformational state of the VcINDY homodimer is influenced by different dicarboxylate substrates. We therefore separately crystallized detergent-solubilized VcINDY with Na$^+$ and three different C$_4$-dicarboxylate substrates (succinate, fumarate, and malate) and solved their structures at 3.09 Å, 3.29 Å, and 3.50 Å, respectively. Regardless of the dicarboxylate substrate identity, VcINDY adopted the same C$_i$-C$_i$ state in all three of these structures. Similarly, as seen in the previous structure of the C$_i$-C$_i$ state (Mancusso et al., 2012; Nie et al., 2017; Sauer et al., 2020), the binding pocket of the dicarboxylate substrate in all three structures was partially comprised of two hairpin loops that coordinated distinct Na$^+$ ions, Na1 and Na2 (Fig. 8). Given this binding geometry, it is possible that tight binding of both co-substrates requires stabilization of these hairpin loops and that their local conformational rearrangements gates binding and/or release of the co-substrates.

Discussion

The VcINDY transport domain undergoes elevator motions to enable alternating access.

The ubiquitous presence in our experiments of stochastic, instantaneous transitions between E_{FRET} states consistent with the C_o-C_o and the C_i-C_i conformations is the first direct evidence for the hypothesis that VcINDY uses an elevator transport mechanism to enable alternating access (Fig. 4) (Mulligan et al., 2016). Because we directly observed these stochastic, elevator motions in the presence of substrates but absence of an electrochemical gradient (Buffer HS) and also in the absence of any co-substrates at all (Buffer LX), it is clear that the VcINDY transport domain can translocate the substrate binding site across the membrane using only thermal energy—as is expected for the conformational motions of a secondary active transporter that are responsible for transport. Moreover, the rate constants for that process are on the order of 10^{-1} s^{-1} (Fig. 5–Figure Supplement 4), suggesting that the energy barrier separating the C_o and C_i state along the translocation reaction coordinate is not prohibitively high. Since the VcINDY transport cycle is much slower and occurs with a rate on the order of 10^{-2} s^{-1} (Mulligan et al., 2014), it seems that multiple back and forth substrate translocation events occur before a productive transport event is completed. This observation is consistent with a model in which a molecular process other than transport domain translocation (*e.g.*, gating dynamics within the substrate binding pocket itself) govern the transport rate.

The conformational dynamics of VcINDY protomers are non-cooperative.

The experimental observation of a mixed C_o-C_i conformation for the VcINDY dimer constrains the role that the two protomers can play in the transport cycle. Previously, proteoliposome-based transport experiments determined that VcINDY transports succinate with a Hill coefficient of 0.88 ± 0.13 (Mulligan et al., 2014). Given the VcINDY transport stoichiometry of 3:1 Na$^+$:dicarboxylate (Fitzgerald et al., 2017), this Hill coefficient measurement suggested that the VcINDY transport cycle is non-cooperative and that each VcINDY protomer can independently transport succinate. We supported this biochemical hypothesis on a molecular level by interpreting our experimental observation of the E_{FRET} state assigned to the mixed C_o-C_i conformation as direct evidence that the VcINDY protomers undergo non-cooperative substrate translocation reactions during vectorial transport. Since the VcINDY dimer can significantly occupy the C_o-C_i state, the two VcINDY protomers must not undergo a strictly cooperative translocation reaction (*i.e.*, C_o-$C_o \rightleftharpoons C_i$-$C_i$). Furthermore, they must not undergo a strictly anti-cooperative translocation reaction either (*i.e.*, C_o-$C_i \rightleftharpoons C_i$-$C_o$), because this precludes the occupation of the C_o-C_o and C_i-C_i conformations that we observe in our smFRET experiments. Therefore, our results provide direct evidence that the VcINDY protomers undergo non-cooperative translocation reactions, and that the VcINDY transport cycle itself can be considered to occur independently in each protomer.

In support of this idea, a recent structure of CitS, another homodimeric elevator transporter in a different structural family, reveals a CitS dimer with one inward-facing and one outward-facing protomer (Kim et al., 2017), and a recent high-speed atomic force microscopy study of the CitS elevator translocation motion reported that

CitS undergoes non-cooperative interprotomer conformational dynamics (Maity et al., 2022). These results, combined with evidence that an elevator-type, glutamate transporter, Glt_{Ph}, also shows independence between the motions of its three protomers (Akyuz et al., 2013; Erkens et al., 2013; Ruan et al., 2017), suggest that such independence may be a general feature of oligomeric elevator transporters (Garaeva and Slotboom, 2020).

Slippage cycles are avoided without inhibiting VcINDY transport domain translocation.

To avoid wasting the free energy provided by the Na^+ gradient that drives transport, the conformational motions of VcINDY that enable it to transport substrates across the membrane must be coordinated. In particular, VcINDY must minimize slippage cycles where Na^+ ions are transported across the membrane without the corresponding transport of a dicarboxylate substrate (Fig. 1). We had hypothesized that VcINDY avoids such slippage cycles by inhibiting the substrate binding pocket translocation reaction (*i.e.*, $C_o \rightleftharpoons C_i$) when sub-stoichiometric amounts of the co-substates are bound, and by promoting those motions only when VcINDY is apo or fully bound. However, the smFRET experiments we recorded under singly bound substrate conditions, yielded E_{FRET} trajectories that exhibited no discernable differences in the E_{FRET} states or their rates of transitions, regardless of the presence of relatively high concentrations of either individual co-substrate in solution (Fig. 5–Figure Supplement 4). Given our determination that those observed transitions could reasonably correspond to apo VcINDY, it is quite possible that transitions between C_o and C_i in partially bound states are directly suppressed and that we were not able to observe those effects, but our smFRET experiments suggest that VcINDY also uses another, additional mechanism to minimize slippage (Fig. 1B).

There are at least two additional mechanisms that VcINDY can use to minimize slippage. One possible explanation is that VcINDY employs a secondary conformational change—independent of $C_o \rightleftharpoons C_i$ translocation—to 'gate' both binding and release of co-substrates. For instance, while VcINDY could be able to translocate substrates across the membrane if the $C_o \rightleftharpoons C_i$ transition is not completely inhibited, a slippage cycle cannot be completed without co-substrate release. Mechanistically, such gating can be achieved with an additional VcINDY conformational equilibrium that is relatively independent of the transport domain translocation reaction $C_o \rightleftharpoons C_i$. Without direct structural insight into such a hypothetical equilibrium, it is attractive to think that it might involve a conformational rearrangement of the binding pocket and, in particular, the hairpin loops (*c.f.,* Fig. 8). In this mechanistic scenario, the transitions we observe in the smFRET data for partially bound VcINDY would reflect transitions between so-called "occluded" states.

Another mechanism that VcINDY could concurrently employ to minimize slippage is to ensure that co-substrate binding is extremely cooperative, such that essentially no binding can occur unless both co-substrates are present. In our smFRET experiments performed in Buffers HX and LS, this mechanism would manifest as transitions in the E_{FRET} trajectories that predominately correspond to transport domain translocation of apo VcINDY. That is consistent with the non-negligible fraction of apo VcINDY we estimated is present under these experimental conditions. Since we observed similar conformational dynamics from apo VcINDY in smFRET experiments in Buffer LX, this could explain the apparent lack of substrate dependence in our smFRET

experiments (Fig. 5-Figure Supplement 4). This cooperative co-substrate binding scenario agrees with our thermodynamic modeling (Fig. 6), which predicts that an optimized transporter that has evolved to minimize slippage cycle flux should have extremely stable apo and fully bound states with very weak binding of the initial co-substrate(s) (Fig. 7). The extent of that optimization reflects how nature has navigated the speed-accuracy tradeoff between rapid transport and minimal slippage. This raises the possibility that some secondary active transporters undergo a non-negligible amount of slippage in order to be efficient at transporting the driven co-substrate as long as they do not significantly waste the free energy of the driving co-substrate.

Experimental evidence is also consistent with a cooperative co-substrate binding mechanism. For instance, our X-ray crystallographic structures show that Na1, Na2, and dicarboxylate substrates all interact with common hairpin loop elements (Fig. 8). That the VcINDY conformation is effectively independent of dicarboxylate identity suggests that this fully bound conformation is very stable—as predicted in our modeling of optimal co-substrate binding. Moreover, recent PEG-modification protection experiments demonstrated that, in the presence of high concentrations of only Na^+, residues associated with the substrate binding pocket are protected from PEG modification (Sampson et al., 2020; Sauer et al., 2022a). This suggests some interaction between Na^+ and the apo VcINDY binding pocket, which is consistent with a Na^+-first binding order and with the prediction that slippage is minimized and transport is optimized by weak binding of the first co-substrate(s) (Fig. 7). Regardless, in order to understand the extent to which VcINDY utilizes cooperative co-substrate binding as a mechanism to avoid slippage cycles during vectorial transport, future work should aim to determine the affinities and binding order of co-substrates, as well as the structural basis for how co-substrate binding and release is achieved by VcINDY.

Conclusion

Previous studies of the structure, transport biochemistry, and alternating access mechanism of VcINDY have established it as a prototypical membrane-bound transporter of the DASS family (Mancusso et al., 2012; Mulligan et al., 2014, 2016; Nie et al., 2017; Fitzgerald et al., 2017; Sauer et al., 2020; Sampson et al., 2020). Here, we have advanced this fundamental understanding of the DASS family and secondary active transport by directly determining how the conformational dynamics of VcINDY enable it to achieve alternating access, and how those dynamic motions are tuned in order to maintain the efficiency of the transport cycle. Specifically, our study provides direct evidence that (i) the VcINDY translocation reaction uses a thermally driven, elevator mechanism; (ii) VcINDY protomer conformational dynamics are non-cooperative; (iii) the VcINDY transport cycle uses cooperative substrate binding to minimizes slippage; and (iv) cooperative substrate binding is a general approach to optimizal transport efficiency for certain secondary active transporters. Altogether, these findings immediately suggest future studies aimed at determining the extent to which highly cooperative co-substrate binding and/or additional conformational dynamics that gate transport (Drew and Boudker, 2016; Lolkema and Slotboom, 2019; Stein and Lieb, 1986) enable VcINDY to avoid slippage cycles during vectorial transport. Overall, the powerful

smFRET-based experimental modality for interrogating the function and regulation of VcINDY established here is easily extended to study the transport cycles of other DASS family transporters, and even dimeric transporters from other families, to further our understanding of secondary active transport.

Methods and Materials

Expression and purification of VcINDY. Expression and purification of wild type VcINDY was carried out according to our previous protocol (Mancusso et al., 2012). Briefly, *E. coli* BL21-AI cells (Invitrogen) were transformed with a modified pET vector encoding N-terminal 10X His tagged VcINDY. Cells were grown for 16 hours post-induction at 19°C. After cell breakage, membranes were solubilized in 1.2% N-decyl-β-maltoside (DM) and the protein was purified on a Ni-NTA affinity column (Qiagen). Following His-tag removal, VcINDY was further purified by size-exclusion chromatography (SEC) in buffer containing 25 mM Tris pH 8, 100 mM NaCl, 5% glycerol, 0.1% DM unless otherwise indicated.

Purification of VcINDY mutants. Expression and purification of VcINDY mutants G211C and S436C was performed as with the wildtype protein, but with the following modifications. All buffers contained 10 mM sodium citrate but not glycerol. Membranes were solubilized in 1.0% N-dodecyl-β-maltoside (DDM). The protein was eluted from Ni-NTA agarose beads using 300 mM imidazole. Finally, the SEC buffer contained 25 mM Tris pH 8, 100 mM NaCl, 0.1% DDM and 0.5 mM TCEP.

Expression, purification and biotinylation of membrane scaffolding protein. Expression and purification of membrane scaffolding protein (MSP1E3D1) was carried out according to a previously published protocol (Denisov et al., 2004). In brief, *E. coli* BL21-gold (DE3) cells (Agilent) were transformed with a modified pET 28a vector encoding N-terminal 7X His tagged MSP with a mutated cysteine at position 277 for biotinylation. Cells were grown at 30°C for 4 hrs post 0.1 mM IPTG induction at OD 0.8. After cell breakage, lysate was bound to Ni-NTA agarose beads (Qiagen) for 4 hours with nutation at 4°C. Protein was eluted with 100 – 250 mM imidazole and dialyzed into 100 mM Tris pH 7.4, 300 mM NaCl for 1 hr at 4°C. The His-tag was then removed by overnight TEV digestion at 12°C. Cleaved MSP protein was subsequently biotinylated using EZ-Link Maleimide-PEG11-Biotin (Promega) according to the manufacturer's instructions. Specifically, a 20-fold molar excess of biotin reagent dissolved in DMSO was incubated with protein overnight at 4°C. Labeled MSP1E3D1-T277C protein was further purified by SEC in 100 mM NaCl, 1 mM TCEP, 20 mM Tris pH 7.4.

Fluorophore Labeling. Stock solutions of Alexa Fluor 555 C2 maleimide (AF555) and Alexa Fluor 647 C2 maleimide (AF647) were prepared in anhydrous DMSO at 3mM and stored at -20°C. Prior to labeling, a 1 mg/ml solution of VcINDY (total of 0.5mg to 1mg) was treated with 10mM DTT for 2hrs at 4°C to reduce the cysteines. DTT was then removed by 3 serial dilutions in SEC buffer (25 mM tris pH 8.0, 100 mM NaCl, 5% glycerol, 0.15% DDM, 1 mM TCEP) followed by re-concentration with an Amicon concentrator (cutoff 50 kDa). This preparation

was incubated with a 10-fold molar excess of AF555 and AF647 overnight at 4°C, protected from light. The resulting solution was purified on Superdex 200 10/300 GL size exclusion column to remove the unreacted dye. Fractions from the elution peak were combined and the labeling stoichiometry was determined with a spectrophotometer using the following wavelengths and extinction coefficients: VcINDY: λ=280 nm with ϵ = 57075 cm^{-1}M^{-1}; AF555: λ= 556 nm with ϵ= 155,000 cm^{-1}M^{-1}; AF647: λ= 651 nm with ϵ= 270,000 cm^{-1}M^{-1}. Typical labeling efficiency was 42% for each dye.

Nanodisc Reconstitution. A thin film of *E. coli* polar lipid (EPL) was prepared by adding 10 mg of chloroform stock solution to a glass test tube. The solvent was evaporated with argon gas and the test tube was placed in a vacuum chamber at ~1 mTorr for ~2 hr. This lipid film was resuspended by adding 285 µl of nanodisc buffer (20mM Tris pH7.4; 100mM NaCl, 0.5mM EDTA, 0.5mM TCEP) followed by vortexing and sonication in an ultrasonic bath (Laboratory Supplies Co. Inc, Model G112SP1T) with 10 sec bursts alternating with 10 sec rests for a total of 3min. This aqueous lipid solution was solubilized by adding 215 µl of 10% DDM to produce 500 µl of final lipid stock concentration at 20mg/ml.

To reconstitute VcINDY into nanodiscs, detergent-solubilized EPL was added to VcINDY at a molar ratio of 280 and incubated on ice for 10 min. Biotinylated MSP was then added to this mixture at an 8-fold molar excess to produce 500 µl of a solution containing 2.5mM EPL, 72 µM MSP, 9 µM VcINDY and 7.2 mM DDM. This solution was incubated at 4°C for 1hr followed by the addition of SM-2 Bio-Beads in three steps: 300mg initially, 200mg added 60 min later, and 100mg added at 90 min later and incubated overnight. During these incubations, solutions were kept at 4°C and gently stirred.

Nanodiscs containing VcINDY were purified by incubating the solution with 0.5 ml of Ni-NTA beads pre-equilibrated with nanodisc buffer (20mM Tris pH7.4; 100mM NaCl; 0.5mM EDTA; 0.5mM TCEP) for 30min at 4°C. The beads were packed into a column and washed with 1 ml of nanodisc buffer. Nanodiscs containing VcINDY were then eluted with 2 ml nanodisc buffer supplemented with 0.3 M imidazole and the peak elution fractions were pooled and concentrated with an Amicon concentrator (cutoff 50 kDa). The concentrated sample was fractionated on Superdex 200 10/300 GL size exclusion column (GE Healthcare) pre-equilibrated and eluted with nanodisc buffer. The fractions were evaluated by SDS-PAGE and those containing both MSP and VcINDY were pooled, and concentrated to ~0.2-0.4 mg/ml. Typical yield for 1mg of VcINDY was ~0.4 mg of reconstituted protein (VcINDY plus MSP). Electron microscopy grids were prepared in 2% (w/v) phosphotungstic acid (pH 7.5) and were examined in a Talos L120C transmission electron microscope (FEI-Thermo Fisher Scientific). Nanodisc samples were frozen in aliquots and stored at –80°C.

TIRF Microscopy. smFRET imaging experiments were performed on a prism-based, wide-field TIRF microscope. Briefly, a 532 nm diode-pumped solid-state continuous-wave laser (Laser Quantum, GEM532) was directed through an acousto-optic modulator (AOM) (Isomet, IMDD-P80L) such that the +1 order peak is directed toward a prism suspended over a 60x, 1.2 NA water immersion objective (Nikon). VcINDY embedded nanodiscs

were tethered to the surface of quartz slides derivatized with dilute biotin-poly(ethylene)-glycol (PEG) in PEG *via* a biotin-streptavidin-biotin bridge, and illuminated with the TIR field generated at the slide/buffer interface. The instantaneous Illumination power incident at the prism entry face was 40 mW (estimated density of 4.3 W/cm^2), and the AOM was modulated with a 50% duty cycle at 1 MHz, such that the time averaged power was 20 mW. Fluorescence was collected through the objective, separated with a multi-wavelength imager (Photometrics, Dual-view), and imaged onto a water-cooled electron-multiplying charge coupled device (Andor, iXon Ultra 888). Movies were collected for 2000 frames at 25 msec exposures with 2x binning, 600 EM gain, 1x pre-amp gain, 30.0 MHz readout mode, 0.60 μs vertical speed, and +4 clock voltage running at –67°C using the open source microscopy software Micro-Manger (Edelstein et al., 2014).

smFRET Imaging Buffers. VcINDY homodimers were imaged in the following buffers, which all contained 100 mM KCl, and 20 mM Tris-Cl (pH$_{25C}$=7.5). **Buffer LX**: 100 mM choline chloride. **Buffer LS**: 100 mM choline chloride, and 1 mM succinic acid. **Buffer HX**: 100 mM NaCl. **Buffer HS**: 100 mM NaCl, and 1 mM succinic acid. All buffers also contained 1% β-D-glucose, 1 mM trolox, 0.0025 U glucose oxidase, 0.02 U catalase, 1 mM 1,3,5,7-cyclooctatetraene, and 1 mM 3-nitrobenzyl alcohol as photostabilizing additives.

smFRET Data Analysis. Diffraction-limited spots of fluorescence intensity within the first 400 frames (10 seconds) were located in both the donor and acceptor fluorescence channels. Intensity versus time trajectories were estimated using the iterative, maximum likelihood formulas for intensity estimation $N = \left(\sum_{xy}(d_{xy} - b)\Psi_{xy}\right)/\left(\sum_{xy}\Psi_{xy}^2\right)$ and $b = \left(\sum_{xy}(d_{xy} - N\Psi_{xy})\right)/\left(\sum_{xy}1\right)$, where (x,y) is a pixel location, N is the fluorescence intensity, b is the background intensity, and Ψ is a fixed width, Gaussian-shaped point spread function with width defined by the TIRF microscope optics. Donor and acceptor fluorescence channels were aligned using a 4th order polynomial transform of ~2800 control points obtained from an image of an array of sub-diffraction limit nanopatterned features localized in both channels. Trajectories were manually classified to select trajectories from only donor-acceptor labeled molecules (see Fig. 2—Figure Supplement 1). The photobleaching point for each E$_{FRET}$ trajectory was inferred using Bayesian inference (see below). A non-linear filter (Haran, 2004) was applied to fluorescence intensity trajectories for easier visualization in intensity and E$_{FRET}$ trajectory plots, but all analyses (*e.g.*, histograms, and HMMs) were performed using the unfiltered data.

Global, variational Bayes hidden Markov modeling (vbHMM). Global vbHMMs were estimated in a manner similar to a previously described single trajectory vbHMM implemented in the software package vbFRET (Bronson et al., 2009). vbFRET is a Bayesian inference-based 'maximum evidence' method (Kinz-Thompson et al., 2021), which allowed one to determine the simplest HMM that sufficiently accounts for the complexity of the smFRET data (Bronson et al., 2009; van de Meent et al., 2014; Hon and Gonzalez, 2019). In this method, one can find the most parsimonious HMM for smFRET data by estimating several HMMs, each with an increasing

number of hidden states, then calculating the 'evidence' for each HMM and choosing the one with the maximum evidence value (Bronson et al., 2009). As seen in the directed acyclic graph (Fig. 5–figure supplement 1), the only difference between vbFRET, and the global vbHMM used here is that an ensemble of E$_{FRET}$ versus time trajectories is assumed to be independent and identically distributed (I.I.D.) from the same HMM parameters in the global vbHMM model. Thus, all data points then contribute to the expectation and maximization equations for the hyperparameters governing the posterior probability distribution, as opposed to only those from a single trajectory is done for vbFRET. This is equivalent to considering one large, concatenated trajectory with vbFRET, where however there is no conditional dependence for the hidden state on the previous hidden state for neighboring data points from different trajectories. Instead, the first data point of each trajectory depends on the initial-state probability distribution denoted here by π. As can be done with vbFRET, model selection can be performed by calculating the evidence lower bound (ELBO) for each model and choosing that model with the largest value.

To demonstrate the utility of using model selection with global vbHMMs, we simulated multiple four-state time series, performed model selection, and compared this to an autocorrelation function (ACF) analysis (Fig. 5–figure supplement 2). The two-point autocorrelation function (Berne and Pecora, 2000) of a stationary stochastic process or signal, y(t), describes how similar the process remains after some time lag, τ, and can be calculated as

$$ACF(\tau) \equiv \langle \delta y(0) \cdot \delta y(\tau) \rangle = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} d\delta y(0) \cdot d\delta y(t) \cdot y(0) \cdot y(\tau) \cdot P(\delta y(0), \delta y(\tau))$$

where $\delta y(\tau) = y(\tau) - \langle y \rangle$, angular brackets denote an expectation value, P(x) is the probability distribution function of x, and

$$P(\delta y(0), \delta y(\tau)) = P(\delta y(\tau)|\delta y(0)) \cdot P(\delta y(0)).$$

For a discrete, zero-mean signal, A(t), composed of N data points, the ACF can be estimated as

$$ACF(\tau) \equiv \langle A(0)A(\tau) \rangle = \lim_{N \to \infty} \frac{1}{N} \sum_{i=1}^{N} A_i A_{i+\tau}$$

Thus, assuming normally distributed values of $A_i A_{i+\tau}$, the ACF and the associated precision is distributed as a student-T distribution, $T_\nu(x|\mu, \sigma^2)$ with ν degrees of freedom, as $T_{N-1}(ACF(\tau)|\bar{x}, (\sum_i (x_i - \bar{x})^2)/(N(N-1)))$, where $x_i = A_i A_{i+\tau}$, and $\bar{x} = \frac{1}{N} \sum_i A_i A_{i+\tau}$. This is derived from the posterior marginal of a normal-distribution likelihood with a normal-gamma prior with κ=0, α=−1/2, and β=0 (Murphy, 2007). The same formulas can be used to calculate an ACF from multiple signal trajectories that are independent and identically distributed; for instance the ACF at time τ would then include terms $A_i^0 A_{i+\tau}^0 + A_i^1 A_{i+\tau}^1 + \cdots + A_i^M A_{i+\tau}^M$, where the superscript denotes an index over *M* number of signal trajectories. We refer to this as the ensemble ACF.

For some specific types of signals, we can calculate an analytical form of the ACF for comparison to observed ACFs. First, if a time-dependent signal was stationary at zero and distributed according to $y(t) \sim \mathcal{N}(0, \sigma^2)$, where $\mathcal{N}$ denotes a normal distribution, and σ^2 is the variance, the $ACF(t) = \sigma^2 \cdot \delta(t)$, where

$\delta(t)$ is a Dirac delta function, which is one when t=0, and zero otherwise. Second, consider a zero-mean Markov chain with N states that are normally distributed according to $\mathcal{N}(\mu_i, \sigma_i^2)$ for the i^{th} state, but where $\langle y \rangle = 0$. Also, consider that the rate constant for the transition from the i^{th} state to the j^{th} state, k_{ij}, can be written as a rate constant matrix, K, with the associated Q matrix (K but where the on-diagonals are the negative sum of the rate constants out the i^{th} state). Notably, a vector of the probability of being in a particular state, P(t), given an initial probability vector, P(0), can then be written $P(t) = e^{Qt} \cdot P(0)$, where e denotes a matrix-exponential. If the Markov chain is in state i, the signal value at a particular time can be written $y(t) \sim \mu_i + N_i$, where $N_i \sim \mathcal{N}(0, \sigma_i^2)$. Thus, the ACF for this model is

$$\langle y(0)y(t) \rangle = \sum_i \sum_j \int_{N_i} \int_{N_j} (\mu_i + N_i)(\mu_j + N_j) \cdot P(j, t; i, 0; N_t; N_0),$$

where

$$P(j, t; i, 0; N_t; N_0) = P(j, t|i, 0) \cdot P_i(t = \infty) \cdot \mathcal{N}(0, \sigma_j^2) \cdot \mathcal{N}(0, \sigma_i^2).$$

Therefore

$$\langle y(0)y(t) \rangle = \sum_{i,j} \mu_i \mu_j \cdot d_j \cdot e^{Qt} \cdot d_i \cdot P_i(t = \infty) + \sum_i \sigma_i^2 \cdot P_i(t = \infty) \cdot \delta(t)$$

$$\equiv ACF_{MC}(t) + \langle \sigma^2 \rangle \cdot \delta(t),$$

where d_i is a vector with a 1 in position i, and ACF$_{MC}$ is the contribution from the Markov chain. Additionally, we note that $P(j, t|i, 0) \cdot P(i) = d_j \cdot e^{Qt} \cdot d_i \cdot P_i(t = \infty)$ can be rewritten with a spectral expansion as $\sum_i C_i e^{-\lambda_i t}$, where C_i is a weighting factor, and λ_i is an eigenvalue of e^{Qt}. Thus, for this Markov chain model with normally distributed noise, the ACF can be written as a weighted sum of N exponential decays with a term at t=0 corresponding to the average amount of uncorrelated, normally distributed noise. Therefore, for this model, the difference between ACF(t=0) and the next data point ACF(t=dt) is mostly $\langle \sigma^2 \rangle$, and the rest of the ACF is described by various exponential decays to zero.

In order to simulate single-molecule trajectories, we used the stochastic simulation algorithm (Gillespie, 1976) with the parameters in Table 1 with additive, normally distributed noise. From these 400 trajectories, we estimated global HMMs with one through six states as described above. As expected, the HMM with four states had the largest ELBO (Fig. 5–figure supplement 2). Additionally, we calculated the ACFs of these HMMs as described above. Notably, of these ACFs, only those for HMMs with four or more states are sufficient to match the individual and ensemble ACFs of the simulated trajectories (Fig. 5–figure supplement 2). Thus, the most parsimonious global HMM determined by this method also recapitulates the kinetic results of a model-free ACF method, but has the additional benefit as a model-based method of providing a mechanistic interpretation of those results.

Table 1. E_{FRET} trajectory simulation parameters

Parameter	Value	Units
μ_1	0.1	A.U.
μ_2	0.3	A.U.
μ_3	0.7	A.U.
μ_4	0.9	A.U.
k_{12}	0.25	sec^{-1}
k_{13}	0.50	sec^{-1}
k_{14}	1.00	sec^{-1}
k_{21}	0.50	sec^{-1}
k_{23}	1.00	sec^{-1}
k_{24}	0.25	sec^{-1}
k_{31}	1.00	sec^{-1}
k_{32}	0.25	sec^{-1}
k_{34}	0.50	sec^{-1}
k_{41}	0.25	sec^{-1}
k_{42}	0.50	sec^{-1}
k_{43}	1.00	sec^{-1}
σ	0.20	A.U.
τ	0.01	sec
n	2000	frames
N	400	trajectories

μ is an emission mean, k is a rate constant, σ is a standard deviation of additive noise, τ Is the time period of each data point, n is the length of each trajectory, N is the number of trajectories

Photobleaching detection. We detect photobleaching by using shape-based Bayesian model selection to choose the photobleaching time, or lack thereof, in a fluorescence intensity trajectory (Ray et al., 2021). Each model, M_T, that we choose between is similar in that before some time, T, the data points are assumed to be distributed according to a normal distribution with some mean, μ₁, and some precision, λ₁ (i.e., inverse variance λ=σ⁻²), which we do not care about, while the data points after time T are distributed according to a normal distribution with mean of zero, and some different precision, λ₂. We then calculate the probability of each M_T, having marginalized out the means and precisions so that the conclusions are relevant regardless of those values, and then use Bayesian inference to select the M_T that gives the best estimate of the photobleaching time as is supported by the available data. Practically, this method assigns all observed fluorescence at the beginning of a trajectory into the same 'on' state, usually with a large variance to account for different intensity states that compose the 'on' state, and then finds the last point where the data can be described as dropping to zero. For an ensemble of intensity trajectories, we perform model selection twice. The first time, we use uniform priors for the photobleaching time, while the second time, we use the independently calculated photobleaching times to infer an ensemble photobleaching rate constant, and then repeat the photobleaching time model selection calculation using this photobleaching rate constant to define the prior probability distribution for T for all of the independent intensity trajectories.

Mathematically, the likelihood for one of these models M_T is

$$P(\{x\}|\mu_1, \lambda_1, \lambda_2, T) = \prod_{t<T} \mathcal{N}(x_t|\mu_1, \lambda_1^{-1}) \prod_{t\geq T} \mathcal{N}(x_t|0, \lambda_2^{-1}),$$

where $\mathcal{N}(x|\mu, \sigma^2)$ is a normal distribution with mean μ, and variance σ². Using a prior probability distribution of

$$P(\mu_1, \lambda_1, \lambda_2) = NG(\mu_1, \lambda_1 | \alpha_1, \beta_1, \kappa_1, m_1) \cdot G(\lambda_2 | \alpha_2, \beta_2)$$

$$= \left[(2\pi)^{-\frac{1}{2}} (\kappa_1 \lambda_1)^{\frac{1}{2}} e^{-\frac{\kappa_1 \lambda_1}{2}(\mu_1 - m_1)^2} \cdot \beta_1^{\alpha_1} \Gamma(\alpha_1)^{-1} \lambda_1^{\alpha_1 - 1} e^{-\beta_1 \lambda_1} \right] \cdot \left[\beta_2^{\alpha_2} \Gamma(\alpha_2)^{-1} \lambda_2^{\alpha_2 - 1} e^{-\beta_2 \lambda_2} \right],$$

where NG is a joint normal-gamma distribution, and G is a gamma distribution,(Murphy, 2007) we obtain a posterior probability distribution of the same form but with updated hyperparameters (denoted by primes)

$$P(\mu_1, \lambda_1, \lambda_2 | \{x\}, T) = NG(\mu_1, \lambda_1 | \alpha_1', \beta_1', \kappa_1', m_1') \cdot G(\lambda_2 | \alpha_2', \beta_2'),$$

where

$$\alpha_1' = \alpha_1 + T/2$$

$$\beta_1' = \beta_1 + \frac{1}{2} \sum_{i=0}^{T} (x_i - \langle x \rangle)^2 + \frac{\kappa_1 T(\langle x \rangle - m_1)^2}{2(\kappa_1 + T)}$$

$$\kappa_1' = \kappa_1 + T$$

$$m_1' = \frac{\kappa_1 m_1 + T\langle x \rangle}{\kappa_1 + T}$$

$$\alpha_2' = \alpha_2 + (N - T)/2$$

$$\beta_2' = \beta_2 + \frac{1}{2} \sum_{i=T}^{N} x_i^2 ,$$

where N is the total number of data points. However, we do not particularly care about the exact values of μ_1, λ_1, or λ_2, so instead, we can marginalize them out to find the probability that of the observed data given the model M_T (i.e., the "evidence"), regardless of these nuisance parameters. In this case, we find

$$P(\{x\}|M_T) = P(\{x\}_{t<T}|M_T) \cdot P(\{x\}_{t\geq T}|M_T)$$

$$= \left[\frac{\Gamma(\alpha_1')}{\Gamma(\alpha_1)} \cdot \frac{\beta_1^{\alpha_1}}{\beta_1'^{\alpha_1'}} \cdot \left(\frac{\kappa_1}{\kappa_1'} \right)^{\frac{1}{2}} \cdot (2\pi)^{-\frac{T}{2}} \right] \left[\frac{\Gamma(\alpha_2')}{\Gamma(\alpha_2)} \cdot \frac{\beta_2^{\alpha_2}}{\beta_2'^{\alpha_2'}} \cdot (2\pi)^{-\frac{(N-T)}{2}} \right].$$

Now, with the evidence for the data given a photobleaching time, $P(\{x\}|M_T)$, we use Bayesian inference to select best photobleaching model (i.e., the model best supported by the data, regardless of the individual parameters) as the largest value of

$$P(M_T|\{x\}) = \frac{P(\{x\}|M_T) \cdot P(M_T)}{\sum_{T=0}^{N} P(\{x\}|M_T) \cdot P(M_T)}.$$

As discussed above, $P(M_T)$ is at first taken to be a constant and all trajectories are analyzed independently. Then, we infer the ensemble photobleaching rate constant using Bayesian inference with an exponential distribution as a likelihood, and a gamma distribution as the conjugate prior, and therefore also the posterior. As a result, we estimate $k_{PB} = (1 + n)/(1 + \sum_{i=0}^{n} T_i)$, where i indexes the different trajectories. Then, we use an exponential distribution as the prior $P(M_T) = k_{PB} e^{-k_{PB} T}$, where k_{PB} is treated as a deterministic value. Finally, we note that these model selection calculations are performed in log-space, and that we subtract the maximum of all the log terms (i.e., scale by a constant) from both the numerator and denominator in order the to avoid computer underflow.

Since the models used for model selection in this algorithm are not comprehensive in the sense that they do not account for every experimental nuance of the data, we believe it is best to manually examine the selected model and the underlying fluorescence intensity trajectory to ensure that they are reasonably well matched, and so that one can manually adjust the photobleaching time when they do not match due to unaccounted for experimental behaviors (*e.g.*, deviations from Gaussian emissions).

Calculation of Cycle Fluxes. For the mechanistic scheme for transport in Fig. 6, the following terms are necessary to calculate cycle fluxes to be calculated. For shorter notation, we use $t \equiv k_t$, $d' \equiv k_d'$, $d'' = k_d''$, $A_o \equiv k_a'[A]_o^m$, $A_i \equiv k_a'[A]_i^m$, $B_o \equiv k_a''[B]_o^n$, and $B_i \equiv k_a''[B]_i^n$. The cycle flux for any cycle is

$$J_\kappa = J_{\kappa+} - J_{\kappa-} = (\Pi_{\kappa+} - \Pi_{\kappa-})\,\Sigma_\kappa/\Sigma,$$

where

$$\Pi_{a+} - \Pi_{a-} = t^2 d'(A_o - A_i),$$
$$\Pi_{b+} - \Pi_{b-} = t^2 d''(B_o - B_i),$$
$$\Pi_{c+} - \Pi_{c-} = t^2 d'd''(A_o B_o - A_i B_i),$$
$$\Sigma_a = 2td'' + d''^2,$$
$$\Sigma_b = t(A_o + A_i) + A_o A_i,$$
$$\Sigma_c = 1,\text{ and}$$

$$
\begin{aligned}
\Sigma = \;& 2t^3[d''(4d' + 2A_i + A_o) + (2A_i B_i + A_o B_o + A_o B_i + B_o A_i)] \\
& + 2t^2\big[d''^2(2d' + A_i + A_o) \\
& + d''\big((2(A_o + A_i) + (B_o + B_i) + 2d')d' + (A_i B_i + B_o A_i + 2A_o A_i + A_o B_o + A_o B_i)\big) \\
& + \big((A_o B_o + A_i B_i)d' + (A_o B_o A_i + A_o A_i B_i + A_o B_o B_i + B_o A_i B_i)\big)\big] \\
& + 2t\big[d''^2\big((A_o + A_i)d' + A_o A_i + d'^2\big) + d''\big((A_o B_o + A_i B_i)d' + (A_o B_o A_i + A_o A_i B_i)\big) + (A_o B_o A_i B_i)\big].
\end{aligned}
$$

From these equations, the functional form of J_c for each first order rate constant, k, in Fig. 6 is $[ak + b + ck^{-1}]^{-1}$ where the coefficients *a*, *b*, and *c* depend on the other rate constants. This functional form has a maximum at $k = \sqrt{c/a}$, so there is an optimal value of k_t that maximizes transport cycle flux under for a given set of substrate concentrations. Finally, we note that Σ has 90 terms, and thus so does J_c. Due to the changes to the energy landscape in Fig. 7, 68 of the terms in J_c increase the magnitude of J_c, 18 decrease it, and 4 do not change it.

To find the optimal parameters to describe previously published succinate transport data, a target energy function of $\left(J^{expt} - (J_b + J_c)\right)^2$ was used for $J^{expt}([X] = K_M) = \frac{V_{max}}{2}$ and $J^{expt}([X] = 1000K_M) = V_{max}$ for the sodium titration in their Figure. 3 ($K_M = 41.7\,mM$, and $V_{max} = 53.5\,nmol\,mg^{-1}min^{-1} = 58.3\,s^{-1}$), and the succinate titration in their Figure. 6 ($K_M = 1.0\,\mu M$, and $V_{max} = 232.6\,nmol\,mg^{-1}min^{-1} = 253.4\,s^{-1}$) (Mulligan et al., 2014). The collective energy function for both experiments was minimized using an affine invariant MCMC ensemble sampler (Foreman-Mackey et al., 2013) with 200 walkers, a 1,000 step burn-in period, and a 10,000 step production period from which every 100th point was taken as a sample to avoid autocorrelation within the samples.

The steady-state probability of apo VcINDY in Buffer HX was calculated from the mechanism in Fig. 6 by setting $[B]_o = [B]_i = 0$, using the estimates for k_a', $\langle k \rangle \equiv k_t$, and k_d' (Fig. 6—Figure Supplement 2) and $[A]_o = [A]_i = [Na^+] = 0.1M$, to calculate the steady state probability $P_{apo} = P_{C_o} + P_{C_i}$ as

$$P_{apo} = \frac{3\langle k \rangle^3 + k_a'[Na^+]^3\langle k \rangle^2}{3\langle k \rangle^3 + 4k_a'[Na^+]^3\langle k \rangle^2 + k_a'^2[Na^+]^6\langle k \rangle}.$$

Crystallization of Wild Type VcINDY with Substrates. Crystals of VcINDY were grown with a protein concentration of 6 mg/ml at 4°C in hanging-drop vapor diffusion. For VcINDY-succinate, crystallization was carried out with 30% PEG 300, 5 mM sodium succinate, 200 mM sodium chloride, and 100 mM MES pH 5.8. VcINDY-fumarate crystals were grown with 37.5% PEG 300, 200 µM sodium fumarate, and 100 mM sodium acetate pH 4.6. For VcINDY-malate, crystals were grown with 25% PEG 400, 20 mM sodium malate, and 100 mM sodium acetate pH 5.0.

X-ray Data Collection and Structure Determination. Crystals were frozen in liquid nitrogen with the crystallization solution serving as the cryoprotectant. X-ray data were collected at the Advanced Photon Source Beamlines 19-ID and 19-BM. Data processing and scaling were performed using the HKL2000 software (Otwinowski and Minor, 1997). All crystals were of space group $P2_1$ with unit cell dimensions around a = 106 Å, b = 102 Å, c = 171 Å, β = 99°, and contained four molecules per asymmetric unit. The structures were determined by molecular replacement using the structure of VcINDY (PDB: 5UL9) (Nie et al., 2017) with sodium and citrate removed as the initial search model, followed by repeated cycles of model building in Coot (Emsley and Cowtan, 2004), and refinement in Phenix (Table 2) (Adams et al., 2010).

Data and Code Availability

VcINDY structures with fumarate, malate, and succinate bound were deposited in the PDB with IDs 6OKZ, 6OL0, and 6OL1, respectively. Code used to process smFRET data is described in a separate manuscript (in preparation, C.D.K. and R.L.G.) and available at https://github.com/GonzalezBiophysicsLab/XXXXXXXXXXXX.

Table 2. X-ray crystallography of VcINDY in complex with substrate

	Parameter	VcINDY-succinate	VcINDY-fumarate	VcINDY-malate
Data collection	Wavelength (Å)	0.9791	0.9794	0.9794
	Resolution range (Å)	50.0-3.09 (3.15-3.09)	50-3.29 (3.35-3.29)	50.0-3.50 (3.56-3.50)
	Space group	$P2_1$	$P2_1$	$P2_1$
	Cell dimensions: a, b, c (Å)	105.5, 102.5, 171.1	105.8, 102.1, 171.2	106.8, 102.4, 171.8
	Cell angle: β (°)	98.808	98.937	97.791
	Total reflections	128,621 (6,360)	99,521 (3,598)	86,871 (4,328)
	Unique reflections	65,778 (3,229)	51,510 (1,913)	44,866 (2,248)
	Multiplicity	5.0 (5.1)	6.6 (5.3)	6.9 (6.4)
	Completeness (%)	99.9 (100)	93.9 (71.0)	96.4 (96.9)
	$I/\sigma(I)$	10.3 (0.625)	10.1 (0.667)	10.8 (0.571)
	Wilson B (Å^2)	80.97	110.64	134.20
	R_{pim}	8.4 (152)	4.4 (54.2)	3.5 (61.5)
	$CC_{1/2}$	0.998 (0.443)	0.102 (0.512)	0.997 (0.604)
Refinement	R_{work}/R_{free} (%)	25.3% / 29.6%	24.9% / 28.5%	26.1% / 30.5%
	No. atoms - Protein/Ligands	13,325 / 40	13,097 / 23	12,792 / 42
	RMSD bond lengths (Å)	0.003	0.003	0.001
	RMSD bond angles (°)	0.66	0.52	0.39
	Ramachandran favored (%)	88.7	91.3	93.2
	Ramachandran allowed (%)	9.99	7.30	5.72
	Ramachandran outliers (%)	1.35	1.44	1.12
	Rotamer outliers (%)	0.07	0.0	0.0
	Clashscore	34.26	29.10	23.49

Numbers in parentheses are statistics for the highest resolution shell.

CRediT Authorship Contribution Statement

Colin D. Kinz-Thompson: Conceptualization, Methodology, Resources, Investigation, Validation, Formal Analysis, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization. **Marisa Lopez-Redondo:** Methodology, Resources, Investigation, Validation. **Christopher Mulligan:** Investigation, Validation, Formal Analysis, Visualization. **David B. Sauer:** Investigation, Validation, Formal Analysis, Data Curation, Visualization. **Jennifer J. Marden:** Resources, Validation. **Jinmei Song:** Resources, Validation. **Emad Tajkhorshid:** Conceptualization, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition. **John F. Hunt:** Conceptualization, Methodology, Resources, Supervision, Funding Acquisition. **David L. Stokes:** Conceptualization, Methodology, Resources, Supervision, Funding Acquisition. **Joseph A. Mindell:** Conceptualization, Methodology, Resources, Data Curation, Writing - Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition. **Da-Neng Wang:** Conceptualization, Methodology, Resources, Data Curation, Writing - Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition. **Ruben L. Gonzalez, Jr.:** Conceptualization, Methodology, Resources, Data Curation, Writing - Review & Editing, Visualization, Supervision, Project Administration, Funding Acquisition.

Acknowledgements

R.L.G. acknowledges support from the National Institutes of Health (NIH) through grants R01-GM084288 and R01-GM137608. D.N.W. acknowledges support from the NIH through grants R01-GM121994, R01-NS108151, R01-DK099023 and the G. Harold and Leila Y. Mathers Foundation. J.M. and C.M. acknowledge support from the National Institute of Neurological Disorders and Stroke (NINDS) Intramural Program, and C.M. acknowledges support from the Biotechnology and Biological Sciences Research Council through grant BB/V007424/1. D.L.S. acknowledges support from NIH through grant R35-GM144109. J.F.H. acknowledges support from NIH through grants R01-GM127883. E.T. acknowledges support from NIH through grants P41-GM104601 and R01-GM123455. D.B.S. acknowledges support, in part, from an American Cancer Society Postdoctoral Fellowship (PF-17-135-01) and from the Office of the Assistant Secretary of Defense for Health Affairs, through the Peer Reviewed Cancer Research Program under Award No. W81XH-16-1-0153. Crystal screening and X-ray data collection were carried out at NSLS-II Brookhaven National Laboratory and at Argonne National Laboratory, Structural Biology Center (SBC) at the Advanced Photon Source. SBC-CAT is operated by Univ. Chicago Argonne, LLC, for the U.S. Department of Energy, Office of Biological and Environmental Research under contract DE-AC02-06CH11357. Electron microscopy was performed at the NYU School of Medicine Microscopy Core.

References

Adams PD, Afonine PV, Bunkóczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung L-W, Kapral GJ, Grosse-Kunstleve RW, McCoy AJ, Moriarty NW, Oeffner R, Read RJ, Richardson DC, Richardson JS, Terwilliger TC, Zwart PH. 2010. PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**:213–221. doi:10.1107/S0907444909052925

Akyuz N, Altman RB, Blanchard SC, Boudker O. 2013. Transport dynamics in a glutamate transporter homologue. *Nature* **502**:114–118. doi:10.1038/nature12265

Bergeron MJ, Clémençon B, Hediger MA, Markovich D. 2013. SLC13 family of Na+-coupled di- and tri-carboxylate/sulfate transporters. *Mol Aspects Med* **34**:299–312. doi:10.1016/j.mam.2012.12.001

Berne BJ, Pecora R. 2000. Dynamic Light Scattering: With Applications to Chemistry, Biology, and Physics, Unabridged edition. ed. Mineola, N.Y: Dover Publications.

Bronson JE, Fei J, Hofman JM, Gonzalez RL, Wiggins CH. 2009. Learning rates and states from biophysical time series: a Bayesian approach to model selection and single-molecule FRET data. *Biophys J* **97**:3196–205. doi:10.1016/j.bpj.2009.09.031

Denisov IG, Grinkova YV, Lazarides AA, Sligar SG. 2004. Directed Self-Assembly of Monodisperse Phospholipid Bilayer Nanodiscs with Controlled Size. *J Am Chem Soc* **126**:3477–3487. doi:10.1021/ja0393574

Denisov IG, Sligar SG. 2017. Nanodiscs in Membrane Biochemistry and Biophysics. *Chem Rev* **117**:4669–4713. doi:10.1021/acs.chemrev.6b00690

Drew D, Boudker O. 2016. Shared Molecular Mechanisms of Membrane Transporters. *Annu Rev Biochem* **85**:543–72. doi:10.1146/annurev-biochem-060815-014520

Edelstein AD, Tsuchida MA, Amodaj N, Pinkard H, Vale RD, Stuurman N. 2014. Advanced methods of microscope control using μManager software. *J Biol Methods* **1**:10. doi:10.14440/jbm.2014.36

Emsley P, Cowtan K. 2004. Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* **60**:2126–2132. doi:10.1107/S0907444904019158

Erkens GB, Hänelt I, Goudsmits JMH, Slotboom DJ, van Oijen AM. 2013. Unsynchronised subunit motion in single trimeric sodium-coupled aspartate transporters. *Nature* **502**:119–123. doi:10.1038/nature12538

Felle H, Porter JS, Slayman CL, Kaback HR. 1980. Quantitative measurements of membrane potential in Escherichia coli. *Biochemistry* **19**:3585–3590. doi:10.1021/bi00556a026

Fersht A. 2017. Structure and Mechanism in Protein Science: A Guide to Enzyme Catalysis and Protein Folding. WORLD SCIENTIFIC. doi:10.1142/10574

Fitzgerald GA, Mulligan C, Mindell JA. 2017. A general method for determining secondary active transporter substrate stoichiometry. *eLife* **6**:e21016. doi:10.7554/eLife.21016

Foreman-Mackey D, Hogg DW, Lang D, Goodman J. 2013. emcee: The MCMC Hammer. *Publ Astron Soc Pac* **125**:306–312. doi:10.1086/670067

Forrest LR, Krämer R, Ziegler C. 2011. The structural basis of secondary active transport mechanisms. *Biochim Biophys Acta Bioenerg* **1807**:167–188. doi:10.1016/j.bbabio.2010.10.014

Garaeva AA, Slotboom DJ. 2020. Elevator-type mechanisms of membrane transport. *Biochem Soc Trans* **48**:1227–1241. doi:10.1042/BST20200290

Gillespie DT. 1976. A general method for numerically simulating the stochastic time evolution of coupled chemical reactions. *J Comput Phys* **22**:403–434. doi:10.1016/0021-9991(76)90041-3

Haran G. 2004. Noise reduction in single-molecule fluorescence trajectories of folding proteins. *Chem Phys* **307**:137–145. doi:10.1016/j.chemphys.2004.05.017

Hill TL. 2005. Free energy transduction and biochemical cycle kinetics. Mineola, N.Y: Dover Publications.

Hon J, Gonzalez RL. 2019. Bayesian-Estimated Hierarchical HMMs Enable Robust Analysis of Single-Molecule Kinetic Heterogeneity. *Biophys J* **116**:1790–1802. doi:10.1016/j.bpj.2019.02.031

Jardetzky O. 1966. Simple allosteric model for membrane pumps. *Nature* **211**:969–970. doi:10.1038/211969a0

Kim JW, Kim Subin, Kim Songwon, Lee H, Lee J, Jin MS. 2017. Structural insights into the elevator-like mechanism of the sodium / citrate symporter CitS. *Sci Rep* **7**:2548. doi:10.1038/s41598

Kinz-Thompson CD, Ray KK, Gonzalez RL. 2021. Bayesian Inference: The Comprehensive Approach to Analyzing Single-Molecule Experiments. *Annu Rev Biophys* **50**:191–208. doi:10.1146/annurev-biophys-082120-103921

Lambert TJ. 2019. FPbase: a community-editable fluorescent protein database. *Nat Methods* **16**:277–278. doi:10.1038/s41592-019-0352-8

Lo C-J, Leake MC, Berry RM. 2006. Fluorescence Measurement of Intracellular Sodium Concentration in Single Escherichia coli Cells. *Biophys J* **90**:357–365. doi:10.1529/biophysj.105.071332

Lolkema JS, Slotboom DJ. 2019. Models to determine the kinetic mechanisms of ion-coupled transporters. *J Gen Physiol* **151**:369–380. doi:10.1085/jgp.201812055

Maity S, Trinco G, Buzón P, Anshari ZR, Kodera N, Ngo KX, Ando T, Slotboom DJ, Roos WH. 2022. High-speed atomic force microscopy reveals a three-state elevator mechanism in the citrate transporter CitS. *Proc Natl Acad Sci* **119**:e2113927119. doi:10.1073/pnas.2113927119

Mancusso R, Gregorio GG, Liu Q, Wang D-N. 2012. Structure and mechanism of a bacterial sodium-dependent dicarboxylate transporter. *Nature* **491**:1–6. doi:10.1038/nature11542

Markovich D. 2012. Sodium-Sulfate/Carboxylate Cotransporters (SLC13)Current Topics in Membranes. Elsevier. pp. 239–256. doi:10.1016/B978-0-12-394316-3.00007-7

Mulligan C, Fenollar-Ferrer C, Fitzgerald GA, Vergara-Jaque A, Kaufmann D, Li Y, Forrest LR, Mindell JA. 2016. The bacterial dicarboxylate transporter VcINDY uses a two-domain elevator-type mechanism. *Nat Struct Mol Biol* **23**:256–263. doi:10.1038/nsmb.3166

Mulligan C, Fitzgerald GA, Wang D-N, Mindell JA. 2014. Functional characterization of a Na+-dependent dicarboxylate transporter from Vibrio cholerae. *J Gen Physiol* **143**:745–759. doi:10.1085/jgp.201311141

Murphy KP. 2007. Conjugate Bayesian analysis of the Gaussian distribution (Technical Report). University of British Columbia.

Nie R, Stark S, Symersky J, Kaplan RS, Lu M. 2017. Structure and function of the divalent anion/Na+ symporter from Vibrio cholerae and a humanized variant. *Nat Commun* **8**:15009. doi:10.1038/ncomms15009

Otwinowski Z, Minor W. 1997. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol* **276**:307–326.

Ray KK, Verma AR, Gonzalez Jr RL, Kinz-Thompson CD. 2021. Inferring the shape of data: A probabilistic framework for analyzing experiments in the natural sciences. *ArXiv210912462 Phys Q-Bio*.

Ruan Y, Miyagi A, Wang X, Chami M, Boudker O, Scheuring S. 2017. Direct visualization of glutamate transporter elevator mechanism by high-speed AFM. *Proc Natl Acad Sci* **114**:1584–1588. doi:10.1073/pnas.1616413114

Sampson CDD, Fàbregas Bellavista C, Stewart MJ, Mulligan C. 2021. Thermostability-based binding assays reveal complex interplay of cation, substrate and lipid binding in the bacterial DASS transporter, VcINDY. *Biochem J* **478**:3847–3867. doi:10.1042/BCJ20210061

Sampson CDD, Stewart MJ, Mindell JA, Mulligan C. 2020. Solvent accessibility changes in a Na+-dependent C4-dicarboxylate transporter suggest differential substrate effects in a multistep mechanism. *J Biol Chem* **295**:18524–18538. doi:10.1074/jbc.RA120.013894

Sauer DB, Marden JJ, Sudar JC, Song J, Mulligan C, Wang D-N. 2022a. Structural basis of ion – substrate coupling in the Na+ -dependent dicarboxylate transporter VcINDY. *bioRxiv*. doi:10.1101/2022.01.11.475879

Sauer DB, Song J, Wang B, Hilton JK, Karpowich NK, Mindell JA, Rice WJ, Wang D-N. 2021. Structure and inhibition mechanism of the human citrate transporter NaCT. *Nature* **591**:157–161. doi:10.1038/s41586-021-03230-x

Sauer DB, Trebesch N, Marden JJ, Cocco N, Song J, Koide A, Koide S, Tajkhorshid E, Wang D-N. 2020. Structural basis for the reaction cycle of DASS dicarboxylate transporters. *eLife* **9**:e61350. doi:10.7554/eLife.61350

Sauer DB, Wang B, Sudar JC, Song J, Marden J, Rice WJ, Wang D. 2022b. The ups and downs of elevator-type di-/tricarboxylate membrane transporters. *FEBS J* **289**:1515–1523. doi:10.1111/febs.16158

Stein WD, Lieb WR. 1986. Transport and diffusion across cell membranes. Orlando: Academic Press.

van de Meent J-W, Bronson JE, Wiggins CH, Gonzalez RL. 2014. Empirical Bayes Methods Enable Advanced Population-Level Analyses of Single-Molecule FRET Experiments. *Biophys J* **106**:1327–1337. doi:10.1016/j.bpj.2013.12.055

Figures

Figure 1. VcINDY transport cycle. (A) Diagram of VcINDY dimer (green) reversibly transporting sodium ions (Na^+) and dicarboxylate substrate (D^{2-}) into and out of a cell with a stoichiometry of 3:1; an inward facing Na^+ electrochemical gradient provides the free energy for vectorial transport. (B) Minimal mechanistic model of VcINDY transport cycle. Outward- and inward-facing conformations of the VcINDY homodimer (C_o, and C_i, respectively) bind Na^+ and D^{2-} in an unknown order along the horizontal transitions; double-prime signifies both co-substrates are bound. VcINDY protomer conformational motions along the vertical transitions are involved in substrate translocation. At least one closed transport cycle, such as shown here in the clockwise direction in blue, is required for vectorial transport. Conformations of each protomer in the C_o and C_i states are unknown. Slippage reactions occur when an incomplete set of co-substrates is moved across the membrane.

Figure 2. Cartoon diagram of smFRET pairs designed to report on the VcINDY translocation reaction for VcINDY^{ext} (top) and VcINDY^{int} (bottom). The distance between each labeling position, *r*, was determined from the C_α-C_α distance, and was measured for a C_o, substrate-bound repeat-swap model (C_o-C_o, PMDB: PM0080216), and a substrate-bound, C_i X-ray crystallography structure (C_i-C_i, PDB: 5ULE). E_{FRET} was estimated using R_0=51 Å. Scaffold domains are shown in gray, and transport domains are shown in green.

Figure 2–Figure Supplement 1. Cartoon schematic of approach to yield donor- and acceptor- fluorophore-labeled VcINDY homodimers. Site-directed mutagenesis is used to remove all codons encoding cysteine from a plasmid-based "wild type" VcINDY gene (1) to yield a "cys-less" variant (2), and used further to introduce a single codon encoding cysteine (3). This is over-expressed in *Escherichia coli* (*E. coli*), and purified to yield homodimers with single-cysteine containing protomers (4). The homodimers are labeled with a 1:1 ratio of maleimide-conjugated donor- : acceptor-fluorophore (5). Only those homodimers that are labeled with both a single donor- and single acceptor-fluorophore are computationally selected after imaging.

Figure 2–Figure Supplement 2. VcINDY biochemical transport assays. Transport rates of proteoliposome-embedded VcINDY homodimers for the external (left) and internal (right) smFRET pair signal. Wild type (*i.e.*, cysteine-containing) assays were performed as control separately for each labeling position.

Figure 3. Schematic diagram of nanodisc-embedded VcINDY homodimer construct. Single cysteine mutant protomers labeled with a 1:1 ratio of donor:acceptor fluorophores are mixed with biotinylated membrane scaffold protein (MSP) and *E. coli* polar lipids to make nanodisc-embedded VcINDY homodimers, which are then tethered to a flow cell surface *via* a biotin-streptavidin-biotin bridge, and imaged using TIRF microscopy. The scaffold domains are shown in gray, and transport domains are shown in green.

Figure 3–Figure Supplement 1. Negative stain electron microscopy of VcINDY reconstituted into lipid nanodiscs. Both top and side views of nanodiscs were seen. The nanodiscs had a diameter of ~110 Å and each contained one VcINDY dimer. The sample was uniform and no aggregation was observed.

Figure 4. Equilibrium smFRET from VcINDY^{ext} (left) and VcINDY^{int} (right) in the presence of sodium and succinate (buffer HS). Example donor (green) and acceptor (red) fluorescence intensity trajectories, with corresponding E$_{FRET}$ trajectories (blue) and idealized trajectories (black) are shown. Several trajectories shown include donor fluorescence intensity after the acceptor fluorophore has photobleached, but this was removed prior to further analysis.

Figure 5. Substrate dependence of VcINDY E_{FRET} distributions. Histograms of E_{FRET} values from all external (left) and internal (right) smFRET signals recorded with VcINDY[ext], and VcINDY[int], respectively, in different buffer conditions. The E_{FRET} data in these histograms were not filtered, and E_{FRET} values corresponding to donor photobleaching have been removed (*i.e.*, the last dwell in an E_{FRET} state of zero). The predictive posterior probability distribution for E_{FRET} from the global HMM with the maximum ELBO is shown overtop (solid black) as a heuristic measure of how well the global HMM explains the entirety of each dataset, but this ignores the kinetic aspects of the HMM. The distributions for the individual, consensus E_{FRET} states from the global HMM are shown as black dashed lines.

Figure 5–Figure Supplement 1. Directed acyclic graph for a global, variational Bayes HMM. Pink nodes represent hyperparameters, grey nodes represent unobserved variables, and blue nodes represent observed variables. T_j is the number of data points in the j^{th} trajectory, and $D_{i,j}$ corresponds to the i^{th} data point of the j^{th} trajectory. Arrows represent conditional dependencies.

Figure 5–Figure Supplement 2. Autocorrelation function analysis of global HMMs with various number of hidden states of simulated, four-state E_{FRET} trajectories. Individual ACFs (black), ensemble ACF (blue), and global HMM ACF (red). The maximum ELBO HMM (n=4; ln(ELBO)= −1.1x10^4) matches the number of E_{FRET} states in the simulation.

Figure 5–Figure Supplement 3. Substrate dependence of evidence lower bound (ELBO) calculations for global HMMs of VcINDY smFRET experiments. The maximum ELBO model is the HMM that best represents the set of E_{FRET} trajectories for those substrate conditions.

Figure 5–Figure Supplement 4. Substrate dependence of the HMM with maximum ELBO for global HMMs of VcINDY smFRET experiments. For each kinetic scheme, states are denoted by the mean E_{FRET} value of that state, which was inferred from the HMM. Rate constants for transitions between states are shown above the arrows.

Figure 6. Free energy transduction model for a secondary active transporter. First order and pseudo-first order rate constants for the parsimonious kinetic model discussed in the text are given over each transition. For simplicity, co-substrate binding has been reduced to a strictly sequential, two-step process with mirror symmetry (Stein and Lieb, 1986). Substrate A is the first co-substrate to bind with stoichiometry m; co-substrate B binds second with stoichiometry n. Prime signifies the first co-substrate is bound; double-prime signifies both co-substrates are bound; C_o and C_i denote the outward and inward facing conformation of the transporter. k_t is the translocation rate constant, k_a is a second order association rate constant, and k_d is a dissociation rate constant. Cycles *a* (red) and *b* (green) are responsible for slippage, while cycle *c* (blue) is responsible for transport. The positive direction is defined to be clockwise.

Figure 6—Figure Supplement 1. MCMC trajectories for optimal kinetic model parameters of VcINDY proteoliposome-based succinate transport assay. (A) Affine invariant ensemble MCMC sampler trajectories for 200 walkers during burn-in period. These samples were discarded. (B) Production period trajectories from 200 walkers. Only every 100th sample of those shown here was used to minimize autocorrelation.

Figure 6—Figure Supplement 2. Corner plot of MCMC samples for optimal kinetic model parameters of VcINDY proteoliposome-based succinate transport assay. Ranges given are the 1σ level containing 39.3% of the sample volume.

Figure 6—Figure Supplement 3. Modeled VcINDY transport rate experiments. Data from Mulligan and coworkers were reproduced from their reported K_M, V_{max}, and Hill coefficients (Mulligan et al., 2014). Modeled data are the 95% credible intervals generated from MCMC samples of k_a', k_a'', and $\langle k \rangle$, and dashed lines are the lowest energy solution.

Figure 7. Schematic of optimized energy landscape for binding co-substrates A and B. Original energy levels are denoted with dashed lines for apo- (C), partially bound (C'), and fully bound (C'') transporter. Direction of perturbations to energy levels by amounts $\Delta G'$ and $\Delta G''$ are shown with arrows. Relative positioning of unperturbed C, C', and C'', and transition states ‡' and ‡'' are arbitrarily chosen, as are the magnitudes of $\Delta G'$ and $\Delta G''$.

Figure 7—Figure Supplement 1. Estimated physiological Na+ and succinate concentrations in a bacterial cell in different environments. External (ext) and internal (int) Na+ concentrations were measured for another gammaproteobacteria, *Escherichia coli* (*E. coli*), at pH=7 (Lo et al., 2006). For E. coli at pH=7, the membrane potential difference ($\Delta\psi$) is approximately -100 mV (Felle et al., 1980). With these numbers, balancing the chemical potential difference and effects of charge for the VcINDY transport reaction at equilibrium provides the internal to external ratio of succinate concentrations for the cell (orange). When $\Delta\psi = 0$, the cell is depolarized (dashed).

Figure 8. View of the VcINDY binding pocket from the cytosol. The X-ray crystallography derived structural model is of succinate-, and Na$^+$-bound VcINDY (green). Electron density for the substrates is shown contoured at 2.5σ.