Integrated viral elements unveil the dual lifestyle of *Tetraselmis* spp.
polinton-like viruses

Chase EE[1,2], Desnues C[1,2], and Blanc G[1].

[1] Microbiologie Environnementale Biotechnologie, Institut Méditerranéen
d'Océanologie, Campus de Luminy, 163 Avenue de Luminy, 13009 Marseille, France

[2] Institut hospitalo-universitaire (IHU) Méditerranée infection, 19-21 Boulevard Jean
Moulin, 13005 Marseille, France

Abstract

In this study, we aimed at exploring horizontal gene transfer between viruses and
Chlorodendraceae green algae (Chlorophyta) using available genomic and
transcriptomic sequences for 20 algal strains. We identified a significant number of
genes sharing a higher sequence similarity with viral homologues, thus signaling their
possible involvement in HGTs with viruses. Further characterization showed that many
of these genes were clustered in DNA regions of several tens to hundreds of kilobases
in size, originally belonging to viruses related to known *Tetraselmis* spp. viruses (TetV
and TsV). In contrast, the remaining candidate HGT genes were randomly dispersed in
the algal genomes, more frequently transcribed and belonged to large multigene
families. The presence of homologs in *Viridiplantae* suggested that these latter were
more likely of algal rather than viral origin. We found a remarkable diversity Polinton-
like virus elements inserted in Tetraselmis genomes, all of which were most similar to
the *Tetrasemis striata* virus TsV. The genes of PLV-like elements are transcriptionally
inactive with the notable exception of the homologue of the TVSG_00024 gene of TsV
whose function is unknown. We suggest that this gene may be involved in a sentinel
process to trigger virus reactivation and excision in response to an environmental
stimulus. Altogether, these results provide evidence that TsV-related viruses have a
dual lifestyle, alternating between a free viral phase (i.e., virion) and a phase
integrated into host genomes.

Keywords: polintons-like viruses, polintons, transposons, microalgae, genome
integration, *Tetraselmis* spp.

Abbreviations: MCP; major capsid protein, mCP; minor capsid protein, PLV; polintons-
like virus, TsV-N1; T. striata virus, PgVV; Phaeocystis globasa virus virophage, ORF;
open reading frame, TIRs; terminal inverted repeats,

Introduction

Polintons (also known as Maverick) were first recognized as large transposable elements (<40Kb in size) encoding up to 10 different proteins, found in the genomes of both unicellular and multi-cellular eukaryotes (1). They generally encode both a DNA polymerase ("POL") and a retrovirus-like integrase ("INT"), thereby how they acquired their name "POLINTons" and their classification as self-synthesizing transposons (2). Many encode a DNA packaging ATPase, major and minor jelly-roll capsid proteins (MCP and mCP respectively) and a capsid maturation protease (PRO) (3). They were therefore predicted to alternate between the transposon and viral lifestyles although virion formation remains to be demonstrated. Polintons are considered the ancient ancestors of most eukaryotic dsDNA viruses (4). There are considerable similarities between polintons and 2 groups of dsDNA viruses: the virophages and polinton-like viruses (PLVs). Virophages involved in tripartite infections with giant viruses (phylum *Nucleocytoviricota)* and their cellular hosts (5, 6). Virophages can parasitize giant DNA viruses and may provide adaptive anti-giant virus defence in unicellular eukaryotes (7–9). Virophage genomes (15–30 kbp) typically encode a MCP, a mCP, a DNA packaging ATPase, and a PRO (10). They also encode DNA replication and integration proteins that were likely acquired independently in different virophage lineages (11). Virophages can integrate into the genome of the cellular or viral host (12–14), in which case they resemble Potintons, and and reactivates upon superinfection with a giant virus (8). Virophages can therefore have a dual life style combining a virus stage with an integrated stage The PLVs are also similar in size and genomic composition to the Polintons. They have mainly been discovered in sequencing datasets and include members integrated into algal genomes (i.e. Monoraphidium neglectum, and Guillardia theta) or associated with giant virus particles (3, 15, 16). The majority of PLVs were however recovered from metagenomes of viral fractions (i.e. filtration has occurred to target particles <0,22 μm, therefore microeukaryotes have been virtually removed while viral particles remain), suggesting that PLVs occur abundantly as free viruses in the environment, especially in aquatic ecosystems (17). The PLV genomes generally do not contain all of the hallmark genes found in polintons, such as the DNA polymerase and retrovirus-like integrase, but can include ATPases, mCPs, and MCPs, and typically terminal inverted repeats at either end of their genome (3). Furthermore, PLVs lack the capsid maturation protease normally present in both polintons and virophages. The presence of mCPs and MCP allude to the concept of a "dual lifestyle" among PLVs as well. The first infectious PLV particle was isolated from the cosmopolitan green microalgae *Tetraselmis striata* (phylum *Chlorophyta*; family *Chlorodendraceae*) (18). The

Tetraselmis striata virus TsV-N1 has a 31-kb genome and is estimated to release
thousands of 60 nm virions per algal host cell. The TsV-N1 host infectivity range was
tested on several *Tetraselmis* spp., and was only found to infect the species it was
isolated from (18), highlighting the narrow host specificity of this PLV. Whether TsV-N1
also exist as an integrated form in the host genome has not yet been tested because
the *T. striata* genome was not sequenced at the time of the PLV discovery. Tentative
classifications of PLVs based on shared gene content and major capsid protein
phylogeny, identified 3 to 5 major groups including the TSV group, which contains TsV-
N1 (3, 17). Furthermore, two other viruses have been isolated from Tetraselmis spp.,
including TetV, a member of Mimiviridae ("giant viruses"; phylum *Nucleocytoviricota*)
with a 668-Kb genome (19). The 39Kb genome of a third potential *Tetraselmis* virus,
named *Tetraselmis viridis* virus S20 (TvV-S20), is available in Genbank since 2010 and
is unrelated to TsV-N1 and TetV (20). The genetic content of this virus resembles that
of an ordinary bacteriophage, suggesting that TvV-S20 could rather correspond to a
bacteriophage associated with bacteria in co-culture with Tetraselmis (21).

Until recently algal genomes of close taxonomic proximity to Tetraselmis were not
available. At present, public databases contain draft genomes for several *Tetraselmis*
species as well as for additional *Chlorodendraceae* genera, including *Prasinocladus*
and *Platymonas* (22, 23). Transcriptomic data for several Tetraselmis strains and
Scherffelia dubia were also made available through various sequencing programs (24–
28). In this study we took the opportunity to explore both algal draft genomes and
transcriptomic data for evidence of TsV-N1 or other closely related PLVs and other
viral elements potentially integrated into algal genomes. More specifically our
motivation was to find clues supporting the hypothesis that PLVs related to TsV-N1
might have a dual lifestyle alternating integrated and free phases as for virophages.
By searching the genomic data for viral-derived coding sequences, we identified an
unexpected diversity of integrated PLV sequences closely related to TsV-N1 in all
sequenced *Chlorodendraceae* strains. In addition, we highlight large genome
fragments of giant TetV-related viruses in the *T. striata* and *T. suecica* genomes with
remarkably different GC contents than the original TetV.

Materials and Methods

Acquisition of genomic and transcriptomic data from family Chlorodendraceae

Six draft genome assemblies for *T. striata* (strains LANL1001 and UTEXSP22), *T.*
suecica (strain CCMP904 sequenced independently by New York University Abu Dhabi
[NYUAD] and Institut Français de Recherche pour l'Exploitation de la MER [IFREMER]),
T. subcordiformis and *Prasinocladus* sp. malaysianus were retrieved from the National

Center for Biotechnology Information (NCBI) GenBank, the Dryad database and the
IFREMER data repository server. Also, transriptomic data (Illumina RNA-seq reads) for
15 Chlorodendraceae strains, including *Scherffelia dubia* and several *Tetraselmis* spp.,
were retrieved from Genbank. Algal strain names and identifying information for
retrieving sequence data are given in **Table S1**. Genbank accession numbers of the
virus genomic sequences used in this study are KY322437 for TetV, NC_020869 for
TsV-N1, and NC_020840 for TvV-S20.

Note that the *T. suecica* CCMP904 assemblies released by NYUAD and IFREMER were
quite dissimilar. The NUYAD assembly was generated using Illumina short reads, and
contained a large number of contigs (N=392,918) of relatively short length (<20Kb,
L50=868bp), totaling 237 Mb (23). In contrast, the IFREMER assembly was generated
using Oxford Nanopore Technologies (ONT) long reads, and contained a much smaller
number of contigs (N=175) of greater sizes (L50=4.3Mb), totaling 100 Mb. Mapping of
39 million NYUAD short reads on the IFREMER assembly using HISAT2 (29) resulted in
30% of overall alignment rate. Likewise, 27% of the NYUAD assembly sequence found
a significant BLASTN match (evalue 1e-100) in the IFREMER assembly. Conversely 65%
of the IFREMER assembly sequence was aligned on the NYUAD assembly. Thus the two
assemblies had only partial overlap and contained a substantial amount of assembly-
specific sequences.

Sequence analysis

Transcriptome raw reads were first trimmed for low-quality and adaptor sequences
using AlienTrimmer (30), then assembled into transcript sequences using SPADES
v3.15.3 (31) with the flag "--rna"and default parameters otherwise. Multiple RNA-seq
datasets from the same strain were pooled before assembly. Finally, reconstructed
transcripts were clustered based on nucleotide similarity (>95%) using CDHIT-EST (32)
to remove redundancy due to alternative splicing and highly conserved paralogues.
The longest sequence of each sequence cluster was kept for further analysis.

For each assembly (transcriptome or genome), we generated a collection of all open
reading frames >150 codons (see **Table S2** for assembly and ORFing statistics). While
the typical structure of eukaryotic algal genes contains introns, the use of an ORFing
approach (i.e. without consideration of introns that interrupt the reading phase) in the
present genomes is not a problem because viral genes do not have introns, which is
generally true for genes brought by viral genome insertions. Translated ORFs were
aligned against the TrEMBL public database using MMSEQS (e-value <1e-5) (33) and
the best matches were recorded. We did not consider best matches to uncultured
organism or environmental sequence because the nature of the corresponding
organism is often uncertain. We also disregarded matches with *Chlorodendraceae*

species to allow detection of putative viral HGT events that occurred in a
Chlorodendraceae ancestor. When a translated ORF returned a best match to one of
these sequences, we recorded the next highest scoring match to a clearly identified
non-*Chlorodendraceae* organism or virus. Finally, ORFs with a recorded best match to
a virus ("VBM" will be used hereafter in place of "viral best match") were considered
as potentially involved in HGT with a virus and were analyzed further. Note that this
approach may miss some candidates when the best match is a viral insertion in the
genome of a cellular organism. Protein family clustering was done using the MMSEQS
easy-cluster workflow by setting the -c parameter to 0.2 to account for truncated
sequences.
Transcriptomic analysis was performed using the HISAT2 program (29) with default
parameters to map RNA-seq reads onto the reference genomes. The resulting
alignments were used to calculate the FPKM values (Fragments Per Kilobase Million)
for gene, which are used as an estimation of the relative transcription level in the cell.

Identification of putative viral elements based on VBM identification

Viral elements were defined as clusters of candidate viral genes co-localized on a
genomic contig or segment of thereof. We applied two types of methods for detecting
these clusters. The sliding-window approach consisted of sliding a 10-Kb window along
the contig with a 1-Kb step and counting the number of VBM ORFs in each window. We
systematically investigated genomic regions returning windows that had at least four
candidate viral genes. This approach allowed to identify large viral inserts of TetV-like
DNA. In the second approach, which proved to be more effective in identifying
conserved PLV elements (shorter insertions), the MCP of TsV-N1 was used as "bait" to
recover all potential MCPs from each of the algal genomes. Then, 20 Kb upstream and
downstream of algal MCP gene were searched for additional ORFs with a best match to
the TsV-N1 genome or to a collection of PLVs genomes gathered in Yutin et al. (3).
Finally, sequences of 3 Kb following the last matches against the PLVs on the left and
on the right were aligned by BLASTN to identify possible direct or reverse repeats (i.e.
terminal inverted repeats; TIRs) likely to mark the limits of the PLV elements.

Phylogenetic placement of relevant algal strains by RBCL and VBMs by MCP genes

A phylogenetic tree of the algal strains used in this study was reconstructed based on
the RBCL coding sequence (large subunit of ribulose bisphosphate carboxylase)
identified in Genbank NR and in the assembled datasets using BLAST (**Figure S1**). A
multiple sequence alignment of the RBCL genes was generated with MAFFT (34),
followed by a maximum likelihood phylogenetic reconstruction using the FastTree
program (35) and the GTR+G substitution model. Statistical support for branches was

estimated using the Shimodaira-Hasegawa test as implemented in FastTree. The RBCL
tree showed 2 apparent incongruences between species names and phylogenetic
position. The first one concerns *T. suecica* PLY305 which clusters within the *T. striata*
clade while other algae referenced under the *T. suecica* species name grouped in their
own clade. The corresponding transcriptomic sequences came from the oneKp project
(25), a large-scale sequencing program, and it is possible that there has been a
confusion in the identification of the sequenced material. There was also a fairly large
phylogenetic distance between *T. subcordiformis* FACHB-1751 and UTEX171, which
suggests that these two organisms did not belong to the same species. The major
capsid protein (MCP) based trees were produced using the same procedure except
that we used the WAG+G substitution model adapted to protein sequences.

Results

Virus-like genes recovered from algal genomes of family Chlorodendraceae

ORFs with a viral best match (VBM) in TrEMBL were identified in the genome
assemblies of *T. striata, T.suecica, P. subcordiformis*, and *Prasinocladus* sp.
malaysianus. The number of these ORFs ranged from 51 for *Prasinocladus* sp.
malaysianus, to 1148 for *T. striata* LANL1001. Overall VBMs were primarily from
proteins of the Tetraselmis infecting viruses including TsV-N1 and TetV (see **Figure**
1A). However, we obtained no best hits to TvV-S20 for all tested species. A smaller
fraction of VBMs were from members of families *Phycodnaviridae*, *Mimiviridae*, and
Lavidaviridae (i.e. virophages) – of which the first two groups contain microalgae
infecting viruses and are giant viruses (phylum *Nucleocytoviricota*). Additionally, there
were few VBMs to other known giant viruses (i.e. families Irodoviridae,
Marseilleviridae, and Pitoviridae). Overall, over 80% of all hits were from TsV-N1 or
TetV, with TetV responsible for 58% of the total. The proportion of TsV-N1 and TetV
hits varied across species and also across strains of the same species, with noticeable
differences between both *T. striata* strains (TsV-N1 was 29% and 41%, and TetV was
52% and 22%, for LANL1001 and UTEXSP22 respectively) and between *T. suecica*
CCMP904 assemblies (TsV-N1 was 10% and 17%, and TetV was 71% and 53% for
NYUAD and IFREMER, respectively). Overall, the *T. suecica* CCMP904 NYUAD assembly
had 4 times more VBM than the *T. suecica* CCMP904 IFREMER assembly. These
incongruences may have different reasons, including the heterogeneous quality of
genome assembly, sequencing depth and technology, and different within-species
history of interaction with viruses. Although *Prasinocladus* sp. malaysianus diverged
early from the *Tetraselmis* genus within the *Chlorodendraceae* family (**Figure S1)** it
also had viral hits (albeit substantially less than other algal genomes analysed), which
included both TsV-N1 and TetV matches; two viruses not known to be associated with

genus *Prasinocladus*.

To complete the analyses we assembled public transcriptomic data from 15 cultured
VBM ORFs versus 149 VBM transcripts, respectively. This suggests that only a fraction
of the LANL1001 VBM ORFs were transcribed under the culture conditions applied for
transcriptome sequencing. Furthermore, the taxonomic distribution of VBMs differed
substantially between ORFs and transcripts as shown in **Figures 1A and 1B**,
respectively. *Phycodnaviridae* was the dominant taxonomic category in transcript
VBMs, recruiting from 31% to 58% of VBMs, whereas it only accounted for 6 to 18% of
the ORF VBMs. TetV also recruited a large portion of the transcript VBMs (21-41%) but
less so than for genomic ORFs (31-71%). Finally, TsV-N1 accounted for 0-10% of
transcript VBMs, which is substantially less than the 10-41% observed in genomic
ORFs. Overall, these results suggest that the propensity of a gene to be transcribed
varies with the taxonomic origin of its VBM.

Algal sequences with a VBM (3431 genomic ORFs and 1532 transcripts; **Dataset S1**)
were translated and clustered into 561 protein families (**Table S3**). About two thirds of
the protein families (N=344) were found in two or more strains. The largest family
included 802 proteins containing a RING-type z-finger domain (FAM001). These
proteins were encoded by highly duplicated genes in all *Chlorodendraceae*. Only one
remote homolog was found in green algae (i.e., Trebouxia sp. A1-2 KAA6422835). Their
most closely related homologs were found in Chloroviruses which contained from 1 to
10 gene copies. Other large protein families include Ankyrin domain proteins (FAM002,
N=367) also found in various Nucleocytoviricota, homologues of the TsV (PLV) DNA
packaging ATPases (FAM003; N=236) and homologues of the TsV MCP (FAM004;
N=200).

We queried the *Chlorodendraceae* sequences with reference major capsid proteins
(MCP) from TetV (Nucleocytoviricota) and from TetV (PLV). This approach is more
sensitive than the former VBM analysis because some algal sequences may be
truncated or match viral insertions in cellular genomes, in which case they cannot be
detected as VBM candidate. By definition, MCPs are encoded by hallmark viral genes
and therefore cannot be of cellular origin; their presence necessarily attests to events
of genetic transfer from viruses to algae. The amino acid sequence divergence
between Nucleocytoviricota MCPs and PLV MCPs is generally so high that they cannot
align against each other at the specified threshold of detection (MMSEQS e-value<1e-
5). In addition to the number of matching ORFs in genome assemblies, Table 1 reports
the number of contigs carrying these ORFs, because sequencing errors or nucleotide
substitutions can lead to splitting of a full length gene into several successive ORFs
(hence multiple counts) due to occurrence of internal stop codons. This contig number

can be considered as a lower bound on the estimate of the number of actual MCP
genes in each genome assembly. Thus, between zero and four ORFs had a significant
protein similarity to the TetV MCP and between two and 247 ORFs (on 203 contigs)
had significant protein similarity to the TsV MCP (**Table 1**). In transcriptomes, only
Tetraselmis sp. KCTC12432BP had a transcript with significant similarity to the TetV
MCP, and 8 distinct Chlorodendraceae strains had between 2 to 10 transcripts with
significant similarity to a PLV MCP.

Phylogenetic reconstruction of the TetV-like MCPs indicated that most algal sequences
clustered with the TetV (Tetraselmis giant virus) homologue, further confirming the
close-knit relationship between this viral clade and the *Tetraselmis* genus (Figure 2).
However, the MCP encoded by the *Tetraselmis* sp. KCTC12432BP transcript was more
divergent and occupied an isolated position in the tree without close viral homologues.
This sequence could represent the signature of a still unknown virus infecting this
Tetraselmis strain. A phylogenetic analysis of the PLV-like MCP is shown in the PLV
elements section below .

Large TetV-like genome insertions in Chlorodendraceae algae

We then analysed the distribution of VBM genes in the genome assemblies using a
sliding window approach. Contigs containing 10-Kb windows with at least four viral
ORFs were identified in all assemblies but *Prasinocladus* sp. and were inspected
further using graphical representations of the ORF best hit taxonomic classifications
and local GC content of contigs. This way we could identify three large regions
(roughly 400Kb to 500Kb) of the *T. striata* LANL1001 assembly that had a higher viral
gene density (**Figure 3**). We confirmed that the regions were flanked by ORFs with
algal matches to rule out viral contamination. Using a phylogenetic analysis, the
relationship of the integrated viral elements to TetV was confirmed using the single
copy of DNA polymerase ORF identified in each fragment. The TetV-like MCP gene
identified in LANL1001 (**Table 1**) is not located in one of these genomic regions;
instead, it is isolated among genes of algal origin suggesting that it results from
another integration event. In regard to the host GC content verses the viral insertions,
one located on contig #2972 is more similar (G+C=58.8%) to the GC content of the
host algal genome (G+C=58.0%), whereas the two others are drastically different at
over 66% GC content (G+C=66.0% and 67.1% for contigs #3242 and #784
respectively) (**Figure 3**). Furthermore, the CG content of the viral genome of TetV is
also drastically different (i.e. 41.2%) than that of all three viral insertion elements and
the host algal genome.

Although other investigated algal genomes contained a great number of best matches

to TetV (i.e. *T. striata* UTEXSP22, *T. suecica* CCMP904 NYUAD and *T. subcordiformis*
UTEX171), we did not identify such large viral insertions in their assemblies (see
Figure S2). One obvious reason is that these assemblies were highly truncated as
reflected by the quasi absence of contigs >20 Kb, which was not suitable for this
analysis. As a result, positive windows were only located on short contigs, which did
not contain additional ORFs matching to green algae. Thus, we could not determine if
these sequences originated from viral contaminants or integrated viral DNA. All of the
contigs containing positive windows shown in **Figure S2** had a majority of best
matches to TetV. Their G+C contents ranged between 47.2% and 63.9%, and were
generally higher than the average host G+C% (from 51.6% for *T. suecica* CCMP904
NYUAD to 56.4% for *T. striata* UTEXSP22).

In contrast, *T. suecica* CCMP904 IFREMER project was sequenced using long ONT
reads, and 97% of the genomic sequences was assembled in chromosome-sized
contigs >100 Kb. Although the number of TetV matches for this genome assembly is
comparatively low, we found three contigs of likely viral origin, generating positive
windows containing exclusively best matches to TetV. The largest contig (#6216) was
383 Kb with a GC content (63.6%) higher than for the algal (53.0%) and TetV
genomes. The two remainders (#5310 and #5311) were 11 Kb and 32 Kb (72.5% and
73.9% G+C, respectively). Five additional high GC content contigs (G+C>70%; contigs
#6197,   #6058,   #1543,   #2055   and   #5034)   containing   best   matches   to
Nucleocytoviricota (including TetV) but too short to generate positive windows were
also observed in the graphical representation of VBM genes in the *T. suecica* CCMP904
IFREMER genome assembly (**Figure S3**). All of these high-GC contigs could originate
from the same genomic region. Again, the absence of match to green algae at the
extremity of the contigs made it impossible to determine if these sequences result
from viral contaminants or integrated viral DNA. However, an analysis of the depth of
coverage of IFREMER contigs by Illumina NYUAD reads may provide some indication
(**Figure S4**). While the IFREMER algal contigs have an average mapping coverage of
43X, the viral contigs have an average coverage of 3.3X. It is therefore possible that
the viral contigs are not part of the algal genome but rather originate from a low
intensity viral infection contained in the CCMP904 cultures distributed by the culture
collection centers. Other genes with a best match in *Nucleocytoviricota* but located in
genomic regions with a background GC content appear to be randomly distributed.
The largest TetV-like *T. suecica* CCMP904 IFREMER contig and the 3 *T. striata*
LANL1001 contigs exhibited residual gene colinearity conservation with the TetV
genome and with each other (**Figure S5**).

In all cases, our results suggest that five strains of Tetraselmis, which are distinct from

the Tetraselmis strain on which TetV was initially isolated (19), were concerned by
infection by TetV relatives at some point in the history of their lineages. Collectively,
the algal genomes contained copies for 285 of the 653 protein genes predicted in the
TetV genome, with *T. suecica* CCMP904 IFREMER having the smallest TetV gene
complement (N=51) and *T. striata* LANL1001 the largest (N=253).

PLV-like elements provide insights into the evolution and mechanisms behind PLVs

A total of 66 integrated viral elements above 2000 bp in length and with similarities to
PLVs were also retrieved from the six draft genomes assessed in this study. We also
identified 16 contigs resembling PLVs in the assembled transcriptomic data of 7
Chlorodendraceae strains including *Tetraselmis* spp. GSL018 and M8, *T.*
subcordiformis, *T. chui* UTEXLB232, *S. dubia*, *T. suecica* CS-187 and CCMP906. These
contigs often contained multiple ORFs, sometimes in opposite directions and/or
corresponding to distinct hallmark PLV genes, which probably reflects sequence
overlaps between successive PLV transcripts. Phylogenetic reconstruction based on
the PLV MCP shows a remarkable diversity of the algal PLV-like elements around the
TsV itself (**Figure 4A**). Four coherent phylogenetic groups could be defined (G1 to
G4), one of which contains the TsV (G2). The closest TsV relative was a short
transcribed fragment assembled from the *T. subcordiformis* FACHB-1751 RNA-seq data
(76% amino acid identity between MCPs). Elements from the same algal species, such
as *T. striata* or *T. suecica*, tended to share higher sequence similarity and clustered
tightly in the tree. This suggests that algal species independently underwent burst of
insertions by specific viral lineages rather than steady accumulation of insertions by a
diversity of PLVs. Structural annotation revealed many repeating viral gene hits across
the PLV-like elements, but their relative order varied especially between phylogenetic
groups (**Figure 4B**). Several genes are found split into several consecutive ORFs due
to interior stop codons in the reading phase, following sequencing errors or sequence
decay. Shared genes included ones from a "typical" PLV repertoire and encode
proteins such as ATPase, minor and major capsid proteins, as well as genes noted by
previous studies of PLVs (3) including lipase, cytosine methylase and transcription
factor VLTF3. Lipases are enzymes that normally break down lipids but have been
recorded without this activity in viruses and instead could be involved in replication
(36). Four shared putative genes without a known or implied function (TVSGs and G-
protein (3); i.e. hypothetical proteins) were also recorded and of significance because
they appear in TsV and other PLVs. Interestingly PLV-like elements had 3 different,
mutually exclusive genes encoding proteins involved in viral genome replication,
including S3H helicase – TVpol (transposon-viral polymerase) fusion, protein-primed
DNA polymerase (pDNAP) and S1H helicase. This suggests that these important genes

can be readily interchanged during PLV evolution but they seemingly cannot coexist in
a same viral genome. The most parsimonious scenario based on conciliation of gene
distribution and tree topology suggests that S3H helicase – TVpol was the ancestral
gene form and was replaced by the S1H helicase in group 2 PLV-like elements and
pDNAP in group 4. No significant hit to protein involved in DNA integration (e.g., RVE
integrase or tyrosine recombinase YREC (OLV11)) was found in *Chlorodendraceae*
sequences, either within or outside the identified PLV-like elements. Thus the
mechanism and enzymatic apparatus responsible for the *Chlorodendraceae* PLV
integration remain unknown. Other genes previously found in PLVs of the TVS group
but missing the *Chlorodendraceae* elements are the ones encoding Tlr6F and
endosialidase. Tlr6F is an uncharacterized protein found in several viruses (e.g. PLVs,
Lavidaviridae, *Nucleocytoviricota*, and bacteriophages) and thought to be a
component of viral reproduction (3).
Some of *Chlorodendraceae* PLV element genes have not been documented in PLVs
yet. For example zinc finger (displayed as Zn-finger) proteins encoded by some group
G1 PLV elements are known to have many different functions in relation to interacting
with DNA or RNA, or other proteins, but in some cases have been shown to inhibit
replication in viruses such as retroviruses, alphaviruses, and filoviruses . These are
RNA virus taxa, however, designed zinc finger proteins have been used to inhibit a
clinical DNA virus (herpes simplex virus 1 (37)). Zinc finger domains have also been
shown to be associated with an integrase domain in eukaryotic transposons (38). It is
possible that these zinc finger genes could have some effect on replication or have
been involved in the PLV integration process. PLV elements of group G2 carry a gene
encoding an adenine-methyltransferase of prokaryotic origin. This gene has never
been identified in a PLV but a distant homologue is found in Yellowstone lake
phycodnavirus 2 and in some bacteriophages. In the latter, adenine methyltransferase
activity renders the DNA resistant to cleavage by restriction enzymes sensitive to
adenine methylation. Moreover, these same PLV elements carry in the vicinity of the
adenine-methyltransferase gene another gene coding for a restriction endonuclease
also of prokaryotic origin. It seems that these PLVs have a complete restriction-
modification system that probably allows them to recognize and degrade the DNA of
unmethylated competitor viruses in case of coinfection. Finally, several PLV elements
of the G2 and G4 groups carry a retrotransposon. It is possible that these transposable
elements were inserted into the PLV elements after the latter were integrated into the
genome. However, the position of the retrotransposon is conserved in PLV elements of
the same group, suggesting that the insertion of the retrotransposon occurred in the
ancestor of these PLVs, before the latter integrated into the host genome. If this
hypothesis is true, the PLVs could then participate in the propagation of transposable

elements from one host to another.

Some PLV-like elements exhibited terminal inverted repeats (TIRs) at both ends of the
sequence (**Figure 4B).** Typically TIRs indicate that the genome of the inserted PLV is
complete (3, 39, 40). There is a total of 13 PLVs retrieved from these datasets that
have TIRs flanking the genes (i.e. on both ends), suggesting that these 13 PLV
insertions are likely complete. Their sizes ranged from 16,954 bp (Tstr-LANL1001-
C0660) to 31,706 bp (Tstr-LANL1001-C2288), with a mean at 19,903bp, which is
smaller than the size of TsV (26,407 bp) but comparable to other PLVs (MED1:
21,217bp, RED1: 19,641 bp, SPO1: 22,704 bp (3)). Another interesting feature is the
apparent uneven distribution of PLV-like elements along chromosomes of *T striata*
CCMP904 IFREMER (**Figure S3**). 56% of the ORFs encoding a PLV MCP, including
group 1 and 2 MCPs (**Figure 4**), were clustered in the central region (between
positions 8,6 Mb and 9.4 Mb) of the largest contig (Tsue-IFRE-C001; 17,2 Mb), which
most likely correspond to the largest chromosome. The remaining PLV MCP genes
were randomly dispersed throughout the rest of the genome. This central region may
represent a hotspot for PLV integration. Unfortunately the highly fragmented
assemblies of the other algal species made it impossible to determine whether such
hotspots also exist in other *Chlorodendraceae*.

VBM gene transcription were low among TsV and TetV categories within
Chlorodendraceae individuals

We sought to study the transcription activity of VBM genes using the available
Chlorodendraceae RNA-seq data. Only *T. striata* LANL1001 had both genomic and
RNA-seq data available to perform a conventional transcriptomic analysis (i.e.,
genome and transcriptome from the same strain). To widen the analysis, we also
aligned RNA-seq data and genome assemblies from the same phylogenetic group (i.e.,
the *T. striata* and *T. suecica* groups as defined in the RBCL phylogenetic tree as shown
in **Figure S1**), assuming that VBM genes are likely to be conserved between closely
related strains. Alignment of the *T. striata* LANL1001 transcriptomic data on the *T.
striata* LANL1001 genome assembly revealed that most VBM genes associated with
the taxonomic categories TetV and TsV were transcriptionally silent (FPKM~0; see
Figure 5A). This was also the case for VBM genes associated with the category "Other
viruses", however these genes were most often located within PLV elements or in
TetV-like genomic regions, suggesting that they have the same evolutionary origin as
the TsV or TetV categories. Only 16, 26 and 2 genes from categories "TsV", "TetV" and
"Other" had FPKM≥1 and were considered as transcribed (accounting for 2.7%, 7.9%
and 3.9% of their categories, respectively). Interestingly 13 of the 16 transcribed TsV-
like genes were most similar to the unknown protein gene TVSG_00024 (found

extensively in G1 and G4; **Figure 4B**), suggesting that they may act as sentinel genes
that orchestrate a possible reactivation of virus following an unknown stimulus or
event. Transcribed genes from the "TetV" and "Other" categories had more diverse
functions. RNA-seq data of four other Tetraselmis strains did not align better with the
LANL1001 VBM genes in categories "TsV", "TetV" and "Other". In contrast, a large
fraction of VBM genes in categories Mimiviridae, Pandoraviruses, Phycodnaviridae and
Pithoviridae were transcribed in all five Tetraselmis strains. The same analysis using
the *T. striata* UTEXSP22 genome as reference yielded similar overall results (**Figure
5B**). A different set of RNA-seq data was aligned against the genome assemblies of
the suecica group. When the *T. suecica* CCMP904 IFREMER genome was used as
reference, the measured levels of transcription were generally null or very low for
genes in the TsV, TetV and "Other" categories in all tested transcriptomes whereas
they were more substantial for genes in the Mimiviridae, Pandoraviruses,
Phycodnaviridae and Pithoviridae categories (**Figure 5C**). This overall pattern was
similar to what was observed in the striata group. In contrast, when the genomes of *T.
suecica* CCMP904 NYUAD and *T. subcordiformis* UTEX171 were used as reference, only
the *Phycodnaviridae* VBM gene category exhibited substantial gene transcription
(**Figures 5D and 5F**). The difference observed between the IFREMER distribution and
that of the other 2 strains of the suecica clade, especially for the categories
Mimiviridae, Pandoravirus and Pithoviridae, is mainly explained by the fact that the
IFREMER strain contained comparatively small numbers of genes in these categories
(**Table S4**). These genes were also mostly transcribed. They also exist in the other
two strains of the suecica group, and are also transcribed, but are outnumbered in
their category by a large number of additional genes that are not transcribed.

Another question is why the measured levels of transcription are globally different
between VBM categories. Phycodnaviridae best matching genes had systematically
relatively high FPKM values (plus to some extent Mimiviridae, Pandoraviruses, and
Pithoviridae) whereas TsV and TetV best matching genes were generally silent. Since
most of TsV and TetV-like genes resided in genomic regions of viral origin, it appears
that viral genes tend to be untranscribed. The origin of the other VBM gene categories
is unknown (i.e., virus to alga HGT or the reverse) but the frequency of VBM genes
that also show significant secondary BLASTP matches in *Viridiplantae* (i.e., includes
green algae (*Chlorophyta*) and land plants (*Streptophyta*)) may give some indication in
this regard. Indeed, secondary matches are consistent with the scenario that the gene
is of algal origin but was recently captured by a virus, ultimately resulting in a best
match in a virus derived from the lineage involved in the HGT. **Figure 6** shows that
for each algal genome, the percentage of genes that have significant matches in

Viridiplantae is higher for Phycodnaviridae, Mimiviridae, Pandoravirus, and Pithoviridae
than for the other categories. This result therefore suggests that genes in these
categories are more likely to have an algal rather than a viral origin. Taken together,
these results lead to a consistent hypothesis that the categories Phycodnaviridae,
Mimiviridae, Pandoravirus and Pithoviridae are characterized by higher transcriptional
activity because they mainly consist of genes of algal origin that have been captured
by viruses and whose expression is naturally required to participate in biological
functions of the host cell.

Discussion

Transcriptionally silent giant virus-like insertion elements (phylum Nucleocytoviricota)
in this study may be actually transcribed under specific conditions

Viral genomic regions in *T. striata* LANL1001 and *T. suecica* CCMP904 IFREMER that
best match the algal giant virus TetV had GC content strikingly different than both the
cellular host's genome and the TetV genome. Positive windows containing TetV best
matches in the other *Chlorodendraceae* genomes also had G+C content substantially
higher than that of the TetV genome. Excluding the hypothesis that these GC content
variations were the result of specific mutational bias within inserted viral regions
induced by host DNA replication or DNA repair, it is highly likely that they reflect the
diversity of GC contents that exists among TetV-related viruses.
Only a fraction of the TetV gene complement were found in the algal genomes. For
instance neither of the two key fermentation genes (i.e., pyruvate formate-lyase and
pyruvate formate-lyase activating enzyme), which have been a surprising discovery in
the TetV genome (19), were found in the viral insertions or elsewhere in the algal
genomes. Although we cannot rule out the viral genes were missed due to gaps in
sequencing or because the viral genome insertions were incomplete, this result
suggests that these accessory genes might have a sporadic distribution in the TetV-
like clade, reflecting different adaptive niches.

In general, TetV-like genes found in algal genomes were not represented in
Chlorodendraceae transcriptomes, suggesting that these genes do not produce mRNA,
at least under the conditions under which the transcriptomes were generated. With
the exception of LANL1001, it was not possible to align transcriptomes and genomes
belonging to the same algal strain. Although overall, the results of the transcriptomic
analyses were similar to those obtained for LANL1001, we cannot exclude that the lack
of transcription measured in heterologous comparisons was the consequence of a
physical absence of the corresponding genes in the algal strains used for
transcriptome sequencing. This is especially true for *T. suecica* CCMP904 where a

background low intensity infection of the stock culture is suspected. A complementary
RNA-seq study of the CCMP904 culture would certainly be able to determine if this
hypothesis is valid (viral genes are actively transcribed upon infection).

Giant virus genomic insertions are relatively common in algae and some protozoa but
their biological role in the host cell remains unknown (13, 23, 41–43). This role could
be nil if the viral DNA is permanently inactivated because the cellular transcription
machinery is unable to recognize viral promoters or as the result of active gene
silencing by the host. The absence of gene transcription in viral regions supports these
hypotheses. Alternatively, viral-derived genes could be expressed under very specific
conditions (e.g not unlike the response of genome integrated virophages to giant virus
infection (8)), namely upon infection with a related virus that brings its transcription
factors capable of recognizing viral promoters. Since viral host-borne genes have
accumulated numerous mutations, their expression could lead to the production of
abnormal viral proteins that would compete with the native viral proteins and hinder
infection and/or the production of new infectious particles. Another possibility is that
the inserted viral regions are transcribed into small interference RNAs (siRNAs), which
are not covered by the transcriptomic sequencing data analyzed in this work. The
siRNA pathway is an antiviral mecanism widespread across the eukaryotic phylogeny,
which can potentially direct viral DNA methylation and transcriptional silencing (44).
Obviously, further studies are needed to investigate and understand the biological
significance of these giant-virus insertions.

Polinton-like viruses likely exhibit dual lifestyles that also facilitate rapid hybridization

Chlorodendraceae genomes also showed an abundance of diverse PLV-like elements
whose closest known relative was TsV infecting *T. striata*. TsV is the only PLV
specimen that is known and isolated as a viral infectious particle. All other PLVs
reported in the literature have been identified either as inserted elements in
eukaryotic genomes or in viral metagenomes. For this reason, PLVs are suspected to
have a dual lifestyle, alternating between a free viral phase and a phase integrated
into host genomes. The PLV-like elements integrated into Chlorodendraceae genomes
provide an additional argument to this hypothesis. It is highly likely that TsV is capable
of inserting itself into the host genome under conditions that remain to be determined.
Transcriptomic data, particularly in the case of LANL1001, suggest that PLV-like
elements are essentially silent in their integrated form except for a gene homologous
to TVSG_00024 of TsV that is transcribed in several elements. The function of this
gene is unknown but given its quasi-systematic transcription in the inserted elements
it is possible that the product of this gene is involved in a sensing mechanism that
allows to trigger the reactivation/excision of the provirus according to a stimulus that

remains to be discovered. In some transcriptomes, assembled sequences contained
multiple ORFs encoding distinct PLV proteins. It is likely that these sequences
correspond to distinct but overlapping transcription units (i.e., adjacent genes) that
were assembled in the same contig. We do not know whether these transcripts were
generated by integrated elements in the genomes of the concerned strains. However,
given the very restricted transcription of the PLV elements of LANL1001, it is more
likely that these highly transcribed genes result from an infection of the culture that
served as a template for transcriptomic sequencing.

In the IFREMER strain, the distribution of elements in the genome was not random and
was concentrated in a central region of the largest chromosome. It is possible that this
region also houses the centromere, but this information is not known. In any case, this
distribution suggests that insertions of PLVs occur in genomic hotspots. Further
sequencing and near-complete assembly of the other algal strains will confirm
whether these hotspots exist in other Chlorodendraceae. It is also striking that the
Chlorodendraceae PLV elements had 3 different types of replication proteins (i.e., S3H
helicase – TVpol fusion, pDNAP and S1H helicase) and that these were mutually
exclusive, so that each element had only one gene. Given the relative phylogenetic
proximity of PLV-elements to each other, this observation suggests that PLVs change
their replication genes quite easily and frequently in evolution. The existence of
element-dense integration hotspots, where new PLVs are likely to integrate into
already established elements, is likely to favor the creation of hybrid elements that
basically result in gene exchanges between PLVs, especially their replication gene.
These new genetic combinations can then be tested by natural selection if the hybrid
genome can be excised and propagated in a neoformed virion. In this context, it is
interesting to note the existence of hybrids between virophages and polintons, as was
found among rumen virophages where a virophage MCP was combined with a polinton
DNA polymerase B (45)

Conclusion

This study explored both genomic and transcriptomic data of previously sequenced
algae (family *Chlorodendraceae*) with close relation to a polinton-like virus (TsV) host
T. striata to further support the dual lifestyle hypothesis among PLVs. Additionally,
integration elements of giant viruses were also explored, given a known *Tetraselmis*
sp. virus (TetV). Both of these analyses uncovered integrated viral elements among
several of the related algal species and gave insights into their behaviors. Given the
substantial evidence of both giant virus and PLV viral integrated elements, we
recommend future studies target the mechanism(s) behind triggering the possible
transcription of these integration elements. More specifically we recommend further

characterization of the unknown protein, TVSG_00024 (found within many PLV integrated viral elements) given its possible role in triggering transcription of PLV viral integrated elements. This work could further reveal the mechanisms behind the implied dual lifestyle of PLVs.

Acknowledgment:

we would like to thank Rossana Sussarellu, Gregory Carrier and Sabine Stachowski-Haberkorn from IFREMER Nantes for providing us access to their CCMP904 data in advance of publication.

Funding:

E.E.C, G.B, and C.D received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No713750, with the financial support of the Regional Council of Provence-Alpes-Côte d'Azur and with the financial support of the A*MIDEX (n° ANR- 11-IDEX-0001-02), funded by the Investissements d'Avenir project funded by the French Government, managed by the French National Research Agency (ANR). The Phycovir project leading to this publication has received funding from Excellence Initiative of Aix-Marseille University - A*MIDEX, a French "Investissements d'Avenir" program.

602   REFERENCES

1.   Pritham EJ, Putliwala T, Feschotte C. 2007. Mavericks, a novel class of giant transposable elements widespread in eukaryotes and related to DNA viruses. Gene 390:3–17.

2.   Krupovic M, Koonin EV. 2016. Self-synthesizing transposons: unexpected key players in the evolution of viruses and defense systems. Current Opinion in Microbiology 31:25–33.

3.   Yutin N, Shevchenko S, Kapitonov V, Krupovic M, Koonin EV. 2015. A novel group of diverse Polinton-like viruses discovered by metagenome analysis. BMC Biol 13:95.

4.   Krupovic M, Koonin EV. 2015. Polintons: a hotbed of eukaryotic virus, transposon and plasmid evolution. Nat Rev Micro 13:105–115.

5.   La Scola B, Desnues C, Pagnier I, Robert C, Barrassi L, Fournous G, Merchat M, Suzan-Monti M, Forterre P, Koonin E, Raoult D. 2008. The virophage as a unique parasite of the giant mimivirus. Nature 455:100–104.

6.   Xu S, Zhou L, Liang X, Zhou Y, Chen H, Yan S, Wang Y. 2020. Novel Cell-Virus-Virophage Tripartite Infection Systems Discovered in the Freshwater Lake Dishui Lake in Shanghai, China. J Virol 94.

7.   Duponchel S, Fischer MG. 2019. Viva lavidaviruses! Five features of virophages that parasitize giant DNA viruses. PLoS Pathog 15:e1007592.

8.   Fischer MG, Hackl T. 2016. Host genome integration and giant virus-induced reactivation of the virophage mavirus. Nature 540:288–291.

9.   Mougari S, Chelkha N, Sahmi-Bounsiar D, Di Pinto F, Colson P, Abrahao J, La Scola B. 2020. A virophage cross-species infection through mutant selection represses giant virus propagation, promoting host cell survival. Commun Biol 3:248.

10.   Krupovic M, Kuhn JH, Fischer MG. 2016. A classification system for virophages and satellite viruses. Arch Virol 161:233–247.

11.   Yutin N, Raoult D, Koonin EV. 2013. Virophages, polintons, and transpovirons: a complex

evolutionary network of diverse selfish genetic elements with different reproduction strategies. Virol J 10:158.

12. Hackl T, Duponchel S, Barenhoff K, Weinmann A, Fischer MG. 2021. Virophages and retrotransposons colonize the genomes of a heterotrophic flagellate. eLife 10:e72674.

13. Blanc G, Gallot-Lavallée L, Maumus F. 2015. Provirophages in the Bigelowiella genome bear testimony to past encounters with giant viruses. PNAS 201506469.

14. Desnues C, Scola BL, Yutin N, Fournous G, Robert C, Azza S, Jardot P, Monteil S, Campocasso A, Koonin EV, Raoult D. 2012. Provirophages and transpovirons as the diverse mobilome of giant viruses. PNAS 109:18078–18083.

15. Santini S, Jeudy S, Bartoli J, Poirot O, Lescot M, Abergel C, Barbe V, Wommack KE, Noordeloos AAM, Brussaard CPD, Claverie J-M. 2013. Genome of Phaeocystis globosa virus PgV-16T highlights the common ancestry of the largest known DNA viruses infecting eukaryotes. PNAS 110:10800–10805.

16. Stough JMA, Yutin N, Chaban YV, Moniruzzaman M, Gann ER, Pound HL, Steffen MM, Black JN, Koonin EV, Wilhelm SW, Short SM. 2019. Genome and Environmental Activity of a Chrysochromulina parva Virus and Its Virophages. Front Microbiol 10:703.

17. Bellas CM, Sommaruga R. 2021. Polinton-like viruses are abundant in aquatic ecosystems. Microbiome 9:13.

18. Pagarete A, Grébert T, Stepanova O, Sandaa R-A, Bratbak G. 2015. Tsv-N1: A Novel DNA Algal Virus that Infects Tetraselmis striata. Viruses 7:3937–3953.

19. Schvarcz CR, Steward GF. 2018. A giant virus infecting green algae encodes key fermentation genes. Virology 518:423–433.

20. Stepanova OA, Boǐko AL, Shcherbatenko IS. 2013. Computational genome analysis of three marine algoviruses. Mikrobiol Z 75:76–81.

21. Kazlauskas D, Krupovic M, Venclovas Č. 2016. The logic of DNA replication in double-

stranded DNA viruses: insights from global analysis of viral genomes. Nucleic Acids Res 44:4551–4564.

22. Steadman Tyler CR, Hovde BT, Daligault HE, Zhang XL, Kunde Y, Marrone BL, Twary SN, Starkenburg SR. 2019. High-Quality Draft Genome Sequence of the Green Alga *Tetraselmis striata* (Chlorophyta) Generated from PacBio Sequencing. Microbiol Resour Announc 8.

23. Nelson DR, Hazzouri KM, Lauersen KJ, Jaiswal A, Chaiboonchoe A, Mystikou A, Fu W, Daakour S, Dohai B, Alzahmi A, Nobles D, Hurd M, Sexton J, Preston MJ, Blanchette J, Lomas MW, Amiri KMA, Salehi-Ashtiani K. 2021. Large-scale genome sequencing reveals the driving forces of viruses in microalgal evolution. Cell Host & Microbe 29:250-266.e8.

24. Keeling PJ, Burki F, Wilcox HM, Allam B, Allen EE, Amaral-Zettler LA, Armbrust EV, Archibald JM, Bharti AK, Bell CJ, Beszteri B, Bidle KD, Cameron CT, Campbell L, Caron DA, Cattolico RA, Collier JL, Coyne K, Davy SK, Deschamps P, Dyhrman ST, Edvardsen B, Gates RD, Gobler CJ, Greenwood SJ, Guida SM, Jacobi JL, Jakobsen KS, James ER, Jenkins B, John U, Johnson MD, Juhl AR, Kamp A, Katz LA, Kiene R, Kudryavtsev A, Leander BS, Lin S, Lovejoy C, Lynn D, Marchetti A, McManus G, Nedelcu AM, Menden-Deuer S, Miceli C, Mock T, Montresor M, Moran MA, Murray S, Nadathur G, Nagai S, Ngam PB, Palenik B, Pawlowski J, Petroni G, Piganeau G, Posewitz MC, Rengefors K, Romano G, Rumpho ME, Rynearson T, Schilling KB, Schroeder DC, Simpson AGB, Slamovits CH, Smith DR, Smith GJ, Smith SR, Sosik HM, Stief P, Theriot E, Twary SN, Umale PE, Vaulot D, Wawrik B, Wheeler GL, Wilson WH, Xu Y, Zingone A, Worden AZ. 2014. The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): illuminating the functional diversity of eukaryotic life in the oceans through transcriptome sequencing. PLoS Biol 12:e1001889.

25. One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant transcriptomes and the phylogenomics of green plants. Nature 574:679–685.

26. Yao C, Ai J, Cao X, Xue S, Zhang W. 2012. Enhancing starch production of a marine green microalga Tetraselmis subcordiformis through nutrient limitation. Bioresource Technology

118:438–444.

27. Lauritano C, De Luca D, Amoroso M, Benfatto S, Maestri S, Racioppi C, Esposito F, Ianora A. 2019. New molecular insights on the response of the green alga Tetraselmis suecica to nitrogen starvation. Sci Rep 9:3336.

28. Lim DKY, Schuhmann H, Thomas-Hall SR, Chan KCK, Wass TJ, Aguilera F, Adarme-Vega TC, Dal'Molin CGO, Thorpe GJ, Batley J, Edwards D, Schenk PM. 2017. RNA-Seq and metabolic flux analysis of Tetraselmis sp. M8 during nitrogen starvation reveals a two-stage lipid accumulation mechanism. Bioresource Technology 244:1281–1293.

29. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nature Biotechnology 37:907–915.

30. Criscuolo A, Brisse S. 2013. AlienTrimmer: a tool to quickly and accurately trim off multiple short contaminant sequences from high-throughput sequencing reads. Genomics 102:500–506.

31. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. Journal of Computational Biology 19:455–477.

32. Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics 22:1658–1659.

33. Steinegger M, Söding J. 2017. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. Nat Biotechnol 35:1026–1028.

34. Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. Molecular Biology and Evolution 30:772–780.

35. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – Approximately Maximum-Likelihood Trees for Large Alignments. PLOS ONE 5:e9490.

36. Kamil JP, Tischer BK, Trapp S, Nair VK, Osterrieder N, Kung H-J. 2005. vLIP, a viral lipase

homologue, is a virulence factor of Marek's disease virus. J Virol 79:6984–6996.

37. Papworth M, Moore M, Isalan M, Minczuk M, Choo Y, Klug A. 2003. Inhibition of herpes simplex virus 1 gene expression by designer zinc-finger transcription factors. Proc Natl Acad Sci U S A 100:1621–1626.

38. Llorens C, Bernet GP, Ramasamy S, Feschotte C, Moya A. 2012. On the transposon origins of mammalian SCAND3 and KRBA2, two zinc-finger genes carrying an integrase/transposase domain. Mob Genet Elements 2:205–210.

39. Haapa-Paananen S, Wahlberg N, Savilahti H. 2014. Phylogenetic analysis of Maverick/Polinton giant transposons across organisms. Mol Phylogenet Evol 78:271–274.

40. Koonin EV, Krupovic M. 2017. Polintons, virophages and transpovirons: a tangled web linking viruses, transposons and immunity. Curr Opin Virol 25:7–15.

41. Moniruzzaman M, Weinheimer AR, Martinez-Gutierrez CA, Aylward FO. 2020. Widespread endogenization of giant viruses shapes genomes of green algae. Nature 588:141–145.

42. Gallot-Lavallée L, Blanc G. 2017. A Glimpse of Nucleo-Cytoplasmic Large DNA Virus Biodiversity through the Eukaryotic Genomics Window. Viruses 9:17.

43. Maumus F, Blanc G. 2016. Study of Gene Trafficking between Acanthamoeba and Giant Viruses Suggests an Undiscovered Family of Amoeba-Infecting Viruses. Genome Biol Evol 8:3351–3363.

44. Erdmann RM, Picard CL. 2020. RNA-directed DNA Methylation. PLoS Genet 16:e1009034.

45. Yutin N, Kapitonov VV, Koonin EV. 2015. A new family of hybrid virophages from an animal gut metagenome. Biol Direct 10:1–9.

603 **Table 1.** Total number of best matches to the virus hallmark MCP

	TetV MCP[*]	TsV MCP[*]
Genome assemblies:		
T. striata LANL1001	1 (1)	126 (98)
T. striata UTEXSP22	1 (1)	247 (203)
T. suecica CCMP904 NYUAD	4 (4)	105 (100)
T. suecica CCMP904 IFREMER	0	39 (9)
T. subcordiformis UTEX171	3 (3)	115 (103)
*Prasinocladus sp.*malaysianus UTEX2601	0	2 (2)
Transcriptome assemblies[**]:		
Scherffelia dubia CCAC0019	0	3
T. chuii PLY429	0	2
T. chuii UTEXLB232	0	4
T. subcordiformis FACHB-1751	0	2
T. suecica CCMP906	0	10
T. suecica CS-187	0	9
Tetraselmis sp. GSL018	0	7
Tetraselmis sp. KCTC12432BP	1	0
Tetraselmis sp. M8	0	4

604 * number of ORFs (number of contigs); ** transcriptome datasets are listed only if they contained a best match to one of the two target MCPs.

Figure 1. Taxonomic classification of TrEMBL viral best matches from *Chlorodendraceae* derived sequences. **(A)** Number of corresponding best matches in ORFs generated from draft genomes of several *Tetraselmis* spp., and a member of genus *Prasinocladus*. **(B)** Viral best matches in *Chlorodendraceae* transcripts. Colours correspond to the best match's family, except in the case of TsV-N1 (a polinton-like virus), and TetV (member of family *Mimiviridae*).

Figure 2. Unrooted phylogenetic tree of major capsid proteins (MCP). Proteins identified in the *Chlorodendraceae* data are indicated in color. Grey circles show branches with >85% local support as calculated with the Shimodaira-Hasegawa test. The branch scale indicates the number of amino-acid substitution per site.

Figure 3. Comparison of three viral insertion elements within a *Tetraselmis striata*
draft genome (LANL1001), with the corresponding viral match; TetV-1. ORFs are
represented in the "protein match" track, where red represent best match in viruses,
green are for best match in green algae (i.e., Chlorophyta), blue are for best match in
eukaryotes and pink represent best match in prokaryotes. GC content is also reported
for each viral elements/TetV, alongside the GC content of the host genome (*T. striata*)
overall. On the left, a phylogenetic tree is shown based on viral DNA polymerase
identified as a single copy ORF in each contig and virus. The phylogenetic tree is
rooted by the Dishui Lake phycodnavirus homologue. The scale bar represents the
number of amino acid substitution per site.

Figure 4. PLV-like elements in algal genomes. **(A)** Phylogenetic tree of the MCPs encoded by PLV-like elements, rooted by several PLVs previously identified as members of the TVS group by Yutin et al. (3). **(B)** Genomic architecture of the corresponding PLV-like elements. Names of elements identified in genomic data are

shown in color whereas names of elements identified in transcriptomic data are in
black and preceded by the term "RNA". We only considered PLV elements >2000 bp
except RNA *T. subcordiformis* C21952 (565 bp) which was conserved owing to its close
phylogenetic proximity with TsV. Structural annotations of each viral elements are
depicted with a colour/pattern legend. Spans of terminal inverted repeats (TIRs) are
identified on relevant viral elements by red lines. Scale of the phylogenetic tree
(amino acid substitutions/site) and scale of the genome lengths are included.
Abbreviations are RED: Red Sea, MED: Mediterranean, INO: Indian Ocean and SPO:
South Pacific Ocean, according in reference to their sampling locations; IFRE, IFREMER.
Tstr: *T. striata*; Tsue: *T. suecica*; Tsub: *T. subcordiformis*; Prasino: *Prasinocladus* sp.
malaysianus. Transcribed PLV-like sequences are shown in italic.

Figure 5. Transcription activity of VBM genes. Five algal genomes **(A—E)** were used as reference to map RNA-seq data from various *Tetraselmis* spp. stains and calculate FPKM values for genes. The graphs show the boxplot distributions of FPKM values for genes sorted by VBM taxonomic categories and transcriptomes. Two different sets of transcriptomes were analysed for the *T. striata* clade (**A—B)** and *T. suecica* clade (**C—E)**, as shown in the legends.

646　**Figure 6.** Percentage of genes with secondary matches in Viridiplantae.