Title

Multiple trade-offs between defense and competitiveness traits in a planktonic

predator-prey system

Journal

Ecology

Authors

Tom Réveillon and Lutz Becks

Affiliations

T. Réveillon (https://orcid.org/0000-0003-1788-8635) and L. Becks (https://orcid.org/0000-
0002-3885-5253), Aquatic Ecology and Evolution Group, Limnological Institute, Department
of Biology, University of Konstanz, Konstanz, Germany.

Corresponding authors

Tom Réveillon (tom.reveillon@uni-konstanz.de)

Data availability

Experimental data and scripts will be uploaded on the open platform for research GitHub

(https://github.com/TomReveillon/Multiple-trade-offs-between-defense-and-competitiveness-

traits-in-a-planktonic-predator-prey-system) upon publication.

Abstract

Predator-prey interactions play a central role in community dynamics and depends on the

covariation of traits of the interacting organisms. Intraspecific trade-off relationships between

defense and competitiveness traits are important for understanding trophic interactions.

However, quantifying the relevant traits forming defense-competitiveness trade-offs and how

these traits determine prey and predator fitness remain major challenges. Here, we conducted

feeding and growth experiments to assess multiple traits related to defense and competitiveness

in 6 different strains of the green alga *Chlamydomonas reinhardtii* exposed to predation by the

rotifer *Brachionus calyciflorus*. We found large differences in defense and competitiveness

traits among prey strains and negative relationships between defense and competitiveness traits.

Because we compared trait differences among strains whose ancestors previously evolved in

controlled environments where selection favored defense or competitiveness, these negative

correlations suggest the presence of a trade-off between defense and competitiveness. This

trade-off was found for multiple combinations of defense and competitiveness traits.

Furthermore, the differences in traits translated into differences in prey and predator fitness,

which demonstrated the contribution of intraspecific trade-offs for predicting the outcome of

predator-prey interactions.

Keywords: fitness, defense, competitiveness, predator-prey interactions, algae, rotifers,

Chlamydomonas reinhardtii, *Brachionus calyciflorus*

Introduction

Predator-prey interactions and their consequences on population dynamics are primarily determined by combinations of traits driving the performance of organisms (Colina et al. 2016). Understanding the mechanisms linking traits of predators and prey to the strength of trophic interactions is therefore crucial for assessing the functioning of food webs (Rossberg et al. 2010; Gravel et al. 2016; Laigle et al. 2018; Litchman et al. 2021). Traits involved in defense against predation (i.e., *sensu* protection) and competitiveness (i.e., *sensu* reproduction) are intimately related to the fitness of organisms and mediate trophic interactions (Colina et al. 2016). Prey can exhibit a large range of defense mechanisms against predators, which interfere with different steps of the predation sequence (Weiss et al. 2012; Bateman et al. 2014). These traits affect predator the predator by reducing prey detection and consumption (Rall et al. 2012) via altered attack rate (i.e., the rate of prey encounter), handling time (i.e., the time spent on prey manipulation and ingestion) or gape-size limitation and thus have consequences on the fitness of predators (Long and Hay 2006).

Organisms cannot optimize all functions contributing to fitness simultaneously due to genetic and/or energetic constraints and trade-off relationships between traits can emerge (Stearns 1989). Trade-offs between defense and competitiveness traits are expected (Yoshida et al. 2003; 2004; Becks et al. 2010; Sunda and Hardison 2010; Kasada et al. 2014; Ehrlich et al. 2017; Pančić and Kiørboe 2018; Cadier et al. 2019) and may explain the diversity of fitness strategies (Agrawal 1998; Strauss et al. 2002). Competitiveness costs can arise from the energy investment in expressing anti-grazing defenses due to a reduction in resource acquisition and utilization (Halsey and Jones 2015), and thus in reproduction (Smith et al. 2014). Therefore, trade-off relationships between defense and competitiveness traits can affect trophic interactions by influencing both prey and predator population growth (Wood et al. 2020).

However, such trade-offs are complex to assess because multiple traits can contribute to these

relationships. Evaluating suitable traits describing defense-competitiveness trade-offs using a

multiple traits approach can give important insights on the presence and the role of these trade-

offs for population dynamics and energy transfers in food webs.

Plankton communities are characterized by strong trophic interactions and trade-offs

between defense and competitiveness traits play an important role in regulating these

interactions (Litchman et al. 2007; 2015; Ehrlich et al. 2020). However, there are few empirical

data on defense-competitiveness trade-offs and the existing data should be interpreted with

caution. First, many studies assume fixed traits and trade-offs at the genus or species level

(Bruggeman 2011). However, this assumption has been shown to be often invalid (Züst and

Agrawal 2017) as intraspecific traits and trade-offs can be ecologically and evolutionary

important (Yoshida et al. 2004; Becks et al. 2010). Second, defensive and competitive abilities

of phytoplankton can be assessed by multiple traits (Fleischer et al. 2018) and the relevant traits

constituting a trade-off are often unknown. Previous studies used different traits as proxies for

defense and competitiveness and thus different trait correlations. This makes comparisons of

trade-offs across studies difficult and raises the question of whether different traits can be used

for studying the role of trade-offs for planktonic communities. Third, correlations between traits

might not reflect the absence or the presence of a causal trade-off relationship due to

confounding factors when traits are independently affected by the same environment

(Bohannan et al. 2002; Fry 2003). False-positive (i.e., negative correlation) or false-negative

(i.e., no or positive correlation) trade-offs could for example arise from variation in a third

correlated trait which was unaccounted (Edwards et al. 2011; 2013). Finally, it remains unclear

whether a negative correlation represents a trade-off or simply a signature of differential

selection without knowing whether evolution could optimize both traits simultaneously (Fry

2003; Fuller et al. 2005). Perhaps one of the most thorough demonstrations of trade-offs is when

a trait under selection is optimized at the expense of a trait not under selection and vice versa

(Kassen 2002).

Here, we investigate the intraspecific trade-off relationship between anti-grazing

defense and competitiveness traits for 6 different prey genotypes (hereafter: strains) of the green

alga *Chlamydomonas reinhardtii* exposed to the rotifer predator *Brachionus calyciflorus*. Algal

strains were isolated from a previous experimental evolution study where they were grown

under selection for either defense or competitiveness (Bernardes et al. 2021). They differed in

their morphologies (i.e., single or colonial cells) as well as defense (i.e., measured as reduction

in predator growth rate) and growth (i.e., measured as the prey growth rate). We quantified prey

defense against a predator by measuring functional responses (i.e., the ingestion rate as a

function of prey density) on prey strains to derive the attack rates and the handling times using

mechanistic models, and also by measuring a range of size and shape characteristics and the

carbon to nitrogen ratio of prey strains to link prey morphology and stoichiometry to the trophic

interaction with the predator. We further quantified prey competitiveness by measuring prey

growth in limiting nitrate conditions to derive the maximum growth rate, the nitrate affinity and

the nitrate half-saturation constants using mechanistic models. We tested for the trade-off

relationship between defense and competitive traits, whether it was present for all trait

combinations, and how the trait and trade-off differences affected population growth (i.e.,

fitness) of the algal prey strains and the rotifer predator.

Material and methods

We used 6 strains of the green alga prey *Chlamydomonas reinhardtii* (Dang 1888) with known recent evolutionary history which were derived from isolates obtained from the Chlamydomonas Resource Center (University of Minnesota, USA). Isolates originated from an experimental evolutionary study in which different monoclonal naive ancestors were selected either for defense (exposed to the rotifer *Brachionus calyciflorus*) or competitiveness (not exposed) for 6 months (~500 generations) in semi-continuous cultures (Tab. S1; Bernardes et al. 2021). For the present study, a single colony from each evolved strain was sampled from solid media and transferred to tissue culture flasks filled with 50 mL of sterile growth medium with 800 µmol NO_3^- L^{-1}. Flasks were exposed to continuous light (200 µmol photon m^{-2} sec^{-1}) and shaking (120 rpm) at 20°C for 2 weeks before the assays, which allowed the cultures to reach relatively high densities. We used a clonal line of the rotifer predator *Brachionus calyciflorus* Pallas (Pallas 1766) derived from an isolate sampled in Milwaukee harbor (Bennett and Boraas 1989). This clonal line lost the ability to reproduce sexually and the possibility for evolutionary changes through genetic mixing (Fussmann et al. 2003). Rotifer stocks were kept in 1 L bottles filled with sterile growth medium with 800 µmol NO_3^- L^{-1} (Barreiro and Hairston 2013) and fed with the nutrient-rich chlorophyte alga *Monoraphidum minutum* obtained from the Culture Collection of Algae (University of Göttingen, Germany).

Prey morphology

To assess the variation in morphological traits among strains, *C. reinhardtii* strains were sampled (100 µL) from the culture flasks and transferred into 1.5 mL tubes. Samples were mixed and imaged using image flow cytometry (Amnis® ImageStreamX Mk II). A collection of 5000 images each showing a single cell or a cell colony was acquired for each strain with a 20X magnification using the Cy5 channel (642 nm) and the autofluorescence signal of algal

cells. We chose 9 morphological features (Tab. S2) that were calculated per image using the

instrument analysis software (Amnis IDEAS®, see *Supplementary Materials Appendix 2*).

Features were divided in 2 categories: cell size and cell shape. We used a principal components

analysis (PCA) on the morphological features estimated on individual cell images to identify

the most relevant morphological traits for characterizing strains. The PCA was run on the image

features and the 2 first dimensions explaining most of the variance in the data were selected (D_1

= 69.43%, D_2 = 18.67%). We performed parametric correlations between 2 relevant features

associated to cell size (feature: *area*) and cell shape (feature: *roundness*) according to

contributions to the PCA dimensions. Statistical differences among strains for these 2 features

were tested using Kruskal-Wallis and Dunn post-hoc tests. The position of features on the PCA

dimensions (Fig. S4) and the correlations (Fig. S5) showed a significant relationship between

area and *roundness* of prey strains. We chose to include the feature *area* representing prey

particle size, which includes single cells or cell colonies, as a proxy trait for anti-grazing defense

(Lürling 2021).

Prey stoichiometry

To assess the variation in stoichiometric traits among strains, *C. reinhardtii* strains were

sampled (10 mL) from culture flasks and centrifuged at 2000 rpm for 10 min at 20 °C. The

culture medium was removed and the pellets were suspended in 5 mL of fresh culture medium

at a density of 5×10^5 cells mL^{-1}. Carbon and nitrogen cell masses were measured following a

combustion oxidation method (Bremner 1965). Algae were filtrated on 0.2 µm-mesh filters that

were pre-calcinated for 2 h in a muffle furnace at 550°C to remove carbon. Filters were dried

for 24 h in an oven at 60°C and stored for 24 h in a desiccator dome at 20°C. Dried filters were

then packed in tin capsules and oxidated by combustion at 1020°C into a gas chromatographer

to estimate carbon and nitrogen masses (Hekatech® Euro Elemental Analyzer). The

experimental designs included 3 replicates for each strain and 3 control treatments of culture

medium. Cell mass of each element (μg cell^{-1}) was calculated as: $M = (M_S - M_C) / (D \times V)$,

where M_S is the total element concentration in the algae solution (μg mL^{-1}), M_C is the total

element concentration in the control solution (μg mL^{-1}), D is the cell density (cells mL^{-1}) and V

is the volume of the filtrate (mL).

Prey defense

Prior to the experiment, *C. reinhardtii* strains were sampled from culture flasks and centrifuged

at 2000 rpm for 10 min at 20 °C. The culture medium was removed and the pellets were

suspended in nitrate-free medium at a density of 1×10^6 cells mL^{-1} for 24 h to significantly slow

down algal growth during the exposure to predation. At the start of the experiment, strain

solutions were diluted to 10 different cell densities (1×10^4 to 1×10^6 cells mL^{-1}). These dilutions

were made in 2 mL tubes by mixing the strains with nitrate-free medium and 200 μL were

transferred into wells of 96 well plates. The experimental design included 3 replicates per cell

density for each strain for a control treatment without exposure to predation to estimate the

initial densities in the wells and for a predation treatment to estimate the final densities after

exposure to predation. For the predation treatment, 4 adult rotifers without carrying eggs were

introduced per well and allowed to feed for 8 h at 20°C under continuous light. The duration of

the experiment did not allow for rotifer reproduction to maintain a constant predation pressure.

Afterwards, cells were fixed by adding 10 μL of Lugol solution per well and plates were stored

at 4°C in the dark to allow cells to settle to the bottom of the wells. Initial and final cell densities

were assessed by taking 21 images per well using a Cy5 filter set and the autofluorescence of

the algal cells (642 nm) under a 10x magnification using a high content microscope

(ImageXpress® Micro 4 High-Content Imaging System). Images were analyzed and cell

densities were calculated using a custom module within an analysis software (MetaXpress®

High Content Image Acquisition and Analysis, *Supplementary Materials Appendix 1*).

Ingestion rates were calculated as $i_R = (C_{RI} - C_{RF}) / (t \times B_C)$, where C_{RI} is the mean initial cell

density of replicate wells from the control treatment (cells mL^{-1}), C_{RF} is the final cell density of

each well from the predation treatment (cells mL^{-1}), t is the time of feeding (sec) and B_C is the

number of rotifers per well (individuals).

To obtain predator funtionnal responses and derive predator attack rates (a_i) and

handling times (h_i), the mean ingestion rate of the 3 replicates was calculated over the range of

cell densities for each strain. Following the disc equation (Holling 1959), functional responses

were expressed as mean ingestion rate of the predator (i_{Ri}) as a function of prey densities (C_{Ri})

and predator ingestion parameters. We fitted the Holling type I, II and III (Holling 1965) and

the Ivlev type II models (Ivlev 1961) for each strain using non-linear least squares regressions

and compared their AIC values (Tab. S2). The best fitting model was the Holling type II model

for strains C_{R2}, C_{R3}, C_{R6}, C_{R7} (Eq. 1) and Holling type III model for strains C_{R1} and C_{R4} (Eq. 2):

$$i_{Ri} = \frac{a_i C_{Ri}}{1 + a_i h_i C_{Ri}} \qquad\qquad\qquad\qquad\qquad\qquad \text{(Eq. 1)}$$

$$i_{Ri} = \frac{a_i C_{Ri}{}^{c_i}}{1 + a_i h_i C_{Ri}{}^{c_i}} \qquad\qquad\qquad\qquad\qquad\qquad \text{(Eq. 2)}$$

where a_i is the attack rate of the predator on strain i (10^{-6} mL sec^{-1}), h_i is the handling time of

the predator on strain i (sec), c_i is the prey density exponent of the predator on strain i and C_{Ri}

is the cell density of strain i (cells mL^{-1}). Because prey density declined over the duration of the

experiment due to predation, we incorporated the decrease in prey density for the calculation of

ingestion rates (Rosenbaum and Rall 2018). For this, we computed of system of differential

equations to simulate dynamics of prey density over time and solved the Holling type II or type

III model for each strain using an iterative maximum likelihood estimation (hereafter: prey

density-corrected ingestion rate). We then derived the attack rates a_i and handling times h_i of

each strain from the fitted models (see *Supplementary Materials Appendix 3*).

Prey competitiveness

Following the same protocol described above, *C. reinhardtii* strains were concentrated 24 h

prior to the experiment. We measured growth under 7 different nitrate concentrations: 6 low

concentrations ($1-10\,\mu mol\ NO_3^-\ L^{-1}$) and 1 high concentration ($100\,\mu mol\ NO_3^-\ L^{-1}$) to calculate

growth rates, nitrate affinity constants and nitrate half-saturation constants. At the start of the

experiment, strain solutions were diluted to an initial cell density of 1×10^3 cells mL^{-1}. Growth

assays were made in 96 well plates in a volume of 200 µL by mixing the strains (20 µL) and

medium (180 µL). We used a disruptive sampling method: a well plate was assigned to each

day and cells were fixed daily in one of the replicated well plates by adding 10 µL of Lugol

solution per well. The experimental design included 3 technical replicates per nitrate

concentration for each strain. Plates with fixed cells were stored at 4°C in the dark for 24 h to

allow cells to settle to the bottom of the wells. Cell densities were assessed as described above

and growth curves were estimated over 8 days.

To derive prey maximum growth rates (r_{Cimax}), nitrate affinity constants (f_{Ci}) and half-

saturation constants (K_{Ci}), we computed non-linear least squares regressions on prey growth

rates as a function of nitrate concentrations for each strain using the Monod model (Eq. 4):

$$r_{Ci} = r_{Cimax}\frac{N}{K_{Ci} + N} \hspace{4cm} (Eq.\,4)$$

where r_{Cimax} is the maximum growth rate of strain i among nitrate concentrations (day^{-1}), K_{Ci} is

the nitrate half-saturation constant of strain i ($\mu mol\ NO_3^-\ L^{-1}$) and N is nitrate concentration

($\mu mol\ NO_3^-\ L^{-1}$). Nitrate affinity constants of strains were calculated as $f_{Ci} = r_{Cimax}\ /\ K_{Ci}$. To

obtain prey growth rates (r_{Ci}), we computed linear regressions on the growth curves for each

strain and nitrate concentration. We applied these regressions on the first 3 days of the growth

curves where nitrate concentrations were expected to be non-limiting for all strains.

To obtain prey growth curves at different nitrate concentrations, the mean cell density

of the 3 replicates was calculated for each strain, nitrate concentration and day. Following

previous studies (Paine et al. 2012; Malerba et al. 2018), we chose 2 non-asymptomatic growth

models which assume an unlimited cell growth over time (i.e., linear and exponential models)

and 3 asymptotic growth models which assume a limited cell growth with a maximum cell

density reached over time (i.e., logistic, gompertz and mortality models). We fitted the growth

models to each combination of strain and nitrate concentration using non-linear least squares

regressions on cell densities as a function of time and compared their AIC values (Tab. S4).

The best fitting model was the logistic model for strains C_{R1}, C_{R2}, C_{R3}, C_{R6} and C_{R7} (Eq. 3) and

the mortality model for strain C_{R4} (Eq. 5):

$$C_{Ri} = \frac{C_{R0i} c_{Ci}}{C_{R0i} + (c_{Ci} - C_{R0i}) e^{-r_{Ci}t}}$$ (Eq. 5)

$$C_{Ri} = c_{Ci} e^{m_{Ci}(t-t_{mi}) - \frac{m_{Ci}(1 - e^{-r_{Ci}(t-t_{mi})})}{r_{Ci}}}$$ (Eq. 6)

where C_{Ri} is the cell density of strain i at time t (cells mL^{-1}), C_{R0i} is the initial cell density of

strain i (cells mL^{-1}), c_{Ci} is the asymptotic cell density of strain i (cells mL^{-1}), r_{Ci} is the intrinsic

growth rate of strain i (cells day^{-1}), m_{Ci} is the decreasing slope following maximum cell density

of strain i (cell day^{-1}), t is the time (day) and t_{mi} is the time of maximum intrinsic growth rate of

strain i (day).

Prey fitness

We used the prey growth curves and the prey growth rates under the highest nitrate conditions

in the experiment described above (100 µmol NO$_3^-$ L^{-1}) to infer the differences in prey fitness

among *C. reinhardtii* strains.

Predator fitness

Prior to the experiment, *C. reinhardtii* strains were sampled from culture flasks and

concentrated in nitrate-free medium at a density of 1×10^6 cells mL^{-1} as described above. At the

start of the experiment, strain densities were estimated and diluted to 5×10^5 cells mL^{-1} in 1 mL

of a nitrate-free medium in 24 well plates. The nitrate-free medium prevented prey growth and

the high initial cell density provided sufficient resource for maximal predator growth. We

introduced 2 juveniles and 3 adults *B. calyciflorus* carrying 1 or 2 eggs per well to maintain a

homogeneous age structure in the initial population. The experimental design included 5

replicates for each strain. Rotifer individuals were daily counted using a stereomicroscope and

growth curves were estimated over 5 days.

To obtain predator growth curves, the mean rotifer density of the 5 replicates was

calculated for each strain and day. We chose 2 non-asymptotic growth models (i.e., linear and

exponential models) and 2 asymptotic growth models (i.e., logistic model and mortality

models). We fitted the growth models using non-linear least squares regressions on rotifer

densities as a function of time and compared their AIC values (Tab. S5). The best fitting model

was the mortality model for strains C_{R1}, C_{R2}, C_{R3} and C_{R4} (Eq. 4) and the logistic model for

strains C_{R6} and C_{R7} (Eq. 3). We computed linear regressions on the growth curves for each

strain to obtain the predator growth rates (r_{Bi}). We used the predator growth curves and the

predator growth rates to infer the differences in predator fitness among *C. reinhardtii* strains.

Statistical analysis

All statistical analyses were computed using the R software v.4.0.4 (R Development Core Team

2020). Linear regressions and non-linear least squares regressions were computed using the *lm*

and *nls* functions from the 'lme4' R package. Confidence intervals for non-linear least squares

regressions were computed using the *predictnls* function from the 'propagate' R package.

Models were compared using the *AIC* function from the 'stats' R package. Maximum likelihood

regressions were computed using the *mle2* function from the 'bbmle' R package. Parametric

correlations were computed using the function *cor* from the 'stats' R package. The principal

components analysis was computed using the *PCA* function from the 'FactoMineR' R package.

Graphics were obtained using the *ggplot* function with associated subfunctions from the

'ggplot2' R package.

Results

Prey traits

The defense of *C. reinhardtii* strains was defined based on the maximum ingestion rate of the predator *B. calyciflorus* and ranged from defended (low ingestion) to undefended (high ingestion). *B. calyciflorus* ingestion rates differed among strains (Fig. 1a). Strains C_{R1}, C_{R2} and C_{R3} were classified as more defended due to the low maximum ingestion rates exhibited by the predator while strains C_{R4}, C_{R6} and C_{R7} were classified as less defended due to the high maximum ingestion rates exhibited by the predator. The comparison of functional response models revealed that the ingestion rates of *B. calyciflorus* on *C. reinhardtii* strains were better described by non-linear saturating models (Tab. S3 and Fig. S1a). The differences in defense between strains were evident by the shapes of the functional responses (Fig. 1a) and lower attack rates but higher handling times of *B. calyciflorus* on more defended strains compared to less defended strains (Tab. 1). The competitiveness of *C. reinhardtii* strains was classified based on the maximum growth rate and ranged from less competitive (low growth) to more competitive (high growth). Growth parameters related to nitrate differed among *C. reinhardtii* strains (Fig. 1b). More defended strains C_{R1}, C_{R2} and C_{R3} were classified as less competitive due to the low nitrate affinity and the high nitrate half-saturation constants while less defended strains C_{R4}, C_{R6} and C_{R7} were classified as more competitive due to the high nitrate affinity and the low half-saturation constants (Tab. 1).

Figure 1: Defense and competitiveness of *C. reinhardtii* strains. (a) Functional responses of *B. calyciflorus* showing ingestion rate as a function of cell density of *C. reinhardtii* strains. Predicted functional response curves (lines) were estimated from the 3 replicates for ingestion rates (dots) for raw data (dotted lines) and prey density-corrected data (solid lines). (b) Monod curves of *C. reinhardtii* strains showing growth rate as a function of nitrate concentration ($1-10\,\mu$mol NO$_3^-$ L^{-1}). Predicted curves were estimated from the growth rate at each nitrate concentration (dots). Predicted curves with lower and upper boundaries using standard errors are shown in the *Supplementary Materials* (Fig. S1a and S1b).

Prey morphology

To investigate whether the observed differences in defensive and competitive traits of *C.*

reinhardtii strains were related to morphological differences among strains, we analyzed

morphological traits in more detail. Strains exhibiting strong clumping morphologies (C_{R1}, C_{R2}

and C_{R3}) displayed populations covering a larger spectrum of the PCA dimensional space (Fig.

S4a) compared to strains exhibiting mostly single cell morphologies (C_{R4}, C_{R6} and C_{R7}).

Clumping strains were positively associated to traits indicating large asymmetric cell

aggregates (e.g., area and lobe count, Fig. S4b) and negatively associated to traits indicating

small symmetric single cells (e.g., aspect ratio and roundness, Fig. S4b). This was confirmed

by the negative correlations found between area and roundness for clumping strains (e.g., C_{R2}:

Pearson correlation, $r = -0.43$, $P < 0.001$) but not for non-clumping strains (e.g., C_{R4}: Pearson

correlation, $r = 0.07$, $P < 0.001$). Accordingly, significant differences among strains were found

for area (Kruskal-Wallis, $\chi^2 = 14.37$, $P = 0.013$) and roundness (Kruskal-Wallis, $\chi^2 = 8177.60$,

$P < 0.001$) indicating that clumping strains exhibited on average larger size and less symmetric

shapes (Fig. S6). Overall, the presence or absence of clumping morphotypes coincided with the

defensive and competitive traits of strains, meaning that strains having large cell clumps were

classified as very defended and poorly competitive and vice versa (Fig. 3).

We found strong negative relationships between anti-grazing defense and competitiveness traits

independent of the trait used (Fig. 2). Significant relationships were found between attack rate,

handling time or predator growth rate and prey maximum growth rate or nitrate affinity constant

(Tab. 2). Except for C:N ratio, non-significant relationships between traits showed persistent

relative positions in trait spaces with non-overlapping standard deviations. The more defended

strains were consistently less competitive and vice versa. More defended strains (C_{R1}, C_{R2} and

C_{R3}) were characterized by large-sized cell clumps (e.g., C_{R1}: 323 ± 230 µm), resulting in low

attack rates (e.g., C_{R1}: 0.19 ± 0.04 10^{-6} mL sec^{-1}) and high handling times (e.g., C_{R1}: 4.64 ± 0.80

sec). Conversely, the more competitive strains (C_{R4}, C_{R6} and C_{R7}) were characterized by small-

sized single cells (e.g., C_{R4}: 108 ± 35 µm), resulting in high attack rates (e.g., C_{R4}: 1.55 ± 0.11

10^{-6} mL sec^{-1}) and low handling times (e.g., C_{R4}: 0.56 ± 0.09 sec). The positions of strains in

the trait space were similar for trait spaces constructed with prey maximum growth rates (Fig.

2a) or growth parameters related to nitrate, such as nitrate affinity constants (Fig. 2b) and half-

saturation constants (Fig. 2c), as traits for competitiveness.

Figure 2: Defense-competitiveness trait spaces of *C. reinhardtii* strains. Shown are the positions for

four anti-grazing defense traits (attack rate a, handling time h, prey particle size s_C and carbon to nitrogen

ratio $C{:}N_C$) as a function of three competitiveness traits (prey maximum growth rate r_{Cmax}, nitrate affinity

constant f_C and nitrate half-saturation constant K_C). Trait estimates (dots) and standard deviations

(horizontal and vertical bars) were represented pairwise for competitiveness traits on each plot: (a) prey

maximum growth rate, (b) prey nitrate affinity constant and (c) prey nitrate half-saturation constant.

Linear regression lines showing the significant correlations between anti-grazing defense and

competitiveness traits were represented (grey lines) and non-significant correlation were indicated by a

label (*NS*).

Prey and predator fitness

The defensive and competitive traits of *C. reinhardtii* strains had an impact on the population

growth of the prey and the predator (Fig. 3). The comparison of prey growth models revealed

that *C. reinhardtii* strains had a logistic growth (Tab. S4 and Fig. S1b). More defended strains

had lower growth rates and asymptotic densities under non-limiting nitrate conditions compared

to less defended strains (Tab. S6). The less defended strain C_{R4} showed a rapid increase

followed by a decrease in cell density probably attributed to depletion of nitrate and/or shading

at high densities (Fig; 3a). The comparison of predator growth models revealed that *B.*

calyciflorus had a logistic growth in the presence of the less defended strains C_{R6} and C_{R7} but a

logistic growth followed by a decline in the presence of the more defended strains C_{R1}, C_{R2} and

C_{R3} and the less defended strain C_{R4} (Fig. 3b; Tab. S5 and Fig. S1c). More defended strains

were associated with lower predator growth rates (e.g., C_{R1}: -0.05 ± 0.18 day^{-1}) and asymptotic

densities while less defended strains were associated with higher predator growth rates (e.g.,

C_{R4}: 2.33 ± 0.27 day^{-1}) and asymptotic densities (Tab. S6). The less defended strain C_{R4} showed

a rapid increase in rotifer density followed by a decrease probably attributed to a strong prey

depletion.

Figure 3: Fitness *C. reinhardtii* strains and *B. calyciflorus*. (a) Growth curves of *C. reinhardtii* strains

shown as cell densities over time for a high nitrate concentration (100 µmol NO_3^- L^{-1}). Predicted growth

curves (lines) were estimated from the 3 replicates for cell densities (dots). (b) Growth curves of *B.*

calyciflorus feeding on *C. reinhardtii* strains shown as rotifer densities over time. Predicted growth

curves (lines) were estimated from the 5 replicates of rotifer densities (dots). Predicted curves with lower

and upper boundaries using standard errors are shown in the *Supplementary Materials* (Fig. S2a and

S2b).

Discussion

Discussion

Defense-competitiveness trade-off relationships are often observed in plankton organisms and

can be expressed with multiple traits (Pančić and Kiørboe 2018). Determining the relevant traits

describing these trade-offs is essential to predict the outcome of trophic interactions (Cadier et

al. 2019). Comparing multiple anti-grazing defense and competitiveness traits, we found

substantial variation among strains of *C. reinhardtii* and overall negative relationships between

these traits. The negative relationships between defense and competitiveness traits suggests a

colimitation in energy investment and/or an evolutionary constrain between survival and

reproduction (Stearns 1989). This observation also shows that multiple morphological and

trophic traits of prey and predator can be suitable proxies to estimate trade-off relationships and

a mechanistic link between prey structural traits and predator ingestion and reproduction traits

(Litchman et al. 2013). The differences in defense and competitiveness traits and their

relationships translated into different fitness consequences for prey and predator and thus their

interaction strength. We found lower population sizes of *B. calyciflorus* in the presence of more

defended *C. reinhardtii* strains, confirming that prey defensive and competitive traits, predator-

prey trophic interactions and predator reproduction are closely connected (Litchman et al.

2013). Our study confirms previous observations comparing trait variation between species or

within species of higher taxonomic groups (Litchman et al. 2013; Ehrlich et al. 2020) and

improves the understanding of planktonic communities (Litchman et al. 2007; 2010) by

demonstrating the mechanistic links between traits, trade-offs and fitness at the intraspecific

level.

Our data suggest a mechanistic relationship at the intraspecific level linking prey

morphological traits to predator trophic traits. *C. reinhardtii* strains classified as defended

against ingestion by the predator were growing as cell groups bound by a viscous extracellular

matrix. These strains were associated with lower predator functional responses which were

mainly the consequence of the gape-size limitation of the predator for ingesting cell groups

within an extracellular matrix (Reynolds 2006). Lower ingestion rates of cell colonies could

also indicate longer manipulation and digestion times for the predator (Verschoor et al. 2007),

and/or a resistance to digestion during gut passage (DeMott et al. 2010). Predator traits such as

attack rate and handling time are reliable indicators of prey defense (Jeschke and Tollrian 2000)

and other traits related to ingestion such as digestion time and conversion efficiency (Jeschke

et al. 2002) should be further explored. Prey morphological traits can also provide relevant

estimates of prey defense (e.g., Becks et al. 2010; Ellner and Becks 2011). Prey size and shape

is a particularly pertinent trait as it is directly associated with gape-size limitation for the

predator. Some caution is still required with prey size and shape as other morphological traits

(e.g., spines) or other physiological (e.g., toxins) and behavioral (e.g., motility) traits can

contribute to prey defense (Pančić and Kiørboe 2018; Lürling 2021) and the relative

contribution of these traits to prey defense is often unclear (Van Donk et al. 2011).

Our data also suggest a mechanistic relationship at the intraspecific level linking prey

morphological traits to prey growth traits. *C. reinhardtii* strains classified as competitive were

associated with higher growth rates and nitrate affinities but lower nitrate half-saturation

constants, which indicated enhanced abilities for nutrient acquisition or conversion to

reproduction (Litchman et al. 2007; 2010; Halsey and Jones 2015; Ward et al. 2017).

Competitive abilities of phytoplankton species mainly depend on surface-area constrains in

terms of resource acquisition and requirements for basal maintenance and functions

contributing to fitness. Single cells tend to have higher nutrient acquisition rates due to a more

efficient diffusion of nutrients (Yoshiyama and Klausmeier 2008) while palmelloid colonies

are more limited by nutrient and light competition among colonial cells (Raven and Kubler

2002), despite larger nutrient storage capacities (Litchman et al. 2009). Prey growth rate and

nutrient acquisition rate are reliable indicators of prey competitiveness and show a relation

between the processes of nutrient absorption, assimilation, and conversion to reproduction

(Litchman et al. 2007). However, complex interplays between processes may complicate

predictions as prey competitiveness can also be driven by other regulating factors such as the

acquisition of colimiting resources (Klausmeier et al. 2004) or trade-offs between nutrient and

light acquisition (Strzepek and Harrison 2004).

The intraspecific covariation in defense and competitiveness traits of phytoplankton

have consequences on the growth of zooplankton. Prey defense, including morphological and

physiological traits, and prey stoichiometry have been shown to affect food quantity and quality

(Bi and Sommer 2020), which prevents the predator from meeting its energetic requirements

and thus reduces its survival and reproduction (Sommer 2008). Prey stoichiometry is often used

as a good predictor of prey defense and competitiveness traits and thus predator fitness (Sterner

and Elser 2002) as C:N (and C:P) ratio increases due to the high carbon requirements when

cells form colonies. Conversely, C:N (and C:P) ratio decreases due to the high nitrogen and

phosphorus requirements of cell replication (Branco et al. 2020). Such differences in prey

stoichiometry can impact predator growth rate (Branco et al. 2010; 2018; Mandal et al. 2018)

but we did not observe differences in C:N ratios among *C. reinhardtii* strains. Whether this is

due to the precise growth conditions (e.g., nutrient concentration or light intensity) or indicates

a difference between intraspecific and interspecific variation should be further studied.

We used 6 strains of *C. reinhardtii* with a known recent evolutionary history (Bernardes

et al. 2021) to examine defense-competitiveness trade-off relationships. The isolates came from

an evolution experiment in which either defense or competitiveness was selected for and a

general trade-off between survival against predation and replication was observed across a large

range of isolates (Bernardes et al. 2021). Previous approaches consisted in split-family

experiments in which clones were evolved in different environments and thus this could affect

the fitness of clones in an opposite way leading to no or positive correlations between traits

(Fry 1993; 1996). The experimental evolution approach has been proven to be more suitable

for detecting trade-offs than the split-family approach (Kassen 2002; Fry 2003; Fuller et al.

2005). Selection experiments can also be used to determine the role of phenotypic plasticity

(Scheiner 2002) and environmental heterogeneity (Kassen 2002) versus genetic adaptation for

the maintenance of trait variation (Roff and Fairbairn 2007). Overall, this approach allows

testing whether 2 traits can be optimized concurrently or whether optimization of one trait leads

to a reduction in the other trait.

460   Investigating intraspecific relationships between defensive and competitiveness traits is

fundamental to understand fitness strategies of interacting organisms (Ehrlich et al. 2017;

Bartlett et al. 2018) and coexistence between resources and consumers (Lichstein et al. 2007;

Petry et al. 2018; Ehrlich et al. 2020). Our study demonstrates intraspecific defense-

competitiveness trade-offs via negative relationships estimated over a range of traits measured

on both prey and predator. Assessing the magnitude and the shape of intraspecific trade-offs

can be decisive as empirical and theoretical studies showed the significant role of trade-offs for

consumer-resource dynamics (e.g., Yoshida et al. 2003; Becks et al. 2010; Cortez and Ellner

2010; Kasada et al. 2014; Ehrlich et al. 2017). Knowledge on the underlying traits involved in

trade-offs and how these trade-offs translate into fitness of interacting organisms at different

trophic levels can provide key insights into the transfers of energy and matter in food webs

(Våge et al. 2014; 2018; Cadier et al. 2019) and the response of populations to environmental

perturbations (Jessup and Bohannan 2008; Frickel et al. 2017; Theodosiou et al. 2019).

Intraspecific trade-offs are for instance a prerequisite for eco-evolutionary feedback dynamics,

which can contribute to the maintenance of trait variation (e.g., Becks et al. 2010; Kasada et al.

2014) and in turn allow the persistence of populations (Yamamichi and Miner 2015; Hermann

and Becks 2022). Investigating intraspecific trade-offs thus offers a promising avenue for

assessing trait variation within population and predict the strength of trophic interactions,

population dynamics and ecosystem processes.

Acknowledgements – We are thankful to U. Gaedke and Z. Zhang for comments on an

earlier version the manuscript, to E. Ehrlich for the advice on the data analysis, and to R.

Hermann and R. Lambrecht for the theoretical and practical contributions to the experiments.

Author contributions – TR and LB compiled the first theoretical ideas. TR and LB

designed the experiments. TR conducted the laboratory experiments and statistical analyses

with the assistance of LB. TR wrote the initial drafts of the manuscript. TR and LB wrote the

final version of the manuscript.

Conflicts of interest – The authors have no conflicts of interest to declare.

Funding – This work was supported by the German Research Foundation (DFG) to LB (BE

4135/4-2) as part of the priority program "Flexibility Matters: Interplay between Trait Diversity

and Ecological Dynamics Using Aquatic Communities as Model Systems – DynaTrait" (SPP

1704).

Tables

Table 1: Estimates of defense and competitiveness traits [± standard deviation] for the 6 different *C. reinhardtii* strains. Traits were combined pairwise to construct the different defense-competitiveness trait spaces (Fig. 2). Anti-grazing defenses were represented by *B. calyciflorus* attack rate (a in 10^{-6} mL sec^{-1}), handling time (h in sec), *C. reinhardtii* particle size (s_C in 10^3 µm^2) and carbon to nitrogen ratio (C:N_C). Competitiveness ability was represented by *C. reinhardtii* maximum growth rate (r_{Cmax} in day^{-1}), nitrate affinity constant (f_C in µmol NO$_3^-$ L^{-1} day^{-1}) and nitrate half-saturation constant (K_C in µmol NO$_3^-$ L^{-1}). Strains exhibited different morphologies in terms of cell clumping: clumping strains were C_{R1} (large clumps [10−30 cells]), C_{R2} (medium clumps [5−10 cells]) and C_{R3} (medium clumps [5−10 cells]) while non-clumping strains were C_{R4} (small single cells), C_{R6} (small single cells) and C_{R7} (small single cells).

Strain	a	h	s_C	r_{Cmax}	f_C	K_C	$C{:}N_C$
C_{R1}	0.21 [0.17–0.26]	7.08 [5.90–8.85]	3.23 [0.93–5.53]	0.12 [0.11–0.13]	0.15 [0.13–0.17]	1.11 [0.84–1.38]	15.62 [15.28–15.95]
C_{R2}	0.54 [0.43–0.65]	6.52 [5.43–8.15]	1.78 [0.60–2.96]	0.10 [0.09–0.11]	0.06 [0.05–0.07]	1.90 [1.12–2.68]	15.31 [14.02–16.60]
C_{R3}	0.39 [0.31–0.47]	4.69 [3.90–5.86]	1.71 [0.60–2.82]	0.11 [0.10–0.12]	0.07 [0.06–0.08]	1.77 [1.50–2.04]	15.33 [14.09–16.57]
C_{R4}	1.68 [1.57–1.78]	0.60 [0.52–0.67]	1.08 [0.73–1.42]	0.33 [0.32–0.34]	0.43 [0.40–0.47]	0.63 [0.44–0.83]	12.90 [12.58–13.23]
C_{R6}	1.49 [1.21–1.75]	2.27 [2.00–2.55]	1.21 [0.67–1.74]	0.34 [0.33–0.35]	0.50 [0.46–0.55]	0.79 [0.64–0.93]	14.97 [14.53–15.41]
C_{R7}	1.53 [1.24–1.80]	2.65 [2.35–2.95]	1.36 [0.91–1.81]	0.32 [1031–0.33]	0.42 [0.40–0.46]	0.74 [0.57–0.92]	15.73 [15.12–16.33]

Table 2: Parameter estimates for parametric correlation tests performed between defense and
competitiveness traits. Correlations were represented by regression lines on the different
defense-competitiveness trait spaces (Fig. 2). Anti-grazing defense was represented by *B.*
calyciflorus attack rate (a in 10^{-6} mL sec^{-1}), handling time (h in sec), *C. reinhardtii* particle size
(s_C in 10^3 µm^2) and carbon to nitrogen ratio ($C{:}N_C$). Competitiveness was represented by *C.*
reinhardtii maximum growth rate (r_{Cmax} in day^{-1}), affinity constant (f_C in µmol NO$_3^-$ L^{-1} day^{-1})
and half-saturation constant (K_C in µmol NO$_3^-$ L^{-1}) for nitrate. Significance was estimated with
a Pearson correlation test (t, df and P values).

Defense trait	Competitiveness trait	r	t	df	P
a		0.97	7.90	4	0.001
h		−0.91	−4.41	4	0.012
s_C	r_{Cmax}	−0.69	−1.89	4	0.132
$C{:}N_C$		−0.47	−1.06	4	0.350
a		0.97	7.85	4	0.001
h		−0.93	−4.86	4	0.008
s_C	f_C	−0.66	−1.76	4	0.154
$C{:}N_C$		−0.58	−1.42	4	0.229
a		−0.79	−2.59	4	0.061
h		0.73	2.14	4	0.099
s_C	K_C	0.30	0.63	4	0.566
$C{:}N_C$		0.42	0.93	4	0.404

References

Agrawal, A. A (1998). Algal defense, grazers, and their interactions in aquatic trophic cascades. *Acta Oecologia*, 19: 331–337.

Barreiro, A. and N. G. Hairston (2013). The influence of resource limitation on the allelopathic effect of *Chlamydomonas reinhardtii* on other unicellular freshwater planktonic organisms. *Journal of Plankton Research*, 35: 1339–1344.

Bartlett, L. J., L. Wilfert and M. Boots (2018). A genotypic trade-off between constitutive resistance to viral infection and host growth rate. *Evolution*, 72: 2749–2757.

Bateman, A. W., M. Vos and B. R. Anholt (2014). When to defend: antipredator defenses and the predation sequence. *The American Naturalist*, 183: 847–855.

Becks, L., S. P. Ellner, L. E. Jones and N. G. Hairston (2010). Reduction of adaptive genetic diversity radically alters eco-evolutionary community dynamics. *Ecology Letters*, 13: 989–997.

Bennett, W. N. and M. E. Boraas (1989). Comparison of population dynamics between slow- and fast-growing strains of the rotifer *Brachionus calyciflorus* pallas in continuous culture. *Oecologia*, 81: 494–500.

Bernardes, J. P., U. John, N. Woltermann, M. Valiadi, R. J. Herrman and L. Becks (2021). The evolution of convex trade-offs enables the transition towards multicellularity. *Nature Communication*, 12: 4222.

Bi, R. and U. Sommer (2020). Food quantity and quality interactions at phytoplankton-zooplankton interface: chemical and reproductive responses in a calanoid copepod. *Frontiers in Marine Science*, 7: 274.

Branco, P., M. Egas, J. J. Elser and J. Huisman (2018). Eco-evolutionary dynamics of ecological stoichiometry in plankton communities. *The American Naturalist*, 192: 1–20.

Branco, P., M. Egas, S. R. Hall, and J. Huisman (2020). Why do phytoplankton evolve large size in response to grazing?. *The American Naturalist*, 195: 20–37.

Branco, P., M. Stomp, M. Egas and J. Huisman (2010). Evolution of nutrient uptake reveals a trade-off in the ecological stoichiometry of plant-herbivore interactions. *The American Naturalist*, 176: 162–176.

Bremner, J. M. (1965). Total Nitrogen. In *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*, edited by A. G. Norman, p. 1149–1178, *Iowa State University Press*.

Bruggeman, J. (2011). A phylogenetic approach to the estimation of phytoplankton traits. *Journal of Phycology*, 47: 52−65.

Cadier, M., K. H. Andersen, A. W. Visser and T. Kiørboe (2019). Competition–defense tradeoff increases the diversity of microbial plankton communities and dampens trophic cascades. *Oikos*, 128: 1027–1040.

Colina, M., D. Calliari, C. Carballo and C. Kruk (2016). A trait–based approach to summarize zooplankton–phytoplankton interactions in freshwaters. *Hydrobiologia*, 767: 221–233.

Cortez, M. H. and S. P. Ellner. (2010). Understanding rapid evolution in predator-prey interactions using the theory of fast-slow dynamical systems. *The American Naturalist*, 176: 109–127.

DeMott, W. R., E. N. McKinney and A. J. Tessier (2010). Ontogeny of digestion in Daphnia:
implications for the effectiveness of algal defenses. *Ecology*, 91: 540–548.

Edwards, K. F., C. A. Klausmeier and E. Litchman (2011). Evidence for a three-way trade-off between
nitrogen and phosphorus competitive abilities and cell size in phytoplankton. *Ecology*, 92: 2085–2095.

Edwards, K.F., E. Litchman and C. A. Klausmeier (2013). Functional traits explain phytoplankton
community structure and seasonal dynamics in a marine ecosystem. *Ecology Letters*, 16: 53–63.

Ehrlich, E. and U. Gaedke (2018). Not attackable or not crackable. How pre- and post-attack defenses
with different competition costs affect prey coexistence and population dynamics. *Ecology and*
Evolution, 8: 6625–6637.

Ehrlich, E., L. Becks and U. Gaedke (2017). Trait–fitness relationships determine how trade-off shapes
affect species coexistence. *Ecology*, 98: 3188–3198.

Ehrlich, E., N. J. Kath and U. Gaedke (2020). The shape of a defense-growth trade-off governs seasonal
trait dynamics in natural phytoplankton. *The ISME Journal*, 14: 1451–1462.

Ellner, S. P. and L. Becks (2011). Rapid prey evolution and the dynamics of two-predator food webs.
Theoretical Ecology, 4: 133–152.

Fleischer, S. R., C. P. ter Horst and J. Li (2018). Pick your trade-offs wisely: predator-prey eco-evo
dynamics are qualitatively different under different trade-offs. *Journal of Theoretical Biology*, 456:
201–212.

Frickel, J., L. Theodosiou, and L. Becks (2017). Rapid evolution of hosts begets species diversity at the
cost of intraspecific diversity. *Proceedings of the National Academy of Sciences*, 114: 11193–11198.

Fry, J. D. (1993). The ''general vigor'' problem: can antagonistic pleiotropy be detected when genetic
covariances are positive? *Evolution*, 47: 327–333.

Fry, J. D. (1996). The evolution of host specialization: are trade-offs overrated? *The American*
Naturalist, 148: 84–107.

Fry, J. D. (2003). Detecting ecological trade-offs using selection experiments. *Ecology*, 84: 1672–1678.

Fuller, R. C., C. F. Baer and J. Travis (2005). How and when selection experiments might actually be
useful. *Integrative and Comparative Biology*, 45: 391–404.

Fussmann, G. F., S. P. Ellner, K. W. Shertzer and N. G. Hairston (2000). Crossing the Hopf bifurcation
in a live predator-prey system. *Science*, 290: 1358–1360.

Fussmann, G. F., S. P. Ellner and N. G. Hairston. (2003). Evolution as a critical component of plankton
dynamics. *Proceedings of the Royal Society B*, 207: 1015–1022.

Gravel, D., C. Albouy and W. Thuiller. (2016). The meaning of functional trait composition of food
webs for ecosystem functioning. *Philosophical Transactions of the Royal Society B*, 371: 20150268.

Halsey, K. H. and B. M. Jones. (2015). Phytoplankton strategies for photosynthetic energy allocation.
Annual Review of Marine Science, 7: 265–297.

Holling, C. S (1959). Some characteristics of simple types of predation and parasitism. *The Canadian*
Entomologist, 91: 385–398.

Holling, C. S (1965). The functional response of predators to prey density. *Memoirs of the*
Entomological Society of Canada, 45: 1–60.

Ivlev, V. S (1961). Experimental ecology of the feeding of fishes. Translated from Russian and edited
by D. Scott, p. 1−302, *Yale University Press*.

Jeschke, J. M. and Tollrian, R. (2000). Density-dependent effects of prey defences. *Oecologia*, 123:
391−396.

Jeschke, J. M., Kopp, M. and Tollrian, R. (2002). Predator functional responses: discriminating between
handling and digesting prey. *Ecological Monographs*, 72: 95–112

Jessup, C. M. and B. J. Bohannan (2008). The shape of an ecological trade-off varies with environment.
Ecology Letters, 11: 947–959.

Kasada, M., M. Yamamichi and T. Yoshida (2014). Form of an evolutionary trade-off affects eco-
evolutionary dynamics in a predator–prey system. *Proceedings of the National Academy of Sciences*,
111: 16035–16040.

Kassen, R (2002). The experimental evolution of specialists, generalists, and the maintenance of
diversity. *Journal of Evolutionary Biology*, 15: 173–190.

Klausmeier, C. A., E. Litchman, T. Daufresne and S. A. Levin (2004). Optimal nitrogen-to-phosphorus
stoichiometry of phytoplankton. *Nature*, 429: 171–174.

Laigle, I., I. Aubin, C. Digel, U. Brose, I. Boulangeat. and D. Gravel (2018). Species traits as drivers of
food web structure. *Oikos*, 127: 316–326.

Lichstein, J. W., J. Dushoff, S. A. Levin and S. W. Pacala (2007). Intraspecific variation and species
coexistence. *The American Naturalist*, 170: 807–818.

Litchman, E (2007). Resource competition and the ecological success of phytoplankton. In *Evolution of*
primary producers in the sea, edited by P. G. Falkowski and A. H. Knoll, p. 351−375, *Elsevier*.

Litchman, E. and C. A. Klausmeier (2008). Trait–based community ecology of phytoplankton. *Annual*
Review of Ecology, Evolution, and Systematics, 39: 615–639.

Litchman, E., C. A. Klausmeier, O. M. Schofield and P. G. Falkowski (2007). The role of functional
traits and trade-offs in structuring phytoplankton communities: scaling from cellular to ecosystem level.
Ecology Letters, 10: 1170–1181.

Litchman, E., C. A. Klausmeier, and K. Yoshiyama. (2009). Contrasting size evolution in marine and
freshwater diatoms. *Proceedings of the National Academy of Sciences*, 106: 2665–2670.

Litchman, E., P. de Tezanos Pinto, C. A. Klausmeier, M. K. Thomas and K. Yoshiyama (2010). Linking
traits to species diversity and community structure in phytoplankton. *Hydrobiologia*, 653: 15–28.

Litchman, E., M. D. Ohman and T. Kiørboe (2013). Trait–based approaches to zooplankton
communities. *Journal of Plankton Research*, 35: 473–484.

Litchman, E., K. F. Edwards and C. A. Klausmeier (2015). Microbial resource utilization traits and
trade-offs: implications for community structure, functioning, and biogeochemical impacts at present
and in the future. *Frontiers in Microbiology*, 6: 1–10.

Litchman, E., K. F. Edwards and P. W. Boyd (2021). Toward trait-based food webs: universal traits and
trait matching in planktonic predator-prey and host-parasite relationships. *Limnology and*
Oceanography, 66: 3857–3872.

Long, J. D. and M. E. Hay (2006). When intraspecific exceeds interspecific variance: Effects of
phytoplankton morphology and growth phase on copepod feeding and fitness. *Limnology and*
Oceanography, 51: 988–996.

Lürling, M (2021). Grazing resistance in phytoplankton. *Hydrobiologia*, 848: 237–249.

Malerba, M. E., M. M. Palacios and D. J. Marshall (2018). Do larger individuals cope with resource
fluctuations better? An artificial selection approach. *Proceedings of the Royal Society B*, 285: 20181347.

Mandal, S., R. A. Wilkins and J. B. Shurin (2018). Compensatory grazing by *Daphnia* generates a trade-
off between top-down and bottom-up effects across phytoplankton taxa *Ecosphere*, 9: e02537.

Pančić, M. and T. Kiørboe (2018). Phytoplankton defence mechanisms: traits and trade-offs. *Biological*
Reviews, 93: 1269–1303.

Paine, C. E., T. R. Marthews, D. R. Vogt, D. Purves, M. Rees, A. Hector and L. A. Turnbull (2012).
How to fit nonlinear plant growth models and calculate growth rates: an update for ecologists. *Methods*
Ecology and Evolution, 3: 245–256.

Petry, W. K., G. S. Kandlikar, N. J. B. Kraft, O. Godoy and J. M. Levine (2018). A competition-defence
trade-off both promotes and weakens coexistence in an annual plant community. *Journal of Ecology*,
106: 1806–1818.

R Development Core Team (2020). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, p. 1–2673.

Rall, B. C., U. Brose, M. Hartvig, G. Kalinkat, F. Schwarzmüller, O. Vucic-Pestic and O. L. Petchey
(2012). Universal temperature and body-mass scaling of feeding rates. *Philosophical Transactions of*
the Royal Society B, 367: 2923–2934.

Reynolds, C. S. (2006). Ecology of phytoplankton. Edited by C. S. Reynolds, p. 1–535, *Cambridge*
University Press.

Rosenbaum, B. and C. M. Rall (2018). Fitting functional responses: direct parameter estimation by
simulating differential equations. *Methods Ecology and Evolution*, 9: 2076–2090.

Rossberg, A. G., A. Brännström and U. Dieckmann (2010). How trophic interaction strength depends
on traits. *Theoretical Ecology*, 3: 13–24.

Scheiner, S. M. (2002). Selection experiments and the study of phenotypic plasticity. *Journal of*
Evolutionary Biology, 15: 889–898.

Smith, S. L., A. Merico, S. Hohn and G. Brandt (2014). Sizing-up nutrient uptake kinetics: combining
a physiological trade-off with size-scaling of phytoplankton traits. *Marine Ecology Progress Series*,
511: 33–39.

Sommer, U. (2008). Trophic cascades in marine and freshwater plankton. *International Review of*
Hydrobiology, 93: 506–516.

Stearns, C. S. (1989). Trade-offs in life-history evolution. *Functional Ecology*, 3: 259−268.

Sterner, R. W. and J. J. Elser (2002). Ecological stoichiometry. Edited by P. M. Vitousek, p. 1–464, *Princeton University Press*.

Strauss, S. Y., J. A. Rudgers, J. A. Lau and R. E. Irwin (2002). Direct and ecological costs of resistance to herbivory. *Trends in Ecology and Evolution*, 17: 278–285.

Strzepek, R. F. and P. J. Harrison (2004). Photosynthetic architecture differs in coastal and oceanic diatoms. *Nature*, 431: 689–692.

Sunda, W. G. and D. R. Hardison (2010). Evolutionary tradeoffs among nutrient acquisition, cell size, and grazing defense in marine phytoplankton promote ecosystem stability. *Marine Ecology Progress Series*, 40: 63–76.

Theodosiou, L., T. Hiltunen and L. Becks (2019). The role of stressors in altering eco-evolutionary dynamics. *Functional Ecology*, 33: 73–83.

Våge, S., Bratbak, G., Egge, J., Heldal, M., Larsen, A., Norland, S., Paulsen, M. L., Pree, B., Sandaa, R. A., Skjoldal, E. F. and Tsagaraki T. M. (2018). Simple models combining competition, defence and resource availability have broad implications in pelagic microbial food webs. *Ecology Letters*, 21: 1440–1452.

Våge, S., J. E. Storesund, J. Giske and T. F. Thingstad. (2014). Optimal defense strategies in an idealized microbial food web under trade-off between competition and defense. *Plos One*, 9: e101415.

Van Donk, E., M. Lürling and W. Lampert (1999). Consumer-induced changes in phytoplankton: inducibility, costs, benefits and impacts on grazers. In *Ecology and Evolution of Inducible Defenses*, edited by R. Tollrian and C. D. Harvell, p. 89–103, *Princeston University Press*.

Van Donk, E., A. Ianora and M. Vos (2011). Induced defences in marine and freshwater phytoplankton: a review. *Hydrobiologia*, 668: 3–19.

Verschoor, A. M., Y. Zadereev and W. M. Mooij (2007). Infochemical-mediated trophic interactions between the rotifer Brachionus calyciflorus and its food algae. *Limnology and Oceanography*, 52: 2109–2119.

Ward, B. A., E. Marañón, B. Sauterey, J. Rault, and D. Claessen (2017). The size dependence of phytoplankton growth rates: a trade-off between nutrient uptake and metabolism. *The American Naturalist*, 189: 170–177.

Weiss, L., C. Laforsch and R. Tollrian (2012). The taste of predation and the defences of prey. In *Chemical ecology in aquatic systems*, edited by C. Brönmark and L. A. Hansson, p. 111–126, *Oxford University Press*.

Wood, Z. T., D. C. Fryxell, E. R. Mofett, M. T. Kinnison, K. S. Simon and E. P. Palkovacs (2020). Prey adaptation along a competition-defense tradeof cryptically shifts trophic cascades from density- to trait-mediated. *Oecologia*, 192:767–778.

Yamamichi, M. and B. E. Miner (2015). Indirect evolutionary rescue: prey adapts, predator avoids extinction. *Evolutionary Applications*, 8: 787–795.

Yoshida, T., L. E. Jones, S. P. Ellner, G. F. Fussmann and N. G. Hairston (2003). Rapid evolution drives ecological dynamics in a predator–prey system. *Nature*, 424: 303−306.

Yoshida, T., N. G. Hairston and S. P. Ellner (2004). Evolutionary trade-off between defence against
grazing and competitive ability in a simple unicellular alga, Chlorella vulgaris. *Proceedings of the Royal*
Society B, 271: 1947–1953.

Yoshiyama, K. and C. A. Klausmeier (2008). Optimal cell size for resource uptake in fluids: a new facet
of resource competition. *The American Naturalist*, 171: 59−70.

Zust, T. and Agrawal, A. A. (2017). Trade-offs between plant growth and defense against insect
herbivory: an emerging mechanistic synthesis. *Annual Review of Plant Biology*, 68: 513–534.