Decoding of Speech Information using EEG in Children with Dyslexia:

Less Accurate Low-Frequency Representations of Speech, Not "Noisy" Representations

Mahmoud Keshavarzi*, Kanad Mandke, Annabel Macfarlane, Lyla Parvez, Fiona

Gabrielczyk, Angela Wilson, Sheila Flanagan, Usha Goswami

Centre for Neuroscience in Education, Department of Psychology, University of Cambridge,

Cambridge, CB2 3EB, United Kingdom

11
12   **Running head: Decoding speech information in dyslexia**

21   *Corresponding Author: mk919@cam.ac.uk, mahmoud.keshavarzi.ir@ieee.org

MK: Conceptualisation, Methodology, Data analysis, Visualisation, Writing original draft
KM: EEG Methodology, Investigation, Writing – review and editing
AM: EEG Investigation, Behavioural Data Curation
LP: EEG Investigation
FG: Behavioural Investigation, Data Curation
AW: Behavioural Investigation, Data Curation
SF: Behavioural Investigation, Writing - review and editing
UG: Conceptualisation, Funding Acquisition, Methodology, Resources, Supervision, Writing
Original Draft

Abstract

The amplitude envelope of speech carries crucial low-frequency acoustic information that

assists linguistic decoding. The sensory-neural Temporal Sampling (TS) theory of

developmental dyslexia proposes atypical encoding of speech envelope information <10 Hz,

leading to atypical phonological representations. Here a backward linear TRF model and story

listening were employed to estimate the speech information encoded in the

electroencephalogram in the canonical delta, theta and alpha bands by 9-year-old children with

and without dyslexia. TRF decoding accuracy provided an estimate of how faithfully the

children's brains encoded low-frequency envelope information. Between-group analyses

showed that the children with dyslexia exhibited impaired reconstruction of speech information

in the delta band. However, when the quality of speech encoding for each child was estimated

using child-by-child decoding models, then the dyslexic children did not differ from controls.

This suggests that children with dyslexia encode neither "noisy" nor "normal" representations

of the speech signal, but different representations.

Keywords

Developmental dyslexia, EEG, speech envelope, speech decoding, backward linear model

1. Introduction

Dyslexia is a disorder of development. Classically, a child has shown apparently typical language acquisition and cognitive development until faced with the task of learning to read. Suddenly, the affected child shows a specific problem with reading and spelling that cannot be accounted for by low intelligence, poor educational opportunities, or obvious sensory or neurological damage. Reading is the process of understanding speech when it is written down. To acquire early reading skills, the child must learn the visual code used by their culture for representing speech as a series of symbols. Logically, individual differences in acquiring reading could be related developmentally to either spoken language processing or visual code processing or both (Snowling, 2000; Stein and Walsh, 1997). However, it is important to recognise that the visual code that must be learned is not a neutral visual stimulus. It is a culturally-specific code that is taught directly using symbol-sound correspondences, such as the alphabetic correspondences used to represent English phonology (the sound system of the language). Learning this visual code typically begins a few years into the development of a spoken language system, and it is now widely recognised that individual differences in children's pre-reading *phonological awareness* (their ability orally to recognise or manipulate phonological units in words such as syllables, rhymes or phonemes) is a causal determinant of how readily their visual code learning progresses (Ziegler and Goswami, 2005, for cross-language review). Indeed classically, 'deficits' in the representation and use of phonological information have been seen as critical in the etiology of developmental dyslexia (Catts, 1993; Stackhouse and Wells, 1997; Stanovich and Siegel, 1994; Swan and Goswami, 1997; Snowling, 2000).

To understand the nature of this 'phonological deficit' in dyslexia, longitudinal studies that begin with infants are required. Evidence from both electrophysiological and behavioural measures suggests that infants at family risk (genetic risk) for dyslexia already exhibit deficits

in the auditory processing of speech and non-speech sounds in their first year of life, which could be expected to affect novel word learning. It is well known that the speech signal provides a rich repertoire of acoustic cues that infants exploit to learn new words (Saffran, 2001). Although very few neurophysiological longitudinal studies of at-risk infants exist, EEG studies suggest atypical detection of changes to the acoustic features of speech and non-speech sounds (e.g., F0 frequencies, vowel length, and consonant voice onset time) in newborns and two-month-olds, and there are also differences in their hemispherical distribution of the neural responses to these acoustic stimuli (Guttorm, Leppänen, Hämäläinen, Eklund, and Lyytinen, 2010; Guttorm et al., 2005; Leppänen et al., 2010; see also van Zuijen, Plakas, Maassen, Maurits, and van der Leij, 2013). Regarding the atypical development of phonological representations for words that characterise older children once they are diagnosed with dyslexia, the most commonly accepted hypothesis is that these phonological representations are somehow "noisy" or "imprecise" (Swan and Goswami, 1997; Snowling, 2000). As explained in detail by Ramus and Svenkovits (2008), the general idea concerning "noisy" representations is that a dyslexic child's neural representations of the speech signal are somehow fuzzier, for example having a lower resolution than they should, or preserving too many acoustic or allophonic details (Adlard and Hazan, 1998; Elbro et al., 1998; Manis et al., 1997; Mody, Studdert-Kennedy, and Brady, 1997; Serniclaes et al., 2004; Bogliotti et al., 2008; Snowling, 2000). Other terms used in the literature to describe dyslexic children's phonological representations are "imprecise" or "under-specified". Ramus and Svenkovits (2008) disputed this idea about "noisy" representations, arguing that the phonological representations of individuals with dyslexia may be quite normal. The phonological 'deficit' might arise instead from problems in *accessing* these normal representations when required for phonological awareness and other oral tasks. Although access is not mechanistically defined, Ramus and Svenkovits expect access to be more difficult when a task requires lots of phonological

information to be stored in short-term memory, or a task requires speeded or repeated retrievals

of phonological information, or it requires the extraction of speech information from noise

(Ramus and Svenkovits, 2008). To our knowledge, however, neural studies have not yet been

able to contrast the "noisy representations" view of the phonological 'deficit' in dyslexia with

the opposing view of Ramus and Svenkovits that the phonological representations developed

by the dyslexic brain are "normal". We describe such a contrast here.

In the last 5 years, neural studies with children with dyslexia have been applying novel

computational methods to EEG and MEG data to try to measure the quality of children's

phonological representations directly. These methods either use original speech features such

as the envelope to predict the brain's neuroelectric responses (forward models, from speech

envelope to brain response, Di Liberto et al., 2018), or reconstruct speech-based representations

of envelope information from the brain's neuroelectric responses to speech input (backward or

speech reconstruction models, from brain response to speech signal, Power et al., 2016;

Destoky et al., 2020). These studies use a linear approach to stimulus reconstruction or to neural

prediction, namely temporal response functions (TRFs, Di Liberto et al., 2015; Crosse et al.,

2016). Backward TRF models reconstruct either the broadband speech envelope in the original

signal or the envelopes in different frequency bands (such as $0.5 - 1.5$ Hz or $2 - 8$ Hz; Destoky

et al., 2020). Such studies are still rare in the literature, and only one of them to date has used

natural speech (Di Liberto et al., 2018) rather than degraded speech (vocoded speech, Power

et al., 2016) or speech-in-noise (Destoky et al., 2020). Nevertheless, all studies to date show

atypical low-frequency encoding of the speech signal by children with dyslexia, with speech

envelope information <10 Hz encoded reliably less accurately by dyslexic children compared

to age-matched control children.

These prior TRF studies also included younger children matched for reading level to

the children with dyslexia. This is an important experimental control for the effects of learning

on the developing brain. It is known that learning to read in itself changes phonological

representations, as orthographic knowledge gained through reading affects performance in oral

listening tasks for both children and adults (Ehri and Wilce, 1980; Ziegler and Ferrand, 1998).

In principle, younger reading-level-matched (RL match) children provide a control for the

effects of reading experience on the developing language system. Both of the prior EEG studies

using both backward and forward TRF models reported significantly greater stimulus

reconstruction accuracy (Power et al., 2016) or neural response accuracy (Di Liberto et al.,

2018) regarding low-frequency acoustic speech information for their RL controls compared to

their dyslexic participants. Accordingly, in EEG studies the dyslexic brain shows less accurate

speech encoding than the brains of younger children matched for reading experience,

suggesting a fundamental difference in encoding linguistic information. The exception was the

MEG study using speech-in-noise tasks, and the authors explained this null result by suggesting

that discriminating speech in noise may be facilitated by learning to read (Destoky et al., 2020).

If speech-in-noise performance depends on the level of reading attained, then children with

dyslexia should perform at the same level as younger children matched for reading, as found

by Destoky et al. (2020).

To date, therefore, data from neural TRF studies do not suggest that the phonological

representations of words developed by children with dyslexia are "normal" regarding the

encoding of low-frequency acoustic information. At the same time, however, it is not clear that

these representations are "noisy" or "fuzzy". Rather, low-frequency acoustic information in the

speech signal appears to be represented *less accurately* in the neural mental lexicons of children

with dyslexia. This suggests that speech itself may be perceived differently by children with

dyslexia. Rather than being a noisy signal for the dyslexic brain, the signal itself seems to be

represented in an atypical manner, with less accurate representation of low-frequency

information (Goswami, 2022). This may possibly be compensated by an over-weighting of

allophonic and other information (Bogliotti et al., 2008). These differences in the encoding of

acoustic information can be assessed with psychophysical methods (Goswami et al., 2011).

Goswami et al. (2011) presented a phonetic contrast (ba/wa) to children with dyslexia and age-

matched and RL controls, changing "ba" to "wa" either by varying amplitude rise time in

synthetic syllables or by varying frequency rise time. The children with dyslexia were

significantly poorer at discriminating "ba" from "wa" when the phonetic change depended on

amplitude rise time, but were significantly better than both age-matched and RL controls when

the phonetic change depended on frequency rise time. These differences in the encoding of

acoustic information appear to impair dyslexic children's performance in classical

phonological awareness tasks while enhancing their performance in some speech-based tasks

such as categorical perception (Serniclaes et al., 2004; Bogliotti et al., 2008). These

representational differences also complicate the process of learning to read, since any visual

symbol system that is being learned will have been designed for learners who hear speech

differently from children with dyslexia. This would apply whether the visual code is an

alphabetic system or any other orthographic system (see Goswami, 2022).

In the current study, we set out to use a stimulus reconstruction approach to tackle both

theoretical issues ("noisy" representations versus "normal" representations) with the same data.

Our reasoning was as follows. Using a backward TRF modelling approach, thereby using

neurophysiological responses to reconstruct speech envelopes < 10Hz, natural speech listening

data can be used either to compare the decoding accuracy of the models that are reconstructed

between groups of children, or within individual children. Decoding accuracy can be taken as

an estimate of how faithfully the children's brains are encoding low-frequency envelope

information in the speech signal. If children with dyslexia are developing phonological

representations that do not encode low-frequency acoustic information as accurately as control

children, then the between-group comparison should reveal significant differences in the

accuracy of the decoding models. In such a case, the phonological representations developed

by the children with dyslexia could not be designated as "normal". If children with dyslexia

are encoding "noisy" or "fuzzy" representations of the speech signal, then the within-child

comparison should also reveal significant differences in the accuracy of the models at the group

level. The averaged within-child models for the children with dyslexia should be less accurate

than the averaged within-child models for the control children, as the speech-based

representations developed by children with dyslexia should be less consistent. To investigate

these possibilities, EEG was recorded while children listened to a 10-minute story presented as

audio-only. Backward TRF models were then computed, and the accuracy of these models was

compared for the canonical delta, theta and alpha (control) bands using both between-group

and within-child comparisons. For the between-group comparisons, we expected to replicate

the finding in the literature that low-frequency acoustic speech information is encoded less

accurately by children with dyslexia in the delta and theta bands. Regarding the "noisy"

representations question, given the frequent observation that children with dyslexia appear to

show no difficulties in speaking and listening tasks that do not tax phonological knowledge, it

may be that the children are operating with perceptually stable phonological representations

that are not "noisy" for their users.

2. Material and Methods

2.1. Participants

Fifty-one children were participated in this study. Twenty-one participants were

typically developing children (mean age of 109.3 ± 5.4 months) and thirty participants had

developmental dyslexia (mean age of 110.7 ± 5.6 months). The unequal group sizes arose due

to Covid-19, which necessitated the cessation of testing part-way through the study, thereby

also preventing testing of the recruited RL control group. One of the dyslexic children only

completed 5 minutes of the EEG session, due to lots of head movement, not listening to the story, and taking out the earphone phone during data collection. We therefore excluded this child from further analysis. The children with dyslexia did not have any additional learning difficulties (e.g., ADHD, dyspraxia, autistic spectrum disorder, developmental language disorder), and were recruited through learning support teachers. The absence of the additional learning difficulties was confirmed based on school and parental reports and our own behavioural testing. All participants had a nonverbal IQ above 84, and their first language spoken at home was English. All children received a short hearing screen across frequency range of 0.25 – 8 kHz (0.25, 0.5, 1, 2, 4, 8 kHz) using an audiometer, and they all were found to be sensitive to sounds within the 20 dB HL range. SES data were not formally collected, but children were attending state schools (equivalent to US public schools) situated in a range of towns and villages near a university town in the United Kingdom. Participants and their parents gave informed consent for the EEG study in accordance with the Declaration of Helsinki, and the study was approved by the Psychology Research Ethics Committee of the University of Cambridge.

2.2. Behavioural tests

A series of standardised tests (see Table 1) of written and spoken language were administered in schools prior to the EEG session to assess cognitive development (please see Keshavarzi et al., 2022, for a full description of each test; the current participants also received the repetitive speech task employed in that study). Reading and spelling were assessed using the British Ability Scales (BAS, Elliot et al., 1996) and the Test of Word Reading Efficiency (TOWRE, Torgesen et al., 1999). Four subscales from the Wechsler Intelligence Scale for Children (WISC-V, Weschler, 2016) including two verbal (vocabulary and similarities) and two non-verbal (block design and matrix reasoning) scales were administered. Full-scale IQ

was then estimated following the approach of Aubry and Bourdin (2018). In addition, the British Picture Vocabulary Scale (BPVS3) measure was used to assess receptive vocabulary and the Phonological Assessment Battery (PhAB, Frederickson et al., 1997; GL Assessment) was administered to assess phonological awareness at the rhyme and phoneme levels, along with rapid naming of objects and digits (RAN). The children's amplitude rise time thresholds were estimated using 3 psychoacoustic threshold tasks based on (1) sine tones; (2) tones made from speech-shaped noise; (3) a synthetic syllable "ba". As shown in Table 1, the children with dyslexia had significantly poorer reading and spelling skills than the control children, significantly poorer phonological skills, and significantly poorer amplitude rise time discrimination in a synthetic syllable task (discriminating differences in rise time of the syllable "ba").

Table 1. Group characteristics (N=51) expressed as mean and (S.D.) for children with dyslexia and age-matched control children.

	Dyslexic	Age-Matched Control
N	30	21
Age (months)	110.7 (5.6)	109.3 (5.4)
WISC FSIQ	101.7 (10.1)	103.6 (12.0)
BAS Reading SS	81.0 (8.0) ***	99.5 (6.2)
BAS Reading Age in months	85.5 (11.0) ***	106.5 (11.7)
BAS Spelling SS	79.9 (7.5) ***	97.1 (6.1)
TOWRE SWE SS	79.5 (12.8) ***	101.1 (7.7)
TOWRE PDE SS	79.2 (10.9) ***	98.0 (8.6)
PhAB Rhyme SS	92.6 (11.7) ***	102.4 (5.9)
PhAB Phoneme SS	97.6 (9.8) **	105.1 (9.4)
PhAB RAN objects SS	91.9 (15.0) *	101.2 (13.0)
PhAB RAN digits SS	85.8 (15.8) **	97.4 (13.3)
Rise time sine tone ms	174.7 (81.7)	139.7 (84.5)
Rise time SSN ms	221.9 (56.7)	215.0 (54.9)
Rise time "ba" ms	101.5 (43.8)*	70.8 (34.6)

Note. ***$p < .001$; **$p < .01$; *$p < .05$; +$p = .052$. WISC FSIQ = Wechsler Intelligence Scale for Children, estimated Full Scale IQ; BPVS = British Picture Vocabulary Scales, SS = standardised mean score = 100; BAS = British Ability Scales, SS = standardized score = 100; TOWRE SWE = Test of Word Reading Efficiency Sight Word Efficiency Scale, SS = standardized score = 100; TOWRE PDE = Phonic Decoding Efficiency Scale, SS = standardized score = 100; PhAB = Phonological Awareness Battery, SS = standardised mean score = 100; RAN = Rapid Automatized Naming; Rise time = threshold in rise time task in ms; SSN = speech-shaped noise.

260 *2.2. Experimental set-up and stimuli*

261   The children were seated in an electrically shielded soundproof room. The auditory

stimuli (through earphones) were presented at a sampling rate of 44.1 kHz to the participant

while EEG data were collected at a sampling rate of 1 kHz using a 128-channel EEG system

(HydroCel Geodesic Sensor Net). The stimuli were natural speech presented as a 10-minute

long story for children, *The Iron Man: A Children's Story in Five Nights* by Ted Hughes. The

story was read in child-directed speech and was presented in ten sections, each of which lasted

about one minute followed by a 2 second gap. During the experiment, participants were

instructed to listen to the speech carefully and to look at a red cross (+) shown on the screen

that was in front of them.

2.3. Auditory Stimuli and EEG Data Pre-processing

272   In this study, we first obtained the broad-band envelopes by calculating the absolute

273 value of the analytical signal of the speech stimuli. The broad-band envelopes were then filtered

274 at frequency range of 0.5 – 8 Hz to extract the low frequency envelopes. The low frequency

275 envelopes were used for the analyses used here.

276   The collected EEG data were referenced to Cz channel and then band-passed filtered

into frequency range of 0.5 – 48 Hz using a zero phase FIR filter with low cutoff (−6 dB) of

0.25 Hz and high cutoff (−6 dB) of 48.25 Hz (EEGLab Toolbox; Delorme and Makeig, 2004).

Bad channels were detected and interpolated through spherical interpolation (EEGLab

Toolbox). The EEG data were downsampled to 100 Hz and filtered to extract delta (0.5 – 4

Hz), theta (4 – 8 Hz) and alpha (8 – 12 Hz) frequency bands. The data were further

downsampled to 50 Hz to reduce the computational costs, and then epoched into the 1-minute

story parts.

2.4. Stimulus Reconstruction Using Backward TRF Model

To investigate how accurately the children were encoding the speech signal, we used a

decoding accuracy method based on the backward TRF model (Crosse et al., 2016). The TRF

model was used to reconstruct the stimuli envelopes from the EEG signals at each frequency

band of interest (delta, theta, alpha). The backward TRF model estimates the stimulus

envelope, *s*(*t*), through the following equation:

$s(t) = g(t,k) * r(t,k) = \sum_{k=1}^{K} \sum_{\tau=-T_{min}}^{T_{max}} g(\tau,k) r(t+\tau,k)$ (1)

where $g(\tau,k)$ is a decoder representing the linear mapping from the neural response to the

stimulus envelope for time lag τ and channel k, parameters $T_{min} = 0\ ms$ and $T_{max} = 500\ ms$

determine the minimum and maximum time lags, respectively, and $r(t,k)$ represents the neural

response at channel k. Accordingly, the backward model can estimate the quality of the child's

encoding of the original speech signal, as the accuracy of decoding estimated by the model

provides a measure of the quality of the neural representation of the low-frequency speech

information in each frequency band. The validation procedure used to regularize the backward

linear model was the "leave-one-out" cross-validation (using *mTRFcrossval* function from the

mTRF Toobox) in which each trial is "left out" or used for testing and the remainder are used

to train the model and this procedure is repeated across all trials. The Pearson correlation

between the estimated envelope and the actual envelope was used to estimate the performance

of the model, with a higher correlation value indicating more accurate decoding by the model.

2.5. Between-Group Backward TRF Analysis

To model normative decoding of the speech signal from the EEG signal at each

frequency band, we used EEG recorded from the typically-developing (control) children. For

each frequency band of interest (delta, theta, alpha), we used a randomly-selected subset of

control children (C_{train} = 11) for training 11 backward TRF models. A final model was obtained

by averaging across all 11 individual trained models. This final averaged model was then tested

separately on the rest of the control children (C_{test} = 10) and on the 29 dyslexic children. We

next computed the mean correlation values for the tested control children and the dyslexic

children, resulting in a single value for each group (control, dyslexic). This procedure was then

repeated 100 times, using random permutations across all the control children, in order to use

different combinations of 11 versus 10 (C_{train} and C_{test}) control children for training and testing

the model, resulting in 100 normative (averaged) models. Accordingly, 100 correlation values

were generated for the control children and 100 correlation values were generated for the

dyslexic children, providing an averaged correlation value for each group. This then enabled

us to assess whether there were differences in speech decoding accuracy between children with

dyslexia and age-matched controls. In essence, the reconstruction accuracy for the typically-

developing children was utilised as a benchmark to compare with the reconstruction accuracy

obtained for the dyslexic children.

2.6. Within-Child Backward TRF Analysis

To discover whether the perceptual experience of the speech signal is consistent (or

stable) for any individual child, we also built a single backward TRF model for each child in

the study based on the data from that child. This within-child approach enabled us to assess

whether the speech-based representations developed by children with dyslexia are less

consistent, "fuzzy" or "noisy" for each frequency band of interest. This was achieved by using

80% of the data (eight story sections) for each child to train the TRF model for that child, and

then using the remaining data from that child (two story sections) to test the model. We

calculated the Pearson correlation between the estimated speech envelope and the actual speech

envelope for each child separately for each frequency band. This resulted in a single correlation

score for each band for each child. These scores were then summed by group for each band

and compared between groups.

2.7. Computation of the Chance Level for Backward TRF

To check the statistical significance of stimulus reconstruction accuracy as estimated

by the backward TRF models (both between-group and within-child), we computed null

models for each frequency band of interest. To obtain the null models, we randomly selected

ten control children and EEG data were permuted across different story sections for each child.

We then built a single model for each child. To achieve this, 80% of the data for each child

were used to train the null model for that child, and the remaining data were used to test the

null model. The Pearson correlation between the estimated speech envelope and the actual

speech envelope for each child separately for each frequency band was then calculated for these

null models, resulting in a single correlation value for each band and for each child. We finally

computed the mean reconstruction accuracy by averaging across the accuracy scores obtained

for the selected children. This was done 100 times for each band, in order to calculate the

probability density functions (PDF, a statistical measure that determines the probability

distribution of a variable) for the null models and thereby establish chance levels. Note that the

area under the PDF curve should be 1, hence for our data the smaller the range of Pearson

correlations on the *x* axis, the higher the value required for the PDF on the *y* axis.

3. Results

3.1. Does the Accuracy of Speech Envelope Decoding Vary Between Control and Dyslexic

Children?

To investigate whether the accuracy of speech envelope decoding estimated from EEG

recorded from the dyslexic brain is different from that recorded from the typically-developing

brain, the Between-Group Backward TRF analysis (see Section 2.5) was applied separately for

each frequency band. To check the statistical significance of the stimulus reconstruction

accuracy obtained for each frequency band by group, we first compared decoding accuracy in

each band to decoding accuracy for the null models (the estimate of chance level for each band,

see Section 2.7). Statistical significance by band and group is shown in Figure 1. The

modelling showed that stimulus-reconstruction accuracy was significantly greater than chance

in the delta band only (Figure 1A). Reconstruction accuracy in the theta and alpha bands was

not statistically different from the noise values in these bands for either group (Figure 1B,C).

Figure 1. Statistical significance of reconstruction accuracy for the delta band (Panel A),
theta band (Panel B) and alpha band (Panel C) for the Between-Group Analyses (Blue,
control children; Red, dyslexic children). The probability density functions of the null models
are shown by the black curves, please note that the scales for the three bands are different. The dashed
black lines denote the correlation values corresponding to statistical significance ($p = 0.05$). The blue
and red dashed lines show the mean correlation values obtained for the control and dyslexic groups,
respectively.

To compare the accuracy of decoding between the control and dyslexic groups in the

delta band, we applied the Wilcoxon rank sum test. This showed that reconstruction accuracy

for the control group in the delta band as derived from 100 TRF models was significantly

greater than reconstruction accuracy for the dyslexic group (Wilcoxon rank sum test, $z = 3.72$,

$p = 0.0002$). This difference in speech envelope decoding suggests that control brains encode

a more accurate neural representation of the speech envelope in the delta band compared to

dyslexic brains, as shown in Figure 2.

Figure 2. Stimulus-reconstruction accuracy for control versus dyslexic children in the
delta band summarising correlation values from 100 backward TRF models. The black
circles show the average accuracy scores. On each box, the central line indicates the median, and the
bottom and top edges of the box denote the 25th and 75th percentiles, respectively. The whiskers also
extend to the most extreme data points. *** denotes $p < 0.001$.
391

3.2. Do Children with Dyslexia Develop "Noisy" Representations of Speech Stimuli?

To explore whether children with dyslexia have "noisy" or "fuzzy" representations of

the speech signal, the Within-Child Backward TRF analysis method (see Section 2.6) was

employed separately for each frequency band. The statistical significance of the stimulus

reconstruction accuracy obtained for each frequency band for the within-child analyses were

computed by considering chance level for that band (the null models, see Section 2.7).

Comparisons with the null models showed that stimulus-reconstruction accuracy was above

chance-levels (alpha = 0.05) for both groups in all three frequency bands (Figure 3).

Figure 3. Statistical significance of within-child stimulus reconstruction accuracy for the delta
band (Panel A), theta band (Panel B) and alpha band (Panel C) summarised by group (Blue,
control children; Red, dyslexic children). The probability density functions of the null models are
again shown by the black curves. The dashed black lines denote the correlation values corresponding
to statistical significance ($p = 0.05$). The blue and red dashed lines show the mean correlation values
obtained from the within-child analysis for control and dyslexic groups, respectively.
408

Figure 4. Stimulus-reconstruction accuracy computed for each individual control and dyslexic child in the delta (Panel A), theta (Panel B), and alpha (Panel C) bands in the within-child analysis. The grey disks denote individual children's reconstruction accuracy scores, and the black circles show the average accuracy scores for each band and each group.

Figure 4 shows the box plots for stimulus reconstruction accuracies for the control and dyslexic groups based on the backward TRF models from individual children (shown as dots) by band, separated by group. To compare reconstruction accuracy between the two groups, we employed the Wilcoxon rank sum test. The results by band showed that there was no difference between the two groups in any band (delta band, Figure 4A; Wilcoxon rank sum test, $z = -0.59$, $p = 0.55$; theta band, Figure 4B; Wilcoxon rank sum test, $z = -1.30$, $p = 0.19$; alpha band, Figure 4C; Wilcoxon rank sum test, $z = -0.81$, $p = 0.42$). Accordingly, as shown in Figure 4, the backward TRF model can reconstruct the stimulus envelopes for each individual child in each band consistently from that child's EEG data. This contrasts with the reconstruction consistency achieved for the between-group analysis, where decoding accuracy was only above chance for both groups in the delta band condition (Figure 1). The within-child group

comparisons (Figure 4) show that the consistency of decoding accuracy is similar irrespective

of whether the child is a child with dyslexia or a control child, indicating that the neural

representation of the speech signal for each child is consistent for that child and provides a

stable percept of speech.

4. Discussion

Here we employed backward TRF models to estimate stimulus reconstruction accuracy

of bands of low-frequency acoustic information in the speech signal (delta- and theta-band

speech envelope information) along with alpha-band speech envelope information (control

band) in children with and without dyslexia matched for age. Two different group comparison

approaches (between-group and within-child) were employed to investigate whether the

speech-based representations being developed by the dyslexic brain were "noisy" or "normal".

If decoding accuracy varies by group for the between-group TRF models, then the speech-

based representations developed by children with developmental dyslexia cannot be considered

"normal". However, if decoding accuracy is equal irrespective of group for within-child TRF

models, then the speech-based representations developed by children with developmental

dyslexia cannot be considered "noisy". The modelling suggested that the dyslexic brain does

not develop "noisy" representations, but neither are these representations "normal". The

between-group models showed that the speech-based representations developed by children

with dyslexia are *less accurate* than the speech-based representations developed by children

without dyslexia regarding low frequency envelope information. These differences were

restricted in the current study to the delta band (Figure 2). The between-group modelling

generated lower decoding accuracies than the within-child modelling, as can be seen by

comparing the correlation values in Figures 2 and 4. Between-group decoding accuracy for the

theta and alpha bands did not exceed chance levels as estimated by the null models. Therefore,

the conservative conclusion is that encoding of low-frequency speech information in the delta

band is atypical in 9-year-old children with developmental dyslexia.

At the level of an individual child, however, the modelling did not support the classical

theoretical view (including our own earlier view, see Swan & Goswami, 1997) that the

phonological representations of speech developed by children with dyslexia are "fuzzy",

"noisy" or "imprecise" (see also Elbro et al., 1998; Snowling, 2000). The quality of the speech-

based representations for each child as indexed by the neural decoding method were of similar

consistency whether the child was dyslexic or not, for all frequency bands explored. This

suggests that even if acoustic information is weighted differently in the speech-based

representations developed by children with dyslexia, the experience of speech processing is

perceptually stable for the children themselves. Nevertheless, a linear decoding approach to

stimulus reconstruction is only one method for assessing whether neural stimulus

representations are "noisy" or not, and converging neural methods are required.

The delta-band between-group difference in decoding accuracy found here was

predicted by TS theory (Goswami, 2011, 2019), which provided the conceptual framework for

the current study. TS theory has proposed that dyslexic difficulties in discriminating amplitude

envelope rise times, difficulties found for dyslexic children in 7 languages to date (see

Goswami, 2015, for a review), are associated with atypical neural encoding of acoustic

information within the amplitude envelope < 10Hz, related to perceptual impairments in

processing speech rhythm. Rise times are one neural trigger for oscillatory phase-resetting, an

automatic process that aids multi-timescale speech-brain alignment in adults (Giraud &

Poeppel, 2012; Doelling et al., 2014; Lizarazu et al., 2021). A recent MEG study with adults

with dyslexia has shown that rise times do not provide an efficient phase-resetting mechanism

in the dyslexic brain during natural speech listening (Lizarazu et al., 2021). The between-group

data presented here suggest that this inefficient phase-resetting may particularly affect the

encoding of speech envelope information in the delta band during childhood.

In conclusion, the current study converges with the three preceding studies of children

with dyslexia that measured speech-based representations directly using neural methods

(Power et al., 2016; Di Liberto et al., 2018; Destoky et al., 2020). These studies all reported

that children with dyslexia show atypical encoding of low-frequency information in the speech

amplitude envelope. Together with the current study, this suggests that the speech-based

representations developed by children with dyslexia are not "normal", but rather encode less

accurate prosodic (delta band) speech-based information in comparison to the representations

developed by typically-developing children. TS theory proposes that these differences affect

phonological awareness at all linguistic levels, which then impacts the process of learning to

read and spell. At the same time, the current study contributes a novel child-by-child neural

decoding approach that reveals that the perceptual world of the dyslexic child regarding speech

is a consistent one. At the level of the individual child, the speech-based representations

developed by children with dyslexia are not "fuzzy" or "noisy". This finding contributes to a

long-standing debate in the psycholinguistic literature regarding whether the phonological

'deficit' that characterises individuals with dyslexia across languages stems from differences

in phonological representation or in phonological access (e.g., Adlard and Hazan, 1998; Elbro

et al., 1998; Manis et al., 1997; Mody, Studdert-Kennedy, and Brady, 1997; Serniclaes et al.,

2004; Bogliotti et al., 2008; Snowling, 2000; Ramus & Svenkovits, 2008). The neural decoding

method employed here suggests that the speech-based representations developed by children

with dyslexia are neither "noisy" nor "normal", which accords well with the linguistic data

gathered by recent longitudinal studies of infants at family risk for developmental dyslexia

(Kalashnikov et al., 2018, 2019a,b, 2020). Nevertheless, the strongest test of this conclusion

would be a demonstration that speech production is also atypical in dyslexia regarding the

specification of speech prosody. If speech prosody is not encoded accurately in the dyslexic

brain, yet this atypical encoding is consistent and perceptually stable for the child, then speech

output of prosodic information should also be affected.

Acknowledgment

The research was funded by a grant awarded to UG by the Foundation Botnar (project number:

6064) and a donation from the Yidan Prize Foundation. The sponsors played no role in the

study design, data interpretation or writing of the report. The authors would like to thank all

the families and schools involved in the study.

References

Adlard, A., & Hazan, V. (1998). Speech perception in children with specific reading difficulties
(dyslexia). *The Quarterly Journal of Experimental Psychology Section A*, 51(1), 153-177.
https://doi.org/10.1080/713755750.

Aubry, A., & Bourdin, B. (2018). Short Forms of Wechsler scales assessing the intellectually
gifted children using simulation data. *Frontiers in psychology*, 9, 830.
https://doi.org/10.3389/fpsyg.2018.00830.

Bogliotti, C., Serniclaes, W., Messaoud-Galusi, S., & Sprenger-Charolles, L. (2008).
Discrimination of speech sounds by children with dyslexia: Comparisons with chronological
age and reading level controls. *Journal of experimental child psychology*, 101(2), 137-155.
https://doi.org/10.1016/j.jecp.2008.03.006.

Catts, H. W. (1993). The relationship between speech-language impairments and reading
disabilities. *Journal of Speech, Language, and Hearing Research*, 36(5), 948-958.
https://doi.org/10.1044/jshr.3605.948.

Crosse, M. J., Di Liberto, G. M., Bednar, A., & Lalor, E. C. (2016). The multivariate temporal
response function (mTRF) toolbox: a MATLAB toolbox for relating neural signals to
continuous stimuli. *Frontiers in human neuroscience*, 10, 604.
https://doi.org/10.3389/fnhum.2016.00604.

Delorme, A., & Makeig, S. (2004). EEGLAB: an open-source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of neuroscience methods*, 134, 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009.

Destoky, F., Bertels, J., Niesen, M., Wens, V., Vander Ghinst, M., Leybaert, J., Lallier, M., Ince, R. A., Gross, J., De Tiège, X., & Bourguignon, M. (2020). Cortical tracking of speech in noise accounts for reading strategies in children. *PLoS biology*, 18(8), p.e3000840. https://doi.org/10.1371/journal.pbio.3000840.

Di Liberto, G. M., O'Sullivan, J. A., & Lalor, E. C. (2015). Low-frequency cortical entrainment to speech reflects phoneme-level processing. *Current Biology*, 25(19), 2457-2465. https://doi.org/10.1016/j.cub.2015.08.030.

Di Liberto, G. M., Peter, V., Kalashnikova, M., Goswami, U., Burnham, D., & Lalor, E. C. (2018). Atypical cortical entrainment to speech in the right hemisphere underpins phonemic deficits in dyslexia. *NeuroImage*, 175, 70-79. https://doi.org/10.1016/j.neuroimage.2018.03.072.

Doelling, K. B., Arnal, L. H., Ghitza, O., & Poeppel, D. (2014). Acoustic landmarks drive delta-theta oscillations to enable speech comprehension by facilitating perceptual parsing. *NeuroImage*, 85, 761–768. https://doi.org/10.1016/j.neuroimage.2013.06.035.

Elbro, C., Borstrøm, I., & Petersen, D. K. (1998). Predicting dyslexia from kindergarten: The importance of distinctness of phonological representations of lexical items. Reading research quarterly, 33(1), 36-60. https://doi.org/10.1598/RRQ.33.1.3.

Elliott, C. D., Smith, P., & McCullogh, K. (1996). British Ability Scales (2nd ed.). Windsor, UK: NFER-Nelson.

Ehri, L. C., & Wilce, L. S. (1980). The influence of orthography on readers' conceptualization of the phonemic structure of words. Applied Psycholinguistics, 1(4), 371-385. https://doi.org/10.1017/S0142716400009802.

Frederickson, N., Frith, U., & Reason, R. (1997). Phonological Assessment Battery: Standardised Edition. NFER-Nelson.

Giraud, A.L., & Poeppel, D. (2012). Cortical oscillations and speech processing: Emerging computational principles and operations. *Nature Neuroscience*, 15(4), 511–517. https://doi.org/10.1038/nn.3063.

Goswami, U. (2011). A temporal sampling framework for developmental dyslexia. *Trends in Cognitive Sciences*, 15(1), 3–10. https://doi.org/10.1016/j.tics.2010.10.001.

Goswami, U. (2015). Sensory theories of developmental dyslexia: Three challenges for research. *Nature Reviews Neuroscience*, 16(1), 43–54. https://doi.org/10.1038/nrn3836.

Goswami, U. (2019). Speech rhythm and language acquisition: An amplitude modulation phase hierarchy perspective. *Annals of the New York Academy of Sciences*, e14137. https://doi.org/10.1111/nyas.14137.

Goswami, U., 2022. Theories of Developmental Dyslexia. To appear in M. Skeide (Ed.), The Cambridge Handbook of Dyslexia and Dyscalculia. Cambridge, UK: Cambridge University Press.

Goswami, U., Fosker, T., Huss, M., Mead, N., & Szücs, D. (2011). Rise Time and Formant Transition Duration in the Discrimination of Speech Sounds: The Ba-Wa Distinction in Developmental Dyslexia. *Developmental Science*, 14, 34-43. https://doi.org/10.1111/j.1467-7687.2010.00955.x.

Guttorm, T. K., Leppänen, P. H., Poikkeus, A. M., Eklund, K. M., Lyytinen, P., & Lyytinen, H. (2005). Brain event-related potentials (ERPs) measured at birth predict later language development in children with and without familial risk for dyslexia. *Cortex*, 41(3), 291-303. https://doi.org/10.1016/S0010-9452(08)70267-3.

Guttorm, T. K., Leppänen, P. H., Hämäläinen, J. A., Eklund, K. M., & Lyytinen, H. J. (2010). Newborn event-related potentials predict poorer pre-reading skills in children at risk for dyslexia. *Journal of learning disabilities*, 43(5), 391-401. https://doi.org/10.1177/0022219409345005.

Kalashnikova, M., Goswami, U., & Burnham, D. (2018). Mothers speak differently to infants at-risk for dyslexia. *Developmental Science*, 21(1), 1–15. https://doi.org/10.1111/desc.12487.

Kalashnikova, M., Goswami, U., & Burnham, D. (2019a). Delayed development of phonological constancy in toddlers at family risk for dyslexia. *Infant Behavior and Development*, 57, 101327. https://doi.org/10.1016/j.infbeh.2019.101327.

Kalashnikova, M., Goswami, U., & Burnham, D. (2019b). Sensitivity to amplitude envelope rise time in infancy and vocabulary development at 3 years: A significant relationship. *Developmental Science*, 22(6), 1–9. https://doi.org/10.1111/desc.12836.

Kalashnikova, M., Goswami, U., & Burnham, D. (2020). Novel word learning deficits in infants at family risk for dyslexia. *Dyslexia*, e1649. https://doi.org/10.1002/dys.1649.

Keshavarzi, M., Mandke, K., Macfarlane, A., Parvez, L., Gabrielczyk F., Wilson, A., & Goswami, U. (2022). Atypical Delta-band Phase Consistency and Atypical Preferred Phase in Children with Dyslexia during Neural Entrainment to Rhythmic Audio-Visual Speech. http://dx.doi.org/10.2139/ssrn.3982171.

Leppänen, P. H., Hämäläinen, J. A., Salminen, H. K., Eklund, K. M., Guttorm, T. K., Lohvansuu, K., Puolakanaho, A., & Lyytinen, H. (2010). Newborn brain event-related potentials revealing atypical processing of sound frequency and the subsequent association with later literacy skills in children with familial dyslexia. *Cortex*, 46(10), 1362-1376. https://doi.org/10.1016/j.cortex.2010.06.003.

Lizarazu, M., Lallier, M., Bourguignon, M., Carreiras, M., & Molinaro, N. (2021). Impaired neural response to speech edges in dyslexia. *Cortex*, 135, 207–218. https://doi.org/10.1016/j.cortex.2020.09.033.

Manis, F. R., McBride-Chang, C., Seidenberg, M. S., Keating, P., Doi, L. M., Munson, B., & Petersen, A. (1997). Are speech perception deficits associated with developmental dyslexia?. *Journal of experimental child psychology*, 66(2), 211-235. https://doi.org/10.1006/jecp.1997.2383.

Mody, M., Studdert-Kennedy, M., & Brady, S. (1997). Speech perception deficits in poor readers: Auditory processing or phonological coding?. *Journal of experimental child psychology*, 64(2), 199-231. https://doi.org/10.1006/jecp.1996.2343.

Power, A. J., Colling, L. J., Mead, N., Barnes, L., & Goswami, U. (2016). Neural encoding of the speech envelope by children with developmental dyslexia. *Brain and Language*, 160, 1-10. https://doi.org/10.1016/j.bandl.2016.06.006.

Power, A. J., Mead, N., Barnes, L., & Goswami, U. (2012). Neural entrainment to rhythmically presented auditory, visual, and audio-visual speech in children. *Frontiers in Psychology*, 3, 216. https://doi.org/10.3389/fpsyg.2012.00216.

Ramus, F., & Szenkovits, G. (2008). What phonological deficit?. *Quarterly journal of experimental psychology*, 61(1), 129-141. https://doi.org/10.1080/17470210701508822.

Saffran, J.R. (2001). Words in a sea of sounds: The output of infant statistical learning. *Cognition*, 81(2), 149-169. https://doi.org/10.1016/S0010-0277(01)00132-9.

Serniclaes, W., Van Heghe, S., Mousty, P., Carré, R., & Sprenger-Charolles, L. (2004). Allophonic mode of speech perception in dyslexia. *Journal of experimental child psychology*, 87(4), 336-361. https://doi.org/10.1016/j.jecp.2004.02.001.

Snowling M.J. (2000). Dyslexia. Blackwell Publishers, Oxford.

Stackhouse, J., & Wells, B. (1997). Children's Speech and Literacy Difficulties, Book1: A Psycholinguistic Framework, 9. Wiley-Blackwell.

663 Stanovich, K. E., & Siegel, L. S. (1994). Phenotypic performance profile of children with
664 reading disabilities: A regression-based test of the phonological-core variable-difference
665 model. *Journal of educational psychology*, 86(1), 24. https://doi.org/10.1037/0022-
666 0663.86.1.24.
667
668 Stein, J., & Walsh, V. (1997). To see but not to read; the magnocellular theory of dyslexia.
669 *Trends in neurosciences*, 20(4), 147-152. https://doi.org/10.1016/S0166-2236(96)01005-3.
670
671 Swan, D., & Goswami, U. (1997). Phonological awareness deficits in developmental dyslexia
672 and the phonological representations hypothesis. *Journal of experimental child psychology*,
673 66(1), 18-41. https://doi.org/10.1006/jecp.1997.2375.
674
675 Torgesen, J., Wagner, R. K., & Rashotte, C. (1999). Test of Word Reading Efficiency
676 (TOWRE). Austin, TX: Pro-Ed.
677
678 van Zuijen, T. L., Plakas, A., Maassen, B. A., Maurits, N. M., & van der Leij, A. (2013). Infant
679 ERPs separate children at risk of dyslexia who become good readers from those who become
680 poor readers. *Developmental science*, 16(4), 554-563. https://doi.org/10.1111/desc.12049.
681
682 Wechsler, D. (2016). Wechsler Intelligence Scale for Children, 5th Edn. London, UK: Pearson
683 Assessment.
684
685 Ziegler, J. C., & Ferrand, L. (1998). Orthography shapes the perception of speech: The
686 consistency effect in auditory word recognition. *Psychonomic Bulletin & Review*, 5(4), 683-
687 689. https://doi.org/10.3758/BF03208845.
688
689 Ziegler, J. C., & Goswami, U. (2005). Reading acquisition, developmental dyslexia, and skilled
690 reading across languages: a psycholinguistic grain size theory. *Psychological bulletin*, 131, 3.
691 https://doi.org/10.1037/0033-2909.131.1.3.