Two Competing Guilds as a Core Microbiome Signature for Health Recovery

Guojun Wu[2,3]*, Ting Xu[1]*, Naisi Zhao[6]*, Yan Y. Lam[7]*, Xiaoying Ding[5]*, Dongqin Wei[4], Jian Fan[4], Yajuan Shi[4], Xiaofeng Li[4], Mi Li[4], Shenjie Ji[4], Xuejiao Wang[5], Huaqing Fu[1], Feng Zhang[8], Yu Shi[4]†, Chenhong Zhang[1]†, Yongde Peng[5]†, Liping Zhao[1,2,3]†

[1]State Key Laboratory of Microbial Metabolism and Ministry of Education Key Laboratory of Systems Biomedicine, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai 200240, China

[2]Department of Biochemistry and Microbiology, School of Environmental and Biological Sciences and Center for Microbiome, Nutrition, and Health, New Jersey Institute for Food, Nutrition, and Health, Rutgers, The State University of New Jersey, New Brunswick, NJ 08901, USA

[3]Rutgers-Jiaotong Joint Laboratory for Microbiome and Human Health, New Brunswick, NJ, USA

[4]Department of Endocrinology and Metabolism, Qidong People's Hospital, Jiangsu 226200, China

[5]Department of Endocrinology and Metabolism, Shanghai General Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200080, China

[6]Department of Public Health and Community Medicine, School of Medicine, Tufts University, Boston, MA, 02111, USA

[7]Gut Microbiota and Metabolism Group, Centre for Chinese Herbal Medicine Drug Development, School of Chinese Medicine, Hong Kong Baptist University, Units 201-207, Building 15W, 15 Science Park West Avenue, Pak Shek Kok, N.T., Hong Kong, China

[8]Nutrition Department (Clinical Study Center of Functional Food), The Affiliated Hospital of Jiangnan University Wuxi, Jiangsu 214122, China

*co-first author

†co-corresponding author

Liping Zhao, liping.zhao@rutgers.edu; Yongde Peng, pengyongde0908@126.com ; Chenhong Zhang zhangchenhong@sjtu.edu.cn; Yu Shi, 2383994147@qq.com

33
34
Summary Paragraph
36

Over eons of co-evolution, the gut microbiota has become an essential organ for humans[1,2].

However, it is unclear what core members and their ecological organization ensures the stable

provision of this organ's essential health-relevant functions to the host. With high quality

metagenome-assembled genomes as network nodes, here we identified two competing guilds[3] of

the most stably and highly connected bacteria that together correlate with a wide range of host

health conditions. Genomes in these two guilds kept their ecological relationship unchanged

despite experiencing profound abundance changes during a 3-month high fiber intervention and

1-year follow-up in patients with type 2 diabetes (T2DM). The genomes of one guild harbored

more genes for plant polysaccharide degradation and butyrate production, while the other guild

had more genes for virulence or antibiotic resistance. A Random Forest regression model showed

that the abundance distributions of these genomes were associated with 41 out of 43 bio-clinical

parameters in the study cohort. With these genomes as reference, Random Forest modeling

successfully classified case and control of T2DM, atherosclerotic cardiovascular disease, liver

cirrhosis, inflammatory bowel diseases, colorectal cancer, ankylosing spondylitis, schizophrenia,

and Parkinson's disease in 12 independent metagenomic datasets from 1,816 participants across

ethnicity and geography. This core microbiome signature may serve as a common target for

health recovery.

Introduction

The gut microbiome supports the host's homeostasis in metabolism, immunity,

development, and behavior, etc.[4] It has been regarded as an essential organ because the

attenuation or loss of health-relevant functions of a dysbiotic gut microbiome has been linked

with the initiation and/or progression of many chronic diseases, including type 2 diabetes

(T2DM)[5-7]. However, the core members of the gut microbiota and their underlying ecological

structure remain to be identified.

The gut microbiota is a complex adaptive system[8], in which the minimum responding units

to environmental perturbations are bacterial genomes[9]. More importantly, genomes are not

independent microbiome features. They form ecological interactions, such as competition or

cooperation, with each other and organize themselves into a higher-level structure called

"guilds"[3]. Each guild is a potential functional group of bacteria in the gut ecosystem. Guild

members may have widely diverse taxonomic backgrounds but thrive or decline together and

thus show co-abundant behavior. Guild-level variations have been positively or negatively

correlated with disease phenotypes and their members have been demonstrated as having causal

role in host disease phenotypes[10,11]. Although a suite of microbiome-wide association studies

(MWAS) has attempted to identify the microbiome signatures (using features such as genes,

pathways, taxa, etc.) that are associated with disease phenotypes [12-15], genomes and their guild-

level organization have not been extensively employed to describe the ecological structure that

supports the gut microbiota to stably provide health-relevant functions to the host. To this end,

we adopt a genome-centric approach which is based on high-quality draft genomes assembled

directly from metagenomic datasets (high-quality metagenome-assembled genomes, HQMAGs).

This approach uses genomes as nodes of ecological networks and their guild-level aggregations

as ecologically meaningful features for identifying microbiome signatures associated with host

phenotypes in health and diseases.

In this study, we hypothesized that bacteria required for providing essential health-relevant

functions to the host[2] should maintain stable ecological interactions with each other to form

robust guilds[16,17]. To identify microbiome signatures that are based on stable interactions among

HQMAGs, we randomized T2DM patients at baseline (M0) to receive either 3 months (M3) of a

high fiber intervention[11] (W group; n = 74) or standard care (U group; n= 36) followed by a one-

year follow-up (M15) in an open label, randomized, controlled trial (Fig. 1A and Fig. S1). The

high fiber intervention was used to exert an environmental perturbation to dramatically change

the abundance of members of the gut microbiome[10,11]. Co-abundance network analysis at each of

the three time points enabled us to identify genome pairs that can keep their correlations

unchanged despite significant community-wide abundance changes caused by the perturbations.

We found that these robust genome pairs were from 141 HQMAGs and these genomes formed

two guilds. These two guilds were organized as the two competing ends of a robust seesaw-like

network, whenever one guild increased, the other decreased in abundance. Together, these

seesaw networked genomes supported Random Forest models for predicting the response of a

wide range of metabolic phenotypes to dietary intervention in the T2DM cohort, as well as for

classifications of case and control of 8 different diseases in 12 independent metagenomic

datasets suggesting that we may have identified a core microbiome signature across different

chronic diseases.

Results

Reversible changes in the gut microbiota associate with reversible changes of host

metabolic phenotypes

Dietary fiber intake in the U group remained unchanged throughout the study, whereas W

group had a significant increase in the intake of dietary fibers from M0 to M3 and a decrease

from M3 to M15 (Fig. 1B). Compared with the U group, fiber intake was significantly higher in

the W group at both M3 and M15 (Fig. 1B), but energy and macronutrient consumption were

similar between the two groups across the study period (Fig. S2).

To investigate the gut microbial responses to the introduction and withdrawal of the high

fiber intervention, we performed shotgun metagenomic sequencing on 315 fecal samples

collected from 110 patients of the W and U group, among whom 95 patients provided samples at

all 3 time points and 15 provided samples at M0 and M3 only (Table S2, Fig S1). To achieve

genome-level resolution, we reconstructed 1,845 non-redundant high-quality draft genomes

(HQMAGs, two HQMAGs were collapsed into one if the average nucleotide identity, ANI,

between them was > 99%) from the metagenomic datasets. These HQMAGs accounted for more

than 70% of the total reads. In the context of beta-diversity based on Bray-Curtis distance, the

overall structure of the gut microbiota in the W group significantly changed from M0 to M3

(PERMANOVA test, $P < 0.001$) and returned to that of M0 at M15; there was no difference in

the U group across the 3 timepoints (Fig. 1C, D). Similar changes in alpha-diversity based on

Shannon and Simpson indices were also observed (Fig. S3). These results showed that the high

fiber intervention induced significant structural changes of the gut microbiota[11], however the gut

microbiota reverted to baseline after the intervention was withdrawn indicating a high resilience

in community structure.

To determine if host metabolic phenotypes would show similar reversible changes as the

gut microbiota, we examined 43 bio-clinical parameters of 9 categories across the 3 time points

(Table S3). Hemoglobin A1c (HbA1c) in the U group showed no significant changes throughout

the trial. The high fiber intervention significantly reduced the level of HbA1c in the W group

from M0 to M3 by 15.22 ± 9.82% (mean ± SD). At one-year follow-up of the W group, HbA1c

was significantly increased from M3 but remained lower than that at M0 (Fig. 1E). The

proportion of patients who achieved adequate glycemic control (HbA1c ≤ 7%) was significantly

higher in the W group (61.6 % versus 33.3% in the U group) at M3, but showed no difference

between the two groups at M15 (Fig. 1F). The level of fasting blood glucose and postprandial

glucose in meal tolerance test followed a similar trend as HbA1c (Fig. 1G, H). Among the rest 40

bio-clinical parameters, 14 also showed an alleviation from M0 to M3 but rebounded at one-year

follow-up in the W group (Table S3). These results indicate that changes of the host metabolic

phenotypes were associated with the reversible changes of the gut microbiota in response to the

introduction or withdrawal of the high fiber intervention.

Genome pairs with stable interactions form a seesaw-like network of two competing guilds

To facilitate the identification of genome pairs that keep their ecological interactions stable

during the trial, particularly in the W group with profound microbiota and host phenotypic

changes, we constructed a co-abundance network for each time point based on the abundance

matrix of the HQMAGs representing the prevalent microbes. Co-abundance network is a data-

driven way to investigate ecological interactions between microbes across habitats[18,19]. A total of

477 HQMAGs were selected for network construction because they were detectable in more than

75% of the samples at each time point in the W group. These 477 HQMAGs also accounted for

~60% of the total abundance of the 1,845 HQMAGs. In the W group, we calculated pairwise

correlations of all 113,526 possible genome pairs among these 477 prevalent HQMAGs based on

their abundance across the patients at each time point and constructed 3 co-abundance networks

(G_{M0}, G_{M3} and G_{M15}) (Figure 2A, Table S4). The three networks were of similar order S, i.e., the

total number of nodes (HQMAGs), S_{M0}(442), S_{M3}(421), and S_{M15}(429), but they varied

considerably in their size L, i.e., the total number of edges (correlations), L_{M0}(4,231), L_{M3}(2,587)

and L_{M15}(4,592). L in G_{M3} decreased to 61.14% of that in G_{M0} and rebounded back in G_{M15} to

108.53% of that in G_{M0}. This pattern was confirmed by changes in connectance, which is defined

as the proportion of realized ecological interactions among the potential ones (in undirected

network, connectance$=\dfrac{L}{S(S-1)/2}$, range: $[0,1]$)[20]. Connectance decreased from 0.043 in G_{M0} to

0.029 in G_{M3} and rebounded to 0.050 in G_{M15}. Changes in L and connectance showed that the

high fiber intervention dramatically reduced the correlations among the prevalent genomes in the

network. In addition, we found that the distributions of degree, i.e. the number of edges a node

has, fit well with a power-law model (Fig. S4, R2 values G_{M0}: 0.79, G_{M3}: 0.82, G_{M15}: 0.79),

indicating the presence of network hubs[21]. Defining hubs as nodes that connect with more than

one-fifth of the total nodes in the network (Fig. S5), we found 24 hubs, in which 10 were in $G_{M0,}$

20 were in G_{M15} but none were in G_{M3}. These results indicate that the overall structure of the gut

microbiome undergone profound changes during the trial, particularly, the high fiber intervention

resulted in the loss of interactions between genome pairs.

We considered two genomes are connected with robust and stable ecological relationship if

they keep the same type of correlation across all the three timepoints. Out of the 113,526

possible genome pairs, 92.39% showed no correlations at any of the three time points, suggesting

that it may be a rare event for two genomes to establish an ecological relationship even

transiently (Fig. 2B). Interestingly, 517 genome pairs showed positive correlations and 118

negative correlations at all the three time points. All the 635 stable correlations involved 184

genomes. Among these 184 HQMAGs, 43 were excluded from subsequent analysis because they

had no interactions with the remaining 141 nodes (Fig. S6) and most of them showed no

response to the high fiber intervention (M0 vs M3: $P > 0.05$, Friedman test followed by Nemenyi

post-hoc). The remaining 141 HQMAGs, which included genome pairs with 468 positive and

118 negative stable correlations throughout the trial were further defined as genomes with stable

ecological interactions (GSEIs) and became our microbiome signature candidates. We then

explored how these 141 GSEIs were connected with each other and with the rest of the nodes in

G_{M0}, G_{M3}, and G_{M15} (Fig. S7A). The 141 GSEIs had significantly higher degree, betweenness

centrality, eigenvector centrality, closeness centrality and stress centrality than the rest of the

genomes in the networks (Fig. S7B-F). This finding indicates that these GSEIs exerted a

relatively large amount of control over the interaction of other nodes (reflected by betweenness

centrality and eigenvector centrality) and the information flow in the network (reflected by

closeness centrality and stress centrality). Removing these GSEIs would lead to the collapse of

the networks since on average 86.08% of the total edges would have been lost. These suggest

that the 141 GSEIs can be considered as the core nodes of the networks as they were highly

connected not only within themselves but also with other nodes.

These 141 GSEIs were also highly prevalent among participants, as 140 of them were in >

90%, and 104 were in 100% of the 74 individuals in the W group (Fig. S8). In addition, most of

these 141 GSEIs were also predominant members of the gut microbiota as the abundance of 111

of them was higher than the median of the 1,845 HQMAGs and accounted for 20.78% of the

total sequencing reads. Based on Bray-Curtis distance, beta-diversity analysis showed significant

correlations between the profiles of the 141 GSEIs and all the 1,845 HQMAGs, as evidenced in

the Mantel test ($R^2 = 0.62$, $P = 0.001$) and Procrustes analysis ($P = 0.001$) (Fig. S9, Fig. 1C, D).

These indicate that the variations of the 141 GSEIs contributed to the major variations of the

whole gut microbial community across the 3 time points.

Based on the average linkage of these robust correlations, the 141 GSEIs fell into two

clusters. There were 50 genomes in cluster 1 and 91 genomes in cluster 2 (Fig.2C). These two

clusters can be considered as guilds as genome in each cluster were highly interconnected with

both robust and transient positive correlations[3] (Fig. 2D and Fig. S10). The two guilds were

connected by negative edges only, indicating a competitive relationship that structures a seesaw-

like network. Our data showed that the two competing guilds of the 141 GESIs existed in all

three ecological networks G_{M0}, G_{M3}, and G_{M15} in the W group. Furthermore, the finding of the

two competing guilds in the W group at M0 suggests that such microbial organization exists

irrespective of the high fiber intervention in our study. Given the similar overall gut microbiota

structure between the W and U groups at M0 and in the U group across 3 timepoints (Fig. 1C, D),

we speculated that two competing guilds can also be observed in the U group across the trial.

Thus, we constructed the co-abundance networks based on the abundance of the 141 GESIs

across the individuals in the U group at each time point. In the co-abundance networks, 99.8%,

99.51% and 99.74% of the total edges agreed with our two competing guilds, i.e., positive

correlations in the guilds and negative between guilds (Fig. S11A). This showed that the

detection of these two competing guilds was independent of the high fiber intervention,

indicating that the such a pattern may be an inherent structure of the gut microbiome in our study.

Functionality of the metagenomes of the two competing guilds associates with host

metabolic phenotypes

We sought to determine whether the balance between the two competing guilds could be

modulated by dietary fiber and describe how the two competing guilds affects the host metabolic

phenotypes. In the W group, the total abundance of Guild 1 increased and Guild 2 decreased

significantly from M0 to M3. Then at M15, Guild 1 decreased to a level similar to that at M0,

and Guild 2 bounced back but remained lower than that at M0. Subsequently, from M0 to M3,

high fiber intervention significantly increased the Guild 1-to-Guild 2 ratio. At one-year follow-

up, the ratio significantly decreased and was not different from M0 (Fig. 3A). Neither the

abundances of the 2 guilds nor their ratio was changed in the U group across the trial (Fig. S11B).

These results showed that the changes of the balance between the two guilds composed of GSEIs

were concomitant with changes in dietary fiber intake, overall gut microbiota and host

phenotypes. To further validate our hypothesis that GSEIs may be essential to host health, we

used the GSEIs as the selected features and applied machine leaning algorithms to explore the

associations between GSEIs and each host bio-clinical parameter. Random Forest regression via

leave-one-out cross-validation based on the 141 GSEIs showed that 41 out of the 43 bio-clinical

parameters had significant Pearson's correlation coefficient ranged from 0.11 to 0.44 (adjusted P

value < 0.05) between the predicted and measured values (Fig. 3B). These results showed that

the 141 genomes, as two competing guilds in a seesaw-like network, constitute an important

microbiome signature for T2DM and the related metabolic phenotypes.

Next, we performed genome-centric analysis of the HQMAGs in the two competing guilds

to explore the genetic basis underlying the association between the dynamic changes of the

seesaw networked microbiome signature and relationships with the host's metabolic phenotypes.

As the balance between the two guilds can be shifted by dietary fibers, we first sought to identify

carbohydrate-active enzyme (CAZy)-encoding genes and genes encoding key enzymes in short-

chain fatty acid (SCFA) production to compare the genetic capacity for carbohydrate utilization

between the two guilds. Compared with genomes in Guild 2, those in Guild 1 enriched CAZy

genes for arabinoxylan ($P < 0.001$), cellulose ($P < 0.01$) and had lower proportion of CAZy

genes for inulin utilization ($P < 0.01$) (Fig. 3C, Table S5). There was no difference in genes for

starch, pectin, and mucin utilization between the two guilds. Our previous study showed that gut

microbiota benefited patients with T2DM via acetic and butyric acid production from

carbohydrate fermentation[11]. Among the terminal genes for the butyrate biosynthetic pathways

from both carbohydrates (i.e., *but* and *buk*) and proteins (i.e., *atoA/D* and *4Hbt*), the copy number

of *but* was significantly higher in Guild 1 and there was no difference in the other terminal genes

between the two guilds (Fig. 3C). More than one-third of the genomes in Guild 1 harbored the

but gene while less then 5% of the genomes in Guild 2 had this gene (Fisher's exact test $P <$

0.001). Compared with Guild 2, Guild 1 also trended higher in its genetic capacity for acetate

production ($P = 0.06$) but had a lower genetic capacity for propionate production ($P < 0.05$) (Fig.

3C). These results showed that compared to Guild 2, Guild 1 had significantly higher genetic

capacity for utilizing complex plant polysaccharides and producing acetate and butyrate.

From the perspective of pathogenicity, 21 out of the 1,845 HQMAGs encoded 750

virulence factor (VF) genes. Among the 21 VF-encoding genomes, 3 were in Guild 1 while 18

were in Guild 2. Three out of the 50 genomes in Guild 1 had one VF gene involved in

antiphagocytosis. In Guild 2, 18 out of the 91 genomes encoded 747 VF genes across 15

different VF classes i.e., acid resistance, adherence, antiphagocytosis, biofilm formation, efflux

pump, endotoxin, invasion, iron uptake, manganese uptake, motility, nutritional factor, protease,

regulation, secretion system, and toxin (Fig. 3C, S12A). Notably, 98.53% of all the VF genes in

Guild 2 were harbored in 8 genomes (1 in *Enterobacter kobei*, 2 in *Escherichia flexneri,* 3 in

Escherichia coli and 2 in *Klebsiella*). The highly enriched genes for virulence factors in genomes

of Guild 2 (Fisher's Exact test, $P < 2.2 \times 10^{-16}$,) indicates that this guild may play an important

role in aggravating the metabolic disease phenotypes. In terms of antibiotic resistance genes

(ARG), in Guild 1, only 1 genome (2.00% of the genomes in this guild) harbored a copy of an

ARG related to phenicol (Fig. 3C, S12B). In Guild 2, 17 genomes (18.68% of the genomes in

this guild) encode 40 ARGs for resistance to 7 different antibiotic classes i.e., aminoglycosides,

beta-lactam, fosfomycin, glycopeptide, quinolone, macrolide, and tetracycline. Thus, Guild 2

may serve as a reservoir of ARGs for horizontal transfer to opportunistic pathogens. Taken

together, our data showed that the two competing guilds had distinct genetic capacity with Guild

1 being potentially beneficial and Guild 2 detrimental.

The seesaw networked microbiome signature exists in cohorts across ethnicity and

geography

We then asked that whether these 141 genomes, organized as two competing guilds in a

stable seesaw-like network, is this a microbiome signature unique in T2DM or it is common in

other diseases. To answer this question, we used the 141 GSEIs in our seesaw-like network as

reference genomes to perform read recruitment analysis, which is a commonly used method to

estimate abundance of reference genomes[22,23] in metagenomes (Fig. S13). In an independent

T2DM study[24], 32.92% of the reads were recruited and 128 of the GSEIs were detected as part of

a co-abundance network based on their estimated abundance across the T2DM patients. In this

co-abundance network, 97.82% of the total edges followed the pattern in our seesaw-like

network (i.e., positive edges within each guild and negative edges between the 2 guilds) (Fig.

4A), which further supported the existence of this seesaw-like network in T2DM patients.

Moreover, 35.28% of the reads were recruited in the metagenomes of 136 healthy controls of the

same study[24] and 119 of the GSEIs were constructed into a co-abundance network in which

99.45 % of the total edges agreed with our seesaw-like network (Fig. 4A). In the context of beta-

diversity based on Bray-Curtis distance, our microbiome signature showed significant

differences (PERMAONVA test $P = 2 \times 10^{-4}$) between T2DM patients and the healthy controls

based on the abundance matrix of the reference genomes (Fig. 4B). This suggests that the

variation of this microbiome signature was associated with T2DM in this independent dataset. To

further validate such associations, using the abundance matrix of the genomes in the microbiome

signature as input features and the phenotype data as target variable, we constructed Random

Forest regression models and found that this microbiome signature was significantly correlated

with BMI, fasting insulin, and HbA1c (Fig S14). Furthermore, we developed a machine learning

classifier based on a Random Forest algorithm to see if we can classify patients and control.

Receiver operating characteristic curve analysis showed a moderate diagnostic power with area

under the curve (AUC) of 0.70 by a leave-one-out cross-validation. Thus, we showed that our

seesaw networked microbiome signature not only existed in an independent T2DM study but

also maintained a similar relationship with the host metabolic phenotypes.

We then extended our hypothesis that the seesaw networked microbiome signature

represents an inherent feature of human gut microbiome and the disruption of which may be

related to diseases in addition to T2DM. We first performed the same validation analysis in

metagenomic datasets of three different types of diseases, including atherosclerotic

cardiovascular disease (ACVD) [25], liver cirrhosis (LC)[26] and ankylosing spondylitis (AS)[27]. In

ACVD patients and their controls, 36.21% and 32.73% of the reads were recruited respectively,

and 134 genomes from the patients and 133 genomes from the controls were constructed into co-

abundance networks with 97.32% and 97.70% of the total edges respectively agreed with our

seesaw-like network (Fig. 4A). There were 33.84%, 35.83% and 41.02% of the reads recruited to

the reference genomes in the metagenomic datasets of the healthy control (the studies on LC and

AS employed the same control cohort), LC and AS patients respectively. With the 141 GSEIs as

reference, 112, 113 and 113 genomes were constructed into co-abundance networks with 99.80%,

98.81% and 98.19% of the total edges agreed with our seesaw-like network in the metagenomic

datasets of the healthy control, LC and AS patients respectively (Fig. 4A). In the PCoA plot

based on Bray-Curtis distance, our microbiome signature showed significant differences

(PERMANOVA test, $P < 0.001$) between control and patients in all 3 datasets (Fig. 4B). In the

LC study, we also used the abundance matrix of the genomes in the microbiome signature as

input features and the phenotype data as target variable to construct Random Forest regression

models and found that our microbiome signature was significantly correlated with total bilirubin,

albumin level, and BMI (Fig. S15). Compared with the T2DM dataset[24], the Random Forest

classifier based on our microbiome signature showed better diagnostic power in distinguishing

case from control for ACVD (AUC = 0.80), LC (AUC = 0.90), and AS (AUC = 0.98) (Fig. 4C).

To further confirm the relevance of this microbiome signature to human diseases, we

estimated the abundances of the genomes from this microbiome signature in datasets from more

disease types and across different ethnicity and geography. These datasets included inflammatory

bowel diseases (IBD) (American cohort and Dutch cohort), colorectal cancer (CRC) (Chinese

cohort, Australian cohort and German cohort), schizophrenia (Chinese cohort), and Parkinson's

disease (PD) (Chinese cohort). On average, 31.32% ± 4.21% (mean ± SD) of reads were

recruited to the reference genomes in these datasets. We validated that this microbiome signature

showed diagnostic power to classify case and control in the metagenomic dataset from studies on

IBD (AUC = 0.71 for IBD dataset 1[28], AUC=0.91 for IBD dataset 2[29] and AUC=0.83 for IBD

dataset 3[29]), CRC (AUC = 0.74 for CRC dataset 1[30], AUC = 0.75 for CRC dataset 2[31] and AUC

= 0.71 for CRC dataset 3[32]), schizophrenia[33] (AUC = 0.68), and PD[34] (AUC = 0.77) (Fig. S16).

In addition, we used MMUPHin[35] to correct batch effects from the different cohorts in IBD and

CRC and applied leave-one-cohort-out (LOCO) analysis[36] to evaluate the universality of the

diagnostic power of this microbiome signature in these two diseases. The AUC values from

LOCO analysis were 0.77 to 0.84 for IBD and 0.68 to 0.70 for CRC (Fig. S17). These results

showed the existence of our microbiome signature in healthy controls and various patient

populations across ethnicity and geography from independent studies. The associations between

the 141 genomes and host phenotypes and their discriminative power as biomarkers to classify

controls vs. patients with various types of diseases indicate that these genomes, organized as two

guilds in a seesaw-like network, represent a common microbiome signature associated with a

wide range of human diseases.

Discussion

In the current study, our genome-centric, reference-free, and ecological-interaction-focused

approach led to the identification of a robust seesaw-like network of two competing guilds of

bacterial genomes, whose changes were associated with a wide range of host phenotypes in

patients with T2DM. Moreover, random forest models based on these genomes classified case

and control across a wide range of diseases, indicating that these genomes may form a core

microbiome signature that exists in populations of diverse ethnicity, geography, and disease

status.

Our novel microbiome signature organizes genomes in a seesaw-like network exhibiting

both cooperative and competitive interactions. Cooperative ecological networks are expected to

promote overall metabolic efficiency, such as the co-operative metabolism that benefits the

host[37]. However, they also create dependency and the potential for mutual downfall that may

bring destabilizing effect on the gut microbial ecosystem. This destabilizing effect of cooperation

can be dampened by introducing ecological competition into the network[37]. Thus, a seesaw-like

network with both cooperative and competitive interactions may represent the key characteristic

of a stable microbiome structure[37]. Interestingly, while the seesaw-like network is stable, the

weight of the two ends i.e., the abundances of Guild 1 and Guild 2, are modifiable and such

changes are associated with host health. When large amount of complex fiber was introduced or

withdrawn, Guilds 1 and 2 showed no change in membership nor in the types of interactions with

each other but experienced dramatic shifts in guild-level abundance in a competing manner.

Members in Guild 1 have higher genetic capacity for degrading complex plant polysaccharides

and produce beneficial metabolites including SCFAs which may suppress populations of

pathobionts in Guild 2[11]. Members of Guild 2 need to be kept low since their overgrowth may

jeopardize host health by increasing inflammation, etc.[38]. However, pathobionts in Guild 2

cannot be eliminated, e.g., they could serve as the necessary agents that train our immune system

from early days[39,40]. Therefore, the balance between Guild 1 and Guild 2 becomes critical in

determining whether the gut microbiome supports health or aggravate diseases. This seesaw-like

network between Guilds 1 and 2 allows the genomes in our microbiome signature to readily

respond to changes of external energy input to the gut microbial ecosystem and mediate its

impact on host health, while simultaneously maintains its structural integrity. Such structural

integrity may be key to long-term ecological stability of the gut microbiome and its ability to

provide essential health-relevant functions to the host.

The fact that this seesaw-like network can be detected in other independent metagenomic

datasets and is shown correlated with different diseases indicates that such an ecological

organization is not specific to T2DM nor dependent on the high fiber diet we used, Such a

seesaw networked structure may have been stabilized by natural selection over a long history of

co-evolution between microbiomes and their hosts[16 41]. A selection pressure may have been

exerted by dietary fibers that interact directly with gut bacteria as external energy source[42,43].

Studies on coprolites showed that dietary fiber intake was much higher in ancient humans and

only reduced significantly in the past 150 years[44,45] (130 g/d of plant fiber intake in prehistoric

diet[46] vs. a median intake of 12–14 g/d in the modern American diet[47]). Such a high fiber intake

over evolutionary history may have favored beneficial bacteria in Guild 1 due to their higher

genetic capacity to utilize plant polysaccharides as an external energy supply, enabling them to

gain competitive advantage over pathobionts in Guild 2[48]. Akin to tall trees as the foundation

species for a closed forest, Guild 1 may work as the "foundation guild" for stabilizing a healthy

gut microbiome and keeping the pathobionts at bay[49]. The dominance of Guild 1 over Guild 2

can increase host fitness as shown by the epidemiologically and clinically proven health benefits

of dietary fibers in both preventing and alleviating a wide range of chronic conditions[11,43,50,51].

Moreover, the two seesaw networked guilds may constitute a core gut microbiome

signature in humans[52,53], since 1) they are commonly shared among populations across ethnicity

and geography; 2) they show temporal stability not only in membership but also in their

interactions with each other and the host; 3) they make up about 10% of the gut microbiome

membership but are disproportionally important for shaping the ecological community; 4) they

397 support the provision of essential health-relevant functions to the host; and 5) such a core

398 microbiome organized in a seesaw-like network may have been established over a long history of

399 co-evolution.

400  As an evolutionarily conserved ecological structure, our seesaw-like network demonstrated

401 remarkably stable relationships both internally within the network and externally with the host.

402 Within the seesaw-like network, it is the imbalance between the two competing guilds that may

403 play a role as the common biological basis for many human diseases. Targeting this core

404 microbiome signature to restore and maintain ecological dominance of the beneficial guild over

405 the detrimental guild could be fundamentally important to human health recovery and

406 maintenance.

407 **Materials and Methods**

408 **Clinical Experiment**

409 **Study design**[11]: This clinical trial, conducted at the Qidong People's Hospital (Jiangsu, China),

410 examined the effect of a high fiber diet in free-living conditions in a cohort of individuals

411 clinically diagnosed T2DM (QIDONG). The study protocol was approved by Ethics Committee

412 of Shanghai General Hospital (2014KY104), and the study was conducted in accordance with the

413 principles of the Declaration of Helsinki. All participants provided written informed consent. The

414 trial was registered in the Chinese Clinical Trial Registry (ChiCTR-IPC-14005346). The study

415 design and participant flow are shown in Fig. S1.

416  T2DM patients of the Chinese Han ethnicity were recruited for the study (age: 37 - 70 years;

417 HbA1c: 6.5% - 12.0%. More detailed description of inclusion and exclusion criteria were shown

418 in Chinese Clinical Trial registry (http://www.chictr.org.cn).

Patients received either a high-fiber diet (WTP diet) as the treatment group (W group) or the

usual care (Usual diet) as the control group (U group) for 3 months. Total caloric and

macronutrients prescriptions were based on age-specific Chinese Dietary Reference Intakes

(Chinese Nutrition Society, 2013). The WTP diet, based on wholegrains, traditional Chinese

medicinal foods and prebiotics, included three ready-to-consume pre-prepared foods[11]. The usual

care included standard dietary and exercise advice that was made according to the Chinese

Diabetes Society guidelines for T2DM[54]. Patients in W group were provided with the WTP diet

to perform a self-administered intervention at home for three months, while patients in U group

accepted the usual care. W group stopped WTP diet intervention at the end of the third month (at

M3). Then W and U continued a one-year follow-up (M15). A meal-based food frequency

questionnaire and 24-h dietary recall were used to calculate nutrient intake based on the China

Food Composition 2009[55]. Patients in both groups continued with their antidiabetic medications

according to their physician prescriptions (Table S1).

Before a 2-week run-in period, all participants attended a lecture on diabetes intervention and

improvements and received diabetes education and metabolic assessments. 119 eligible

individuals were enrolled based on the inclusion and exclusion criteria and assigned into two

groups in a 2:1 ratio (n = 79 in W group, n = 40 in U group) determined by SAS software.

Physical examinations were carried out at M0, M3, and M15 in Qidong People's Hospital

(Jiangsu, China). Sample collection instructions were provided to the participants at the day

before. The participants provided the feces and first early morning urine as requested. After

collecting fasting venous blood sample, a 3-h meal tolerance test (Chinese buns containing 75 g

of available carbohydrates; MTT test) was conducted and the postprandial venous blood samples

at 30, 60, 120, and 180 min were collected. All the blood samples were centrifuged at 3000 rpm

for 20 min at 4□ after standing at room temperature for 30 min to obtain serum. The fasting

blood serum were divided into two parts, one used for hospital tests and the other used for lab

tests. The feces, urine, and serum samples were stored in dry ice immediately then transported to

lab and frozen at -80℃. Subsequently, anthropometric markers and diabetic complication

indexes were measured. Ewing test[56] and 24-h dynamic electrocardiogram were conducted to

estimate diabetic autonomic neuropathy (DAN). B-mode carotid ultrasound was conducted to

estimate atherosclerosis. Michigan Neuropathy Screening Instrument[57] was conducted to

estimate diabetic peripheral neuropathy (DPN). In addition, A meal-based food frequency

questionnaire and the 24-h dietary review were recorded for nutrient intake calculation. The drug

use was self-reported and presented in table S1.

The fasting venous blood was used to measure HbA1c, fasting blood glucose, fasting insulin,

fasting C-Peptide, C-reactive protein (CRP), blood routine examination, blood biochemical

examination and five analytes of thyroid. The venous blood samples at 30, 60, 120, and 180 min

of MTT were used to measure the postprandial blood glucose, insulin, and C-Peptide. The

fasting early morning urine was used to measure the routine urine examination and urinary

microalbumin creatinine ratio. The measurements above were completed at Qidong People's

Hospital. Fasting venous blood was used to quantify TNF-α (R&D Systems, MN, USA),

lipopolysaccharide-binding protein (Hycult Biotech, PA, USA), leptin (P&C, PCDBH0287,

China) and adiponectin (P&C, PCDBH0016, China) by enzyme-linked immunosorbent assays

(ELISAs) at Shanghai Jiao Tong University.

The homeostatic model assessments of insulin resistance (HOMA-IR) and islet β-cell function

(HOMA-β) were calculated based on fasting blood glucose (mmol/L) and fasting C-Peptide

(pmol/L)[58]: HOMA-IR = 1.5 + FBG * Fasting-C-Peptide / 2800;

HOMA-β = 0.27 * Fasting-C-Peptide / (FBG - 3.5). Glomerular Filtration Rate was estimated by

formula GFR (ml/min per 1.73 m^2) = 186 * Scr$^{-1.154}$ * age$^{-0.203}$ * 0.742 (if female) * 1.233 (if

Chinese)[59], where Scr (serum creatinine) is in mg/dl and age is in years.

<u>Gut microbiome analysis</u>

Metagenomic sequencing. DNA was extracted from fecal samples using the methods as

previously described[10]. Metagenomic sequencing was performed using Illumina Hiseq 3000 at

GENEWIZ Co. (Beijing, China). Cluster generation, template hybridization, isothermal

amplification, linearization, and blocking denaturing and hybridization of the sequencing primers

were performed according to the workflow specified by the service provider. Libraries were

constructed with an insert size of approximately 500 bp followed by high-throughput sequencing

to obtain paired-end reads with 150 bp in the forward and reverse directions. Table S3 shows the

number of raw reads of each sample.

Data quality control. Prinseq[60] was used to: 1) trim the reads from the 3′ end until reaching the

first nucleotide with a quality threshold of 20; 2) remove read pairs when either read was < 60 bp

or contained "N" bases; and 3) de-duplicate the reads. Reads that could be aligned to the human

genome (H. sapiens, UCSC hg19) were removed (aligned with Bowtie2[61] using --reorder --no-hd

--no-contain --dovetail). Table S3 shows the number of high-quality reads of each sample for

further analysis.

De novo assembly, abundance calculation, and taxonomic assignment of genomes. De novo

assembly was performed for each sample by using IDBA_UD[62] (--step 20 --mink 20 --maxk 100

--min_contig 500 --pre_correction). The assembled contigs were further binned using

MetaBAT[63] (--minContig 1500 --superspecific -B 20). The quality of the bins was assessed

using CheckM[64]. Bins had completeness > 95%, contamination < 5% and strain heterogeneity <

5% were retained as high-quality draft genomes (Table S6). The assembled high-quality draft

genomes were further dereplicated by using dRep[65]. DiTASiC[66], which applied kallisto for

pseudo-alignment[67] and a generalized linear model for resolving shared reads among genomes,

was used to calculate the abundance of the genomes in each sample, estimated counts with P-

value > 0.05 were removed, and all samples were downsized to 36 million reads (One sample

with read mapping ratio < 25%, which could not be well represented by the high quality

genomes, were removed in downstream analysis). Taxonomic assignment of the genomes was

performed by using GTDB-Tk[68] with default parameters (Table S7).

Gut microbiome functional analysis. Prokka[69] was used to annotate the genomes. KEGG

Orthologue (KO) IDs were assigned to the predicted protein sequences in each genome by

HMMSEARCH against KOfam using KofamKOALA with default parameters[70]. Antibiotic

resistance genes were predicted using ResFinder[71] with default parameters. The identification of

virulence factors were based on the core set of Virulence Factors of Pathogenic Bacteria

Database (VFDB[72], download July 2020). The predicted proteins sequences were aligned to the

reference sequence in VFDB using BLASTP (best hist with E-value < 1e-5, identity > 80% and

query coverage > 70%). Genes encoding carbohydrate-active enzymes (CAZys) were identified

using dbCAN (releasee 6.0)[73], and the best-hit alignment was retained. Genes encoding formate-

tetrahydrofolate ligase, propionyl-CoA: succinate-CoA transferase, propionate CoA-transferase,

4Hbt, AtoA, AtoD, Buk and But were identified as described previously[11].

Gut microbiome network construction and analysis. In W group, prevalent genomes shared

by more than 75% of the samples at every timepoint were used to construct the co-abundance

network at each timepoint. Fastspar[74], a rapid and scalable correlation estimation tool for

microbiome study, was used to calculate the correlations between the genomes with 1,000

permutations at each time point based on the abundances of the genomes across the patients and

the correlations with $P \leq 0.001$ were retained for further analysis. The networks were visualized

with Cytoscape v3.8.1[75]. The layout of the nodes and edges was determined by Edge-weighted

Spring Embedded Layout using the correlation coefficient as weights. The links between the

nodes are treated as metal springs attached to the pair of nodes. The correlation coefficient was

used to determine the repulsion and attraction of the spring[75]. The layout algorithm sets the

position of the nodes to minimize the sum of forces in the network. We defined robust stable

edges as the unchanged positive/negative correlations between the same two genomes across all

the 3 networks at M0, M3, and M15. Stable genome pairs were clustered based on robust

positive (set as 1) and negative (set as -1) edges with average clustering. We used iTOL[76], an

online tool for display, manipulation, and annotation for various trees, to integrate and visualize

the clustering tree, taxonomy information, and abundance changes of the 141 genomes.

Validation in independent cohorts. Twelve independent metagenomic datasets were

downloaded from SRA or ENA database. The group information was collected from the

corresponding papers or from curatedMetagenomicData[77] (Table S8). DiTASiC was used to

recruit reads and estimate the abundance of the 141 genomes in each sample, estimated counts

with P-value > 0.05 were removed and further converted to relative abundance divided by the

total number of reads. To reduce false positive in the validation dataset, relative abundance <

0.001% were further removed. A random forest classification model to classify case and control

was constructed based on the estimated abundances of the genomes in each dataset with leave-

one-out cross-validation.

MMUPHin[35] was used to adjust the estimated abundances of the genomes by correcting

batched effects from the different cohorts in IBD and CRC studies. Random forest classification

models with leave-one-cohort-out analysis were further performed on the adjusted abundance

matrix[36].

Datasets from 4 studies were included to validate the commonality of the seesaw-like

network. These datasets were from 136 control and 136 T2DM individuals in Qin et al., 2012[24];

171 control and 214 atherosclerotic cardiovascular disease individuals in Jie et al., 2017[25]; 83

control and 84 liver cirrhosis individuals in Qin et al., 2014[26]; and 83 control and 97 ankylosing

spondylitis individuals in Wen et al., 2017[27]. Fastspar was used to calculate the correlations

between the genomes with 1,000 permutations and the correlations with $P \leq 0.001$ were

remained for constructing the networks. 30 repeat 5-fold cross-validation was used and the

correlations shared by more than 95% of the 150 networks constructed from the cross-validation

process were remained in the final network.

Statistical Analysis.

Statistical analysis was performed in the R environment (R version3.6.1). Friedman test

followed by Nemenyi post-hoc test was used for intra-group comparisons for repeat

measurements. Mann-Whitney test (two-sided) was used for comparisons between W and U at

the same time point. Pearson Chi-square tests was performed to compare the differences of

categorical data between groups or timepoints. PERMANOVA test (9,999 permutations) was

used to compare the groups of gut microbiota structure.

Mann-Whitney test (two-sided) and Fisher's exact test (two sided) were used to compare

the functions between Guild 1 and Guild 2. Random Forest with leave-one-out cross-validation

was used to perform regression and classification analysis based on this microbiome signature

and clinical parameters/groups.

Acknowledgements

This work was supported by grants from National Natural Science Foundation of China

(31930022, 81871091, 81870594 and 81870596), the National Key Research and Development

Project (2019YFA0905600), the School of Environmental and Biological Sciences and the New

Jersey Institute for Food, Nutrition, and Health (seed pilot grant cycle 1, awarded in 2019),

Canadian Institute for Advanced Research, Notitia Biotechnologies Company, Clinical Research

Plan of SHDC [No. SHDC2020CR1016B], and the Project of Songjiang District Municipal

Health Commission (0702N18003).

Author contributions

L.Z., Y.P., C.Z. and Y.S. designed the supervised the study; T.X., F.Z., X.D. and Y.S. managed

the clinical research; T.X., X.D., D.W., J.F., Y.S., X.L., S.J., X.W. and F.Z. conducted the

clinical study; T.X., X.W., H.F. and C.Z. performed clinical testing and data analysis; H.F.

provided dietary counselling to the participants and conducted dietary analysis; T.X. and X.W.

prepared the fecal DNA for sequencing. G.W., X.T., C.Z. performed microbial data analysis.

G.W. performed data validation in independent datasets. L.Z., C.Z., G.W., N.Z. and Y.Y.L.

wrote and revised manuscript. G.W., T.X., N.Z., Y.Y.L. and X.D. contributed equally to this

work.

Competing financial interests

L.Z. is a co-founder of Notitia Biotechnologies Company.

Data availability

The metagenomic sequencing data has been deposited under accession numbers PRJEB15179

and PRJEB52732

Code availability

Parameters of the bioinformatic tools applied in the study were showed in the method section.

Scripts and command lines related to the current study are freely available from the

corresponding author (L.Z., liping.zhao@rutgers.edu) upon request.

Ethics & Inclusion statement

We have carefully considered research contributions and authorship criteria when involved in

multi-region collaborations involving local researchers so as to promote greater equity in

research collaborations.

Reference

1   Dominguez-Bello, M. G., Godoy-Vitorino, F., Knight, R. & Blaser, M. J. Role of the
602   microbiome in human development. *Gut* **68**, 1108-1114, doi:10.1136/gutjnl-2018-
603   317503 (2019).
604   2   Kundu, P., Blacher, E., Elinav, E. & Pettersson, S. Our Gut Microbiome: The Evolving
605   Inner Self. *Cell* **171**, 1481-1493, doi:10.1016/j.cell.2017.11.024 (2017).

3   Wu, G., Zhao, N., Zhang, C., Lam, Y. Y. & Zhao, L. Guild-based analysis for understanding
gut microbiome in human health and diseases. *Genome Medicine* **13**, 22,
doi:10.1186/s13073-021-00840-y (2021).
4   O'Hara, A. M. & Shanahan, F. The gut flora as a forgotten organ. *Embo Rep* **7**, 688-693,
doi:10.1038/sj.embor.7400731 (2006).
5   Koh, A. & Backhed, F. From Association to Causality: the Role of the Gut Microbiota and
Its Functional Products on Host Metabolism. *Mol Cell* **78**, 584-596,
doi:10.1016/j.molcel.2020.03.005 (2020).
6   Sanna, S. *et al.* Causal relationships among the gut microbiome, short-chain fatty acids
and metabolic diseases. *Nat Genet* **51**, 600-+, doi:10.1038/s41588-019-0350-x (2019).
7   Meijnikman, A. S., Gerdes, V. E., Nieuwdorp, M. & Herrema, H. Evaluating Causality of
Gut Microbiota in Obesity and Diabetes in Humans. *Endocr Rev* **39**, 133-153,
doi:10.1210/er.2017-00192 (2018).
8   Levin, S. A. Ecosystems and the biosphere as complex adaptive systems. *Ecosystems* **1**,
431-436 (1998).
9   Zhang, C. & Zhao, L. Strain-level dissection of the contribution of the gut microbiome to
human metabolic disease. *Genome Med* **8**, 41, doi:10.1186/s13073-016-0304-1 (2016).
10   Zhang, C. H. *et al.* Dietary Modulation of Gut Microbiota Contributes to Alleviation of
Both Genetic and Simple Obesity in Children. *Ebiomedicine* **2**, 968-984,
doi:10.1016/j.ebiom.2015.07.007 (2015).
11   Zhao, L. *et al.* Gut bacteria selectively promoted by dietary fibers alleviate type 2
diabetes. *Science* **359**, 1151-1156, doi:10.1126/science.aao5774 (2018).
12   Tierney, B. T., Tan, Y., Kostic, A. D. & Patel, C. J. Gene-level metagenomic architectures
across diseases yield high-resolution microbiome diagnostic indicators. *Nat Commun* **12**,
2907, doi:10.1038/s41467-021-23029-8 (2021).
13   Wang, J. & Jia, H. Metagenome-wide association studies: fine-mining the microbiome.
Nat Rev Microbiol **14**, 508-522, doi:10.1038/nrmicro.2016.83 (2016).
14   Duvallet, C., Gibbons, S. M., Gurry, T., Irizarry, R. A. & Alm, E. J. Meta-analysis of gut
microbiome studies identifies disease-specific and shared responses. *Nat Commun* **8**,
1784, doi:10.1038/s41467-017-01973-8 (2017).
15   Jackson, M. A. *et al.* Gut microbiota associations with common diseases and prescription
medications in a population-based cohort. *Nat Commun* **9**, 2655, doi:10.1038/s41467-
018-05184-7 (2018).
16   Foster, K. R., Chluter, J. S., Oyte, K. Z. C. & Rakoff-Nahoum, S. The evolution of the host
microbiome as an ecosystem on a leash. *Nature* **548**, 43-51, doi:10.1038/nature23292
(2017).
17   Sommer, F., Anderson, J. M., Bharti, R., Raes, J. & Rosenstiel, P. The resilience of the
intestinal microbiota influences health and disease. *Nat Rev Microbiol* **15**, 630-638,
doi:10.1038/nrmicro.2017.58 (2017).
18   Ma, B. *et al.* Earth microbial co-occurrence network reveals interconnection pattern
across microbiomes. *Microbiome* **8**, 82, doi:10.1186/s40168-020-00857-2 (2020).
19   Bauer, E. & Thiele, I. From Network Analysis to Functional Metabolic Modeling of the
Human Gut Microbiota. *mSystems* **3**, doi:10.1128/mSystems.00209-17 (2018).

20 Poisot, T. & Gravel, D. When is an ecological network complex? Connectance drives
degree distribution and emerging network properties. *PeerJ* **2**, e251,
doi:10.7717/peerj.251 (2014).
21 Barabasi, A. L. Network science. *Philos Trans A Math Phys Eng Sci* **371**, 20120375,
doi:10.1098/rsta.2012.0375 (2013).
22 Nayfach, S. *et al.* A genomic catalog of Earth's microbiomes. *Nat Biotechnol* **39**, 499-509,
doi:10.1038/s41587-020-0718-6 (2021).
23 Reji, L., Cardarelli, E. L., Boye, K., Bargar, J. R. & Francis, C. A. Diverse ecophysiological
adaptations of subsurface Thaumarchaeota in floodplain sediments revealed through
genome-resolved metagenomics. *ISME J*, doi:10.1038/s41396-021-01167-7 (2021).
24 Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2 diabetes.
Nature **490**, 55-60, doi:10.1038/nature11450 (2012).
25 Jie, Z. *et al.* The gut microbiome in atherosclerotic cardiovascular disease. *Nat Commun*
8, 845, doi:10.1038/s41467-017-00900-1 (2017).
26 Qin, N. *et al.* Alterations of the human gut microbiome in liver cirrhosis. *Nature* **513**, 59-
64, doi:10.1038/nature13568 (2014).
27 Wen, C. *et al.* Quantitative metagenomics reveals unique gut microbiome biomarkers in
ankylosing spondylitis. *Genome Biol* **18**, 142, doi:10.1186/s13059-017-1271-6 (2017).
28 Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel
diseases. *Nature* **569**, 655-662, doi:10.1038/s41586-019-1237-9 (2019).
29 Franzosa, E. A. *et al.* Gut microbiome structure and metabolic activity in inflammatory
bowel disease. *Nat Microbiol* **4**, 293-305, doi:10.1038/s41564-018-0306-4 (2019).
30 Yu, J. *et al.* Metagenomic analysis of faecal microbiome as a tool towards targeted non-
invasive biomarkers for colorectal cancer. *Gut* **66**, 70-+, doi:DOI 10.1136/gutjnl-2015-
309800 (2017).
31 Feng, Q. *et al.* Gut microbiome development along the colorectal adenoma-carcinoma
sequence. *Nat Commun* **6**, 6528, doi:10.1038/ncomms7528 (2015).
32 Wirbel, J. *et al.* Meta-analysis of fecal metagenomes reveals global microbial signatures
that are specific for colorectal cancer. *Nat Med* **25**, 679-689, doi:10.1038/s41591-019-
0406-6 (2019).
33 Zhu, F. *et al.* Metagenome-wide association of gut microbiome features for
schizophrenia. *Nat Commun* **11**, 1612, doi:10.1038/s41467-020-15457-9 (2020).
34 Qian, Y. W. *et al.* Gut metagenomics-derived genes as potential biomarkers of
Parkinson's disease. *Brain* **143**, 2474-2489 (2020).
35 Ma, S. *et al.* Population Structure Discovery in Meta-Analyzed Microbial Communities
and Inflammatory Bowel Disease. *bioRxiv*, 2020.2008.2031.261214,
doi:10.1101/2020.08.31.261214 (2020).
36 Wu, Y. *et al.* Identification of microbial markers across populations in early detection of
colorectal cancer. *Nat Commun* **12**, 3063, doi:10.1038/s41467-021-23265-y (2021).
37 Coyte, K. Z., Schluter, J. & Foster, K. R. The ecology of the microbiome: Networks,
competition, and stability. *Science* **350**, 663-666, doi:10.1126/science.aad2602 (2015).
38 Kamada, N., Seo, S. U., Chen, G. Y. & Nunez, G. Role of the gut microbiota in immunity
and inflammatory disease. *Nat Rev Immunol* **13**, 321-335, doi:10.1038/nri3430 (2013).

39   Vatanen, T. *et al.* Variation in Microbiome LPS Immunogenicity Contributes to
693   Autoimmunity in Humans. *Cell* **165**, 1551, doi:10.1016/j.cell.2016.05.056 (2016).
694   40   Bach, J. F. The hygiene hypothesis in autoimmunity: the role of pathogens and
695   commensals. *Nat Rev Immunol* **18**, 105-120, doi:10.1038/nri.2017.111 (2018).
696   41   Risely, A. Applying the core microbiome to understand host–microbe systems. *Journal of*
697   *Animal Ecology* **89**, 1549-1558 (2020).
698   42   Makki, K., Deehan, E. C., Walter, J. & Backhed, F. The Impact of Dietary Fiber on Gut
699   Microbiota in Host Health and Disease. *Cell Host Microbe* **23**, 705-715,
700   doi:10.1016/j.chom.2018.05.012 (2018).
701   43   Reynolds, A. *et al.* Carbohydrate quality and human health: a series of systematic
702   reviews and meta-analyses. *The Lancet* **393**, 434-445 (2019).
703   44   Eaton, S. B. The ancestral human diet: what was it and should it be a paradigm for
704   contemporary nutrition ? *P Nutr Soc* **65**, 1-6, doi:10.1079/Pns2005471 (2006).
705   45   Jew, S., AbuMweis, S. S. & Jones, P. J. Evolution of the human diet: linking our ancestral
706   diet to modern functional foods as a means of chronic disease prevention. *J Med Food*
707   **12**, 925-934, doi:10.1089/jmf.2008.0268 (2009).
708   46   Spiller, G. A. & Amen, R. J. *Topics in dietary fiber research*. (Springer, 1978).
709   47   Thompson, H. J. & Brick, M. A. Perspective: Closing the dietary fiber gap: An ancient
710   solution for a 21st century problem. *Advances in Nutrition* **7**, 623-626 (2016).
711   48   Deehan, E. C. *et al.* Modulation of the gastrointestinal microbiome with nondigestible
712   fermentable carbohydrates to improve human health. *Microbiology spectrum* **5**, 5.5. 04
713   (2017).
714   49   Prevey, J. S., Germino, M. J. & Huntly, N. J. Loss of foundation species increases
715   population growth of exotic forbs in sagebrush steppe. *Ecol Appl* **20**, 1890-1902,
716   doi:10.1890/09-0750.1 (2010).
717   50   Anderson, J. W. *et al.* Health benefits of dietary fiber. *Nutr Rev* **67**, 188-205,
718   doi:10.1111/j.1753-4887.2009.00189.x (2009).
719   51   Kaczmarczyk, M. M., Miller, M. J. & Freund, G. G. The health benefits of dietary fiber:
720   beyond the usual suspects of type 2 diabetes mellitus, cardiovascular disease and colon
721   cancer. *Metabolism* **61**, 1058-1066, doi:10.1016/j.metabol.2012.01.017 (2012).
722   52   Risely, A. Applying the core microbiome to understand host-microbe systems. *J Anim*
723   *Ecol* **89**, 1549-1558, doi:10.1111/1365-2656.13229 (2020).
724   53   Berg, G. *et al.* Microbiome definition re-visited: old concepts and new challenges.
725   *Microbiome* **8**, 103, doi:10.1186/s40168-020-00875-0 (2020).
726   54   Society, C. D. China guideline for type 2 diabetes (2013 edition). *Chin J Diabetes* **22**
727   (2014).
728   55   yuexin, Y., guangya, W. & xingchang, P. China Food Composition (Book 1, Beijing Medical
729   Univ. Press, ed. 2, 2009). (2009).
730   56   Ewing, D. & Clarke, B. Diagnosis and management of diabetic autonomic. *British Medical*
731   *Journal* **285** (1982).
732   57   Feldman, E. L. *et al.* A Practical Two-Step Quantitative Clinical and Electrophysiological
733   Assessment for the Diagnosis and Staging of Diabetic Neuropathy. *Diabetes Care* **17**,
734   1281-1289 (1994).

58 Li, X., Zhou, Z. G., Qi, H. Y., Chen, X. Y. & Huang, G. Replacement of insulin by fasting C-
peptide in modified homeostasis model assessment to evaluate insulin resistance and
islet beta cell function. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* **29**, 419-423 (2004).
59 Ma, Y. C. *et al.* Modified glomerular filtration rate estimating equation for Chinese
patients with chronic kidney disease. *J Am Soc Nephrol* **17**, 2937-2944,
doi:10.1681/ASN.2006040368 (2006).
60 Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic datasets.
Bioinformatics **27**, 863-864, doi:10.1093/bioinformatics/btr026 (2011).
61 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9, 357-359, doi:10.1038/nmeth.1923 (2012).
62 Peng, Y., Leung, H. C., Yiu, S. M. & Chin, F. Y. IDBA-UD: a de novo assembler for single-
cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics* **28**,
1420-1428, doi:10.1093/bioinformatics/bts174 (2012).
63 Kang, D. D., Froula, J., Egan, R. & Wang, Z. MetaBAT, an efficient tool for accurately
reconstructing single genomes from complex microbial communities. *PeerJ* **3**, e1165,
doi:10.7717/peerj.1165 (2015).
64 Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043-1055, doi:10.1101/gr.186072.114 (2015).
65 Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate
genomic comparisons that enables improved genome recovery from metagenomes
through de-replication. *ISME J* **11**, 2864-2868, doi:10.1038/ismej.2017.126 (2017).
66 Fischer, M., Strauch, B. & Renard, B. Y. Abundance estimation and differential testing on
strain level in metagenomics data. *Bioinformatics* **33**, i124-i132,
doi:10.1093/bioinformatics/btx237 (2017).
67 Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol* **34**, 525-527, doi:10.1038/nbt.3519 (2016).
68 Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. *Bioinformatics*,
doi:10.1093/bioinformatics/btz848 (2019).
69 Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-
2069, doi:10.1093/bioinformatics/btu153 (2014).
70 Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and
adaptive score threshold. *Bioinformatics* **36**, 2251-2252,
doi:10.1093/bioinformatics/btz859 (2020).
71 Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J Antimicrob*
Chemother **67**, 2640-2644, doi:10.1093/jac/dks261 (2012).
72 Liu, B., Zheng, D., Jin, Q., Chen, L. & Yang, J. VFDB 2019: a comparative pathogenomic
platform with an interactive web interface. *Nucleic Acids Res* **47**, D687-D692,
doi:10.1093/nar/gky1080 (2019).
73 Yin, Y. *et al.* dbCAN: a web resource for automated carbohydrate-active enzyme
annotation. *Nucleic Acids Res* **40**, W445-451, doi:10.1093/nar/gks479 (2012).

74   Watts, S. C., Ritchie, S. C., Inouye, M. & Holt, K. E. FastSpar: rapid and scalable
correlation estimation for compositional data. *Bioinformatics* **35**, 1064-1066,
doi:10.1093/bioinformatics/bty734 (2019).
75   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res* **13**, 2498-2504, doi:10.1101/gr.1239303
(2003).
76   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Res* **47**, W256-W259, doi:10.1093/nar/gkz239 (2019).
77   Pasolli, E. *et al.* Accessible, curated metagenomic data through ExperimentHub. *Nat*
Methods **14**, 1023-1024, doi:10.1038/nmeth.4468 (2017).

Fig.1 Reversible changes of gut microbiota associates with reversible shifts of metabolic phenotypes in
patients with T2DM. (A) Study design. Before Run-in, written informed consent, questionnaire of personal
information and measuring HbA1c at screening. After Run-in, medical checkup and sample collection at baseline
(M0), three months after on the high fiber intervention or usual diet (M3) and one year after the high fiber
intervention stopped (M15). (B) Changes of fiber intake. (C) Global changes of the gut microbiome as shown by the
principal coordinate analysis based on the Bray-Curtis distance for the 1845 genomes and (D) Average Bray-Curtis
distance between the groups. PERMANOVA test (9,999 permutations) was performed to compare the groups. * P <
0.05 and *** P < 0.001. The color of the square showed the magnitude of average Bray-Curtis distance. (E) Change
of HbA1c, (F) The percentage of participants with adequate glycemic control, (G) Fasting blood glucose, and (H)
The glucose area under the curve (AUC) in a meal tolerance test (MTT). For (E), (G) and (H), data shown as percent
changes from baseline (± S.E.M). Friedman test followed by Nemenyi post-hoc test was used for comparison in the
same group, compact letters reflect significance (P < 0.05). n = 67 in W group and n = 28 in U group. Mann-
Whitney test (two-sided) was used for comparison between W and U at the same time point, * P < 0.05, ** P < 0.01
and *** P < 0.001. n = 74 in W (M0) (For panel H, n=72), n = 74 in W (M3), n = 67 in W (M15), n = 36 in U (M0),
n = 36 in U (M3) and n = 28 in U (M15).

Fig. 2. Two competing guilds of bacteria constitute a robust seesaw-like network despite the profound global
changes in the gut microbial ecosystem induced by introduction and withdrawal of the high fiber intervention.
(A) Co-abundance networks of the prevalent genomes in W group at M0, M3 and M15 during the trial, denoted as
G_{M0}(442; 4231), G_{M3}(421; 2587) and G_{M15}(429; 4592), numbers in parenthesis are order and size of the network.
The correlations between genomes were calculated using FastSpar, n = 67 patients. All significant correlations with
$P \le 0.001$ were included. Edges between nodes represent correlations. Red and blue colors indicate positive and
negative correlations, respectively. Node size indicates the average abundance of the genomes. The layout of the
nodes and edges was determined by Edge-weighted Spring Embedded Layout with correlation efficient as weight.
(B) The distribution of different types of correlations of the genome pairs during the trial. The 3 letters show the
correlations of the genome pairs at M0, M3 and M15 subsequently. Stable correlations, NNN and PPP, were
highlighted (C) Average clustering of the 141 nodes based on their robust positive and negative correlations showed
two clusters (green and purple range). The bar plots show the abundance changes of each node throughout the trial,
which is expressed as median abundance with Z-score transformation. The differences of each node over time were
tested using the Friedman test followed by Nemenyi post-hoc test. $P < 0.05$ was considered as significant. This
panel was plotted using iTOL. (D) The seesaw-like network with the 141 nodes in two polarizing clusters. Edges
between nodes represent correlations. Red and blue colors indicate positive and negative correlations, respectively.
For (C) and (D), the color of the node represents the members in the two guilds: green for Guild 1 and purple for
Guild 2.

834
Fig. 3. The balance between the two competing guilds in the seesaw-like network was associated with the
metabolic health of patients with type 2 diabetes. (A) Change of the total abundance of Guild 1, Guild 2, and
their ratio across the trial in the W group. Friedman test followed by Nemenyi test was used to analyze the
difference between time points. Compact letters reflect the significance at P < 0.05. (B) Random Forest regression
with leave-one-out cross-validation was used to explore the associations between the 141 genomes and the clinical
parameters. The bar plot shows the Pearson's correlations coefficient between the predicted and measured values.
The asterisk before the parameter's name shows the significance of the Pearson's correlations. P values were
adjusted by Benjamini & Hochberg's method. * adjusted P < 0.05, ** adjusted P < 0.01 and *** adjusted P < 0.001.
BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; WC, waist circumference; HP,
hip circumference; TNF-α, tumor necrosis factor-α; WBC, white blood cell count; CRP, C-reactive protein; LBP,
lipopolysaccharide-binding protein; TC, total cholesterol; TG, triglyceride; Lpa, lipoprotein a; HDL, high-density
lipoprotein; APOA, apolipoprotein A; LDL, low-density lipoprotein; APOB, apolipoprotein B; GFR (MDRR),
glomerular filtration rate; CysC, Cystatin C; ACR, urinary microalbumin to creatinine ratio; IMT, intima-media
thickness; DAN, diabetic autonomic neuropathy score; MHR, mean heart rate; SDNN, standard deviation of NN
intervals; SDANN, standard deviation of the average NN intervals calculated over 5 minutes; SDNNIndex, mean of
standard deviation of NN intervals for 5-minute segments; rMSSD, root-mean-square of the differences of
successive NN intervals; pNN50, percentage of the interval differences of successive NN intervals greater than 50
ms; TP, total power; VLF, very low frequency power; LF, low frequency power; HF, high frequency power; DPN,
diabetic peripheral neuropathy score. (C) Differences in genetic capacity of carbohydrate substrate utilization
(CAZy), short-chain fatty acid production (SCFA), number of antibiotic resistance genes (ARG) and number of
virulence factor genes (VF). (C) The heatmaps show the proportion (CAZy) or gene copy numbers (SCFA, ARG
and VF) of each category in each genome. For carbohydrate substrate utilization, CAZy genes were predicted in
each genome. The proportion of CAZy genes for a particular substrate was calculated as the number of the CAZy
genes involved in its utilization divided by the total number of the CAZy genes. Arabinoxylan-related CAZy
families: CE1, CE2, CE4, CE6, CE7, GH10, GH11, GH115, GH43, GH51, GH67, GH3 and GH5; cellulose-related:

GH1, GH44, GH48, GH8, GH9, GH3 and GH5; inulin-related: GH32 and GH91; mucin-related families: GH1, GH2,
GH3, GH4, GH18, GH19, GH20, GH29, GH33, GH38, GH58, GH79, GH84, GH85, GH88, GH89, GH92, GH95,
GH98, GH99, GH101, GH105, GH109, GH110, GH113, PL6, PL8, PL12, PL13 and PL21;pectin-related: CE12,
CE8, GH28, PL1 and PL9; starch-related: GH13, GH31 and GH97. For short chain fatty acid production, FTHFS:
formate-tetrahydrofolate ligase for acetate production; ScpC: propionyl-CoA succinate-CoA transferase and Pct:
propionate-CoA transferase for propionate production; But: Butyryl–coenzyme A (butyryl-CoA): acetate CoA
transferase, Buk: butyrate kinase, 4Hbt: butyryl- CoA: 4-hydroxybutyrate CoA transferase, Ato: butyryl-
CoA:acetoacetate CoA transferase (AtoA: alpha subunit, AtoD: beta subunit) for butyrate production. Mann-
Whitney test (two-sided) was used to analyze the difference between Guild 1 and Guild 2. # $P < 0.1$, * $P < 0.05$, **
$P < 0.01$ and *** $P < 0.001$. Guild 1 (green bar): n = 50, Guild 2 (purple bar): n = 91.

885
886
887
Fig.4. The seesaw networked microbiome signature exists in other independent human cohorts and supports
classification models for different diseases. (A) Members of the two competing Guilds in the seesaw networked
microbiome signature showed similar ecological interactions in four independent human gut metagenomic datasets.
The correlations between the genomes were calculated using FastSpar. All significant correlations (P ≤ 0.001) were
included. Lines between nodes represent correlations, and red and blue colors indicate positive and negative
correlations, respectively. The color of the node represents the members in the two competing guilds: green for
Guild 1 and purple for Guild 2. The percentage of correlations followed the pattern in the seesaw networked
microbiome signature (i.e., positive edges within each guild, negative edges between the 2 guilds) was in yellow,
and the ratio of correlations that were negative within each guild and positive between the guilds was in black of the
100% stacked bar. (B) The composition of the microbiome signature was different between control and patients in
each dataset as shown in the Principal Coordinates Analysis plot based on Bray-Curtis distance. 95% confidence
ellipses were projected for control and patients respectively. The p values of the PERMANOVA test were indicated.
(C) The microbiome signature supports classification models for the four different diseases. The area under the ROC
curve (AUC) of the Random Forest classifier based on the 141 genomes in the microbiome signature to classify
control and patients in each dataset. Leave-one-out cross validation was applied. Type 2 diabetes (T2DM): Control n
= 136, T2D n=136; Atherosclerotic cardiovascular disease (ACVD): Control n = 171 and ACVD n = 214; Liver
cirrhosis (LC): control n = 83 and LC n =84; Ankylosing spondylitis: Control n = 83 and AS n = 97.
905