Original Article

Title:

Relationships of Worldwide *Onobrychis* species: from seed & pod morphology and molecular
perspective

Authors:

Mohammad Zarrabian[1, 2], Azam Nikbakht Dehkordi[3], Mohammad Hossein Ehtemam[1], Mohammad Mahdi
Majidi[1]

Affiliations:
1) Department of Plant Production and Genetics, College of Agriculture, Isfahan University of
Technology, Isfahan, Iran.
2) Corresponding author (Email: Moh8340@Gmail.com)
3) Research and Technology Institute of Plant Production (RTIPP), Shahid Bahonar University of
Kerman, Kerman, Iran

Highlight:

This paper discusses a worldwide phylogenic analysis and population structures of *Onobrychis* genus.
The results of this paper provide a comprehensive answer to complexity of subgenus and section
discrimination.

Abstract

The present study aimed to investigate the phylogeny of 31 *Onobrychis* species with the help of
seed and fruit macro & micro-morphological traits and ISSR molecular markers. The species were found
to be highly variable in their seed and fruit morphological traits. Also, a new type of fruit (*O. crista* galli
type), in addition to the three predefined types of fruit, was introduced. From molecular perspective, a set
of 22 ISSR primers were shown to have a good efficiency for genetic discrimination among the species.
Bayesian model based STRUCTURE analysis grouped the genotypes into 4 genetic clusters, and the high
value of F_{st} calculated for these clusters indicated a high level of genetic differentiation among them. On
the other hand, based on the Maximum Likelihood phylogenetic tree, our results are in agreement with the
subgenus classification (*Onobrychis* and *Sisyrosema*), while contrary to *Sisyrosema*, the *Onobrychis*
subgenus (the *Lophobrychis* and *Onobrychis* sections) appears not to be monophyletic. Furthermore, in
contrast to the traditional taxonomic classification, species belonging to the *Lophobrychis* section do not
form a coherent group, indicating that this section could be considered as a heterogeneous unit and
perhaps could be merge into the *Onobrychis* section.

Key words: ISSR, *Onobrychis*, Phylogeny, Seed coat, Species, SEM

٤٥
٤٦
Introduction

The genus *Onobrychis Mill.,* which belongs to the Fabaceae family, includes about 170
species around the world. This genus is widely distributed in northern temperate regions of the
world whereas Iran and Turkey are considered to be the center of diversity for this species.
(Rechinger, 1969; Yildiz *et al*., 1999; Hayot-Carbonero, 2011). This genus is divided into two
subgenera: *Onobrychis* and *Sisyrosema,* which are distinguished from each other through
deciduous flowers, glabrous corollas, erect fruits, and the epidermis of a calyx with crystals
within (Yildiz *et al*., 1999). Moreover, subgenus *Onobrychis* includes four sections
(*Dendrobrychis* DC, *Lophobrychis*, *Onobrychis*, and *Laxiflorae*) while the subgenus *Sisyrosema*
comprises five sections (*Anthyllium*, *Afghanicae*, *Heliobrychis*, *Hymenobrychis*, and *Insignes*)
(Rechinger, 1969).
Up until now, several approaches have been adopted to provide an appropriate
classification of the *Onobrychis* genus. Yildiz *et al*. (1999) recognized 170 species based on
fruit morphological characters and re-evaluated the sectional delimitation of the *Onobrychis*
genus. However, later studies suggested that it would be necessary to re-examine the taxonomic
position of the *Onobrychis* taxa (Cenci *et al*., 2000; Abou-el-Enain, 2002). A karyological study
within *Onobrychis* clade found that sub-genus *Sisyrosema*, unlike subgenus *Onobrychis*, is
monophyletic (Hejazi *et al*., 2010). Arslan and Ertugrul (2010) maintained that a greater
similarity could be established on the basis of seed storage protein between the *Heliobrychis* and
Hymenobrychis sections rather than the *Onobrychis* section.
Unlike studies on protein and characteristics of seeds, DNA fingerprinting revealed that a
classification of the *Onobrychis* genus weakly supported the taxonomical sections and many
species would be synonyms (Hayot-Carbonero, 2011; Safaei Chaei Kar *et al*., 2014; Duan *et al*.,
2015; Zarrabian and Majidi, 2015; Amirahmadi *et al*., 2016). In the previous nrDNA ITS-based

phylogenetic studies (Ahangarian *et al*., 2007; Safaei Chaei Kar *et al*., 2012), the non-monophyly
of the *Onobrychis* subgenus indicated the subgenus *Sisyrosema* was retained as monophyletic.
On the other hand, Amirahmadi *et al.* (2014) suggests that none of these two subgenera is
monophyletic. More studies based on three non-coding chloroplast sequences and the nuclear
ribosomal DNA internal transcribed spacer revealed that section *Heliobrychis* was retrieved as a
well-supported monophyletic group sister to *Hymenobrychis* (Kaveh *et al*. 2018). It seems that
the challenges and uncertainties are the results of the multitude of recognized species and the
high variability of morphological characters due to their wide geographical dispersion (Zarrabian
et al., 2013). To expand our knowledge on classification and genetic structure of the *Onobrychis*
genus, the present study was conducted with the following objectives: 1) To illustrate the
structures of the fruit (legume) surface and the external morphology of mature seeds, 2) To
evaluate the diagnostic value of the investigated characters in terms of their systematic
implications, and 3) To clarify phylogenetic relationships within the genera.
٨٤
Methods and martial
Plant material
Eighty-eight accessions belonging to 31 species of the genus *Onobrychis* were used in
this study (Table 1). The germplasms were obtained from the Leibniz Institute of Plant Genetics
and Crop Plant Research (IPK), United States Department of Agriculture (USDA), and some
species were collected from different geographical regions nationwide by the authors. All the
accessions used for molecular experiments were germinated and grown at 25 °C with a 16/8 h
day/night cycle in the greenhouse.

Seed Morphological analysis
For seed Macro-morphological observations, seven seeds and legumes were placed under
a binocular stereoscopic microscope (Osk 1844-b) (Table S1&S2). Moreover, fifteen seeds and
legumes of each accession were measured for dimensions using a Digital Caliper (Osk 9888) and
their average was used for each species (Table S1 and Fig. S1). Also, mature seeds of the species
under investigation were selected for SEM study. For this purpose, the seeds were mounted on
stubs using a double-face carbon tape and covered with gold (Au) at 15 mA for 3 min under high
vacuum in an Ion Sputter Coating Unit (Baltek SCD-005). The samples were subsequently
detected and photographed using SEM (Philips XL 130). The terminology of Botanical Latin
Stearn (Stearn, 1983; Barthlott,1984; and Punt et al.,1994) was used to describe the SEM aspects
of seed coat ornamentations (Figure S2&S3). For the cross-section of seed coats, three seeds
were randomly selected from each accession, and the seed coats were separated before they were
soaked in distilled water for 45 min. For convenience and uniformity of the cuts, a razor blade
was used for making cross-sections. Transverse sections were cleaned with sodium hypochlorite
for 1 min and slides were examined under a Nikon ECLIPSE E600 microscope (×400). For the
seed coat cross-sections, four traits were investigated including the Palisade layer (or Malpighian
cells), Osteosclereids, inner Parenchyma tissue, and total seed coat thickness was measured with
a Zoom browser ex. Ink program (Table S1&S2; Fig. S4).
ISSR analysis
For ISSR analysis, 31 species were planted and young leaves (0.5 g) from each accession
were chosen. Total genomic DNA was extracted according to the Murray and Thomson method
(1980) and the extracts from all the accessions belonging to one species were mixed. Moreover,
the quality and quantity of DNA were checked by agarose gel (0.7%) electrophoresis. Of the 45
ISSR primers screened, 22 produced a higher number of reproducible bands, which were

selected for ISSR analysis (Table 2). The ISSR reaction mixture (total volume = 15 ml)
contained 20 ng total DNA, 1.5 mM 10x PCR buffer, 1.5 mM MgCl$_2$, 0.3 mM dNTP, 2 pM of
each primer, and 1 U Taq DNA polymerase. Amplifications were performed using a Bio-Rad
Thermocycler (58BR 08334) programmed for 4 min at 94 ℃ as the initial denaturation
temperature followed by 35 cycles at 94 ℃ for 1 min, 45 S at an appropriate annealing
temperature (Table 2), 2 min at 72 ℃, and 7 min at 72 ℃ as the last synthesis step. Amplified
DNA fragments were separated in a 1.5% agarose gel at 100 W for 2 h in 1× TBE buffer (100
mM Tris–Borate, pH 8.0, 2 mM EDTA) and stained with ethidium bromide.
١٢٥
١٢٦ **Statistical analysis**
١٢٧
The fruit and seed morphological and anatomical characteristics were coded between 0
(absent) and 1 (present) to create the data matrix for computation (Table S2). For molecular
analysis, only the sharp, distinct, and well-resolved fragments were scored as 1 designating
present and 0 designating absent as part of a binary matrix (Table S3). Moreover, to increase the
ISSR band accuracy, the fragment without any missing data was chosen. So, out of 252 bands,
only 223 were used for downstream analysis. To calculate different parameters related to the
primer performance, the iMEC program (https://irscope.shinyapps.io/iMEC/) was used
(Amiryousefi *et al*., 2018). These parameters are Information Content (PIC), Resolving Power
(Rp), Discriminating Power (Dp), expected heterozygosity (H), arithmetic mean heterozygosity
(H.avp), and Effective multiplex ratio (E) (Table 2). Moreover, genetic similarity among all the
species was calculated according to the Jaccard similarity index with NTSYS (ver. 2.02) (Rohlf,
1998) (Table S5) and principal component analysis (PCA) was carried out with NCSS software
(2021).

To measure the genetic divergence between *Onobrychis* species, Euclidean's genetic
distances matrix was used and tree were implemented in the factoextra package in R version
3.4.2 (R Project for Statistical Computing, www.r-project.org/). Furthermore, to better show the
relationships between the *Onobrychis* species, both seed morphological traits and ISSR marker
data were combined. The phylogenic tree-based of joint data was carried out with Maximum
Likelihood methods with 1000 bootstrap in MEGA version 6.10 for Windows (Tamura *et al*.,
2013). In this study, parameters related to the genetic structure were evaluated by the POPGENE
program (ver. 1.32) (Yeh *et al*., 1999). In addition, Arlequin (ver. 3.5.2.2) (Excoffier and
Lischer, 2010) was used to investigate the diversity between subgroups and sections. For genetic
structure and admixture detection, a Bayesian model clustering algorithm by STRUCTURE,
version 2.3.4 was performed (Pritchard *et al*., 2010). For estimation of the most likely number of
genetic groups (K), the output files of structure analysis were compressed into a single file then
uploaded online using Structure Harvester (ver. 0.6.93) (http://taylor0.biology.ucla.edu/structure)
(Earl., 2012).
١٥٥
Result
Seed and fruit Morphological Study
Seed and fruit morphological characteristics of the species are shown in Tables S1&S2.
The pod was acanaceous in all the species, except in *O. pulchella*. On the other hand, the only
species with short teeth on spines (component spines) was *O. crista-galli* while other species had
simple spines (Table S2). Among the members of the *Onobrychis* genus studied, only two
species (*O. hajastana* and *O. crista-galli*) had thorns with triangular cross-sections on the disc

while other species had conical or both conical and triangular cross section spines on disc (Table
S2 and Fig. S1). The marginal spine cross-sections in all the species were conical except those
belonging to the *Lophobrychis* and *Onobrychis* sections. The surface of the pod in all the species
was villous except for the *O. persica*, *O. pulchella*, and *O. crista-galli* while the only species
with silky hair was *O. melanotricha* (Table S2). The result showed that all the species belonging
to the *Hymenobrychis* section had sub-marginal veins and borders. On the other hand, only two
species (*O. melanotricha* and *O. argyrea*) had no crest on fruits (Table S2&Fig. S1).
Furthermore, all the species of the *Hymenobrychis* sections had fully-curved crests. The size of
the legumes ranged between 1.49 mm (*O. pulchella*) and 2.91 mm (*O. arenaria*) in length, 9.74
mm (*O. pulchella*) and 1.98 mm (*O. arenaria*) in width, and 2.37 (*O. caput-galli*) and 1.125 (*O.
petolemaica*) in length/width ratio. Among the members of the *Onobrychis* genus studied, only
the *O. crista-galli* and species belonging to the *Hymenobrychis* section were found to have two
seeds in fruit while the other species contained a single seed in fruit (Table S2). The seed shapes
were distinguished as Trapeziform, Orbiculate, Rhombic, Reniform, and Ovate. However, the
general shape of the seeds in the four sections was reniform (Table S2).
Based on 40 species of the *Onobrychis* genus, Yildiz *et al.* (1999) suggested three
morphology types for pods (*O. radiata* type, *O. beata* type, and *O. ornata* type) in the
Onobrychis genus. However, the species in this study, especially those having the same pod
shapes as the *O. beata* type were varied in legume shape, spines density, crest length, and spine
type. Hence, based on our results, recognition of another pod type seems to be necessary for the
Onobrychis genus; we propose the *O. crista galli* type to be recognized in addition to the three
already recognized fruit types for this genus (Table S1&S2 and Fig. S1). So, a key is now
provided for distinguishing between fruit (legume) types. Type 1 (*O. beata* type) involved the
simple spine type, both triangular and conical transverse spine sections on margin and disk,
curved crest as a half circle with a short crest, and one seed. This was observed in all the species
of the *Onobrychis* section and in some of the *Lophobrychis* sections such as *O. aequidentata*, *O.
pulchella*, and *O. caput-galli* (Table S2& Fig. S1). Although *O. pulchella* was the only species
without hair and spines, the general shape was the same as type one. It may, therefore, be
categorized with type one species (Table S2 and Fig. S1). Type 2 (*O. radiata* type) consisted of
the simple spine type, conical surface spine section, conical marginal spine section, a fully-
curved crest with a long crest, marginal veins, and 2 ovulate ovate. All the species of the
Hymenobrychis section were found to have this fruit type. Type 3 (*O. crista galli* type) included
the compound spine type with a high surface spine density; long spines; glabrous, curved crest as
a half circle with a short crest, and 2 ovulate ovates. Based on our results, this type is only
observed in *O. crista-galli*. Type 4 (*O. ornate* type) contained the one with the simple spine type
with a conical surface spine section, a conical marginal spine section without a crest but with
marginal veins and borders. This was observed in the species of the *Heliobrychis* section (Table
S2).
٢٠١
Seed micro-morphological study
Light microscopy study
The general anatomy of the *Onobrychis* seed coat was found to be similar to that
described by Cenci et al. (2000) who reported the outer layer represented by radially elongated
cells to form a palisade-like (Malpighian cells) layer that constitutes the outer epidermis. Below
the Malpighian cells, there are bone-shaped and thick-walled cells, termed 'osteosclereids',
which underlie the inner tissue with thin-walled parenchyma cells which is followed by the
cuticle layer (Fig. S4). In this study, *O. pulchella* showed the highest values for the Malpighian

cells while *O. melanotricha* had the lowest values for the osteosclerosis layer. Significant
differences were also observed in the total thickness of the parenchyma layer, with *O. michauxii*
having the highest (0.88 mm) and *O. pallasii* having the lowest value (0.024 mm). The total seed
coat thickness ranged between 0.142 mm and 0.052 mm in *O. pulchella* and *O. melanotricha*,
respectively.
Scanning electron microscopy study (SEM)
The seed coat sculpture patterns were classified into six types: Reticulate (having a
network wrinkle), Undulate (having wavy wrinkles), Druse (having aggregation of irregular
wrinkles), Convolutus (having tortuous wrinkles), Flat (having flat wrinkles), and Stellate
(Asteroid elevations) (Table S1 and Fig. S2). The majority of the studied species showed the
reticulate (14 species) followed by Undulate (5 species), Convolutus (8 species), flat (2 species)
and the remaining two types each represented by only one single species (Table S2 and Fig. S2).
Cluster analysis of seed & fruit morphological data
Based on 75 seed morphological traits, the species were divided into two main groups
(Fig. 1A). Group A consisted of all the species from the *Onobrychis*, *Lophobrychis* sections
except *O. aequidentata* while the second group contains all the species from the *Hymenobrychis*
and *Heliobrychis* sections along with *O. pulchella* (Fig. 1A). As a complementary test in
grouping species with minimal data loss, we performed principal components analysis in which
the first three principal components covered 41.79% of the cumulative variation (Table S4).
Two-dimensional visual illustrations of the first three PCA are shown in Figs. 2A & S5. Based
on the first two components, two district groups were found. The first group contains all the
Onobrychis subgenera's species, despite the small distance of the *O. pulchella* due to first PC.
The second group contains all *Sisyrosema* subgenus species in which almost all the species
related to the section *Heliobrychis* were separated from the section *Hymenobrychis* according to
the second PC (Fig.2A).
٢٣٥

٢٣٦ **ISSR analysis**
٢٣٧
٢٣٨ Marker efficiency
٢٣٩
The 22 primers selected amplified 252 bands (Table S3), of which 223 (88.5%) were
polymorphic among 31 *Onobrychis* species. The amplified DNA ranged from 150 to 1400 bp.
The maximum and minimum numbers of bands were observed by the primers $(TC)_8$-C and
$(CA)_8$-RT, respectively. The percentage of polymorphic bands ranged from 75% to 100% (Table
2). The heterozygosity (H) varied from 0.423 (ISSR 16) to 0.5 for ISSR 06&10&11 with a mean
of 0.484 per primer. The polymorphism information content (PIC) value ranged from 0.33%
$((AC)8-T)$ to 0.38% $((GA)8- RT)$ with an average of 0.37%. Moreover, the marker index (MI)
with an average value of 0.008 ranged from 0.006 to 0.01 (Table 2). For resolving power index
(Rp), the lowest and highest values belonged to $(CA)8- RT$ (3.16) and $(TC)8C$ (10), respectively
(Table 2). The arithmetic means of heterozygosity (H.av) ranged between 0.001 and 0.0031 with
the mean of 0.0016 per primer. To determine the judicious profundity of primer, we calculate
discriminative power (D) with a mean index of 0.723 and extend it from 0.515 $((AC)8-T)$ to
0.855 $((TC)8G)$. The effective multiplex ratio (E), which is a conditional factor on the magnitude
of primer polymorphism, is spanned from 2.9 $((CA)8- RT)$ to 7.71 $((GA)8 SG)$ with an average
of 5.312 per primer.
٢٥٥
٢٥٦ ISSR based Genetic similarity (GS)

Genetic distances and similarity coefficients for different species (Jaccard methods) and
sections (Ni & Li method) are calculated and are shown in the Table S5 and Table S6,
respectively. The findings of the present study revealed the significant genetic differentiation
among the *Onobrychis* species which ranged from 0.199 (*O. bobrovii* vs. *O. megataphrose*) to
0.833 (*O. petraea* vas *O. inermis*) (Table S5). Moreover, from a sectional classification point of
view, the highest similarity was observed between *Onobrychis* and *Lophobrychis* sections (0.87)
while the lowest similarity was calculated for *Lophobrychis* and *Heliobrychis* (0.61) (Table S6).

ISSR Cluster analysis
٢٦٥

The cluster analysis based on ISSR marker was performed for the 31 species as shown in
Fig. 1B. The species were divided into two main groups in which all species belong to
Onobrychis and *Sisyrosema* subgenus grouped in the first and second clades, respectively (Fig.
1B); however, in the first group, all species belonging to the *Onobrychis* subgenera are mixed,
while in the second group *Hymenobrychis* and *Heliobrychis* sections are mostly separated (Fig.
1B).
       For more validation, the principal component analysis was run. The first three principal
٢٧٣   components cumulatively explained 32.36 % of all the variation (Table S4). Like the cluster
٢٧٤   analysis, we observed complete discrimination of subgenus. However, unlike cluster analysis,
٢٧٥   *Sisyrosema* sectional classification was not observed, and also, based on the first component *O.*
٢٧٦   *pallasii,* few were grouped far from their other subgenus member (Fig. 2B and Fig. S5).
٢٧٧

٢٧٨ Genetic Structure
٢٧٩

Estimates of population parameters in each subgroup are shown in Table 3. The results
showed that there was a slight difference between the calculated parameters for both subgenera.
However, the calculation of the parameter for each section showed that the greatest variation was
related to *Onobrychis* and *Hymenobrychis* sections equally (Na: 1.8, Ne: 1.5, H: 0.3, and I: 0.44)
while the lowest variation belongs to *Heliobrychis* section (Na: 1.34, Ne: 1.24, H: 0.14, and I:
0.21).
In addition, to study the structure of the existing diversity in more detail, molecular
analysis of variance (AMOVA) was used (Table 4). Based on the AMOVA result, 7.8% of the
total variation was related to the subgenus difference, while 24.51% of the diversity was related
to the differences between the sections in each subgenus and 68.41% of the variations were
related to the differences between the species in each section.
For evaluation of population structure and assigning species to specific populations based
on ISSR marker, Structure version 2.2.3 was used based on a Bayesian model. The structure
analysis was initially performed based on the maximum number of (K = 1 to 7), as the original
population order and the most probable value of population were calculated to the maximum
peak at $\Delta K = 4$ (K value =3.25; Lnprob (K) = -3669.8). The rate of change of the likelihood
distribution (mean), the absolute value of the 2nd order of the change of the likelihood
distribution (mean) for $\Delta K = 4$ are shown in Fig. 3. Based on best K = 4, it was determined that
all the evaluated species could be positioned into four major groups visualized with four distinct
colors of blue, green, yellow, and red. Each individual is represented by one vertical bar plot and
the labels below the bar plots are the corresponding numbers for each species in Table 1. More
than one color in each bar plot reveals the genetic complexity of that individual that, in this case,

the species belongs to a group that has the largest color width of that cluster. The estimated mean
F_{st} value for the species under clusters 1 and 2 was 0.2412 and 0.279, respectively while this
value for clusters 3 and 4 was 0.515 and 0.823, respectively (Table S7). Due to the membership
likelihood, described by We *et al*., (49), some of the species showed unique structure. Based on
Q > 0.60 as the purity standard, in the red zone 7 species, almost 5 species in the blue zone, and
3 species in the yellow zone shows some level of purity and the rest of the species were noted as
admixture units based on Q < 0.60 (Fig. 3B).

Seed and fruit morphological and ISSR combination analysis
To better discriminate between the *Onobrychis* species, both seed morphological traits
and ISSR marker data were combined. So, based on joint data, Jaccard's Genetic similarity
coefficients (GS) between the studied species were calculated and were in the range between
0.185 (*O. melanotricha* vs *O. persica*) and 0.806 (*O. inermis* vs *O. petraea*) (Table S5). The GS
mean value for each section was 0.5, 0.43, 0.52, and 0.49 for *Onobrychis*, *Lophobrychis*,
Heliobrychis, and *Hymenobrychis,* respectively (Table S5).
The hierarchical clustering analysis based on Joint data is shown in Fig. 4. The
dendrogram illustrated that all the 31 species were clustered into two main groups. Group A
consisted of all *Onobrychis* subgenus species in which *O. arenaria* was classified in a separate
subgroup. Furthermore, all the *Hymenobrychis* and *Heliobrychis* section species were clustered
into the second group with complete separation (Fig. 4).
To lead the clustering investigation, principal component analysis was used. The
distribution of eigenvalues, percentage of genetic variation, and cumulative percentage of genetic
variation were displayed in Table S4. The PCA analysis revealed that the share of the first three
PC is 41.77 % (PC1 = 18.85%, PC2 =13.22% and PC3 = 9.7%) of total variation. In the case of
PCA analysis from the first principal component, the highest value was for *O. megataphrose* and
O. transcaucasia (0.93 and 0.92 respectively) while the least value (-1.96) was found for *O.*
michauxii which has the least contribution to total diversity (Table S4). The first two PC bi-plots
showed the complete separation of the subgenus classification (Fig. 2C). However, as far as the
sections are concerned, this discrimination has not happened.
Discussion
Seed and pod morphological characteristics
Legume morphological characteristics, seed-coat sculpturing, and seed-coat anatomy
have been shown to provide valuable characters in the delimitation of selected species of the
Onobrychis genus. Norri and colleagues (2014) showed that morpho-biometerical characters are
useful in distinguishing among species and pod characters are the most valuable issue for
separation of the *Onobrychis* genus. A great variation in the morphological characteristics in this
study showed they could be useful tools for classifying and identifying species in the *Onobrychis*
genus. However, none of the fruit morphological traits are specific to one section or subgenus.
For example, we can refer to the fully carve crest in the fruit, which can be considered as an
indicator for the *Hymenobrychis* section, nonetheless, this indicator did not exist in the *O.*
pallasii. These traits in combination with each other might be used as an identification key for
the corresponding section or subgenre. Therefore, in addition to the different types of fruit that
were suggested by Yildiz *et al*. (1999), we also introduced a new fruit type called *O. crista* galli.
Apart from the differences between new fruit type with *O. beata* type mentioned earlier (legume
shape, spines density, crest length, and spin type), a fundamental difference between these two
types is the number of seeds in the fruit, which for *O. beata* type this number is one, while the
newly introduced type has ovary 2-ovulate (Table S2). Due to the characteristics related to fruit,

a very interesting result is the high variability in the *Lophobrychis* section. Two species in this
section had unique characteristics. The first species was *O. pulchella*, which generally has fruit
like *O. beata* type, but it did not have any thorns on the fruit or long crest length (Table S2 and
Fig. S1). The second species was *O. crista* galli, which was the only species in this study that had
a compound spin and unlike all species of *Lophobrychis* section, had the ovary with 2-ovulate
(Table S2 and Fig. S1).

For the seed transverse section, no significant variation was observed in the osteosclerotic
layer in any of the species while the difference in seed coat thickness was mainly due to
variations in the length of the Malpighian cells and the inner parenchyma tissues (data not
shown). Contrary to Cenci *et al*. (2000), who reported differences in the shape of Malpighian
cells between the perennial and annual species, we did not observe any significant differences
between them (Table S2).

For seed coat patterns, the variation observed was generally species-specific rather than
genus-specific and almost invariant across the different populations of a given species (Koul *et*
al., 2000 and Moazzeni *et al*., 2007). Our results revealed six main types of seed-coat patterns
among the investigated species (Table S2 and Fig. S3). Although it seems that the seed-coat
patterns observed in this study were variable, it was not sections or subgenus specific. Therefore,
it cannot be used as a unique trait to classify species of this genus.

Another interesting result in our study is the differentiation between annual and perennial
species in *Onobrychis* genus. Based on Flora Iranica (Rechinger, 1969), Flora of Turkey (Davis
et al. 1988), and Flora Europaea (Ball, 1968), all the species belonging to the *Lophobrychis*
section are annual while all the other sections have perennial species. However, we did not find
any differences among the annual and perennial species, neither by seed anatomical and fruit
morphological traits nor by ISSR markers (Fig. 1). Similar results have been reported by Pavlova
and Monova (2000) who found no differences in pollen morphology among the annual and
perennial species in the *Onobrychis* genus.

ISSR marker

It has been well documented that ISSR markers are highly polymorphic and are useful
tools for studying genetic diversity, phylogeny, gene tagging, genome mapping and evolutionary
biology (Pradeep *et al*., 2002). Our ISSR marker results represented a good level of
polymorphism suggesting the efficacy of these markers for the evaluation of the phylogenic
relationship among the *Onobrychis* species. There are many factors to check the efficacy of
primers, some of which have great importance. Two major indexes for the informativeness of
genomic marker polymorphism are expected heterozygosity (H) and the polymorphic
information content (PIC). PIC and H measures the ability of a marker to detect polymorphisms,
and is important in selecting markers for genetic studies (Khan *et al*., 2021). Considering, the
range of H and PIC value for a dominant marker (0 for monomorphic primer to 0.5 (for primer
with multiple alleles in an identical frequency), the value for these two indexes in studied
population was more than average, which indicated the advanced discriminatory capacity of the
marker system. The MI index, in addition to the benefits of PIC, also considers the total number
of bands and the polymorphic ratio which shows the potential of each primer to produce more
bands. Moreover, Rp measures the ability of primers to recognize and differentiate individuals.
So, all the represented values for these indexes showed the high polymorphism and efficiency of
the used primer for genetic discrimination analysis on this genus (Table 2). Overall, the greater

RP and MI indices refer to the greater efficacy of the respective primers (Zarei *et al*., 2021).
Therefore, the most effective primers are (TC)8C (Rp= 10, MI = 5.51), (GA)8- RC (Rp = 8.7,
MI= 3.77), (GA)8 WT (Rp=8.12, MI= 4.05) and (GA)8 SG (Rp= 8, MI = 4.42)

Genetic structure based on ISSR marker
The population structure of *Onobrychis* genus assessed by Bayesian admixture analysis
indicated 4 clusters (Fig. 4). Based on Q > 0.60, out of 31 *Onobrychis* species 15 were almost
pure and the other species were highly complex, indicating these species were genetically
admixture. Interestingly all the species belonging to *Lophobrychis* section (number 18, 19, 20
and 21) have different color in each bar plot which reveals their genetic complexity. Moreover,
the bar related to *O. viciifolia* had three colors (red, blue and yellow) in almost equal proportions,
which indicates the genetic complexity of this species too. Perhaps, the cross-pollinated nature of
cultivated sainfoin (*O. viciifolia*) as well as vast diversity regain make this species so complex.
For better understanding of differentiation among studied population, fixation index (Fst) was
calculated. Due to Hartl and Clark (1997), a standard scale of fixation index categorized in: little
genetic difference (< 0.05), moderate genetic difference (0.05–0.15), great genetic difference
(0.15–0.25) and very great genetic difference (> 0.25). The estimated Fst suggested the species
under cluster 1 (Fst = 0.241) showed little genetic difference while the species under cluster 2
(Fst = 0.279), 3 (Fst = 0.515) and 4 (Fst = 0.823) showed very great genetic difference. Over all,
these Fst values indicated high level of genetic differentiation with greater genetic distance
between clusters.
٤١٢
٤١٣ Sectional and Subgenus classification
٤١٤
In traditional classification, characterizing and assigning species to sections or sub-genera
seem to have often been based on morphological traits such as flower and leaf characteristic or
seeds and fruits related traits. So, some previous researchers, such as Yildiz and colleagues
(1999), by using seeds and fruits morphology claim that no evidence can be found in support of
subgenera classification while sectional discrimination was supported. On the contrary, some
phylogenic studies based on molecular markers, had completely opposite results in the
supporting subgenus classification and no verification of sectional discrimination, especially in
the subgenus *Onobrychis*. (Ahangarian *et al*., 2007; Safaei Chaei Kar *et al*., 2012) So,
considering the morphological traits of this study, even though the key was provided to identify
the fruits and seeds of each section and species, not only the subgenus, but also the sectional
discrimination for *Onobrychis* subgenus was not confirmed. These results show that usage of
morphological characters alone in phylogenetic analysis was restricted. Such restriction
happened because of the open pollination with a high allogamy rate (Avci *et al*., 2016), along
with wide geographical dispersion (Zarrabian *et al*., 2013) of the *Onobrychis* taxa. Therefore, the
morphological characters are more affected by environmental factors. Contrary to the
morphological traits, most of the molecular data supported the subgenera classification, however,
they did not support the sectional discrimination in *Onobrychis* subgenus (Ahangarian *et al*.,

2007; Chaei Kar *et al*., 2014 and Duan *et al*., 2015). So, to increase the accuracy of finding
relationships between species, especially relationships between sections and subgenus, joint data
was used and downstream analysis was conducted. Our study with sufficient species sampling
for each subgenre suggests that *Sisyrosema* is the only monophyletic subgenera. And also,
contrary to *Onobrychis* subgenre, *Sisyrosema* was divided according to its traditional sectional
classification. Such classification may not have occurred by principal component analysis for
joint data, but at least two subgenres were clearly separated from each other (Fig. 2C). Other
researchers have used seed storage protein (Arslan and Ertugrul, 2010) and sequence data of ITS,
matK, trnL-F and psbA-trnH (Duan *et al*., 2015), suggesting that the subgenus *Sisyrosema* might
be derived from the subgenus *Onobrychis*. In the nrDNA ITS-based phylogenic studies
(Ahangarian *et al*., 2007; Safaei Chaei Kar *et al*., 2012) the non-monophyly of subgenus
Onobrychis was also indicated, but subgenus *Sisyrosema* was retained as monophyletic. Also,
Hayot-Carbonero and colleagues (2012) supported the paraphyly of *Onobrychis* subgenus. They
suggested, not only from a molecular point of view, but also from morphological perspective, all
species from the *Onobrychis* section are very similar to *O. viciifolia*. They also found that *O.*
petraeais is more closely related to the *Lophobrychis* section, which our result completely agrees
with (Table S4B).
Lophobrychis section as a heterogeneous unit
Generally, the species were positioned more closely in the generated phylogenetic tree,
suggesting that they were genetically more similar or even have identical genes. On the other
hand, species possessing the distant group suggest that they were genetically dissimilar, even
though they come from the same origin (Khan *et al*., 2021). Such a trend has not been observed
for the species related to *Lophobrychis* section. In this study, no independent clade containing all
species of the *Lophobrychis* section was found, neither through molecular markers nor with
morphological traits. Therefore, in contrast to the traditional taxonomic classification, species
belonging to the *Lophobrychis* section were not coherently classified in one group (Fig.1 and
Fig. 4), indicating the *Lophobrychis* section not only has a comparatively highly derived
organization, but also can be considered as a heterogeneous unit in the *Onobrychis* genus (Abou-
el-Enain, 2002; 2004). Although a previous study (Safaei Chaei Kar *et al*., 2014) showed
Onobrychis and *Lophobrychis* sections were separated, the high similarity found in this study
between *Lophobrychis* and other sections, especially the *Onobrychis* (Fig. S6 and Table S5),
gives rise to the probability that the concept of section applied to *Lophobrychis* is meaningless or
unrealistic. Bandara *et al*. (2013), based on ITS sequences, supported the sectional treatment and
may require some changes. They assumed that, in order to form a monophyletic clade
corresponding to a section, part of *Lophobrychis*, *Dendrobrychis* and the subsection
Macropterae of *Onobrychis* section should be kept together. Hayot-Carbonero *et al*. (2012)
declared the *Lophobrychis* and *Onobrychis* sections were less clearly distinct. By looking at
similarity matrix based on the joint data (Table S5), O. *caput-galli* had a higher similarity to the
species belonging to *Onobrychis* section rather than the rest of *Lophobrychis* species. Elena
(2006) reported that *O. caput-galli* was cyto-taxonomically closer to the section *Onobrychis*

rather than other species of its own section. Hence, it seems that this species could be the
connecting link between the *Onobrychis* and *Lophobrychis* sections of this genus. Overall, there
is a general difficulty in discrimination of the *Lophobrychis* and *Onobrychis* sections. However,
it is possible that these two sections have a relatively recent common ancestor (Hayot-Carbonero
et al., 2012).

Conclusion
Investigations of seed macro- and micro-morphology of the genus *Onobrychis* showed
that it could be a useful tool (especially macro-morphological traits) for differentiating species.
Hence, these data alone are insufficient for phylogenic analysis. Such conflict may be due to the
fact that species were assigned to different sections based on only one or few morphological
characters and some characters may have converged in the course of evolutionary history (Duan
et al., 2015). The results obtained from this study support the subgenus (*Onobrychis* and
Sisyrosema) classification while the monophyly of *Onobrychis* subgenus was not supported.
Moreover, it seems that the section *Lophobrychis* has a comparatively highly derived
organization and can be considered as a heterogeneous unit in the *Onobrychis* genus. Our results
also showed that combination of seed macro- and micro-morphological traits with ISSR data
provide complementary information for the *Onobrychis* genus classification. However, more
study with larger number of molecular markers will be needed to obtain an appropriate
phylogenetic classification.

Supplementary data:

The following supplementary data are available at *JXB* online.

Fig. S1. : Fruit shape of different *Onobrychis* species.

Fig. S2: Transverse sections of *Onobrychis* species seed coats.

Fig. S3. Technical shape which discripe the seed surface pattern

Fig. S4: Scanning electron microscopy photographs of seed coat

Fig. S5: Two-dimensional representation of genetic relationships based on principal components analysis

Table S1. Seed and fruit macro and micro morphological characters.

Table S2: Data matrix of the fruit and seed characteristics of *Onobrychis* species in this study.

Table S3: ISSR Data matrix of species in this study.

Table S4 A: Jaccard's Similarity matrix between the 31 *Onobrychis* species.

TableS5: Nei's genetic identity and genetic distance for each sections based on ISSR.

Table S6: The eigenvalues, % of variation and cumulative % of variation of 30 PCs and PCoA

Table S7: species contribution in each component in different experiment

Table S8: Inferred ancestry of species and Fst value for each cluster

Acknowledgements:

I would like to thank Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), and
United States Department of Agriculture (USDA) for providing plant material

Author contributions

MZ designed, analyzed and preformed the experiments; MZ, AND, MHE, MMM wrote the paper.

Conflict of interest:

The authors declare no conflicts of interest.

Funding:

This work was supported by funding through the Isfahan University of Technology, college of agriculture
engineering, Department of production, and plant genetics Graduate Student Fellowship

Data availability:

The raw data used in this study are available from the corresponding author upon request.

References

1) Abou-El-Enain MM. 2002. Chromosomal criteria and their phylogenetic implications in the genus Onobrychis Mill, Sect. Lophobrychis (Leguminosae) with special reference to Egyptian species. Botanical Journal of the Linnean Society 139: 409- 414.

2) Abou-El-Enain MM. 2004. SDS-PAGE of seed protein criteria in relation to taxonomy of *Onobrychis* sect. *Lophobrychis* s. str. and the Egyptian species. Cytologia 69: 351-358.

3) Al-Ghamdi FA, Al-Zahrani RM 2012. Seed morphology of some species of Tephrosia Pers. (Fabaceae) from Saudi Arabia identification of species and systematic significance. Feddes Repertorium 121: 59–65.

4) Amirahmadi A, Kazempour-Osaloo S, Kaveh A, Maassoumi AA, Naderi R. 2016. The phylogeny and new classification of the genus *Onobrychis* (*Fabaceae-Hedysareae*): evidence from molecular data. *Plant* Systematics and *Evolution* 302:1445–1456.

5) Arslan E, Ertugrul K. 2010. Genetic relationships of the genera Onobrychis, Hedysarum, and Sartoria using seed storage proteins. Turkish Journal of Biology 34: 67-73.

6) Attar F, Keshvari A, Ghahreman A, Zarre S, Aghabeigi F. 2007. Micromorphological studies on Verbascum (Scrophulariaceae) in Iran with emphasis on seed surface, capsule ornamentation and trichomes. Flora 202: 169–175.

7) Avcı A, Ilhan E, Erayman E, Sancak C 2014. Analysis of Onobrychis genetic diversity using SSR markers from related legume species. The Journal of Animal & Plant Sciences 24(2): 556-566

8) Avci S, Tekin N, Sancak C, Özcan S, Marang AO. 2016. Phylogenetic relationship of some Onobrychis taxa naturally grown in Turkey based on morphology and nuclear ribosomal DNA ITS Sequences. Legume Research 39(5): 665-67.

9) Ball PW 1968. Onobrychis Miller. In: Tutin TG, Heywood VH, Burges NA, Moore DM, Valentin SM (ed) Flora Europaea, Vol 2. Cambridge: Cambridge University, pp 187-191.

10) Bandara NL, Papini A, Mosti S, Brown T, Smith LMJ. 2013. A phylogenetic analysis of genus *Onobrychis* and its relationships within the tribe Hedysareae (Fabaceae). Turk J Bot 37: 981-992.

11) Barthlott W (1984). Microstructural features of seed surfaces. In: Heywood VH, Moore DM (Ed) Current concepts in plant taxonomy. London.

12) Bhattarai S, Coulman B, Fu YB, Beattie AD, Biligetu B 2018. Genetic diversity and relationship of sainfoin (*Onobrychis viciifolia* Scop.) germplasm as revealed by amplified fragment length polymorphism markers. Canadian Journal of Plant Science 98: 543–551

13) Cenci CA, Bassi G, Ferranti E, Romano B. 2000. Some morphometric, anatomical and biochemical characteristics of fruits and seeds of *Onobrychis* spp. in Italy. Plant Biosystems 134: 91-98.

14) Davis PH, Mill RR, Tan K. 1988. Flora of Turkey and the East Aegean Islands (Supplement), Vol 10. Edinburgh: Edinburgh University, pp 561-588.

15) Duan L, Wen J, Yang X, Liu PL, Arslan E, Ertuğrul K, Chang ZY. 2015. Phylogeny of *Hedysarum* and tribe Hedysareae (Leguminosae: Papilionoideae) inferred from sequence data of ITS, matK, trnL-F and psbA-trnH. Taxon 64 (1):49–64.

16) Earl DA. 2012. STRUCTURE HARVESTER: A website and program for visualizing STRUCTURE output and implementing the Evanno method. Conservation Genetics Resources 4(2), 359–361.

17) Elena T. 2006. Cytological aspects of the Onobrychis genus. Buletin USAMV 62: 154-158.

18) Emre I, Turgut-bahk D, Sahm A, Kursat M. 2007. Total Electrophoretic Band Patterns of Some Onobrychis Species Growing in Turkey. Journal of *Agriculture* and *Environmental* Sciences 2: 123-126.

19) Excoffier L, Lischer HEL. 2010. Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. Molecular Ecology Resources 10: 564-567.

20) Gontcharova SB, Gontcharov AA, Yakubov VV, Kondo K. 2009. Seed surface morphology in some representatives of the Genus Rhodiola sect. Rhodiola (Crassulaceae) in the Russian Far East. Flora 204: 17–24.

21) Hartl, D. L., Clark, A. G., & Clark, A. G. (1997). Principles of Population Genetics. Vol. 116. (Sinauer Associates,).

22) Hayot-Carbonero C. 2011. Sainfoin (Onobrychis viciifolia), a forage legume with great potential for sustainable agriculture, an insight on its morphological, agronomical, cytological and genetic characterization. PhD Thesis,, Univercity of Manchester, Manchester, U.K.

23) Hayot-Carbonero C, Carbonero F, Smith LMJ, Brown TA. 2012. Phylogenetic characterisation of Onobrychis species with special focus on the forage crop Onobrychis viciifolia Scop. Genetic Resources and Crop Evolution 59: 1777-1788.

24) Johnson LA, Huish KH, Porter JM. 2004. Seed surface sculpturing and its systematic significance in Gilia (Polemoniaceae) and segregated genera. International Journal of Plant Sciences 165: 153–172.

25) Karcz J, Ksiazazyk T, Maluszynska J. 2005. Seed coat patterns in rapid-cycling Brassica forms. Acta Biologica Cracoviensia s. Botanica 47: 159–165.

26) Khan MMH, Rafii MY, Ramlee SI, Josoh M, Mamum DA, Halidu J. 2021. DNA fingerprinting, fixation-index (Fst), and admixture mapping of selected Bambara groundnut (Vigna subterranea [L.] Verdc.) accessions using ISSR markers system. Scientific Reports 11:14527. https://doi.org/10.1038/s41598-021-93867-5

27) Koul KK, Nagpal R, Raina SN. 2000. Seed Coat Microsculpturing in Brassica and Allied Genera (Subtribes Brassicinae, Raphaninae, Moricandiinae). Annals of Botany 86: 385-397.

28) Moazzeni H, Zarre S, Al-Shehbaz IA, Mummenhoff K. 2007. Seed-coat microsculpturing and its systematic application in Isatis (Brassicaceae) and allied genera in Iran. Flora 202: 447–454.

29) Murray MG, Thompson WF. 1980. Rapid isolation of high molecular weight plant DNA. Nucleic Acids Research 8: 4321-4325.

30) Pavlova DK, Manova VI. 2000. Pollen morphology of the genera Onobrychis and Hedysarum (Hedysarea, Fabaceae) in Bulgaria. Annales Botanici Fennici 37: 207-217.

31) Powell W, Morgante M, Ander C, Hanafey M, Vogel J, Tingy S, Rafalaski A. 1996. The comparision of RFLP, RAPD, AFLP and SSR (microsatellite) marker for germplasm analysis. Molecular Breeding 2: 225-238.

32) Pradeep Reddy M, Sarla N, Siddiq E. 2002. Inter simple sequence repeat (ISSR) polymorphism and its application in plant breeding. Euphytica 128: 9–17. https://doi.org/10.1023/A:1020691618797

33) Prevost A, Wilkinson MJ. 1999. A new system of comparing PCR primers applied to ISSR fingerprinting of potato cultivars. Theoretical *and* Applied Genetics 98: 107-112.

34) Pritchard JK, Wen W, Falush D. 2010. Documentation for STRUCTURE Software: Version 2 (University of Chicago).

35) Punt W, Blackmore S, Nilsson S, Thomas A. 1994. Glossary of pollen and spore terminology. LPP Foundation, Utrecht.

36) Rechinger KH. 1969. Flora Iranica. In: Reshinger KH (ed). Vol. 157. Akademische Druck- und Verlgsantalt, Graz, Austria. p. 387.

37) Rohlf FJ (1998). NTSYS-pc numerical taxonomy and multivariate analysis system. Version 2.00. Exeter Software, Setauket, NY.

38) Roldan-Ruiz I, Dendauw J, Bockstaele EV, Depicker A, Loose MD. 2000. AFLP markers reveal high polymorphic rates in ryegrasses (*Lolium* spp.). Molecular Breeding 6: 125-134.

39) Safaei Chaei Kar S, Ghanavati F, Naghavi MR, Amirabadi-zade H, Rabiee R. 2014. Molecular phylogenetics of the Onobrychis genus (Fabaceae: Papilionoideae) using ITS and trnL–trnF DNA sequence data. *A*ustralian Journal of Botany 62: 235–250.

40) Stearn WT (1983). Botanical Latian. Nelson. London.

41) Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. Molecular Biology and Evolution 30(12): 2725–2729.

42) Toluei Z, Atri M, Ranjbar M, Wink M. 2013. Iranian Onobrychis carduchorum (Fabaceae) populations: morphology, ecology and phylogeography.   Plant Ecology *and* Evolution 146: 53–67

43) Vural C, Ekici M, Akan H, Aytac Z. 2008. Seed morphology and its systematic implications for genus Astragalus L. sections Onobrychoidei DC., Uliginosi Gray and Ornithopodium Bunge (Fabaceae). Plant Systematics and Evolution 274: 255–263 .

44) Yeh FC, Yang R, Boyle T. 1999. Popgene version 1.32: Microsoft Windows-based freeware for population genetic analysis. University of Alberta, Edmonton.

45) Yildiz B, Ciplak B, Aktoklu E. 1999. Fruit morphology of sections of the genus Onobrychis Miller (Fabaceae) and its phylogenetic implications. Israel Journal of Plant Sciences 47: 269–282.

46) Zarei A, Erfani-Moghadam J. 2021. SCoT markers provide insight into the genetic diversity, population structure and phylogenetic relationships among three Pistacia species of Iran. Genetic Resources and Crop Evolution. 68(4), 1625–1643.

47) Zarrabian M, Majidi M, Ehtemam MH. 2013. Genetic Diversity in a Worldwide Collection of Sainfoin Using Morphological, Anatomical, and Molecular Markers. Crop Science 53: 1–14.

48) Zarrabian, M, Majidi, M.M. (2015). Genetic diversity and relationships within and among *Onobrychis* species using molecular markers. Turkish Journal of Botany. 39: 681–692.

Num.	Sub-genus	Section	Species	Number	Origin
					Table 1. Information of species and accessions investigated in this study
1	*Onobrychis*	*Onobrychis*	*O. transcaucasia*	9	Armenia, Iran, Turkey, Georgia and Uzbekistan
2	*Onobrychis*	*Onobrychis*	*O. arenaria*	10	Soviet Union, Russia, Romania and Azerbaijani
3	*Onobrychis*	*Onobrychis*	*O. iberica*	3	Soviet Union and Pakistan
4	*Onobrychis*	*Onobrychis*	*O. cyri-* Grossh	2	Russia
5	*Onobrychis*	*Onobrychis*	*O. altissima*	7	Soviet Union, Russia, Azerbaijani, Georgia and Iran
6	*Onobrychis*	*Onobrychis*	*O. alba*	3	Bulgaria and Soviet Union
7	*Onobrychis*	*Onobrychis*	*O. inermis*	4	Russia
8	*Onobrychis*	*Onobrychis*	*O. petraea*	5	Germany, Iran and Russia
9	*Onobrychis*	*Onobrychis*	*O. oxyodonta*	1	Soviet Union
10	*Onobrychis*	*Onobrychis*	*O. gracilis*	1	Bulgaria
11	*Onobrychis*	*Onobrychis*	*O. persica*	1	Iran
12	*Onobrychis*	*Onobrychis*	*O. peduncularis*	1	Spain
13	*Onobrychis*	*Onobrychis*	*O. hajastana*	1	Soviet Union
14	*Onobrychis*	*Onobrychis*	*O. megataphrose*	1	Turkey
15	*Onobrychis*	*Onobrychis*	*O. montana*	1	France
16	*Onobrychis*	*Onobrychis*	*O. viciifolia*	10	Iran, China, USA, Czech Republic, Kyrgyz Republic, Spain, Ukraine, England, Morocco, Russia
17	*Onobrychis*	*Onobrychis*	*O. kemulariae*	1	Soviet Union
18	*Onobrychis*	*Lophobrychis*	*O. caput-galli*	3	Turkey, Israel and Unknown
19	*Onobrychis*	*Lophobrychis*	*O. crista-galli*	4	Iran, Israel and Unknown
20	*Onobrychis*	*Lophobrychis*	*O. aequidentata*	3	France
21	*Onobrychis*	*Lophobrychis*	*O. pulchella*	1	Turkmenistan
22	*Sisyrosema*	*Heliobrychis*	*O. melanotricha*	2	Iran
23	*Sisyrosema*	*Heliobrychis*	*O. argyrea*	1	Turkey
24	*Sisyrosema*	*Hymenobrychis*	*O. ptolemaica*	2	Iraq and unknown
25	*Sisyrosema*	*Hymenobrychis*	*O. hypargyrea*	2	Turkey
26	*Sisyrosema*	*Hymenobrychis*	*O. michauxii*	2	Turkey and Iran
27	*Sisyrosema*	*Hymenobrychis*	*O. chorassanica*	1	Soviet Union
28	*Sisyrosema*	*Hymenobrychis*	*O. sintenisii*	3	Soviet Union and Iran
29	*Sisyrosema*	*Hymenobrychis*	*O. bobrovii* Grossh	1	Russia
30	*Sisyrosema*	*Hymenobrychis*	*O. pallasii*	1	Soviet Union
31	*Sisyrosema*	*Hymenobrychis*	*O. radiata*	1	Armenia

Num.	Sequence (3'-5')	T_a	NB	NPB	PP%	H	PIC	E	H.av	MI	D	Rp
Table 2: ISSR primers information used in this study.												
1	(CA)8G	52	14	11	0.79	0.494	0.372	6.097	0.0014	3.23	0.694	7.419
2	(TC)8C	56	15	15	1.00	0.484	0.367	6.161	0.0010	5.51	0.832	10.000
3	(TC)8G	54	14	12	0.86	0.472	0.361	4.581	0.0013	3.73	0.855	7.548
4	(AC)8 G	48	10	9	0.90	0.499	0.375	4.355	0.0018	3.04	0.767	7.032
5	(CA)8- RT	46	7	5	0.71	0.487	0.368	2.903	0.0031	1.31	0.664	3.161
6	(GA)8- RT	51	14	11	0.79	0.500	0.375	5.452	0.0015	3.26	0.755	7.097
7	(AC)7 DBD	50	12	10	0.83	0.475	0.362	6.742	0.0014	3.66	0.625	7.226
8	(AG)7C	52	9	9	1.00	0.487	0.368	3.774	0.0017	3.31	0.825	4.000
9	(GA)8SC	57	12	11	0.92	0.494	0.372	6.097	0.0014	3.76	0.694	7.613
10	(AC)8C	48	8	7	0.88	0.500	0.375	3.548	0.0023	2.31	0.744	4.258
11	(AG)8 SG	56	10	10	1.00	0.500	0.374	4.903	0.0016	3.74	0.760	6.387
12	(GA)8 SG	58	14	12	0.86	0.483	0.366	7.710	0.0012	4.42	0.649	8.065
13	(GA)8 WT	47	11	11	1.00	0.486	0.368	6.419	0.0014	4.05	0.660	8.129
14	(CT)8-RG	51	10	9	0.9	0.487	0.368	4.194	0.0016	3.68	0.825	5.548
15	(GA)8-C	50	12	10	0.83	0.437	0.342	6.774	0.0014	2.84	0.542	5.677
16	(AC)8-T	54	13	10	0.77	0.423	0.333	6.968	0.0014	2.56	0.515	5.871
17	(GA)8-YT	52	11	10	0.91	0.490	0.370	5.710	0.0016	3.37	0.675	7.032
18	(GA)8-YC	54	12	11	0.92	0.473	0.361	4.226	0.0014	3.65	0.853	6.968
19	(AG)8-YT	54	10	10	1.00	0.485	0.367	4.129	0.0016	3.67	0.830	6.000
20	(GACA)4	50	12	9	0.75	0.498	0.374	4.774	0.0018	2.52	0.720	7.355
21	(GA)8- RC	51	12	11	0.92	0.497	0.373	5.935	0.0015	3.77	0.710	8.710
22	(GACA)5	55	10	10	1.00	0.496	0.373	5.419	0.0016	3.73	0.707	5.742

T_a = Annealing temperature, NB: Number of total band, NPB: Number of polymorphic band, PP%: Percentage of polymorphic band, H: Heterozygosity index, PIC: Polymorphic Information Content, E: Effective multiplex ratio, H.av: Arithmetic mean of Heterozygosity index, MI: Marker Index, D: Discriminating power, and R: Resolving power

Table 3 : genetic parameter of *Onobrychis* subgenera and sections								
subgenus	section	No. of species	Na	Ne	H	I	Number of polymer	PPB%
Onobrychis		21	1.84	1.54	0.31	0.47	189	84.75
	Onobrychis	17	1.8	1.51	0.3	0.44	178	79.82
	Lophobrychis	4	1.68	1.47	0.27	0.4	153	68.61
Sisyrosema		10	1.86	1.57	0.33	0.49	192	86.1
	Heliobrychis	2	1.34	1.24	0.14	0.21	77	34.53
	Hymenobrychis	8	1.81	1.5	0.3	0.44	182	81.61
Na: Observed number of alleles , Ne: Effective number of alleles ,H: gene diversity , I: Shannon's index , PPB%: The percentage of polymorphic loci								

Table 4 : The Analysis of Molecular Variance(AMOVA) among 31 *Onobrychis* species					
source of variation	Df	sum of squares	variance components	percentage of variation	p-value
among subgenus	1	416.42	3.87	7.08	<0.001***
among section within subgenus	2	355.79	13.39	24.51	0.0001***
among species within sections	59	2205.56	37.38	68.41	0.0001***
Total	62	2977.77	54.64		

Figure legends:

Fig 1. Association between *Onobrychis* species reveled by complete linkage model and Euclidean's genetic distance based on seed and fruit macro & micro-morphological traits (A) and ISSR marker (B).

Fig 2. PCA (correlation's measure) case scores using NCSS program. A: Seed and Fruit macro & micro-morphological traits, B: ISSR marker, and C: Joint data.

Fig 3. Structure harvester and Delta K value Bayesian model based assessment of population structure for 31 *Onobrychis* species based on ISSR markers. A: Bar Plot based on original population order, B: Bar plot based on estimated membership coefficients values, C: Bar Plot based on species ID (population) order, D: ΔK = mean ($|L"(K)|$)/sd(L(K)) and ΔK = 4 indicates the best K value, E: absolute value of the 2^{nd} order rate of change of the likelihood distribution (mean), F: rate of change of the likelihood distribution (mean), and G: mean of estimated Ln probability.

Fig 4. The Maximum Likelihood phylogenetic tree illustrating the genetic relationship among *Onobrychis* species based on 1000 bootstrap based on Maximum Likelihood model.

Figure 1: Association between *Onobrychis* species reveled by complete linkage model and Euclidean's genetic distance based on seed and fruit macro & micro-morphological traits (A) and ISSR marker (B).

Figure2: PCA (correlation's measure) case scores using NCSS program. A: Seed and Fruit macro & micro-morphological traits, B: ISSR marker, and C: Joint data.

Figure 3: Structure harvester and Delta K value Bayesian model based assessment of population structure for 31 *Onobrychis* species based on ISSR markers. A: Bar Plot based on original population order, B: Bar plot based on estimated membership coefficients values, C: Bar Plot based on species ID (population) order, D: ΔK = mean (|L"(K)|)/sd(L(K)) and ΔK = 4 indicates the best K value, E: absolute value of the 2nd order rate of change of the likelihood distribution (mean), F: rate of change of the likelihood distribution (mean), and G: mean of estimated Ln probability.

Figure 4: The Maximum Likelihood phylogenetic tree illustrating the genetic relationship among *Onobrychis* species based on 1000 bootstrap based on Maximum Likelihood model.

Figure S1: Fruit shape of different *Onobrychis* species. A: *O. transcaucasia*, B: *O. arenaria*, C: *O. iberica*, D: *O. cyri*-Grossh, E: *O. altissima*, F: *O. alba*, G: *O. inermis*, H: *O. petraea,* I: *O. oxyodonta*, J: *O. gracilis*, K: *O. persica*, L: *O. peduncularis* (Scale bar: 1 mm)

Figure S1: Continued. M: *O. hajastana*, N: *O. megataphrose*, O: *O. montana*, P: *O. viciifolia,* Q: *O. kemulariae*, R: *O. caput-galli*, S: *O. crista-galli*, T: *O. aequidentata*, U: *O. pulchella*, V: *O. melanotricha*, W: *O.argyrea*, and X: O. ptolemaica (Scale bar: 1 mm).

Figure S1: Continued. Y: *O. hypargyrea*, Z: *O. michauxii*, AB: *O. sintenisii*, AC: *O. chorassanica*, AD: *O. bobrovii-Grossh*, AE: *O. pallasii*, and AF: *O. radiate* (Scale bar: 1 mm).

Figu

re S2: Transverse sections of *Onobrychis* species seed coats: I) *O. caput-galli,* II) *O. altissima* and III) scanning electron microscope (SEM) cross section of *O. transcaucasia*, (C= Cuticle layer; M = Malpighian cells; O= Osteosclereids; P= Inner parenchyma tissue) (Scale bar = 40 µm for Nikon ECLIPSE E600 microscope and 100 µm for SEM).

Figure S3. Technical shape which discripe the seed surface pattern : A) Druse, B) Stellate, C) Undulate,

D) Flat, E) Reticulate, and F) Convolutus.

Figure S4: Scanning electron microscopy photographs of seed coat. A: *O. transcaucasia*, B: *O. arenaria*, C: *O. iberica*, D: *O. cyri*-Grossh, E: *O. altissima*, F: *O. alba*, G: *O. inermis*, and H: *O. petraea*.

Figure S4: Continued. I: *O. oxyodonta*, J: *O. gracilis*, K: *O. persica*, L: *O. peduncularis*, M: *O. hajastana*, N: *O. megataphrose*, O: *O. montana*, and P: *O. viciifolia.*

Figure S4: Continued. Q: *O. kemulariae*, R: *O. caput-galli*, S: *O. crista-galli*, T: *O. aequidentata*, U: *O. pulchella*, V: *O. melanotricha*, W: *O.argyrea*, and X: O. ptolemaica.

Figure S4: Continued. Y: *O. hypargyrea*, Z: *O. michauxii*, AB: *O. sintenisii*, AC: *O. chorassanica*, AD: *O. bobrovii-Grossh*, AE: *O. pallasii*, and AF: *O. radiate.*

B1   B2

C1   C2

Figure S5: Two-dimensional representation of genetic relationships based on principal components analysis among the 31 *Onobrychis* species. A: seed and fruit macro & micro-morphological traits, B: PCA based on ISSR data, and C: based on joint data.