Highly dense and scalable protein arrays for single-molecule studies

Tural Aksel, Hongji Qian, Pengyu Hao, Pierre F. Indermuhle, Christina Inman, Shubhodeep Paul, Kevin Chen, Ryan Seghers, Julia K. Robinson, Mercedes De Garate, Brittany Nortman, Jiongyi Tan, Steven Hendricks, Subra Sankar, Parag Mallick*

*To whom correspondence should be addressed
[1]Nautilus Biotechnology, 835 Industrial Rd, Suite 200, San Carlos, CA 94070, USA
Corresponding author at: 835 Industrial Rd, Suite 200, San Carlos, CA 94070, USA.
Email Address: paragm@nautilus.bio (Parag Mallick)

ABSTRACT

Single-molecule protein studies are challenging because both single-molecule sensitivity and a high dynamic range are needed to comprehensively analyze the complex mixture of proteins present in biologically relevant samples. One way to achieve both high sensitivity and dynamic range would be to create a protein array that arranges single molecules with uniform spacing to achieve billions of well-separated molecules on a patterned surface. So far, no approach has combined both a uniformly patterned surface and the possibility to deposit a single protein at each spot. In addition, current patterning approaches cannot be scaled to the size needed to interrogate billions of molecules. Here, we present a scalable method for fabricating dense protein arrays using brushy-DNA origami deposition onto patterned surfaces. Brushy-origami structures are made large enough to occupy the features on the array singly and we developed an efficient process to ensure single protein conjugation to the brushy-origami structures. We hypothesize that our technology will ultimately enable the development of highly scalable proteomics platforms that address the need for both single-molecule sensitivity and high dynamic range.

INTRODUCTION

Tools that can quantify proteome dynamics with single-molecule resolution[1-3] are critically needed to address a myriad of challenges in biomedicine including the measurement of low abundance proteins in therapeutic development[4] and biomarker discovery[5,6] as well as for single-cell proteomics studies[7,8]. Additionally, such tools could provide insights into the molecular heterogeneity of populations of proteoforms that would be masked by bulk measurements. Building tools to measure proteome dynamics with single-molecule resolution is challenging because both single-molecule sensitivity and a high dynamic range are needed to comprehensively analyze the complex mixture of proteins present in biologically relevant samples. Sensitivity arises by using single-molecule counting methodologies, while dynamic range relies on the measurement of a very large number of single molecules. Ideally, it would be possible to build a platform that could measure tens-of-billions of individual molecules. One possible approach to address this challenge is to create high-density, large-scale, single-molecule protein arrays that allow massively parallel detection/identification. To date, no such platform exists that can be readily scaled to allow the interrogation of billions of individual protein molecules.

The current gold standard for ultrasensitive protein detection is Simoa (Single Molecule Arrays), or digital ELISA, whereby single target molecules are captured on antibody-coated beads, enabling the detection of a few molecules in a sample[9,10]. However, these approaches rely on limiting dilution: digital measurements are possible due to a large excess number of beads over the number of target molecules in the sample, so that each bead has a small number of captured target molecules. The actual number of molecules per bead will follow a Poisson distribution and there is no way to know how many target proteins were actually captured per bead. Such techniques are therefore unsuitable for multiplexed quantification of several different proteins and are unable to achieve the dynamic range needs of complex protein samples.

There are several potential ways to create a patterned surface. Simoa techniques generally use microwells created using soft lithography techniques, whereby each well can hold only one bead[9]. Photolithography techniques can create a patterned surface with uniformly separated positively charged wells where each well is occupied by a large DNA scaffold[11]. Similarly, using e-beam lithography and nanoimprinting, flat DNA origami structures can be placed on a glass surface with a regular spacing[12]. However, these techniques are often challenging to execute robustly, and cost-effectively, at scale. Generally, the smaller the feature-size required, the more challenging or expensive the fabrication process. Consequently, to-date, no approach has combined both a high quality, low-cost, uniformly patterned surface and the possibility to scale to billions of molecules.

Since it is possible to conjugate a single protein at a specific location on a DNA origami nanoparticle structure[13], we hypothesized that DNA origami could be used as the scaffold to create single protein arrays on patterned surfaces. A defined and consistent number of fluorescent labels can be attached to origami particles, enabling their visualization and making it possible to determine if multiple particles are co-localized. Unfortunately, many origami designs are constrained in size due to limitations in the size of the commonly used scaffolds. While these limitations can be overcome using supramolecular assemblies, fabrication of such assemblies is challenging and error prone. Here we introduce an alternative approach to generate large tile-shaped origami particles, without requiring multiple units. When coupled to an efficient process that ensures single protein conjugation to the origami structures, this approach generates a self-assembling single-molecule protein array. We hypothesize that our technology will ultimately enable the development of highly scalable platforms that have both single-molecule sensitivity and address a wide dynamic range.

RESULTS

Overview of approach

We set out to design an approach whereby every protein in a biological sample is placed singly on a patterned surface (**Fig. 1**). Our approach uses large DNA origami-based particles that we refer to as 'structured nucleic acid particles' (SNAPs) and that have a single site for protein conjugation. In our overall workflow, we first extract the proteins from a biological sample using standard isolation and precipitation methods. Next, to conjugate the proteins to SNAPs, we functionalize their lysines via an N-hydroxysuccinimide-polyethylene glycol-methyltetrazine (NHS-PEG-MtZ) crosslinker. MtZ-modified proteins are mixed with SNAPs that carry single trans-cyclooctene (TCO) molecules. TCO and MtZ react to create a covalent bond between the SNAP and the protein[14]. Single-stranded DNA overhangs surrounding the SNAP are extended using terminal deoxynucleotidyl transferase (TdT), which increases the effective size of the particle{Yang, 2021 #7} (see Methods for details). We term the extended particles brushy-SNAPs. The negatively charged brushy-SNAPs are then deposited onto positively charged patterns on a glass/silicon surface.

SNAPs are conjugated to proteins with high efficiency

For the base component of brushy-SNAPs, we used a single-layer, square DNA origami tile design (**Fig. 2a, SFig. 1**) with an estimated diameter of 88 nm. The DNA origami structure was folded from a mix of ssDNA oligos and the m13mp18 ssDNA. All the oligos including the one with TCO were mixed in excess with the scaffold DNA. Folding was achieved by heating up the mix and slowly bringing the sample to room temperature. We verified that the TCO-modified oligo in the SNAP is functional and conjugates to MtZ-modified proteins with high efficiency and specificity. Analytical HPLC results of the SNAP mixed with excess MtZ-Cy5 showed that the fraction of SNAPs with the functional TCO group was above 95% (**SFig. 2, Table 1**). Similarly, analytical HPLC results of the SNAP, with and without the TCO group, mixed with MtZ-modified proteinA-647 demonstrated that the protein conjugation to the TCO group is specific and the conjugation efficiency is above 95% (**Fig. 2b, Table 1**). Agarose gel analysis showed that the extension of

brushes for the proteinA-647 conjugated SNAPs does not impact protein retention on the structure (**SFig. 3**).

Brushy SNAPs are large and monodispersed

To characterize the sizes of the base-tile and brushy-SNAPs we used both dynamic light scattering (DLS) and atomic force microscopy (AFM). DLS measurements showed that, in solution, the base SNAPs are monodispersed, with an average diameter of 73.6 nm. Brushy SNAPs were monodispersed with an average diameter of 215 nm (**Fig. 2c, Table 2**). We used AFM to obtain high-resolution, single-particle structural information on the integrity and size of the particles on a 2D surface (**Fig. 2d, top-left panel**). The measured SNAP dimensions in liquid matched the expected SNAP edge length (85-90 nm) and SNAP height (1.5-2 nm) (**SFig. 4**). We imaged brushy-SNAPs on mica in liquid (**Fig. 2d, right panels**) and as dry particles (**Fig. 2d, bottom-left panel**). In liquid, the seed tile structure was visible, but the brushes were not visible (**Fig. 2d, top-right panel**). In phase images, we observed a halo around the SNAPs where we expect to observe the brushes (**Fig. 2d, bottom-right panel**). When the brushy-SNAPs were dried on mica, we observed uniformly sized particles with a larger diameter of 579.8 nm (**Fig. 2d, bottom-left panel, SFig. 5**). Given the low density of the extended brushy arms, we did not expect to be able to clearly measure the sizes of the brushes via DLS. As such, the SNAP and the brushy-SNAP diameters measured by DLS are lower than the diameters measured from AFM images since SNAPs and brushy-SNAPs are not perfectly spherical in 3D. Overall, these analyses assured us that the particle structures were as expected, that brush extension did not negatively impact the SNAP fold, and that the brushy-SNAPs were substantially larger than the base-tile SNAPs, suggesting they were likely to exclude a larger area around themselves.

The brushy-SNAPs created a regularly spaced pattern on mica due to exclusion of particles from each other. To estimate the exclusion diameter for the brushy-SNAPs, we measured the nearest-neighbor distances between the particles. The nearest-neighbor distances yielded a tight distribution centered at 360 nm (**Fig. 2e**). To show that the observed regular pattern is unique to brushy-SNAPs, we calculated the nearest-neighbor distances for simulated SNAP depositions on

mica at the brushy-SNAP density for objects 90 nm wide. The nearest-neighbor distance distribution for simulated SNAP depositions was wider and peaked around 200 nm (**Fig. 2e**) suggesting that the observed 360 nm spacing of brushy-SNAPs on mica is a result of sub-Poissonian deposition[16]. These data suggest that the brushy-SNAPs can effectively exclude a much larger area than their central structure, and suggest that they may be suitable for single-SNAP deposition on chips with feature sizes of approximately 360 nm.

Patterned chip has high single-molecule occupancy with brushy-SNAPs

To create positively-charged landing pads for high-density single-molecule deposition of brushy-SNAPs, we patterned glass wafers using photolithography and deposited 3-aminopropyltrimethoxysilane (APTMS) on the surface. We expect the brushy structures to lay flat on the positively charged surface. As observed from AFM images, the pad diameter and the pitch of the APTMS patterned surface averaged 370 nm with a standard deviation of 21 nm and 1.599 μm with a standard deviation of 18 nm, respectively (**Fig. 3a-c, Table 3, SFig. 6**). The patterning is composed of square regions, 200 μm by 200 μm, each containing 15,129 landing pads. A region the size of a traditional 25 mm × 75 mm microscope slide contains 37,632 of these square regions, allowing deposition of nearly 600 million protein molecules. Incubation of the patterned surface with Cy5-labeled DNA oligo yielded patterned deposition of Cy5-oligo demonstrating that the patterning and APTMS coating of the pads were successful (**Fig. 3d**). These results suggest that it is possible, using straightforward photolithography and vapor deposition techniques, to generate large-scale patterned surfaces for SNAP deposition, with tight tolerances on landing pad size and pitch.

To assess single-molecule occupancy on the chips, we used two-color brushy SNAP mixing experiments (**Fig. 3e**). We produced Alexa488- and Alexa647-labeled brushy SNAPs separately, mixed them at equal ratio, and deposited them onto the chip at 100 pM. By counting the features that fluoresce by a single wavelength (indicating only one brushy-SNAP) and those that fluoresce by both wavelengths (indicating more than one brushy-SNAP), we can assess the degree of

exclusion. We note that this method is unable to capture multiple-localizations of same-colored SNAPs. We address this concern with a second method described below. Base-SNAP samples yielded no exclusion leading to 100% co-localization of 488 and 647 labeled species (**Fig. 3e left panel**). On the other hand, at 98% deposition, brushy-SNAPs led to less than 7% co-localization suggesting that brushy-SNAPs are large enough to occupy a pad singly, excluding the deposition of a second one (**Fig. 3e right panel**) and thus leading to large-scale super-Poisson loading of the patterned surface.

We additionally examined single-molecule occupancy using pad intensities. This approach, though less visual, can distinguish multiple SNAP landings regardless of their color. As base-SNAPs and brushy-SNAPs are decorated with 40 dye molecules, we expected the fluorescent intensities of multiple SNAPs to scale with the number of colocalized SNAPs. To identify single vs. multiple brushy-SNAPs on the pads, we performed deposition leading to low and high pad occupancy and compared the intensity histograms (**Fig. 3f, top two panels**). At brushy-SNAP concentrations above 100 pM, we observed a second intensity peak with a mean and standard deviation twice that of the first peak (**SFig. 7**). At concentrations above 100 pM for base-SNAPs, we observed a multi-modal wide distribution indicating that there could be multiple SNAPs per pad (**Fig. 3f, bottom panel**). To quantify colocalization from single color intensities, we fit a multimodal Gaussian to the intensity histogram and quantified the contribution of multi brushy-SNAP peaks to the intensity distribution (**SFig. 7**). Using single color intensities, we verified that co-localization rates for brushy-SNAPS, even at high deposition concentrations (above 100 pM), are not significantly impacted by increasing deposition concentrations (**Fig. 3g**). In these studies, the maximum average co-localization rate was approximately 2%, even at 400 pM deposition for brushy-SNAPs, but was above 90% at 100 pM deposition of base-SNAPs.

DISCUSSION

We have used a brushy-SNAP, fabricated using a DNA origami approach, alongside protein click-conjugation, photolithography and surface functionalization to generate a scalable method for

fabricating dense protein arrays with super-Poisson loading, while minimizing the number of loci in which multiple proteins are deposited. Additionally, it is possible to estimate which locations on the chip are likely to have multiple deposited SNAPs. Notably, the brushy-SNAPs, though not a supramolecular assembly, have diameters of greater than 200 nm, and can exclude adjacent deposition such that the average distance between adjacent particles is 360 nm. The features required to arrange such large particles are easily within reach of conventional lithographic approaches, reducing the complexity and cost of chip fabrication. Our approach is not reliant upon limiting dilution to minimize multiple occupancy, which was less than 2% even at >90% occupancy.

The high occupancy and dense packing will ultimately enable efficient measurement by either optical or non-optical methods. We showed chips with a pitch of 1.6 μM giving approximately 600 million protein molecules per microscope-slide sized chip. Fabricating chips in a hexagonal grid with a pitch of 900 nm would be straightforward and would allow for 3x as many landing pads, thereby generating surfaces capable of holding billions of molecules within the space of a conventional microscope slide. Using a few such chips in concert, deposition of 10^{10} protein molecules could readily be achieved. Simulations suggest measurement of 10^{10} protein molecules would enable assays to reach dynamic ranges greater than 11 orders of magnitude, far exceeding the dynamic range of existing immuno-assays while also creating the possibility of single-molecule sensitivity.

Potential limitations of the approach will be addressed in future work. For instance, our method does not allow orientational control of the brushy-SNAPs on the APTMS functionalized pads via electrostatic interactions and this may kinetically impact interactions between molecules in solution and the immobilized target protein. To achieve orientational control, one side of the particles could be functionalized with a chemistry that specifically interacts or couples with the surface chemistry on the pads. In addition, the accessibility of the target protein could be further enhanced by flexible and long linkers or rigid DNA origami extensions keeping the target away from the brushy-SNAP surface. Another potential limitation is the lack of tight control on the size

distribution of the brushy-SNAPs. The size of the brushy-SNAPs is controlled by the dTTP monomer concentration added in the terminal transferase extension reaction. Although our current process yields relatively tight distributions based on the DLS measurements and gel electrophoresis, we could use density-based centrifugation methods to fractionate the sample further to achieve a tighter distribution for a lower brushy-SNAP co-localization rate.

Ultimately, we envisage protein arrays such as these having a wide variety of uses, including becoming a key component in proteomics platforms. By immobilizing samples on the array, it may be possible to probe iteratively with a series of affinity reagents. Additionally, such arrays would enable a direct link between the identification and quantification of immobilized proteins. Further potential applications include the screening of protein-protein interactions or protein-ligand interactions, such as for compound screening. Kinetics assays could also be performed either through single-molecule imaging of interactions over time, or using non-imaging methods such as surface plasmon resonance or impedance/conductance. Alternately, by immobilizing affinity reagents, either patterned, or non-patterned, on the array, it may be possible to develop ELISA-like assays with sensitivity far exceeding existing digital ELISA approaches. We note, however, that immobilizing affinity reagents, rather than sample proteins, will mean the method is not a true counter of single molecules present in the sample, as detections will ultimately be governed by the binding on-rate of immobilized affinity reagents. In summary, we expect our technology to revolutionize single-molecule protein studies by providing highly scalable arrays for a wide range of applications.

MATERIALS and METHODS

Structured nucleic acid particle (SNAP) production

The SNAP was designed and modified using CaDNAno and folded as described previously {Tikhomirov, 2017 #27}. Briefly, 10 nM m13mp18 ssDNA scaffold (Bayou Biolabs, Metairie, LA) was folded with 75 nM staples (IDT, Coralville, IA). Reaction volume was 50 µl per tube. The SNAP folding buffer was 5 mM Tris-Cl, 5 mM NaCl, 1 mM EDTA, 12.5 mM $MgCl_2$ at pH 8.0. SNAP folding was performed on Biorad Tetrad by temperature ramp starting from 90°C and ending at 20°C, decreasing one °C every minute. The folded SNAP was concentrated to nearly 100 nM using Amicon Ultra (Millipore Sigma, Burlington, MA) spin filter with 100 kDa cutoff. The SNAP was purified from excess staples using Agilent Bio SEC-5, 5 µm, 2000Å, 4.6 x 300 mm (P/N: 5190-2543) size exclusion column on an Agilent 1200 HPLC system equipped with a Diode Array Detector and Fraction Collector. Method for SNAP purification consisted of isocratic flow at 0.3 mL/min of 5 mM Tris-Cl, 200 mM NaCl, 1 mM EDTA, 12.5 mM $MgCl_2$ (1X O.B.). Monomeric SNAP peak fractions were combined to prepare SNAPs decorated with Alexa488 or Alexa647 dye-labeled oligos (IDT Coralville, IA) and TCO- (in-house) or biotin-modified oligo (IDT Coralville, IA). Dye-labeled and TCO- or biotin-modified oligos were annealed to SNAP handles in 2X excess with respect to SNAP handle concentration. Dye oligos were hybridized to 40 complementary handles on the SNAP surface. Biotin- or TCO-modified oligo was annealed to a single complementary handle on the SNAP surface. Annealing was performed with a temperature ramp starting from 40°C and ending at 20°C, decreasing one °C every minute on Biorad Tetrad. SNAPs decorated with dye-oligos and TCO- or biotin-modified oligos were purified from excess oligos using Agilent Bio SEC-5 column on an Agilent 1200 HPLC system.

Brushy-SNAP production

For 10 nM brushy-SNAP-3000 production, 10 nM SNAP sample was mixed with 3.6 U terminal transferase enzyme (NEB, Ipswich, MA) and 1.32 mM dTTP (NEB, Ipswich, MA) in 40 µl reaction volume. The dTTP amount required for 3000 bases extension of the 7T-3' handles surrounding the SNAP edges was calculated via 3000 x 44 x [SNAP]. The reaction was kept at 37°C for 16 h to reach completion. The excess enzyme was removed, and the buffer was exchanged to 1X O.B. using Amicon Ultra (Millipore Sigma, Burlington, MA) spin filters with 100 kDa cutoff.

SNAP and brushy-SNAP-3000 characterization

SNAP and brushy-SNAP samples were run on 0.5% Agarose Gel made using Seakem LE Agarose (Lonza, Basel, Switzerland) in 0.5X TBE supplemented with 11 mM $MgCl_2$. For visualization of the samples, SybrSafe was added to the gel from a 10,000X stock (Thermofisher, Waltham, MA). The gel was run at 70V for 3 h in an ice-water bath. The gel was visualized using Biorad Chemidoc gel imager (Hercules, CA).

Dynamic light scattering measurements of the SNAP and brushy-SNAP samples were performed on Malvern Zetasizer Nano. A 15 µl sample at around 10-20 nM concentration was added to quartz cuvette for measurements. Data was acquired with automatic measurement duration settings at room temperature.

AFM images were collected on Park systems NX-20 (Santa Clara, CA) using USC-F0.3-k0.3 (0.3 N/m) for liquid and NSC14/Cr-Au (5 N/m) for dry imaging. The data were acquired in manual mode at 1.89 Hz for high-resolution, and at 0.35 Hz for low-resolution imaging. The raw AFM images were processed using Gywddion (http://gywddion.net).

Chip fabrication

200 mm photoresist patterned wafers were sourced from CEA-Leti (Grenoble, France). The glass (Borofloat 33, Schott) wafers were modified with an HMDS adhesion layer, followed by a

lift off Layer (LOL), bottom anti reflecting coating (BARC), and photoresist stack. The photoresist was patterned in a square packed array of 320 nm round openings to reveal the silicon dioxide surface with a pitch of 1.625 microns. The patterned wafers were then modified with a water-alcohol-based plasma followed by 3-aminopropyltrimethoxysilane (APTMS) in an RPX-540 CVD system (Integrated Surface Technologies, Menlo Park, CA). A silicon monitor with 100 nm thermal oxide was placed in the chamber alongside the wafers as a proxy to measure the resulting APTMS film thickness and contact angle. Typical film thickness was about 4 Å to 5 Å and the contact angle (after an initial water rinse) ranged from 35 to 50 degrees. Following the CVD process, an additional protective layer of resist (SPR 220, Kayaku, Westborough, MA) was applied before dicing into 1-inch square die. The photoresist stack was then removed from the surface of each die by sonicating in a N-Methyl-2-pyrrolidone (NMP) bath at 50°C. The dice were then rinsed with isopropyl alcohol followed by DI water and dried under a stream of nitrogen. The contact angle was measured before and after the photoresist removal and each die was inspected under a 100X objective microscope to confirm resist removal. Representative dice were sent for AFM characterization to confirm feature size and surface roughness (Evans Analytical Group, Sunnyvale, CA). The dice were then assembled into a flow cell using a PSA gasket and a polyethylene glycol coated (Integrated Surface Technologies, Menlo Park, CA) glass slide with drilled fluidic ports (Schott).

Flow cell assembly

The flow cell consisted of a 25 mm x 75 mm glass backer with a glass-based nanoarray chip joined by a film of pressure-sensitive adhesive (PSA). The backer had drilled holes to form the inlet and the outlet ports for the flow cell lanes. The PSA was cut in a design to define the lanes of the flow cells. The PSA layer also acted as the spacer to form the height of the channels. The flow cell assembly consisted of two main steps. First, the sub-assembly was done where the PSA was attached to the underside of the glass backer, and second, the subassembly was

attached to the array chip. The PSA was placed onto the assembly fixture adhesive side facing up, using the alignment pins to register. The glass backer was then placed down on top of the PSA and pressed to complete the sub-assembly process. The assembly fixture helps align the glass backer holes to the PSA cut lanes. The final step in the flow cell assembly process was to attach the chip to the backer. The liner of the PSA was removed to expose the adhesive surface. The nanoarray chip was pressed against the backer to complete the flow cell assembly.

TCO oligo production

5' amine modified DNA oligo with C12 linker was purchased from IDT (Coralville, IA). Amino modified oligo was resuspended in 20% DMSO/ 0.8x PBS pH 8.0 to 2 mM. 50 molar equivalents of diisopropylethylamine (DiPEA) and 30 molar equivalents of TCO-PEG4-NHS ester (CAS 1621096-79-4) were added to the resuspended oligo in neat DMSO. Additional 20% DMSO / 0.8x PBS was added to fully solubilize all components. Final reaction composition was 1 mM amino-oligo, 30 mM TCO-PEG4-NHS ester, 50 mM DiPEA, in 59% DMSO, 0.4x PBS. Reaction was shaken at 25°C for approximately 1 h in the dark before the reaction was checked for completion (complete conversion of the amino oligo) by analytical RP-HPLC. The TCO-modified DNA oligo was purified on an Agilent 1260 HPLC system equipped with a binary pump, PDA and Fraction Collector. Purification was performed using a Phenomenex Jupiter 5 µm C5 300Å RP column (250 x 10 mm) (P/N: 00G-4052-N0). HPLC Mobile Phase A = 400 mM hexafluoroisopropanol, 16.3 mM triethylamine in water; HPLC Mobile Phase B = 100% methanol. Column temperature = 30°C. Fractions were collected in 16 s intervals from 20 to 30 min. HPLC method was as follows:

Time (min)	A%	B%	Flow (mL/min)
0	90	10	3.5
5	90	10	3.5

10	62	38	3.5
15	62	38	3.5
25	52	48	3.5
30	52	48	3.5
35	40	60	3.5
37	40	60	3.5
40	90	10	3.5
60	90	10	3.5

Following HPLC purifications, fractions were checked for purity using analytical RP-HPLC coupled to a Q-TOF MS. Pure fractions (>90% *trans* isomer product by A260 chromatogram peak area) were pooled and evaporated to dryness using a Labconco Centrivap.

Modification of ubiquitin and proteinA-647

Recombinant ubiquitin with N-terminal Histag (BPS Biosciences, P/N: 79293) and Alexa647-proteinA conjugate (Thermofisher, Waltham, MA) were modified with methyltetrazine-PEG4-NHS ester (mTz) using the following conditions. Ubiquitin was buffer-exchanged to 1x PBS pH 7.4 using BioRad Econo-Pac 10DG Desalting Columns (Biorad, Hercules, CA); 10 molar equivalents of mTz were added (from a stock 10 mM solution in dimethylformamide (DMF)), along with additional 1x PBS pH 7.4, to the protein such that the final protein concentration was 40 µM in a final solvent composition of 4% DMF in PBS pH 7.4. Reaction was incubated for 3 h at 25°C in the dark. Modification reaction was quenched by adding a sufficient volume of stock 800 mM L-Arginine in 1x PBS (not pH adjusted after arginine addition) such that the final concentration of arginine in the reaction mixture was 50 mM. Quench was performed for 2 h at 25°C, in the dark. Alexa647 protein A conjugate was modified and quenched with mTz in a

similar fashion but with a final protein concentration (in the modification) of 80 µM and 30 molar equivalents of the mTz-PEG4-NHS ester, 8% DMF in 0.92x PBS pH 8.0 final solvent composition. Quenched modification reactions were purified using the manufacturer's procedure for desalting on BioRad Econo-Pac 10DG Desalting Columns (Biorad, Hercules, CA) to remove excess mTz modifier. 10DG columns were equilibrated and eluted with 1x PBS, 200 mM L-Arginine pH 8.7. Purified proteins were concentrated to >15 µM using Sartorius Vivaspin 3k (ubiquitin) or 10k (proteinA) 3 mL centrifugal spin filters with hydrosart membranes. The concentrated proteins were rinsed in the spin filters with an additional 2 mL of 1x PBS, 200 mM L-Arginine pH 8.7. Modified proteins were stored at 4°C or directly taken to the conjugation step.

Conjugation of MtZ-modified proteins to SNAPs

SNAPs with a single TCO (*trans*-cyclooctene) functional group, at a minimum of 100 nM, were incubated with approximately 3 molar equivalents of mTz-modified proteins in 1x O.B. at 25°C overnight (approximately 16 h) on a tube-shaker protected from light with foil. Protein-conjugated SNAP sample was purified from excess unconjugated protein via HPLC-SEC (Size Exclusion Chromatography) using an Agilent 1200 HPLC system equipped with DAD and fraction collector. The SEC column used was an Agilent Bio SEC-5, 5 µm, 2000Å, 4.6 x 300 mm (P/N: 5190-2543). Method for purification consisted of isocratic flow at 0.3 mL/min of 1x HPLC O.B. (200 mM NaCl, 11 mM $MgCl_2$, 5 mM Tris-HCl, 1 mM EDTA) for 1.5 column volumes. Fractions corresponding to the SNAP peak were pooled and concentrated using Amicon Ultra 100k MWCO Regenerated Cellulose Spin Filters. Purified, concentrated sample was stored in DNA Low-Bind tubes at 4°C.

Quantification of TCO oligo and MtZ reagent conjugation efficiency

Multiple tools were used to quantify the conjugation efficiency of MtZ reagents to SNAPs with a single TCO group.

For fluorescent MtZ reagents, HPLC-SEC chromatography was used to derive the relative absorbances of SNAP at 260 nm and the conjugated fluorescent molecule's absorbance at its max absorbance wavelength. The 260 nm SNAP extinction coefficient used to quantify the relative molar ratio of SNAP to the conjugated particle was 116,898,548/ (cm*M). For Mtz-Cy5 conjugation, the extinction coefficient at 652 nm was 250,000/ (cm*M). For Mtz-ProteinA-647 conjugation, the extinction coefficient at 652 nm was 956,000/ (cm*M). The conjugation efficiency was derived using Equation 1.

$$\text{Conjugation efficiency} = \frac{(\text{SNAP Abs. at 652 nm}) \times (\text{SNAP Ex. Coeff. At 260 nm})}{(\text{SNAP Abs. at 260 nm}) \times (\text{FL mol Ex. Coeff. At 652 nm})} \qquad \textbf{Eq. 1}$$

For non-fluorescent protein targets, the conjugation efficiency was assessed by the difference in MtZ-Cy5 conjugation efficiency of the starting SNAP with TCO group and SNAP coupled to MtZ-modified protein. In addition, a protein conjugated SNAP sample was run on denaturing SDS-page gel to observe the protein conjugated oligo band.

SNAP and brushy-SNAP fluorescent imaging

In each lane of a dry patterned chip, SNAPs and brushy-SNAPs with or without the conjugated protein were deposited at 50-400 pM in running buffer (10 mM HEPES, 120 mM NaCl, 10 mM MgCl$_2$, and 5 mM KCl, pH 7.4, 0.1% Tween-20 (Sigma P/N 44112) and 0.001% Lipidure (NOF P/N: CM5206)) for 15 min at room temperature. Excess nanoparticles were washed with 1mL of running buffer. In lanes where dark SNAPs or brushy-SNAPs were deposited, 1 μM of dye-labeled oligos (18.1 in sequence, Alexa Fluor 488 or Alexa Fluor 647) in hybridization buffer (running buffer + 100mg/mL dextran sulfate) were incubated for 2 min before rinsing out by 1 mL of running buffer. Images were taken on a home-built fluorescence microscope. The microscope objective was Olympus UCPLFLN 20x, 0.7 NA, with a correction collar for thick cover glass. The camera was Photometrics Prime BSI (2048 x 2048 pixels, 6.5 mm pixel size,

sCMOS). Effective pixel size in the focal plane of the microscope objective was 0.325 mm. The excitation light source was a Necsel NovaLum 3-laser module (488 nm, 525 nm, 638 nm). The three lasers were independently controlled. The fluorescence filter set was Chroma 89402 (DAPI/FITC/TRITC/Cy5).* Laser light was focused to a 2048-pixel x 200-pixel (666 mm x 65 mm) rectangular area that was galvo-scanned (galvo scanner from Thorlabs GVS201) across the camera's field of view synchronously with the camera's rolling shutter. A B&K Precision 4052 waveform generator generated the sawtooth wave that controlled the galvo. The camera's exposure time was set so that 200 rows of the sCMOS sensor detected light simultaneously. These 200 rows were in effect a confocal slit that moves across the 2048 rows of the sCMOS sensor at a rate of 1 row per 11.28 ms when the camera is operated at its maximum line rate. The line rate was reduced to 1 row per 112.8 ms when increased sensitivity was required. A total of 30 images in a 2-by-15 layout was acquired for each lane of the flowcell.

Image processing

The input images were corrected for spherical aberration using the patterned surface array dimensions and the pad pitch as ground truth. Patterned array and single pad boundaries were determined using the subarray fluorescent patterns excited by the 488 laser. The mapping between 488 and 635 laser acquired images was constructed using a SNAP decorated with 44 Alexa488 and 20 Alexa647 modified oligos. The occupancy of the pads was determined based on the difference between the pad intensities and the background intensity. Briefly, the whole image was normalized against the background, where the background intensity estimate of the images was calculated by tiling the image (64x64 pixel) and computing the 5th percentile for each tile. The computed background intensity was fitted with paraboloid. The image was then normalized against the fitted background. The primary feature score was computed for each candidate object by convolving the pads with a 5x5 gaussian kernel on the normalized image.

Occupancy for each pad was classified using a one-feature clustering method to determine threshold the occupancy status of the candidate pad.

Quantification of brushy-SNAP co-localization from fluorescent microscopy images

Two types of experiments were used to estimate brushy-SNAP co-localization on the pads. For the first experiment, Alexa488 and Alexa647 labelled brushy-SNAPs were mixed at equimolar ratio and deposited on the chips. The co-localization rate was estimated by the ratio of pads occupied by Alexa488 and Alexa647 brushy-SNAPs and all occupied pads. For the co-localization analysis, six microscopy images taken at the center of the flow cell away from fluidic inlet and outlet were used. For the SNAP co-localization analysis, we used the double color mixing data, because the SNAP single color deposition intensity distribution is wide multi-modal, not allowing easy separation of single and multi-tile peaks. For the second experiment, Alexa488 labelled brushy-SNAPs were deposited onto the surface and the co-localization rate was estimated from the fluorescence intensity histogram analysis. In this analysis, the bimodal gaussian distribution was fitted to the intensity distribution where the mean intensity and the standard deviation of the second peak were constrained to twice the mean intensity and the standard deviation of the first peak respectively. In most of the intensity distributions with Brushy-SNAPs, the second peak in the distribution had low amplitudes; therefore, constraining the mean intensity and the standard deviation of the second peak provided robust results that are comparable among images from the same experimental condition and images across different experiment conditions. The co-localization rate was quantified by Equation 2.

$$\text{Co-localization rate} = \frac{2 \times (\text{Amplitude of Peak2})}{(\text{Amplitude of Peak1}) + 2 \times (\text{Amplitude of Peak2})} \qquad \textbf{Eq. 2}$$

Table 1. TCO SNAP and MtZ reagent conjugation efficiency determined from the HPLC-SEC chromatogram absorbances.

Conjugated molecule	Conjugation efficiency (%)
MtZ-Cy5	*95±10
ProteinA-647	*96±1
Ubiquitin	**90±5

* Conjugation efficiencies for the fluorescent molecules to SNAP are measured as described in the methods.

** Ubiquitin conjugation efficiency is estimated from the difference between the MtZ-Cy5 conjugation efficiency of the starting TCO SNAP and the ubiquitin-conjugated SNAP.

Table 2. SNAP and brushy-SNAP diameters measured from AFM and DLS data.

Particle type	Measurement type	Average particle diameter (nm)
Brushy-SNAP	*AFM (aqueous)	361.0 ± 35.8
Brushy-SNAP	AFM (dry)	579.8 ± 38.1
SNAP	AFM (aqueous)	85.3 ± 1.8
SNAP	DLS	73.6 ± 0.6
Brushy-SNAP	DLS	214.5 ± 6.3

*Average AFM brushy-SNAP diameter in aqueous imaging is calculated from the nearest neighbor distances between the brushy particles deposited at maximum density.

Average AFM particle diameters are calculated from at least 20 particles. DLS diameters are averaged over 3 to 6 measurements.

Table 3. Chip pad diameter and the pattern pitch data from AFM images

Feature	Length (nm)
Pad diameter	370 ± 21
Pitch	1599 ± 18

Average pad diameter and pitch lengths are calculated from at least 10 measurements from AFM chip images.

Pitch is the pad-center to pad-center distance along the horizontal and vertical axis along the square grid.

Figure 1. Overview of the single protein array sample prep workflow. The method presented here would be used to create single protein arrays from biological samples for single molecule studies. The workflow would consist of 1) MtZ modification of proteins separated from biological samples, 2) Production of origami tiles (SNAPs) with single TCO for click chemistry, 3) Conjugation of MtZ protein mixture to SNAPs bearing single TCO group, 4) Extension of ssDNA overhangs on the SNAP edges by terminal transferase. 5) Deposition of protein-bearing brushy-SNAPs onto APTMS functionalized patterned surface.

Figure 2. Structural and functional characterization of the SNAPs and brushy-SNAPs. A) Protein-conjugated brushy-SNAP production workflow. The oligo that bears the TCO moiety and the remaining staples are designed using CaDNAno (Douglas et al. 2009). The staples including the TCO-modified DNA oligo are combined with m13mp18 ssDNA scaffold in excess and the SNAP is folded over a temperature ramp on a thermocycler. The TCO-modified SNAP is coupled to MtZ protein, then the ssDNA overhangs flanking the SNAP edges are extended by terminal transferase to increase the particle size. Brushy-SNAPs are then deposited onto a patterned surface to occupy each pad with a single brushy particle. **B)** HPLC-SEC chromatograms of TCO-bearing SNAP conjugated to proteinA-647 (top) and the SNAP with no TCO group incubated with proteinA-647 (bottom). Blue curve is for absorbance at 260 nm and the red curve is for absorbance at 652 nm. The dashed lines mark the tile peak boundaries. **C)** DLS diameter distribution for SNAP (black dashed line) and brushy-SNAP (black solid line). **D)** AFM images of SNAP and brushy-SNAP in aqueous and dry conditions. The SNAPs adopt the fold aimed at by the design (top left, scale bar is 100 nm). At saturating concentrations (3-5 nM) the brushy-SNAPs create a pattern excluding each other on the mica surface (top right, scale bar is 1 µm). The dashed lines are for the nearest-neighbor distances. The brushes are not clearly visible in aqueous imaging conditions, but there is a halo observed around the brushy particles in the phase images (bottom right, scale bar is 400 nm). When brushy-SNAPs are rinsed and dried on the mica, the full size of brushy particles becomes visible (bottom left, scale bar is 400 nm). **E)** Brushy-SNAP to brushy-SNAP nearest neighbor (NN) distance distribution (blue) on mica (panel B, top right) vs. nearest neighbor distance distribution derived from simulated SNAP depositions at the same density as the brushy-SNAP density (black).

Figure 3. Single-molecule deposition studies with brushy-SNAPs. A) 2D cartoon representation for the APTMS patterned chip surface. The expected pad diameter is 370 nm and the square grid pitch on the horizontal and vertical axis is 1.6 µm. **B)** Landing pad diameter (average and standard deviation shown). **C)** Landing pad spacing (average and standard deviation shown). **D)** Alexa647 labeled DNA oligo deposition fluorescence microscopy image reveals the gridded pattern. Zoomed view of the area enclosed by yellow square is on the right. **E)** Two-color SNAP (left) and brushy-SNAP (right) co-localization images. Green is for the Alexa488 channel (top left), and purple is for the Alexa 647 channel (top right). The white pad counts on the composite image show the degree of brushy-SNAP co-localization (bottom left, bottom right is for the zoomed view). **F)** Brushy-SNAP co-localization is less sensitive to increase in deposition concentration than the base-SNAP co-localization. An increase in brushy-SNAP (top: 5 pM, middle: 200 pM) deposition concentration leads to higher occupancy (left) while not causing a major impact on the intensity distributions (top right and middle). 50 pM concentration deposition of the SNAP (bottom left) leads to a multimodal pad intensity distribution correlated with high co-localization observed with two-color brushy-SNAP mixing experiments (Panel C, left). The red circled pads on the microscopy images (left) are for the pads that have a higher intensity than the red line on the intensity histograms (right). **G)** Pad co-localization and occupancy rates. Blue is for 50 pM, green is for 100 pM, red is for 200 pM and purple is for 400 pM brushy-SNAP deposition concentration. Base-SNAP (black) deposition concentration is 100 pM. The shaded areas are a visual aid to show the data points belonging to different deposition concentrations.

REFERENCES

Boskovic, F. & Keyser, U. F. Toward single-molecule proteomics. *Science* **374**, 1443-1444, doi:10.1126/science.abn0001 (2021).

Alfaro, J. A. *et al.* The emerging landscape of single-molecule protein sequencing technologies. *Nat Methods* **18**, 604-617, doi:10.1038/s41592-021-01143-1 (2021).

Brady, M. M. & Meyer, A. S. Cataloguing the proteome: Current developments in single-molecule protein sequencing. *Biophysics Reviews* **3**, 011304, doi:10.1063/5.0065509 (2022).

Lill, J. R., Mathews, W. R., Rose, C. M. & Schirle, M. Proteomics in the pharmaceutical and biotechnology industry: a look to the next decade. *Expert Rev Proteomics* **18**, 503-526, doi:10.1080/14789450.2021.1962300 (2021).

Landegren, U. *et al.* Opportunities for sensitive plasma proteome analysis. *Anal Chem* **84**, 1824-1830, doi:10.1021/ac2032222 (2012).

Hoheisel, J. D., Alhamdani, M. S. & Schroder, C. Affinity-based microarrays for proteomic analysis of cancer tissues. *Proteomics Clin Appl* **7**, 8-15, doi:10.1002/prca.201200114 (2013).

Davis-Marcisak, E. F. *et al.* From bench to bedside: Single-cell analysis for cancer immunotherapy. *Cancer Cell* **39**, 1062-1080, doi:10.1016/j.ccell.2021.07.004 (2021).

Slavov, N. Counting protein molecules for single-cell proteomics. *Cell* **185**, 232-234, doi:10.1016/j.cell.2021.12.013 (2022).

Rissin, D. M. *et al.* Single-molecule enzyme-linked immunosorbent assay detects serum proteins at subfemtomolar concentrations. *Nat. Biotechnol.* **28**, 595-599, doi:10.1038/nbt.1641 (2010).

Cutler, J. I., Auyeung, E. & Mirkin, C. A. Spherical nucleic acids. *J. Am. Chem. Soc.* **134**, 1376-1391, doi:10.1021/ja209351u (2012).

Drmanac, R. *et al.* Human genome sequencing using unchained base reads on self-assembling DNA nanoarrays. *Science* **327**, 78-81, doi:10.1126/science.1181498 (2010).

Gopinath, A. & Rothemund, P. W. K. Optimized assembly and covalent coupling of single-molecule DNA origami nanoarrays. *ACS Nano* **8**, 12030-12040, doi:10.1021/nn506014s (2014).

Derr, N. D. *et al.* Tug-of-war in motor protein ensembles revealed with a programmable DNA origami scaffold. *Science* **338**, 662-665, doi:10.1126/science.1226734 (2012).

Devaraj, N. K., Weissleder, R. & Hilderbrand, S. A. Tetrazine-based cycloadditions: application to pretargeted live cell imaging. *Bioconjug. Chem.* **19**, 2297-2299, doi:10.1021/bc8004446 (2008).

Yang, Y. *et al.* Programmable Site-Specific Functionalization of DNA Origami with Polynucleotide Brushes. *Angew. Chem. Int. Ed Engl.* **60**, 23241-23247, doi:10.1002/anie.202107829 (2021).

Berkson, J. Tests of Significance Considered as Evidence. *J. Am. Stat. Assoc.* **37**, 325-335, doi:10.1080/01621459.1942.10501760 (1942).

1. Supplementary Figures

SFigure 1. CaDNAno design diagram for DNA origami tile.

SFigure 2. HPLC-SEC chromatograms of the (A) TCO-modified SNAP and (B) ubiquitin-conjugated SNAP incubated with Mtz-Cy5. A) TCO-modified SNAP incubated with excess MtZ-Cy5. The peak between the dashed line is for the SNAP. Blue is for absorbance at 260 nm and red is for absorbance at 652 nm. The right peak is for the free MtZ-Cy5. The conjugation efficiency of the TCO with MtZ-Cy5 is quantified from the ratios of the 260 nm and 652 nm absorbance peak areas. **B)** Ubiquitin-conjugated SNAP incubated with excess MtZ-Cy5. Ubiquitin-conjugated SNAP does not absorb at 652 nm suggesting that there is no SNAP with free (unconjugated) TCO, therefore all the SNAPs are conjugated to ubiquitin.

SFigure 3. 0.5% Agarose gel image for proteinA-647-conjugated SNAP and brushy-SNAP. Samples from left to right are 1) 1 kb NEB DNA Ladder, 2) m13mp18 scaffold, 3) TCO-SNAP, 4) TCO-SNAP conjugated with MtZ-proteinA-647, 5) Sample 4 extended with 1000 fold excess dTTP per initiation site, 6) Sample 4 extended with 2000 fold excess dTTP per initiation site, 7) Sample 4 extended with 3000 fold excess dTTP per initiation site, 8) Control SNAP incubated with MtZ-proteinA-647 and purified by HPLC-SEC, 9) Sample 8 extended with 1000 fold excess dTTP per initiation site, 10) Sample 8 extended with 2000 fold excess dTTP per initiation site, 11) Sample 8 extended with 3000 fold excess dTTP per initiation site. Top image is acquired with SybrSafe laser and filter settings. Middle image is acquired with Alexa647 laser and filter settings. Bottom image is the composite of the top (green) and middle (red) images.

SFigure 4. SNAP dimensions from AFM in aqueous conditions. A) High-resolution zoomed-in view of a SNAP by AFM. The scale bar is 50 nm. The z-height profile of the SNAP along the dashed line is in C. **B)** 3D view of the SNAP in A. **C)** The Z-height profile of the SNAP in A. The length of the SNAP along the dashed line is 90 nm. The maximum height is 1.5 nm.

SFigure 5. Brushy-SNAP dimensions from AFM in dry condition. A) AFM image of brushy-SNAPs in dry condition. **B)** Zoomed-in view of a brushy tile. The radially averaged profile of the brushy tile along the green line is in C. **C)** Radially averaged z-height profile of the brushy tile. The radius at which the z-height flatness is marked by a dashed line. The effective diameter of the brushy structure is estimated by multiplying the radius (244 nm) by two.

SFigure 6. Patterned chip pad size and array pitch characterization. A) AFM image of the patterned chip surface. APTMS functionalized pads are enclosed by circular rims. **B)** Height profile along the green dashed line in A. The numbers show the width of the pads and the inter-pad distance in nm along the line.

SFigure 7. Quantification of brushy-SNAP co-localization per pad on the patterned chip from fluorescence intensities. At concentrations higher than 100 pM, the intensity distribution becomes multimodal with two distinct peaks. The brushy-SNAP titration results suggest that the second minor peak with higher mean intensity corresponds to pads occupied with two brushy-SNAPs. The co-localization rate is quantified from two modal gaussian fit to the intensity distribution where the second gaussian is constrained to have twice the mean intensity and the standard deviation of the major peak.