Inner hair cell synapse density influences auditory processing

Lingchao Ji[1,&]; Beatriz C. Borges[1]; David T. Martel[1]; Calvin Wu[1]; M. Charles Liberman[2]; Susan E. Shore[1,3, 4,*], Gabriel Corfas[1,*]

[1]Kresge Hearing Research Institute and Dept. of Otolaryngology, Head and Neck Surgery, University of Michigan

[2]Mass Eye and Ear Infirmary and Harvard Medical School

[3]Biomedical Engineering, University of Michigan

[4]Molecular and Integrative Physiology, University of Michigan

[*] co-senior authors

[&] Current affiliation: Dept. of Otolaryngology, Peking University Shenzhen Hospital

Corresponding author:

Gabriel Corfas, Ph.D.
Kresge Hearing Research Institute
The University of Michigan
Medical Sciences I Building, Rm. 5428
1150 West Medical Center Drive
Ann Arbor, MI 48109-5616
Tel: 734-615-2222
Email: corfas@med.umich.edu

Acknowledgements: This work was supported in part by NIH/NIDCD R01 DC018500 (GC), NIH/NIDCD R01 DC000188 (MCL), and a grant from the American Tinnitus Association (GC).

Conflict of interest: GC and MCL are scientific founders of Decibel Therapeutics, have equity interest in the company and have received compensation for consulting. SES and DM are scientific founders of Auricle, Inc and have equity interest in the company. Neither company was involved in this study.

Abstract

It has been proposed that inner hair cell (IHC) synaptopathy, i.e., loss of synapses between spiral ganglion neurons and IHCs, leads to hidden hearing loss (HHL), i.e, poor performance on challenging hearing tasks despite a normal audiogram. However, this has only been tested in animals after exposure to noise or ototoxic drugs, which can cause deficits beyond synaptopathy. Furthermore, the impact of supernumerary synapses on auditory processing has not been evaluated. Here, we studied mice in which IHC synapse counts were increased or decreased by altering neurotrophin 3 (Ntf3) expression in IHC supporting cells. As we previously showed, postnatal Ntf3 knockdown or overexpression reduces or increases, respectively, IHC synapse density and suprathreshold ABR amplitude, without changing cochlear thresholds. Here, we show that IHC synapse density does not influence the magnitude of the acoustic startle reflex or its prepulse inhibition. In contrast, gap-prepulse inhibition, a behavioral test for auditory temporal processing, was enhanced or reduced according to Ntf3 expression levels. These results indicate that cochlear synaptopathy causes temporal processing deficits, which may be a key contributor to the hearing impairments that define HHL. Furthermore, the improvement in temporal acuity achieved by increasing Ntf3 expression and synapse density suggests a therapeutic strategy for improving hearing in noise for individuals with synaptopathy of various etiologies.

Introduction

The inner hair cells (IHCs) of the mammalian organ of Corti transduce sounds into neural activity through glutamatergic excitatory synapses onto primary auditory neurons, the type I spiral ganglion neurons. These synapses are vulnerable to aging and noise exposure (1), i.e.; even moderate noise exposure and/or normal aging result in loss of IHC synapses (2, 3). This type of deafferentation, a.k.a. cochlear synaptopathy, has been well documented in various mammalian species, including mice (1), guinea pigs (4), rats (5), gerbils (6), chinchillas (7) and humans (8, 9). Animal studies have shown that moderate cochlear synaptopathy caused by noise overexposure or aging causes "hidden hearing loss," i.e. an auditory neuropathy that manifests as a reduction in suprathreshold amplitudes of the first peak (peak I) of the auditory-brainstem response (ABR) without a shift in cochlear thresholds (1).

It has been hypothesized that cochlear synaptopathy leads to neural coding deficits that impair speech discrimination and intelligibility, especially in noisy environments, and to compensatory increases in the "gain" of central circuits that cause hyperacusis and tinnitus (10). Indeed, experiments on animals with noise- or drug-induced cochlear synaptopathy show evidence for persistent gain in central circuits, either in the enhancement of acoustic startle responses (11), or of sound-evoked response rates in neurons from the inferior colliculus or cortex (12, 13). Cortical responses from synaptopathic animals also show threshold elevations for detectability of tones in noise without changes in tone detection in quiet (13). However, other studies have failed to find such correlation (14). In humans, several studies have shown a relationship between noise-exposure history and reductions in ABR suprathreshold amplitudes among those with normal audiometric thresholds (15), or a correlation between ABR peak I amplitudes and performance on speech-in-noise tasks (16, 17), but these studies did not provide direct demonstration of cochlear synaptic loss. Furthermore, other studies have found no evidence for a relationship between noise history or ABR amplitudes and impaired speech perception (18).

To explore the impact of cochlear synapse density on auditory function, and to probe the effects of cochlear synaptopathy in the absence of cochlear insults like acoustic overexposure, ototoxic drugs or aging, where damage may be more widespread than the loss of cochlear synapses, we used transgenic mice in which IHC synapse density can be controlled via altering neurotrophin 3 (Ntf3) expression in the IHC's supporting cells (19). We previously showed that altering the levels of supporting-cell Ntf3 expression starting in the neonatal period permanently changes IHC synapse density, i.e., supporting-cell Ntf3 over-expression increases IHC synapse numbers and enhances ABR peak I amplitudes, whereas supporting-cell specific Ntf3 knock-out decreases them (19). Here, we tested the behavioral phenotypes of these transgenic mice, focusing on the acoustic startle reflex and its modulation. For the latter, we measured prepulse inhibition (PPI) and gap-inhibition of the acoustic startle (GPIAS), behaviors used to assess stimulus salience, sensory gating and temporal processing in both animals and humans (20-26).

We found that, compared to control animals, GPIAS is stronger in animals with increased IHC synapse density (and Ntf3 overexpression) and weaker in animals with decreased synapse density (and Ntf3 knockdown), without any changes in the strength of the acoustic startle itself or its PPI. As seen with noise- or age-related loss of synapses (5, 11), the later ABR peaks, originating from auditory brainstem and midbrain nuclei, are unchanged after Ntf3 knockdown, despite a decrease in peak I amplitudes, suggesting a compensatory central gain. In Ntf3 overexpressors, ABR peaks I-IV are enhanced, suggesting there are no mechanisms for central gain reduction after a peripheral gain of function. Together, these results are consistent with the notion that enhancing IHC synapse density could lead to a general enhancement of temporal processing abilities.

Results

Regulating IHC synapse density and sound-evoked auditory nerve activity via Ntf3 knock-down or over-expression.

To modify the levels of Ntf3 expression in IHC supporting cells, we used cell-specific inducible gene recombination as in prior studies (19). Briefly, the Plp1-CreERT transgenic line (27) was used to drive gene recombination in IHC supporting cells via tamoxifen treatment during the neonatal period (28). This CreERT transgene was combined with either conditional Ntf3 KO alleles (Ntf3$^{flox/flox}$)(29) to reduce Ntf3 expression in these cells, or an inducible Ntf3 over-expression transgene (Ntf3^{STOP}) to increase it (19). Since Ntf3 is expressed by both IHCs and their surrounding supporting cells in the cochlea (30), knockout of the Ntf3 gene from IHC supporting cells in Ntf3$^{flox/flox}$:Plp1-CreER^{T} reduces cochlear Ntf3 level but does not eliminate it (19). As controls, we used mice with the conditional Ntf3 alleles without the CreERT transgene.

Between the ages of 8 and 15 weeks, mutant and control mice underwent a variety of behavioral and physiological tests (Fig. 1). At 16 weeks, cochlear tissues were harvested to measure Ntf3 expression levels and the number of synapses per IHC (synapse density). We present the latter analyses first, which show that the Ntf3 manipulations had the expected effects on the cochlea.

Quantitative RT-PCR (Fig. 2A, B) confirmed that cochlear Ntf3 levels were reduced in the Ntf3$^{flox/flox}$:Plp1-CreER^{T}, i.e., Ntf3 Knockdown (Ntf3-KD) mice and increased in the Ntf3^{STOP}:Plp1-CreER^{T}, i.e., Ntf3 Overexpressor (Ntf3-OE), mice. Similarly, expression of VGF mRNA, a gene downstream of neurotrophin receptor signaling (31), was decreased in the cochleas of Ntf3-KD mice (Fig. 2A) and increased in Ntf3-OE mice (Fig. 2B). Furthermore, there was a clear correlation between the mRNA levels of Ntf3 and VGF (Fig. 2C), indicating that the changes in Ntf3 expression impact TrkC signaling in the inner ear. Since the Plp1 gene is also expressed in oligodendrocytes in the brain, and auditory-driven behaviors such as GPIAS are modulated by cortical circuits (32), we also measured the levels of Ntf3 and VGF in the cerebral cortex. We found a small decrease in cortical Ntf3 mRNA levels in Ntf3-KD mice (Fig. 2D), no change in Ntf3-OE mice (Fig. 2E), and most importantly, no changes in VGF mRNA levels in either of the

mutants (Fig. 2D, E), indicating that the manipulation of Ntf3 expression in Plp1-expressing cells is unlikely to have a direct effect on the central nervous system.

We also immunostained cochleas to assess the numbers of synapses between IHCs and auditory-nerve fibers. Whereas each auditory-nerve fiber contacts a single IHC, each IHC is contacted by numerous auditory-nerve fibers. Each of these glutamatergic synaptic contacts can be identified as a closely apposed pair of puncta in cochleas immunostained for CtBP2, a major component of the presynaptic ribbon, and GluA2, a subunit of the glutamate receptors localized at the post-synaptic terminals (Fig. 3A, E). In a normal mouse, the mean number of synapses per IHC follows an inverted U-shaped function, peaking in mid-cochlear regions at a value of roughly 20 synapses per IHC (2, 19). Here, after normalizing the synaptic counts to the mean peak value in controls, we observed that, as we previously reported (19), reduced supporting-cell Ntf3 expression levels decreased IHC synapse density in the basal half of the cochlea by as much as 20%, while increased supporting-cell Ntf3 expression levels increased IHC synapse density in the same cochlear regions by as much as 30% (Fig. 3 B-D, F-H). The stronger impact of supporting-cell derived Ntf3 on synapse density in the basal half of the cochlea likely reflects the fact that endogenous Ntf3 levels are lower in this region (30), making it more sensitive to changes in expression.

At 8 weeks of age, we assessed cochlear function by measuring distortion product otoacoustic emissions (DPOAEs), which reflect outer hair cell function, and ABRs, which reflect the summed responses of the auditory nerve and several higher auditory centers (33). While peak I reflects synchronous sound-evoked activity of auditory-nerve fibers, the second peak (peak II) is dominated by contributions from cochlear-nucleus bushy cells, and later waves represent bushy-cell targets in the superior olivary complex and inferior colliculus (33-37).

These recordings confirmed that, as we reported earlier (19), reduced Ntf3 expression by IHC supporting cells reduces ABR peak I amplitudes without changing ABR and DPOAE thresholds

(Fig. 4A-C), while increasing Ntf3 leads to normal thresholds with increased peak I amplitudes (Fig. 4D-F). Like the changes in synapse density, the effects of Ntf3 levels on peak I amplitudes is stronger in the middle and high frequencies, which reflect responses arising from the middle/basal cochlear regions.

Whereas peak I amplitude was reduced in mice with reduced Ntf3 cochlear expression, later ABR peak amplitudes were normal in these mice (Fig. 5A), suggesting that decreased sound-evoked activity of the auditory nerve leads to central gain in several higher auditory centers, as seen after noise-induced or age-related synaptic loss (5, 11). In contrast, Ntf3 overexpression resulted in increased amplitudes for ABR peaks I-IV (Fig. 5B). All peak latencies were normal in both mutants (Fig. 6A and B), suggesting that conduction velocity was not affected by the altered Ntf3 levels and synapse densities.

IHC synapse density influences auditory processing but not the startle reflex or sensory gating.

To examine the impact of IHC synapse density on auditory processing, we tested three auditory-driven behaviors; the acoustic startle response (ASR) (38), and two behaviors that asses modification of the ASR, Prepulse Inhibition of the ASR (PPI) (39), and Gap-Prepulse Inhibition of the ASR (GPIAS) (40). These tests have been a mainstay of studies on temporal processing and hearing in noise deficits in animal models (22, 41-43).

The ASR is a reflexive and rapid burst of muscular activity in response to a sudden, brief, and intense sound. This is a robust and consistent behavior, easily quantified by measuring the whole-body startle response (44). To investigate the impact of alterations in IHC synapse density on the ASR, we measured the response to a high-intensity startle stimulus (20 ms 120 dB SPL broadband noise burst) (Fig. 7B1). We also measured responses to moderate-intensity narrowband stimuli (50 ms 65 dB SPL) centered at different frequencies (8, 12, 16, 24 and 40 kHz) (Fig. 7A1), to detect hyper-responsiveness to innocuous sound and to determine if

responses to the prepulse stimuli are altered by changes in Ntf3 levels. As expected, the moderate-intensity prepulse stimulus alone did not induce startle responses in either mutant or control mice (Fig. 7A2, A3). The high-intensity startle stimulus elicited strong responses with amplitudes that were not different between control and mutants(Fig. 7B2, B3). These results indicated that that changes in Ntf3 expression, synapse density, and ABR peak I amplitudes, do not affect reflexive motor responses to sound.

PPI is commonly used to assess sensorimotor gating, i.e., the ability of a sensory stimulus to suppress a motor response (26). The PPI assay is quantified as the decrease in the ASR when a prepulse stimulus is presented a few milliseconds before the startle stimulus (45). We used the 50 ms pre-pulse stimulus described above ending 50 ms prior to the startle stimulus (Fig. 7C1) and quantified the magnitude of PPI as the fractional ASR reduction, i.e., 1 minus the ratio of startle magnitude with and without prepulse (45). On average, the prepulse inhibited the startle response by 40%. There was no significant difference between control and mutant mice in PPI (Fig. 7C2, C3), suggesting that changes in synapse density and associated changes in auditory-nerve activity do not affect sensorimotor gating.

Finally, we measured GPIAS, which is used to examine auditory temporal processing (20, 22, 25, 46), and correlates with speech recognition in humans (47, 48). In GPIAS, the suppression of the ASR is induced by the presentation of a brief silent gap in a continuous background noise instead of a mild sound stimulus in silence. As with the PPI, we quantified GPIAS as the fractional reduction of startle, i.e., 1 minus the ratio of the startle magnitude with and without a gap.

To quantify gap detection threshold in GPIAS, we presented animals in broadband background noise (BBN) with gaps of various durations (3-50 ms) ending 50 ms before the startle stimulus (Fig. 8A). Conventionally, the gap duration that elicits 50% of the maximal inhibition level is defined as gap-detection threshold (41). Gap-detection threshold is usually used to measure the

temporal acuity for acoustic transients. Consistent with previous reports (22), the GPIAS suppression level increased with increasing gap duration (Fig. 8B, C). Remarkably, gap detection threshold was increased in Ntf3-KD mice and reduced in Ntf3-OE mice (Fig. 8D, E), indicative of deficits in temporal acuity in Ntf3-KD mice and enhancement in Ntf3-OE mice.

To investigate if the changes in Ntf3 expression levels and IHC synapse density affect GPIAS, we measured gaps of 50-ms duration in narrowband noise (NBN) around different center frequencies (6-10 kHz, 10-14 kHz, 14-18 kHz, 22-26 kHz, and 38-42 kHz; spectral width: 4 kHz) (Fig. 9A). Whereas Ntf3 expression levels did not alter the startle responses at any frequency (Fig. 9B, C), there were clear significant differences in GPIAS, i.e., Ntf3-KD mice showed reduced gap inhibition and Ntf3-OE showed stronger gap inhibition with NBN background (Fig. 9D, E). These results indicate that the difference in suppression level caused by altered Ntf3 expression cannot be attributed to impact on startle amplitudes, and that the influence of Ntf3 expression levels on gap inhibition level is not restricted to any specific frequency.

Discussion

Self-reporting studies indicate that 10% to 20% of adult humans with normal audiometric thresholds have hearing difficulties (49-52). Furthermore, humans with sensorineural hearing loss have longer gap-detection thresholds that correlate with poorer speech perception (53-55). It has been proposed that cochlear synaptopathy could be the cause of this highly prevalent auditory dysfunction (3, 11, 15). However, since the presence of synaptopathy in living human subjects cannot be assessed directly, and the impact of synapse density on auditory processing had not been specifically tested in animal models without confounders such as noise-exposure or aging, the causal link between IHC synatopahty and HHL remained hypothetical. Our findings using mice with reduced or enhanced IHC synapse density due to changes in Ntf3 expression provide direct evidence that auditory processing is influenced by cochlear synapse density and that IHC synaptopathy is sufficient to degrade auditory processing and temporal acuity.

Furthermore, our results suggest that increased IHC density leads to enhanced temporal acuity, raising the possibility that therapies that increase IHC synapse density could improve auditory performance in humans.

Analysis of ABR waveforms in our mouse models also provides new insights into the impact of IHC synapse density on the function of the ascending auditory pathway. We found that ABR peaks II-V are normal in mice with IHC synaptopathy induced by Ntf3 knockdown despite the reduced peak I amplitude. This is similar to the finding in animals with cochlear damage (3, 11-13, 56), which has been interpreted as reflecting homeostatic compensation in the auditory brainstem and midbrain to partial peripheral deafferentation. Our results provide further support for this view. Most surprising was the observation that ABR peak I-IV amplitudes are increased in mice with supernumerary IHC synapses. This finding indicates that increased IHC synapse density enhances sound-evoked signaling to supranormal levels along the ascending auditory pathway. Our prior work showed that Ntf3-induced increase in synaptic counts is not associated with a proliferation of auditory nerve fibers (19). Thus, we presume that a fraction of fibers now must make more than just one synaptic contact with an IHC. This, in turn, might lead to better synchronization of onset responses and higher ABR amplitudes, which are then propagated through the ascending central circuitry. It is possible that increased synchrony contributes to the improved detection of noise gaps in the Ntf3 over-expressing mice, suggesting that creating supernumerary IHC-SGN synapses could also improve auditory performance in a wider range of stimulus contexts.

Since the Plp1-CreERT transgene also drives gene recombination in oligodendrocytes in the central nervous system (CNS) (27), and auditory cortical circuits contribute to gap detection (32), it was important to determine if the mice had significant changes in Ntf3 expression and signaling in the CNS. Finding little or no change in Ntf3 and VGF expression in the cortex of Ntf3 KD and OE mice, together with data showing that few CNS neurons express the Ntf3

receptor TrkC in the brain (see Expression Summary in (57)) suggests that the Ntf3 manipulations we created did not directly affect cortical circuits. Thus, the phenotypes seen here are most likely dominated by changes originating with the altered IHC synapse density.

Interestingly, mice with IHC synaptopathy due to Ntf3 knockdown did not show enhanced magnitude of the baseline acoustic startle or PPI. This contrasts with observations in mice with synaptopathy induced by noise (11) or aging (45), where the hyper-responsiveness has been interpreted as a sign of hyperacusis, a clinical disorder associated with noise-induced or age-related hearing loss, in which moderate-intensity sounds become intolerably loud (23, 45, 58-60). We see several potential explanations for this apparent discrepancy. First, whereas the noise- or age-induced acoustic trauma occurs after the circuits are fully formed, the synapse reduction in our model originates during the neonatal period, when central circuits are still developing, raising the possibility that some aspects of auditory processing adapt to the early onset of the synaptopathy, preventing the development of hyper-responsiveness. Second, whereas the extent of synaptic loss in our model is 20–30%, the other studies involved more extensive cochlear damage (11, 45), which could lead to more dramatic central dysfunction. Third, noise exposure induces stress or fear, which can play a role in hyperacusis-like behavior (61).

The observation that alterations in IHC synapse density influence performance in GPIAS but not in PPI likely reflects the different neural mechanism responsible for these sound-driven modifications of the ASR, which are mediated by higher order nuclei (62, 63). During ASR, the auditory-nerve inputs activate cochlear root neurons that project to nucleus reticularis pontis caudalis (PnC) neurons, which then excite the spinal motor neurons that elicit the whole-body startle response (64, 65). Prepulses and gaps are relayed from the cochlear root neurons to the inferior colliculus (IC) (66), but their pathways diverge thereafter. Pre-pulse processing has been shown to involve the lateral globus pallidus (LGP) whereas gap processing involves the auditory

cortex (AC) (67, 68). Our results suggest that alterations in synapse density might affect primarily the latter. Further studies on the impact of synapse density on these circuits could provide new insights into the mechanisms by which peripheral responses influence central auditory processing.

We previously demonstrated that increasing cochlear Ntf3 availability after noise exposure induces IHC synapse regeneration and recovery of ABR peak I amplitudes (19, 69, 70). More recently we showed that increasing cochlear Ntf3 expression levels in the middle-aged mouse acutely increases ABR peak I amplitudes and slows the progression of age-related synaptopathy, preserving the IHC synapse density of middle age until the end of the lifespan (71). The results from the current study suggest that Ntf3-based therapies could not only promote IHC synapse health and numbers, but also improve auditory processing after noise trauma or in aging.

In summary, our finding that cochlear synaptopathy elicited without cochlear insults such as noise or ototoxic drugs results in temporal processing deficits further supports the notion that synaptopathy is a key contributor to the impaired speech perception experienced by many with hidden hearing loss. Furthermore, the improvement in temporal acuity achieved by increasing Ntf3 expression and synapse density suggests a therapeutic strategy for improving hearing in noise for individuals with synaptopathy of various etiologies.

Materials and methods

Animals

All experimental procedures complied with the National Institutes of Health guidelines and were approved by the Institutional Animal Care and Use Committee of University of Michigan, MI, USA. Cochlear supporting-cell specific Ntf3 knock-down mice (Ntf3 KD) derived from $Ntf3^{flox/flox}$:Plp1/CreER^T mice by Cre-recombination) and Ntf3 overexpressing mice (Ntf3 OE derived from $Ntf3^{stop}$:Plp1/CreER^T mice by Cre-recombination) were generated as previously described (19). *Ntf3 KD* mice and their controls were maintained on C57BL/6 background. *Ntf3 OE* and their controls were on FVB/N background. Both male and female mice were included in this study.

Tamoxifen administration

Tamoxifen was injected into intraperitoneal cavity of P 3-10 Ntf3^{stop}: Plp1/CreER^T mice or P1-3 Ntf3$^{flox/flox}$:Plp1/CreER^T mice as previously described (19). A 10 mg/ml solution of tamoxifen was obtained by dissolution in corn oil. Injection was 33 mg/kg for Ntf3^{stop}: Plp1/CreER^T mice and 50 mg/kg for Ntf3$^{flox/flox}$:Plp1/CreER^T mice.

Real-time quantitative RT-PCR.

Total RNA was isolated from the cortical brain and cochlea samples from 1-month-old mice using RNA extraction kit and Qiazol Reagent (RNeasy mini kit; Qiagen, Germany), and DNase treatment was performed (RNase-free; Qiagen). The complementary DNA was synthesized using iScript cDNA synthesis kit (Bio-Rad, #1708891, USA), according to the manufacturers' protocol. Quantitative RT-PCR was performed on a CFX-96 Bio-Rad reverse transcription polymerase chain reaction detection system (Hercules, CA, USA) using iTaq Universal SYBR® Green supermix (Bio-Rad, # 172-5121, USA) and primer pairs were synthesized by IDT (Coralville, IA, USA). All samples and standard curves were run in triplicate. Water instead of

complementary DNA was used as a negative control. The 10 µl reaction contained 5 µl of SYBR

Green supermix, 6 pmol of each forward and reverse primer (0.6 µl), 1.9 µl nuclease-free of

water and 2.5 µl of cDNA sample. The mRNA expression levels in PLP1cre ERT: NT3fl or

PLP1cre ERT: NT3 STOP versus their control mice were determined by a comparative cycle

threshold (Ct) method and relative gene copy number was calculated as normalized gene

expression, defined as described previously (72). Ribosomal protein L19 (RPL19) was used as

the housekeeping gene. The following specific oligo primers were used for the target genes:

Rpl19, F: 5'ACCTGGATGAGAAGGATGAG 3'; R: 5'ACCTTCAGGTACAGGCTGTG 3'; Ntf3, F

5'GCCCCCTCCCTTATACCTAATG 3'; R: 5'CATAGCGTTTCCTCCGTGGT 3'; Vgf: F:

5'GGTAGCTGAGGACGCAGTGT 3'; R: 5'GTCCAGTGCCTGCAACAGT 3'. Changes in mRNA

expression were calculated as relative expression (arbitrary units) respective to the control

group for each mouse line.

Immunostaining and synaptic counts

Cochleas from 16-week-old mice were prepared for whole-mount imaging as described in (19).

In brief, the samples were fixed with 4% formaldehyde for 2 hours and then decalcified with 5%

ethylenediaminetetraacetic acid (EDTA) for 3-5 days. Cochlear epithelia were micro-dissected

into five segments for whole mount processing. The cochlea segments were permeabilized by

freeze–thawing in 30% and then blocked with 5% normal horse serum for 1 hour. Afterwards,

the following primary antibodies were used: 1) CtBP2 to visualize synaptic ribbons (mouse anti-

CtBP2 at 1:200, BD Biosciences, catalog # 612044, RRID: AB_399431), 2) GluR2 to visualize

postsynaptic receptors (mouse anti-GluR2 at 1:2000, Millipore, catalog # MAB397, RRID:

AB_2113875) and 3) Myosin VIIa to visualize IHCs (rabbit anti-Myosin VIIa at 1:200; Proteus

Biosciences, catalog # 25-6790, RRID: AB_10015251). Secondary antibodies used were Alexa

Fluor 488 conjugated anti-mouse IgG2a (1:1000, Invitrogen), Alexa Fluor 568 conjugated anti-

mouse IgG1 (1:1000, Invitrogen) and Alexa Fluor 647 conjugated anti-rabbit (1:1000, Life

Technologies). Frequency maps were created using a custom ImageJ plug-in. Images were captured from 5.6 to 45.2 kHz using a Leica SP8 with a 1.4 NA 63x oil immersion objective at 3 digital zoom. Offline image analysis was performed using Amira (Visage Imaging). Quantifications of synapses were done by a blind investigator.

Distortion Product Otoacoustic Emissions (DPOAE) and Auditory Brainstem Responses (ABR)

DPOAEs and ABRs were performed as previously described (19). Mice were anaesthetized by i.p. injections of xylazine (20 mg kg^{-1}, i.p.) and ketamine (100 mg kg^{-1}, i.p.). The DPOAEs were induced by two primary tones (f1 and f2) and recorded at (2 × f1)–f2. f1 level was 10 dB higher than the f2 level and frequency ratio f2/f1 was 1.2. The ear-canal sound pressure was amplified and averaged at 4 μs intervals. DPOAE thresholds were defined as the f2 level that produced a response 10 dB SPL higher than the noise floor. For ABR measurement, subdermal electrodes were placed (a) at the dorsal midline of the head, (b) behind the left earlobe and (c) at the base of the tail (for a ground electrode). ABRs were evoked with 5 ms tone pips (0.5 ms rise–fall) delivered to the eardrum. The frequencies of tone pips were 5.6, 8, 11.3, 16, 22.6, 32 and 45.2 kHz, with 15 sound levels from 10 to 80 dB SPL for each frequency. The signals were amplified 10,000 times and filtered through a 0.3 - 3 kHz passband. At each level, the average of 1,024 signals were taken after 'artifact rejection'. Both recordings were performed using National Instruments input/output boards hardware. Offline analysis was performed using excel and ABR peak analysis software.

Pre-pulse inhibition (PPI) and gap inhibition of the acoustic startle (GPIAS)

Mice were tested in a 10 x 4.5 x 4 cm cage inside a sound-isolation chamber that were placed within a sound attenuating room. Acoustic sound source was located in the upper part of this chamber. The piezoelectric motion sensor attached to the cage detected the vertical force of the

startle reflex. All ASR, PPI and GPIAS stimuli and responses were generated and recorded with Kinder Scientific Startle Monitor (Kinder Scientific, Poway, CA).

PPI tests were used for assessing sensorimotor gating on 8-15 old week mice, twice a week, each 2 days apart. PPI tests were conducted quiet. The startle stimuli were BBN bursts at 120 dB SPL, 20 ms in duration, 0.1 ms rise-fall times. The prepulse was a narrow band sound centered at 8, 12, 16, 24 and 40 kHz, 50 ms in duration, with 2-ms rise-fall ramps (Fig. 7). PPI test consisted of prepulse trials and startle-only trials, which were delivered alternatively. In prepulse trials, a prepulse ended 50 ms before the startle stimulus. Startle-only trials were similar to the prepulse trials, but no prepulse was delivered. PPI startle ratio is the ratio of the startle magnitude in prepulse trials over the startle magnitude in startle-only trials.

The GPIAS paradigm has been used for measuring auditory temporal processing (20, 25). Gap inhibition was assessed on 8-15 old week mice, twice a week, 2 days apart. The testing consists of two types of trials, gap trials and no-gap trials (Fig. 8A, 9A) that were delivered alternatively. In both trials, the startle stimulus was 20 ms BBN at 120 dB with 0.1 ms rise/fall times. The startle was preceded either by gaps with varied durations (3-, 5-, 15-, 25-, or 50-ms long) embedded in BBN or by a 50-ms gap embedded in a narrow bandpass background sound centered at 8, 12, 16, 24, and 40 kHz at 65 dB. For gaps with varied durations, the response was normalized to the longest gap (50 ms) on the presumption that the maximum level of inhibition (100%) was elicited at the longest gap. The responses and gap durations were fitted with three-parameter logistic regression models. The gap-detection threshold was defined as the value of the fitted curve that elicited 50% of the maximal inhibition achieved per mouse. Gap-startle ratio is the ratio of the startle magnitude in gap trials over the startle magnitude of the paired no-gap trials.

For each mouse, PPI and gap-startle ratios were averaged from 11 sessions. Each session of PPI and gap-PPI test included 60 pairs of prepulse and startle-only trials for PPI or gap and no-

gap trials for gap detection (5 prepulse or background sound frequencies, 12 pairs for each frequency). The interval between trials was randomly varied between 5 and 15s. Each session began with a 2-min acclimatization period in the cage before startle testing began, and all tests were conducted in darkness. Startle-only trials and no-gap trial amplitudes greater than or equal to mean ± 2.5 standard deviations were eliminated. When a trial was eliminated, its paired trial was also eliminated. PPI ratio bigger than 1 or gap-startle ratios bigger than 1.1 were excluded (73). PPI and gap-startle ratios were calculated as the average with prepulse startle amplitude divided by the mean without prepulse startle amplitude.

Statistical analyses

Analyses were performed using GraphPad Prism 6 (GraphPad Software Inc., La Jolla, CA) and RStudio packages. Data are shown as mean and standard error of the mean (S.E.M.). The number of replicates (n) is indicated in the results section and figure legends. No explicit power analysis was used to predetermine sample sizes, but our sample sizes are similar to those reported in our previous publications. Statistical differences in auditory physiology (DPOAE threshold, ABR threshold, amplitude, and latency), ribbon synapse counts, behavioral background movement, PPI ratio, gap-startle ratios were analyzed using two-way ANOVA, followed by Bonferroni multiple comparisons test. mRNA expression, ASR amplitude and gap detection threshold were compared using unpaired Student's *t*-test. Correlations were computed using Pearson's correlation. Statistical threshold was set to alpha = 0.05.

Figure legends

Figure 1. **Timeline of the experiments.** Experimental timeline showing the ages of mice for tamoxifen treatments, ABR measurements, behavioral assay (ASR, PPI, GPIAS), and sample collections for quantitative RT-PCR and immunostaining (P = postnatal day, wk = weeks).

Figure 2. Altering Ntf3 expression levels in Plp1-expressing cells impacts TrkC signaling in the cochlea, **but not in the CNS.** mRNA level of Ntf3 and VGF, a gene downstream of TrkC signaling, are reduced in Ntf3-KD cochleas (**A**) and increased in Ntf3-OE cochleas (**B**). Furthermore, cochlear of Ntf3 and VGF mRNA levels are correlated (**C**). In contrast, cortical Ntf3 mRNA level is slightly decreased in Ntf3-KD mice (**D**) and unchanged in Ntf3-OE mice (**E**). No changes in VGF mRNA levels are observed in the brains of either Ntf3-KD or Ntf3-OE mice (**D,E**). n=5-8, ns = p > 0.05, * p < 0.05, ** p < 0.01, mRNA levels were compared by two-tailed unpaired t test. Error bars represent SEM.

Figure 3. Ntf3 regulates IHC synapse density. Representative confocal images of IHC synapses at the 16 kHz cochlear region from Ntf3-KD (**A**) and Ntf3-OE (**E**) mice and their respective controls immunolabeled for pre-synaptic ribbons (CtBP2 - red), post-synaptic receptor patches (GluA2 - green) and hair cells (Myo7a - blue). Mean counts (± SEM) of ribbons (**B, F**), GluA2 patches (**C, G**) and colocalized markers (**D, H**) in Ntf3 KDs and OEs. n = 5, ns = p > 0.05, * p < 0.05, ** p < 0.01, *** p < 0.001, *** p < 0.0001. Synaptic markers were compared by two-way ANOVA.

Figure 4. Ntf3 knockdown or overexpression influence ABR peak I amplitudes without effecting cochlear thresholds. DPOAE (**A, D**) and ABR (**B, E**) thresholds in Ntf3-KD and Ntf3-

OE mice are not different than their controls. In contrast, Ntf3 knockdown reduces ABR P1 amplitudes (**C**) whereas overexpression leads to increased peak I amplitudes (**F**). n = 15-24. Normalized ABR P1 amplitudes were assessed at 80 dB SPL. ns = p > 0.05, *p < 0.05, **p < 0.01, ***p < 0.001, ***p < 0.0001 by two-way ANOVA. Error bars represent SEM.

Figure 5. Ntf3 knockdown or overexpression have different effects on the input-output function of ABR peaks I to IV. Mean amplitude-vs level functions for ABR peaks I-IV in Ntf3-KD (**A**) and Ntf3-OE (**B**) mice and their respective controls at 16 kHz. Whereas peak I amplitudes are reduced in Ntf3-KD mice, the amplitudes of the other peaks remain normal, indicative of central compensation (**A**). In contrast, Ntf3 overexpression increases amplitudes of ABR peaks I to IV (**B**). N = 14-20, ns = p > 0.05, *p < 0.05, **p < 0.01, ***p < 0.001, ***p < 0.0001 by two-way ANOVA. Error bars represent SEM.

Figure 6. Ntf3 expression levels do not influence the latencies of the ABR waveform peaks. Plots of peak latency recorded at 16 kHz against sound stimulus level show that latencies of ABR peaks I-V are not altered by Ntf3-KD (**A**) and Ntf3-OE (**B**). n=14-20, ns = p > 0.05 by two-way ANOVA. Error bars represent SEM.

Figure 7. IHC synapse density does not influence the acoustic startle response or prepulse inhibition. Schematics of the protocols for prepulse stimulus (**A1**), startle stimulus (**B1**) and PPI stimuli (**C1**). (**A1**) The noise prepulse stimulus is a narrowband noise (4 kHz width around variable center frequencies, 65 dB SPL, 50 ms duration). (**B1**) The startle stimulus is a broadband noise (120 dB SPL, 20 ms duration). (**C1**) PPI consists of a noise prepulse and a startle stimulus that starts 50 ms after the prepulse. (**A2**, **A3**) Reactivity to prepulse is not

significantly different between mutant and control mice. (**B2, B3**) Loud sound (120 dB SPL) elicits startle responses with amplitudes that were similar in mutant mice and their control littermates. (**C2, C3**) The degree of prepulse inhibition of the startle response by a prepulse was determined using the formula $\% \, inhibition = \frac{startle \, response - prepulse \, response}{startle \, response} X100$. On average, the prepulse inhibit the startle response by 40%. There is no significant difference between control and mutant mice. N=18-20 mice/group for response to prepulse (**A2**, **A3**); n=20-24 mice/group for startle response (**B2**, **B3**); n=15-20 mice/ group for PPI (**C2**, **C3**). ns = p > 0.05 by two-tailed unpaired t tests (**B2**, **B3**) or two-way ANOVA (**C2**, **C3**). Mean ± SEM are shown.

Figure 8. Ntf3 expression levels influence gap detection thresholds in broadband background noise. (**A**) Schematic depiction of No gap trials (left) and gap trials (right). No gap trials consisted of a startle sound (120 dB SPL, 20 ms duration) in continuous noise background (broadband noise, BBN, 65 dB SPL). Gap trials consisted of a continuous broadband noise background and a gap of variable length (0-50 ms) ending 50 ms before the startle stimulus (S). (**B**, **C**) show the level of gap inhibition vs. gap length for Ntf3 KD and OE mice, respectively. The inhibition of the startle reflex increases as the gap duration increases. Dotted line represents the gap threshold (defined as the gap length that elicits half-maximal inhibition) in control mice. **(D, E)** show gap detection thresholds. Gap detection threshold is increased in Ntf3-KD mice (**D**) and decreased in Ntf3-OE mice **(E)**. n=9-17 mice/group, *p < 0.05, **p < 0.01 by t-test. Mean ± SEM are shown.

Figure 9. Ntf3 expression levels influence gap inhibition in narrowband background noise. (**A**) A schematic view of No gap trials (left) and gap trials (right). No gap trials consist of a startle sound (120 dB SPL, 20 ms duration) presented in continuous noise background

(narrowband noise, NBN, 4 kHz width around variable center frequencies, 65 dB SPL). Gap-prepulse inhibition of the acoustic startle (GPIAS) was tested in gap trials with the same background noise with a 50 ms gap included as a prepulse followed 1 ms later by the startle-eliciting stimulus. (**B**, **C**) Responses to startle stimulus in continuous NBN background are unaffected in the Ntf3-KD or Ntf3-OE mice. (**D**, **E**) GPIAS in narrowband noises were significantly weakened in Ntf3-KD mice and strengthened in Ntf3-OE mice in a two-way ANOVA. Sidak multiple comparison tests revealed a significant reduction at frequency of background noise band 10-14, 14-18 and 22-26 kHz in Ntf3-KD mice and no frequency-specific changes in Ntf3-OE mice (not shown in the figure). n=10-15 mice/group, *p < 0.05, **p < 0.01 by two-way ANOVA. mean ± SEM are shown.

References

1. Kujawa SG, Liberman MC. Synaptopathy in the noise-exposed and aging cochlea: Primary neural degeneration in acquired sensorineural hearing loss. Hearing Research. 2015;330:191-9.

2. Kujawa SG, Liberman MC. Adding Insult to Injury: Cochlear Nerve Degeneration after "Temporary" Noise-Induced Hearing Loss. J Neurosci. 2009;29(45):14077-85.

3. Sergeyenko Y, Lall K, Liberman MC, Kujawa SG. Age-Related Cochlear Synaptopathy: An Early-Onset Contributor to Auditory Functional Decline. J Neurosci. 2013;33(34):13686-94.

4. Lin HW, Furman AC, Kujawa SG, Liberman MC. Primary Neural Degeneration in the Guinea Pig Cochlea After Reversible Noise-Induced Threshold Shift. J Assoc Res Otolaryngol. 2011;12(5):605-16.

5. Möhrle D, Ni K, Varakina K, Bing D, Lee SC, Zimmermann U, et al. Loss of auditory sensitivity from inner hair cell synaptopathy can be centrally compensated in the young but not old brain. Neurobiology of Aging. 2016;44:173-84.

6. Gleich O, Semmler P, Strutz J. Behavioral auditory thresholds and loss of ribbon synapses at inner hair cells in aged gerbils. Experimental Gerontology. 2016;84:61-70.

7. Hickman TT, Smalt C, Bobrow J, Quatieri T, Liberman MC. Blast-induced cochlear synaptopathy in chinchillas. Scientific Reports. 2018;8(1):10740.

8. Viana LM, O'Malley JT, Burgess BJ, Jones DD, Oliveira CACP, Santos F, et al. Cochlear neuropathy in human presbycusis: confocal analysis of hidden hearing loss in post-mortem tissue. Hearing Research. 2015;327:78-88.

9. Wu PZ, Liberman LD, Bennett K, de Gruttola V, O'Malley JT, Liberman MC. Primary neural degeneration in the human cochlea: evidence for hidden hearing loss in the aging ear. Neuroscience. 2019;407:8-20.

10. Liberman MC, Kujawa SG. Cochlear synaptopathy in acquired sensorineural hearing loss: Manifestations and mechanisms. Hearing Research. 2017;349:138-47.

11. Hickox AE, Liberman MC. Is noise-induced cochlear neuropathy key to the generation of hyperacusis or tinnitus? Journal of Neurophysiology. 2014;111(3):552-64.

12. Asokan MM, Williamson RS, Hancock KE, Polley DB. Sensory overamplification in layer 5 auditory corticofugal projection neurons following cochlear nerve synaptic damage. Nat Commun. 2018;9(1):2468.

13. Resnik J, Polley DB. Cochlear neural degeneration disrupts hearing in background noise by increasing auditory cortex internal noise. Neuron. 2021;109(6):984-96 e4.

14.   Chen H, Shi L, Liu L, Yin S, Aiken S, Wang J. Noise-induced Cochlear Synaptopathy and Signal Processing Disorders. Neuroscience. 2019;407:41-52.

15.   Liberman MC, Epstein MJ, Cleveland SS, Wang H, Maison SF. Toward a Differential Diagnosis of Hidden Hearing Loss in Humans. PLOS ONE. 2016;11(9):e0162726.

16.   Bramhall N, Ong B, Ko J, Parker M. Speech Perception Ability in Noise is Correlated with Auditory Brainstem Response Wave I Amplitude. J Am Acad Audiol. 2015;26(5):509-17.

17.   Grant KJ, Mepani AM, Wu P, Hancock KE, de Gruttola V, Liberman MC, et al. Electrophysiological markers of cochlear function correlate with hearing-in-noise performance among audiometrically normal subjects. J Neurophysiol. 2020;124(2):418-31.

18.   Guest H, Munro KJ, Prendergast G, Millman RE, Plack CJ. Impaired speech perception in noise with a normal audiogram: No evidence for cochlear synaptopathy and no relation to lifetime noise exposure. Hearing Research. 2018;364:142-51.

19.   Wan G, Gómez-Casati ME, Gigliello AR, Liberman MC, Corfas G. Neurotrophin-3 regulates ribbon synapse density in the cochlea and induces synapse regeneration after acoustic trauma. Elife. 2014;3.

20.   Fournier P, Hebert S. The gap prepulse inhibition of the acoustic startle (GPIAS) paradigm to assess auditory temporal processing: Monaural versus binaural presentation. Psychophysiology. 2021;58(3):e13755.

21.   Jafari Z, Kolb BE, Mohajerani MH. Prepulse inhibition of the acoustic startle reflex and P50 gating in aging and alzheimer's disease. Ageing Res Rev. 2020;59:101028.

22.   Moore S, Meschkat M, Ruhwedel T, Trevisiol A, Tzvetanova ID, Battefeld A, et al. A role of oligodendrocytes in information processing. Nat Commun. 2020;11(1):5497.

23.   Thomas ME, Guercio GD, Drudik KM, de Villers-Sidani É. Evidence of Hyperacusis in Adult Rats Following Non-traumatic Sound Exposure. Front Syst Neurosci. 2019;13:55.

24.   Salloum RH, Yurosko C, Santiago L, Sandridge SA, Kaltenbach JA. Induction of enhanced acoustic startle response by noise exposure: dependence on exposure conditions and testing parameters and possible relevance to hyperacusis. PLOS ONE. 2014;9(10):e111747.

25.   Turner JG, Brozoski TJ, Bauer CA, Parrish JL, Myers K, Hughes LF, et al. Gap detection deficits in rats with tinnitus: A potential novel screening tool. Behavioral Neuroscience. 2006;120(1):188-95.

26.   Cryan FJ, Reif A. Behavioral Neurogenetics: Humana New York, NY; 2012.

27.   Doerflinger NH, Macklin WB, Popko B. Inducible site-specific recombination in myelinating cells. genesis. 2003;35(1):63-72.

28.   Gómez-Casati ME, Murtie J, Taylor B, Corfas G. Cell-Specific Inducible Gene Recombination in Postnatal Inner Ear Supporting Cells and Glia. JARO. 2010;11(1):19-26.

29.   Bates B, Rios M, Trumpp A, Chen C, Fan G, Bishop JM, et al. Neurotrophin-3 is required for proper cerebellar development. Nat Neurosci. 1999;2(2):115-7.

30.   Sugawara M, Murtie JC, Stankovic KM, Liberman MC, Corfas G. Dynamic patterns of neurotrophin 3 expression in the postnatal mouse inner ear. J Comp Neurol. 2007;501(1):30-7.

31.   Eagleson KL, Fairfull LD, Salton SR, Levitt P. Regional differences in neurotrophin availability regulate selective expression of VGF in the developing limbic cortex. J Neurosci. 2001;21(23):9315-24.

32.   Weible AP, Moore AK, Liu C, DeBlander L, Wu H, Kentros C, et al. Perceptual gap detection is mediated by gap termination responses in auditory cortex. Curr Biol. 2014;24(13):1447-55.

33.   Buchwald JS, Huang C. Far-field acoustic response: origins in the cat. Science. 1975;189(4200):382-4.

34.   Melcher JR, Guinan JJ, Knudson IM, Kiang NYS. Generators of the brainstem auditory evoked potential in cat. II. Correlating lesion sites with waveform changes. Hearing Research. 1996;93(1):28-51.

35.   Melcher JR, Kiang NYS. Generators of the brainstem auditory evoked potential in cat III: identified cell populations. Hearing Research. 1996;93(1):52-71.

36.   Melcher JR, Knudson IM, Fullerton BC, Guinan JJ, Norris BE, Kiang NYS. Generators of the brainstem auditory evoked potential in cat. I. An experimental approach to their identification. Hearing Research. 1996;93(1):1-27.

37.   Starr A, Hamilton AE. Correlation between confirmed sites of neurological lesions and abnormalities of far-field auditory brainstem responses. Electroencephalography and clinical neurophysiology. 1976;41(6):595-608.

38.   Curzon P, Zhang M, Radek RJ, Fox GB. The Behavioral Assessment of Sensorimotor Processes in the Mouse: Acoustic Startle, Sensory Gating, Locomotor Activity, Rotarod, and Beam Walking. In: Buccafusco JJ, editor. Methods of Behavior Analysis in Neuroscience. Frontiers in Neuroscience. 2nd ed. Boca Raton (FL): CRC Press/Taylor & Francis; 2009.

39.   Hoffman HS, Ison JR. Reflex modification in the domain of startle: I. Some empirical findings and their implications for how the nervous system processes sensory input. Psychol Rev. 1980;87(2):175-89.

40.   Fournier P, Hébert S. The gap-startle paradigm to assess auditory temporal processing: Bridging animal and human research. Psychophysiology. 2016;53(5):759-66.

41.	Barsz K, Ison JR, Snell KB, Walton JP. Behavioral and neural measures of auditory temporal acuity in aging humans and mice. Neurobiology of Aging. 2002;23(4):565-78.

42.	Fournier P, Hebert S. Gap detection deficits in humans with tinnitus as assessed with the acoustic startle paradigm: does tinnitus fill in the gap? Hear Res. 2013;295:16-23.

43.	Truong DT, Che A, Rendall AR, Szalkowski CE, LoTurco JJ, Galaburda AM, et al. Mutation of Dcdc2 in mice leads to impairments in auditory processing and memory ability. Genes Brain Behav. 2014;13(8):802-11.

44.	Davis M. The mammalian startle response. In: Eaton RC, editor. Neural Mechanisms of Startle Behavior: Springer New York, NY; 1984.

45.	Ison JR, Allen PD, O'Neill WE. Age-related hearing loss in C57BL/6J mice has both frequency-specific and non-frequency-specific components that produce a hyperacusis-like exaggeration of the acoustic startle reflex. J Assoc Res Otolaryngol. 2007;8(4):539-50.

46.	Felix RA, 2nd, Chavez VA, Novicio DM, Morley BJ, Portfors CV. Nicotinic acetylcholine receptor subunit alpha7-knockout mice exhibit degraded auditory temporal processing. J Neurophysiol. 2019;122(2):451-65.

47.	Heeke P, Vermiglio AJ, Bulla E, Velappan K, Fang X. The Relationship between Random Gap Detection and Hearing in Noise Test Performances. J Am Acad Audiol. 2018;29(10):948-54.

48.	Nakajima Y, Matsuda M, Ueda K, Remijn GB. Temporal Resolution Needed for Auditory Communication: Measurement With Mosaic Speech. Front Hum Neurosci. 2018;12:149.

49.	Tremblay KL, Pinto A, Fischer ME, Klein BE, Klein R, Levy S, et al. Self-Reported Hearing Difficulties Among Adults With Normal Audiograms: The Beaver Dam Offspring Study. Ear Hear. 2015;36(6):e290-9.

50.	Spankovich C, Gonzalez VB, Su D, Bishop CE. Self reported hearing difficulty, tinnitus, and normal audiometric thresholds, the National Health and Nutrition Examination Survey 1999-2002. Hear Res. 2018;358:30-6.

51.	Saunders GH, Haggard MP. The clinical assessment of obscure auditory dysfunction--1. Auditory and psychological factors. Ear Hear. 1989;10(3):200-8.

52.	Gates GA, Cooper JC, Jr., Kannel WB, Miller NJ. Hearing in the elderly: the Framingham cohort, 1983-1985. Part I. Basic audiometric test results. Ear Hear. 1990;11(4):247-56.

53.	Blankenship C, Zhang F, Keith R. Behavioral Measures of Temporal Processing and Speech Perception in Cochlear Implant Users. J Am Acad Audiol. 2016;27(9):701-13.

54.   Li B, Guo Y, Yang G, Feng Y, Yin S. Effects of Various Extents of High-Frequency Hearing Loss on Speech Recognition and Gap Detection at Low Frequencies in Patients with Sensorineural Hearing Loss. Neural Plast. 2017;2017:8941537.

55.   Yuvaraj P, Jayaram M. Audiological Profile of Adult Persons with Auditory Neuropathy Spectrum Disorders. J Audiol Otol. 2016;20(3):158-67.

56.   Knipper M, Van Dijk P, Nunes I, Ruttiger L, Zimmermann U. Advances in the neurobiology of hearing disorders: recent developments regarding the basis of tinnitus and hyperacusis. Prog Neurobiol. 2013;111:17-33.

57.   Allen Brain Map. Neurotrophic tyrosine kinase, receptor, type 3. 2020 June.

58.   Hayes SH, Radziwon KE, Stolzberg DJ, Salvi RJ. Behavioral models of tinnitus and hyperacusis in animals. Front Neurol. 2014;5:179.

59.   Sun W, Deng A, Jayaram A, Gibson B. Noise exposure enhances auditory cortex responses related to hyperacusis behavior. Brain Res. 2012;1485:108-16.

60.   Sun W, Lu J, Stolzberg D, Gray L, Deng A, Lobarinas E, et al. Salicylate increases the gain of the central auditory system. Neuroscience. 2009;159(1):325-34.

61.   Du Y, Wu X, Li L. Differentially organized top-down modulation of prepulse inhibition of startle. J Neurosci. 2011;31(38):13644-53.

62.   Meloni EG, Davis M. The dorsal cochlear nucleus contributes to a high intensity component of the acoustic startle reflex in rats. Hear Res. 1998;119(1-2):69-80.

63.   Yeomans JS, Frankland PW. The acoustic startle reflex: neurons and connections. Brain Res Brain Res Rev. 1995;21(3):301-14.

64.   Lee Y, Lopez DE, Meloni EG, Davis M. A primary acoustic startle pathway: obligatory role of cochlear root neurons and the nucleus reticularis pontis caudalis. J Neurosci. 1996;16(11):3775-89.

65.   Lingenhohl K, Friauf E. Giant neurons in the rat reticular formation: a sensorimotor interface in the elementary acoustic startle circuit? J Neurosci. 1994;14(3 Pt 1):1176-94.

66.   Fendt M, Li L, Yeomans JS. Brain stem circuits mediating prepulse inhibition of the startle reflex. Psychopharmacology (Berl). 2001;156(2-3):216-24.

67.   Bowen GP. Auditory Cortex Lesions in the Rat Impair Both Temporal Acuity and Noise Increment Thresholds, Revealing a Common Neural Substrate. Cerebral Cortex. 2003;13(8):815-22.

68.   Moreno-Paublete R, Canlon B, Cederroth CR. Differential Neural Responses Underlying the Inhibition of the Startle Response by Pre-Pulses or Gaps in Mice. Front Cell Neurosci. 2017;11:19.

69.	Suzuki J, Corfas G, Liberman MC. Round-window delivery of neurotrophin 3 regenerates cochlear synapses after acoustic overexposure. Sci Rep. 2016;6:24907.

70.	Hashimoto K, Hickman TT, Suzuki J, Ji L, Kohrman DC, Corfas G, et al. Protection from noise-induced cochlear synaptopathy by virally mediated overexpression of NT3. Sci Rep. 2019;9(1):15362.

71.	Cassinotti LR, Ji L, Borges BC, Cass ND, Desai AS, Kohrman DC, et al. Cochlear Neurotrophin-3 overexpression at mid-life prevents age-related inner hair cell synaptopathy and slows age-related hearing loss. Aging Cell. 2022;21(10):e13708.

72.	Stankovic KM, Corfas G. Real-time quantitative RT-PCR for low-abundance transcripts in the inner ear: analysis of neurotrophic factor expression. Hearing Research. 2003;185(1):97-108.

73.	Fernandez KA, Jeffers PWC, Lall K, Liberman MC, Kujawa SG. Aging after noise exposure: acceleration of cochlear synaptopathy in "recovered" ears. J Neurosci. 2015;35(19):7509-20.

A Control   Ntf3-KD

Myo7a CtBP2 GluA2

B
CtBP2 Puncta (% re Control Max)
○ Control
○ Nft3-KD
Cochlear Region (kHz)
*

C
GluA2 Puncta (% re Control Max)
Cochlear Region (kHz)
**

D
Synapses (% re Control Max)
Cochlear Region (kHz)
**

E Control   Ntf3-OE
10 μm

F
CtBP2 Puncta (% re Control Max)
□ Control
□ Ntf3-OE
Cochlear Region (kHz)
**

G
GluA2 Puncta (% re Control Max)
Cochlear Region (kHz)
***

H
Synapses (% re Control Max)
Cochlear Region (kHz)
***

A

Peak I

Control
Ntf3-KD

****

Peak II

ns

Peak III

ns

Peak IV

ns

Peak V

ns

B

Control
Ntf3-OE

****

****

***

***

ns

Intensity (dB SPL)