1 **Theory and application of an improved species richness estimator**
2

E. W. Tekwa[1,2*], Matthew A. Whalen[1,2,3], Patrick T. Martone[3], Mary I. O'Connor[1]

[1]Department of Zoology, University of British Columbia, Vancouver, BC, Canada.

[2]Hakai Institute, Heriot Bay, BC, Canada

[3]Department of Botany, University of British Columbia, Vancouver, BC, Canada.

*Corresponding author email: ewtekwa@gmail.com, ORCID: 0000-0003-2971-6128

Emails for MW: mawhal@gmail.com, PM: patrick.martone@botany.ubc.ca, MO:

oconnor@zoology.ubc.ca.

Keywords: species richness, biodiversity, bias, occupancy, clustering, bootstrapping

Statement of authorship: ET and MO designed the study, ET performed the derivations and

simulations, MW and PM collected the data, ET performed the analyses, ET and MW wrote the

code, ET wrote the first draft of the manuscript, and all authors contributed to editing.

Data accessibility statement: All code and data are provided on Github at

https://github.com/EWTekwa/RichnessEstimator and will be assigned a DOI through Zenodo

upon journal acceptance.

Abstract

Species richness is an essential biodiversity variable indicative of ecosystem states and rates of invasion, speciation, and extinction both contemporarily and in fossil records. However, limited sampling effort and spatial aggregation of organisms mean that biodiversity surveys rarely observe every species in the survey area, which introduces bias to the estimated richness and inaccuracy to comparisons of communities across space and time. Here we present a nonparametric, asymptotic, and bias-corrected richness estimator, Ω_T, by modelling how spatial abundance characteristics affect observation of species richness. We conduct simulation tests and applied Ω_T to a tree census and a seaweed survey. Ω_T consistently outperforms common estimators in balancing bias, precision, and difference detection accuracy. Our results provide theoretical insights into how natural and observer-induced variation affects species observation and support Ω_T as a promising and application-ready richness estimator for a wide variety of data.

Introduction

A central objective of biodiversity science is to understand how species richness changes across space and time [1]. Not only is species richness important for today's ecosystem and human health assessments [2–4], it is essential to inferring paleoecological patterns and mass extinction events [5]. Species richness estimation techniques are among the most highly cited works in ecology [6–8]. Further, estimating missing species or classes from limited observations is a fundamental problem in other fields, including genomics [9], cryptography, machine learning, and linguistics [10].

While it is clear that global extinction rate exceeds baseline levels [11,12], synthesis of

species richness timeseries have suggested both gain and loss of local species richness at local

scales [13–16]. Observed species richness and detection of differences across communities or

time are prone to data bias [17] and undocumented errors resulting from varying survey and

sampling effort, design, and observer skill [18]. These problems are more acute for species

richness than for other biodiversity metrics [19], in part because species richness is sensitive to

detection of rare species. Data-intensive methods such as multi-species occupancy models can

estimate richness of targeted species groups, but they require repeated field surveys and are

sensitive to models of abiotic and biotic drivers [20,21]. For many general applications and meta-

analyses, it is desirable to have an asymptotic (finite and effort-independent) richness estimator

that can both correct bias (uncover true absolute richness) and accurately detect spatial and

temporal differences across communities.

In most places, the best biodiversity data we can hope for are individual counts identified

to the species level. One obvious way to improve biodiversity estimation is to increase sampling

effort, but this is costly and not possible for some taxa, habitats, and historical samples.

Insufficient sampling effort can reduce diversity estimation precision and undermine efforts to

detect change over time, but a robust, unbiased estimator can recover missing observations to

some extent. The most robust asymptotic richness estimator that uses abundance data remains the

Chao1 method [6,22]. Chao1 and related estimators are based on Turing's sampling theory that

relates the number of unobserved classes (codes or species) to observed rarity [22,23]. These

methods are expected to be inaccurate with spatial heterogeneity, low sampling effort, and

heterogeneous observation probabilities. The latter issue has been tackled using models requiring

tuning, but these methods can be unstable and still under development [24]. Further, bias to both

estimates and confidence bounds remain intractable problems [25]. There is a need to develop

new statistical methods that fully utilize information in common survey data, without having to

separately estimate additional quantities that may be less reliable than raw richness itself [20].

Here we outline five technical problems that a new asymptotic estimator should account

for. The first two are human-induced obstacles to species observation: low fraction of individuals

observed and low number of patches (sites, transects, or quadrats) sampled. The last three are

natural properties of biological spatial distributions that also reduce the number of species

observed: low abundance, patch occupancy, and spatial clustering within patch (Figure 1). These

variables will be precisely defined, and we use them to mechanistically derive observation

probabilities and a set of what we will call Ω bias-corrected asymptotic richness estimators. This

approach departs from both Turing's reliance on rarity, which is prone to noisy observations, and

from earlier parametric extrapolations that have weak justifications [22]. Using the derivations,

we highlight how distributions of the variables affect richness observations. Next, we evaluate

the performance of two proposed Ω-estimators relative to five other best-established asymptotic

estimators (Chao1, Chao2, ACE, Jackknife1-abundance, Jackknife2-incidene) using simulations

designed to challenge each estimator through the five problems introduced above. Six statistics

measure bias (under- or over-estimation tendency), precision (repeatability of estimates), and

accuracy (error magnitude and spatial/temporal trend identification). We additionally devise an

improved bootstrapping procedure for confidence bounds that account for the spatial- and

species-dependence between individuals. To test our method with empirical applications, we

used two multi-year, multi-patch datasets: the Barro Colorado Island tree census [26] and the

British Columbia intertidal seaweed survey [27], where the probable true temporal trends are,

respectively, decreasing and not changing. We conduct spatial subsampling and local

downsampling experiments on the real datasets to explore how much different estimators can

recover absolute richness and detect temporal trends. Finally, we create Matlab and R functions

for estimating richness via a suite of asymptotic methods.

Methods

Richness Estimator Derivation

Here we define variables that can be measured from a spatially partitioned abundance

dataset, with the aim of arriving at a bias-corrected richness estimator. Our strategy involves the

following steps (Figure 1B). We begin at the finest spatial resolution sampled ("patches") where

individuals are counted, across which we track how both spatial heterogeneities and observation

errors (through spatial abundance statistics) affect the probability that a species is observed at

least once at the larger community level across patches. We then use spatial abundance statistics

of observed species to estimate true richness in the underlying community, which is the observed

richness divided by some expected observation probability.

In a community defined by patches (indexed x) belonging to the spatial set M, each

species i has a true mean density n_i, which is the total number of individuals at scale M divided

by the number of patches X, or equivalently the individuals per patch in k_i randomly sampled

patches. The fraction of individuals observed at x, m_{ix}, is also understood as the individual

catchability of species i [28]. Thus, $m_{ix}n_{ix}$ is the local "observable abundance", which will serve

as the first (non-spatial) random state variable in our subsequently developed richness estimator.

In contrast to the above community scale, let us descend to the local patch scale where

observation occurs. At each local patch ("observation neighbourhood") where an individual of

species i occurs, there tends to be $n_{i:i}$ individuals (counting the focal individual itself) on average

because of spatial variance [29]. Note "on average" here means per individual, not per occupied

patch, the latter being equal to or smaller than the former because some patches have more than

one individual. The two subscript i's indicate that $n_{i:i}$ is the local density of species i from the

perspective of an individual of species i, which contrasts from the global density n_i. The quantity

$n_{i:i}$ can be understood as the Lagrangian description of spatial distribution (i.e. tracking other

individuals from a focal individual's perspective), which can be precisely translated to

abundance at patch x: n_{ix}, the Eulerian description (i.e. tracking individuals from a fixed spatial

perspective) in fluid dynamic terminologies that gave rise to modern spatial ecology [30]. The

spatial variance $var_x(n_{ix})$ is the variance in the true number of individuals of species i across all

patches x. It can be shown that the mean local density $n_{i:i}$ is a function of spatial variance, with

variance having a normalization factor k_i, the number of patches sampled, rather than the usual

k_i-1: $var_x(n_{ix}) = \frac{1}{k_i}\sum_x^{k_i}|n_{ix} - n_i|^2$ [31,32].

Equation 1. $n_{i:i} = n_i + var_x(n_{ix})/n_i$

This local density is the true expected local abundance that the human observer samples

from in an occupied patch, because in such a patch the observer takes the same perspective as an

individual of species i. We define the first spatial heterogeneity quantity that will serve as our

second random state variable – the within-species clustering across space C_i – as [29,32]:

Equation 2. $C_i = \frac{n_{i:i}}{n_i} = 1 + \frac{var_x(n_{ix})}{n_i^2}$

Note the following identity allows us to simplify Equation 2 in terms of the observable

abundance $m_{ix}n_{ix}$ across patches x instead of the generally unobserved true abundance n_{ix}, if the

fraction of individuals observed does not differ across patches ($m_{ix=}m_i$) – an assumption we will

carry forward:

Equation 3. $\dfrac{\text{var}_x(n_{ix})}{n_i^2} = \dfrac{\text{var}_x(m_i n_{ix})}{(m_i n_i)^2}$

The expected local observable abundance in an occupied patch is thus $m_i n_{i:i} = C_i m_i n_i$.

Using this mean the local observation process can be described by a non-spatial Poisson

sampling process (sampling with replacement). This local Poisson sampling process should not

be confused with a Poisson spatial distribution across patches, which was not assumed. Let the

local observation probability of species i at an occupied patch (the spatial set X_{io}) be $D_{ix \in X_{io}}$.

This probability is one minus the rate of failure to observe:

Equation 4. $D_{ix \in X_o} = 1 - exp(-C_i m_i n_i)$

Now we zoom back out to the community level M and consider spatial processes across

multiple patches that may be occupied or not. The second spatial heterogeneity quantity – our

third state variable – relevant to this level is occupancy P_i, which is the number of occupied

patches k_{oi} by species i divided by k_i:

Equation 5. $P_i = k_{oi}/k_i$

Occupancy is related to clustering through spatial variance, but this relationship is

mediated by true mean abundance [33], which we generally cannot observe from biodiversity

surveys because of imperfect observations. Thus, we treat occupancy and clustering as separate

dynamic variables but, as we will see, our statistical method accounts for the effect of their

covariance. The observation probability of species i per random patch x is the occupancy

(Equation 5) times the conditional local observation probability (Equation 4) when the patch is

occupied.

Equation 6. $D_{ix} = P_i D_{ix \in X_{io}}$

The observation probability of species i within a community M is one minus the

cumulative probability of failure to observe, the latter being the probability of failure to observe

within patch raised to the power k_i (our fourth state variable) if patches are randomly picked. The

observation probability at scale M (D_{iM}) is:

Equation 7. $D_{iM} = 1 - (1 - D_{ix})^{k_i}$

If patches are not randomly picked, then k_i should be the effective sample size. As an

extreme example where only one patch is repeatedly sampled at the same time in exactly the

same way, effectively $k_i=1$. In practice however, even sampling the same patch repeatedly would

generate different individual and species counts when catchability m is below 1 and/or

individuals move between patches during the finite time separating samples. Ignoring the non-

independence of spatial samples would inflate D_{iM}, but defining what is meant by effective

sample size will require nuanced considerations of how the spatial sampling design relates to the

community's spatial distribution. We will assume k_i samples are independent and leave effective

sample size as a venue for future improvements.

The raw observed species richness S can be written without additional assumptions as the

true richness Ω multiplied by the expectation of D_{iM} across truly present species i as a function of

$\boldsymbol{\Phi}$, which includes R state variables ϕ_r. In our observation process model, $R=4$ and ϕ_1 to ϕ_4 are

$m_i n_i$, C_i, P_i, and k_i.

Equation 8. $S = \Omega E_i[D_{iM}(\boldsymbol{\Phi})]$

We can stop here and define Ω_o (Equation 9) as our first proposed bias-corrected richness

estimator using the average observation probabilities based on state variables $\widehat{\boldsymbol{\Phi}}$ measured from

the set S_o of observed (rather than truly present) species (see Figure 1 for numerical example).

All quantities with a $\widehat{hat}$ denote observed rather than true values in the subset S_o. The estimator

Ω_o contains survivorship bias [34] – a type of sampling bias stemming from the difference

between $\widehat{\boldsymbol{\Phi}}$ and $\boldsymbol{\Phi}$, because $\widehat{\boldsymbol{\Phi}}$ is only measurable for observed species. For intuition, observed

species will likely have higher abundances than the average of truly present species because rare

species tend to be missed.

Equation 9. $\Omega_o = S/E_{i \in S_o}[D_{iM}(\widehat{\boldsymbol{\Phi}})]$

An alternative is to write $E_i[D_{iM}(\boldsymbol{\Phi})]$ in Equation 8 in terms of a second-order Taylor

expansion (with r and q being variable indices), which provides insights into how means and

variances in different statistics affect the observation process:

Equation 10. $S \approx \Omega \left(D_{iM}(\bar{\phi}) + \sum_{r=1}^{R} \left(\frac{d^2 D_{iM}(\bar{\phi})}{2d\phi_r^2} \mathrm{var}_i(\phi_r) + \sum_{q>r}^{R} \frac{d^2 D_{iM}(\bar{\phi})}{d\phi_q \phi_r} \mathrm{cov}_i(\phi_q, \phi_r) \right) \right)$

$\bar{\phi}$ stands for the true average values of the R state variables across truly present species. If

there are observation errors, higher moments are increasingly harder to estimate and may

destabilize the estimator. We thus propose an idealized estimate for Ω, the second-order Taylor

approximated richness Ω_{TC}, with TC indicating that the spatial abundance statistics are the

"corrected" or unbiased statistics for all truly present species and not just observed species:

Equation 11. $\Omega_{TC} =$

$S \bigg/ \begin{array}{l} D_{iM \in S_o}(\bar{\phi}) + \dfrac{d^2 D_{iM}(\bar{\phi})}{2d(mn)^2} \mathrm{var}_i(m_i n_i) + \dfrac{d^2 D_{iM}(\bar{\phi})}{2dC^2} \mathrm{var}_i(C_i) + \dfrac{d^2 D_{iM}(\bar{\phi})}{2dP^2} \mathrm{var}_i(P_i) \\[3mm] + \dfrac{d^2 D_{iM}(\bar{\phi})}{d(mn)dC} \mathrm{cov}_i(m_i n_i, C_i) + \dfrac{d^2 D_{iM}(\bar{\phi})}{d(mn)dP} \mathrm{cov}_i(m_i n_i, P_i) + \dfrac{d^2 D_{iM}(\bar{\phi})}{dCdP} \mathrm{cov}_i(C_i, P_i) \end{array}$

Ω_{TC} contains a different set of bias than Ω_o. The potential bias for Ω_{TC} comes from the

Taylor moment truncation beyond the second order terms but contains no survivorship bias (the

spatial statistics are from $\boldsymbol{\Phi}$). However, Ω_{TC} cannot be implemented in real data because we do

not know $\boldsymbol{\Phi}$ containing $\bar{\phi}$ and ϕ_r across all truly present species. We thus propose a Taylor

approximated richness Ω_T that uses directly measured variables $\widehat{\boldsymbol{\Phi}}$ in place of $\boldsymbol{\Phi}$ in Equation 11,

which is operational on real biodiversity survey datasets but contains survivorship bias. This

estimator is:

Equation 12. $\Omega_T =$

$$S \Bigg/ \begin{array}{l} D_{iM \in S_o}(\hat{\phi}) + \frac{d^2 D_{iM}(\hat{\phi})}{2d(mn)^2}\operatorname{var}_{i \in S_o}(\widehat{m_i n_i}) + \frac{d^2 D_{iM}(\hat{\phi})}{2dC^2}\operatorname{var}_{i \in S_o}(\widehat{C_i}) + \frac{d^2 D_{iM}(\hat{\phi})}{2dP^2}\operatorname{var}_{i \in S_o}(\widehat{P_i}) \\ + \frac{d^2 D_{iM}(\hat{\phi})}{d(mn)dC}\operatorname{cov}_{i \in S_o}(\widehat{m_i n_i}, \widehat{C_i}) + \frac{d^2 D_{iM}(\hat{\phi})}{d(mn)dP}\operatorname{cov}_{i \in S_o}(\widehat{m_i n_i}, \widehat{P_i}) + \frac{d^2 D_{iM}(\hat{\phi})}{dCdP}\operatorname{cov}_{i \in S_o}(\widehat{C_i}, \widehat{P_i}) \end{array}$$

$\hat{\phi}$ stands for the observed across-species averaged value set $(\widehat{mn}, \hat{C}, \hat{P}, k)$, and $\widehat{m_i n_i}$, $\widehat{C_i}$,

and $\widehat{P_i}$ are the observed values for each species i. We assumed that k is always known and

identical for all species in a survey, hence there is not a second-order correction term associated

with variance in k. We define the seven terms in the denominator of Equation 12 as "correction

factors": 0^{th}-order, V_{mn}, V_C, V_P, $C_{mn,C}$, $C_{mn,P}$, and $C_{C,P}$. Ω_T looks complicated but is

computationally trivial and effectively takes no more time or data than Ω_o or other abundance-

based estimators. In summary, Ω_T (Equation 12) contains both survivorship bias and truncation

bias; Ω_{TC} contains only truncation bias (Equation 11) but this is because it is idealized (it can

only be obtained from simulation data); Ω_o (Equation 9) contains only survivorship bias. In

simulations, Ω_{TC} serves to illustrate the upper performance limit of Ω_T if the survivorship bias

$\hat{\boldsymbol{\Phi}} - \boldsymbol{\Phi}$ can be corrected.

The data and code used in this paper are available on a Github/Zenodo repository at

https://github.com/EWTekwa/RichnessEstimator. The Matlab/R function, *RichnessEstsCov*,

takes in a spatial abundance matrix and returns all estimators.

Confidence Bounds

We produced 95% confidence bounds of each estimator from 2000 the full empirical datasets

and 50 for each of 40 subsampled replicates within each scenario. We use non-parametric

bootstrapping [35] because the richness distribution is unknown. Individual occurrences are not

independent both within species and patches because of the intraspecific tendency to cluster

[31,36]. In contrast, previous bootstrapping procedures for richness estimates assumed that

sampled individuals are independent (as did the estimators themselves) [6], an assumption that is

also used for rarefaction and extrapolation techniques that control for sampling effort when

comparing communities [18]. To account for spatial and species dependencies of sampled

individuals, for each bootstrap we first randomly sampled with replacement k transects (Figure

2). Then from this spatially randomized set we randomly sampled with replacement as many

species as the point estimator predicts from the original dataset (eg. Ω_T or Chao1). All species are

drawn from the observed set of S species but the bootstrap community now consists of a

potentially larger number of pseudospecies. We used this bootstrap sample size instead of the

raw S, because we know S underestimates the true community richness and bootstrapping is

intended to recreate sampling variations from that community, not from the observed subset. We

further used a mean bias-corrected percentile method to obtain the final confidence bounds. This

involves first centering the bootstrap estimates so that the mean of the corrected bootstrap

estimates matches the point estimate. Then the 2.5th and 97.5th percentiles of the corrected

bootstrapped estimates serve as the 95% confidence bounds. For subsample schemes, the

confidence bounds were plotted as means of the 2.5th and 97.5th quantiles across replicates,

which showed average confidence bounds (not uncertainties in the richness point estimates

across replicates). The bootstrapping function in Matlab/R is *BootRichnessEsts_all*.

Simulation Setup

To test the performance of asymptotic richness estimators, we simulated spatially

partitioned communities under four scenarios that present different challenges that arise from

imperfect local observation and spatial heterogeneities. For each simulation scenario, we created

2000 unique communities, each with 100 patches and 1 to 100 species. Communities in a

scenario have different characteristics but converge on scenario mean values in abundance E[n],

clustering E[C], and occupancy E[P], which simulate natural variations in spatial biodiversity

patterns that we may want to discern. Further, each community was sampled at different efforts,

converging on mean spatial sampling effort (patches sampled E[k]) and fraction of individuals

observed E[m]. Thus, each of the 2000 communities represent a unique set of biological and

sampling conditions.

Specifically, each simulated community's mean fraction of individuals observed $\tilde{m}$ was

~$1/(1+(\text{E}[1/m]\text{-}1)U(0,1))$ where U is a uniform random value. Each species in a community in

turn had a $1/m_i$ value drawn from a normal distribution centered at the community mean $\tilde{m}$ with

a variance of 10 and a minimum of 1 (perfect observation). Each community's mean abundance

$\tilde{n}$ was ~E[n]$U(0,1)$ and species n_i was drawn from a lognormal distribution with a standard

deviation of 0.5. Each community's mean clustering $\tilde{C}$ was ~E[C]$U(0,1)$, and species C_i was

drawn from a lognormal distribution with a standard deviation of 1 and a minimum of 1 (no

clustering). Each species' n_i and C_i implied a P_i value according to the identity provided in [33].

We assigned individuals to each patch by drawing from a Poisson distribution with the mean

being the species' local density (Equation 1) multiplied by 1 if ~$U(0,1)<P_i$ and 0 otherwise. This

led to true species abundance distributions that satisfied the scenario means E[n], E[C], and E[P].

Finally, subsampling biodiversity surveys were simulated by randomly picking $k\in[2,10]$ patches

in each community. Within each patch and for each species, individuals were observed by

drawing from a Poisson distribution with the mean being the species' local abundance times m.

This implied sampling individuals with replacement within patches, such that locally observed

abundance may be higher than what exists as m approaches 1, but even in this case the average

observed individuals will converge to n. Sampling with replacement is likely in non-destructive

field surveys when individuals move and their identities are not tracked; on the other hand the

scheme approximates sampling without replacement when m is small. Our main results are from

two simulation scenarios with E[n]=2: imperfect local observation (E[m]<1) (Figure 3A) and

spatial heterogeneity (E[C]>1 and E[P]<1) (Figure 3B), each representing unique challenges to

richness estimates. The n distributions for these scenarios are shown in Figure S1A-B. We also

tested the estimators in scenarios with mixtures of imperfect local observation and spatial

heterogeneity (Figure S2).

Performance Evaluation

The evaluation of estimator' performances in the simulations was conducted with six

statistics that cover different aspects of bias, precision, and detection accuracy. A simple least-

squares linear regression was performed on the estimated richness vs. true richness within each

estimator. A slope less than 1 indicates that more diverse communities are more severely

underestimated than less diverse communities, which is a prevalent bias among richness

estimators (sometimes expressed as 1-slope [37]). In contrast to slope deviating from 1 as bias,

precision is how close are multiple estimates given the same true richness; this can be measured

by a low standard deviation (S.D.) in the regression slope. In addition, R^{2*} was computed for the

1:1 line, which describes the variance in estimated richness explained by true richness. A small

R^{2*} indicates either underestimates or overestimates, with large errors capable of generating

negative values. R^{2*} is an overall measure of the estimator's accuracy. R^{2*} contrasts from the R^2

of the true:estimate linear regression, the latter having been used to measure precision [37] but

we believe the measure is confounded with bias. Even if an estimator always returns the same

richness value for a given true richness, bias would reduce R^2 below 1, thus we chose to use

slope and S.D. in slope as our measures of bias and precision. Third, richness difference

detections were checked between 2000 randomly picked community pairs that were different in

richness by at most 2, 10, or 20 but were not identical. These tests provide more nuanced

measures of detection accuracy relevant to uncovering spatial or temporal trends, which are

major objectives of biodiversity studies. Detection accuracies above 0.5 is better than a coin-flip.

Low bias and high precision should improve accuracy, but there are many possible

interpretations of accuracy. For instance, the detection accuracies we defined (richness difference

detections in the paragraph above) refer to point estimates from the full dataset, whereas in

practice we may look at whether spatial or temporal trend in richness change has a low average p

value across bootstraps (when incorporating uncertainties). In this sense trend detection would be

affected by an estimator's precision more than pairwise comparison accuracy based on point

estimates. We desire asymptotic richness estimators to both reduce bias and increase accuracy,

but these components may exhibit tradeoffs given limited information [37]. Thus, the

performance of an estimator cannot be summarized by a single measure; we therefore present all

six measures introduced above to assess whether certain estimators rank higher in one aspect and

lower in others.

Simulations from the worst data conditions led Ω_{TC} correction factors to estimate

negative observation probabilities in up to 3% of the communities (Figure S2). These cases

indicate conditions where variances and higher moments were unstable – even though Ω_{TC} had

access to the true values of mn, C, and P, the data was still poor because m and P were small.

Based on where outliers occurred, we set a threshold of 10% estimated observation probability

below which Ω_T and Ω_{TC} switched to using only the 0^{th}-order correction factor (first correction

term in denominator of Equation 12), which cannot be negative and dominated among correction

factors. This threshold did not affect Ω_T in the simulations but serves as a safeguard against

unforeseen scenarios.

Empirical Datasets

The first dataset that we will use to test estimators is the Barro Colorado Island tree

census from Panama [26]. Start in 1982, every stem was identified by species every 5 years, for a

total of 8 census. Over 300 species was observed across 1250 quadrats that form a continuous

grid over 50 hectares. A negative temporal richness trend seems likely from the raw data. The

census nature of the dataset allows us to conduct spatial subsampling and local downsampling

experiments to test whether an estimator can recover the known true richness and whether the

negative trend can be detected under poor sampling conditions.

The second dataset is an extensive seaweed diversity survey collected annually from

2012 to 2019 on the coast of British Columbia, Canada, and provide a protocol for obtaining

bootstrapped confidence bounds. The dataset features relatively high standards in sampling effort

and taxonomic identification for the marine realm [27]. Still, we expect uncertainty to persist

because of imperfect sampling and spatial heterogeneity. We also conducted post-hoc

subsampling experiments to explore how various scenarios of reduced sampling effort affect the

estimators.

The survey tracked seaweed species and their abundance (estimated as percent cover)

across steep gradients in abiotic stress associated with rapid changes in elevation and immersion

time. The researchers established nine transects running parallel to shore at three intertidal

elevations. Each year, we randomly selected ten quadrats on each transect and surveyed them for

percent cover of all seaweed species. Between 73 and 82 species were observed each year, and

no temporal trend can be detected from the raw data. Transects likely represent independent

spatial samples of an underlying community, whereas quadrats were less spatially independent,

so we aggregated quadrat data within transects for this analysis. A 10 x 10 grid was overlaid on

the 0.5m x 0.5m quadrat, and any organism filling each square of this grid added 1 percent to the

species' total cover within the quadrat [38]. A species that was partially present in only one

square of the grid was assigned a % cover of 0.5 (or trace). We therefore divided all percent

cover data by 0.5 to obtain abundance.

Abundance within the context of sampling has no real biological meaning beyond being

discrete; the abundance unit does not have to coincide with the individual organism, a concept

that we note is open to biological debates and subject to evolution [39,40] but has no

consequences for richness estimation. The only important feature of count data is that the

observable abundance mn, not the biological n, approximates a Poisson process; clustering C and

occupancy P are not affected by a change of unit and are thus insensitive to the individual

definition (Equation 3 & Equation 5). The other consideration for converting percent cover to

observed local abundance (Cmn) is that it is capped and thus not quite Poisson, but in practice

recording abundances beyond this cap will likely not affect the estimated observation probability

(eg. failure to observe, exp($-Cmn$), is 4×10^{-44} for $Cmn =100$ in Equation 4).

In addition to obtaining point estimates of richness on both full datasets, we subsampled

transects and quadrats in different combinations and produced replicated subsample estimates

(plus bootstrap confidence bounds). For each subsampling replicate for the Barro Colorado

Island tree census, we randomly pick a subset of quadrats (patches) without replacement and

sampled these same quadrats every year. For each subsampling replicate for the British

Columbia seaweed survey, we randomly pick transects and quadrats (without replacement) and

sampled these same quadrats every year. Quadrat counts are summed to obtain transect data, so

transects were treated as patches here. Each spatial effort level also receives a complement local

downsampling scheme in which only 10% of individuals in each selected transect was observed.

For each scheme, we replicate the subsampling 40 times since different transects and quadrats

could be randomly picked. An ideal estimator would predict the same true richness regardless of

spatial subsampling and local downsampling.

For each full dataset and subsampling/downsampling scheme, and for each subsampled

replicate and bootstrap, we perform a linear regression of estimated richness on year or census.

The slope of the regression is the detected temporal trend, the S.D. is the standard deviation in

slope from the regression, and the p-value of the slope is a measure of how frequent are more

extreme slopes expected given the null hypothesis of no trend. When the regression is performed

on the point estimates (using the non-bootstrapped full data or subsampled data), the statistics are

related to the detection accuracy that we measured in the simulation (presented in brackets in

Figure 4 and Figure 5). For subsampled schemes, the point estimate regressions are performed

across replicates and averaged to represent the expected statistics when sampling effort is lower

than in reality. When the regression is performed on a bootstrapped dataset, the statistics include

uncertainties in the estimates given the data, so the average p-value across bootstraps should be

higher than for point estimates (presented as the main statistics in brackets in Figure 4 and Figure

5).

For the Barro Coloardo Island tree census, we measure the false negative rate β (type II

error) for detecting a temporal trend using each estimator (defined as $p > 0.05$), assuming a trend

is real. Per standard practice, $\beta \lesssim 0.1$ is considered an acceptable false negative rate; otherwise the

study and estimator are statistically underpowered. For point estimates, β is obtained from the

frequency of regressions across subsampled replicates whose slope $p>0.05$ (presented in brackets

in Figure 4). These statistics do not exist for point estimates on the full dataset (because there is

only one sample). For bootstrapped estimates, β is obtained from the mean frequency of $p>0.05$

across bootstraps (and all replicates under subsampling schemes). These bootstrapped β

statistics are presented as the main statistics in Figure 4.

For the British Columbia seaweed survey, we measured the false positive rate α (type I

error) for detecting a temporal trend using each estimator (defined as $p\leq0.05$), assuming a trend

does not exist. This null assumption is more tenuous than the negative trend assumption for the

Barro Colorado dataset because the seaweed survey, like most marine studies, are not census and

only covers a small fraction of the community habitat, so an estimated $\alpha>0.05$ may either be a

true false positive identification by an estimator or evidence for an actual temporal change. For

point estimates, α is obtained from the frequency of regressions across subsampled replicates

whose slope $p\leq0.05$ (presented in brackets in Figure 5). These statistics do not exist for point

estimates on the full dataset (because there is only one sample). For bootstrapped estimates, α is

obtained from the mean frequency of $p\leq0.05$ across bootstraps (and all replicates under

subsampling schemes). These bootstrapped α statistics are presented as the main statistics in

Figure 5.

Results

Theoretical Insights

Here we describe how sampled patches (k) and means and variances in observable

abundance (mn), clustering (C), and occupancy (P) affect species observation and induce bias to

the observed richness. First, we found that the effects of variable means $\bar{\phi}$ (across species and

patches) were captured by the 0^{th}-order correction factor $D_{iM}(\bar{\phi})$ in the Taylor expansion of the

richness estimator (Equation 11), which is the single-species observation probability (Equation

7) computed at variables' mean values (as if all species in the community share the same

characteristics). Equation 7 states that observation probability increases with means of *k, n, m, C,*

and *P*.

Across species variance $(\text{var}(\phi))$ effects depend on the signs of the second-order partial

derivatives (Equation 11). The analytical expressions are long but always negative (see

numerical outcomes in Figure 3C-D). Therefore, variances always decrease the overall

observation probability.

These results clarify two main ways in which non-richness biodiversity components

induce bias to observed richness. Clustering and occupancy are related to spatial turnover and

dissimilarity, yet their mean values affect species observation (Equation 7). Variance in

observable abundance, which is likely inversely proportional to community evenness (assuming

m is not anti-correlated with *n*), reduces observed richness even though true richness should not

be affected by evenness. These effects illustrate the need for biodiversity metrics to be bias-

corrected if they are to represent their definitions.

Simulation Tests

We tested the proposed estimators (Ω_o: Equation 9 and Ω_T: Equation 11) on a simulated

dataset of species abundance across a heterogeneous landscape. We compared Ω_o and Ω_T to other

leading estimators including: the unbiased Chao1 [6,22], Chao2 [28], the Abundance-based

coverage estimator (ACE) [41], Jackknife1 for abundance data, and Jackknife2 for incidence

data [42]. We also tested the idealized version of Ω_T that is only opertional in simulations where

the true spatial statistics are known for all species to establish theoretical limits of Ω-type

estimators (Ω_{TC}: Equation 11). For incidence-based estimators including Chao2 and Jackknife2,

abundance within patches were converted to presence/absence. The six performance statistics in

Table 1 are described in the Methods: Performance Evaluation section.

The simulations show that our proposed Ω_o and Ω_T estimators were either the best or

close to the best across observation and spatial heterogeneity scenarios in terms of detection

accuracy (across all true difference levels of 2, 10, or 20), while their biases (true:estimated

richness slope) were generally lower than other abundance-based estimators but higher than

incidence-based estimators (Figure 3, Figure S2, Table 1). All estimators eroded precision when

compared to raw, but Ω_o and Ω_T had the highest precision after raw (in term of standard deviation

of slope estimate). Incidence-based estimators performed better under spatial heterogeneity [37]

but worse under imperfect local observations. In particular, Chao2 and Jackknife2 for incidence

data contained the least bias as a function of true richness (in term of regression slope) but

performed poorly in detecting small differences under poor conditions (Table 1 C&D).

Jackknife2 consistently ranked near the top along with Ω_o and Ω_T in term of overall accuracy (1:1

R^{2*}), but Jackknife2 detected actual differences much less accurately. All estimators other than

Chao1, Ω_o, and Ω_T were near-worst performers under some scenarios and measures so are prone

to catastrophic failures when the underlying community characteristics are unknown. Between

the three most robust estimators, Ω_T always out-ranked Chao1 and either out-ranked or equaled

Ω_o across all scenarios and measures. In particular, under the poorest scenario only Ω_o and Ω_T

were better than coin flips in detecting richness differences were ≤ 2 (scenario D in Table 1).

451   Between Ω_o and Ω_T, we found that Ω_T exhibited less bias overall while losing little in

452 precision and detection accuracy, therefore we will focus on Ω_T as the main product of our paper

453 (Equation 12). The proposed Ω_T gained 2-8% in detection accuracy compared to using raw

454 observed richness across scenarios. The idealized Ω_{TC} (only operational in simulations)

455 performed best overall, further improving difference detection accuracy (up to 15% higher than

456 raw) while almost eliminating bias (slope close to one, grey rows in Table 1).

457   The room for improvement to Ω_T comes from survivorship bias in raw measurements of

458 mn, C, and P. This bias can be observed in the differences between the Ω_T and Ω_{TC} correction

459 factors in each simulation scenario (Figure 3C-D). Across both test sets, observed mean and

460 variance in mn were biased upward because rare species were missed and observation noise

461 added to differences (Figure S1C, E). The upward bias in mean mn overestimated observation

462 probability, but the upward bias in mn variance underestimated observation probability (Figure

463 3C-D). For the imperfect local observation scenario, mean observed P was biased downward,

464 and mean observed C and variances in C and P were all biased upward (Figure S1D). The

465 downward bias in mean P and upward biases in variances in C and P led to underestimates in

466 observation probability (Figure 3C), while the upward biases in mean C overestimated

467 observation probability. For the scenario with spatial heterogeneity, the biases in means and

468 variances of C and P were the exact opposite of those caused by low fractions of individuals

469 observed (Figure 3D, Figure S1F). The opposing effects of biases across variables partly

470 cancelled each other, reducing net bias in the Ω_T richness estimate and causing the net correction

471 to approach that of the idealized estimator Ω_{TC} (Figure 3C-D). The covariance correction factors

472 were generally smaller than the other factors and similar in Ω_o and Ω_T. In comparison, other

473 estimators depended only on the observation of rare species and did not benefit from opposing

spatial or variational biases [6]. However, the net bias in our method depended on the relative

magnitudes of these survivorship biases, which explains why unbiased versions of the variables

(inaccessible in real datasets) improved the proposed method (up to Ω_{TC}). Consistent directions

in the survivorship biases associated with different biotic and abiotic sources suggest that

corrections are possible to some extent; we identify these as future research objectives.

Next we examine how well the proposed estimators perform in reducing bias and

detecting differences in real applications where sampling is incomplete.

Barro Colorado Island Tree Census

We first computed the raw observed richness, Chao1, Chao2, and the proposed Ω_T point

estimates for each year using the full census data. For Ω_T, observed spatial statistics used to make

corrections are shown in Figure S3. Based on the raw data, a negative trend of four species lost

per census (per 5 years) was observed (Figure 4A, $p=2\times10^{-5}$ from a least-squares linear

regression). Bootstrapped regressions on the raw data (containing possible variations of the true

community represented by the census plot) continued to support a negative temporal trend (mean

$p=0.002$). We therefore recorded the bootstrapped false negative rate β for all estimators across

subsampling schemes assuming a true negative trend (assuming a significance level $\alpha=0.05$ and

a standard acceptable false negative rate being $\beta<0.1$). Applications to bootstrapped datasets

reveal that Chao1 and Chao2 overcorrected and obscured the temporal trend, as seen by

increased p values and $\beta>0.1$. In contrast, Ω_T determined no correction was necessary given the

data quality and retained the original temporal trend, p values, and β.

Next we examined how the estimators perform when only 12 or 3 quadrats out of 1250

were sampled through the census years. Across 40 replications for each of the spatial

subsampling schemes, all estimators except for the raw data at 12 sampled quadrats indicate no

evidence of temporal trend and high false negative rates when bootstrapped uncertainty was

included (high main p and β values in Figure 4B, C). However, when only the point estimates are

considered (excludes uncertainty in richness estimates), both raw and Ω_T consistently indicated a

decreasing temporal trend at 12 sampled quadrats, and raw continued to do so at 3 sampled

quadrats (low bracketed p and β values in Figure 4B, C). When spatial subsampling was coupled

with local downsampling of individuals observed, all except raw failed to detect a temporal trend

(Figure 4E, F), although Ω_T came close to recovering the trend at 3 sampled quadrats based on

point estimates ($p=0.09$ in Figure 4F). While a generally lower β for Ω_T when compared to

Chao1 and Chao2 across sampling schemes indicates a greater derived statistical power (lower

data requirement to infer the true temporal trend), regressions based on raw data consistently

indicated the lowest β. In this dataset raw data correctly identified the trend even with

subsampling and downsampling, which appears to oppose the simulation conclusion that raw

data is worst at detection. Ω_T's correction at the intermediate spatial sample size was likely

suppressed by the non-independence of quadrat samples, especially without downsampling

(effective k_i in Equation 7 is likely smaller than the number of quadrats sampled). We will revisit

the apparent contradiction in detection accuracy in the Discussion.

Overall, Ω_T exhibited similar or less downward bias compared to other estimators

(including raw) under all subsampling and downsampling, recovering close to the raw absolute

richness from the full dataset when only 12 quadrats and 10% of individuals were sampled

(Figure 4E). Precision decreased in this case, but detection accuracy was no worse than other

estimators except when compared to raw. In all subsampled and downsampled schemes, only

Chao2 (Figure 4D and F) and Ω_T (Figure 4E) had confidence bounds that sometimes covered the

raw richness from the full dataset. These results are consistent with the simulation analyses,

which revealed Chao2 and Ω_T should be less biased than Chao1.

British Columbia Seaweed Survey

In this marine seaweed diversity dataset, the raw observed richness across years indicated

a marginal signal of temporal change but was inconclusive (average p=0.5 across bootstraps or

0.2 for point estimate, α=0.05 fraction of times bootstrapped regression showed $p \leq 0.05$, Figure

5A). For Ω_T, observed spatial statistics used to make corrections are shown in Figure S4. Chao1

and Chao2 indicated no temporal trend, while Ω_T indicated marginal support for increasing

richness based on point estimates (p=0.07, α=0.06 across bootstraps). Given these statistics, the

true temporal trend is likely weak and the null hypothesis of no change prevails. Ω_T indicated

that between 3 to 9 seaweed species remain unobserved in the survey throughout the years, while

Chao1 indicated none or few species unobserved. Chao2 made larger upward corrections that

were uneven across years. When the full spatial sample was locally downsampled, all estimators

indicated no temporal trends (Figure 5D).

Under the spatial subsampling scheme of 9 transects with 2 quadrats each, all estimators

including raw showed no evidence of temporal trends (Figure 5B). Confidence bounds for Chao2

and Ω_T covered the raw richness from the full dataset, but the upper bound for Chao2 was well

beyond the probable truth (>200 in some years). Under the subsampling scheme of 2 transects

with 9 quadrats each, raw and Chao2 were more prone to false positives than Chao1 and Ω_T (α in

Figure 5C). Only Chao2's confidence bounds covered the full dataset's raw richness but again

the upper bound appears too high. With the addition of local downsampling on top of spatial

subsampling, raw data continued to show the highest false positive rates (α in Figure 5E-F).

Overall, in this dataset Chao2 exhibited the least bias but the lowest precision, while Ω_T

was the most precise, less biased than Chao1, and potentially detected a stronger signal of

temporal trend in the full dataset.

Discussion

The proposed Ω richness estimators for species richness achieve a superior balance of

low bias while retaining precision and accuracy in detecting richness differences compared to

abundance-based estimators including Chao1 [6], ACE [41], and Jackknife1 [22]. In particular,

our second-order Taylor-expansion estimator Ω_T (Equation 12) accounts for the mean and

variance effects of patches sampled (sample size), fractions of individuals observed within each

species (catchability), abundance (rarity), clustering (spatial variance), and occupancy, all readily

computed from spatial abundance data and without the need to independently assess survey

effectiveness. Ω_T harnesses information already present in common surveys, thus it is no more

data-hungry than other methods we assessed. The proposed method can be used to study spatial

and temporal trends [17] as well as their relationships with other environmental covariates,

without the confounding effects of these covariates on richness estimates (i.e. richness and other

covariates may spuriously influence each other through endogeneities introduced if richness is

first estimated with the aid of those covariates) [20,21].

There remains a downward bias (increasing underestimation when true richness

increases) to all richness estimators we examined with simulations including Ω_T, with the

exception of the incidence-based Chao2 [28] and Jackknife2 [22], which surprisingly had the

minimal bias when looking across all replicates despite ignoring abundances (they only relied on

spatial occupancy). However, these incidence-based estimators were poor at resolving small

differences between communities under data-poor conditions. In contrast, across the different

scenarios that simulations datasets represent, Ω_T detected trends correctly more often than no

correction and any other estimators under any condition while maintaining the lowest bias

among abundance-based estimators. Nevertheless, Ω_T contains survivorship bias [34] because

observed species tend to have higher abundance, lower clustering, and higher occupancy than

unobserved species. If survivorship bias can be corrected (as represented by Ω_TC in simulations)

– and this may very well be possible with further research because the bias directions are known

– the proposed approach would come very close to eliminating bias while increasing detection

accuracy drastically. Like Chao1, Ω_T is best understood as a lower-bound richness estimator but

for slightly different reasons. Chao1 was explicitly derived as a lower-bound estimator regardless

of moment truncations [6], whereas the downward bias for Ω_T came from survivorship bias. In

addition, non-independence of spatial samples (k) adds downward bias to all Ω estimators by

inflating inferred observation probabilities; a careful treatment of effective sample size in the

context of these estimators would help improve performance further.

Interestingly, the second-order Taylor-expansion estimator Ω_T outperformed the exact

estimator Ω_o derived directly from the inferred observation probabilities across observed species.

In other words, the Taylor series truncation provided a better description of the underlying

community than assuming that the observed species were the best representations of the hidden

community. This is surprising because Ω_o contains only survivorship bias (since the spatial

abundance statistics are only available for observed species), whereas Ω_T contained both

survivorship bias and truncation error (from excluding statistical moments beyond the second

order). While it is well-known that higher moments are individually difficult to estimate and

would thus introduce instability to the estimator, it remains to be mathematically explained why

the sum of all higher moments up to infinite order, which are theoretically contained in Ω_o, put

Ω_o at a slight disadvantage when compared to Ω_T. However, the theme of multiple opposing

biases also operates among the six different correction factors within the second-order Taylor

approximation of species observation probability in Ω_T. The magnitudes and directions of these

biases are sensitive to underlying non-spatial and spatial heterogeneities in the community,

which always partially cancel each other, which helps explain why the sum outperforms existing

estimators.

Empirical applications of the estimators to the Barro Colorado Island tree census and the

British Columbia seaweed survey revealed some expected patterns and other surprising results.

Using spatial subsampling and local downsampling of individuals, richness estimates and

bootstrapped confidence bounds from Chao1, Chao2, and Ω_T were compared to those obtained

using the full datasets. These comparisons showed that Ω_T and Chao2 were less biased than

Chao1 and raw; under different conditions either Ω_T or Chao2 was the least biased (had estimates

from subsampled data that were closest to richness observed in the full dataset). Ω_T was more

precise than Chao1 and Chao2, the latter being very imprecise with poor data. Application to the

Barro Colorado Island data showed that Ω_T was superior to Chao1 and Chao2 in detecting the

presumably true negative temporal trend across census years either with the full data or with

subsampled and downsampled data (lower false negative rate β); in other words Ω_T gains more

statistical power than other corrective estimators and thus requires less data despite the

estimator's apparent complexity. However, the raw data was surprisingly more consistent than

Ω_T in detecting a negative trend under poor data conditions, which contradicts our simulation

findings that raw data should be the worst at detection. But we should keep in mind this is only

one counterexample of possible communities; experiments with different datasets containing

clear temporal or spatial trends will be necessary to answer whether our simulation conclusions

hold on average. The related question of false positive rate associated with estimators was more

clearly answered with the British Columbia seaweed survey, where the true temporal trend is

likely constant across years. There the raw data was most prone to falsely detecting a trend under

subsampling and downsampling. A caveat to the assumption of no true change is that Ω_T was the

only estimator that generated marginal support for a slight richness increase in the full dataset.

Overall, the empirical applications demonstrated some superior qualities of Ω_T that parallel the

simulation results, but our uncertainty about the true states of the communities, as well as the

paucity of different communities examined, leave room for further empirical and simulation

assessments.

We note that Ω_T is not restricted to counting species and individuals – the number of any

taxonomic unit or non-biological class can be estimated instead of species (e.g., order, family,

functional, or morphological richness, alphabets, words, etc.), and any discrete abundance unit

can replace the biological individual counts. The method partly relies on individual units being

sampled within-patch through a Poisson process, which is likely satisfied even by survey

methods that appear to carry continuous values. Beyond this, the method should even handle

spatial incidence data. With local observable abundance $C_i m_i n_i$ fixed to a sufficiently high value

for observed species (set to >10 when preparing the data), the observation probability per

occupied patch (Equation 4) is effectively one, meaning that the only effect on Ω_T is equating the

observation probability per patch D_{ix} with occupancy P_i (Equation 6). This simple extension

illustrates the potential for the current approach to handle a variety of data qualities including

surveys designed for biodiversity assessment, unstructured observational data, and mixed data

meta-analyses. We leave it for future research to quantify and improve the breadth and

performance of Ω-type richness estimators across different data types. One can further examine

whether rarefaction and extrapolation techniques that control for sample effort or coverage

[18,23] can improve difference detection accuracy when it is not a priority to estimate absolute

richness (i.e. to reduce bias). This would not be an easy endeavour however, as the spatially

heterogeneous setting that our estimator explicitly confronts also poses unaddressed problems on

how to properly measure sample coverage.

While we have offered an improved asymptotic richness estimator that already performs

well in balancing bias, precision, and accuracy, we identified many more open questions and

precise research venues that may be large undertakings on their own. These complications reveal

the inherent difficulties in estimating species richness and detecting biodiversity change, but also

point to exciting possibilities that come with an approach that deviates completely from past

methods. Ultimately, any richness estimator will be sensitive to whether the observed species are

representative of the underlying community; in fact, the meaning of the underlying community in

terms of taxonomic and geographic inclusion will be defined by the organisms that we choose to

observe and record in the first place. Thus, even when armed with an ideal richness estimator,

biological and field-methodological considerations are indispensable to improving the accuracy

of estimated richness and interpreting its meaning across different communities.

Acknowledgement

We thank the Hakai Biodiversity Synthesis group for discussions. ET and MW were

funded by the Tula Foundation. ET was additionally funded by a Mitacs Accelerate Internship.

PM was funded by a NSERC Discovery Grant (RGPIN 2019-06240).

References

1. Loreau M *et al.* 2021 Biodiversity as insurance: from concept to measurement and
application. *Biol Rev* **96**, 2333–2354. (doi:10.1111/brv.12756)

2. Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B. 2011 How Many Species Are There
on Earth and in the Ocean? *PLoS Biology* **9**, e1001127. (doi:10.1371/journal.pbio.1001127)

3. Costello MJ, May RM, Stork NE. 2013 Can We Name Earth's Species Before They Go
Extinct? *Science* **339**, 413–416. (doi:10.1126/science.1230318)

4. Gopalakrishnan V *et al.* 2018 Gut microbiome modulates response to anti–PD-1
immunotherapy in melanoma patients. *Science* **359**, 97–103. (doi:10.1126/science.aan4236)

5. Jackson JBC, Johnson KG. 2001 Measuring Past Biodiversity. *Science* **293**, 2401–2404.
(doi:10.1126/science.1063789)

6. Chao A. 1984 Nonparametric Estimation of the Number of Classes in a Population.
Scandinavian Journal of Statistics **11**, 265–270.

7. Magurran AE. 2004 *Measuring biological diversity*. Malden, Ma: Blackwell Pub.

8. Pielou EC. 1966 The Measurement of Diversity in Different Types of Biological Colledions.
Journal of Theoretical Biology **13**, 131–144.

9. Wang J-P. 2010 Estimating species richness by a Poisson-compound gamma model.
Biometrika **97**, 727–740. (doi:10.1093/biomet/asq026)

10. Orlitsky A, Santhanam NP, Zhang J. 2003 Always Good Turing: Asymptotically Optimal
Probability Estimation. *Science* **302**, 427–431. (doi:10.1126/science.1088284)

11. Barnosky AD *et al.* 2011 Has the Earth's sixth mass extinction already arrived? *Nature* **471**,
51–57. (doi:10.1038/nature09678)

12. Kolbert E. 2014 *The sixth extinction: an unnatural history*. First edition. New York: Henry
Holt and Company.

13. Dornelas M, Gotelli NJ, McGill B, Shimadzu H, Moyes F, Sievers C, Magurran AE. 2014
Assemblage Time Series Reveal Biodiversity Change but Not Systematic Loss. *Science* **344**,
296–299. (doi:10.1126/science.1248484)

14. Elahi R, O'Connor MI, Byrnes JEK, Dunic J, Eriksson BK, Hensel MJS, Kearns PJ. 2015
Recent Trends in Local-Scale Marine Biodiversity Reflect Community Structure and Human
Impacts. *Current Biology* **25**, 1938–1943. (doi:10.1016/j.cub.2015.05.030)

15. Chase JM *et al.* 2019 Species richness change across spatial scales. *Oikos* **128**, 1079–1091.
(doi:10.1111/oik.05968)

16. Loreau M, Cardinale BJ, Isbell F, Newbold T, O'Connor MI, de Mazancourt C. 2022 Do not
downplay biodiversity loss. *Nature* **601**, E27–E28. (doi:10.1038/s41586-021-04179-7)

17. Gonzalez A *et al.* 2016 Estimating local biodiversity change: a critique of papers claiming no
net loss of local diversity. *Ecology* **97**, 1949–1960. (doi:10.1890/15-1759.1)

18. Chao A, Jost L. 2012 Coverage-based rarefaction and extrapolation: standardizing samples
by completeness rather than size. *Ecology* **93**, 2533–2547. (doi:10.1890/11-1952.1)

19. Roswell M, Dushoff J, Winfree R. 2021 A conceptual guide to measuring species diversity.
Oikos **130**, 321–338. (doi:10.1111/oik.07202)

20. Welsh AH, Lindenmayer DB, Donnelly CF. 2013 Fitting and Interpreting Occupancy
Models. *PLoS ONE* **8**, e52015. (doi:10.1371/journal.pone.0052015)

21. Devarajan K, Morelli TL, Tenan S. 2020 Multi-species occupancy models: review, roadmap,
and recommendations. *Ecography* **43**, 1612–1624. (doi:10.1111/ecog.04957)

22. Gotelli NJ, Colwell RK. 2011 Estimating species richness. In *Biological Diversity: Frontiers
in Measurement and Assessment*, pp. 39–54. Oxford University Press.

23. Good IJ. 1953 The Population Frequencies of Species and the Estimation of Population
Parameters. *Biometrika* **40**, 237–264. (doi:10.1093/biomet/40.3-4.237)

24. Bunge J, Willis A, Walsh F. 2014 Estimating the Number of Species in Microbial Diversity
Studies. *Annu. Rev. Stat. Appl.* **1**, 427–445. (doi:10.1146/annurev-statistics-022513-115654)

25. Mao CX, Lindsay BG. 2007 Estimating the number of classes. *Ann. Statist.* **35**.
(doi:10.1214/009053606000001280)

26. Condit R, Pérez R, Aguilar S, Lao S, Foster R, Hubbell S. 2019 Complete data from the
Barro Colorado 50-ha plot: 423617 trees, 35 years. , 286722098 bytes. (doi:10.15146/5XCP-
0D46)

27. Whalen M, Starko S, Lindstrom S, Martone P. 2021 Extreme warming restructures habitat
distribution and productivity along local gradients in stress and biodiversity.
(doi:10.22541/au.162192715.58945198/v1)

28. Chao A. 1987 Estimating the Population Size for Capture-Recapture Data with Unequal
Catchability. *Biometrics* **43**, 783. (doi:10.2307/2531532)

29. Bolker B, Pacala SW. 1997 Using Moment Equations to Understand Stochastically Driven
Spatial Pattern Formation in Ecological Systems. *Theoretical Population Biology* **52**, 179–
197. (doi:10.1006/tpbi.1997.1331)

30. Levin SA, Pacala SW. 1997 Theories of Simplification and Scaling of Spatially Distributed
Processes. In *Spatial Ecology*, pp. 271–295. Princeton, NJ: Princeton University Press.

31. Lloyd M. 1967 `Mean Crowding'. *The Journal of Animal Ecology* **36**, 1. (doi:10.2307/3012)

32. Tekwa EW, Nguyen D, Loreau M, Gonzalez A. 2017 Defector clustering is linked to
cooperation in a pathogenic bacterium. *Proceedings of the Royal Society B: Biological*
Sciences **284**, 20172001. (doi:10.1098/rspb.2017.2001)

33. He F, Gaston KJ. 2003 Occupancy, Spatial Variance, and the Abundance of Species. *The*
American Naturalist **162**, 366–375. (doi:10.1086/377190)

34. Brown SJ, Goetzmann W, Ibbotson RG, Ross SA. 1992 Survivorship Bias in Performance
Studies. *Rev. Financ. Stud.* **5**, 553–580. (doi:10.1093/rfs/5.4.553)

35. Efron B, Tibshirani R. 1986 Bootstrap Methods for Standard Errors, Confidence Intervals,
and Other Measures of Statistical Accuracy. *Statistical Science* **1**, 54–75.

36. Taylor LR. 1961 Aggregation, Variance and the Mean. *Nature* **189**, 732–735.
(doi:10.1038/189732a0)

37. Brose U, Martinez ND, Williams RJ. 2003 Estimating species richness: Sensitivity to sample
coverage and insensitivity to spatial patterns. *Ecology* **84**, 2364–2377. (doi:10.1890/02-0558)

38. Dethier M, Graham E, Cohen S, Tear L. 1993 Visual versus random-point percent cover
estimations: 'objective' is not always better. *Mar. Ecol. Prog. Ser.* **96**, 93–100.
(doi:10.3354/meps096093)

39. Heyward AJ, Collins JD. 1985 Fragmentation in Montipora ramosa: the genet and ramet
concept applied to a reef coral. *Coral Reefs* **4**, 35–40. (doi:10.1007/BF00302202)

40. Tekwa EW, Gonzalez A, Loreau M. 2015 Local densities connect spatial ecology to game,
multilevel selection and inclusive fitness theories of cooperation. *Journal of Theoretical*
Biology **380**, 414–425. (doi:10.1016/j.jtbi.2015.06.016)

41. Chao A, Lee S-M. 1992 Estimating the Number of Classes via Sample Coverage. *Journal of*
the American Statistical Association **87**, 210–217. (doi:10.1080/01621459.1992.10475194)

42. Burnham KP, Overton WS. 1979 Robust Estimation of Population Size When Capture
Probabilities Vary Among Animals. *Ecology* **60**, 927–936. (doi:10.2307/1936861)

Tables

Table 1. Simulation results. Performance was measured as the slope of estimated richness when regressed on true richness (downward bias when <1), standard deviation in regression slope (precision), R^{2*} of the 1:1 line (overall accuracy), and pairwise difference detection accuracy when randomly picked communities (out of 2000) were within a richness difference of 2, 10, or 20 (detection accuracy). For each scenario **(A-D)** and statistics, ranks higher than raw are indicated by dark shades (with black indicating top ranks); ranks lower than raw are unshaded. Ω_{TC} is not ranked (shaded grey).

Estimator	A. Imperfect local observation (m=0.2, n=2, C=1, P=1, k=6)			B. Spatial heterogeneity (m=1, n=2, C=5, P=0.4, k=6)		
	Slope±S.D.	1:1 R^{2*}	Detection ($\leq\Delta2,\leq\Delta10,\leq\Delta20$)	Slope±S.D.	1:1 R^{2*}	Detection ($\leq\Delta2,\leq\Delta10,\leq\Delta20$)
Ω_{TC}	0.93 ± 0.26	0.95	0.59, 0.79, 0.88	1.0 ± 0.15	0.98	0.64, 0.84, 0.91
Ω_{o}	0.92 ± 0.43	0.88	0.59, 0.73, 0.80	0.87 ± 0.31	0.89	0.61, 0.76, 0.83
Ω_{T}	0.95 ± 0.46	0.87	0.58, 0.73, 0.79	0.86 ± 0.30	0.89	0.62, 0.75, 0.83
Chao1	0.93 ± 0.43	0.80	0.56, 0.70, 0.78	0.74 ± 0.30	0.76	0.58, 0.71, 0.77
Chao2	1.0 ± 0.58	0.80	0.56, 0.70, 0.78	0.97 ± 0.37	0.88	0.60, 0.76, 0.84
ACE	1.0 ± 0.46	0.79	0.56, 0.70, 0.79	0.74 ± 0.30	0.76	0.58, 0.72, 0.78
Jackknife1	1.0 ± 0.33	0.89	0.54, 0.68, 0.79	0.75 ± 0.31	0.76	0.56, 0.69, 0.76
Jackknife2	1.1 ± 0.47	0.88	0.52, 0.63, 0.72	1.1 ± 0.35	0.88	0.55, 0.72, 0.81
Raw	0.65 ± 0.41	0.64	0.52, 0.65, 0.73	0.73 ± 0.32	0.75	0.57, 0.70, 0.76

Estimator	C. Mixed conditions (m=0.2, n=2, C=5, P=0.4, k=6)			D. Mixed poor conditions (m=0.1, n=2, C=8, P=0.3, k=6)		
	Slope±S.D.	1:1 R^{2*}	Detection ($\leq\Delta2,\leq\Delta10,\leq\Delta20$)	Slope±S.D.	1:1 R^{2*}	Detection ($\leq\Delta2,\leq\Delta10,\leq\Delta20$)
Ω_{TC}	1.2 ± 0.76	0.51	0.55, 0.70, 0.78	1.1±1.1	0.01	0.53, 0.60, 0.69
Ω_{o}	0.75 ± 0.48	0.66	0.54, 0.63, 0.71	0.52 ± 0.51	0.04	0.53, 0.55, 0.64
Ω_{T}	0.78 ± 0.51	0.68	0.54, 0.64, 0.71	0.55 ± 0.55	0.09	0.53, 0.54, 0.64
Chao1	0.67 ± 0.46	0.49	0.54, 0.62, 0.68	0.53 ± 0.54	-0.06	0.50, 0.53, 0.62
Chao2	1.0 ± 0.79	0.51	0.52, 0.64, 0.72	0.97 ± 1.0	-0.08	0.51, 0.54, 0.63
ACE	0.69 ± 0.53	0.51	0.54, 0.62, 0.69	0.60 ± 0.65	0.04	0.51, 0.53, 0.61
Jackknife1	0.74 ± 0.44	0.64	0.51, 0.61, 0.69	0.55 ± 0.46	0.14	0.49, 0.53, 0.61
Jackknife2	1.0 ± 0.54	0.79	0.50, 0.60, 0.70	0.74 ± 0.73	0.39	0.46, 0.46, 0.56
Raw	0.51 ± 0.39	0.22	0.52, 0.58, 0.67	0.33± 0.37	-0.62	0.49, 0.50, 0.61

Figures

- observable mean abundance: $E_x[m_i n_{ix}]$=1,1/3
- occupancy: P_i=2/3,1/3
- clustering: C_i=1+var($m_i n_{ix}$)/$E_x[m_i n_{ix}]$=1.67,1.67

- observation probability per patch:
  D_{ix}=2/3·(1-exp(-1·1.67)),
  1/3·(1-exp(-1/3·1.67))=0.541,0.142
- observation probability across
  k(=3) sampled patches: D_{iM}=1-(1-D_{ix})³=0.903,0.368
- richness estimate Ω_o=2/$E_i[D_{iM}]$=3.2

Figure 1. From spatial statistics to richness estimate. Spatial statistics are used to infer observation probabilities, leading to a bias-corrected richness estimate Ω_o. The statistics are observable mean abundance *mn* (blue), occupancy *P* (orange), clustering *C* (green), and sites sampled *k* (cyan) from the two observed species (black and outlined grey numbers).

Figure 2. Bootstrapping, subsampling, and downsampling procedures. Any richness estimator produces a point estimate (eg. Ω_T, here 5) from an original dataset (here 4 species S labelled by colours and 4 patches q labelled by grey shades). Each bootstrap first resamples with replacement patches from the original dataset, then resamples with replacments species for all resampled patches to create Ω_T pesudospecies. Multiple bootstrap estimates generate confidence bounds. Subsampling on empirial data tests how estimators perform with reduced spatial coverage, while downsapling is used to test how estimators perform with reduced individual coverage.

Figure 3. Simulated richness estimates. 2000 communities were populated with 1 to 100
species over 100 spatial patches, and 2 to 10 patches (k) were sampled for richness estimates,
including raw (black dots), Chao1 (blue circles), and Ω_T (red circles). **A.** Imperfect local
observation scenario, with mean parameter values of $m=0.2$, $n=2$, $C=1$, $P=1$, $k=6$. **B.** Spatial
heterogeneity scenario, with mean parameter values of $m=1$, $n=2$, $C=5$, $P=0.4$, $k=6$. **C, D.**
Corrections (Δ) to cross-species observation probability (with a baseline of one) by the proposed
estimator (Ω_T, red) and the corresponding idealized estimator where spatial abundance statistics
were corrected for survivorship bias (Ω_{TC}, grey), with the sum of correction factors (first column)
broken into the individual factors (Equation 12). Each dot represents an estimate from the 2000
replicates, and ideally the Ω_T corrections would approach the Ω_{TC} corrections.

Figure 4. Richness estimates and bootstrap confidence bounds for the Barro Colorado Island tree census. For each year, raw (black dotted line), Chao1 (blue solid line), Chao2 (cyan dotted line), and Ω_T (red solid line) richness estimates were computed without **(A)** and with quadrat subsampling (40 times, each without replacement, same quadrats through years, **B**, **C**). Additionally we downsampled the portion of individuals observed in sampled patches **(D-F)**. Lines represent point estimates or mean point estimates across subsamples. 2000 total bootstraps were performed (50 for each subsampled dataset each census) for 95% confidence bounds. Shades represent upper (97.5th) and lower (2.5th) percentiles (or their means across subsamples). Displayed Δ are estimated changes in richness per census either from point estimates (bracketed statistics – ignoring observation and estimation errors) or from bootstrapped estimates. β is the observed false negative rate in detecting a temporal trend (frequency of $p>0.05$ across bootstraps and across subsamples) assuming there is a true trend (based on full census data).

Figure 5. Richness estimates and bootstrap confidence bounds for the British Columbia seaweed survey. For each year, raw (black dotted line), Chao1 (blue solid line), Chao2 (cyan dotted line), and Ω_T (red solid line) richness estimates were computed without **(A, D)** and with transect and quadrat subsampling (40 times, each without replacement, **B, C, E, F**). Additionally we downsampled the portion of individuals observed in sampled patches in **(D-F)**. Transects were considered spatially independent samples of the community of species (*s*), and quadrat data were summed within transects. Lines represent point estimates or mean point estimates across subsamples. 2000 total bootstraps were performed (50 for each subsampled dataset each year) for 95% confidence bounds. Shades represent means of upper (97.5th) and lower (2.5th) percentiles across subsamples (or their means across subsamples). Displayed Δ are estimated changes in richness per census either from point estimates (bracketed statistics – ignoring observation and estimation errors) or from bootstrapped estimates. α is the observed false positive rate in detecting a temporal trend (frequency of p≤0.05 across bootstraps and across subsamples) assuming there is no true trend (based on full data).