Hunter-gatherer oral microbiomes are shaped by contact network structure

Federico Musciotto[1,2+], Begoña Dobon[2,3+], Michael Greenacre[4,5], Alex Mira[6,7], Nikhil Chaudhary[8], Gul Deniz Salali[9], Pascale Gerbault[11], Rodolph Schlaepfer[2], Leonora H. Astete[12], Marilyn Ngales[12], Jesus Gomez-Gardenes[13,14], Vito Latora[15,16,17], Federico Battiston[2,19], Jaume Bertranpetit[3#], Lucio Vinicius[2,9#], Andrea Bamberg Migliano[2,9#]

[1]Dipartimento di Fisica e Chimica, Università di Palermo; Palermo, Italy.

[2]Department of Anthropology, University of Zurich; Zurich, Switzerland.

[3]Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra); Barcelona, Spain.

[4]Department of Economics and Business, Universitat Pompeu Fabra & Barcelona Graduate School of Economics; Barcelona, Spain.

[5]Faculty of Biosciences, Fisheries and Economics, University of Tromsø; Norway

[6]Department of Health and Genomics, Center for Advanced Research in Public Health, FISABIO Foundation; Valencia, Spain.

[7]CIBER Center for Epidemiology and Public Health; Madrid, Spain

[8]Department of Archaeology, University of Cambridge; Cambridge, United Kingdom.

[9]Department of Anthropology, University College London; London, United Kingdom.

[11]Department of Genetics and Evolution, University of Geneva, Geneva, Switzerland.

[12]Lyceum of the Philippines University, Intramuros, Manila, Philippines.

[13]GOTHAM Lab, Institute for Biocomputation and Physics of Complex Systems, and Department of Condensed Matter Physics, University of Zaragoza; Zaragoza, Spain.

[14]Center for Computational Social Science (CCSS), Kobe University; Kobe, Japan.

[15]School of Mathematical Sciences, Queen Mary University of London; London, United Kingdom.

[16]Dipartimento di Fisica ed Astronomia, Università di Catania and INFN; Catania, Italy.

[17]Complexity Science Hub Vienna (CSHV); Vienna, Austria.

[18]Department of Network and Data Science, Central European University; Vienna, Austria.

[#]equal contribution

+equal contribution

Corresponding author: andrea.migliano@uzh.ch jaume.bertranpetit@upf.edu

Abstract

Ancestral humans evolved a complex social structure still observed in extant hunter-

gatherers. Here we investigate the effects of extensive sociality and mobility on the oral

microbiome of 138 Agta hunter-gatherers from the Philippines. Comparisons of

microbiome composition showed that the Agta are more similar to Central African Bayaka

hunter-gatherers than to neighboring farmers. We also defined the Agta social microbiome

as a set of 137 oral bacteria (only 7% of 1980 amplicon sequence variants) significantly

influenced by social contact (quantified through wireless sensors of short-range

interactions). We show that interaction networks covering large areas, and their strong

links between close kin, spouses, and even unrelated friends, can significantly predict

bacterial transmission networks across Agta camps. Finally, more central individuals to

social networks are also bacterial supersharers. We conclude that hunter-gatherer social

microbiomes, which are predominantly pathogenic, were shaped by evolutionary tradeoffs

between extensive sociality and disease spread.

Introduction

Hominins have significantly diverged from other African apes regarding social behaviour and structure[1]. Compared to polygynous mating and male philopatric residence patterns typically found in chimpanzees, bonobos and gorillas, archaeological and ethnographic evidence point to a stepwise emergence of features such as pair bonding, multilocal residence, high mobility between residential camps and increased co-residence with unrelated individuals[2,3]. Such traits were the foundations of multilevel social structuring appearing in ancestral *Homo sapiens* and possibly earlier hominins. The niche of extant hunter-gatherers may offer a window into past human adaptations as it still exhibits features prevalent before the advent of agriculture, such as a high-quality diet including meat and tubers, and multilevel sociality. Multilevel organization results in interconnected social networks covering large areas and multiple residential camps[4], and in frequent interactions between individuals differing by sex, age and relatedness level. Interconnected networks may have accelerated the evolution of cultural innovations in humans compared to other apes[5,6]. However, efficient networks may also facilitate the spread of infectious diseases[7], potentially affecting the structure and composition of hunter-gatherer microbiomes. Previous studies have investigated the role of diet, ecology and environment in hunter-gatherer oral, gut and milk microbiomes[8–16] and revealed higher oral microbiome diversity in hunter-gatherers than in farming populations[17]. However, they have not been able to isolate the contribution of high sociality and individual mobility to microbial transmission from other factors such as shared environments or diet. Although the more fluid and complex sociality of hunter-gatherers results in high levels of camp coresidence[2], cooperation and social interactions among unrelated individuals[18], its potential effects on microbiome transmission have been mostly neglected. We conducted a comprehensive investigation of the oral microbiome of

Agta hunter-gatherers to analyse the specific effect of sociality and social network structure on

the composition of the Agta oral microbiome, with a companion article examining the separate

role of environmental (diet) and biological (age, sex, host genotype) factors[19].

We obtained both oral microbiome sequences and high-resolution social network data

from the same 138 Agta hunter-gatherers from the Philippines. We also collected oral

microbiome data for 21 Bayaka hunter-gatherers from the Congo, and 14 Palanan farmers

neighboring the Agta territory. We sequenced the 16S rRNA region and identified 6409

amplicon sequence variants (ASVs)[20], later reduced to 1980 ASVs (with at least 10 counts and

present in at least two individuals), to detect fine-scale variation between individuals. We also

collected data on proximity interactions and social networks using radio sensor technology

recording close-range dyadic interactions every two minutes for 5-7 days[6,18] from four Agta

camps, and from two longer multi-camp experiments (interactions recorded every hour for one

month). Proximity data were supplemented with information on household composition, kinship

and affinal relationships from all Agta individuals.

Our extensive dataset on oral microbiome composition and social interactions from the

same individuals allowed us to investigate in more depth the possible effects of sociality on oral

microbiome transmission and composition in Agta hunter-gatherers. Our aims were to

investigate the roles of hunter-gatherer niche and geography on oral microbiome diversity in

hunter-gatherers from two continents and a neighboring farming population from the Philippines;

to determine which fraction of the Agta oral microbiome specifically responds to levels of social

interaction; to identify levels of pathogenicity of the oral microbiome transmitted through social

contact; to investigate any potential tradeoffs between increased sociality and the spread of

infectious disease; and to verify potential tradeoffs at individual level by testing whether 'hyper

social' individuals also shared more bacteria. In the following, we provide evidence that the oral

microbiome of extant hunter-gatherers was partially shaped by tradeoffs between extensive

sociality and the spread of infectious disease.

Results

Hunter gatherer niches shape the oral microbiome. To investigate the contributions of

lifestyle versus environment to the hunter-gatherer oral microbiome, we compared the Agta

(n=138) to smaller samples of Bayaka hunter-gatherers from the Congo (n=21) and neighboring

Palanan farmers from the Philippines (n=14) (see Methods). Both Agta (mean of 252.1±90 ASVs

per individual) and Bayaka (280.1±83) exhibited significantly more ASVs than Palanan farmers

(163.4±34) (P<0.0001; Figure 1a), and higher levels of ASV diversity as measured by Faith's

Phylogenetic Diversity index (Figure 1b). Comparisons based on the total set of ASVs in each

population (controlling for differences in sample size through subsampling) revealed that the

Agta shared more bacteria with African Bayaka (471.2±33.9) than with neighboring Palanan

farmers (423.4±23.8) (Figure 1c). Finally, Agta and Bayaka resampled groups showed

respectively 651.5±93.7 and 688.1±61.9 exclusive ASVs, against only 285.1±16.9 in Palanan

farmers (Figure 1d). In summary, the two hunter-gatherer populations show higher microbiome

diversity and uniqueness than Palanan farmers, consistent with findings that farming

significantly reduced gut microbiome diversity[21]. Results therefore demonstrate the precedence

of niche over geography in shaping hunter-gatherer oral microbiomes.

box limits represent first and third quartiles (****: FDR-adjusted P<0.0001; ***: P<0.001; **:

P<0.01).

The social microbiome is a socially transmitted fraction of the oral microbiome. Primate

social 'pan-microbiomes' were recently defined as the totality of microorganisms present in a

host population or species[22], but this definition also includes microorganisms acquired due to

common diet or environment. Here, we define the 'social microbiome' as the oral microbiome

specifically transmitted through social interactions. To identify the socially transmitted fraction

of the Agta oral microbiome, we used the contact network recorded through radio sensor devices

and split all Agta dyadic social interactions into a strong (top 25% from the distribution of dyadic

link weights) and a weak set (the remaining 75% links; see Methods). We then tested for

differences in the proportion of each of the 1980 ASVs between the strong and weak sets. We

identified 137 ASVs (7% of the Agta oral microbiome; see Supplementary Figure 1 and

Supplementary Table 1) whose presence was significantly higher in the strong set, and therefore

statistically associated with higher frequencies of social interactions. In the following we

investigate the transmission patterns and composition of the hunter-gatherer social microbiome.

The hunter-gatherer social microbiome is predominantly pathogenic. Human sociality is

associated with multiple fitness benefits, including increased reproductive success[7], reputation[23],

food sharing[24], cooperation[3,25] and cultural transmission[26], but may also facilitate pathogen

transmission[7,18]. In our dataset, from the 18 ASVs that could be classified at species level, 14 are

socially transmitted, 9 of which (64.3%) are typically pathogenic, and 10 (71.4%) are typically

oral. By contrast, all four non-socially transmitted species were non-pathogenic and typically

oral.

We were able to classify 1886 of the 1980 ASVs at genus level, resulting in 36 socially

(those representing ASVs included in the social microbiome) and 62 non-socially transmitted

genera (the remaining ones). Among the social genera, 61.8% were classified as typically or

exclusively pathogenic (21 out of 34; two genera could not be classified), against only 16.4%

among non-socially transmitted genera (10 out of 61; one genus could not be classified). We

identified many socially transmitted genera either typically (*Aggregatibacter*, *Capnocytophaga*)

or uniquely (*Corynebacterium*) associated with dental plaque formation, gingivitis and calculus,

the full red complex of periodontal disease (*Porphyromonas*, *Treponema* and *Tannerella*), and

other potential periodontal pathogens (*Prevotella*, *Desulfobulbus*, *Fusobacterium*)[27–30]. The

classification of bacterial genera as pathogenic is not unequivocal for those cases where some

species within the genus can be pathogenic and others non-pathogenic. Thus, practical criteria

were applied in these cases for assigning a genus to the pathogenic group (see Methods).

Following these criteria, the social microbiome clearly has a higher proportion of pathogenic

organisms than the non-socially transmitted portion.

We also found pathogenic bacteria typical of the gut (*Rickenellaceae*), non-human

environments (*Tetragenococcus*, *Comamonas*), respiratory tract (*Staphylococcus*, *Moraxella*,

Streptococcus pneumoniae), and both urogenital and respiratory tracts (*Mycoplasma*), suggesting

that their spread may be facilitated by oral transmission[31–33]. In summary, the predominantly

pathogenic nature of the social microbiome suggests a trade-off between benefits of hunter-

gatherer sociality and costs associated with disease transmission.

Hunter-gatherer multilevel social structure shapes social microbiome sharing. Hunter-

gatherer sociality is characterised by specific interaction channels not found in non-human apes,

such as long-term pair bonding and households, extended families, friendships among unrelated

individuals, and frequent between-camp relocation. We estimated the effect of relatedness level,

residence camp and friendships on the probability of sharing socially transmitted bacteria. First,

we built a bacterial sharing network, where the weight of each Agta dyadic link is given by how

many of the 137 social bacteria are shared by the two Agta individuals (rather than by the

strength of its social bond, as in the social network). Next, we classified all dyadic links in this

network into: i) levels of kinship (mother-offspring, father-offspring, siblings: r=0.5, other kin:

r=0.25 or r=0.125, non-kin: r=0.0625 or lower, spouses, friends: defined as non-kin at the top

25% distribution of social dyadic weights, and other non-kin) and ii) residence (same or different

camp, same or different household). Finally, we compared the mean weight of each type of

dyadic link in our bacterial sharing network to its mean weight in a sample of 1000 networks of

the same size and topology, but where the dyadic classification was randomised (Figure 2 and

Supplementary Tables 2 and 3).

Figure 2. Effect of kinship, friendship and residence on dyadic bacterial sharing. Dyads were classified into kinship levels; same or different households; same or different camps; and between friends in the same or different camps. Dots show the z-score, or the standardised ratio of the mean link weight in real to randomised networks, in either social (orange) or non-socially transmitted bacteria (purple). Vertical red dashed line indicates a ratio of 1, or no difference between the number of shared bacteria in real and randomised networks. For socially transmitted bacteria, kinship, friendship and residence in the same household or camp are associated with significantly higher bacterial sharing than predicted from randomised networks of the same size and structure. By contrast, dyads from different camps or non-kin share significantly fewer bacteria than expected by chance; bacterial sharing in dyads from different households do not differ from randomised networks. For non-socially transmitted bacteria, the only dyadic categories significantly increasing bacterial sharing were siblings, spouses, and dyads from the same household (all of which share the same close environment). See Supplementary Tables 2 and 3 for values on mean weights for real and randomised networks. (****: FDR-adjusted $P<0.0001$; ***: $P<0.001$; **: $P<0.01$).

Results showed that some dyadic categories share significantly more socially transmitted bacteria than expected by chance. First, we observed higher bacterial sharing within the same household and camp than between different households and camps, an expected consequence of the Agta multilevel social structure. Kinship effects were also clear, with the highest levels of social microbiome sharing found in mother-offspring pairs, followed by siblings known interact every day in households and playgroups. High sharing between spouses also confirmed the importance of human pair bonding in microbial transmission. In addition, strong friendship links were also associated with increased bacterial sharing. Social bacterial sharing between friends in the same camp is as high as between close kin or within households. Friends in different camps also share a higher proportion of social bacteria than expected by chance, which is possibly a consequence of high between-camp mobility. By contrast, non-kin or individuals from different camps share fewer social bacteria than expected, further demonstrating the role of Agta friendships in the transmission of social bacteria across households and whole camps.

The same analysis performed instead on non-socially transmitted ASVs did not reveal

significant effects on bacterial sharing from most dyadic categories, except for three types:

spouses, siblings, and same household. A possible explanation is that some non-socially

transmitted bacteria may be shared due to a common environment and diet in the same

household. For example, we have shown in our parallel study[19] that the proportion of meat

versus rice in individual diets affects the composition of the oral microbiome. Therefore, similar

diets may explain the presence of the same ASVs within the individuals of a household

irrespective of social interaction levels. However, the effects of sociality and shared diets seem to

be independent. This is shown by the fact that socially transmitted bacteria are equally likely to

be related or not to diet: 13 socially transmitted genera were found also associated to diet (41

genera), whereas 23 were not (57 genera) (proportion test: chi-squared=0.44, P=0.51; See

companion paper[19] for further data and analysis). There we also show that host genotype

correlates with the presence of certain ASVs19. While high genetic relatedness may play a role

in bacteria sharing within households, none of the ASVs associated with the host genotype were

present in the social microbiome. Therefore, our analyses seem to distinguish between the effects

of social contact from shared environment or genes within dyadic types. Overall, the results

show the roles of mobility and the multiple interaction channels created by multilevel sociality in

social microbiome sharing, similarly to what is also observed in cultural transmission[6,18,26,34].

Frequency of social contact predicts social microbiome sharing. Although previous studies

have investigated patterns of bacterial sharing in human groups, they have often been unable to

comprehensively characterize transmission patterns due to limited information on social

contact[35]. In order to obtain a full picture of individual contact and exposure levels, we built

social networks based on proximity data from four camps and two multi-camp locations (Figure

3a-b). Overall, Agta social networks reveal a multilevel structure of households (mostly

consisting of strong kin links) connected by a few strong links (mostly among unrelated friends)

in each camp, and in the case of multi-camp groups, camps interconnected mostly due to visits

among friends. We also observed equality of interactions within and between sexes and across

age groups[6,18]. This pattern creates multiple channels for social transmission of bacteria both

within and between camps, between close kin and unrelated individuals, and finally across whole

multi-camp structures.

Figure 3. Characterisation of the social microbiome. a) Recording networks of social
interactions using radio sensor technology. b) Reinforcement analysis estimates the probability
of a link occurring in the bacterial sharing network (top layer, purple) based on the weight of the
same link in the social contact network (bottom layer, blue). Network nodes (circles) represent
the same Agta individuals in the bacterial sharing and social contact network. Panel displays
networks from multi-camp 1 (23 individuals). Map shows geographical location of four camps

interconnected by frequent migration. c) Probabilities of links in the Agta bacterial sharing
network increase with their weights in the social contact network. Curves estimated by
generalised additive modelling (binomial option). Data from four Agta camps and two multi-
camps. d) Eigenvector centralities in bacterial sharing and social contact networks. Linear
regression based on pooled data from four Agta camps and two multi-camp structures. Virtually
similar results were obtained by including camp either as a fixed factor in a multiple regression
(with or without interactions), or as a random factor (on intercept and slope) in a mixed effects
linear regression.

To further assess whether increasing levels of social contact predict higher levels of

sharing of socially transmitted bacteria, we applied reinforcement analysis[36] (Figure 3b) to assess

whether the social network predicts (or reinforces) the bacterial sharing network in each Agta

camp. We calculated the conditional probability of each link between two individuals A and B in

the bacterial network provided the same weighted link is present in the social network

(Supplementary Table 4 and 5). For all four camps and two multi-camp structures, results

showed that the weight of a dyadic link in the social network significantly predicts the

probability of the same link occurring in the bacterial sharing network (Figure 3c and

Supplementary Figure 2). Specifically, a larger dyadic weight in the social network implies a

higher probability that the same individuals also share at least one socially transmitted ASV. For

example, for multi-camp 1, while weak social network links (with weights under 10 recorded

social interactions) show a probability below 20% of sharing any socially transmitted bacteria,

strong links (over 200 recorded contacts) are associated with a probability above 70%. Overall,

the results confirm that the Agta social microbiome is shaped by their social interactions.

Hypersocial individuals are supersharers. We also investigated whether more socially interactive individuals exhibited higher social microbiome diversity. We calculated eigenvector centralities for all individuals in the bacterial sharing network, resulting in a significant and positive slope in a regression on eigenvector centralities from the same individuals in the social network (b=0.32, P=0.0001, R^2 = 0.1, n=138, Figure 3d). We also identified 16 individuals ranked at the top quartile of eigenvector centralities in both networks as potential microbial 'superspreaders' or 'superacquirers', that is, "supersharers". They do not stem from a specific age (7 to 68 years) or sex (six males, ten females), which is compatible with the egalitarian social structure of hunter-gatherers allowing individuals from any age or sex to be potentially central to social networks.

Discussion

We have identified and characterized a socially transmitted fraction of the Agta hunter-gatherer oral microbiome. This fraction (7%) is surprisingly small, since in principle all 1980 identified ASVs could be orally transmitted between closely interacting people. Nonetheless, our results demonstrate a significant and independent role of social interactions on the transmission of oral microbiome, in addition to other factors such as shared environment (household and diet) and host characteristics (age, sex and genes) previously investigated in other hunter-gatherer populations[8–16] and in the same Agta population[19]. The transmission of the 137 bacteria classified into the social microbiome seem to be facilitated by the extended sociality of hunter-gatherers and its various transmission channels, ranging from spouses to unrelated friends often residing in different camps. Together, reinforcement analysis, multiple channels of social interaction, and supersharers show that social microbiome sharing is strongly shaped by hunter-gatherer multilevel sociality. From an evolutionary perspective, sociality has considerably changed from

our closest ape relatives to ancestral humans, when adopting a hunter-gathering lifestyle meant

exhibited higher frequency of social contact with unrelated individuals, larger networks of

extended kin across large geographical regions, and more egalitarian interactions between and

within sexes and across ages. Such changes may have affected patterns of pathogen transmission

and affected the human microbiome as observed in current hunter-gatherers. As with our study

of the Agta, future research should collect data on both social networks and social microbiomes

from the same populations of non-human apes; such a dataset would provide a comparative basis

for analysing of the role of social evolution on the human social microbiome. Hunter-gatherer

social networks are efficient systems of cultural transmission, and its specific channels organised

around kinship, friendship and camp interconnectivity are central for the organisation of

between-household cooperation, food sharing and social learning[24,26]. Agta mothers with higher

social network centrality enjoy increased access to help and reproductive success[7], but our results

have shown that efficient networks may also facilitate the spread of infectious diseases, and

hence significantly affect the structure and composition of the Agta microbiome. Crucially the

frequency of pathogenic bacteria is much higher in the socially than in the non-socially

transmitted fraction of the Agta oral microbiome. Together with the association between

hypersocial individuals and increased bacterial sharing, this suggests a tradeoff between potential

fitness benefits and costs of increased pathogen transmission. We conclude that the

predominantly pathogenic oral social microbiome we identified in hunter-gatherers may be at

least partially the outcome of a tradeoff between the advantages of multilevel sociality and the

cost of infectious disease.

Methods

Ethnographic data collection

Agta demography. Ethnographic data collection took place over two seasons in April-June 2013

and February-October 2014. We censused 915 Agta individuals (54.7% male) across 20 camps.

For the current study we selected four camps and two multi-camp structures where we collected

data both on proximity networks and saliva samples. Accurate ages were estimated following

relative aging protocols[37]. Relatedness (biological and affinal) was based on household

genealogies. To resolve inconsistencies, we took either the genealogy from the most

knowledgeable individual (i.e. mother over aunt) or the genealogy that reduced other

inconsistencies (i.e. discarding six-month interbirth intervals). Genealogies contained 2953 living

and dead Agta. We used the *R* packages *pedigree*, *kinship2*, and *igraph* to measure

consanguineous relatedness (r)[34,38]. For comparative purposes, we obtained 14 saliva samples

from neighbouring Palanan farmers, making sure individuals were unrelated by directly asking.

Bayaka demography. Ethnographic data collection took place over two seasons in April-June

2013 and February-October 2014. We collected saliva samples from 21 individuals for

microbiome analyses.

Ethics. This study was approved by UCL Ethics Committee (UCL Ethics code 3086/003) and

carried out with permission from local government and community members. Informed consent

was obtained from all participants, after group and individual explanation of research objectives

in the indigenous language. A small compensation (usually a thermal bottle or cooking utensils)

was given to each participant. The National Commission for Indigenous Peoples (NCIP), advised

us that the process of Free Prior Informed Consent with the tribal leaders, youth and elders

would be necessary to validate our data collection under their supervision. This was done in2017

with the presence of all tribal leathers, elders and youth representatives at the NCIP regional

office, with the mediation of the regional officer and the NCIP Attorney. The validation process

was approved unanimously by the tribal leaders, and the NCIP, and validated the full 5 years of

data collection.

Oral microbiome analysis

Microbial DNA extraction and 16S rRNA gene sequencing. A total of 190 saliva samples were

selected from Agta hunter-gatherers (n = 155) and Palanan farmers in the Philippines (n = 14),

and Bayaka hunter-gatherers from the Congo (n = 21). Microbial DNA was extracted following

the protocol for manual purification of DNA for Oragene·DNA/saliva samples. The 16S rRNA

gene V3-V4 region was amplified by PCR with primers containing Illumina adapter overhang

nucleotide sequences. All PCR products were validated through an agarose gel and purified with

magnetic beads. Index PCR was then performed to create the final library also validated through

an agarose gel. All samples were pooled together at equimolar proportions and the final pool was

qPCR-quantified before MiSeq loading. Raw Illumina pair-end sequence data were

demultiplexed and quality-filtered with *QIIME* 2 2019.1[39] and *DADA2*[40], which generates single

nucleotide exact amplicon sequence variants (ASV or ESV). ASVs are biologically meaningful

as they identify a specific sequence and allow for higher resolution than operational taxonomic

units (OTUs)[41] or clusters of sequences above a similarity threshold, and thus an ASV is

equivalent to a 100% similar OTU. Taxonomic information was assigned to ASVs using a naïve

Bayes taxonomy classifier against the *SILVA* database release 132 with a 99% identity

sequence[42].

Reads outside the kingdom Bacteria or assigned to mitochondria or chloroplasts were

removed. Phylogenetic analyses aligned sequences with *MAFFT*[43] and generated a rooted

phylogenetic tree with *FastTree2*[44] using default settings via *QIIME* 2. We generated an Alpha

rarefaction curve with R package *vegan* to confirm that sample richness had been fully observed

(Supplementary Figure 3).

Samples with extremely low number of reads (8000) were removed. This resulted in

6409 ASVs (later reduced to 1980 ASVs present in at least two individuals and abundance of at

least 10 counts per individual) and 173 individuals: 138 Agta, 21 Bayaka, and 14 Palanan

farmers.

Identification of the Agta social microbiome. In our Agta sample, we first selected a set of strong

social links (top 25% of the weight distribution from each camp and multi-camp). For each ASV,

we calculated the proportion of strong links (f_A^s) where a given ASV A was present. Next, we

calculated the same proportion in the complementary set of 75% weak social links (f_A^W). We then

computed for each ASV A the score $s_A^S = \frac{f_A^S - f_A^W}{f_A^S + f_A^W}$, or normalised difference between the two

proportions. This score can be paired with the z-score $\frac{f_A^S - f_A^W}{\sqrt{f_A(1-f_A)(\frac{1}{n^S}+\frac{1}{n^W})}}$ which quantifies the

deviation from the null hypothesis that the two proportions are equal, with n^S and n^W as

respectively the numbers of strong and weak links and f_A as the proportion of total links that

share ASV A. We then selected as affected by social interaction 137 ASVs with s >0.5 and P<

0.05. P-values were adjusted by False Discovery Rate (FDR).

We performed a sensitivity analysis to investigate the consequences of varying the

threshold defining strong versus weak social links in the Agta social network. Instead of 137

ASVs resulting from selecting dyads at top 25% of the weight distribution, we obtained:

Top 45%: 166 ASVs

Top 35%: 156 ASVs

Top 25%: 137 ASVs

Top 15%: 138 ASVs

Top 5%: 99 ASVs

The list shows that whether the strong set consists of a very reduced number dyads with

very strong weights (top 5%), or instead includes nearly half the dyads (top 45%), the number of

ASVs significantly responsive to social contact varies from 99 (5%) to 166 (8%), representing a

small fraction of the total of 1980 ASVs found in the Agta. Therefore, setting the threshold at the

top 25% did not affect our results and conclusions.

Agta social network data, construction and analysis

Mote devices. Motes are wireless sensing devices storing all between-device communications

within a specified distance and have been described in detail elsewhere[6,7,18,45]. We used the

UCMote Mini (with a TinyOS operating system) sealed into wristbands or belts, labelled with a

unique number, and identified with coloured string to avoid accidental swaps. Motes require no

grounded infrastructure and collect interactions even when individuals are away from camps.

Individuals arriving at a camp after the start of data collection were given a mote and entry time

was recorded, while those leaving a camp before the end of data collection had their exit time

recorded. To prevent swaps individuals were checked twice daily, and mote numbers were

checked upon return. Any swaps were later corrected by reassigning data to the correct

individuals.

Data were later downloaded via a PC side application in *Java*. Data were limited to 5am-

8pm. We ran raw data through a stringent data-processing system in *Python* to prevent data

corruption. Data were matched to ID numbers and start-stop times of each mote. The result was a

matrix with the number of recorded beacons for all possible dyads and their weights.

For the camp-level experiment, all individuals from four camps wore motes from five to

seven days. Each device sent a message every two minutes that contained its unique ID, a time

stamp and the signal strength. Messages are stored by any other mote within a three-meter

radius, a frequently used threshold[46,47]. For the multi-camp experiment, adult individuals from

two areas (consisting of seven and three camps respectively) wore motes for one month.

Effect of dyad category on bacterial sharing. The bacterial sharing network was constructed by

defining link weights as the number of social bacteria shared by two individuals. Dyads in the

network were classified into: i) levels of kinship (mother-offspring, father-offspring, siblings:

r=0.5, other kin: r=0.25 or r=0.125, non-kin: r=0.0625 or lower, spouses, friends: defined as

non-kin at the top 25% distribution of social dyadic weights, and other non-kin) and ii) residence

(same or different camp, same or different household). Mean weights were calculated for each

dyadic category (Supplementary Table 2). Then, we produced 1000 network randomisations

based on a single-step ID swap between nodes. For example, if dyad 1 consisted of two spouses

in the real network, randomisation preserved dyad 1 and its weight, but randomly replaced the

two nodes (potentially changing the dyadic classification to siblings, friends, etc.). We calculated

the mean weights for each dyadic category in the 1000 randomised networks, and then calculated

one-sample t-tests with the mean weight in the real network as the test value. We repeated the

analysis for non-social bacteria (Supplementary Table 3).

Reinforcement analysis. In a multilayer network, reinforcement analysis measures the overlap in

links between different layers to quantify the probability of finding a link on a layer conditioned

on the weight of the same link on another[48]. Reinforcement between two network layers α and

α', or $P(\alpha'|\alpha)$, is defined as

$P(\alpha'|\alpha) = \frac{\sum_{ij} a_{ij}^{[\alpha']} w_{ij}^{[\alpha]}}{\sum_{ij} w_{ij}^{[\alpha]}}$ (1)

where $a_{ij}^{[\alpha']}$ is the adjacency matrix of conditioned layer α', and $w_{ij}^{[\alpha]}$ is the adjacency matrix of

the conditioning layer α. We split the weight of social links into three tertiles, and computed

equation (1) for each. We obtained increasing values of reinforcement from the lower to the

higher tertile, providing evidence of an effect of social contact on bacterial sharing.

ASV classification

ASV diversity metrics. To distinguish between the effects of lifestyle and shared ecology on the

microbiome, we compared the diversity of the oral microbiome of Agta hunter-gatherers with

neighbouring Palanan farmers in the Philippines and with Bayaka hunter-gatherers in Congo.

Using the 6409 ASVs dataset we calculated the number of observed ASVs in each population

with *R* package *Phyloseq* (version 1.30.0) and the Faith's Phylogenetic Diversity index was

calculated with *R* package *picante* using the generated rooted phylogenetic tree. To estimate the

number of shared ASVs in the Agta, Bayaka and Palanan farmers, we sampled a random subset

of 10 samples for each population without replacement, calculated the shared ASVs between the

populations and repeated this procedure 100 times. Global differences between groups and

pairwise comparisons were assessed by Kruskal-Wallis and Wilcoxon Rank Sum tests

respectively and plotted by the *R* package *ggpubr*. Pairwise p-values were adjusted by False

Discovery Rate (FDR).

Classification of oral bacteria as pathogens. ASVs were classified as oral pathogens if they have

been reported as etiological agents of periodontitis[49,50] or dental caries[51,52]. For gum disease

pathogens, these included the classical "red" and "orange" complex of periodontal pathogens and

the recent update by Pérez-Chaparro and cols[50] based on systematic review and metaanalysis.

For caries pathogens, the list of active microorganisms detected through metatranscriptomics of

cavities was used. Common oral commensals potentially causing endocarditis or systemic

infections in immunocompromised patients only were not considered pathogens. Bacteria

reported as etiological agents of lower respiratory infections (e.g. pneumonia, whooping cough,

bronchitis, or sinusitis) and biofilm-mediated infections (e.g. lactational mastitis, medical

implant biofilm infections, chronic lung infections, osteomyelitis or chronic wounds) were also

considered pathogens, including organisms present in healthy carriers[53,54]. Bacteria causing

urinary tract infections or sexually transmitted diseases transiently found in the oral cavity were

also considered pathogens[55]. Bacteria were classified as "oral" if detected in more than 10% of

the population in oral samples according to the Human Oral Microbiome database. If a bacterial

species or genus had been isolated from the oral cavity of an animal, it was also classified as

oral.

For assignment of bacteria to pathogenic or non-pathogenic, we used species-level ASVs,

given that there are multiple cases where different species from the same genus had a different

assignment. If taxonomic classification of the ASV was only possible at the genus level, it was

considered a pathogen if: i) >90% of named species within the genus were pathogenic, or ii) the

genus included a major pathogenic species but the remaining species within the genus were not

classified as oral by the Human Oral Microbiome Database[56]. ASV with a top hit to a sequence

classified as "Oral taxa" in databases but without a species assignment were not considered

named species and were discarded from the analysis. Cases where taxonomic classification of the

ASV was only possible at the family level or higher were also discarded.

Acknowledgements

This project was funded by Leverhulme Trust Grant RP2011-R 045 to A.B.M. J.B. received

grant PID2019-110933GB-I00/AEI/10.13039/501100011033 by the Agencia Estatal de

Investigación (AEI), Ministerio de Ciencia, Innovación y Universidades (MCIU, Spain) and

support from Secretaria d'Universitats i Recerca del Departament d'Economia i Coneixement de

la Generalitat de Catalunya (GRC 2017 SGR 702). Part of the "Unidad de Excelencia María de

Maeztu", funded by the AEI (CEX2018-000792-M), that granted the microbiome analysis by the

Servei de Genòmica, Universitat Pompeu Fabra. A.M is funded by grant RTI2018-102032-B-100

from Ministerio de Ciencia, Innovación y Universidades (MCIU, Spain).

Data availability

16S amplicon data (EGAS00001005317) are deposited at the European Genome-phenome

Archive (EGA), which is hosted at the EBI and the CRG. Data at the individual level on age,

kinship relationships, household composition, camp assignation and social contacts that support

the findings of this study are available on request from the corresponding author (A.B.M.). The

individual data are not publicly available due to information that could compromise research

participant privacy.

Code availability

Source code and data for visualization are available at https://doi.org/10.5281/zenodo.6338840

References

1. Migliano, A. B., Vinicius, L. & Lahr, M. M. Life history trade-offs explain the evolution of human pygmies. *Proc. Natl. Acad. Sci. U. S. A.* **104**, (2007).

2. Hill, K. R. *et al.* Co-residence patterns in hunter-gatherer societies show unique human social structure. *Science (80-.).* **331**, 1286–1289 (2011).

3. Apicella, C. L., Marlowe, F. W., Fowler, J. H. & Christakis, N. A. Social networks and cooperation in hunter-gatherers. *Nature* **481**, 497–501 (2012).

4. Smith, K. M., Larroucau, T., Mabulla, I. A. & Apicella, C. L. Hunter-gatherers maintain assortativity in cooperation despite high levels of residential change and mixing. *Curr. Biol.* **28**, 3152–3157 (2018).

5. Derex, M. & Boyd, R. Partial connectivity increases cultural accumulation within groups. *Proc. Natl. Acad. Sci.* **113**, 2982–2987 (2016).

6. Migliano, A. B. *et al.* Hunter-gatherer multilevel sociality accelerates cumulative cultural evolution. *Sci. Adv.* **6**, (2020).

7. Page, A. E. *et al.* Hunter-Gatherer Social Networks and Reproductive Success. *Sci. Rep.* **7**, (2017).

8. Nasidze, I. *et al.* High Diversity of the Saliva Microbiome in Batwa Pygmies. *PLoS One* **6**, e23352 (2011).

9. Schnorr, S. L. *et al.* Gut microbiome of the Hadza hunter-gatherers. *Nat. Commun.* **5**, 3654 (2014).

10. Rampelli, S. *et al.* Metagenome Sequencing of the Hadza Hunter-Gatherer Gut Microbiota. *Curr. Biol.* **25**, 1682–1693 (2015).

11. Smits, S. A. *et al.* Seasonal cycling in the gut microbiome of the Hadza hunter-gatherers of Tanzania. *Science* **357**, 802–806 (2017).

12. Fragiadakis, G. K. *et al.* Links between environment, diet, and the hunter-gatherer microbiome. *Gut Microbes* **10**, 216–227 (2019).

13. Schnorr, S. L. The Diverse Microbiome of the Hunter-Gatherer. *Nature* **518**, S14–S15 (2015).

14. Gomez, A. *et al.* Gut Microbiome of Coexisting BaAka Pygmies and Bantu Reflects Gradients of Traditional Subsistence Patterns. *Cell Rep.* **14**, 2142–2153 (2016).

15. Moeller, A. H. The shrinking human gut microbiome. *Curr. Opin. Microbiol.* **381. A. H.**, 30–35 (2017).

16. Meehan, C. L. *et al.* Social networks, cooperative breeding, and the human milk microbiome. *Am. J. Hum. Biol.* **30**, e23131 (2018).

17. Lassalle, F. *et al.* Oral microbiomes from hunter⬜gatherers and traditional farmers reveal shifts in commensal balance and pathogen load linked to diet. *Mol. Ecol.* **27**, 182–195 (2018).

18. Migliano, A. B. *et al.* Characterization of hunter-gatherer networks and implications for cumulative culture. *Nat. Hum. Behav.* **1**, 1–6 (2017).

19. Dobon, B. *et al.* The making of the oral microbiome in Agta hunter-gatherers. (2021).

20. Callahan, B. J., McMurdie, P. J. & Holmes, S. P. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J.* **11**, 2639–2643 (2017).

21. Schnorr, S. L. *et al.* Gut microbiome of the Hadza hunter-gatherers. *Nat. Commun.* **5**, 3654 (2014).

22. Sarkar, A. *et al.* Microbial transmission in animal social networks and the social microbiome. *Nat. Ecol. Evol.* **4**, 1020–1035 (2020).

23. Chaudhary, N. *et al.* Polygyny without wealth: popularity in gift games predicts polygyny
in Bayaka Pygmies. *R. Soc. open Sci.* **2**, 150054 (2015).
24. Dyble, M. *et al.* Networks of Food Sharing Reveal the Functional Significance of
Multilevel Sociality in Two Hunter-Gatherer Groups. *Curr. Biol.* **26**, (2016).
25. Smith, D. *et al.* Camp stability predicts patterns of hunter-gatherer cooperation. *R. Soc.*
Open Sci. **3**, (2016).
26. Salali, G. D. *et al.* Knowledge-sharing networks in hunter-gatherers and the evolution of
cumulative culture. *Curr. Biol.* **26**, 2516–2521 (2016).
27. Socransky, S. S., Haffajee, A. D., Cugini, M. A., Smith, C. & Kent Jr., R. L. Microbial
complexes in subgingival plaque. *J. Clin. Periodontol.* **25**, 134–144 (1998).
28. Pérez-Chaparro, P. J. *et al.* Newly Identified Pathogens Associated with Periodontitis: A
Systematic Review. *J. Dent. Res.* **93**, 846–858 (2014).
29. Simón-Soro, A., Guillen-Navarro, M. & Mira, A. Metatranscriptomics reveals overall
active bacterial composition in caries lesions. *J. Oral Microbiol.* **6**, 25443 (2014).
30. Simón-Soro, A. & Mira, A. Solving the etiology of dental caries. *Trends Microbiol.* **23**,
76–82 (2015).
31. Jung, H.-S., Ehlers, M. M., Lombaard, H., Redelinghuys, M. J. & Kock, M. M. Etiology
of bacterial vaginosis and polymicrobial biofilm formation. *Crit. Rev. Microbiol.* **43**, 651–
667 (2017).
32. Natsis, N. E. & Cohen, P. R. Coagulase-Negative Staphylococcus Skin and Soft Tissue
Infections. *Am. J. Clin. Dermatol.* **19**, 671–677 (2018).
33. Haq, I. J., Battersby, A. C., Eastham, K. & McKean, M. Community acquired pneumonia
in children. *BMJ* **356**, j686 (2017).
34. Dyble, M. *et al.* Sex equality can explain the unique social structure of hunter-gatherer
bands. *Science (80-.).* **348**, 796–798 (2015).
35. Brito, I. L. *et al.* Transmission of human-associated microbiota along family and social
networks. *Nat. Microbiol.* **4**, 964–971 (2019).
36. Battiston, F., Nicosia, V. & Latora, V. Structural measures for multiplex networks. *Phys.*
Rev. E **89**, 32804 (2014).
37. Diekmann, Y. *et al.* Accurate age estimation in small-scale societies. *Proc. Natl. Acad.*
Sci. U. S. A. **114**, (2017).
38. Page, A. E., Myers, S., Dyble, M. & Migliano, A. B. Why so many Agta boys? Explaining
'extreme' sex ratios in Philippine foragers. *Evol. Hum. Sci.* **1**, (2019).
39. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data
science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).
40. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat. Methods* **13**, 581–583 (2016).
41. Callahan, B. J., McMurdie, P. J. & Holmes, S. P. Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *ISME J.* **11**, 2639–2643 (2017).
42. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2013).
43. Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid
multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**,
3059–3066 (2002).
44. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood
trees for large alignments. *PLoS One* **5**, e9490–e9490 (2010).

45.   Page, A. E. *et al.* Testing adaptive hypotheses of alloparenting in Agta foragers. *Nat.*
594   *Hum. Behav.* **3**, 1154–1163 (2019).
595   46.   Brent, L. J. N., Lehmann, J. & Ramos-Fernández, G. Social network analysis in the study
596   of nonhuman primates: A historical perspective. *Am. J. Primatol.* **73**, 720–730 (2011).
597   47.   Isella, L. *et al.* What's in a crowd? Analysis of face-to-face behavioral networks. *J. Theor.*
598   *Biol.* **271**, 166–180 (2011).
599   48.   Battiston, F., Nicosia, V. & Latora, V. Structural measures for multiplex networks. *Phys.*
600   *Rev. E* **89**, 32804 (2014).
601   49.   Socransky, S. S., Haffajee, A. D., Cugini, M. A., Smith, C. & Kent Jr., R. L. Microbial
602   complexes in subgingival plaque. *J. Clin. Periodontol.* **25**, 134–144 (1998).
603   50.   Pérez-Chaparro, P. J. *et al.* Newly Identified Pathogens Associated with Periodontitis: A
604   Systematic Review. *J. Dent. Res.* **93**, 846–858 (2014).
605   51.   Simón-Soro, A., Guillen-Navarro, M. & Mira, A. Metatranscriptomics reveals overall
606   active bacterial composition in caries lesions. *J. Oral Microbiol.* **6**, 25443 (2014).
607   52.   Simón-Soro, A. & Mira, A. Solving the etiology of dental caries. *Trends Microbiol.* **23**,
608   76–82 (2015).
609   53.   Leung*, A. K. C. & Hon, A. H. C. W. and K. L. Community-Acquired Pneumonia in
610   Children. *Recent Patents on Inflammation & Allergy Drug Discovery* vol. 12 136–144
611   (2018).
612   54.   Natsis, N. E. & Cohen, P. R. Coagulase-Negative Staphylococcus Skin and Soft Tissue
613   Infections. *Am. J. Clin. Dermatol.* **19**, 671–677 (2018).
614   55.   Jung, H.-S., Ehlers, M. M., Lombaard, H., Redelinghuys, M. J. & Kock, M. M. Etiology
615   of bacterial vaginosis and polymicrobial biofilm formation. *Crit. Rev. Microbiol.* **43**, 651–
616   667 (2017).
617   56.   Chen, T. *et al.* The Human Oral Microbiome Database: a web accessible resource for
618   investigating oral microbe taxonomic and genomic information. *Database (Oxford).* **2010**,
619   baq013–baq013 (2010).
620
621
622