Genome Mining and Biochemical Characterization of a Bifunctional Type I Diterpene Synthase from a Marine-Derived Fungus

Peng Zhang,[†,‡] Guangwei Wu,[†,‡, §,ǁ] Stephanie C. Heard,[†] Changshan Niu,[†] Yonghui Zhang,[#] and Jaclyn M. Winter[†,*]

†Department of Medicinal Chemistry, University of Utah College of Pharmacy, Salt Lake City, Utah, 84112, USA;

#Hubei Key Laboratory of Natural Medicinal Chemistry and Resource Evaluation, School of Pharmacy, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China.

KEYWORDS. fungal type I diterpene synthase, genome mining, biosynthesis, marine-derived fungi

ABSTRACT: We report the identification of the *tnd* biosynthetic cluster from the marine-derived fungal strain *Aspergillus flavipes* CNL-338 and *in vivo* characterization of a cryptic type I diterpene synthase. Heterologous expression of the bifunctional terpene synthase TndC in *Saccharomyces cerevisiae* led to the discovery of a new diterpene backbone, talarodiene **8**, harboring a benzo[*a*]cyclopenta[*d*]cyclooctane tricyclic fused ring system. The cyclization mechanism that converts geranylgeranyl diphosphate to the tricyclic hydrocarbon skeleton was investigated using [13]C-labeling studies and stable isotope tracer experiments showed the biotransformation of **8** into the natural product talaronoid C.

Terpenes are the largest class of natural products and are produced by all kingdoms of life. These compounds possess enormous structural diversity and exhibit various biological activities ranging from anticancer and antimalarial to carcinogens and mycotoxins.[1] Despite their structural complexity, all terpenes are derived from the universal C_5 hemiterpene precursors dimethylallyl diphosphate (DMAPP) and isopentenyl diphosphate (IPP). Coupling of these C_5 precursors, facilitated by prenyltransferases (PTs), generates linear, achiral polyprenyl diphosphates that can be transformed by terpene cyclases (TCs) into complex scaffolds containing multiple fused rings and stereogenic centers.[2-6] The structural diversity associated with terpenes often originates from the cyclization step, and TCs catalyze some of the most complex reactions in natural products chemistry.

In fungi, although condensation and cyclization reactions occur independently, bifunctional terpene synthases have been characterized where the C-terminal half is responsible for producing the polyprenyl diphosphate and the N-terminal half catalyzes the cyclization reaction. The first fungal type I diterpene synthase, PaFS, was characterized in 2007 from *Phomopsis amygdali* and shown to produce fusicoccin **1**.[7] AcOS was the first type I sesterterpene synthase characterized in 2013 from *Aspergillus clavatus* and is responsible for the biosynthesis of ophiobolin F **2**.[8] Because of their potential in synthesizing diverse hydrocarbon skeletons, subsequent genome mining efforts focused on identifying additional cryptic type I bifunctional terpene synthases. As a result, a number of fungal type I sesterterpene synthases were characterized.[9-17] However,

since the discovery of PaFS, only a limited number of type I diterpene synthases have been identified and include those

Figure 1. Structures of selected fungal diterpenes and sesterterpenes produced by type I bifunctional terpene synthases.

responsible for the production of variediene **3**,[18] phomopsene **4**,[19] brassicicene **5**,[20] a precursor to the cyclopiane-type diterpenes **6**,[21] and dolasta-1(15),8-diene **7**[22] (Figure 1). Given our limited knowledge of type I

diterpene synthases, the discovery and biochemical characterization of new enzymes would bring to light cryptic natural products, unveil novel cyclization reactions, and allow for more informed bioinformatic predictions. In this work, we describe the discovery and *in vivo* characterization of a cryptic bifunctional type I diterpene synthase from a marine-derived fungus that synthesizes a new tricyclic 5-8-6 hydrocarbon skeleton. The use of stable tracer isotope experiments also allowed us to show the biotransformation of the diterpene backbone into the talaronoid class of natural products.

It is known that marine organisms are prolific producers of bioactive natural products and often produce molecules not observed in their terrestrial counterparts.[23] The previously characterized type I bifunctional terpene synthases were identified exclusively from terrestrial fungi, and given the tremendous promise that marine organisms hold for characterizing novel biosynthetic enzymes, we turned to marine-derived fungi as an underexplored resource for identifying and characterizing type I terpene synthases. Recently, our group sequenced the genome of the marine-derived fungus *Aspergillus flavipes* CNL-338,[24] and using the PaFS and AcOS sequences as probes, we scanned the genome for bifunctional terpene synthases. A 21 kb biosynthetic cluster harboring a cryptic chimeric synthase was identified (Figure 2A), and bioinformatic analysis of TndC revealed that the 764 amino acid protein possessed both PT and TC domains. Multiple sequence alignment also showed that TndC contains the conserved aspartate-rich DDxxD motif for Mg^{2+} binding in both the PT and TC domains, as well as a second NSE Mg^{2+}-binding motif in the TC domain indicative of type I cyclases (Figure S2). Phylogenetic comparison of the cryptic chimeric synthase with known fungal-derived diterpene and sesterterpene synthases showed that TndC cladded between PaFS and the

sesterterpene synthase EvAS (Figure S1), suggesting that TndC could be producing a new terpene skeleton; however, it was not clear if the product was a diterpene or sesterterpene.

Figure 2. Characterization of the type I diterpene synthase *tnd*C from *A. flavipes* CNL-338. (**A**) Organization of the *tnd* biosynthetic gene cluster in *A. flavipes* CNL-338. (**B**) GC-MS analysis (TIC) of extracts from *S. cerevisiae* ZXM144 transformed with (i) a plasmid-borne *tnd*C or (ii) empty vector. (**C**) The structure identification of compound **8** and key 2D NMR correlations.

Initial efforts at expressing recombinant TndC from *Escherichia coli* and *Saccharomyces cerevisiae* failed to generate any soluble protein. Thus, to elucidate the product

Figure 3. Proposed biosynthesis of the talarodiene backbone. (**A**) Biosynthesis of the linear achiral precursor geranylgeranyl diphosphate using the C-terminal prenyltransferase domain of TndC. (**B**) Formation of the 5-8-6 tricyclic talarodiene backbone **8** by the N-terminal cyclization domain of TndC. [1,2-$^{13}C_2$]acetate labeling patterns are shown as black bold lines and dots to signify double and single enrichments, respectively. Red dots indicate the C–C bond breakage of an intact acetate unit.

of TndC, we heterologously expressed intron-free *tnd*C in *Saccharomyces cerevisiae* ZXM144.[25] GC-MS analysis of crude extracts of *S. cerevisiae* ZXM144 transformed with *tnd*C compared to an empty vector control revealed the presence of a new product, **8**, with a *m/z* 272 [M]⁺ (Figures 2B and S7), supporting the production of a diterpene over a sesterterpene. HREIMS coupled 1D and 2D NMR spectroscopy (Figures S7 and S11-S15 and Table S2) identified that the planar structure of **8**, which was named talarodiene, contained a benzo[*a*]cyclopenta[*d*]cyclooctane tricyclic hydrocarbon backbone (Figure 2C). NOESY correlations were used to assign the relative configuration of **8** (Figures S16-S22), and ECD calculations (Figure S8) determined the absolute configuration to be (2*S*,3*S*,6*R*,11*R*).

With the isolation of **8**, the cyclization mechanism converting geranylgeranyl diphosphate (GGPP) into the 5-8-6 tricyclic hydrocarbon skeleton was investigated using ¹³C-labeling studies. [1-¹³C]acetate, [2-¹³C]acetate and [1,2-¹³C₂]acetate were administered independently to *tnd*C-transformed *S. cerevisiae* ZXM144, and the corresponding labeling patterns of ¹³C-enriched **8** were analyzed by NMR (Figures S23-S25 and Table S3). From the [1,2-¹³C₂]acetate labeling patterns, a cyclization mechanism is proposed in Figure 3. Cleavage of diphosphate followed by 1,11-10,14-cyclization converts GGPP to the bicyclic tertiary cation intermediate **8a⁺**. Ring expansion of **8a⁺** from a 1,2 alkyl shift forms the cation intermediate **8b⁺**, which is transformed to the tertiary cation intermediate **8c⁺** following a transannular proton transfer. A 1,2-hydride shift and 2,6-cyclization forms intermediate **8d⁺**, and deprotonation at C-8 yields **8**.

After heterologous expression of the cryptic *tnd*C gene led to the isolation of **8**, we turned back to the original host and evaluated *A. flavipes* CNL-338 for its production of this new tricyclic diterpene (Figure 4). Unfortunately, we were unable to detect the presence of **8** in crude extracts using GC-MS and LC-MS analyses, which suggests that **8** is an intermediate in the pathway and is modified by tailoring enzymes encoded in the *tnd* biosynthetic cluster. In addition to the diterpene synthase, the *tnd* cluster encodes several oxidative enzymes including the cytochrome P450 (*tnd*B), an aldehyde reductase (*tnd*E), and an alcohol dehydrogenase (*tnd*F) (Table S4). Given the type of tailoring enzymes present, we speculated that the cytochrome P450 TndB would be the next enzyme in the biosynthetic pathway. Indeed, GC-MS analysis of the Δ*tnd*B mutant showed the accumulation of **8** (Figure 4 and S10).

While the gene inactivation experiments unequivocally linked the *tnd* biosynthetic cluster to **8** in *A. flavipes* CNL-338, the final natural product(s) produced by the pathway were unknown. Recently, a group of diterpenoids, talaronoid A **9**, B **10**, and C **11**, containing a 5-8-6 fused ring system were isolated from *Talaromyces stipitatus* (Figure 3).[26] Using the amino acid sequence of TndC, we identified a gene cluster in *T. stipitatus* harboring an assortment of genes similar to those in the *tnd* biosynthetic cluster (Figures S3-S5). However, when we scanned crude extracts of *A. flavipes* CNL-338 for the presence of **9**, **10**, and **11**, the compounds were not detected. Based on these data, coupled with the Δ*tnd*B results, we speculated that low production

of **8** by *A. flavipes* CNL-338 could be impeding our ability to detect its conversion to the talaronoids.

Figure 4. GC-MS chromatograms (TIC) of (i) a standard of compound **8**, (ii) crude extract of Δ*tnd*B strain and (iii) crude extract from wild-type *A. flavipes* CNL-338.

To determine if **8** is indeed an intermediate in talaronoid biosynthesis, we biosynthetically prepared ¹³C-enriched **8** in *S. cerevisiae* using [1-¹³C]acetate. Labeled material was administered to *A. flavipes* CNL-338 and HREIMS inspection of the crude extract showed production of a new compound not observed in the DMSO control. The isotopic fragmentation pattern of the new compound also indicated it was derived from labeled material (Figure 5a). Closer inspection of the new compound showed that its retention time and *m/z* matched to that of an authentic standard of talarodiene C **11**, thereby confirming that **8** had been biotransformed to **11**. Thus, the stable tracer isotope experiment confirmed **8** as an intermediate in the talaronoid biosynthetic pathway and that TndB, TndE and TndF modify the 5-8-6 tricyclic hydrocarbon skeleton to afford talaronoid C.

Figure 5. *In vivo* conversion of the talarodiene backbone **8** in *A. flavipes* CNL-338. (**A**) High-resolution LC-MS analysis (EIC = 343 *m/z*) for the [M+Na]⁺ adduct of the talaronoid C standard **11** compared to *A. flavipes* CNL-338 administered [1-¹³C]acetate-labeled **8** or a DMSO control. All traces are shown in the same scale. (**B**) HREIMS fragmentation pattern of the talaronoid C standard compared to the product observed after administering [1-¹³C]acetate-labeled **8** to *A. flavipes* CNL-338.

In summary, we identified and characterized the *tnd* biosynthetic cluster responsible for the production of talaronoid C from the marine-derived fungus *A. flavipes* CNL-338. Heterologous expression of a cryptic type I bifunctional terpene synthase led to the discovery of a new diterpene possessing a benzo[*a*]cyclopenta[*d*]cyclooctane ring system and demonstrated that a single enzyme is responsible for the synthesis of this complex hydrocarbon scaffold. ^{13}C-labeling studies helped elucidate a possible cyclization mechanism that would convert geranylgeranyl diphosphate to the 5-8-6 tricyclic hydrocarbon skeleton, and stable tracer isotope experiments validated **8** as an intermediate in talaronoid biosynthesis. Our work therefore brought to light the product of a cryptic terpene synthase and information gleaned from the characterization of TndC can assist with genome mining predictions.

ASSOCIATED CONTENT

Supporting Information Available. Experimental details and NMR spectroscopic data. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

*Corresponding Author: Jaclyn.Winter@utah.edu

ORCID

Stephanie C. Heard: 0000-0003-4312-5105
Yonghui Zhang: 0000-0002-7222-2142
Jaclyn M. Winter: 0000-0001-6273-5377

Present Addresses

§ Jiangsu Co-Innovation Center of Efficient Processing and Utilization of Forest Resources and College of Chemical Engineering, Nanjing Forestry University, Nanjing 210037, China
‖ Jiangsu Key Lab of Biomass-Based Green Fuels and Chemicals, Nanjing, 220013, China

Author Contributions

P.Z., G. W., and J.M.W. designed the project; P.Z., G.W., S. C. H., C. N. and J. M. W. carried out the experiments; all authors analyzed data; and all authors wrote manuscript. ‡These authors contributed equally.

Funding Sources

This work was supported by an ALSAM and ARUP laboratory graduate research fellowship to S.C.H, the Gordon and Betty Moore Foundation (GBMF7621) to J.M.W. and in part by the US National Institutes of Health (1R01AI155694) to J.M.W.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We thank Drs William Fenical (Scripps Institution of Oceanography) for *A. flavipes* CNL-338, Joseph Chappell (University of Kentucky) for *S. cerevisiae* ZXM144, Jack Skalicky (University of Utah) for helpful NMR advice, and John Alan Maschek (University of Utah) for GCMS assistance.

ABBREVIATIONS

TC, terpene cyclase; PT, prenyltransferase; *tnd*, talaronoid biosynthetic cluster; HREIMS, high resolution electrospray ionization mass spectroscopy; GC-MS, gas chromatography mass spectroscopy; ECD, electronic circular dichroism; TIC, total ion chromatogram; EIC, extracted ion chromatogram; the GenBank Accession number for the *tnd* cluster is MW248390.

REFERENCES

(1) Gershenzon, J.; Dudareva, N. The function of terpene natural products in the natural world. *Nat. Chem. Biol.* **2007**, *3*, 408-414.
(2) Christianson, D. W. Structural and chemical biology of terpenoid cyclases. *Chem. Rev.* **2017**, *117*, 11570-11648.
(3) Faylo, J. L.; Ronnebaum, T. A.; Christianson, D. W. Assembly-Line catalysis in bifunctional terpene synthases. *Acc. Chem. Res.* **2021**, *54*, 3780-3791.
(4) Mitsuhashi, T.; Abe, I. Chimeric terpene synthases possessing both terpene cyclization and prenyltransfer activities. *Chembiochem.* **2018**, *19*, 1106-1114.
(5) Minami, A.; Ozaki, T.; Liu, C. W.; Oikawa, H. Cyclopentane-forming di/sesterterpene synthases: widely distributed enzymes in bacteria, fungi, and plants. *Nat. Prod. Rep.* **2018**, *35*, 1330-1346.
(6) Dickschat, J. S. Bacterial diterpene biosynthesis. *Angew. Chem. Int. Ed.* **2019**, *58,*15964-15976.
(7) Toyomasu, T.; Tsukahara, M.; Kaneko, A.; Niida, R.; Mitsuhashi, W.; Dairi, T.; Kato, N.; Sassa, T. Fusicoccins are biosynthesized by an unusual chimera diterpene synthase in fungi. *Proc. Natl. Acad. Sci. USA.* **2007**, *104*, 3084-3088.
(8) Chiba, R.; Minami, A.; Gomi, K.; Oikawa, H. Identification of Ophiobolin F synthase by a genome mining approach: a sesterterpene synthase from *Aspergillus clavatus. Org. Lett.* **2013**, *15*, 594-597.
(9) Okada, M.; Matsuda, Y.; Mitsuhashi, T.; Hoshino, S.; Mori, T.; Nakagawa, K.; Quan, Z. Y.; Qin, B.; Zhang, H. P.; Hayashi, F.; Kawaide, H.; Abe, I. Genome-based discovery of an unprecedented cyclization mode in fungal sesterterpenoid biosynthesis. *J. Am. Chem. Soc.* **2016**, *138*, 10011-10018.
(10) Ye, Y.; Minami, A.; Mandi, A.; Liu, C. W.; Taniguchi, T.; Kuzuyama, T.; Monde, K.; Gomi, K.; Oikawa, H. Genome mining for sesterterpenes using bifunctional terpene synthases reveals a unified intermediate of di/sesterterpenes. *J. Am. Chem. Soc.* **2015**, *137*, 11846-11853.
(11) Narita, K.; Sato, H.; Minami, A.; Kudo, K.; Gao, L.; Liu, C.; Ozaki, T.; Kodama, M.; Lei, X.; Taniguchi, T.; Monde, K.; Yamazaki, M.; Uchiyama, M.; Oikawa, H. Focused genome mining of structurally related sesterterpenes: enzymatic formation of enantiomeric and diastereomeric products. *Org. Lett.* **2017**, *19*, 6696-6699.
(12) Bian, G. K.; Han, Y. C.; Hou, A. W.; Yuan, Y. J.; Liu, X. H.; Deng, Z. X.; Liu, T. G. Releasing the potential power of terpene synthases by a robust precursor supply platform. *Metab. Eng.* **2017**, *42*, 1-8.
(13) Chen, R.; Jia, Q.; Mu, X.; Hu, B.; Sun, X.; Deng, Z.; Chen, F.; Bian, G.; Liu, T. Systematic mining of fungal chimeric terpene synthases using an efficient precursor-providing yeast chassis. *Proc. Natl. Acad. Sci. USA.* **2021**, *118*, e2023247118.
(14) Matsuda, Y.; Mitsuhashi, T.; Quan, Z.; Abe, I. Molecular basis for stellatic acid biosynthesis: a genome mining approach for discovery of sesterterpene synthases. *Org. Lett.* **2015**, *17*, 4644-4647.
(15) Guo, J.; Cai, Y.-S.; Cheng, F.; Yang, C.; Zhang, W.; Yu, W.; Yan, J.; Deng, Z.; Hong, K. Genome mining reveals a multiproduct sesterterpenoid biosynthetic gene cluster in *Aspergillus ustus. Org. Lett.* **2021**, *23*, 1525-1529.
(16) Yuan, Y.; Cheng, S.; Bian, G.; Yan, P.; Ma, Z.; Dai, W.; Chen, R.; Fu, S.; Huang, H.; Chi, H.; Cai, Y.; Deng, Z.; Liu, T. Efficient exploration of terpenoid biosynthetic gene clusters in filamentous fungi. *Nat. Cat.* **2022**, *5*, 277-287.

(17) Qin, B.; Matsuda, Y.; Mori, T.; Okada, M.; Quan, Z. Y.; Mitsuhashi, T.; Wakimoto, T.; Abe, I. An unusual chimeric diterpene synthase from *Emericella variecolor* and its functional conversion into a sesterterpene synthase by domain swapping. *Angew. Chem. Int. Ed.* **2016**, *55*, 1658-1661.

(18) Matsuda, Y.; Mitsuhashi, T.; Lee, S.; Hoshino, M.; Mori, T.; Okada, M.; Zhang, H.; Hayashi, F.; Fujita, M.; Abe, I. Astellifadiene: structure determination by NMR spectroscopy and crystalline sponge method, and elucidation of its biosynthesis. *Angew. Chem. Int. Ed.* **2016**, *55*, 5785-5788.

(19) Toyomasu, T.; Kaneko, A.; Tokiwano, T.; Kanno, Y.; Kanno, Y.; Niida, R.; Miura, S.; Nishioka, T.; Ikeda, C.; Mitsuhashi, W.; Dairi, T.; Kawano, T.; Oikawa, H.; Kato, N.; Sassa, T. Biosynthetic gene-based secondary metabolite screening: a new diterpene, methyl phomopsenonate, from the fungus *Phomopsis amygdali*. *J. Org. Chem.* **2009**, *74*, 1541-1548.

(20) Minami, A.; Tajima, N.; Higuchi, Y.; Toyomasu, T.; Sassa, T.; Kato, N.; Dairi, T. Identification and functional analysis of brassicicene C biosynthetic gene cluster in *Alternaria brassicicola*. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 870-874.

(21) Mitsuhashi, T.; Kikuchi, T.; Hoshino, S.; Ozeki, M.; Awakawa, T.; Shi, S. P.; Fujita, M.; Abe, I. Crystalline sponge method enabled the investigation of a prenyltransferase-terpene synthase chimeric enzyme, whose product exhibits broadened NMR signals. *Org. Lett.* **2018**, *20*, 5606-5609.

(22) Bian, G.; Rinkel, J.; Wang, Z.; Lauterbach, L.; Hou, A.; Yuan, Y.; Deng, Z.; Liu, T.; Dickschat, J. S. A clade II-D fungal chimeric diterpene synthase from *Colletotrichum gloeosporioides* produces dolasta-1(15),8-diene. *Angew. Chem. Int. Ed.* **2018**, *57*, 15887-15890.

(23) Fenical, W.; Jensen, P. R. Developing a new resource for drug discovery: marine actinomycete bacteria. *Nat. Chem. Biol.* **2006**, *2*, 666-673.

(24) Heard, S. C.; Wu, G. W.; Winter, J. M. Discovery and characterization of a cytochalasan biosynthetic cluster from the marine-derived fungus *Aspergillus flavipes* CNL-338. *J. Antibiot.* **2020**, *73*, 803-807.

(25) Zhuang, X.; Chappell, J. Building terpene production platforms in yeast. *Biotechnol. Bioeng.* **2015**, *112*, 1854-1864.

(26) Zhang, M.; Yan, S.; Liang, Y.; Zheng, M. J.; Wu, Z. D.; Zang, Y.; Yu, M. Y.; Sun, W. G.; Liu, J. J.; Ye, Y.; Wang, J. P.; Chen, C. M.; Zhu, H. C.; Zhang, Y. H. Talaronoids A-D: four fusicoccane diterpenoids with an unprecedented tricyclic 5/8/6 ring system from the fungus *Talaromyces stipitatus*. *Org. Chem. Front.* **2020**, *7*, 3486-3492.