A hippocampal-anterior thalamic circuit associated with learning

rate in ageing

Darya Frank*[1], Marta Garo-Pascual*[1,2,3], Pablo Alejandro Reyes Velasquez[1], Belén Frades[2],

Noam Peled[4,5], Linda Zhang[2], Bryan A. Strange[1,2]

[1]*Laboratory for Clinical Neuroscience, CTB, Universidad Politécnica de Madrid, 28223*
Madrid, Spain

[2]*Alzheimer's Disease Research Unit, CIEN Foundation, Queen Sofia Foundation Alzheimer*
Center, 28031 Madrid, Spain.

[3]*PhD Program in Neuroscience, Autonoma de Madrid University, 28049 Madrid, Spain*

[4]*Athinoula A. Martinos Center. for Biomedical Imaging, Department of Radiology,*
Massachusetts General Hospital, 02129 Charlestown, MA, USA

[5]*Harvard Medical School, 02115 Boston, MA, USA*

* Contributed equally to this work

Corresponding authors: Marta Garo-Pascual (marta.garo@ctb.upm.es), Darya Frank
(darya.frank@ctb.upm.es) and Bryan A. Strange (bryan.strange@upm.es).

Abstract

Memory normally declines with ageing and these age-related cognitive changes are

associated with changes in brain structure. Episodic memory retrieval has been widely

studied during ageing, whereas learning has received less attention. Here we examined the

neural correlates of learning rate in ageing. Our study sample consisted of 982 cognitively

healthy female and male older participants from the Vallecas Project cohort, without a

clinical diagnosis of mild cognitive impairment or dementia. The learning rate across three

consecutive verbal memory (Free and Cued Selective Reminding Test) recall trials was used

as a predictor of grey matter (GM) and white matter (WM) volume using voxel-based

morphometry, and WM microstructure using tract-based spatial statistics on fractional

anisotropy (FA) and mean diffusivity (MD) measures. Immediate recall improved by 1.4

items per trial on average, and this learning rate was faster in women and negatively

associated with age. Structurally, hippocampal and anterior thalamic GM volume correlated

positively with learning rate. Learning also correlated with the integrity of WM microstructure

(high FA and low MD) in an extensive network of tracts including bilateral anterior thalamic

radiation, fornix, and long-range tracts. These results suggest that structural GM and WM

characteristics, centred on a hippocampal-anterior thalamic circuit, support learning capacity

in ageing. Specifically, reduced volume and microstructure may explain some of the age-

related memory deficits that result from impaired learning.

Significance statement

A detailed understanding of age-related memory changes is crucial in an increasingly ageing

population, in which memory decline is prevalent. Whilst memory performance is usually

quantified by the ability to correctly retrieve information, memory impairment could be

caused by a reduced ability to learn or encode information. In a cohort of cognitively healthy

ageing participants, we found an association between verbal learning rate and grey matter

volume and white matter microstructure in the limbic system. These findings establish the

neural underpinnings of the often-overlooked learning phase of the memory process and

have important implications for our understanding of neurobiological changes in healthy

ageing and their cognitive phenotypes.

Introduction

Ageing is accompanied by a decline in cognition, most characteristically in memory

performance (Glisky, 2007). Memory impairments in ageing can manifest in different ways,

therefore providing insight into different structural and functional age-related changes.

Furthermore, exploring the neural underpinnings of these various manifestations (e.g.,

learning versus retention) could inform dissociations between normal age-related decline

and decline driven by neurodegenerative diseases such as dementia. Memory encapsulates

encoding, consolidation and retrieval processes, and deficits in either three may affect

performance independently. Memory decline in ageing is often measured in terms of

retention, however, impairments could also be driven by a diminished ability to learn

information, rather than to retrieve it. We therefore aimed to elucidate the structural brain

properties underlying learning rate in ageing.

Ageing has been associated with a global reduction in grey matter (GM) volume (Grieve et

al., 2007; Farokhian et al., 2017), although to different extents across brain regions (Cox et

al., 2018). Numerous studies have found a specific GM volume loss in prefrontal, temporal

and parietal cortices (Elliott, 2020; Cox et al., 2021), associated with general cognitive and

memory-specific decline (Gorbach et al., 2017; Cox et al., 2021). White matter (WM) age-

related changes in fractional anisotropy (FA) and mean diffusivity (MD), have also been

reported (Bennett et al., 2010; Madden and Parks, 2017). FA and MD are negatively

correlated such that reduced WM integrity is indexed by a decrease in FA and an increase in

MD. Like GM, age-related WM changes are apparent throughout the brain (Grieve et al.,

2007; Farokhian et al., 2017), although with greater effects in anterior than posterior tracts

(Bennett et al., 2010). Whilst these structural changes contribute to our understanding of

brain ageing, it is vital to also consider their cognitive correlates.

A commonly observed form of age-related cognitive decline is impaired memory, which has

been associated with reduced hippocampal volume (Persson et al., 2012; Hedden et al.,

2016; Gorbach et al., 2017), as well as with damage to the microstructure of frontal and

temporal WM tracts (de Mooij et al., 2018; Rizvi et al., 2020), and specifically limbic tracts

(Bennett et al., 2015). Furthermore, recognition performance on neuropsychological episodic

memory tests has been shown to correlate with FA and MD measures in the fornix,

cingulum, and superior and inferior longitudinal fasciculi (Sasson et al., 2013). However,

other studies have not found correlations between WM microstructure and episodic retrieval

in ageing (Salami et al., 2012; Laukka et al., 2013; Gorbach et al., 2017).

Learning, which is potentially dissociable from retrieval processes (Kwok and Buckley, 2010;

Bennett et al., 2015), is often overlooked in the context of ageing. Whilst memory

performance is usually quantified by the ability to correctly recognise or recall information, a

retrieval impairment could be caused by reduced ability to learn information. Indeed, there is

evidence for reduced error-driven (Nassar et al., 2016) and probabilistic learning (Samanez-

Larkin et al., 2012; Herff et al., 2020) in older adults, but evidence for similar deficits in

episodic learning is lacking. Relatedly, the microstructure of WM limbic tracts has been

related to episodic and navigational learning across species (Aggleton et al., 1995; Buckley

et al., 2008; Hofstetter et al., 2013; Hodgetts et al., 2020), but it is still unclear whether such

changes in these tracts can explain memory deficits in ageing.

We investigated whether the learning rate in the Free and Cued Selective Reminding Test

(FCSRT) is associated with age as well as its neural manifestation in GM and WM volume

and WM tract microstructure. In a large cross-sectional cohort of healthy older adults, we

first calculated the learning rate across three FCSRT trials (i.e., how many additional words

are successfully encoded in each trial) and tested for an association with age. To examine

brain-cognition associations, we used learning rate as a predictor to examine 1) GM and WM

volume using voxel-based morphometry (VBM), and 2) WM microstructure using tract-based

spatial statistics on FA and MD measures. Given the critical role of hippocampus in episodic

memory and correlation with structural changes in ageing, we hypothesised that

hippocampal GM volume, and associated WM limbic tract microstructure would correlate

with learning rate.

Methods

Participants. All participants in this study were part of the Vallecas Project, a single-centre

longitudinal study with annual visits. Inclusion and exclusion criteria for participants in this

project have been described elsewhere (Olazarán et al., 2015). From this cohort, data from

the baseline visit of 982 cognitively normal participants (mean age = 74.8, SD = 3.9, 637

(64.9%) females) were included in the current study. Any subject with a diagnosis of mild

cognitive impairment or Alzheimer's disease at this first visit was excluded. All participants

provided written informed consent and the Vallecas Project was approved by the Ethics

committee of the Instituto de Salud Carlos III.

Neuropsychological assessment. Participants completed a battery of neuropsychological

assessments as part of the Vallecas Project protocol. In this study we reported the total

score of the Mini Mental State Examination (MMSE; Folstein et al., 1975) and we mainly

focused on the Free and Cued Selective Reminding Test (FCSRT; Buschke, 1984),

assessing learning and retention of verbal memory, with immediate and delayed recall

components. The test was administered using standard procedures (Peña-Casanova et al.,

2009). During the study phase, participants were shown a card with four words and asked to

identify the word corresponding to a specific category, cycling through all four words, on four

different cards (16 words in total). Following the study phase, three recall trials were used

for the learned items (immediate recall), each one followed by 20 seconds of interference

counting backwards. The same procedure was done after a 30-minute delay (delayed

recall). To assess the learning rate across trials, we fit a linear-mixed effects model of the

number of items recalled in each trial, as a function of the recall trial (first, second, and third)

using the *lme4* package in R 4.0.2 (https://www.r-project.org/). The model also included a

random slope of the recall trial, and a random intercept per participant, capturing inter-

individual variability in learning rate (across the three trials). The learning rate coefficient for

each participant was extracted for subsequent analyses. Next we built a multiple regression

model where the learning rate was the dependent variable, sex, age and level of education

were the predictors, and delayed free recall score of the FCSRT was included as a covariate

to rule out the retrieval phase of the memory process.

MRI Data acquisition. Images were acquired using a 3T MRI (Signa HDxt GE) with a phased

array eight-channel head coil. T1-weighted images (3D fast spoiled gradient echo with

inversion recovery preparation) were collected using a repetition time (TR) of 10ms, echo

time (TE) of 4.5ms, field of view (FOV) of 240mm and a matrix size of 288x288 with slice

thickness of 1mm, yielding a voxel size of 0.5 x 0.5 x 1 mm^3. Diffusion-weighted images

were single-shot spin echo echo-planar imaging (SE-EPI), with TR 9200ms, TE 80ms, b-

value 800s/mm^2 and 21 gradient directions, FOV 240mm and matrix size 128 x 128 with

slice thickness of 3mm.

Grey and white matter VBM. The analysis was carried out in SPM12 (version r6225;

https://www.fil.ion.ucl.ac.uk/spm). T1-weighted images were segmented into grey matter,

white matter and cerebrospinal fluid and then aligned and normalised to MNI space using

the DARTEL algorithm (Ashburner, 2007). Prior to statistical modelling, the normalised

images were smoothed using a 6mm FWHM Gaussian kernel. The pre-processed grey and

white matter maps were entered into separate general linear models (GLM) with learning

rate from the memory task as the predictor of interest, and total intracranial volume (TIV),

sex, and the delayed free recall score of the FCSRT as covariates. Age and education were

not used in the model as additional covariates, because they are correlated with FCSRT

delayed free recall so the effects of age and level of education are already accounted for by

FCSRT delayed free recall. We conducted whole-brain analyses using a threshold-free

cluster enhancement (TFCE) approach with 5000 permutations and default parameters (E =

0.5 and H = 2) using the TFCE tool (version r223) for CAT12 toolbox in SPM

(http://dbm.neuro.uni-jena.de/tfce). The AAL3 atlas neuroanatomical labels were used to

describe neuroanatomical loci (Rolls et al., 2020) and Mango software was used to produce

the figure (http://rii.uthscsa.edu/mango/). These analyses assessed which regions were

positively associated with the immediate recall learning slope. Significant results are

reported at a family-wise error (FWE) corrected level of $p < 0.05$.

White matter tract-based spatial statistics (TBSS). Of the 982 participants, seven were

excluded as they did not have diffusion data. For preprocessing these images, the FSL

toolbox (http://fsl.fmrib.ox.ac.uk/fsl/fslwiki) was used for motion and eddy current correction,

the extraction of non-brain voxels and, lastly, the calculation of voxel-wise diffusion maps

(FA and MD) for each participant. Individual FA and MD maps were then used in the FSL

TBSS pipeline (http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/TBSS/UserGuider), detailed methods

described by Smith et al. (2006). The general outline of the process is 1) FA individual maps

were non-linearly registered to standard space (FMRIB58_FA template) (Andersson et al.,

2007); 2) a mean FA image was created by averaging all co-registered FA maps; and 3)

individually aligned images were projected onto the mean FA skeleton—representing the

centers of all tracts common to the study sample—and skeletonized images were used for

voxel-wise analysis. Diffusivity maps for MD were generated by applying the same steps

detailed above. The same GLM design matrix as the VBM analysis was used along with the

TFCE approach with 5000 permutations (default parameters E = 0.5 and H = 2). Significant

results are reported at a family-wise error (FWE) corrected level of $p < 0.05$. To visualise our

TBSS results we used the multimodal analysis and visualisation tool (MMVT; Felsenstein et

al., 2019). The pipeline follows these steps: 1) Binary masking: all the voxels in the TBSS

volume below the threshold (0.95) were set to zero. 2) Outlier voxels removal using the

Open3D python package (Zhou et al., 2018). 3) Smoothing the volumetric data using a 3D

Gaussian filter (Virtanen et al., 2020). 4) Surface creation from the volume's TBSS surfaces

using the marching cubes algorithm (Lorensen and Cline, 1987). For that, we re-calculate

the threshold to give us the same number of voxels after the smoothing step. 5) Translation

for the surfaces' vertices coordinates. 6) Projection of the volumetric data on the surfaces.

Results

Memory and neuropsychological performance

On average, across three trials, participants correctly remembered 7.9 items (SD = 2.6).

When looking at individual trials, performance improved as trials progressed, reflecting a

positive learning slope (see **Table 1** for number of items recalled, and **Figure 1A** for learning

rates). Our linear model predicting the learning rate as a function of age, sex, and level of

education revealed significant effects of the three predictors after correcting for FCSRT

delayed free recall score. Learning rate and delayed free recall FCSRT are positively

correlated (Pearson's r = 0.7; $p < 2.2 \times 10^{-16}$). Age has a negative effect on learning rate

($F(1,965) = 10.45$, $p = 0.001$) (**Figure 1B**), sex also has an effect ($F(1,965) = 4.66$, $p =$

0.031) and being a woman is positively associated with learning rate (**Figure 1C**). Finally,

having a higher level of education is positively associated with learning rate ($F(3,965) = 5.17$

, $p < 0.002$) (**Figure 1D**). There is no significant interaction between the three predictors

(age, sex and years of education).

	Total sample (n = 982) Mean (SD)	**Women** (n = 637) Mean (SD)	**Men** (n = 345) Mean (SD)
Demographics			
Age, years	74.8 (3.9)	74.8 (3.9)	74.8 (3.9)

Levels of education, count (%)	Less than primary	185 (18.8)	137 (21.5)	48 (13.9)
	Primary	295 (30.0)	222 (34.9)	73 (21.1)
	Secondary	245 (24.9)	151 (23.7)	94 (27.2)
	Higher education	257 (26.2)	127 (19.9)	130 (37.7)
Neuropsychological performance				
MMSE, total score		28.6 (1.6)	28.6 (1.6)	28.7 (1.4)
Trial 1 immediate free recall FCSRT, items recalled		6.5 (2.1)	6.7 (2.1)	6.3 (2.1)
Trial 2 immediate free recall FCSRT, items recalled		7.9 (2.4)	7.9 (2.4)	7.8 (2.4)
Trial 3 immediate free recall FCSRT, items recalled		9.2 (2.5)	9.4 (2.6)	9.0 (2.6)
Learning rate immediate free recall FCSRT, items recalled/trial		1.4 (0.3)	1.4 (0.3)	1.3 (0.3)
Delayed free recall FCSRT, items recalled		9.4 (2.6)	9.5 (2.7)	9.3 (2.6)

Table 1. Demographic and neuropsychological profile of the total sample and split by

sex. MMSE: Mini Mental State Examination total score, FCSRT: Free and Cued Selective

Reminding Test.

Figure 1. Learning rates across FCSRT trials are related to age, sex and level of

education. A Mean and individual learning rates across the three free immediate recall

FCSRT trials. After controlling for the rest of the model predictors, **B** learning rate decreases

with age, **C** females learn faster than males and **D** level of education is positively associated

with learning rate. Error bars represent 95% confidence intervals.

Grey and white matter volume (VBM)

We found a positive correlation between learning slope and grey matter volume in the

bilateral hippocampus, with more pronounced effects on the left side (**Figure 2A**), and the

right anterior thalamic nucleus with some extension to adjacent nuclei (right ventroanterior

and ventrolateral thalamic nuclei) (**Figure 2B**, **Table 2-1**). There were no significant

associations between learning rate and white matter volume (data not shown).

Figure 2. Grey matter volume of hippocampal-anterior thalamic circuit correlates with learning rate in the older adults. The positive correlation has been overlaid on a canonical T1 image (thresholded at *p* < 0.05 FWE-corr) to show a significant effect in **A** hippocampus bilaterally and **B** right anterior (red), ventroanterior (green) and ventrolateral thalamic nuclei (orange) (thalamic ROIs in the inset come from the AAL3 atlas (Rolls et al., 2020). The coordinates of the sections are given in mm. L: left, R: right, A: anterior, P: posterior.

White matter microstructure (TBSS)

We first examined FA as a marker of WM integrity. We found a bilateral network of temporal,

parietal and occipital tracts showing a positive association with learning rate. Among tracts

showing significant positive correlations were the bilateral anterior thalamic radiation (ATR),

fornix, inferior fronto-occipital fasciculus (IFOF), inferior longitudinal fasciculus (ILF), superior

longitudinal fasciculus (SLF), and uncinate (**Figure 3A-B, Table 3-1**). Next, we examined

MD and found a negative association between a similar network of bilateral tracts and

FCSRT learning rate, including bilateral ATR, corticospinal tract, forceps major and minor,

cingulum (cingulate), IFOF, ILF, SLF, uncinate and fornix (**Figure 3C-D, Table 3-1**).

Figure 3. Extensive network of white matter microstructure integrity is related to

learning rate in the older adults. A. Positive correlation between FA and learning rate

(warm colours; $p < 0.05$ FWE-corr). **B.** FA effects overlaid on the fornix. **C.** Negative

correlation between MD and learning rate (cold colours; $p < 0.05$ FWE-corr). **D.** MD effect

overlaid on the fornix.

Discussion

Learning rate is faster in women, declines with age and is associated with neuroanatomical

structural properties. We found a positive correlation between GM volume and learning rate,

where participants with greater hippocampal and anterior thalamic nucleus volume learned

at a faster rate than those with reduced volume. Furthermore, we found that FA was

positively associated with learning rate in an extended network including limbic tracts,

indicating that structural integrity of these tracts indexed learning ability. A complementary

negative association was observed for MD, in similar tracts, such that increased MD was

associated with a slower learning rate. These findings suggest that structural properties of

hippocampal-anterior thalamic circuit contribute to learning ability in ageing. Specifically,

reduction in volume and WM microstructure may explain some age-related memory deficits

stemming from slower learning, as opposed to deficits in consolidation or retrieval.

Previous research on neural substrates of cognitive decline in ageing has shown a

hippocampal volume deterioration with age that correlated with memory performance

(Persson et al., 2012; Hedden et al., 2016; Gorbach et al., 2017). The more pronounced

effect we observed in the left hippocampus is in accordance with previous VBM findings of

verbal memory tasks (Ezzati et al., 2016), and the general lateralisation of verbal functions.

Our results therefore extend previous research on the relationship between hippocampal

volume and memory decline in ageing, showing learning ability is also indexed by

hippocampal volume. Note that it is unlikely that these effects reflect memory function in

general, given that delayed free recall was included as a covariate in our model.

Structural properties of extra-hippocampal limbic regions were also associated with learning

ability. Our GM thalamic findings indicate a correlation between learning rate and the right

anterior thalamic nucleus, extending to right ventroanterior and right ventrolateral nuclei. The

anterior thalamic nuclei have been suggested to play an important role in learning and

memory (Winocur, 1985; Aggleton et al., 2010), beyond navigation (Wolff and Vann, 2019;

Nelson, 2021). Specifically, and in line with our results, Leszczyński and Staudigl (2016)

posited that the anterior thalamus might modulate information flow, via attention allocation,

to support learning. Taken together with increased hippocampal volume related to better

learning rate, our results indicate that the limbic system plays an important role in learning

ability in ageing, and might explain some of the impairments in navigating in a novel

environment (Grzeschik et al., 2021), and impaired learning strategies observed in MCI

(Ribeiro et al., 2007)The fornix is a major hippocampal input/output pathway and has been

associated with visuo-spatial learning across species (Buckley et al., 2008; Hofstetter et al.,

2013; Hodgetts et al., 2020). The fornix links the hippocampus with the anterior thalamic

nuclei (via mammillary bodies), with both regions showing grey matter volume relationship

with learning rate. Furthermore, we found that fornix integrity, as captured by bilateral FA

and MD, correlated with learning rate in older adults. This finding suggests that the fornix

also plays a role in verbal episodic learning, as captured in our study. We also found that the

WM integrity of the ATR was correlated with learning rate. The ATR, which connects the

anterior and dorsomedial thalamus with the prefrontal cortex (Grodd et al., 2020), may play a

role in learning (McGuire et al., 2014).

In addition to changes in limbic GM volume and WM microstructure, we found learning rate

was associated with broader changes within bilateral WM tracts connecting occipital-

temporal-frontal regions (ILF, SLF, IFOF). This result might point toward an overall WM

microstructure effect, as previously noted in age-related cognitive decline (Grieve et al.,

2007; Farokhian et al., 2017; de Mooij et al., 2018; Rizvi et al., 2020; Molloy et al., 2021).

Importantly, given that we control for age differences by inclusion of the delayed free recall

score as a covariate in the regression model, this result cannot be attributed to age

differences in our cross-sectional cohort. With respect to specific cognitive functions, ILF and

SLF have been shown to relate to memory performance in normal ageing (Sasson et al.,

2013). ILF and IFOF facilitate flow of visual information up the visual stream (Rokem et al.,

2017), and IFOF has been associated with semantic processing (Duffau, 2008), potentially

supporting learning performance in our task. Therefore, the observed relationship between

microstructure of these tracts and learning ability might reflect a more general aspect of

cognitive ability. Whilst we found significant correlations between learning rate and FA and

MD, no correlations were found for WM volume, potentially due to reduced sensitivity of WM

volume to age-related decline (Wozniak and Lim, 2006).

Finally, it is important to note some limitations of the current study; we analysed data from a

cross-sectional cohort of healthy older adults. As GM and WM properties and memory

function both deteriorate with age, future longitudinal studies would be needed to better

understand the relationship between learning ability and structural changes as ageing

progresses and eliminate age-related confounds in cross-sectional studies (Elliott, 2020).

We used the learning rate across trials in an established neuropsychological memory task

(FCSRT) as a measure for episodic learning; it would be interesting to examine neural

correlates of other forms of learning such as error-driven and statistical learning (Samanez-

Larkin et al., 2012; Nassar et al., 2016; Herff et al., 2020), as well as consider learning ability

as a potential cognitive phenotype in pathological ageing.

In conclusion, in a cross-sectional cohort of healthy older adults, we found learning rate on

the FCSRT task was positively associated with GM and WM structural changes in the limbic

system, including the hippocampus, fornix and anterior thalamic nucleus. Furthermore, there

was a positive correlation between learning rate and long-range WM tracts (ILF, SLF, IFOF).

Our findings suggest that some age-related memory deficits might be caused due to

impaired learning underpinned by such structural change.

References

Aggleton JP, Neave N, Nagle S, Hunt PR (1995) A comparison of the effects of anterior

thalamic, mamillary body and fornix lesions on reinforced spatial alternation. Behav

Brain Res 68:91–101.

Aggleton JP, O'Mara SM, Vann SD, Wright NF, Tsanov M, Erichsen JT (2010)

Hippocampal–anterior thalamic pathways for memory: uncovering a network of direct

and indirect actions. Eur J Neurosci 31:2292–2307 Available at:

https://onlinelibrary.wiley.com/doi/full/10.1111/j.1460-9568.2010.07251.x [Accessed

March 7, 2022].

Ashburner J (2007) A fast diffeomorphic image registration algorithm. Neuroimage 38:95–

113 Available at: http://www.ncbi.nlm.nih.gov/pubmed/17761438 [Accessed December

27, 2018].

Bennett IJ, Huffman DJ, Stark CEL (2015) Limbic tract integrity contributes to pattern

separation performance across the lifespan. Cereb Cortex 25:2988–2999.

Bennett IJ, Madden DJ, Vaidya CJ, Howard D V., Howard JH (2010) Age-related differences

in multiple measures of white matter integrity: A diffusion tensor imaging study of

healthy aging. Hum Brain Mapp 31:378–390.

Buckley MJ, Wilson CRE, Gaffan D (2008) Fornix Transection Impairs Visuospatial Memory

Acquisition More Than Retrieval. Behav Neurosci 122:44–53.

Buschke H (1984) Cued recall in amnesia. J Clin Neuropsychol 6:433–440 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/6501581 [Accessed January 16, 2019].

Cox SR, Bastin ME, Ritchie SJ, Dickie DA, Liewald DC, Muñoz Maniega S, Redmond P,

Royle NA, Pattie A, Valdés Hernández M, Corley J, Aribisala BS, McIntosh AM,

Wardlaw JM, Deary IJ (2018) Brain cortical characteristics of lifetime cognitive ageing.

Brain Struct Funct 223:509–518.

Cox SR, Harris MA, Ritchie SJ, Buchanan CR, Valdés Hernández MC, Corley J, Taylor AM,

Madole JW, Harris SE, Whalley HC, McIntosh AM, Russ TC, Bastin ME, Wardlaw JM,

Deary IJ, Tucker-Drob EM (2021) Three major dimensions of human brain cortical

ageing in relation to cognitive decline across the eighth decade of life. Mol Psychiatry

26:2651–2662 Available at: http://dx.doi.org/10.1038/s41380-020-00975-1.

de Mooij SMM, Henson RNA, Waldorp LJ, Kievit RA (2018) Age differentiation within gray

matter, white matter, and between memory and white matter in an adult life span

cohort. J Neurosci 38:5826–5836.

Duffau H (2008) The anatomo-functional connectivity of language revisited. New insights

provided by electrostimulation and tractography. Neuropsychologia 46:927–934.

Elliott ML (2020) MRI-based biomarkers of accelerated aging and dementia risk in midlife:

how close are we? Ageing Res Rev 61:101075 Available at:

https://doi.org/10.1016/j.arr.2020.101075.

Ezzati A, Katz MJ, Zammit AR, Lipton ML, Zimmerman ME, Sliwinski MJ, Lipton RB (2016)

Differential association of left and right hippocampal volumes with verbal episodic and

spatial memory in older adults. Neuropsychologia 93:380–385.

Farokhian F, Yang C, Beheshti I, Matsuda H, Wu S (2017) Age-related gray and white

matter changes in normal adult brains. Aging Dis 8:899–909.

Felsenstein O, Peled N, Hahn E, Rockhill AP, Folsom L, Gholipour T, Macadams K,

Rozengard N, Paulk AC, Dougherty D, Cash SS, Widge AS, Hämäläinen M,

Stufflebeam S (2019) Multi-Modal Neuroimaging Analysis and Visualization Tool

(MMVT). Available at: http://arxiv.org/abs/1912.10079.

Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". J Psychiatr Res 12:189–

198 Available at: https://linkinghub.elsevier.com/retrieve/pii/0022395675900266.

Glisky EL (2007) Changes in Cognitive Function in Human Aging.

Gorbach T, Pudas S, Lundquist A, Orädd G, Josefsson M, Salami A, de Luna X, Nyberg L
(2017) Longitudinal association between hippocampus atrophy and episodic-memory
decline. Neurobiol Aging 51:167–176 Available at:
https://doi.org/10.1016/j.neurobiolaging.2016.12.002.

Grieve SM, Williams LM, Paul RH, Clark CR, Gordon E (2007) Cognitive aging, executive
function, and fractional anisotropy: A diffusion tensor MR imaging study. Am J
Neuroradiol 28:226–235.

Grodd W, Kumar VJ, Schüz A, Lindig T, Scheffler K (2020) The anterior and medial thalamic
nuclei and the human limbic system: tracing the structural connectivity using diffusion-
weighted imaging. Sci Rep 10:1–25 Available at: https://doi.org/10.1038/s41598-020-
67770-4.

Grzeschik R, Hilton C, Dalton RC, Konovalova I, Cotterill E, Innes A, Wiener JM (2021) From
repeating routes to planning novel routes: the impact of landmarks and ageing on route
integration and cognitive mapping. Psychol Res 85:2164–2176 Available at:
https://link.springer.com/10.1007/s00426-020-01401-5.

Hedden T, Schultz AP, Rieckmann A, Mormino EC, Johnson KA, Sperling RA, Buckner RL
(2016) Multiple Brain Markers are Linked to Age-Related Variation in Cognition. Cereb
Cortex 26:1388–1400.

Herff SA, Zhen S, Yu R, Agres KR (2020) Age-dependent statistical learning trajectories
reveal differences in information weighting. Psychol Aging 35:1090–1104 Available at:
http://doi.apa.org/getdoi.cfm?doi=10.1037/pag0000567.

Hodgetts CJ, Stefani M, Williams AN, Kolarik BS, Yonelinas AP, Ekstrom AD, Lawrence AD,

Zhang J, Graham KS (2020) The role of the fornix in human navigational learning.

Cortex 124:97–110.

Hofstetter S, Tavor I, Moryosef ST, Assaf Y (2013) Short-term learning induces white matter

plasticity in the fornix. J Neurosci 33:12844–12850.

Kwok SC, Buckley MJ (2010) Long-term visuospatial retention unaffected by fornix

transection. Hippocampus 20:889–893.

Laukka EJ, Lövdén M, Kalpouzos G, Li TQ, Jonsson T, Wahlund LO, Fratiglioni L, Bäckman

L (2013) Associations between white matter microstructure and cognitive performance

in old and very old age. PLoS One 8:1–8.

Leszczyński M, Staudigl T (2016) Memory-guided attention in the anterior thalamus.

Neurosci Biobehav Rev 66:163–165.

Lorensen WE, Cline HE (1987) Marching cubes: A high resolution 3D surface construction

algorithm. In: Proceedings of the 14th annual conference on Computer graphics and

interactive techniques - SIGGRAPH '87, pp 163–169. New York, New York, USA: ACM

Press. Available at: http://portal.acm.org/citation.cfm?doid=37401.37422.

Madden DJ, Parks EL (2017) Age differences in structural connectivity: Diffusion tensor

imaging and white matter hyperintensitie. In: Cognitive neuroscience of aging: Linking

cognitive and cerebral aging, 2nd ed. (Cabeza R, Nyberg L, Park DC, eds), pp 71–103.

New York: Oxford.

McGuire JT, Nassar MR, Gold JI, Kable JW (2014) Functionally Dissociable Influences on

Learning Rate in a Dynamic Environment. Neuron 84:870–881 Available at:

http://dx.doi.org/10.1016/j.neuron.2014.10.013.

Molloy CJ, Nugent S, Bokde ALW (2021) Alterations in diffusion measures of white matter

integrity associated with healthy aging. Journals Gerontol - Ser A Biol Sci Med Sci

76:945–954.

Nassar MR, Bruckner R, Gold JI, Li SC, Heekeren HR, Eppinger B (2016) Age differences in

learning emerge from an insufficient representation of uncertainty in older adults. Nat

Commun 7:1–13 Available at: http://dx.doi.org/10.1038/ncomms11609.

Nelson AJD (2021) The anterior thalamic nuclei and cognition: A role beyond space?

Neurosci Biobehav Rev 126:1–11 Available at:

https://doi.org/10.1016/j.neubiorev.2021.02.047.

Olazarán J, Valentí M, Belén Frades, Zea-Sevilla MA, Ávila-Villanueva M, Fernández-

Blázquez MÁ, Calero M, Dobato JL, Hernández-Tamames JA, León-Salas B, Agüera-

Ortiz L, López-Álvarez J, Larrañaga P, Bielza C, Álvarez-Linera J, Martínez-Martín P

(2015) The Vallecas Project: A cohort to identify early markers and mechanisms of

Alzheimer's disease. Front Aging Neurosci 7 Available at:

https://pubmed.ncbi.nlm.nih.gov/26483681/ [Accessed January 13, 2022].

Peña-Casanova J, Gramunt-Fombuena N, Quiñones-Úbeda S, Sánchez-Benavides G,

Aguilar M, Badenes D, Molinuevo JL, Robles A, Barquero MS, Payno M, Antúnez C,

Martínez-Parra C, Frank-García A, Fernández M, Alfonso V, Sol JM, Blesa R (2009)

Spanish Multicenter Normative Studies (NEURONORMA Project): norms for the Rey-

Osterrieth complex figure (copy and memory), and free and cued selective reminding

test. Arch Clin Neuropsychol 24:371–393 Available at:

https://pubmed.ncbi.nlm.nih.gov/19661107/ [Accessed January 20, 2022].

Persson J, Pudas S, Lind J, Kauppi K, Nilsson L-G, Nyberg L (2012) Longitudinal Structure-

Function Correlates in Elderly Reveal MTL Dysfunction with Cognitive Decline. Cereb

Cortex 22:2297–2304.

Ribeiro F, Guerreiro M, De Mendonça A (2007) Verbal learning and memory deficits in Mild

Cognitive Impairment. J Clin Exp Neuropsychol 29:187–197 Available at:

https://www.tandfonline.com/doi/full/10.1080/13803390600629775.

Rizvi B, Lao PJ, Colón J, Hale C, Igwe KC, Narkhede A, Budge M, Manly JJ, Schupf N,

Brickman AM (2020) Tract-defined regional white matter hyperintensities and memory.

NeuroImage Clin 25:102143.

Rokem A, Takemura H, Bock AS, Scherf KS, Behrmann M, Wandell BA, Fine I, Bridge H,

Pestilli F (2017) The visual white matter: The application of diffusion MRI and fiber

tractography to vision science. J Vis 17:4.

Rolls ET, Huang CC, Lin CP, Feng J, Joliot M (2020) Automated anatomical labelling atlas 3.

Neuroimage 206:116189.

Salami A, Eriksson J, Nilsson LG, Nyberg L (2012) Age-related white matter microstructural

differences partly mediate age-related decline in processing speed but not cognition.

Biochim Biophys Acta - Mol Basis Dis 1822:408–415 Available at:

http://dx.doi.org/10.1016/j.bbadis.2011.09.001.

Samanez-Larkin GR, Levens SM, Perry LM, Dougherty RF, Knutson B (2012) Frontostriatal

white matter integrity mediates adult age differences in probabilistic reward learning. J

Neurosci 32:5333–5337.

Sasson E, Doniger GM, Pasternak O, Tarrasch R, Assaf Y (2013) White matter correlates of

cognitive domains in normal aging with diffusion tensor imaging. Front Neurosci.

Virtanen P et al. (2020) SciPy 1.0: fundamental algorithms for scientific computing in Python.

Nat Methods 17:261–272 Available at: http://www.nature.com/articles/s41592-019-

0686-2.

Winocur G (1985) THE HIPPOCAMPUS AND THALAMUS: THEIR ROLES IN SHORT-AND

LONG-TERM MEMORY AND THE EFFECTS OF INTERFERENCE.

Wolff M, Vann SD (2019) The cognitive thalamus as a gateway to mental representations. J

Neurosci 39:3–14.

Wozniak JR, Lim KO (2006) Advances in white matter imaging: A review of in vivo magnetic

resonance methodologies and their applicability to the study of development and aging.

Neurosci Biobehav Rev 30:762 Available at: /pmc/articles/PMC2895765/ [Accessed

January 25, 2022].

Zhou Q-Y, Park J, Koltun V (2018) Open3D: A Modern Library for 3D Data Processing.

Available at: http://arxiv.org/abs/1801.09847.