Brain Structure and Episodic Learning Rate in Cognitively Healthy Ageing

Darya Frank*[1], Marta Garo-Pascual*[1,2,3], Pablo Alejandro Reyes Velasquez[1], Belén Frades2,
Noam Peled[4,5], Linda Zhang[2], Bryan A. Strange[1,2]

[1]Laboratory for Clinical Neuroscience, CTB, Universidad Politécnica de Madrid, 28223
Madrid, Spain

[2]Alzheimer's Disease Research Unit, CIEN Foundation, Queen Sofia Foundation Alzheimer
Center, 28031 Madrid, Spain.

[3]PhD Program in Neuroscience, Autonoma de Madrid University, 28049 Madrid, Spain

[4]Athinoula A. Martinos Center. for Biomedical Imaging, Department of Radiology,
Massachusetts General Hospital, 02129 Charlestown, MA, USA

[5]Harvard Medical School, 02115 Boston, MA, USA

* Contributed equally to this work

Corresponding authors: Marta Garo-Pascual (marta.garo@ctb.upm.es), Darya Frank
(darya.frank@ctb.upm.es) and Bryan A. Strange (bryan.strange@upm.es).

<u>Abstract</u>

Memory normally declines with ageing and these age-related cognitive changes are

associated with changes in brain structure. Episodic memory retrieval has been widely

studied during ageing, whereas learning has received less attention. Here we examined the

neural correlates of episodic learning rate in ageing. Our study sample consisted of 982

cognitively healthy female and male older participants from the Vallecas Project cohort,

without a clinical diagnosis of mild cognitive impairment or dementia. The learning rate

across the three consecutive recall trials of the verbal memory task (Free and Cued

Selective Reminding Test) recall trials was used as a predictor of grey matter (GM) using

voxel-based morphometry, and WM microstructure using tract-based spatial statistics on

fractional anisotropy (FA) and mean diffusivity (MD) measures. Immediate Recall improved

by 1.4 items per trial on average, and this episodic learning rate was faster in women and

negatively associated with age. Structurally, hippocampal and anterior thalamic GM volume

correlated positively with learning rate. Learning also correlated with the integrity of WM

microstructure (high FA and low MD) in an extensive network of tracts including bilateral

anterior thalamic radiation, fornix, and long-range tracts. These results suggest that episodic

learning rate is associated with key anatomical structures for memory functioning, motivating

further exploration of the differential diagnostic properties between episodic learning rate and

retrieval in ageing.

Abstract word count: 219

Introduction

Ageing is accompanied by a decline in cognition, most characteristically in episodic memory performance (Glisky, 2007; Tromp et al., 2015), the ability to remember personal experiences. Episodic memory impairments in ageing can manifest in different ways depending on the studied phase (i.e., encoding, consolidation, or retrieval process). It is difficult to study these phases independently in behavioural studies, although previous work has reported that distinct processes may be affected unequally during ageing. For example, more prominent deficits have been found for encoding relative to retrieval in older adults (Friedman et al., 2007; Morcom et al., 2003). Furthermore, exploring the neural underpinnings of these various manifestations (e.g., learning versus retention) could inform dissociations between normal age-related decline and decline driven by neurodegenerative diseases such as dementia. Memory decline in ageing is often measured in terms of retention. However, impairments could also be driven by a diminished ability to learn information over a period of time, rather than to retrieve it. We therefore aimed to elucidate the structural brain properties underlying episodic learning rate in ageing.

Ageing has been associated with a global reduction in grey matter (GM) volume (Farokhian et al., 2017; Grieve et al., 2007), although to different extents across brain regions (Cox et al., 2018; Resnick et al., 2003). Numerous studies have found a specific GM volume loss in prefrontal, temporal and parietal cortices (Cox et al., 2021; Elliott, 2020), associated with general cognitive and memory-specific decline (Cox et al., 2021; Fjell and Walhovd, 2010; Gorbach et al., 2017). White matter (WM) age-related differences in fractional anisotropy (FA) and mean diffusivity (MD) have also been reported (Bennett et al., 2010; Fjell and Walhovd, 2010; Madden and Parks, 2017). FA and MD are negatively correlated such that reduced WM integrity is indexed by a decrease in FA and an increase in MD. Like GM, age-related WM effects are apparent throughout the brain (Farokhian et al., 2017; Grieve et al., 2007), although with greater effects in anterior than posterior tracts (Bennett et al., 2010).

Whilst these structural differences contribute to our understanding of brain ageing, it is vital

to also consider their cognitive manifestations.

A commonly observed form of age-related cognitive decline is impaired memory, which has

been associated with reduced hippocampal volume (Gorbach et al., 2017; Hedden et al.,

2016; Persson et al., 2012), as well as with damage to the microstructure of frontal and

temporal WM tracts (de Mooij et al., 2018; Kennedy and Raz, 2009; Rizvi et al., 2020), and

specifically limbic tracts (Bennett et al., 2015). Furthermore, recognition performance on

neuropsychological episodic memory tests has been shown to correlate with FA and MD

measures in the fornix, cingulum, and superior and inferior longitudinal fasciculi (Sasson et

al., 2013). However, other studies have not found correlations between WM microstructure

and episodic retrieval in ageing (Gorbach et al., 2017; Laukka et al., 2013; Salami et al.,

2012).

Memory performance is usually quantified by the ability to recognise or recall information

correctly, a retrieval impairment could be caused by a reduced ability to encode or learn

information (Boujut and Clarys, 2016; Cadar et al., 2018). Encoding, which is potentially

dissociable from retrieval processes (Bennett et al., 2015; Kwok and Buckley, 2010), has

been shown to underlie several memory deficits observed in ageing (Grady, 2012). Whilst

learning rate is part of the encoding process, in the current context it specifically refers to an

improvement in learning over time (or repetitions). Indeed, there is evidence for reduced

error-driven (Nassar et al., 2016) and probabilistic learning rates (Herff et al., 2019;

Samanez-Larkin et al., 2012) in older adults, but evidence for similar deficits in episodic

learning rate is lacking.

A potential way to probe episodic learning rate is through the Free and Cued Selective

Reminding Test (FCSRT). The FCSRT is one of the most commonly used free-recall

paradigms for episodic memory assessment, including immediate and delayed free- and

cued-recall (Buschke, 1984). Worse recall performance of cognitively normal older adults on

the FCSRT has been associated with reduced hippocampal GM volume (Zammit et al.,

2017), reduced fornix FA (Hartopp et al., 2019; Metzler-Baddeley et al., 2011) and increased

frontal MD (Nicolas et al., 2020). In addition to such retrieval effects, using the immediate

free recall components across three consecutive trials, the FCSRT enables investigating

episode learning by examining how many additional words are successfully recalled on each

trial.

We investigated whether the learning rate in FCSRT is associated with age, as well as its

neural manifestation in GM volume and WM tract microstructure. In a large cross-sectional

cohort of healthy older adults, we first calculated the learning rate across the three

consecutive FCSRT trials and tested for an association with age. To examine brain-cognition

associations, we used learning rate as a predictor to examine 1) GM volume using voxel-

based morphometry (VBM), and 2) WM microstructure using tract-based spatial statistics on

FA and MD measures. Given the critical role of the hippocampus in episodic memory and its

correlation with structural changes in ageing, we hypothesised that hippocampal GM volume

and associated WM limbic tract microstructure would correlate with learning rate.

Methods

Participants. All participants in this study were part of the Vallecas Project, a single-centre

longitudinal study of community-dwelling volunteers aged 69-86 without any cognitive or

psychiatric disorder that compromised their daily functioning at the time of recruitment.

Inclusion and exclusion criteria have been further described elsewhere (Olazarán et al.,

2015). From this cohort, data from the baseline visit of 982 cognitively normal participants

(mean age = 74.8, SD = 3.9, 637 (64.9%) females) were included in the current study. Any

subject with a diagnosis of mild cognitive impairment or Alzheimer's disease at this first visit

was excluded. All participants provided written informed consent and the Vallecas Project

was approved by the Ethics committee of the Instituto de Salud Carlos III.

Neuropsychological assessment. Participants completed a battery of neuropsychological

assessments as part of the Vallecas Project protocol. In this study, we report the total score

of the Mini Mental State Examination (MMSE; Folstein et al., 1975) and we mainly focused

on the Free and Cued Selective Reminding Test (FCSRT; Buschke, 1984), assessing

learning and retention of verbal memory, with immediate and delayed recall components.

The test was administered using standard procedures (Peña-Casanova et al., 2009).

Participants were presented with cards containing four words and asked to identify the word

corresponding to a specific semantic category, going through all four words, on four different

cards (16 words in total). The words presented are not the most obvious member of each

semantic category. Following the presentation phase, participants were asked to recall as

many words as possible in three consecutive recall trials each one followed by 20 seconds

of interference counting backwards (Figure 1A). For each trial, participants were asked to

freely recall as many words as possible with a time limit of 90 seconds, then examiners

provided the semantic category clue for the forgotten items. These three free and cued

recalls constitute the three immediate recall trials of the task. This immediate recall phase is

followed by a 30-minute delay, after which the delayed phase of the test starts. Participants

were asked on a single trial to freely recall as many words as possible otherwise cues were

provided (Figure 1A). To assess the learning rate across trials, we fit a linear mixed-effects

model of the number of items freely recalled in each immediate trial, as a function of the

recall trial (first, second, and third) using the lme4 package in R 4.0.2 (https://www.r-

project.org/). The model also included a random slope of the recall trial, and a random

intercept per participant, capturing inter-individual variability in learning rate (across the three

trials). The learning rate coefficient for each participant was extracted using the coef()

function for subsequent analyses. Next, we built a multiple regression model where the

learning rate was the dependent variable, sex, age and level of education were the

predictors and the delayed free recall score of the FCSRT was included as a covariate to

rule out the retrieval phase of the memory process. Extraction and plotting of the effects

reported below were conducted using the effects (Fox, 2003) and ggplot2 (Wickham, 2009)

packages in R.

MRI Data acquisition. Images were acquired using a 3T MRI (Signa HDxt GE) with a phased

array eight-channel head coil. T1-weighted images (3D fast spoiled gradient echo with

inversion recovery preparation) were collected using a repetition time (TR) of 10ms, echo

time (TE) of 4.5ms, field of view (FOV) of 240mm and a matrix size of 288x288 with slice

thickness of 1mm, yielding a voxel size of 0.5 x 0.5 x 1 mm3. Diffusion-weighted images

were single-shot spin echo echo-planar imaging (SE-EPI), with TR 9200ms, TE 80ms, b-

value 800s/mm2 and 21 gradient directions, FOV 240mm and matrix size 128 x 128 with

slice thickness of 3mm.

Grey matter VBM. The analysis was carried out in SPM12 (version r6225;

https://www.fil.ion.ucl.ac.uk/spm). T1-weighted images were segmented into grey matter,

white matter and cerebrospinal fluid and then aligned and normalised to MNI space using

the DARTEL algorithm (Ashburner, 2007). Prior to statistical modelling, the normalised

images were smoothed using a 6mm FWHM Gaussian kernel. The pre-processed grey

matter maps were entered into a general linear model (GLM) with learning rate from the

memory task as the predictor of interest, and total intracranial volume (TIV), sex, and the

delayed free recall score of the FCSRT as covariates. Age and education were not used in

the model as additional covariates since FCSRT delayed free recall is sensitive to the effects

of age and level of education. Nonetheless, to ensure the model is capturing variance

associated with these variables we devised a second model without FCSRT delayed free

recall and including TIV, sex, age and education as covariates and the same results were

obtained (see Supplementary Materials). We conducted whole-brain analyses using a

threshold-free cluster enhancement (TFCE) approach with 5000 permutations and default

parameters (E = 0.5 and H = 2) using the TFCE tool (version r223) for CAT12 toolbox in

SPM (http://dbm.neuro.uni-jena.de/tfce). Therefore, our analyses fully correct for mass-

univariate testing (and associated multiple-comparisons problem) by employing a whole-

brain FWE correction. Furthermore, we used the TFCE approach to overcome cluster-based

inference issues.   The AAL3 atlas neuroanatomical labels were used to describe

neuroanatomical loci (Rolls et al., 2020) and Mango software was used to produce the figure

(http://rii.uthscsa.edu/mango/). These analyses assessed which regions were positively

associated with the immediate recall learning rate. Significant results are reported at a

family-wise error FWE) corrected level of $p < 0.05$.

White matter tract-based spatial statistics (TBSS). Of the 982 participants, seven were

excluded as they did not have diffusion data. For preprocessing these images, the FSL

toolbox (http://fsl.fmrib.ox.ac.uk/fsl/fslwiki) was used for motion and eddy current correction,

the extraction of non-brain voxels and, lastly, the calculation of voxel-wise diffusion maps

(FA and MD) for each participant. Individual FA and MD maps were then used in the FSL

TBSS   pipeline   (http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/TBSS/UserGuider;   detailed   methods

described by Smith et al. (2006)). The general outline of the process is: 1) FA individual

maps were non-linearly registered to standard space (FMRIB58_FA template) (Andersson et

al., 2007). 2) A mean FA image was created by averaging all co-registered FA maps. 3)

Individually aligned images were projected onto the mean FA skeleton—representing the

centers of all tracts common to the study sample—and skeletonised images were used for

voxel-wise analysis. Diffusivity maps for MD were generated by applying the same steps

detailed above. The same GLM design matrix as the VBM analysis was used along with the

TFCE approach with 5000 permutations (default parameters E = 0.5 and H = 2). Significant

results are reported at a family-wise error (FWE) corrected level of $p < 0.05$. To visualise our

TBSS results we used the multimodal analysis and visualisation tool (MMVT; Felsenstein et

al., 2019). The pipeline follows these steps: 1) Binary masking: all the voxels in the TBSS

volume below the threshold (0.95) were set to zero. 2) Outlier voxels removal using the

Open3D python package (Zhou et al., 2018). 3) Smoothing the volumetric data using a 3D

Gaussian filter (Virtanen et al., 2020). 4) Surface creation from the volume's TBSS surfaces

using the marching cubes algorithm (Lorensen and Cline, 1987). For that, we re-calculate

the threshold to give us the same number of voxels after the smoothing step. 5) Translation

for the surfaces' vertices coordinates. 6) Projection of the volumetric data on the surfaces.

<u>Results</u>

Memory and neuropsychological performance

On average, across three trials, participants correctly remembered 7.9 items (SD = 2.6).

When looking at individual trials, performance improved as trials progressed, reflecting a

positive episodic learning rate (see Table 1 for number of items recalled, and Figure 1B for

learning rates). Our linear model predicting the learning rate as a function of age, sex, and

level of education revealed significant effects of the three predictors after correcting for

FCSRT delayed free recall score. Learning rate and delayed free recall FCSRT were

positively correlated (Pearson's $r = 0.7$; $p < 2.2{\times}10{-}16$). Age had a negative effect on

learning rate ($F(1,965) = 10.45$, $p = 0.001$) (Figure 1C), sex also had an effect ($F(1,965) =$

4.66, $p = 0.031$) and being a woman was positively associated with learning rate (Figure

1D). Finally, having a higher level of education was positively associated with learning rate

($F(3,965) = 5.17$, $p < 0.002$) (Figure 1E). There was no significant interaction between the

three predictors (age, sex and years of education).

196
197

	Total sample (n = 982) Mean (SD)	Women (n = 637) Mean (SD)	Men (n = 345) Mean (SD)

Demographics

Age, years		74.8 (3.9)	74.8 (3.9)	74.8 (3.9)
Levels of education, count (%)	Less than primary	185 (18.8)	137 (21.5)	48 (13.9)
	Primary	295 (30.0)	222 (34.9)	73 (21.1)
	Secondary	245 (24.9)	151 (23.7)	94 (27.2)
	Higher education	257 (26.2)	127 (19.9)	130 (37.7)
Neuropsychological performance				
MMSE, total score		28.6 (1.6)	28.6 (1.6)	28.7 (1.4)
Trial 1 immediate free recall FCSRT, items recalled		6.5 (2.1)	6.7 (2.1)	6.3 (2.1)
Trial 2 immediate free recall FCSRT, items recalled		7.9 (2.4)	7.9 (2.4)	7.8 (2.4)
Trial 3 immediate free recall FCSRT, items recalled		9.2 (2.5)	9.4 (2.6)	9.0 (2.6)
Learning rate immediate free recall FCSRT, items recalled/trial		1.4 (0.3)	1.4 (0.3)	1.3 (0.3)

Delayed free recall FCSRT, items recalled	9.4 (2.6)	9.5 (2.7)	9.3 (2.6)

Table 1. Demographic and neuropsychological profile of the total sample and split by sex. MMSE: Mini Mental State Examination total score, FCSRT: Free and Cued Selective Reminding Test.

Figure 1. Learning rate across FCSRT trials is related to age, sex and level of education. A Diagram of the FCSRT protocol used to assess memory in this study in which the learning rate is calculated from the free recall of the three immediate trials (blue). B Mean and individual learning rates across the three free immediate recall FCSRT trials. After controlling for the rest of the model predictors, C learning rate decreases with age, D females learn faster than males and E level of education is positively associated with learning rate. Error bars represent 95% confidence intervals.

Grey matter volume (VBM)

We found a positive correlation between episodic learning rate and grey matter volume in

the bilateral hippocampus, with more pronounced effects on the left side and the left superior

temporal gyrus (Figure 2A, Supplementary Figure 1), and the right anterior thalamic nucleus

with some extension to adjacent nuclei (right ventroanterior and ventrolateral thalamic nuclei;

Figure 2B, Supplementary Table 1, Supplementary Figure 1).

Figure 2. Grey matter volume correlates with episodic learning rate in older adults.
The positive correlation has been overlaid on a canonical T1 image (thresholded at p < 0.05
FWE-corr) to show a significant effect in A) hippocampus bilaterally and left superior
temporal gyrus and B) right anterior (red) and ventroanterior (green) (thalamic ROIs in the
inset come from the AAL3 atlas (Rolls et al., 2020)). The coordinates of the sections are
given in mm. L: left, R: right, A: anterior, P: posterior.

White matter microstructure (TBSS)

We first examined FA as a marker of WM integrity. We found a bilateral network of temporal,

parietal and occipital tracts showing a positive association with episodic learning rate.

Among tracts showing significant positive correlations were the bilateral anterior thalamic

radiation (ATR), fornix, inferior fronto-occipital fasciculus (IFOF), inferior longitudinal

fasciculus (ILF), superior longitudinal fasciculus (SLF), and uncinate (Figure 3A-B,

Supplementary Table 2, Supplementary Figure 2). Next, we examined MD and found a

negative association between a similar network of bilateral tracts and FCSRT learning rate,

including bilateral ATR, corticospinal tract, forceps major and minor, cingulum (cingulate),

IFOF, ILF, SLF, uncinate and fornix (Figure 3C-D, Supplementary Figure 2, Supplementary

Table 2).

Figure 3. Extensive network of white matter microstructure integrity is related to
episodic learning rate in older adults. A. Positive correlation between FA and learning
rate (warm colours; p < 0.05 FWE-corr). B. FA effects overlaid on the fornix. C. Negative
correlation between MD and learning rate (cold colours; p < 0.05 FWE-corr). D. MD effect
overlaid on the fornix.

Discussion

Our results show that women and individuals with more years of formal education had a

faster episodic learning rate, that it declined with age, and that this rate was associated with

neuroanatomical structural properties. We found a positive correlation between GM volume

and episodic learning rate, where participants with greater volume in hippocampus, anterior

thalamic nucleus and left superior temporal gyrus learned at a faster rate than those with

lower volume. Furthermore, we found that FA was positively associated with episodic

learning rate in an extended network including limbic tracts, indicating that the structural

integrity of these tracts indexed learning ability. A complementary negative association was

observed for MD, in similar tracts, such that decreased MD was associated with a faster

episodic learning rate. The converging GM and WM findings suggest that structural

properties of the hippocampal-anterior thalamic circuit contribute to learning ability in ageing

and may potentially inform age-related decline in encoding (Friedman et al., 2007; Morcom

et al., 2003).

Previous research on neural substrates of cognitive decline in ageing has shown a

hippocampal volume decline with age that correlated with memory performance and with

FCSRT recall specifically (Zammit et al., 2017). The presence of hippocampal volume

findings with relation to both episodic learning rate and recall components of the FCSRT task

suggests the hippocampus may be involved in these two separate processes, both of which

are impaired in ageing. The more pronounced effect we observed in the left hippocampus is

in accordance with previous VBM findings of verbal memory tasks (Ezzati et al., 2016), and

the general lateralisation of verbal functions. Our results, therefore, extend previous

research on the relationship between hippocampal volume and memory decline in ageing,

showing episodic learning rate is also indexed by hippocampal volume. Note that it is

unlikely that these effects reflect memory function in general, given that delayed free recall

was included as a covariate in our model.

Structural properties of extra-hippocampal limbic regions were also associated with learning

ability. Our GM thalamic findings indicate a correlation between episodic learning rate and

the right anterior thalamic nucleus, extending to right ventroanterior and right ventrolateral

nuclei. The anterior thalamic nuclei have been suggested to play an important role in

learning and memory (Aggleton et al., 2010; Sweeney-Reed et al., 2021; Winocur, 1985),

that extend beyond its established role in spatial processing (Nelson, 2021; Wolff and Vann,

2019). For example, fMRI studies in younger adults suggest that the activation of the anterior

thalamic nuclei supports recognition memory performance (Pergola et al., 2013) and

evidence from intracranial EEG studies indicates theta-synchronisation between anterior

thalamus and frontal and parietal regions supporting successful memory formation

(Sweeney-Reed et al., 2014). Furthermore, and in line with our results, Leszczyński and

Staudigl (2016) posited that the anterior thalamus might modulate information flow, via

attention allocation, to support learning. Taken together with the increased hippocampal

volume, which was related to a better learning rate, our results indicate that the limbic

system may play an important role in learning ability in ageing and might explain some of the

impairments in navigating in a novel environment (Grzeschik et al., 2021), and impaired

learning strategies observed in mild cognitive impairment (Ribeiro et al., 2007).

The fornix is a major hippocampal input/output pathway and has been associated with visuo-

spatial learning across species (Buckley et al., 2008; Hodgetts et al., 2020; Hofstetter et al.,

2013). The fornix links the hippocampus with the anterior thalamic nuclei directly and via the

mammillary bodies (Aggleton et al., 2010, 1986), with both the hippocampus and the anterior

thalamic nuclei showing grey matter volume relationships with episodic learning rate.

Furthermore, we found that fornix integrity, as captured by bilateral FA and MD, correlated

with episodic learning rate in older adults. Together with previous findings linking fornix

integrity to recall performance on the FCSRT task (Hartopp et al., 2019; Metzler-Baddeley et

al., 2011), our results extend its role in memory processes, indicating that the fornix also

supports verbal episodic learning. We also found that the WM integrity of the ATR was

correlated with learning rate. The ATR connects the anterior and dorsomedial thalamus with

the prefrontal cortex (Grodd et al., 2020), which has been suggested to play a role in

learning rate (McGuire et al., 2014).

In addition to changes in limbic GM volume and WM microstructure, we found episodic

learning rate was associated with broader changes within bilateral WM tracts connecting

occipital-temporal-frontal regions (ILF, SLF, IFOF). This result might point toward an overall

WM microstructure effect, as previously noted in age-related cognitive decline (de Mooij et

al., 2018; Farokhian et al., 2017; Grieve et al., 2007; Molloy et al., 2021; Rizvi et al., 2020).

With respect to specific cognitive functions, ILF and SLF have been shown to relate to

memory performance in normal ageing (Sasson et al., 2013). ILF and IFOF facilitate the flow

of visual information up the visual stream (Rokem et al., 2017), and IFOF has been

associated with semantic processing (Duffau, 2008), potentially supporting learning

performance in our task. Therefore, the observed relationship between microstructure of

these tracts and learning ability might reflect a more general aspect of cognitive ability.

Finally, it is important to note some limitations of the current study; we analysed data from a

cross-sectional cohort of healthy older adults. As GM and WM properties and memory

function both deteriorate with age, future longitudinal studies would be needed to better

understand the relationship between learning ability and structural changes as ageing

progresses and eliminate age-related confounds in cross-sectional studies (Elliott, 2020).

We used the learning rate across trials in an established neuropsychological memory task

(FCSRT) as a measure for episodic learning; it would be interesting to examine neural

correlates of learning rates in tasks such as error-driven and statistical learning (Herff et al.,

2019; Nassar et al., 2016; Samanez-Larkin et al., 2012), as well as consider learning ability

as a potential cognitive phenotype in pathological ageing. Finally, future research with

hippocampal subfield resolution could examine their differential contribution to episodic

learning rate. It would be interesting to explore whether volumetric effects are more

pronounced in the subiculum, the principal source of hippocampal projections to the anterior

thalamus and mammillary bodies (Hartopp et al., 2019).

In conclusion, in a cross-sectional cohort of healthy older adults, we found learning rate on

the FCSRT task was positively associated with extensive GM and WM structural effects

including the hippocampus, fornix and anterior thalamic nucleus, structures part of the limbic

system. Furthermore, there was a positive correlation between episodic learning rate and

long-range WM tracts (ILF, SLF, IFOF). Our findings indicate that episodic learning rate is

associated with key anatomical structures implicated in memory function, and therefore may

inform further exploration of the relationship between episodic learning rate and retrieval in

ageing.

References

Aggleton, J.P., Desimone, R., Mishkin, M., 1986. The origin, course, and termination of the

hippocampothalamic projections in the macaque. J. Comp. Neurol. 243, 409–421.

https://doi.org/10.1002/cne.902430310

Aggleton, J.P., O'Mara, S.M., Vann, S.D., Wright, N.F., Tsanov, M., Erichsen, J.T., 2010.

Hippocampal–anterior thalamic pathways for memory: uncovering a network of direct

and indirect actions. Eur. J. Neurosci. 31, 2292–2307. https://doi.org/10.1111/J.1460-

9568.2010.07251.X

Ashburner, J., 2007. A fast diffeomorphic image registration algorithm. Neuroimage 38, 95–

113. https://doi.org/10.1016/j.neuroimage.2007.07.007

Bennett, I.J., Huffman, D.J., Stark, C.E.L., 2015. Limbic tract integrity contributes to pattern

separation performance across the lifespan. Cereb. Cortex 25, 2988–2999.

https://doi.org/10.1093/cercor/bhu093

Bennett, I.J., Madden, D.J., Vaidya, C.J., Howard, D. V., Howard, J.H., 2010. Age-related

differences in multiple measures of white matter integrity: A diffusion tensor imaging

study of healthy aging. Hum. Brain Mapp. 31, 378–390.

https://doi.org/10.1002/hbm.20872

Boujut, A., Clarys, D., 2016. The effect of ageing on recollection: the role of the binding

updating process. Memory 24, 1231–1242.

https://doi.org/10.1080/09658211.2015.1091893

Buckley, M.J., Wilson, C.R.E., Gaffan, D., 2008. Fornix Transection Impairs Visuospatial

Memory Acquisition More Than Retrieval. Behav. Neurosci. 122, 44–53.

https://doi.org/10.1037/0735-7044.122.1.44

Buschke, H., 1984. Cued recall in amnesia. J. Clin. Neuropsychol. 6, 433–40.

Cadar, D., Usher, M., Davelaar, E.J., 2018. Age-Related Deficits in Memory Encoding and

Retrieval in Word List Free Recall. Brain Sci. 8, 211.

https://doi.org/10.3390/brainsci8120211

Cox, S.R., Bastin, M.E., Ritchie, S.J., Dickie, D.A., Liewald, D.C., Muñoz Maniega, S.,

Redmond, P., Royle, N.A., Pattie, A., Valdés Hernández, M., Corley, J., Aribisala, B.S.,

McIntosh, A.M., Wardlaw, J.M., Deary, I.J., 2018. Brain cortical characteristics of

lifetime cognitive ageing. Brain Struct. Funct. 223, 509–518.

https://doi.org/10.1007/s00429-017-1505-0

Cox, S.R., Harris, M.A., Ritchie, S.J., Buchanan, C.R., Valdés Hernández, M.C., Corley, J.,

Taylor, A.M., Madole, J.W., Harris, S.E., Whalley, H.C., McIntosh, A.M., Russ, T.C.,

Bastin, M.E., Wardlaw, J.M., Deary, I.J., Tucker-Drob, E.M., 2021. Three major

dimensions of human brain cortical ageing in relation to cognitive decline across the

eighth decade of life. Mol. Psychiatry 26, 2651–2662. https://doi.org/10.1038/s41380-

020-00975-1

de Mooij, S.M.M., Henson, R.N.A., Waldorp, L.J., Kievit, R.A., 2018. Age differentiation

within gray matter, white matter, and between memory and white matter in an adult life

span cohort. J. Neurosci. 38, 5826–5836. https://doi.org/10.1523/JNEUROSCI.1627-

17.2018

Duffau, H., 2008. The anatomo-functional connectivity of language revisited. New insights

provided by electrostimulation and tractography. Neuropsychologia 46, 927–934.

https://doi.org/10.1016/j.neuropsychologia.2007.10.025

Elliott, M.L., 2020. MRI-based biomarkers of accelerated aging and dementia risk in midlife:

how close are we? Ageing Res. Rev. 61, 101075.

https://doi.org/10.1016/j.arr.2020.101075

Ezzati, A., Katz, M.J., Zammit, A.R., Lipton, M.L., Zimmerman, M.E., Sliwinski, M.J., Lipton,

R.B., 2016. Differential association of left and right hippocampal volumes with verbal

episodic and spatial memory in older adults. Neuropsychologia 93, 380–385.

https://doi.org/10.1016/j.neuropsychologia.2016.08.016

Farokhian, F., Yang, C., Beheshti, I., Matsuda, H., Wu, S., 2017. Age-related gray and white

matter changes in normal adult brains. Aging Dis. 8, 899–909.

https://doi.org/10.14336/AD.2017.0502

Felsenstein, O., Peled, N., Hahn, E., Rockhill, A.P., Folsom, L., Gholipour, T., Macadams,

K., Rozengard, N., Paulk, A.C., Dougherty, D., Cash, S.S., Widge, A.S., Hämäläinen,

M., Stufflebeam, S., 2019. Multi-Modal Neuroimaging Analysis and Visualization Tool

(MMVT).

Fjell, A.M., Walhovd, K.B., 2010. Structural Brain Changes in Aging: Courses, Causes and

Cognitive Consequences. Rev. Neurosci. 21.

https://doi.org/10.1515/REVNEURO.2010.21.3.187

Fox, J., 2003. Effect Displays in R for Generalised Linear Models. J. Stat. Softw. 8.

https://doi.org/10.18637/jss.v008.i15

Friedman, D., Nessler, D., Johnson, R., 2007. Memory Encoding and Retrieval in the Aging

Brain. Clin. EEG Neurosci. 38, 2–7. https://doi.org/10.1177/155005940703800105

Glisky, E.L., 2007. Changes in Cognitive Function in Human Aging.

Gorbach, T., Pudas, S., Lundquist, A., Orädd, G., Josefsson, M., Salami, A., de Luna, X.,

Nyberg, L., 2017. Longitudinal association between hippocampus atrophy and episodic-

memory decline. Neurobiol. Aging 51, 167–176.

https://doi.org/10.1016/j.neurobiolaging.2016.12.002

Grady, C., 2012. The cognitive neuroscience of ageing. Nat. Rev. Neurosci. 13, 491–505.

https://doi.org/10.1038/nrn3256

Grieve, S.M., Williams, L.M., Paul, R.H., Clark, C.R., Gordon, E., 2007. Cognitive aging,

executive function, and fractional anisotropy: A diffusion tensor MR imaging study. Am.

J. Neuroradiol. 28, 226–235.

Grodd, W., Kumar, V.J., Schüz, A., Lindig, T., Scheffler, K., 2020. The anterior and medial

thalamic nuclei and the human limbic system: tracing the structural connectivity using

diffusion-weighted imaging. Sci. Rep. 10, 1–25. https://doi.org/10.1038/s41598-020-

67770-4

Grzeschik, R., Hilton, C., Dalton, R.C., Konovalova, I., Cotterill, E., Innes, A., Wiener, J.M.,

2021. From repeating routes to planning novel routes: the impact of landmarks and

ageing on route integration and cognitive mapping. Psychol. Res. 85, 2164–2176.

https://doi.org/10.1007/s00426-020-01401-5

Hartopp, N., Wright, P., Ray, N.J., Evans, T.E., Metzler-Baddeley, C., Aggleton, J.P.,

O'Sullivan, M.J., 2019. A key role for subiculum-fornix connectivity in recollection in

older age. Front. Syst. Neurosci. 12, 1–11. https://doi.org/10.3389/fnsys.2018.00070

Hedden, T., Schultz, A.P., Rieckmann, A., Mormino, E.C., Johnson, K.A., Sperling, R.A.,

Buckner, R.L., 2016. Multiple Brain Markers are Linked to Age-Related Variation in

Cognition. Cereb. Cortex 26, 1388–1400. https://doi.org/10.1093/cercor/bhu238

Herff, S., Rashid, N.A.B.A., Keong, J.L.C., Tih-Shih, L., Agres, K., 2019. Statistical Learning

Ability as a Measure of Cognitive Function. https://doi.org/10.31234/osf.io/u4ry6

Hodgetts, C.J., Stefani, M., Williams, A.N., Kolarik, B.S., Yonelinas, A.P., Ekstrom, A.D.,

Lawrence, A.D., Zhang, J., Graham, K.S., 2020. The role of the fornix in human

navigational learning. Cortex 124, 97–110. https://doi.org/10.1016/j.cortex.2019.10.017

Hofstetter, S., Tavor, I., Moryosef, S.T., Assaf, Y., 2013. Short-term learning induces white

matter plasticity in the fornix. J. Neurosci. 33, 12844–12850.

https://doi.org/10.1523/JNEUROSCI.4520-12.2013

Kennedy, K.M., Raz, N., 2009. Aging white matter and cognition: Differential effects of

regional variations in diffusion properties on memory, executive functions, and speed.

Neuropsychologia 47, 916–927. https://doi.org/10.1016/j.neuropsychologia.2009.01.001

Kwok, S.C., Buckley, M.J., 2010. Long-term visuospatial retention unaffected by fornix

transection. Hippocampus 20, 889–893. https://doi.org/10.1002/hipo.20733

Laukka, E.J., Lövdén, M., Kalpouzos, G., Li, T.Q., Jonsson, T., Wahlund, L.O., Fratiglioni, L.,

Bäckman, L., 2013. Associations between white matter microstructure and cognitive

performance in old and very old age. PLoS One 8, 1–8.

https://doi.org/10.1371/journal.pone.0081419

Leszczyński, M., Staudigl, T., 2016. Memory-guided attention in the anterior thalamus.

Neurosci. Biobehav. Rev. 66, 163–165. https://doi.org/10.1016/j.neubiorev.2016.04.015

Lorensen, W.E., Cline, H.E., 1987. Marching cubes: A high resolution 3D surface

construction algorithm, in: Proceedings of the 14th Annual Conference on Computer

Graphics and Interactive Techniques - SIGGRAPH '87. ACM Press, New York, New

York, USA, pp. 163–169. https://doi.org/10.1145/37401.37422

Madden, D.J., Parks, E.L., 2017. Age differences in structural connectivity: Diffusion tensor

imaging and white matter hyperintensitie, in: Cabeza, R., Nyberg, L., Park, D.C. (Eds.),

Cognitive Neuroscience of Aging: Linking Cognitive and Cerebral Aging. Oxford, New

York, pp. 71–103.

McGuire, J.T., Nassar, M.R., Gold, J.I., Kable, J.W., 2014. Functionally Dissociable

Influences on Learning Rate in a Dynamic Environment. Neuron 84, 870–881.

https://doi.org/10.1016/j.neuron.2014.10.013

Metzler-Baddeley, C., Jones, D.K., Belaroussi, B., Aggleton, J.P., O'Sullivan, M.J., 2011.

Frontotemporal connections in episodic memory and aging: A diffusion MRI

tractography study. J. Neurosci. 31, 13236–13245.

https://doi.org/10.1523/JNEUROSCI.2317-11.2011

Molloy, C.J., Nugent, S., Bokde, A.L.W., 2021. Alterations in diffusion measures of white

matter integrity associated with healthy aging. Journals Gerontol. - Ser. A Biol. Sci.

Med. Sci. 76, 945–954. https://doi.org/10.1093/gerona/glz289

Morcom, A.M., Good, C.D., Frackowiak, R.S.J., Rugg, M.D., 2003. Age effects on the neural

correlates of successful memory encoding. Brain 126, 213–229.

https://doi.org/10.1093/brain/awg020

Nassar, M.R., Bruckner, R., Gold, J.I., Li, S.C., Heekeren, H.R., Eppinger, B., 2016. Age

differences in learning emerge from an insufficient representation of uncertainty in older

adults. Nat. Commun. 7, 1–13. https://doi.org/10.1038/ncomms11609

Nelson, A.J.D., 2021. The anterior thalamic nuclei and cognition: A role beyond space?

Neurosci. Biobehav. Rev. https://doi.org/10.1016/j.neubiorev.2021.02.047

Nicolas, R., Hiba, B., Dilharreguy, B., Barse, E., Baillet, M., Edde, M., Pelletier, A., Periot, O.,

Helmer, C., Allard, M., Dartigues, J.F., Amieva, H., Pérès, K., Fernandez, P., Catheline,

G., 2020. Changes Over Time of Diffusion MRI in the White Matter of Aging Brain, a

Good Predictor of Verbal Recall. Front. Aging Neurosci. 12, 1–9.

https://doi.org/10.3389/fnagi.2020.00218

Olazarán, J., Valentí, M., Belén Frades, Zea-Sevilla, M.A., Ávila-Villanueva, M., Fernández-

Blázquez, M.Á., Calero, M., Dobato, J.L., Hernández-Tamames, J.A., León-Salas, B.,

Agüera-Ortiz, L., López-Álvarez, J., Larrañaga, P., Bielza, C., Álvarez-Linera, J.,

Martínez-Martín, P., 2015. The Vallecas Project: A cohort to identify early markers and

mechanisms of Alzheimer's disease. Front. Aging Neurosci. 7.

https://doi.org/10.3389/fnagi.2015.00181

Peña-Casanova, J., Gramunt-Fombuena, N., Quiñones-Úbeda, S., Sánchez-Benavides, G.,

Aguilar, M., Badenes, D., Molinuevo, J.L., Robles, A., Barquero, M.S., Payno, M.,

Antúnez, C., Martínez-Parra, C., Frank-García, A., Fernández, M., Alfonso, V., Sol,

J.M., Blesa, R., 2009. Spanish Multicenter Normative Studies (NEURONORMA

Project): norms for the Rey-Osterrieth complex figure (copy and memory), and free and

cued selective reminding test. Arch. Clin. Neuropsychol. 24, 371–393.

https://doi.org/10.1093/ARCLIN/ACP041

Pergola, G., Ranft, A., Mathias, K., Suchan, B., 2013. The role of the thalamic nuclei in

recognition memory accompanied by recall during encoding and retrieval: An fMRI

study. Neuroimage 74, 195–208. https://doi.org/10.1016/j.neuroimage.2013.02.017

Persson, J., Pudas, S., Lind, J., Kauppi, K., Nilsson, L.-G., Nyberg, L., 2012. Longitudinal

Structure-Function Correlates in Elderly Reveal MTL Dysfunction with Cognitive

Decline. Cereb. Cortex 22, 2297–2304. https://doi.org/10.1093/cercor/bhr306

Resnick, S.M., Pham, D.L., Kraut, M.A., Zonderman, A.B., Davatzikos, C., 2003.

Longitudinal Magnetic Resonance Imaging Studies of Older Adults: A Shrinking Brain.

J. Neurosci. 23, 3295–3301. https://doi.org/10.1523/JNEUROSCI.23-08-03295.2003

Ribeiro, F., Guerreiro, M., De Mendonça, A., 2007. Verbal learning and memory deficits in

Mild Cognitive Impairment. J. Clin. Exp. Neuropsychol. 29, 187–197.

https://doi.org/10.1080/13803390600629775

Rizvi, B., Lao, P.J., Colón, J., Hale, C., Igwe, K.C., Narkhede, A., Budge, M., Manly, J.J.,

Schupf, N., Brickman, A.M., 2020. Tract-defined regional white matter hyperintensities

and memory. NeuroImage Clin. 25, 102143. https://doi.org/10.1016/j.nicl.2019.102143

Rokem, A., Takemura, H., Bock, A.S., Scherf, K.S., Behrmann, M., Wandell, B.A., Fine, I.,

Bridge, H., Pestilli, F., 2017. The visual white matter: The application of diffusion MRI

and fiber tractography to vision science. J. Vis. https://doi.org/10.1167/17.2.4

Salami, A., Eriksson, J., Nilsson, L.G., Nyberg, L., 2012. Age-related white matter

microstructural differences partly mediate age-related decline in processing speed but

not cognition. Biochim. Biophys. Acta - Mol. Basis Dis. 1822, 408–415.

https://doi.org/10.1016/j.bbadis.2011.09.001

Samanez-Larkin, G.R., Levens, S.M., Perry, L.M., Dougherty, R.F., Knutson, B., 2012.

Frontostriatal white matter integrity mediates adult age differences in probabilistic

reward learning. J. Neurosci. 32, 5333–5337.

https://doi.org/10.1523/JNEUROSCI.5756-11.2012

Sasson, E., Doniger, G.M., Pasternak, O., Tarrasch, R., Assaf, Y., 2013. White matter

correlates of cognitive domains in normal aging with diffusion tensor imaging. Front.

Neurosci. https://doi.org/10.3389/fnins.2013.00032

Sweeney-Reed, C.M., Buentjen, L., Voges, J., Schmitt, F.C., Zaehle, T., Kam, J.W.Y.,

Kaufmann, J., Heinze, H.J., Hinrichs, H., Knight, R.T., Rugg, M.D., 2021. The role of

the anterior nuclei of the thalamus in human memory processing. Neurosci. Biobehav.

Rev. 126, 146–158. https://doi.org/10.1016/j.neubiorev.2021.02.046

Sweeney-Reed, C.M., Zaehle, T., Voges, J., Schmitt, F.C., Buentjen, L., Kopitzki, K.,

Esslinger, C., Hinrichs, H., Heinze, H.J., Knight, R.T., Richardson-Klavehn, A., 2014.

Corticothalamic phase synchrony and cross-frequency coupling predict human memory

formation. Elife 3, e05352. https://doi.org/10.7554/eLife.05352

Tromp, D., Dufour, A., Lithfous, S., Pebayle, T., Després, O., 2015. Episodic memory in

normal aging and Alzheimer disease: Insights from imaging and behavioral studies.

Ageing Res. Rev. 24, 232–262. https://doi.org/10.1016/j.arr.2015.08.006

Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D.,

Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S.J., Brett, M.,

Wilson, J., Millman, K.J., Mayorov, N., Nelson, A.R.J., Jones, E., Kern, R., Larson, E.,

Carey, C.J., Polat, İ., Feng, Y., Moore, E.W., VanderPlas, J., Laxalde, D., Perktold, J.,

Cimrman, R., Henriksen, I., Quintero, E.A., Harris, C.R., Archibald, A.M., Ribeiro, A.H.,

Pedregosa, F., van Mulbregt, P., Vijaykumar, A., Bardelli, A. Pietro, Rothberg, A.,

Hilboll, A., Kloeckner, A., Scopatz, A., Lee, A., Rokem, A., Woods, C.N., Fulton, C.,

Masson, C., Häggström, C., Fitzgerald, C., Nicholson, D.A., Hagen, D.R., Pasechnik, D.

V., Olivetti, E., Martin, E., Wieser, E., Silva, F., Lenders, F., Wilhelm, F., Young, G.,

Price, G.A., Ingold, G.-L., Allen, G.E., Lee, G.R., Audren, H., Probst, I., Dietrich, J.P.,

Silterra, J., Webber, J.T., Slavič, J., Nothman, J., Buchner, J., Kulick, J., Schönberger,

J.L., de Miranda Cardoso, J.V., Reimer, J., Harrington, J., Rodríguez, J.L.C., Nunez-

Iglesias, J., Kuczynski, J., Tritz, K., Thoma, M., Newville, M., Kümmerer, M.,

Bolingbroke, M., Tartre, M., Pak, M., Smith, N.J., Nowaczyk, N., Shebanov, N., Pavlyk,

O., Brodtkorb, P.A., Lee, P., McGibbon, R.T., Feldbauer, R., Lewis, S., Tygier, S.,

Sievert, S., Vigna, S., Peterson, S., More, S., Pudlik, T., Oshima, T., Pingel, T.J.,

Robitaille, T.P., Spura, T., Jones, T.R., Cera, T., Leslie, T., Zito, T., Krauss, T.,

Upadhyay, U., Halchenko, Y.O., Vázquez-Baeza, Y., 2020. SciPy 1.0: fundamental

algorithms for scientific computing in Python. Nat. Methods 17, 261–272.

https://doi.org/10.1038/s41592-019-0686-2

Wickham, H., 2009. ggplot2. Springer New York, New York, NY. https://doi.org/10.1007/978-

0-387-98141-3

Winocur, G., 1985. THE HIPPOCAMPUS AND THALAMUS: THEIR ROLES IN SHORT-

AND LONG-TERM MEMORY AND THE EFFECTS OF INTERFERENCE, Behavioural

Bra#z Research.

Wolff, M., Vann, S.D., 2019. The cognitive thalamus as a gateway to mental representations.

J. Neurosci. 39, 3–14. https://doi.org/10.1523/JNEUROSCI.0479-18.2018

Zammit, A.R., Ezzati, A., Zimmerman, M.E., Lipton, R.B., Lipton, M.L., Katz, M.J., 2017.

Roles of hippocampal subfields in verbal and visual episodic memory. Behav. Brain

Res. 317, 157–162. https://doi.org/10.1016/j.bbr.2016.09.038

Zhou, Q.-Y., Park, J., Koltun, V., 2018. Open3D: A Modern Library for 3D Data Processing.

Funding sources

We thank the participants of the Vallecas Project and the staff of the CIEN Foundation. This

work was supported by the CIEN Foundation and the Queen Sofia Foundation, as well as by

a grant from the Spanish Ministry of Science and Innovation (PID2020-119302RB-I00) to

BAS. DF was supported by a Marie Skłodowska-Curie fellowship project PCI2021-122046-

2B, financed by the Spanish Ministry of Science and Innovation and the Spanish State

Research Agency MCIN/AEI/10.13039/501100011033 and the European Union

"NextGenerationEU"/PRTR. MGP was supported by a MAPFRE-Queen Sofia Foundation

scholarship. LZ was supported by a grant from the Alzheimer's Association (2016-NIRG-

397128) to BAS.

Supplementary Materials

Number of voxels in cluster	p (FWE-corr)	TFCE	p (unc)	x	y	z	AAL3 labels
2287	0.007	1337.9	0	-28	-18	-12	Hippocampus_L Extended to the left superior temporal gyrus
	0.008	1318.49	0	-40	-30	-8	Hippocampus_L
	0.008	1308.5	0	-26	-31	-8	Hippocampus_L
184	0.032	999.29	0.002	4	-3	12	Thal_VL_R
	0.038	959.07	0.001	4	-5	-2	Thal_VA_R
168	0.033	994.55	0.001	30	-19	-12	Hippocampus_R
	0.045	922.07	0.002	20	-23	-16	Parahippocampal_R
46	0.044	928.22	0.001	20	-67	22	Cuneus_R

	0.049	904.13	0.001	18	-59	16	Calcarine_R
64	0.046	915.85	0.002	52	-19	-12	Temporal_Mid_R
	0.047	911.21	0.002	44	-11	-18	Hippocampus_R
22	0.048	909.89	0.002	-24	18	-18	OFCpost_L
7	0.049	903.11	0.001	14	-28	-24	Cerebellum_3_R

570 **Supplementary Table 1. Coordinates of grey matter volume effects in the VBM**
571 **analysis.** The MNI coordinates for the global maximum and local maxima of each cluster are
572 indicated in mm for the three sections in space (x, y and z). Neuroanatomical labels from the
573 AAL3 atlas are indicated. P(FWE-corr): Family Wise Error corrected p-value, TFCE:
574 Threshold-Free Cluster Enhancement local spatial support, p (unc): uncorrected p-value.

Number of voxels in cluster	p (FWE-corr)	Main tracts
Positive correlation with FA values		
4917	0.025	IFOF, SLF, fornix, ATR_R
3123	0.036	IFOF, ILF, SLF, fornix, ATR_L
1310	0.036	IFOF, ILF, forceps major_R
359	0.047	ILF, forceps major_L

239	0.044	SLF_L
128	0.047	Corpus callosum
61	0.049	Forceps minor
8	0.05	Corpus callosum
Negative correlation with MD values		
9921	0.013	IFOF, ILF, SLF, ATR_R, forceps minor, corpus callosum
7467	0.015	IFOF, ILF, SLF, fornix, ATR_L, corpus callosum
3	0.05	Corticospinal tract
1	0.05	

576 **Supplementary Table 2. Brain clusters for correlation between episodic learning rate**
577 **and FA and MD values in the main analysis.** P(FWE-cor): Family Wise Error corrected p-
578 value. Neuroanatomical labels from the JHU white matter atlas are indicated.

Supplementary Figure 1. Grey matter volume correlates with learning rate in older adults with an alternative statistical model. For this model, age and education were introduced in the model whereas the delayed FCSRT score was removed. The positive correlation has been overlaid on a canonical T1 image (thresholded at p < 0.05 FWE-corr) to show a significant effect in A hippocampus bilaterally and B thalamus, right anterior (red), ventroanterior (green) and ventrolateral thalamic nuclei (blue) (thalamic ROIs in the inset come from the AAL3 atlas (Rolls et al., 2020). The coordinates of the sections are given in mm. L: left, R: right, A: anterior, P: posterior.

Supplementary Figure 2. Extensive network of white matter microstructure integrity is related to learning rate in older adults with an alternative statistical model. For this model, age and education were introduced in the model whereas the delayed FCSRT score was removed. A. Positive correlation between FA and learning rate (warm colours; p < 0.05 FWE-corr). B. FA effects overlaid on the fornix. C. Negative correlation between MD and learning rate (cold colours; p < 0.05 FWE-corr). D. MD effect overlaid on the fornix.

596
597