The making of the oral microbiome in Agta hunter-gatherers

Begoña Dobon[1,2+], Federico Musciotto[1,3+], Alex Mira[4,5], Michael Greenacre[6,7], Rodolph Schlaepfer[1], Gabriela Aguileta[2], Leonora H. Astete[8], Marilyn Ngales[8], Vito Latora[9,10,11], Federico Battiston[1,12], Lucio Vinicius[1,13#], Andrea B. Migliano[1,13#*], Jaume Bertranpetit[2#*]

[1]Department of Anthropology, University of Zurich; Zurich, Switzerland.

[2]Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra); Barcelona, Spain

[3]Dipartimento di Fisica e Chimica, Università di Palermo; Palermo, Italy.

[4]Department of Health and Genomics, Center for Advanced Research in Public Health, FISABIO Foundation; Valencia, Spain.

[5]CIBER Center for Epidemiology and Public Health; Madrid, Spain

[6]Department of Economics and Business, Universitat Pompeu Fabra & Barcelona Graduate School of Economics; Barcelona, Spain.

[7]Faculty of Biosciences, Fisheries and Economics, University of Tromsø; Norway.

[8]Lyceum of the Philippines University, Intramuros, Manila, Philippines.

[9]School of Mathematical Sciences, Queen Mary University of London; London, UK.

[10]Dipartimento di Fisica ed Astronomia, Università di Catania and INFN; Catania, Italy.

[11]Complexity Science Hub Vienna (CSHV); Vienna, Austria.

[12]Department of Network and Data Science, Central European University; Vienna 1100, Austria.

[13]Department of Anthropology, University College London; London, UK.

[+]These authors contributed equally to this work.

[#]These authors contributed equally to this work.

[*]Corresponding authors. Email: jaume.bertranpetit@upf.edu, andrea.migliano@uzh.ch

Abstract

Ecological and genetic factors have influenced the composition of the human

microbiome during our evolutionary history. We analyzed the oral microbiota of

the Agta, a hunter-gatherer population where part of its members is adopting an

agricultural diet. We show that age is the strongest factor modulating the

microbiome, likely through immunosenescence as there is an increase of

pathogenicity with age. Biological and cultural processes generate sexual

dimorphism in the oral microbiome. A small subset of oral bacteria is influenced

by the host genome, linking host collagen genes to bacterial biofilm formation. Our

data also suggests that shifting from a fish/meat to a rice-rich diet transforms their

microbiome, mirroring the Neolithic transition. All these factors have implications

in the epidemiology of oral diseases. Thus, the human oral microbiome is

multifactorial, and shaped by various ecological and social factors that modify the

oral environment.

Introduction

The composition and diversity of the human oral microbiota has been influenced by

several factors during our evolutionary history[1,2]. Some are intrinsic biological

characteristics of the host, such as age, sex, and genetic composition, while others such

as diet, drinking water sources, oral hygiene, lifestyle and social interactions are

external factors[2–4]. These factors modulate the physiological conditions of the oral

cavity and affect the composition and diversity of the oral microbiota. While the oral

microbiota is one of the most diverse sites in the human body and shows high variability

between individuals, it remains stable within individuals over time[5]. Little is known

about how the composition of the oral microbiome is modulated in populations adapted

to the hunting and gathering niche, where the fully mature oral biofilm microbiome can

be studied without the confounding effects of tooth brushing or professional dental

cleaning, similar conditions to how the human oral microbiome must have evolved in

the past[6].

To investigate the multiple ecological and genetic factors shaping the human oral

microbiome, we have analysed the oral microbiome of the Agta hunter-gatherers from

the Philippines. The Agta are predominantly hunter-gatherers (fishing, hunting, and

gathering)[7], and while their main source of animal protein is obtained by riverine and

marine spearfishing or by hunting, other activities such as inter-tidal foraging, wild food

gathering, low-intensity cultivation, wage labour and trade complement their

economy[8,9]. Interestingly, there is high variability among the Agta on the amount of

hunting, gathering and sea foraging products that is traded for rice and other items (such

as tobacco) with farming neighbours[7], causing the Agta lifestyle to range from

completely mobile foragers with a protein-rich diet, to settled low intensity farmers,

with a rice-rich diet[7,9,10].

To detect fine-scale variation in the oral microbiome of Agta hunter-gatherers, we

collected saliva samples from 138 Agta, aged 5 to 65 years, sequencing the 16S rRNA

region, and identified 5430 amplicon sequence variants (ASVs)[11] belonging to 110

genera. To study the genetic host factors associated with the microbiome composition

we also genotyped all individuals with the Axiom Genome-wide Human Origins array.

We combined this information with additional individual data on household

composition, age, sex, and diet, measured as the proportion of meals including meat/fish

(any animal protein), and proportion of meals that consisted of only agricultural

products (rice) (Supplementary Figure 1). By using this rich dataset, we have been able

to discern the different contributions of age, sex, diet, and host genetics in the making of

the oral environment.

Results

Factors influencing Agta oral microbiome composition. The Agta oral microbiome is

mostly composed by *Firmicutes* (mean ASV prevalence = 33.1%, sd = 11.4),

Proteobacteria (27±14.6%), *Actinobacteria* (15.5±8.3%) and *Bacteroidetes*

(14.5±7.7%) (Supplementary Figure 2). To compare and identify the main ecological

and social factors contributing to microbiome variation, we performed a constrained

logratio analysis (LRA)[12] on the bacterial genus abundance. Marginally, age explains

7.2% of the total logratio variance (P < 0.0001, based on 9999 permutations), sex

explains 2.2% (P = 0.018), and diet 3.6% (P = 0.015). Altogether they explain 13.0% of

the total logratio variance. We also applied a bipartite stochastic block model (biSBM)

approach[13] at the ASV level, where we assigned each bacterium to an individual, and

then clustered the individuals according to the bacteria they have in common. We

restricted the analysis to the Core Measurable Microbiota (CMM), that we define as

ASVs present in at least 10% of the Agta to reduce random errors due to low-prevalent

taxa (Supplementary Figure 3). The best model produced two clusters of people and

three clusters of bacteria (Figure 1a). While we did not find differences in diet or

proportions of sexes between the clusters of individuals (Supplementary Figure 4), they

strongly differ in their age distribution: adults (mean age = 38 years old) and youth

(mean age = 18 years old) (Welch t-test, t = 5.78, df = 71.35, P < 0.0001), with 55.48%

of ASVs in the CMM being more associated with one of the clusters of individuals.

Thus, while age, diet, and sex influence the composition of Agta microbiome, the

biSBM singles out age as the main modulator of the hunter-gatherer core oral

microbiome.

Figure 1. Age and sex-related effects in the hunter-gatherer oral microbiome. a)
Network representation of the hunter-gatherer CMM. ASVs (triangles) are colour-coded
as: putatively pathogenic (purple), non-pathogenic (orange) or unclassified (white).
Inset shows age distribution for the two clusters of individuals (squares). b) Logratio
analysis constrained to age and sex differences on the bacterial composition at genus

level. The effects of diet were partialed out. Only genera statistically significant in at
least 20 (for age) or 10 (for sex) logratios are displayed (p-value < 0.05 after Benjamini-
Hochberg correction). Dashed lines enclose all individuals (dots) within a category of
sex, with 95% confidence ellipses for their means. Taxa are colour-coded depending on
the associated variable: age, sex, or both. The starting point of the grey arrow indicates
the mean age of the population (30 years old). Log ratio of c) *Haemophilus* and
Selenomonas abundance and d) *Moraxella* and *Bacteroides* according to age. Line and
shaded area indicate the 95% confidence interval of the mean. Relative abundance of e)
Bifidobacterium and f) *Comamonas* according to age and sex. Lines and shaded areas
indicate the 95% confidence interval of the mean for each sex.

Old age is associated with increased frequency of oral pathogens. To investigate the

independent effects of ageing on the oral microbiome, we performed a LRA constrained

with age and sex after partialing out the effects of diet. The resulting ordination shows

that the effects of age and sex are mostly independent, with only few genera being

affected by both variables (Supplementary Figure 5): as expected, the first dimension is

associated with the age of the individuals, while the second dimension separates them

according to sex (Figure 1b).

There is a clear change in the composition and frequency of certain bacteria with

age (Figure 1c-d). At young age, we observe organisms that typically live in mucosa,

such as *Haemophilus* and *Moraxella*, that infect the upper and lower respiratory tract

but are detected in the oral cavity and saliva which are their vehicles of transmission.

Other genera found at younger ages include bacteria normally associated to good oral

health, such as *Bergeyella* and *Rothia*[14]. However, at older ages we observe a marked

decline in the abundance of those genera and an increase of important pathogens related

with periodontitis including the "red complex" periodontal pathogen *Tannerella*, as well

as other periodontitis-related bacteria (*Filifactor*, *Fretibacterium, Saccharimonas*,

Selenomonas, and *Phocaeicola*), consistent with a higher incidence of this disease with

older age[15]. We also found organisms associated with cavities (*Olsenella*), with dental

plaque and dental calculus formation (*Corynebacterium*), with pulmonary infections,

sepsis, or bacteremia, and with chronic diseases (*Acholeplasma*) (see Methods for in-

depth bacteria pathogenic classification). Another sign of ageing was the presence in the

oral cavity of gut bacteria (*Bacteroides*) indicating a potential age-related decline in

immunological function and filtering[16]. However, such changes are not associated with

a decrease in the alpha-diversity of the total oral microbiome as measured by the

number of bacteria observed or their phylogenetic complexity (Supplementary Figure

6a-b), suggesting that the overall effect of aging is a replacement of protective and

commensal bacteria by pathogenic ones. This is supported by an increase in the number

of potential pathogenic bacteria in the CCM in bacterial clusters associated with older

ages (Fisher exact test, $P < 0.001$) (Figure 1a).

Sex differences shape composition but not diversity of the Agta oral microbiome.

We found no differences in alpha-diversity in the Agta oral microbiome between males

and females (Supplementary Figure 6c-d), which may be explained by sex equality

within the Agta hunter-gatherer society regarding diet and social interactions[17,18].

Nevertheless, the LRA constrained to age and sex shows sex-related differences in the

composition of the oral microbiome (Figure 1b). For example, *Stomatobaculum* and

Eubacterium yurii, present in the oral cavity of smokers[19], are associated with males,

consistent with Agta men chewing tobacco more frequently than women. It is also

interesting to mention *Comamonas* (Figure 1f), which even if it has been reported as a

possible contaminant in microbiome studies[20], its presence in females could be related

to its capacity of degrading the female hormone progesterone[21]. This bacterium has

been found in subgingival samples, where female hormones could be present either in

saliva or in gingival crevicular fluid.

Age and sex interactions in microbiome composition. Some bacteria are significantly

associated to both age and sex-related differences, such as *Gemella*, which is a prevalent

inhabitant of the respiratory mucosa such as *Haemophilus* and *Moraxella*, supporting

the idea that mucosa-associated and/or respiratory-tract organisms are more frequently

acquired in young individuals, especially males. At older ages, the

Bifidobacterium/*Saccharibacteria* ratio distinguishes between sexes: while

Bifidobacterium is associated to females, the periodontal pathogen *Saccharibacteria* is

associated to males. Thus, the observed trend of increase of periodontal pathogens with

age is stronger in males, as expected by the global epidemiology of the disease[22,23]. On

the other hand, we found an increase of the caries-related pathogens *Scardovia* and

Bifidobacterium associated to reproductive age females. Caries incidence increases with

age and is more prevalent in females[24], a more saccharolytic or acidic salivary

environment in older women, together with hormonal fluctuations and lower salivary

flow[25] could facilitate the proliferation of saccharolytic bacteria. The strong association

of *Bifidobacterium* with adult females could also be explained by its presence in

breastmilk[26]. Also, its proliferation coincides with the start of the reproductive age and

the increase of childcare[10] (Figure 1d).

Influence of variation in rice consumption on the Agta oral microbiome. While the

impact of diet on gut microbiome has been clearly established[27–30], its role in the oral

microbiome is still unclear. Some studies have found little or no effect, whereas others

have found associations with specific nutrients[2,25,31,32]. The variation in rice

consumption in the Agta allows us to assess both the relationship between the hunter-

gatherer diet on the oral microbiome and the effects of the recent introduction of

farming products. We performed a LRA on the bacterial genus abundance after

partialing out the effects of age and gender (Figure 2). The first dimension of the

ordination shows the gradient of the transition from a diet where all meals include meat

to where most consist of only rice. Agta following a hunter-gatherer diet, where most

meals contain meat, have large quantities of *Actinobacillus*, *Alphaproteobacteria* and

Streptobacillus and lower abundance of *Selenomonas*, *Atopobium*, *Peptoanaerobacter*,

and *Pyramidobacter*. The higher abundance of *Actinobacillus* in individuals ingesting a

protein-rich diet is particularly interesting, given the extraordinary proteolytic potential

of *A. actinomycetencomitans*, a well-known oral pathogen with destructive effects in the

gingival tissue and in aggressive forms of periodontitis[33]. At the other extreme, in

individuals with a rice-rich diet, there is an increase of the highly saccharolytic dental

caries pathogen *Scardovia*, of *Treponema*, of gut organisms like *Butyrivibrio* and

Erysipelotrichaceae, and of *Eggerthia*, a rare organism isolated from dental abscesses.

We also ranked the ASV based on whether they are more present than expected in

individuals with high or low proportion of meals with only rice or with meat/fish. We

found that the scores associating each bacterial species with these two nutrients are

negatively correlated (Spearman's rho = -0.47, P < 0.0001). This fits with a general

separation of oral microorganisms in saccharolytic (caries-related, acidogenic and

acidophilic) and proteolytic (gum-disease and halitosis related, alkalophilic and NH_4

generators), as suggested in a metabolome-based study[25]. Our results suggest that more

settled Agta, which consume more rice, experience a decline in oral health, confirming a

general pattern of health decline due to a Neolithic-like diet and a more farming-derived

lifestyle[9,34,35].

Figure 2. Effect of diet on the oral microbiome in the Agta. Logratio analysis
constrained to diet differences on the bacterial composition at genus level. The effects
of age and sex were partialed out. Only genera statistically significant in more than five
(for rice) or three (for meat) logratios are displayed (p-value < 0.05 after Benjamini-
Hochberg correction). Taxa are colour-coded based on the variable they are associated:
proportion of meals with meat (%Meat), proportion of meals with only rice (%Rice), or
both. The original result was slightly rotated so that the dashed vector indicating the
difference between %Meat and %Rice was horizontal, without any change in explained
variance.

Pathogenic oral bacteria are associated with host collagen genes. The interaction

between the host genetic makeup and microbiome composition differs across body

sites[36,37], and seems especially weak in the oral cavity[37,38], making it difficult to assess

the co-evolution of our genome and the oral microbiome. To overcome this, we

performed a genome-wide association study (GWAS) using a mixed model approach in

a population that evolved in a hunter-gatherer niche and without the confounding

influence of antibiotics or brushing. We treated the relative abundance of each

bacterium as an independent trait, adding age, sex, and household as covariates and

kinship as a random effect. Household membership was used as proxy for the strength

of social interactions between individuals, as social interactions predict microbiome

sharing (Musciotto *et al. companion paper*). These analyses were performed using the

CMM, and then using 92 genera present in at least 10 Agta. All bacteria identified in the

Agta (Supplementary Table 1 and Supplementary Figure 7) overlap with those of other

oral microbiome GWAS[3,36,37] pointing to a small subset of oral bacteria influenced by

the human genome (Figure 3). A pathway enrichment analysis linked this subset of the

oral microbiome to several biological host functions (body fat metabolism, wound

healing, and collagen trimmers) (Supplementary Table 2). Of relevance is an association

between the pathogenic bacteria *Aggregatibacter* and *Selenomonas* with genetic

variation in collagen genes. The ability to bind collagen is a vital feature in the oral

cavity, as many oral bacteria require collagen-binding proteins to attach to oral tissues[39]

suggesting a genetic basis for the predisposition of biofilm formation by those bacteria.

We further tested if we could detect signatures of positive selection in the host genomic

regions associated with the oral microbiome, but we found no signals indicating recent

selective pressures caused by oral bacteria.

Figure 3. Genome-wide association study on bacteria abundance. Aggregated
Manhattan plot of the GWAS results of the a) seven ASV and b) eight genera with non-
zero PVE ("chip heritability") estimates with at least one significant genetic association.
Each dot is a SNP and significant SNPs-bacteria associations (q < 0.1) are color-coded
according to the associated bacteria.

Discussion

The Agta microbiome is influenced by external factors such as social interactions

(Musciotto *et al. companion paper*) as well as intrinsic and ecological factors such as

age, sex, diet and host genetics. Among the latter, we have shown that age has the

strongest effect, with commensal or beneficial microbiota being replaced by potentially

pathogenic ones with ageing. The proliferation of oral pathogenic bacteria exhibits

sexual dimorphism, with caries-related (in females) and periodontitis-related (in males)

bacteria increasing with age, likely associated with immunosenescence[40] and with a sex-

specific oral environment due to biological and cultural factors. In the Agta, the increase

of farming-derived novel foods such as rice influences their microbiome composition

and health. The relatively small subset of bacteria linked to the host genome, which are

also found associated to other factors, suggests that the Agta oral microbiome is mainly

affected by environmental (diet) and intrinsic factors (age), with little influence of

individual host genetic variation (Supplementary Figure 5). Thus, environmental factors

and not host genetics are the main driving force for oral microbiota acquisition, in

agreement with Mukherjee *et. al*[41]. Based on the case study of the Agta hunter-

gatherers, we conclude that the human oral microbiome is multifactorial with distinct

subsets of bacteria shaped by specific ecological and social factors, reflecting multiple

adaptations in the domains of life history, sociality, and diet.

Methods

Ethics approval

This study was approved by UCL Ethics Committee (UCL Ethics code 3086/003) and

carried out with permission from local government and community members. Informed

consent was obtained from all participants, after group and individual explanation of

research objectives in the indigenous language. A small compensation (usually a

thermal bottle or cooking utensils) was given to each participant. The National

Commission for Indigenous Peoples (NCIP), advised us that the process of Free Prior

Informed Consent with the tribal leaders, youth and elders would be necessary to

validate our data collection under their supervision. It was done in 2017 with the

presence of all tribal leathers, elders and youth representatives at the NCIP regional

office, with the mediation of the regional officer and the NCIP Attorney. The validation

process was approved unanimously by the tribal leaders, and the NCIP, and validated

the full 5 years of data collection.

Saliva sample collection

Saliva samples from 155 Palanan Agta were collected over two field seasons: April-

June 2013 and February-October 2014. For comparative genetic studies we also used

saliva samples from 21 Mbendjele Bayaka, an African hunter-gather population,

collected in 2014, and 14 Palanan farmers collected in 2007-2009. In all cases saliva

was collected using the Oragene·DNA/saliva kit and participants were asked to rinse

their mouth with water and to spit into the vial until half full. After collection and

transportation, saliva samples were stored at the UCL Department of Anthropology,

London, UK at -20ºC.

Microbial DNA extraction and 16S rRNA gene sequencing

A total of 155 Agta saliva samples were selected to study their microbiome

composition. DNA was extracted following the protocol for manual purification of

DNA for Oragene·DNA/saliva samples. The 16S rRNA gene V3-V4 region was

amplified by PCR with primers containing Illumina adapter overhang nucleotide

sequences. All PCR products were validated through an agarose gel and purified with

magnetic beads. Index PCR was then performed to create the final library which was

also validated through an agarose gel. All samples were pooled together at equimolar

proportions and the final pool was qPCR quantified prior to the MiSeq loading. Raw

Illumina pair-end sequence data were demultiplexed, quality filtered and denoised with

QIIME 2 2019.1[42] and DADA2[43]. DADA2 generates single nucleotide exact amplicon

sequence variants (ASVs). ASV are biological meaningful entities as they identify a

specific DNA sequence and allow for higher resolution than using operational

taxonomic units (OTUs)[11]. Taxonomic information was assigned to ASVs using a naïve

Bayes taxonomy classifier against SILVA database release 132 with a 99% identity

sequence[44]. ASVs that did not belong to the kingdom Bacteria, or that were classified as

mitochondrial or chloroplast sequences and samples with an extremely low number of

sequences (8000) were excluded from further analyses. ASVs were aligned with

MAFFT[45] and a rooted phylogenetic tree was constructed with FastTree2[46] using default

settings via QIIME 2. This resulted in a total of 5430 ASVs and 138 Agta (67 women

and 71 men). We generated a rarefaction curve with R package vegan (version 2.5-7)[47]

to determine that the richness of the samples had been fully observed (Supplementary

Figure 8). The number of observed ASVs and Shannon Diversity index were calculated

with R package Phyloseq (version 1.30.0)[48]. Faith's Phylogenetic Diversity index[49] was

calculated with R package picante (version 1.8.2)[50] using the rooted phylogenetic tree

generated in R[51]. To determine the set of microbial traits to be included in the analyses,

we selected ASVs with at least 10 reads in at least 2 individuals (n = 1980), then we

aggregated those with a taxonomic assignment at a genus level, resulting in 110 genera.

At the ASV level we also defined a Core Measurable Microbiota (CMM), consisting of

ASVs that appear in at least 10% of the Agta individuals (14 or more) resulting in 575

ASVs (out of 1980) that represent 90% of the composition of the Agta microbiome.

Genotype data

A total of 190 saliva samples were genotyped with the Affymetrix Axiom Genome-

Wide Human Origins 1 array. DNA extraction was carried out following the protocol

for manual purification of DNA for Oragene·DNA/saliva samples in the same

laboratory that sequenced the 16S rRNA data. Samples were analyzed with Axiom

Analysis Suite v4.0 following the Axiom genotyping best-practices workflow for saliva

samples. 618810 markers and 177 samples passed initial quality control. Single

nucleotide polymorphisms (SNPs) with less than 95% genotyping rate and samples

where the estimated gender from the genotypes did not match the recorded gender were

excluded from the analysis. Duplicated samples were identified with KING[52] and

removed. This resulted in a total of 617063 markers and 174 samples: 141 Agtas, 19

Bayaka and 14 Palanan farmers.

Ethnographic data collection

Ethnographic data collection occurred over two field seasons from April-June 2013 and

February-October 2014. In the first season we censused 915 Agta individuals (54.7%

which were male) across 20 camps, collecting basic information on household

composition, sex, and estimated ages. Following relative aging protocols[53], accurate

ages were established for all individuals post data collection.

Diet data collection

Dietary recall data was collected at the household level over a period of 10 days. We

asked the mother and the father at the end of the day (between 17:00 – 18:00) what

foods they had eaten that day, including agricultural produces from trade with nearby

farmers. We counted the total amount of meals we had recorded for a household and

established what proportion of these consisted of meat, vegetables, fruits, honey, and

rice. Therefore, this is only a rough guide to dietary composition and does not take

calorific intake or absolute weighs of the different food types into account. Dietary data

for 80 individuals (37 males and 43 females) was annotated based on the proportion of

meals that consisted of only rice, and the proportion of meals that included meat

(primarily fish and other marine resources and game).

Classification of oral bacteria as pathogens

Bacteria were classified as potential oral pathogens if they have been reported as

etiological agents of periodontitis or dental caries. Assignment as periodontal pathogen

was performed according to the systematic review of Perez-Chaparro *et al.*[54] and

Socransky *et al.*[55], or if they have been previously associated with this gum disease[56,57].

Bacteria were classified as caries pathogens if they were described in transcriptomic

studies of human cavities, according to Simon-Soro *et al.*[58] and Simon-Soro and Mira[59],

previously associated with caries[60,61], with cavities[62] or with dental plaque and dental

calculus formation[63,64]. Bacteria reported as etiological agents of respiratory infections

and biofilm-mediated infections were also considered pathogens, including organisms

that can be present in healthy carriers. These included species described in Leung *et*

al.[65], Bellussi *et al.*[66], and Natsis and Cohen[67]. Bacteria causing urinary tract infection

or sexually transmitted diseases which can transiently be found in the oral cavity were

also considered as potential pathogens and included microorganisms described in Lanao

et al.[68] and Jung *et al.*[69]. Common oral commensals potentially causing endocarditis or

systemic infections in immunocompromised patients were not considered pathogens. If

a bacterium was isolated from the oral cavity of an animal, it was considered an oral

inhabitant for the sake of our classification. If taxonomic classification in our dataset

could be assigned at the genus level only, it was considered a pathogen if: i) > 90 of

species within the genus were pathogenic, or ii) it included a major pathogenic species

but the rest of species within the genus were not oral inhabitants, according to the

Human Oral Microbiome Database (http://www.ehomd.org/)[70]. Bacteria with taxonomic

assignments at higher levels than genus (family, order, class) were excluded from this

analysis.

For assignment of bacteria to pathogenic or non-pathogenic, we used species-

level ASVs, given that there are multiple cases where different species from the same

genus had a different assignment. If taxonomic classification of the ASV was only

possible at the genus level, it was considered a pathogen if: i) >90% of named species

within the genus were pathogenic, or ii) the genus included a major pathogenic species

but the remaining species within the genus were not classified as oral by the Human

Oral Microbiome Database[70]. ASV with a top hit to a sequence classified as "Oral taxa"

in databases but without a species assignment were not considered named species and

were discarded from the analysis. Cases where taxonomic classification of the ASV was

only possible at the family level or higher were also discarded.

Multivariate compositional data analysis on microbial composition

We performed a constrained logratio analysis (LRA) using the package easyCODA[12] in

R[51] on the Agta oral microbiome at the genus level using as constraining covariates age

(as continuous variable), sex (male and female), and diet (both proportion of meals with

meat and proportion of meals with only rice). The microbiome abundance counts of

each Agta individual were treated as compositional data[71] and transformed to logarithms

of ratios (logratios). Constrained LRA is a special case of redundancy analysis[12] where

the total logratio variance is decomposed into parts explained by the covariates (the

"constrained variance") and a residual part (the "unconstrained variance", unrelated to

the covariates). Then, the ordination resulting from the LRA explains a maximum of the

constrained variance in a reduced two-dimensional solution. The statistical significance

of the three covariates was assessed using a multivariate permutation test (999

permutations) in the R package vegan[47]. There is no correlation between these three

covariates, except within the diet covariate, where the two variables are negatively

correlated (Spearman's rho = -0.54, P < 0.0001) (Supplementary Figure 1). To focus on

the genera affected only by internal factors (age and gender), we performed a

constrained LRA on the microbial composition after partialing out the effects of diet.

Similarly, to identify genera affected exclusively by the diet, we performed a

constrained LRA after partialing out the effects of age and gender. Taxon-covariate

association was ranked by counting the number of significant logratios for each of the

taxa, with p-value < 0.05 controlling for the false discovery rate (FDR) at level α =

0.05[72,73].

Community detection

To model the relationship between the Agta and the CMM, we used a stochastic block

model (SBM) approach specifically suited for bipartite networks[13]. SBM infers the

community structure[74] that better fits the existing graph, by building a prior distribution

for edges that holds no information on real data and using it in the framework of

Bayesian inference (biSBM) to find a partition of the two types of nodes whose

associated entropy is maximal. In this framework, the absence of links between nodes of

the same type or set is not considered informative for the model, as it is expected given

the bipartite nature of the graph, different from the general version of SBM. We selected

the number of clusters in the two sets that minimize the description length[75]. Robustness

of the clustering was assessed by calculating the average Adjusted Rand Index (ARI)

between iterations (n = 100), finding a mean ARI on Agta = 0.90 and a mean ARI on

ASV = 0.70. ARI measures the similarity of two partitions against a null hypothesis of

random assignment maintaining the size of the different clusters; the closer to 1 the

more robust is the classification[76]. The resulting clusters were plotted with graph-tool[77].

Ranking of bacteria associated with diet

ASVs present in the Agta were ranked from -1 to 1 based on whether that ASV is more

present than expected in individuals from a given category: low proportion versus high

proportion of meals with only rice, and low proportion versus high proportion of meals

with meat, based on the median value of the population for each variable. Thus, a meat

associated score towards 1 indicates that an ASV is present more than expected in

individuals with high proportion of meals with meat (above the median of the

population), and a score towards -1 indicates that is present more in individuals with

low proportion of meals with meat (below the median of the population). A rice

associated score towards 1 indicates that an ASV is present more than expected in

individuals with high proportion of meals that consist of only rice, and a score towards -

1 indicates that is present more in individuals with low proportion of meals with only

rice.

Microbiome genome-wide association studies

To study the relationship between host genetics and the microbiome in the Agta, we

used a genome-wide association study (GWAS) approach to identify specific SNPs

associated with microbial abundance using GEMMA (version 0.94)[78]. GWAS were

performed using the relative abundance of a given taxon in the Agta as a phenotype

trait, adding as covariates age, sex, and household (as a proxy for diet and shared

environment, as members of the same household share a hearth and their food on daily

bases). A kinship matrix calculated by KING using identical by descent segment

inference[52] was included as random effects. For the GWAS analyses, we applied the

following quality control steps to the Agta genotypes. First, to detect ancestry outliers in

the dataset, we filtered the samples to keep only bi-allelic autosomal SNPs with Minor

Allele Frequency (MAF) > 5% and without missing data with PLINK 1.9[79]. This dataset

was pruned for linkage disequilibrium (LD) using --indep-pairwise 50 5 0.2, and we

performed a principal component analysis (PCA) with EIGENSOFT (version 7.2.1)[80] to

identify ancestry outliers and exclude them from the analysis (Supplementary Figure 9).

Second, per sample heterozygosity was calculated with PLINK and samples with

overall increased/decreased heterozygosity rates (±3 s.d. from the mean of the

population) were removed. A total of 129 Agta samples passed microbiome and

genotype data quality controls and were included in the microbiome GWAS analyses.

The analyses were done at the genus taxonomic level and on the CMM to study the

effect of host genetics at different taxonomic levels. Depending on the taxon analyzed,

as we included only samples with non-zero abundance, and SNPs with MAF < 10% and

with more than 5% missing data were excluded, the number of individuals tested ranged

from 10 to 129 and the number of SNPs tested ranged from 270569 to 313198 markers.

When we performed the GWAS at the genus level, we only included in the analysis 92

genera that are present in at least 10 Agta individuals to exclude low prevalent genera.

P-values were adjusted for multiple testing by FDR, and SNP-taxa associations were

considered significant at q-value < 0.1 on the cases where the proportion of variance in

the bacterial abundance explained by the genotypes (PVE or "chip heritability") was

non-zero. The proportion of variance in the phenotype (bacterial abundance) explained

by the genotypes tested (PVE or "chip heritability") was estimated for each taxon and

was considered non-zero if the standard error measurements did not intersect zero. We

applied genomic control to correct for cryptic relatedness and population stratification

and minimize false positives induced by inflated association test statistics[81]. To do so,

we estimated the genomic inflation factor as the median value of the likelihood ratio test

(LRT) values divided by 0.456 (median of a $\chi 2(1)$ distribution) and recalculated the p-

values after dividing the LRT values by the genomic inflation factor[82]. Threshold of

significance was set at FDR 10%, and only genomic positions having at least three

samples for the major homozygous genotype and for the heterozygous genotype were

considered. SNPs were annotated with ANNOVAR[83] in GRCh37 (hg19) using

RefSeqGene and dbSNP 147. For the enrichment analyses, we extracted the genes

associated to all non-intergenic SNPs and classified the genes in the background set,

that consisted in all genes present in the Axiom Human origin array; and the set to test,

that consisted of all genes that had a non-intergenic SNP significantly associated with

an ASV or a genus. We performed a gene ontology enrichment analysis with

ViSEAGO[84] and TopGo[85] R packages (Fisher exact test) and FUMA

GENE2FUNCTION module (Functional Mapping and Annotation of Genome-Wide

Association Studies)[86] to perform pathway enrichment analysis (hypergeometric test)

with FDR 5%.

Selection analyses

To test whether the GWAS SNPs showed any signal of recent positive selection we

performed a genome-wide scan of selection. We phased the Agta and Palanan farmer

populations independently. For each population, samples that were identified as

ancestry outliers by a PCA or with overall increased/decreased heterozygosity rates (± 3

s.d. from the mean of the population) were excluded from phasing. A total of 138 Agta

samples were phased using SHAPEIT2 version v2 (r900)[87] with the duoHMM method

to improve the phasing by integrating the known pedigree information. SNPs with

missing data were removed and window size was set to 5Mb for phasing. Due to the

small sample size, to phase the 14 unrelated Palanan farmers we used SHAPEIT2 with

default parameters and the 1,000 Genomes Phase 3 panel of haplotypes[88] as a reference

dataset. SNPs with missing data were removed. For the selection analyses, we excluded

one of each pair of related individuals by removing the sample in the pair with the

lowest call rate in the Agta phased dataset. This resulted in 38 unrelated Agta

individuals and 14 unrelated Palanan farmers. We ran the Integrated Haplotype Score

(iHS)[89] in the Agta phased dataset and the Cross-population Extended Haplotype

Homozygosity (XP-EHH) test[90] comparing the Agta against Palanan farmers as

implemented in selscan version v1.3.0[91] to identify signals of positive selection in

GWAS SNPs. Both tests were run with default parameters and with the genetic map

provided by the 1,000 Genomes Phase 3[88]. To identify regions under selection, for each

test we selected markers with scores in the 95th percentile that had at least 3 markers in

the 99th percentile in the surrounding area ($\pm$ 10 Kb). For iHS we used absolute values,

while only positive scores were analyzed for XP-EHH.

Acknowledgements

A.B.M was funded by Leverhulme Trust Grant RP2011-R 045 and PLP-2017-323. J.B.

received grant PID2019-110933GB-I00/AEI/10.13039/501100011033 from the

Agencia Estatal Investigación (AEI), Spain, grant GRC 2017 SGR 702 from Secretaria

d'Universitats i Recerca del Departament d'Economia i Coneixement de la Generalitat

de Catalunya, as well as grant CEX2018-000792-M, part of the "Unidad de Excelencia

María de Maeztu" funded by the AEI, which granted the microbiome analysis by the

Servei de Genòmica at Universitat Pompeu Fabra. A.M. is funded by grant RTI2018-

102032-B-100 from AEI (Spain).

Data availability

16S amplicon data (EGAS00001005317) are deposited at the European Genome-

phenome Archive (EGA), which is hosted at the EBI and the CRG. Genome data

generated in this study has been deposited at EGA under accession number

EGAS00001005315. Data at the individual level on age, household composition and

diet that support the findings of this study are available on request from the

corresponding authors (JB and ABM). The individual data are not publicly available

due to them containing information that could compromise research participant privacy.

Code availability

Source code and data for visualization are available at

https://doi.org/10.5281/zenodo.6342212

References

1. Cornejo Ulloa, P., van der Veen, M. H. & Krom, B. P. Review: modulation of the oral microbiome by the host to promote ecological balance. *Odontology* **107**, 437–448 (2019).

2. Weyrich, L. S. The evolutionary history of the human oral microbiota and its implications for modern health. *Periodontol. 2000* **85**, 90–100 (2021).

3. Gomez, A. *et al.* Host Genetic Control of the Oral Microbiome in Health and Disease. *Cell Host Microbe* **22**, 269-278.e3 (2017).

4. Willis, J. R. *et al.* Citizen science charts two major "stomatotypes" in the oral microbiome of adolescents and reveals links with habits and drinking water composition. *Microbiome* **6**, 218 (2018).

5. Costello, E. K. *et al.* Bacterial Community Variation in Human Body Habitats Across Space and Time. *Science (80-.).* **326**, 1694–1697 (2009).

6. Velsko, I. M. *et al.* Microbial differences between dental plaque and historic

dental calculus are related to oral biofilm maturation stage. *Microbiome* **7**, 102
(2019).

7. Page, A. E., Minter, T., Viguier, S. & Migliano, A. B. Hunter-gatherer health and
development policy: How the promotion of sedentism worsens the Agta's health
outcomes. *Soc. Sci. Med.* **197**, 39–48 (2018).

8. Minter, T. The Agta of the Northern Sierra Madre: Livelihood Strategies and
Resilience Among Philippine Hunter-gatherers. (Faculty of Social Sciences,
Leiden University, 2010).

9. Page, A. E. *et al.* Reproductive trade-offs in extant hunter-gatherers suggest
adaptive mechanism for the Neolithic expansion. *Proc. Natl. Acad. Sci.* **113**,
4694–4699 (2016).

10. Dyble, M., Thorley, J., Page, A. E., Smith, D. & Migliano, A. B. Engagement in
agricultural work is associated with reduced leisure time among Agta hunter-
gatherers. *Nat. Hum. Behav.* **3**, 792–796 (2019).

11. Callahan, B. J., McMurdie, P. J. & Holmes, S. P. Exact sequence variants should
replace operational taxonomic units in marker-gene data analysis. *ISME J.* **11**,
2639–2643 (2017).

12. Greenacre, M. *Compositional data analysis in practice*. (Chapman and
Hall/CRC, 2018).

13. Larremore, D. B., Clauset, A. & Jacobs, A. Z. Efficiently inferring community
structure in bipartite networks. *Phys. Rev. E* **90**, 012805 (2014).

14. Rosier, B. T., Moya-Gonzalvez, E. M., Corell-Escuin, P. & Mira, A. Isolation
and Characterization of Nitrate-Reducing Bacteria as Potential Probiotics for
Oral and Systemic Health. *Front. Microbiol.* **11**, 2261 (2020).

15. Kassebaum, N. J. *et al.* Global Burden of Severe Periodontitis in 1990-2010. *J.*
Dent. Res. **93**, 1045–1053 (2014).

16. De Maeyer, R. P. H. & Chambers, E. S. The impact of ageing on monocytes and
macrophages. *Immunol. Lett.* **230**, 1–10 (2021).

17. Dyble, M. *et al.* Networks of Food Sharing Reveal the Functional Significance of
Multilevel Sociality in Two Hunter-Gatherer Groups. *Curr. Biol.* **26**, 2017–2021
(2016).

18. Migliano, A. B. *et al.* Characterization of hunter-gatherer networks and
implications for cumulative culture. *Nat. Hum. Behav.* **1**, 0043 (2017).

19. Duan, X. *et al.* Smoking May Lead to Marginal Bone Loss Around Non-
Submerged Implants During Bone Healing by Altering Salivary Microbiome: A
Prospective Study. *J. Periodontol.* **88**, 1297–1308 (2017).

20. Eisenhofer, R. *et al.* Contamination in Low Microbial Biomass Microbiome
Studies: Issues and Recommendations. *Trends Microbiol.* **27**, 105–117 (2019).

21. Liu, S., Ying, G.-G., Liu, Y.-S., Peng, F.-Q. & He, L.-Y. Degradation of
Norgestrel by Bacteria from Activated Sludge: Comparison to Progesterone.
Environ. Sci. Technol. **47**, 130829113920003 (2013).

22. Shiau, H. J. & Reynolds, M. A. Sex Differences in Destructive Periodontal
Disease: A Systematic Review. *J. Periodontol.* **81**, 1379–1389 (2010).

23.   Eke, P. I. *et al.* Update on Prevalence of Periodontitis in Adults in the United
States: NHANES 2009 to 2012. *J. Periodontol.* **86**, 611–622 (2015).

24.   Ferraro, M. & Vieira, A. R. Explaining Gender Differences in Caries: A
Multifactorial Approach to a Multifactorial Disease. *Int. J. Dent.* **2010**, 1–5
(2010).

25.   Zaura, E. *et al.* On the ecosystemic network of saliva in healthy young adults.
ISME J. **11**, 1218–1231 (2017).

26.   Fernández, L., Pannaraj, P. S., Rautava, S. & Rodríguez, J. M. The Microbiota of
the Human Mammary Ecosystem. *Front. Cell. Infect. Microbiol.* **10**, 689 (2020).

27.   David, L. A. *et al.* Diet rapidly and reproducibly alters the human gut
microbiome. *Nature* **505**, 559–563 (2014).

28.   Turnbaugh, P. J. *et al.* The Effect of Diet on the Human Gut Microbiome: A
Metagenomic Analysis in Humanized Gnotobiotic Mice. *Sci. Transl. Med.* **1**,
6ra14-6ra14 (2009).

29.   Schnorr, S. L. *et al.* Gut microbiome of the Hadza hunter-gatherers. *Nat.*
Commun. **5**, 3654 (2014).

30.   Smits, S. A. *et al.* Seasonal cycling in the gut microbiome of the Hadza hunter-
gatherers of Tanzania. *Science (80-.).* **357**, 802–806 (2017).

31.   Belstrøm, D. *et al.* Bacterial profiles of saliva in relation to diet, lifestyle factors,
and socioeconomic status. *J. Oral Microbiol.* **6**, 23609 (2014).

32.   De Filippis, F. *et al.* The Same Microbiota and a Potentially Discriminant
Metabolome in the Saliva of Omnivore, Ovo-Lacto-Vegetarian and Vegan
Individuals. *PLoS One* **9**, e112373 (2014).

33.   Fives-Taylor, P. M., Meyer, D. H., Mintz, K. P. & Brissette, C. Virulence factors
of Actinobacillus actinomycetemcomitans. *Periodontol. 2000* **20**, 136–67 (1999).

34.   Adler, C. J. *et al.* Sequencing ancient calcified dental plaque shows changes in
oral microbiota with dietary shifts of the Neolithic and Industrial revolutions.
Nat. Genet. **45**, 450–455 (2013).

35.   Sabbatani, S. & Fiorino, S. Dental worm disease. *Le Infez. Med.* **24**, 349–358
(2016).

36.   Kolde, R. *et al.* Host genetic variation and its microbiome interactions within the
Human Microbiome Project. *Genome Med.* **10**, 6 (2018).

37.   Blekhman, R. *et al.* Host genetic variation impacts microbiome composition
across human body sites. *Genome Biol.* **16**, 191 (2015).

38.   Shaw, L. *et al.* The Human Salivary Microbiome Is Shaped by Shared
Environment Rather than Genetics: Evidence from a Large Family of Closely
Related Individuals. *MBio* **8**, e01237-17 (2017).

39.   Mira, A., Artacho, A., Camelo-Castillo, A., Garcia-Esteban, S. & Simon-Soro, A.
Salivary Immune and Metabolic Marker Analysis (SIMMA): A Diagnostic Test
to Predict Caries Risk. *Diagnostics* **7**, 38 (2017).

40.   Preshaw, P. M., Henne, K., Taylor, J. J., Valentine, R. A. & Conrads, G. Age-
related changes in immune function (immune senescence) in caries and
periodontal diseases: a systematic review. *J. Clin. Periodontol.* **44**, S153–S177

(2017).

41.   Mukherjee, C. *et al.* Acquisition of oral microbiota is driven by environment, not host genetics. *Microbiome* **9**, 54 (2021).

42.   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

43.   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583 (2016).

44.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2012).

45.   Katoh, K. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002).

46.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately Maximum-Likelihood Trees for Large Alignments. *PLoS One* **5**, e9490 (2010).

47.   Oksanen, J. *et al.* Vegan: community ecology package. Ordination methods, diversity analysis and other functions for community and vegetation ecologists. *R Packag. version 2.5-7* (2020).

48.   McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217 (2013).

49.   Faith, D. P. Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* **61**, 1–10 (1992).

50.   Kembel, S. W. *et al.* Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* **26**, 1463–1464 (2010).

51.   R Core Team. R: A language and environment for statistical computing. (2020).

52.   Manichaikul, A. *et al.* Robust relationship inference in genome-wide association studies. *Bioinformatics* **26**, 2867–2873 (2010).

53.   Diekmann, Y. *et al.* Accurate age estimation in small-scale societies. *Proc. Natl. Acad. Sci.* **114**, 8205–8210 (2017).

54.   Pérez-Chaparro, P. J. *et al.* Newly Identified Pathogens Associated with Periodontitis. *J. Dent. Res.* **93**, 846–858 (2014).

55.   Socransky, S. S., Haffajee, A. D., Cugini, M. A., Smith, C. & Kent, R. L. Microbial complexes in subgingival plaque. *J. Clin. Periodontol.* **25**, 134–144 (1998).

56.   Camelo-Castillo, A. *et al.* Relationship between periodontitis-associated subgingival microbiota and clinical inflammation by 16S pyrosequencing. *J. Clin. Periodontol.* **42**, 1074–1082 (2015).

57.   Khemwong, T. *et al.* Fretibacterium sp. human oral taxon 360 is a novel biomarker for periodontitis screening in the Japanese population. *PLoS One* **14**, e0218266 (2019).

58.   Simón-Soro, A., Guillen-Navarro, M. & Mira, A. Metatranscriptomics reveals overall active bacterial composition in caries lesions. *J. Oral Microbiol.* **6**, 25443 (2014).

695   59.   Simón-Soro, A. & Mira, A. Solving the etiology of dental caries. *Trends*
696   *Microbiol.* **23**, 76–82 (2015).

697   60.   Tanner, A. C. R. Anaerobic culture to detect periodontal and caries pathogens. *J.*
698   *Oral Biosci.* **57**, 18–26 (2015).

699   61.   Kressirer, C. A. *et al.* Scardovia wiggsiae and its potential role as a caries
700   pathogen. *J. Oral Biosci.* **59**, 135–141 (2017).

701   62.   Wolff, D. *et al.* Amplicon-based microbiome study highlights the loss of
702   diversity and the establishment of a set of species in patients with dentin caries.
703   *PLoS One* **14**, e0219714 (2019).

704   63.   Mark Welch, J. L., Rossetti, B. J., Rieken, C. W., Dewhirst, F. E. & Borisy, G. G.
705   Biogeography of a human oral microbiome at the micron scale. *Proc. Natl. Acad.*
706   *Sci.* **113**, E791–E800 (2016).

707   64.   Ferrer, M. D. & Mira, A. Oral Biofilm Architecture at the Microbial Scale.
708   *Trends Microbiol.* **24**, 246–248 (2016).

709   65.   Leung, A. K. C., Wong, A. H. C. & Hon, K. L. Community-Acquired Pneumonia
710   in Children. *Recent Pat. Inflamm. Allergy Drug Discov.* **12**, 136–144 (2018).

711   66.   Bellussi, L. M. *et al.* An overview on upper respiratory tract infections and
712   bacteriotherapy as innovative therapeutic strategy. *Eur. Rev. Med. Pharmacol.*
713   *Sci.* **23**, 27–38 (2019).

714   67.   Natsis, N. E. & Cohen, P. R. Coagulase-Negative Staphylococcus Skin and Soft
715   Tissue Infections. *Am. J. Clin. Dermatol.* **19**, 671–677 (2018).

716   68.   Lanao, A. E. & Pearson-Shaver, A. L. Mycoplasma infections. *StatPearls*
717   *[Internet]* (2020).

718   69.   Jung, H.-S., Ehlers, M. M., Lombaard, H., Redelinghuys, M. J. & Kock, M. M.
719   Etiology of bacterial vaginosis and polymicrobial biofilm formation. *Crit. Rev.*
720   *Microbiol.* **43**, 651–667 (2017).

721   70.   Escapa, I. F. *et al.* New Insights into Human Nostril Microbiome from the
722   Expanded Human Oral Microbiome Database (eHOMD): a Resource for the
723   Microbiome of the Human Aerodigestive Tract. *mSystems* **3**, e00187-18 (2018).

724   71.   Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V. & Egozcue, J. J.
725   Microbiome Datasets Are Compositional: And This Is Not Optional. *Front.*
726   *Microbiol.* **8**, 2224 (2017).

727   72.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical
728   and powerful approach to multiple testing. *J. R. Stat. Soc. Soc. Ser. B* **57**, 289–
729   300 (1995).

730   73.   Storey, J. D. A direct approach to false discovery rates. *J. R. Stat. Soc. Ser. B*
731   *(Statistical Methodol.* **64**, 479–498 (2002).

732   74.   Fortunato, S. Community detection in graphs. *Phys. Rep.* **486**, 75–174 (2010).

733   75.   Peixoto, T. P. Nonparametric Bayesian inference of the microcanonical stochastic
734   block model. *Phys. Rev. E* **95**, 012317 (2017).

735   76.   Hubert, L. & Arabie, P. Comparing partitions. *J. Classif.* **2**, 193–218 (1985).

736   77.   Tiago, P. P. The graph-tool python library. *figshare* (2014).

737   doi:10.6084/m9.figshare.1164194

738 78. Zhou, X. & Stephens, M. Genome-wide efficient mixed-model analysis for
739   association studies. *Nat. Genet.* **44**, 821–824 (2012).

740 79. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger
741   and richer datasets. *Gigascience* **4**, 7 (2015).

742 80. Patterson, N., Price, A. L. & Reich, D. Population Structure and Eigenanalysis.
743   *PLoS Genet.* **2**, e190 (2006).

744 81. Devlin, B. & Roeder, K. Genomic Control for Association Studies. *Biometrics*
745   **55**, 997–1004 (1999).

746 82. Bacanu, S.-A., Devlin, B. & Roeder, K. The Power of Genomic Control. *Am. J.*
747   *Hum. Genet.* **66**, 1933–1944 (2000).

748 83. Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of
749   genetic variants from high-throughput sequencing data. *Nucleic Acids Res.* **38**,
750   e164–e164 (2010).

751 84. Brionne, A., Juanchich, A. & Hennequet-Antier, C. ViSEAGO: a Bioconductor
752   package for clustering biological functions using Gene Ontology and semantic
753   similarity. *BioData Min.* **12**, 16 (2019).

754 85. Alexa, A. & Rahnenfuhrer, J. topGO: Enrichment Analysis for Gene Ontology.
755   (2019).

756 86. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional
757   mapping and annotation of genetic associations with FUMA. *Nat. Commun.* **8**,
758   1826 (2017).

759 87. O'Connell, J. *et al.* A General Approach for Haplotype Phasing across the Full
760   Spectrum of Relatedness. *PLoS Genet.* **10**, e1004234 (2014).

761 88. Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–
762   74 (2015).

763 89. Voight, B. F., Kudaravalli, S., Wen, X. & Pritchard, J. K. A Map of Recent
764   Positive Selection in the Human Genome. *PLoS Biol.* **4**, e72 (2006).

765 90. Sabeti, P. C. *et al.* Genome-wide detection and characterization of positive
766   selection in human populations. *Nature* **449**, 913–918 (2007).

767 91. Szpiech, Z. A. & Hernandez, R. D. selscan: An Efficient Multithreaded Program
768   to Perform EHH-Based Scans for Positive Selection. *Mol. Biol. Evol.* **31**, 2824–
769   2827 (2014).