Manufacturing CD20/CD19-targeted iCasp9 regulatable CAR-T$_{SCM}$ cells using *qCART*, the *Quantum pBac*-based CAR-T system

Running Title: Manufacturing CAR-T$_{SCM}$ cells using *qCART*

Peter S. Chang[1,*], Yi-Chun Chen[1,*], Wei-Kai Hua[1,*], Jeff C. Hsu[1], Jui-Cheng Tsai[1], Yi-Wun Huang[1], Yi-Hsin Kao[1], Pei-Hua Wu[1], Yi-Fang Chang[2,3,4], Ming-Chih Chang[2], Yu-Cheng Chang[2,3], Shiou-Ling Jian[5], Jiann-Shiun Lai[5], Ming-Tain Lai[5], Wei-Cheng Yang[6], Chia-Ning Shen[6,7], Kuo-Lan Karen Wen[1,†], Sareina Chiung-Yuan Wu[1,†]

[1]GenomeFrontier Therapeutics, Inc. Taipei City, Taiwan (R.O.C.)

[2]Division of Hematology and Oncology, Department of Internal Medicine, Mackay Memorial Hospital, Taipei, Taiwan (R.O.C.)

[3]Department of Medical Research, Laboratory of Good Clinical Research Center, Mackay Memorial Hospital, Tamsui District, New Taipei City, Taiwan (R.O.C.)

[4]Department of Medicine, Mackay Medical College, New Taipei City, Taiwan (R.O.C.)

[5]OBI Pharma, Inc.

[6]Biomedical Translation Research Center, Academia Sinica, Taipei, Taiwan (R.O.C.)

[7]Genomics Research Center, Academia Sinica, Taipei, Taiwan (R.O.C.)

[*]These authors contributed equally to this work and their authorship order is interchangeable

[†]To whom correspondence may be addressed. Sareina Chiung-Yuan Wu, Kuo-Lan Karen Wen; 18F-1, No. 3, Park St., Nangang Dist., Taipei City 11503, Taiwan (R.O.C.); +886-2-26558766;

Email: sareina@genomefrontier.com, klwen@genomefrontier.com

Keywords: multiplex CAR-T | *piggyBac* | T$_{SCM}$ | transposon

List of Abbreviations:

aAPC, artificial antigen presenting cells; CAR, chimeric antigen receptor; CLL, chronic lymphocytic leukemia; CRS, cytokine release syndrome; CT, threshold cycle; DLBCL, diffuse large B-cell lymphoma; EFS, event-free survival; GF-CART01, patient-derived bicistronic CD20/CD19-targeted iCasp9 regulatable CAR-T cells; GMP, good manufacturing practice; GVHD, graft-versus-host disease; HL, Hodgkin lymphoma; ICANS, immune effector cell-associated neurotoxicity syndrome; iCasp9, inducible caspase 9; i.v., intravenously; MM, multiple myeloma; NHL, Non-Hodgkin lymphoma; PBMCs, peripheral blood mononuclear cells; PE, R-phycoerythrin; PFS, progression-free survival; *qCART*, *Quantum CART*; *qPB*, *Quantum pBac*; T_{CM}, central memory T cell; T_{EFF}, effector T cell; T_N, naïve T cell; T_{SCM}, T stem cell memory.

DECLARATIONS:

Ethics approval and consent to participate

Fresh/frozen peripheral blood mononuclear cells (PBMCs) from adult healthy donors and patients were obtained from Chang Gung Memorial Hospital (IRB approval #201900578A3) and MacKay Memorial Hospital (IRB approval #20MMHIS330e), respectively. *In vivo* mouse xenograft model studies were conducted at the Development Center for Biotechnology (Taiwan) using Taiwan Mouse Clinic IACUC (2020-R501-035) approved protocols.

Consent for publication

All authors consent to publication of this manuscript.

Availability of data and material

The authors confirm that the data supporting the findings of this study are available within the article and/or its supplementary materials.

Competing interests

S.C.-Y.W. is the founder of GenomeFrontier Therapeutics, Inc., P.S.C., Y.-C.C. (Chen), W.-K.H., J.C.H., J.-

C.T., Y.-W.H., Y.-H.K., P.-H.W., and K.-L.K.W. are affiliated with GenomeFrontier Therapeutics, Inc.

Funding

This study is funded by GenomeFrontier Therapeutics, Inc.

Authors' contributions

S.C.-Y.W. designed research. Y.-C. C. (Chen), J.-C.T., Y.-W.H., Y.-H.K., P.-H.W., Y.-F.C., M.-C.C., and Y.-C. C.

(Chang) performed research. P.S.C., W.-K.H., Y.-C. C. (Chen), J.C.H., K.-L.K.W., and S.C.-Y.W. analyzed

data. P.S.C., J.C.H., and S.C.-Y.W. wrote the paper.

Acknowledgements

The authors thank Ms. Jin-hwa Yang for her assistance with IRB preparation and approval process. The

authors also thank Dr. Pei-Yi Tsai for assistance with the animal experiments.

ABSTRACT

Background: CD19-targeted chimeric antigen receptor therapies (CAR19) have driven a paradigm shift in the treatment of relapsed/refractory B-cell malignancies. However, >50% of CAR19-treated patients experienced progressive disease mainly due to antigen escape and low persistence. Clinical prognosis is heavily influenced by CAR-T cell function and systemic cytokine toxicities. Furthermore, it remains a challenge to efficiently, cost-effectively, and consistently manufacture clinically relevant number of virally engineered CAR-T cells.

Methods: Using a highly efficient *piggyBac* transposon-based vector, *Quantum pBac*, we developed a virus-free cell engineering system, *Quantum CART (qCART™)*, for development and production of multiplex CAR-T therapies.

Results: Here, we demonstrated *in vitro and in vivo* that consistent, robust, and functional CD20/CD19 dual-targeted CAR-T stem cell memory (T_{SCM}) cells can be efficiently manufactured using the *qCART™* system for clinical application. *qCART™*-manufactured CAR-T cells from cancer patients expanded efficiently, rapidly eradicated tumors, and can be safely controlled via an iCasp9 suicide gene-inducing drug.

Conclusions: The *qCART™* system is an elegant system for the manufacturing of CAR-T products having all the desired CAR-T therapy attributes. We believe that the simplicity of manufacturing multiplex CAR-T cells using the *qCART™* system will not only significantly enhance the accessibility of CAR-T therapy but also unlock the full potential of armored CAR-T therapy for the treatment of solid tumors in the future.

What is already known on this topic

Despite the considerable success which has been achieved with CD19-targeted chimeric antigen receptor therapies (CAR19), >50% of CAR19-treated patients still experienced progressive disease. Therefore, there

is a need to further improve CAR19 therapies. Current CAR19 therapies commonly utilize virus-based cell

engineering methods. CAR-T production using these methods face multiple hurdles, including difficulties

to efficiently, cost-effectively, and consistently manufacture clinically relevant number of CAR-T cells. We

have previously used a highly efficient *piggyBac* transposon-based vector, *Quantum pBac,* to establish

Quantum CART (*qCART™*) which is a virus-free cell engineering system for development and production

of multiplex CAR-T therapies.

What this study adds

In this report, we further demonstrate *in vitro* and *in vivo* that consistent, robust, and functional iCasp9-

regulatable, CD20/CD19 dual-targeted CAR-T stem cell memory (T_{SCM}) cells can be efficiently

manufactured using the *qCART™* system for clinical application. These cells possess all the desired

attributes for ensuring therapeutic efficacy in CAR-T therapy, including high CAR-T_{SCM}, balanced CD8/CD4

ratio, low exhaustion and senescence marker expressions, and high *ex vivo* and *in vivo* expansion capacity.

Importantly, we show that *qCART™*-manufactured CAR-T cells from hematological cancer patients

expanded efficiently, effectively eradicated tumors, and can be safely controlled via an iCasp9 suicide

gene-inducing drug. We believe that the simplicity of manufacturing multiplex CAR-T cells using the

qCART™ system will not only significantly enhance the accessibility of CAR-T therapy but also unlock the

full potential of armored CAR-T therapy for the treatment of solid tumors in the future.

How this study might affect research, practice or policy

Our findings demonstrate that *qCART™* is a virus-free CAR-T engineering system for manufacturing CAR-

T_{SCM} cells from either healthy donors or hematological cancer patients, that possess all the desired

attributes for a successful CAR-T therapy. These cells expanded efficiently, rapidly eradicated tumors, and

can be safely controlled via activation of iCasp9. We expect that this simple yet robust system for

manufacturing multiplex CAR-T cells will advance the CAR-T field.

BACKGROUND

Over the last decade, chimeric antigen receptor (CAR)-T therapy has become one of the most promising cancer treatment, but the heavy reliance on lentiviral and retroviral vectors for CAR-T cell manufacturing makes the therapy expensive and inaccessible to the masses.[1] Frequent cytokine release syndrome (CRS) and immune effector cell-associated neurotoxicity syndrome (ICANS) associated with clinical patients treated with conventional CAR-T therapies have raised safety concerns.[2] Furthermore, virus-based gene therapies have limited gene payload capacity, making them less suitable for engineering multiplex armored CAR-T cells.[3,4]

To date, B-cell malignancies remain the most actively-studied cancers for CAR-T therapy development. Since *Brentjens et al. (2003)* first demonstrated the eradication of B-cell tumors by CD19-targeted second-generation CAR-T cells, several commercial CAR-T products have been approved by FDA.[5,6] However, treatment with CAR19 therapy resulted in only 30-40% long-term progression-free survival (PFS) in aggressive Non-Hodgkin lymphoma (NHL) patients, and the median event-free survival (EFS) of adult B-cell ALL patients was 6.1 months.[7,8] Evidence suggest that downregulation of CD19 may in part be responsible for patient relapse.[9,10] As demonstrated by recent clinical studies, bispecific CD19/CD20-targeted CAR-T therapy improved patient survival compared to CAR19 therapy.[11,12]

Transposon system, such as *Sleeping Beauty* and *piggyBac*, is a virus-free alternative for generating CAR-T cells but suffers from low gene transfer efficiency and limited expansion of engineered cells.[3,13,14] We recently developed the most advanced version of *piggyBac* known as *Quantum pBac (qPB)* that is 15 times more active than *hyperactive piggyBac* in human T cells.[15,16] We previously developed a novel genetic engineering system known as *Quantum CART (qCART™)* that combines efficient transgene design, evaluation, and identification of lead construct (*GTailor*); virus-free vector (*Quantum pBac*); effective gene delivery (*Quantum Nufect*); and optimized cell expansion (*iCellar*).[15,16] We reported that *qCART™* could be used to engineer highly potent T stem cell memory (T_{SCM}) CAR-Tcells. Here,

we further validate that the *qCART*™ system can be used to rapidly manufacture highly potent patient-

derived bicistronic CD20/CD19-targeted iCasp9 regulatable CAR-T cells (GF-CART01) for treating B-cell

malignancies. We demonstrate enhanced anti-tumor efficacy of GF-CART01 cells both *in vitro* and in a

xenotransplant immunodeficient mouse model.

METHODS

Cell engineering

CD19-expressing K562 cells were generated by using a *piggyBac*-based cell engineering

system. K562 cells were nucleofected using *Quantum Nufect*™ (GenomeFrontier Therapeutics)

with a plasmid carrying CD19 and hygromycin genes followed by hygromycin selection for 14

days. To select for clones that homogeneously express CD19, limiting-dilution was carried out to

reach a concentration of one cell/well. Following clonal expansion, the clone containing the

highest % of CD19$^+$ cells was identified and expanded for storage.

Generation/expansion of CAR-T cells

Frozen peripheral blood mononuclear cells (PBMCs) from adult healthy donors and

patients were obtained from Chang Gung Memorial Hospital (IRB approval #201900578A3) and

MacKay Memorial Hospital (IRB approval #20MMHIS330e), respectively. PBMCs were extracted

using Ficoll-Hypaque gradient separation (Cytiva). Primary human CD3$^+$ T cells were isolated from

PBMCs by negative selection using EasySep Human T Cell Isolation Kit (StemCell Technologies) or

by positive selection using Dynabeads Human T-Expander CD3/CD28 (Thermo Fisher), followed

by activation of isolated T cells with Dynabeads (3:1 bead-to-cell ratio) for two days. An iCasp9-

CD20/CD19-CAR transposon minicircle carrying CD20 scFv[17] and CD19 scFv[18] and a *qPBase*

plasmid were delivered into activated T cells using 4D-Nucleofector device (Lonza) and *Quantum*

Nufect Kit (GenomeFrontier Therapeutics) according to the manufacturer's instructions.

Irradiated artificial antigen presenting cells (aAPC), CD19-expressing K562 cells, were added to

electroporated T cells at 1:1 ratio on day 3. Cells were cultured in OpTmizer medium (Thermo

Fisher) supplemented with *Quantum Booster* (GenomeFrontier Therapeutics) in G-Rex 24- or 6M-

well plates (Wilson Wolf) or conventional plates for 10 days. For lentiviral CAR-T production,

cryopreserved human PBMCs were thawed and primary human CD3$^+$ T cells were isolated from

PBMCs by using human Pan T Cell Isolation Kit (Miltenyi Biotec, 130-096-535). Cells were

activated and expanded with Dynabeads at 1:1 ratio. After 2 days, cells were transduced with 1

MOI of concentrated virus containing 8 ug/mL Polybrene (Merck, TR-1003-G). Two days later,

Dynabeads were removed and the medium (supplemented with IL-7 and IL-15) was changed

every 2-3 days until day 10 when CAR-T cells were cryopreserved.

Cell culture conditions

Fed-batch: On day 1, approximately 1×10^5 nucleofected cells were seeded into a well of 24-well

G-Rex containing 3 ml of medium. Fresh medium (2ml) was added on days 3 and 5. On day 7,

culture was replaced with fresh medium (2ml).

Perfusion: On day 1, nucleofected cells were seeded into conventional 24-well plates at a

concentration of 2.5×10^5 cells/ml. Every 2 to 3 days, the cell numbers were counted, medium was

added to adjust the concentration to 2.5×10^5 cells/ml, and cells were placed back into culture

(depending on the volume, up to a 10 ml flask was used to culture these cells).

For both of these culture approaches, cells were harvested on day 10 to be used for subsequent

experiments.

Flow cytometry

CAR expression on T cells was analyzed by co-incubating cells with a Biotin-SP (long spacer) AffiniPure F(ab')$_2$ Fragment Goat Anti-Mouse IgG antibody (Jackson ImmunoResearch) for 30 min at 4°C and R-phycoerythrin (PE)-conjugated streptavidin (Jackson ImmunoResearch) for 30 min at 4°C. Expression of surface molecules of T cells were determined by labeling T cells with fluorochrome conjugated monoclonal antibodies: CD4-AlexaFlour532 (Thermo Fisher), CD3-Pacific Blue, CD3-PE-Cy5, CD4-AlexaFluor532, CD4-AlexaFlour700, CD8-BV605, CD8-PE-Cy7, CD8-Pacific Blue, CD27-PE-Cy7, CD28-APC, CD45RA-BV421, CD45RO-PE, CD62L-PE-Cy5, CD62L-PE-Dazzle 594, CD95-BV711, CD197-BV510, CD279/PD-1-PE/Cy7, CD366/Tim-3-BV650, CD223/LAG-3-BV711, KLRG-1-BV-421, CD57-PerCP-Cy5.5, CD19-PE, CD56-BV711 (BioLegend), CD45RA-BUV737, and CD95-BUV395 (BD Biosciences). Ghost dye Red 780 fixable viability dye (Cytek), 7AAD or propidium iodide (PI) was used to distinguish live cells from dead cells for data analysis. Mouse blood cell samples were stained with anti-hCD45-APC antibody (BioLegend). For counting the absolute cell number, CountBright™ Absolute Counting Beads (Invitrogen, C36950) were added into FACS samples before acquisition by flow cytometry. Flow cytometric data were acquired on an SA3800 Spectral Analyzer (Sony), a BD FACSCantoTM II (BD Bioscience), or a BD LSRFortessa analyzer (Sony). The graphs were generated using GraphPad Prism or FlowJo software (BD Biosciences or Tree Star).

Processing mouse blood samples

100μl of blood/mouse was collected in EDTA-containing tubes by submandibular blood collection. After centrifugation at 3000 x g for 15 minutes, plasma (upper layer) was collected for ELISA. The cell pellets/red blood cells (lower layer) were then resuspended with the RBC Lysis

Buffer (Biolegend, cat.420301) for 15 minutes at room temperature. Cell pellets were then centrifuged (350 x g for 5 minutes) and washed once with PBS. These cells were used for DNA extraction. For NCI-N87 tumor model experiments, submandibular blood was collected from mice in MiniCollect® tube (K3EDTA, Greiner) and RBC lysed by ACK buffer (Gibco). These cells were used for FACS analysis.

Enzyme-linked immunosorbent assay (ELISA)

For *in vitro* assay, supernatant was collected 24 hours after CAR-T and Raji-GFP/Luc cell (Creative Biogene, Cat. No: CSC-RR0320) co-culture (E:T=1:1). For *in vivo* assay, mouse plasma samples were collected on days 2, 5, 8, and 14 after T cell injection. IFN-γ (Thermo Fisher), TNF-α (Thermo Fisher), IL-6 (Thermo Fisher), and IL-2 (BioLegend) were quantified by ELISA according to manufacturer's instructions.

Genomic DNA extraction and quantitative PCR (qPCR)

Genomic DNA from mouse blood was extracted using a DNeasy Blood & Tissue Kit (Qiagen) following the manufacturer's instructions. For qPCR analysis, CAR-T cells and Raji-GFP/Luc cells were assessed using the following primer sequences: CAR: fwd-5'-ACGTCGTACTCTTCCCGTCT-3', rev-5'-GATCACCCTGTACTGCAACCA-3'; Luciferase: fwd-5'-GGACTTGGACACCGGTAAGA-3', rev-5'-GGTCCACGATGAAGAAGTGC-3'. All qPCR assays were performed using a 7500 fast real-time PCR system (Applied Biosystems). Absolute transgene levels were calculated using the standard curve method. The threshold cycle (CT) value of each target was converted to a concentration using the appropriate standard curve and calculating the % of cells in mouse blood based on the cell copy number.

In vitro cytotoxicity assay

Cytotoxicity was assessed using Celigo image cytometry (Nexcelom). Raji-GFP/Luc target

cells were seeded in 96-well culture plates. CAR-T cells were then added at E:T ratios of 5:1, 1:1,

and 1:5. At 0, 24, 48, 72, and 96 hours after cell co-culture, Raji-GFP/Luc cells were quantified.

The % specific lysis was calculated from each sample using the formula [1-(live fluorescent cell

count in target-effector cell co-culture/ live fluorescent cell count in target cell only culture)] x

100.

Anti-tumor efficacy in mouse tumor models

In vivo mouse xenograft model studies were conducted at the Development Center for

Biotechnology (Taiwan) using Taiwan Mouse Clinic IACUC (2020-R501-035) approved protocols.

Six to eight-week-old male immunodeficient (NOD.Cg-Prkdc^{scid}Il2rg^{tm1Wjl}/YckNarl) mice (National

Laboratory Animal Center, Taiwan) were engrafted with $1.5x10^5$ Raji-GFP/Luc tumor cells

intravenously (*i.v.*) or with Matrigel (1:1, BD Bioscience) containing $2x10^6$ NCI-N87-Luc cells

subcutaneously (*s.c.*) on the right flank. For Raji tumor model experiments, mice were infused

one week later with $7.5x10^5$ (low), $3x10^6$ (medium), or $1x10^7$ (high) CAR-T cells from donor PBMC-

41. After 35 days, $1.5x10^5$ Raji-GFP/Luc tumor cells were injected (*i.v.*) to half of the mice in each

group. Raji-GFP/Luc tumor cells were quantified using the Xenogen-IVIS Imaging System (Caliper

Life Sciences). For NCI-N87 tumor model experiments, mice were injected *i.v.* 11 days later with

$4x10^6$ control or CAR-T cells. Mice were imaged on an Ami-HT optical imaging system (Spectral

Instruments Imaging), following an intraperitoneal (*i.p.*) injection of D-Luciferin (BIOSYNTH, L-

8220)

***In vitro* iCasp9 safety assay**

255  Activation of inducible caspase 9 (iCasp9) was performed *in vitro* with 2.5, 5, or 10 nM of

256 dimerizer (AP1903), and CAR-T cell elimination was assessed by flow cytometry 24 hours later.

257 **ddPCR copy number assays**

258  Genomic DNA (gDNA) was extracted from GF-CART01 cells using a Qiagen DNeasy Blood

259 and Tissue Kit (Qiagen). Minicircle DNA (positive control) was produced by Aldevron. Before

260 ddPCR, the minicircle DNA was digested for 1 min at 37°C with BsrGI. The reaction for ddPCR

261 Copy Number Variation Assay was set up in 20 µl sample volume containing 10 µl of 2X ddPCR

262 Supermix (no dUTP, BioRad), 1 µl of FAM-labeled target primers/probe (BioRad), 1 µl of HEX-

263 labeled reference primers/probe (BioRad), 0.1 µl of BsrGI (Thermo Fisher), and 50 ng of gDNA or

264 minicircle DNA (volume 2 µl). Nuclease-free water was used to adjust the sample to final volume.

265 *RPP30* was used as housekeeping gene. BsrGI directly digests in the ddPCR reaction during setup.

266 Droplets were then generated in a QX200 droplet generator (BioRad) according to

267 manufacturers' instructions with a final volume of 40 µl. DNA amplification was carried out and

268 droplets were analyzed in a QX200 droplet reader (BioRad). Data was analyzed with QuantaSoft

269 software (BioRad).

270 **Statistical analysis**

271  Statistical analysis was performed using GraphPad Prism 7 software. Two-tailed unpaired

272 t-tests were used for comparison of two groups. One-way ANOVA was used for comparison of

273 three or more groups in a single condition. *P* values of less than 0.05 were considered statistically

274 significant, $*p < 0.05$, $**p < 0.01$, and $***p < 0.001$.

                                                RESULTS

Comparison of fed-batch (G-Rex) and perfusion (conventional)-cultured CAR-T cells

To assess the potential of *qCART™* system for CAR-T therapy development, we constructed a

second-generation bicistronic CAR (GF-CART01) containing binding domains that recognize CD20 and

CD19, human 4-1BB and CD3ζ signaling domains, and iCasp9 suicide gene driven by the EF1α promoter

(Figure 1A). Following nucleofection of GF-CART01 DNA, we expanded CAR-T cells in G-Rex (fed-batch)

and conventional (perfusion) plates. The percentage of CAR$^+$ T cells in the fed-batch-cultured (46%) and

perfusion-cultured (51%) CAR-T cells were comparable **(Figure 1B)**. However, we observed significantly

higher expansion of perfusion-cultured CAR-T cells (836-fold) compared to fed-batch-cultured CAR-T cells

(189-fold). Notably, both the perfusion- and fed-batch-cultured CAR-T cells had a high percentage of

CD45RA$^+$CD62L$^+$CD95$^+$ T$_{SCM}$ among the CAR$^+$ CD4 and CD8 populations. Despite noticeable lower

proliferation of fed-batch compared to perfusion-cultured CAR-T cells, we observed enhanced anti-tumor

cytotoxicity in fed-batch-cultured CAR-T cells **(Figure 1C)**. As G-Rex will be adopted for good

manufacturing practice (GMP) CAR-T manufacturing, all subsequent experiments were conducted in G-

Rex.

Safety assessment of GF-CART01 cells

To better manage CAR-T therapy-associated adversities, we included an iCasp9 suicide gene in

the GF-CART01 construct. We demonstrated that AP1903 treatment, which activates iCasp9, resulted in

a significant reduction in GF-CART01 cell survival **(Supplementary Figure S1A)**. To comply with GMP safety

standards for CAR-T manufacturing, we need to ensure that GF-CART01 cells have low integrant (CAR)

copy number per cell (<5). We have previously demonstrated that increasing the concentration of DNA

for electroporation increased CAR$^+$ T cells in a dose-dependent manner (data not shown). Here, we

modified CAR-T cells using different amounts of GF-CART01 DNA and determined that the CAR copy

number remained low (<5) when 15 μg of total DNA was used for nucleofection **(Supplementary Figure**

S1B). All subsequent experiments were conducted using 15 μg of DNA per 5×10^6 cells to maximize generation of CAR$^+$ T cells while maintaining safety levels of CAR DNA copy numbers.

GF-CART01 cells are dominated by T$_{SCM}$ cells with high proliferative capacity and low expression of exhaustion and senescence markers

We next examined the proliferative capacity of GF-CART01 cells generated from 9 independent healthy donors and monitored CAR expression on days 1, 8, and 10 following nucleofection. We observed a mean basal level of 21.49% CAR$^+$ T cell population on day 1, which subsequently increased and stabilized at 55.7% and 56.75% on days 8 and 10, respectively **(Figure 2A)**. As the presence of transposase (*Quantum PBase*, or *qPBase*) in modified T cells may lead to genotoxicity long-term, we also monitored the *qPBase* levels following nucleofection. While the *qPBase* levels were high on day 1 (47.31%), only a minimal (<0.2%) fraction of *qPBase*$^+$ cells was observed on days 8 and 10, suggesting that concerns regarding *qPBase-*induced integrant remobilization should be minimal **(Figure 2B).** We also observed an average of 178.86-fold GF-CART01 cell expansion **(Figure 2C)**.

Among the donors, we consistently observed a higher CD8 (73.58%) compared to CD4 (21.98%) CAR$^+$ population **(Figure 2D)**. To assess the composition of GF-CART01 cells, we determined the percentage of CD45RA$^+$CD62L$^+$CD95$^-$ naïve T cell (T$_N$), CD45RA$^+$CD62L$^+$CD95$^+$ stem cell memory T cell (T$_{SCM}$), CD45RA$^-$CD62L$^+$CD95$^+$ central memory T cell (T$_{CM}$), CD45RA$^-$CD62L$^-$CD95$^+$ effector memory T cell (T$_{EM}$), and CD45RA$^-$CD62L$^-$CD95$^-$ effector T cell (T$_{EFF}$) subsets in the CD4 and CD8 CAR$^+$ populations. We observed high percentage of CAR-T$_{SCM}$ cells in both the CD4 (68.94%) and CD8 (79.84%) populations compared to the other T cell subsets **(Figure 2E)**. Furthermore, these CAR-T cells displayed low exhaustion (PD-1, TIM-3, and LAG-3) markers and the senescence marker CD57 expression **(Figure 2F and G)**. Interestingly, a small but significant population (~9.92%) expressed the senescence marker KLRG-1.

GF-CART01 cells display robust cytotoxic function and distinct proliferative responses to CD19- and CD20-expressing target cells

We next tested the anti-tumor function of GF-CART01 cells in two independent donors, PBMC-41

and PBMC-40. Raji-GFP/Luc tumor cells were co-cultured with GF-CART01 cells at effector to target (E:T)

ratios of 5:1, 1:1, and 1:5, and cytotoxicity was assessed at 24, 48, 72, and 96 hours after co-culture **(Figure**

3A and B). While GF-CART01 cells lysed Raji-GFP/Luc tumors in a dose-dependent manner at early time

points (24 to 48 h), complete tumor lysis was seen in all E:T ratios at 96 hours. Furthermore, GF-CART01

cells secreted significant levels of pro-inflammatory cytokines IFN-γ, TNF-α, and IL-2 compared to pan-T

cells **(Figure 3C-E)**. Collectively, these data suggest that functional GF-CART01 cells can be generated using

the *qCART*™ system. Next, we determined how encountering different antigen levels may affect the

proliferation capacity of GF-CART01 cells. To address this question, eF450 dye-labeled GF-CART01 cells

were co-cultured with K562 cells overexpressing different levels of CD19 or CD20 and proliferation

capacity assessed at 48 and 72 hours (see **Supplementary Figure S2** for gating strategy). As shown in

Table 1, the percentage of proliferated GF-CART01 cells was inversely associated with CD19 expression,

but positively associated with CD20 expression, on K562 cells. No such association was observed with CAR⁻

T cells in the corresponding groups and CAR-T cell viability did not appear to be negatively affected by

CD19 or CD20 expression (data not shown).

Table 1. Proliferation capacity of GF-CART01 cells in response to K562 expressing different levels of CD19

or CD20 antigens. Percentages of proliferated live CAR+ and CAR- GF-CART01 cells following 48 h and 72

h co-culture with different levels of CD19- or CD20-overexpressing K562 clones are shown.

		CAR⁺				CAR⁺			
		% Proliferated				% Proliferated			
	Target cell	K562	K562-CD19 #4-B10 (low)	K562-CD19 #7-B8 (medium)	K562-CD19 #2D-4 (high)	K562	K562-CD20 #4-2 (low)	K562-CD20 #5-2 (medium)	K562-CD20 #3-2 (high)
Time of	48 hours	4.8 ± 1.7	28.7 ± 2.4	24.9 ± 0.6	19.5 ± 1.5	7.6 ± 0.8	12.5 ± 1.2	13.5 ± 0.7	11.9 ± 1.3
co-culture	72 hours	27.6 ± 1.0	77.5 ± 1.3	69.3 ± 1.9	63.9 ± 0.1	10.4 ± 0.2	60.4 ± 3.0	74.7 ± 2.3	72.7 ± 2.0
		CAR⁻				CAR⁻			
		% Proliferated				% Proliferated			
	Target cell	K562	K562-CD19 #4-B10 (low)	K562-CD19 #7-B8 (medium)	K562-CD19 #2D-4 (high)	K562	K562-CD20 #4-2 (low)	K562-CD20 #5-2 (medium)	K562-CD20 #3-2 (high)
Time of	48 hours	2.2 ± 2.0	12.6 ± 1.6	13.9 ± 0.9	11.3 ± 0.3	4.7 ± 0.9	5.5 ± 2.2	5.6 ± 0.7	7.2 ± 3.2
co-culture	72 hours	19.0 ± 1.4	44.9 ± 2.6	50.3 ± 1.3	46.3 ± 0.3	6.9 ± 0.8	28.1 ± 2.1	31.9 ± 0.7	25.6 ± 0.5

% of CAR+ cells (stained by anti-mouse IgG Fab') were detected by flow cytometry (SonySA3800) All results were

obtained in triplicates

GF-CART01 cells display enhanced and persistent anti-tumor function *in vivo*

We next evaluated the anti-tumor efficacy of GF-CART01 cells in an *in vivo* xenotransplant

immunodeficient mouse model using donor PBMC-41. We injected 1.5×10^5 Raji-GFP/Luc tumor cells

intravenously (*i.v.*) followed by 7.5×10^5 (low), 3.0×10^6 (medium), or 1×10^7 (high) GF-CART01 cells (*i.v.*) 7

days later as depicted in **Figure 4A**. As assessed by bioluminescence, all mice in the vehicle group

succumbed to tumor by day 21 **(Figure 4B-C)**. In contrast, Raji-GFP/Luc tumors were dose-response

dependently cleared and all tumors were completely eradicated by day 13 in all three treatment groups

and remained tumor-free **(Figure 4B)**. Minimal cytokine production was detected in all three treatment

groups on days 2, 8, and 14 **(Supplementary Figure S3A)**. On day 26, no detectable levels of Raji-GFP/Luc

tumors were observed in all three treatment groups **(Supplementary Figure S3B)**. Moreover, residual CAR

gene was observed in the blood of the medium and high groups, suggesting that many GF-CART01 cells

persisted to form memory T cells **(Supplementary Figure S3C)**. We also obtained data from mice injected

with non-transfected control Pan-T cells that were cultured under the same conditions as the GF-CART01

cells (**Supplementary Figure S4**). These cells exhibited anti-Raji activity approximately two weeks

following injection, which likely resulted from *in vivo* expansion of cells that recognized (and killed) Raji

tumor cells via a CAR-independent, TCR-mediated mechanism. However, such a killing effect would not

be expected with autologous T cell therapy in patients.

While residual GF-CART01 cells (as determined by qPCR for presence of CAR gene in genomic DNA

of mouse blood samples) can be detected up to day 26, it remained unclear whether these residual cells

were memory T cells and retained their anti-tumor function. To address this, we rechallenged half of the

mice in each treatment group with 1.5×10^5 Raji-GFP/Luc tumor cells on day 35. Due to the lack of CAR-T

cell persistence, most of the rechallenged mice in the low group eventually succumbed to tumor **(Figure**

4B-C). In contrast, the rechallenged mice in the medium and high groups remained tumor-free throughout

the experiment. Some mice in the medium and high groups did not survive, but this was likely due to graft-

versus-host disease (GVHD) as severe ruffling were observed in these mice. More importantly, GF-CART01

cell treatment significantly improved the overall survival of tumor-bearing mice compared to vehicle

control **(Figure 4D)**.

372   To confirm that our non-viral gene engineering approach produces CAR-T cells of high quality and

quantity, we next directly compared our non-viral *qCART™* system with a conventional lentiviral

engineering approach. CAR-T cells derived from the same healthy donor PBMC source carrying a CAR

gene targeting to a pan-cancer antigen was produced utilizing both approaches. The anti-tumor effects of

these CAR-T cells were determined in mice harboring NCI-N87 gastric carcinoma xenografts. The rationale

for choosing a single CAR gene targeting to a solid tumor model system was two-fold: First, since the

transgene size of GF-CART01used in the Raji tumor model animal experiments was 5.2 Kb (**Figure 4**), CAR-

T production using a viral vector system expressing such a sizable transgene would be difficult and may

introduce additional variables influencing the results. Second, comparisons carried out in a solid tumor

model would be expected to better differentiate between the differences in quality of the CAR-T cells

produced using *qCART™* and the lentivirus systems, since solid tumors are more challenging to be

eliminated. We demonstrated that *qCART™* produced CAR-T cells were superior in performance

compared to lentivirally-engineered CAR-T cells, characterized by greater *in vitro* expansion capacity,

higher CD8/CD4 ratio, and enrichment of CAR-T$_{SCM}$ cells **(Supplementary Table S1)**. More importantly, in

the solid tumor NCI-N87 xenotransplant model **(Figure 4E)**, we observed enhanced antitumor efficacy as

well as *in vivo* expansion of *qCART™*-produced CAR-T cells. This was characterized by complete clearance

of tumors at Day 28 **(Figure 4F)** in mice treated with *qCART™*-produced CAR-T cells but not those treated

with lentivirus-engineered counterparts. Furthermore, between 10 and 37 days following CAR-T injection

(CAR-T cells injected at 11 days following tumor inoculation), circulating CAR-T cells robustly proliferated

in mice (10 to 100 fold increase in circulating CAR-T cells), while lentivirus-produced CAR-T cells either

minimally proliferated or did not proliferate at all **(Figure 4G)**. This difference may reflect the above-

mentioned higher CD8/CD4 ratio and percentage of T_N/T_{SCM} population present in both the CD4$^+$ and CD8$^+$

CAR-T populations of *qCART™*-produced CAR-T cells **(Supplementary Table S1)**.

Functional GF-CART01 T$_{SCM}$ cells can be generated in sufficient numbers from cancer patients

We next assessed whether sufficient high-quality GF-CART01 cells can be generated for clinical

applications. In our previous experiments thus far, we expanded all GF-CART01 cells with CD19-expressing

aAPC. To streamline the manufacturing process, we tested whether we can still generate highly potent

patient-derived GF-CART01 T$_{SCM}$ cells in the absence of aAPC selective expansion. We isolated PBMCs from

6 healthy donors, 3 diffuse large B-cell lymphoma (DLBCL) patients, 3 chronic lymphocytic leukemia (CLL)

patients, one Hodgkin lymphoma (HL) patient, and one multiple myeloma (MM) patient to generate GF-

CART01 cells. **Figure 5** and **Table 2** summarize the characteristics of the healthy donor- and patient-

derived CAR-T cells.

Table 2. Comparison of GF-CART01 cells generated from healthy donors and cancer patients

Disease		Healthy Donor	DLBCL1	DLBCL2	DLBCL3	CLL1	CLL2	CLL3	HL	MM
Copy Number		1.85-3.47	2.51	5.46	2.21	3.36	2.81	3.76	2.04	1.82
% of CAR$^+$		32.5–49.6	26.3	50.1	34.23	40.6	33.4	53.22	26.2	19.9
Fold Expansion		43.54–157.55	67.22	67.3	60.38	104.6	58.0	198.96	149.75	139.35
Cell Number (x10^6)		17.34–50.60	7.29	4.95	5.8	15.8	7.71	4.58	24.08	27.54
% of B Cells (CD19$^+$)		0.01-0.36	0.77	1.18	0.35	0	0	0.61	0	0
% of NK Cells (CD3$^-$, CD56$^+$)		0.04-0.27	0.06	2.69	0.63	0.03	0.02	0.02	0.08	0.05
% of Exhaustion Marker$^+$ Cells	PD-1	0.40–1.96	5.4	4.2	4.2	1.0	2.4	6.0	1.7	7.8
	TIM-3	0.10–5.34	1.8	3.0	2.0	1.4	2.3	2.4	1.4	1.1
	LAG-3	0.90–7.00	9.9	12.4	22.8	3.2	4.0	2.6	2.4	2.6
% of Senescence Marker$^+$ Cells	KLRG-1	2.24–14.61	11.8	14.7	8.2	2.8	4.2	6.1	4.1	5.3
	CD57	0.03–0.68	5.8	1.5	2.5	1.1	1.8	0.6	1.1	1.3
Potency (% of lysis) (fresh cells, E/T=5, 48 h)		89.74–95.38	99.66	99.7	99.7	99.88	99.45	99.07	98.96	99.42
Potency (% of lysis) (frozen cells, E/T=5, 48 h)		44.99–61.96	73.35	74.21	75.21	84.30	75.70	45.46	77.49	70.31

CAR-T cells generated from PBMCs of 6 healthy donors and 8 cancer patients were cultured in 24-well G-Rex for 10 days. The cell number of 24-well G-Rex culturing times 50 is equal to cell number of 1 L G-Rex culturing (2.29×10^8-1.38×10^9 cells in 1 L G-Rex). DLBCL: diffuse large B-cell lymphoma; CLL: chronic lymphocytic leukemia; HL: Hodgkin lymphoma; MM: multiple myeloma.

Over the course of 10 days, we expanded the CAR-T cells to 4.58×10^6–2.76×10^7 cells in a 24-well G-Rex culture, which would be equivalent to clinically relevant number of 2.29×10^8–1.38×10^9 cells in a 1 L G-Rex culture vessel. The CAR expression (19.9–53.22%) and the fold expansion (58.0–198.96 folds) of the patient cells were comparable to those from healthy donors (32.5–49.6% and 43.54–157.55 folds) **(Table 2)**. The CAR-T cell purity was confirmed by the low percentage of CD19$^+$ B cells and CD3$^-$CD56$^+$ NK cells in both the healthy donors (0.01-0.36% B; 0.04-0.27% NK) and patient cells (0-1.18% B; 0.02-2.69% NK)

(Table 2). The proportion of CD8 and CD4 were balanced in both the healthy donors (CD8/CD4=0.85) and

the patients (CD8/CD4=1.17) prior to nucleofection **(Figure 5A)**. After nucleofection, the CAR-T cells from

healthy donors had higher CD8/CD4 ratio (5.64). However, the increase in the proportion of CAR$^+$ CD8 T

cells was only observed in some patients **(Figure 5A)**. Notably, compared to pre-nucleofection (PBMC)

levels, the percentages of T$_{SCM}$ subset at 10 days post-nucleofection increased in all patients among the

CD4 (from 13.75% to 76.25%) and CD8 (from 7.17% to 81.90%) populations, confirming the unique quality

of *qCART*™ system in supporting T$_{SCM}$ differentiation **(Figure 5B,** see **Supplementary Figure S5** for gating

strategy). To further confirm T$_{SCM}$ status, we determined the expression of additional surface antigen

biomarkers whose expression have previously been reported to be associated with T$_{SCM}$ and compared

their expression among the T$_{SCM}$ (CD45RA$^+$CD62L$^+$CD95$^+$) population of PBMC T cells and CAR-T cells

derived from three patients and healthy volunteers (**Supplementary Figure S6**). Consistent with data

presented in **Figure 2E** and **Figure 5B**, after engineering and expansion, both the patients' and healthy

volunteers' (CD4 and CD8) CAR-T cells were highly enriched for T$_{SCM}$ (**Supplementary Figure S6A**). High

percentage of these T$_{SCM}$ cells also co-expressed CD45RO, CCR7, CD27 and CD28, confirming these cells'

T$_{SCM}$ status (**Supplementary Figure S6C-S6D**). Notably, the CD45RO^{dim} population in these T$_{SCM}$ cells may

represent cells that were in transition into a more differentiated state. Furthermore, we also found that

the T$_N$/T$_{SCM}$ population of (particularly healthy donor) PBMCs contained only low percentage of CD95$^+$ cells

(**Supplementary Figure S6B**), suggesting that the T$_N$/T$_{SCM}$ population of PBMCs were dominated by naïve

T cells. The GF-CART01 cells were also assessed for expression of exhaustion markers PD-1 and TIM-3 and

senescence marker CD57, and only a low percentage of cells were found to express these markers **(Table**

2). In addition, the percentage of cells expressing the exhaustion marker LAG-3 was low in CLL, HL, and

MM patients (2.4-4%) with a slightly elevated percentage of cells expressing these markers among the

DLBCL patients (9.9-22.8%). While KLRG-1 expression (~7.25%) was detectable in GF-CART01 patient cells,

it was significantly lower than KLRG-1 expression prior to nucleofection (~21.27%; p=0.0087)

(Supplementary Figure S7A). The decrease in KLRG-1 expression was especially prominent in the CD8

population (52.50% vs. 14.59%; p=0.0004) **(Supplementary Figure S7B)**. A 48-hour cytotoxicity assay

further confirmed the functionality of the GF-CART01 cells **(Table 2)**. Together, our data suggest that the

qCART™ system is a promising technology for generating functional multiplex armored CAR-T$_{SCM}$ cells for

cancer immunotherapy application. GF-CART01 therapy is a safe, potent, and cost-effective treatment for

B-cell malignancies.

DISCUSSION/ CONCLUSION

qPB is currently the most efficient transposon for gene integration, and GF-CART01 cell therapy

was the first CAR-T therapy developed using the *qPB*-based *qCART*™ system.[15,16] The *qCART*™ system

can easily produce CAR-T cells with transgene size >7.6 kb, which is not achievable by conventional

lentiviral/retroviral CAR-T engineering.[16] The limited cargo capacity of lentiviral/retroviral vector may

account for the lack of current CAR-T clinical trials targeting B-cell malignancies with both dual-targeting

and iCasp9 safety switch gene (5.2 kb transgene size) in the CAR design. In this report, we presented

evidence on the feasibility of manufacturing highly potent iCasp9-regulatable CD20/CD19-targeted CAR-

T$_{SCM}$ cells using the virus-free *qCART*™ system for clinical application. We demonstrated the purity,

potency, and consistency profiles of CAR-T cells manufactured using the *qCART*™ system.

Maintaining sterility and controlling mycoplasma and endotoxin contamination in the CAR-T

product are important safety concerns. G-Rex is a GMP-compliant medical device used for cell

expansion.[19,20] The minimal manual operation of this device minimizes contamination risks.

Importantly, manufacturing GF-CART01 cells with G-Rex can be achieved without compromising CAR-T

efficacy. Consistent with our previous study, we observed low risk of genotoxicity in GF-CART01 cells,

given the safe integration profile, low enhancer activity, and low residual *qPBase*.[16] Acute toxicities in

CAR-T therapy-treated patients, such as CRS and ICANS, have been major concerns.[21–23] Furthermore,

a rare case of CAR[+] CD4 T cell lymphoma has been reported recently in patients treated with CD19 CAR-T

cells.[24,25] To safely manage toxicities, CAR-T cells can be eradicated by targeting CAR-T cells (e.g. anti-

EGFRt antibody) via antibody-dependent cellular cytotoxicity (ADCC) mechanism or using a small molecule

that activates a suicide gene introduced in the CAR design (e.g. iCasp9).[21] The ADCC approach has slower

onset of killing compared to the latter. We included an iCasp9 safety switch in our GF-CART01 design to

mitigate the adverse effects of CRS and ICANS. Indeed, Foster et al (2021) has recently demonstrated that

acute ICANS can be effectively controlled by treating CAR-T therapy patients with the iCasp9-activating

agent, rimiducid (AP1903).[26]

Manufacturing clinically relevant number of a pure high-quality CAR-T product is the first critical

step towards successful CAR-T therapy. One approach to expanding the CAR[+] T cell population is by

culturing with aAPC, but this often leads to a more differentiated and short-lived T_{EFF} or exhaustive CAR-T

phenotype.[27] Another approach is to include markers within the construct to enrich the CAR[+] T cell

population either via drug selection or antibody-mediated cell isolation during or after CAR-T *ex vivo*

expansion, respectively. However, such approaches increase the payload size and introduce

therapeutically irrelevant genes into the genome.[4,22,23] While we also used aAPC in our initial

experiments, we could generate sufficient quantity of patient-derived GF-CART01 cells without aAPC-

induced enrichment by addition of *Quantum Booster*™, a serum-free supplement which contains cellular

proteins and a set of cytokines with defined concentrations. These components of *Quantum Booster*™ act

to promote proliferation of the CAR-T cells while maintaining stemness as well as minimal levels of

senescence and exhaustion.[16] Notably, we have shown that *Quantum Booster*™ highly enriches CD4

T_{SCM}, which may be critical in maintaining CD8[+] CAR-T cells at the T_{SCM} differentiation stage while being

robustly expanded. Our recent study further confirmed that *qCART*™-manufactured CAR-T cells may be

more effective in tumor control without aAPC enrichment.[16] Here, we demonstrated that 2.29×10^8–

1.38×10^9 T cells could be generated in a 1 L-G-Rex within a short time-frame of 10 days using cancer

patient-derived T cells. Even at a low CAR$^+$ percentage of 20%, we could still produce sufficient number of

CAR$^+$ CAR-T cells (4.58×10^6–2.76×10^7) for treatment, and we can easily enrich CAR$^+$ T cells with aAPC should

the need arise.

The high potency of GF-CART01 cells can be attributed to the balanced CD8/CD4 CAR-T ratio and

the high enrichment of T$_{SCM}$ in the engineered cells. It has been reported that a balanced CD8/CD4 CAR-T

ratio resulted in high remission rates.[28,29] To achieve this, others have engineered CD4 and CD8 CAR-T

cells separately before adoptive transfer.[28] While *qCART*™ cells derived from healthy donors generally

favored CD8 enrichment, cancer patient-derived *qCART*™ cells have a balanced CD8/CD4 CAR-T ratio

(**Figure 5**). Thus, the *qCART*™ system naturally achieves the desired CD8/CD4 balance without the need to

engineer CD4 and CD8 CAR-T cells separately as seen in SB100X-modified CAR-T clinical trials.[30]

Our observation that, in contrast to a positive dose-dependent relationship between CD20 target

cell expression level and proliferation capacity of GF-CART01 cells, there appeared to be an inverse

association between CD19 target cell expression level and CAR-T proliferation when encountering K562

cells overexpressing different levels of CD19. However, potential toning of GF-CART01 cells would be

unlikely since we found CD19 expression level of Raji cell line was much lower than the K562 clone which

expressed the lowest level of CD19. In other words, since tumor's endogenous CD19 expression level was

generally lower than the level that can inhibit CAR-T proliferation, we do not expect to see a decrease in

CAR-T proliferation similar to that observed with the K562 CD19 overexpression clone which expressed

high levels of CD19.

T$_{SCM}$ are a class of immunological memory critical for successful adoptive T cell therapies.[31–35]

The *qCART*™ system is unique in its ability to enrich T$_{SCM}$ population without compromising the cell quality,

expansion, and production time of CAR-T cell product.[16] One approach to harness the potency of less-

differentiated T cells such as T$_{SCM}$ is to shorten the *ex vivo* expansion time of CAR-T product. *Ghassemi et*

al. (2022) demonstrated that shortening the lentiviral CAR-T manufacturing process to 24-hours resulted

in a higher percentage of CAR-T cells with a memory phenotype.[36] Though these CAR-T cells displayed

stronger *in vivo* anti-tumor activity than their conventional counterpart, successful clinical translation of

this process remains to be demonstrated. Another approach is to culture CAR-T cells with specific

cytokines or chemical reagents.[34,35,37–39] Additionally, CAR-T cell production from defined T cell

subsets has been shown to be beneficial in enriching the T_{SCM} population.[40] However, none of these

approaches enrich T_{SCM} population to a high level seen in our *qCART™*-modified T cells. While T_{SCM} cells

make up a small percentage of T cells derived from cancer patients, genetically modifying these cells using

the *qCART™* system resulted in enrichment of T_{SCM} to as high as >80% in both the CD4 and CD8 CAR^+ T cell

populations.

In conjunction with the ability of T_{SCM} to self-renew and persist long-term, we also observed low

expression of exhaustion (PD-1, TIM-3, and LAG-3) and senescence (CD57) biomarkers in our GF-CART01

cells **(Figure 2F-G; Table 2)**. KLRG-1 is a senescence marker. While we observed a small but significant

population of $KLRG-1^+$ cells in GF-CART01 cells, we also observed a significantly lower percentage of KLRG-

1^+ cells after CAR-T production compared to freshly isolated PBMCs **(Supplementary Figure S2)**. Herndler-

Brandstetter *et al.* (2018) recently described a class of memory T cell population developed from $KLRG-1^+$

effector CD8 T cells that have lost KLRG-1 expression.[31] These so-called "exKLRG-1 memory cells" have

been demonstrated to inhibit tumor growth more efficiently compared to $KLRG-1^+$ cells in an OT-I

melanoma mouse model. T_{SCM} cells generally lack KLRG-1 expression, and $KLRG-1^+$ T_{SCM} cells have been

associated with cancer patients undergoing relapse.[32,33] Given the significant decrease in KLRG-1

expression in GF-CART01 cells, it is possible that a significant proportion of CAR-T_{SCM} cells in our model

were actually exKLRG-1 memory cells. Thus, one possible mechanism that made it possible for *qCART™*

system to generate highly potent CAR-T_{SCM} cells may be through the conversion of $KLRG-1^+$ effector CD8

T cells to exKLRG-1 memory cells.

One of the major challenges of CAR-T cell manufacturing is the variation and inconsistency of CAR-

T cell products.[34] In one study of ten CLL and ALL patients, CAR-T cells expanded anywhere between

23.6 and 385-folds with retroviral CAR transduction of 4-70%.[35] Lentiviral transduction of 5.5-45.3% was

reported by other studies of CLL and ALL patients.[36,37] In our GF-CART01 study, we observed tighter

range of CAR$^+$ percentage (19.9-53.22%) and expansion (58.0-198.96 folds) among the eight cancer

patients **(Table 2)**. Furthermore, these low levels of variation were comparable to the CAR$^+$ percentage

(32.5-49.6%) and expansion (43.54-157.55 folds) among the healthy donors in our study, respectively.

More importantly, GF-CART01 cells derived from both healthy donors and patients can effectively

eradicate tumor cells.

In this study, we directly compared between the performance and anti-tumor efficacy of *qCART™*-

and lentivirally-engineered CAR-T cells in a solid tumor xenograft model. The results clearly demonstrated

superior performance, anti-tumor efficacy, and *in vivo* expansion of *qCART™*-produced CAR-T cells over

lentivirally-produced CAR-T cells (**Figure 4E-4G**). These observations suggest that *qCART™*-produced CAR-

T cells, including GF-CART01 cells, are indeed highly potent.

In summary, the *qCART™* system is an elegant system for the manufacturing of CAR-T products.

In this study, we demonstrated that the *qCART™* system can be used to produce patient-derived GF-

CART01 cells with all the desired attributes of CAR-T products to ensure therapeutic success, including

high CAR-T$_{SCM,}$ balanced CD8/CD4 ratio, low exhaustion and senescence marker expressions, high

expansion capacity, and excellent anti-tumor efficacy. We believe that the simplicity and robustness of

manufacturing multiplex CAR-T cells with enhanced product consistency using the *qCART™* system will

not only significantly enhance the accessibility of CAR-T therapy but also unlock the full potential of

armored CAR-T therapy for the treatment of solid tumors in the future.

REFERENCES

1. Larson RC, Maus M V. Recent advances and discoveries in the mechanisms and functions of CAR
T cells. Nat Rev Cancer. Springer US; 2021;

2. Sheth VS, Gauthier J. Taming the beast: CRS and ICANS after CAR T-cell therapy for ALL. Bone
Marrow Transplant. Springer US; 2021;56:552 – 66.

3. Meir YJJ, Wu SCY. Transposon-based vector systems for gene therapy clinical Trials: Challenges
and considerations. Chang Gung Med J. 2011;34:565 – 79.

4. Singh H, Moyes JSE, Huls MH, Cooper LJN. Manufacture of T cells using the Sleeping Beauty
system to enforce expression of a CD19-specific chimeric antigen receptor. Cancer Gene Ther. Nature
Publishing Group; 2015;22:95 – 100.

5. Brentjens RJ, Latouche JB, Santos E, Marti F, Gong MC, Lyddane C, et al. Eradication of systemic
B-cell tumors by genetically targeted human T lymphocytes co-stimulated by CD80 and interleukin-
15. Nat Med. 2003;9:279 – 86.

6. Cronk RJ, Zurko J, Shah NN. Bispecific chimeric antigen receptor T cell therapy for b cell
malignancies and multiple myeloma. Cancers (Basel). 2020;12:1 – 15.

7. Levin A, Shah NN. Chimeric antigen receptor modified T cell therapy in B cell non-Hodgkin
lymphomas. Am J Hematol. 2019;94:S18 – 23.

8. Park JH, Rivière I, Gonen M, Wang X, Sénéchal B, Curran KJ, et al. Long-Term Follow-up of
CD19 CAR Therapy in Acute Lymphoblastic Leukemia. New England Journal of Medicine.
2018;378:449 – 59.

9. Spiegel JY, Dahiya S, Jain MD, Nastoupil LJ, Ghobadi A, Lin Y, et al. Outcomes in large B-cell
lymphoma progressing after axicabtagene ciloleucel (Axi-cel): Results from the U.S. Lymphoma
CAR-T Consortium. Journal of Clinical Oncology. 2019;37:7517 – 7517.

10. Neelapu SS, Rossi JM, Jacobson CA, Locke FL, Miklos DB, Reagan PM, et al. CD19-Loss with
Preservation of Other B Cell Lineage Features in Patients with Large B Cell Lymphoma Who
Relapsed Post-Axi-Cel. Blood. 2019;134:203 – 203.

11. Shah NN, Johnson BD, Schneider D, Zhu F, Szabo A, Keever-Taylor CA, et al. Bispecific anti-
CD20, anti-CD19 CAR T cells for relapsed B cell malignancies: a phase 1 dose escalation and
expansion trial. Nat Med. 2020;26:1569 – 75.

12. Tong C, Zhang Y, Liu Y, Ji X, Zhang WW, Guo Y, et al. Optimized tandem CD19/CD20 CAR-
engineered T cells in refractory/relapsed B cell lymphoma. Blood. 2020;136:1632 – 44.

13. Wu SCY, Meir YJJ, Coates CJ, Handler AM, Pelczar P, Moisyadi S, et al. piggyBac is a flexible
and highly active transposon as compared to Sleeping Beauty, Tol2, and Mos1 in mammalian cells.
Proc Natl Acad Sci U S A. 2006;103:15008 – 13.

14. Querques I, Mades A, Zuliani C, Miskey C, Alb M, Grueso E, et al. A highly soluble Sleeping
Beauty transposase improves control of gene insertion. Nat Biotechnol. Springer US; 2019;37:1502 –
12.

15. Hua W-K, Hsu JC, Chen Y-C, Chang PS, Wen KK-L, Wang P-N, et al. Quantum pBac: An
effective, high-capacity piggyBac-based gene integration vector system for unlocking gene therapy

potential. https://doi.org/101101/20220429490002 [Internet]. 2022; Available from:

https://doi.org/10.1101/2022.04.29.490002

16. Chen Y-C, Hua W-K, Hsu JC, Chang PS, Wen KK-L, Huang Y-W, et al. Quantum CART

(qCART), a piggyBac-based system for development and production of virus-free multiplex CAR-T

cell therapy. bioRxiv [Internet]. 2022; Available from: https://doi.org/10.1101/2022.05.03.490469

17. Hultin LE, Hausner MA, Hultin PM, Giorgi J v. Cd20 (pan-B cell) antigen is expressed at a low

level on a subpopulation of human T lymphocytes. Cytometry. 1993;14:196 – 204.

18. Nicholson IC, Lenton KA, Little DJ, Decorso T, Lee FT, Scott AM, et al. Construction and

characterisation of a functional CD19 specific single chain Fv fragment for immunotherapy of B

lineage leukaemia and lymphoma. Mol Immunol. 1997;34:1157 – 65.

19. Gotti E, Tettamanti S, Zaninelli S, Cuofano C, Cattaneo I, Rotiroti MC, et al. Optimization of

therapeutic T cell expansion in G-Rex device and applicability to large-scale production for clinical

use: GMP-compliant polyclonal T cell expansion in G-Rex. Cytotherapy. 2022;24:334 – 43.

20. Gagliardi C, Khalil M, Foster AE. Streamlined production of genetically modified T cells with

activation, transduction and expansion in closed-system G-Rex bioreactors. Cytotherapy. Elsevier Inc.;

2019;21:1246 – 57.

21. Rafiq S, Hackett CS, Brentjens RJ. Engineering strategies to overcome the current roadblocks in

CAR T cell therapy. Nat Rev Clin Oncol. Springer US; 2020;17:147 – 67.

22. Wang X, Chang WC, Wong CLW, Colcher D, Sherman M, Ostberg JR, et al. A transgene-

encoded cell surface polypeptide for selection, in vivo tracking, and ablation of engineered cells.

Blood. 2011;118:1255 – 63.

23. Wang Q, He F, He W, Huang Y, Zeng J, Zi F, et al. A transgene-encoded truncated human

epidermal growth factor receptor for depletion of anti- B-cell maturation antigen CAR-T cells. Cell

Immunol. Elsevier Inc.; 2021;363:104342.

24. Micklethwaite KP, Gowrishankar K, Gloss BS, Li Z, Street JA, Moezzi L, et al. Investigation of

product-derived lymphoma following infusion of piggyBac-modified CD19 chimeric antigen receptor

T cells. Blood. American Society of Hematology; 2021;138:1391 – 405.

25. Bishop DC, Clancy LE, Simms R, Burgess J, Mathew G, Moezzi L, et al. Development of CAR

T-cell lymphoma in 2 of 10 patients effectively treated with piggyBac-modified CD19 CAR T cells.

Blood. American Society of Hematology; 2021;138:1504 – 9.

26. Foster MC, Savoldo B, Lau W, Rubinos C, Grover N, Armistead P, et al. Utility of a safety switch

to abrogate CD19.CAR T-cell – associated neurotoxicity. Blood. 2021;137:3306 – 9.

27. Nukala U, Rodriguez Messan M, Yogurtcu ON, Wang X, Yang H. A Systematic Review of the

Efforts and Hindrances of Modeling and Simulation of CAR T-cell Therapy. AAPS Journal.

2021;23:1 – 33.

28. Turtle CJ, Hanafi LA, Berger C, Gooley TA, Cherian S, Hudecek M, et al. CD19 CAR-T cells of

defined CD4+:CD8+ composition in adult B cell ALL patients. Journal of Clinical Investigation.

2016;126:2123 – 38.

29. Sommermeyer D, Hudecek M, Kosasih PL, Gogishvili T, Maloney DG, Turtle CJ, et al. Chimeric

antigen receptor-modified T cells derived from defined CD8+ and CD4+ subsets confer superior

antitumor reactivity in vivo. Leukemia. 2016;30:492 – 500.

30. Prommersberger S, Reiser M, Beckmann J, Danhof S, Amberger M, Quade-Lyssy P, et al.
CARAMBA: a first-in-human clinical trial with SLAMF7 CAR-T cells prepared by virus-free
Sleeping Beauty gene transfer to treat multiple myeloma. Gene Ther. Springer US; 2021;28:560–71.
31. Herndler-Brandstetter D, Ishigame H, Shinnakasu R, Plajer V, Stecher C, Zhao J, et al. KLRG1+
Effector CD8+ T Cells Lose KLRG1, Differentiate into All Memory T Cell Lineages, and Convey
Enhanced Protective Immunity. Immunity. Elsevier Inc.; 2018;48:716-729.e8.
32. Liu Q, Sun Z, Chen L. Memory T cells: strategies for optimizing tumor immunotherapy. Protein
Cell. Higher Education Press; 2020;11:549–64.
33. Noviello M, Manfredi F, Ruggiero E, Perini T, Oliveira G, Cortesi F, et al. Bone marrow central
memory and memory stem T-cell exhaustion in AML patients relapsing after HSCT. Nat Commun.
2019;10:1–15.
34. Vormittag P, Gunn R, Ghorashian S, Veraitch FS. A guide to manufacturing CAR T cell therapies.
Curr Opin Biotechnol. Elsevier Ltd; 2018;53:164–81.
35. Brentjens RJ, Rivière I, Park JH, Davila ML, Wang X, Stefanski J, et al. Safety and persistence of
adoptively transferred autologous CD19-targeted T cells in patients with relapsed or chemotherapy
refractory B-cell leukemias. Blood. 2011;118:4817–28.
36. Porter DL, Hwang WT, Frey N V., Lacey SF, Shaw PA, Loren AW, et al. Chimeric antigen
receptor T cells persist and induce sustained remissions in relapsed refractory chronic lymphocytic
leukemia. Sci Transl Med. 2015;7.
37. Dai H, Zhang W, Li X, Han Q, Guo Y, Zhang Y, et al. Tolerance and efficacy of autologous or
donor-derived T cells expressing CD19 chimeric antigen receptors in adult B-ALL with
extramedullary leukemia. Oncoimmunology. 2015;4.

                                     FIGURE LEGENDS

Figure 1. Comparison of perfusion and fed-batch-cultured CAR-T cells. (A) Schematic diagram of the GF-

CART01 CAR construct. A T2A sequence enables co-expression of iCasp9 and CD20/CD19-targeted scFvs

under the EF1α promoter. (B) Characterization of GF-CART01 cells cultured in conventional plates

(perfusion) or G-Rex (fed batch) at day 10 after nucleofection. (C) Cytolytic activities of CAR-T cells after

48 or 72 h of co-culture with Raji-GFP/Luc target cells (E:T ratio of 5:1) were assessed by Celigo image

cytometry. (B-C) Data represent mean ± SD in 6 healthy donors, n=6. *p<0.05, **p<0.01, ***p<0.001.

Figure 2. Characteristics of GF-CART01 cells. Percentage of (A) CAR$^+$ and (B) transposase$^+$ (*qPBase$^+$*) cells

were analyzed by flow cytometry on days 1, 8, and 10 after nucleofection. (C) Fold change of GF-CART01

cells after 10 days of culture in G-Rex are shown. (D) Percentage of CD4 and CD8 T cells in the CAR$^+$

population on day 10 after nucleofection. (E) Distribution of T_N, T_{SCM}, T_{CM}, T_{EM}, and T_{EFF} subsets in the CD4

(upper panel) and CD8 populations (lower panel). (F) Expression of exhaustion markers PD-1, TIM-3, and

LAG-3 in GF-CART01 cells. (G) Expression of senescence markers KLRG-1 and CD57 in GF-CART01 cells. (A-

G) Data represent mean ± SD in 9 healthy donors, n=9. Each set of histogram plots shown in (D), (E), (F),

and (G) represents one donor.

Figure 3. *In vitro* functional analysis of GF-CART01 cells. (A-B) CAR-T cells derived from two healthy

donors PBMC-41 and PBMC-40 were assessed for cytotoxicity against Raji-GFP/Luc cells by Celigo image

cytometry. Statistical differences were calculated by One-way ANOVA with Tukey multiple comparison,

***p<0.001, **p<0.01, *p<0.05. (C) IFN-γ, (D) TNF-α, and (E) IL-2 secretion by GF-CART01 cells following

antigen stimulation was determined by ELISA. Pan-T cells (non-modified cells) served as a control group.

Data represent mean ± SD, n=3. Statistical differences are calculated by Student's t-test, ***p<0.001.

Figure 4. Anti-tumor activity of GF-CART01 cells in a B-cell lymphoma immunodeficient xenograft mouse

model. (A) Schematic diagram of *in vivo* Raji xenograft model experimental design. (B-C) Bioluminescent

imaging was performed to monitor tumor cell persistence (n= 4/group) and tumor growth were quantified

by total flux (photon/s). (D) Kaplan-Meier survival curves for the different treatment groups are shown.

(E) Schematic diagram of *in vivo* NCI-N87 xenograft model experimental design. (F) Bioluminescent

imaging was performed to monitor tumor cell persistence (n= 3/group). (G) Human T cell counts of blood

samples taken from the indicated group of mice at Day 21 and Day 48 post-tumor inoculation.

Figure 5. Assessment of cancer-patient-derived GF-CART01 cells. T cells derived from healthy donors and

cancer patients were nucleofected with GF-CART01. (A) CD8/CD4 ratios were assessed before (PBMC) and

after (CAR-T) nucleofection. (B) Distribution of T_N, T_{SCM}, T_{CM}, T_{EM}, and T_{EFF} subsets in CD4$^+$ (upper panel)

and CD8$^+$ (lower panel) CAR-T cell populations. DLBCL, diffuse large B-cell lymphoma; CLL, chronic

lymphocytic leukemia; HL, Hodgkin's lymphoma; MM, multiple myeloma.

Figure 1.

A

GF-CART01

B

Culture Vessel	Conventional Plate (perfusion)	G-Rex (fed batch)
% CAR⁺ T cells	51%* (44%~61%, n=6)	46%* (42%~49%, n=6)
Expansion capacity of CAR⁺ T cells	836 fold* (406~2090 fold, n=6)	189 fold* (139~262 fold, n=6)
Distribution of T cell subsets in CAR⁺ T cells

 | |

*median

C

Figure 2.

Figure 3.

Figure 4.

Figure 5.