***CHRNA5* links chandelier cells to protection against amyloid pathology in human aging and Alzheimer's Disease**

Jonas Rybnicek[1], Yuxiao Chen[2], Milos Millic[2], JoAnne McLaurin[3], Philip L De Jager[4], Julie A Schneider[5], Yanling Wang[6], David A Bennett[6], Shreejoy Tripathy*[1, 2, 7, 8], Daniel Felsky*[2, 7, 8], Evelyn K Lambe*[1, 7, 9]

1. Department of Physiology, University of Toronto, Toronto ON, Canada
2. Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health, Toronto ON, Canada
3. Biological Sciences, Sunnybrook Research Institute, Toronto, ON M4N 3M5, Canada.
4. Center for Translational & Computational Neuroimmunology, Department of Neurology and the Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University Irving Medical Center, New York NY, USA
5. Department of Pathology, Rush University, Chicago IL, USA
6. Department of Neurological Sciences, Rush University, Chicago IL, USA
7. Department of Psychiatry, University of Toronto, Toronto ON, Canada
8. Institute of Medical Science, University of Toronto, Toronto ON, Canada
9. Department of OBGYN, University of Toronto, Toronto ON, Canada

*Co-corresponding authors
Co-Corresponding Authors email addresses:
Shreejoy Tripathy: shreejoy.tripathy@camh.ca
Dan Felsky: daniel.felsky@camh.ca
Evelyn Lambe: evelyn.lambe@utoronto.ca

Number of Figures: 5
Number of Tables: 1

Abstract

Changes in high-affinity nicotinic acetylcholine receptors are intricately connected to neuropathology in Alzheimer's Disease (AD). Protective and cognitive-enhancing roles for the nicotinic α5 subunit have been identified, but this gene has not been closely examined in the context of human aging and dementia. Therefore, we investigate the nicotinic α5 gene *CHRNA5* and the impact of relevant single nucleotide polymorphisms (SNPs) in prefrontal cortex from 922 individuals with matched genotypic and *post-mortem* RNA sequencing in the Religious Orders Study and Memory and Aging Project (ROS/MAP). We find that a genotype robustly linked to expression of *CHRNA5* (rs1979905A2) predicts significantly reduced cortical β-amyloid load. Yet, co-expression analysis shows a clear dissociation between expression of *CHRNA5* and other cholinergic genes, suggesting a distinct cellular expression profile for the human nicotinic α5 subunit. Consistent with this prediction, single nucleus RNA sequencing from 22 individuals reveals disproportionately-elevated *CHRNA5* expression in chandelier cells. These interneurons are enriched in amyloid-binding proteins and also play a vital role in excitatory/inhibitory (E/I) balance. Cell-type proportion analysis from 549 individuals demonstrates chandelier cells have increased amyloid vulnerability in individuals homozygous for the missense *CHRNA5* SNP (rs16969968A2) that impairs function/trafficking of nicotinic α5-containing receptors. These findings suggest that *CHRNA5* and its nicotinic α5 subunit exert a neuroprotective role in aging and Alzheimer's disease potentially centered on chandelier interneurons.

Keywords: Alzheimer's disease, prefrontal cortex, acetylcholine, nicotinic receptors, chandelier cells, interneurons, amyloid, attention

<u>Introduction</u>
The cholinergic system plays a critical role in the pathology of Alzheimer's disease (AD) (1), a
neurodegenerative disease marked by the accumulation of β-amyloid peptide (β-amyloid) and
neurofibrillary tangles of phosphorylated tau in the brain (2). In AD, there are well-characterized
disturbances in the excitation/inhibition (E/I) balance in cerebral cortex (3,4) arising from the
disruption of inhibitory signalling. The shift toward higher excitation in the cortex is associated
with cognitive impairment in AD.
41
The cholinergic system is an important regulator of E/I balance in the prefrontal cortex (PFC)
(5,6) and is central to one of the first mechanistic explanations of cognitive deficits in AD; the so-
called *cholinergic hypothesis* (7). In AD, there is a decrease of cortical nicotinic acetylcholine
receptor binding (8,9), and β-amyloid binding to nicotinic receptors has been postulated as a
potential mediator of AD pathology (10,11) possibly via blockade of these receptors (12,13). By
contrast, stimulation of neuronal nicotinic receptors has been found to improve survival of
primary neuronal culture exposed to β-amyloid (14), and to promote neurogenesis and improve
cognition (15,16) in preclinical models. Promoting nicotinic signalling using acetylcholinesterase
inhibitors is one of the mainstay AD treatments (17).
51
High-affinity nicotinic acetylcholine receptors are hetero-pentamer cation channels most
commonly composed of α4 and β2 subunits (α4β2) (18–20). Deep layer PFC pyramidal cells
express nicotinic receptors also containing the auxiliary α5 subunit (21–24). Nicotinic α5
subunits do not contribute to the acetylcholine binding site and cannot form functional receptors
on their own (25), requiring the binding sites provided by partner subunits α4 and β2, and
forming the α4β2α5 nicotinic receptor. The α5 subunit alters the kinetics of nicotinic receptors
(21,22) and increases their permeability to calcium ions (26,27). Importantly, β-amyloid binds
less readily to α4β2α5 than α4β2 nicotinic receptors *in vitro* (13), which raises the question of a
possible protective role of the α5 subunit in AD pathology.
61
The nicotinic α5 subunit has previously been linked to cognitive performance, with loss or
disruption of this subunit impairing performance in attentional tasks in rodents (23,28). In
humans, single nucleotide polymorphisms (SNPs) affecting the expression and
function/trafficking of the *CHRNA5* gene, which codes for the α5 subunit, have been linked to
attentional and cognitive deficits (29,30). These SNPs are also linked to smoking (31), a major
AD risk factor (32). However, the role of *CHRNA5* in aging and AD is unknown.
68
To address this critical gap, we built a multi-step model of the connections between SNPs
affecting the expression and function of *CHRNA5*, and age-related cognitive and
neuropathological phenotypes using detailed clinical and post-mortem data from the Religious
Order Study and Memory and Aging Project (ROS/MAP) (33). Next, we leveraged single-
nucleus RNAseq to determine the cell-type expression pattern of *CHRNA5* in the PFC. We then
used a gene ontology analysis of cortical patch-seq data (34) to elucidate the functional makeup
of the cell type with the highest *CHRNA5* levels in the PFC, the chandelier cells. Finally, we
probed an estimated cell type proportion dataset from the PFC (35) to assess the interaction effect
of *CHRNA5* SNPs and Alzheimer's disease pathology on this *CHRNA5*-enriched cell type. The
results of our study suggest a novel role for *CHRNA5* in maintaining the E/I balance in the
forebrain and as a potential new target for therapies aiming to promote neuronal survival in AD.
80

Methods
Study cohort

We accessed data from 2004 deceased individuals from the ROS/MAP cohort study (33), of
whom 1732 were autopsied. Both studies enrolled individuals without known dementia. ROS
enrols elderly nuns, priests, and members of clergy, whereas MAP enrols individuals from
community facilities and individual homes. Both studies were approved by an Institutional
Review Board of Rush University Medical Centre. Participants gave informed consent for annual
clinical evaluation, completed a repository consent allowing their resources to be shared, and
signed an Anatomic Gift Act for brain donation at the time of death. Most individuals assessed
were female (68%). The average age at study entry was 80.5 ± 0.16 years and the average age at
death was 89.2 ± 0.2 years. All data was retrieved from the Synapse AMP-AD Knowledge Portal
(Synapse ID: syn2580853).

Selection of candidate single nucleotide polymorphisms (SNPs)

Nicotinic α5 subunits (encoded by *CHRNA5*) do not contribute to the acetylcholine binding site
and cannot form functional receptors on their own (25). In prefrontal cortex, the nicotinic α5
subunits participate in pentameric receptors with two binding sites contributed by partner
subunits: α4 (encoded by *CHRNA4*) and β2 (encoded by *CHRNB2*) (18). Together, these subunits
form α4β2α5 nicotinic receptors. While the current focus is *CHRNA5*, we also probed the impact
of polymorphisms relevant to its receptor partners *CHRNA4* or *CHRNB2* for perspective. The
specific polymorphisms were selected based on their reported effects on gene expression,
function, or on their clinical associations. For *CHRNA5*, we selected the non-synonymous SNP
rs16969968 (minor allele frequency (MAF) = 0.33) in the coding region of *CHRNA5* and a
cluster of 6 SNPs in linkage disequilibrium (LD) in the regulatory region of *CHRNA5* (36)
represented in our study by rs1979905 (MAF = 0.43). These SNPs have previously been linked to
altered *CHRNA5* expression (36) and altered risk of nicotine addiction (31,37). For *CHRNB2,* the
T allele of the rs2072660 polymorphism (MAF = 0.23) has previously been linked to altered risk
for nicotine dependence (38), and the T allele of the rs4292956 (MAF = 0.07) with a clinical
response to treatment for smoking cessation (39). For *CHRNA4*, the G allele of the rs1044396
polymorphism (MAF = 0.45) has previously been associated with altered performance in
attention tasks as well as with the regulation of *CHRNA4* expression (40).

Genotype data preparation and imputation, quality control, generation of bulk gene
expression residuals

Details on the ROS/MAP cohort genotyping and handling of the post-mortem samples have been
previously published (41) and are described briefly together with the quality control approaches
and generation of the gene expression residuals in **Supplemental Methods**.

Final consensus diagnosis

We included the final consensus diagnosis as a covariate where relevant in our models. At the
time of death, all available clinical data were reviewed by a neurologist with expertise in
dementia, and a summary diagnostic opinion was rendered regarding the most likely clinical
diagnosis at the time of death. The methodology of the consensus diagnosis has been previously
published (42). Summary diagnoses were made blinded to all postmortem data. Case conferences
including one or more neurologists and a neuropsychologist were used for consensus on some

cases. The diagnoses we coded as: 1) NCI: No cognitive impairment (No impaired domains), 2)
MCI: Mild cognitive impairment (One impaired domain) and NO other cause of CI, 3) MCI:
Mild cognitive impairment (One impaired domain) AND another cause of CI, 4) AD:
Alzheimer's dementia and NO other cause of CI (NINCDS PROB AD), 5) AD: Alzheimer's
dementia AND another cause of CI (NINCDS POSS AD), 6) Other dementia: Other primary
cause of dementia.

Neuropathology and cognitive scores

130   A detailed description of the neuropathology and cognitive variables in ROS/MAP is included in
131   **Supplemental Methods** and on the RADC Research Resource Sharing Hub.

Single-nucleus RNA sequencing data processing

The single-nucleus gene counts and metadata available from Cain et al. 2020 (35) on synapse
(ID: syn16780177) were converted into a Seurat object (43) in R Studio for further processing.
Potential doublets were removed by filtering out cells with over 2500 detected features, and
potential dead or dying cells were removed by excluding cells expressing over 5% mitochondrial
genes. Cell type annotations were indicated in the data as described in the metadata downloaded
from the Cain et al. 2020 Synapse repository (ID: syn16780177). The snRNAseq data was log-
normalized and matched with genotype data (n = 22 individuals). *CHRNA5* expression was
averaged per cell type per individual and *CHRNA5* levels in different cell types were then
compared between cell types by one-way ANOVA with Tuckey's post-hoc t-test. To prevent bias
for rare cell types in calculating average *CHRNA5* expression per cell type per individual, cell
types with fewer than 100 individual cells represented in the original data (not aggregated) were
removed (the layer 5 FEZF2 ET cell type was excluded from analysis as only 93 cells of this type
were present in the genotype-matched snRNAseq dataset). The number of cells per cell-subtype
cluster in the genotype-matched snRNAseq dataset can be viewed in the supplementary materials
(**Supplemental Table S1**). The identity of chandelier cells was further validated by assessing the
expression of SCUBE3, a chandelier cell marker (44), across the cell subtypes.

Gene ontology of chandelier cell genes

A set of genes which are upregulated in PVALB+ chandelier cells versus PVALB+ non-
chandelier cells (basket cells) was previously generated by Bakken and colleagues (34). In short,
the authors utilized the patch-seq method (combines electrophysiological patch-clam recording
with RNA sequencing and morphological analysis) to characterize the transcriptional differences
between cortical PVALB+ chandelier cells, and basket cells across multiple species (human,
mouse, and marmoset). In our gene ontology analysis, we filtered the list of differentially
expressed genes (DEGs) for those specifically upregulated in human PVALB+ chandelier cells
versus human basket cells (remaining DEG n = 222). To determine the ontology of this
upregulated gene set we used the Gene Ontology Resource (geneontology.org), querying
specifically for molecular function.

Estimates of relative cell type proportions from bulk DLPFC RNAseq

Estimates of cell-type proportions from bulk DLPFC RNAseq data from 640 ROS/MAP
participants was performed and described by Cain et al. 2020 (35). In brief: The authors
developed a custom regression-based consensus model, CelMod, to extract cell cluster specific

genes from the snRNAseq dataset from 24 ROS/MAP participants, and then used these genes to
estimate the proportions of different cell subtypes in the bulk DLPFC RNAseq dataset from 640
ROS/MAP individuals. The proportions were estimated using a large set of marker genes
extracted by CelMod for each cell subtype. This method was used to ensure that changes in the
expression of a small number of major marker genes would not skew the proportions of any
given cell type. The deconvolved cell-type proportion data from the DLPFC was available on
request from the research group (Cain et al. 2020 personal communication). We matched the cell
type proportion data with genotype, bulk DLPFC RNAseq, and neuropathology (brain levels of
β-amyloid and tau) data (final n = 549 individuals). At the time of analysis full cell-type
annotations (like that in the snRNAseq data) were only available for the different classes of
inhibitory neuron proportions. The estimated proportions of a GABAergic neuron subtype,
marked by its co-expression of (among other marker genes) PVALB+/LHX6+/THSD7A+, were
determined (Cain et al. 2020 personal communication) to represent chandelier cell proportions in
DLPFC. Observed differences in the estimated proportions of this cell subtype represent a
difference in the proportion of this cell subtype in the broad cell class (GABAergic neurons).
Further information on the cell type proportions is provided in **Supplemental Methods**.

Statistical approaches

EQTL and conditional eQTL analyses of candidate SNPs and gene expression in DLPFC

All analyses were performed in R (R version 4.0.2) (R Core Team, 2020) unless otherwise
indicated. For eQTL analyses (n = 924), linear regression was used to model batch- and technical
covariate-corrected gene expression residuals against SNP allele counts, co-varying for the first
10 genomic principal components (fine population structure), biological sex, age at death, and
PMI. Bonferroni correction was applied across tests (6 cis-eQTL tests performed). To assess the
effects of the *CHRNA5* haplotypes tagged by rs16969968 and rs1979905 on *CHRNA5* expression
we used a nested ANOVA analysis with Sidak's post-hoc test for multiple comparisons in
GraphPad Prism.

Association analysis between SNPs, gene expression, AD neuropathology, and cognition

To map the multi-omic relationships between candidate SNPs (n_{SNPs}= 6), cholinergic receptor
gene expression (n_{genes}= 3), post-mortem AD-related neuropathologies ($n_{pathologies}$=2), and global
cognitive performance at the population level, we fit a series of independent linear models,
maximizing sample sizes for each combination of data types, and including appropriate technical
and biological covariates. Only individuals with genotype data were included in these models
(**Supplemental Fig. S1**). SNPs were modelled using additive allelic dosage coding (i.e., 0,1,2).
For the associations of SNPs with β-amyloid and tau pathology (n = 1317), sex, age at death,
post-mortem interval (PMI), genotyping batch, the first 10 genomic PCs, final clinical consensus
diagnosis and *APOE* genotype were included as covariates. For the association of amyloid with
tangle pathology (n = 1317) sex, age at death, PMI, final clinical consensus diagnosis and *APOE*
genotype were included as covariates. For the associations of SNPs with global cognitive
performance at time of death (n = 1448), sex, age at death, years of education, genotyping batch,
and the first 10 genomic PCs were included as covariates. To aid in the interpretation of SNP and
gene expression effects in the context of co-occurring neuropathologies and cognitive
dysfunction, we also modelled the effects of amyloid and tau measures on the last cognition score
in our dataset (n = 1253); sex, age at death, PMI and years of education were used as covariates.

For the association of gene expression residuals with neuropathology (n = 914), sex, age at death,
PMI, final clinical consensus diagnosis and *APOE* genotype were used as covariates. And for
association with global cognition at time of death, we used a linear regression model of
CHRNA4, CHRNA5 or *CHNB2* against the last global cognition score using matched cognition
and gene expression data (n = 887). Sex, age at death, PMI, years of education and *APOE*
genotype were used as covariates.

In total, we fit 36 models for the network analysis (**Fig. 1F**) All two-sided *p*-values for
terms of interest were corrected using the Benjamini and Hochberg false discovery rate (FDR)
approach (45) of the p.adjust function in R, with corrected $p_{FDR}<0.05$ considered significant.

We used a Chi-square test in GraphPad Prism to study the association between *CHRNA5*
SNPs genotype and participant smoking status at baseline.

Supplemental Fig. S1 describes the specific numbers and exclusions of ROS/MAP in
each stage of our analyses.

Network visualization of multi-omic mapping

221 To integrate our multi-level association analyses and aid global interpretation, we constructed a
222 network plot summarizing relationships among the different data types analysed (**Fig. 1F**).
223 Network nodes correspond to individual independent and dependent variables, and each edge
224 represents an association derived from one of the linear models described above.

Cross-correlation of cholinergic gene expression and network construction

To assess the correlation of *CHRNA5* to other major cholinergic genes in the bulk DLPFC
dataset, we performed a series of Pearson-correlation analyses on the residualized expression of
the different cholinergic genes and used the resulting correlations to construct a correlation
network (**Fig. 1G**). Network nodes correspond to the different cholinergic genes and the edges
represent the size of the correlation coefficient (r). Only correlations significant after adjustment
for false discovery rate (56 comparisons) are shown in the network graph.

Analysis of single-nucleus RNAseq data from DLPFC

The single-nucleus RNAseq dataset published by Cain et al. 2020 contained cells clustered into
37 subtypes. As we were interested in genotypic effects on *CHRNA5* expression in neurons
specifically, to narrow the scope of the analysis, non-neuronal cell subsets (microglia : Micro.1 –
Micro.6, oligodendrocyte:Olig.1 – Olig.4, endothelial cells: Endo.1 – Endo.4, astrocyte cells:
Astr.1 – Astr.5) were grouped into cell-subtype categories (Microglia, Oligodendrocytes,
Endothelial cells, Astrocytes) while neuronal cells were kept in their subtypes (except the L5
RORB IT subtype which was formed by grouping the Exc.Exc.RORB_L5_IT_1 and
Exc.Exc.RORB_L5_IT_2 subtypes) as defined in Cain et al. 2020. Using the aggregate function
in R, the log-normalized DLPFC snRNAseq *CHRNA5* expression (n = 24) was averaged per cell
type (subtype) per individual (using the unique participant "projid" IDs) as follows:
[aggregate(*CHRNA5* ~ projid + subtype, data, mean)].

To assess the expression of *CHRNA5* across the different cell types, we used a one-way
ANOVA with Tukey's post-hoc t-test on the *CHRNA5* expression previously averaged per cell
type per individual.

To assess the association between the rs1979905 genotype and *CHRNA5* expression in the
DLPFC snRNAseq data, we matched the *CHRNA5* expression data averaged per cell type per
individual with genotype data (final = 22). We then used general linear models covaried for sex,
age at death, PMI and genotyping batch to assess the association between the rs1979905
genotype and *CHRNA5* expression in the different cell types.

252   Interaction modelling of neuropathology, CHRNA5 genotype, and bulk-estimated cell type
253   proportions

To assess the association between amyloid and tau neuropathology, and chandelier cell
proportions, we used a linear regression model of these variables covaried for sex, age at death
and PMI. A linear regression model with the same covariates was also used to assess the
association between *CHRNA5* expression and chandelier cell proportions. To test the association
between rs1979905 and rs16969968 genotype with chandelier cell proportions, we used a linear
regression model covaried for sex, age at death, post-mortem interval, *APOE* genotype and final
clinical consensus diagnosis. Finally, we used a linear interaction model to test the association
between genotype-neuropathology interaction on the proportion of chandelier cells, covaried for
sex, age at death, post-mortem interval, *APOE* genotype, final clinical consensus diagnosis and
the first 10 genomic PCs: [Cell type proportion ~ Genotype * Neuropathology + sex + age at
death + PMI + *APOE* genotype + diagnosis + PC1 + … + PC10].
265
Results
Expression of α4β2α5 receptor component genes is affected by single nucleotide
polymorphisms

To identify effects in aging and dementia of gene variants previously shown in younger adults to
influence *CHRNA5* expression (36) and α5 coding (26,46), we examined brain expression
quantitative trait loci (eQTL). The variants were in weak-moderate linkage disequilibrium in our
European ancestry sample (r^2=0.34), in agreement with previous work (36). We also found that
dosage of the A allele of the missense SNP rs16969968 (minor allele frequency (MAF) = 0.33) in
the coding region of *CHRNA5* (**Fig 1B**) was associated with lower *CHRNA5* expression (t = -
13.61 *p* = $1.63*10^{-38}$), consistent with existing data (47). A different SNP haplotype in the
regulatory region upstream of *CHRNA5* (**Fig 1B**), denoted here by the A allele of the tag SNP
rs1979905 (MAF = 0.43), was associated with higher *CHRNA5* expression (t= 27.47, *p* =
$5.36*10^{-121}$) (**Fig. 1C**). Furthermore, analyses of the coding-SNP rs16969968 and the regulatory-
SNP rs1979905 together (**Fig. 1D**) showed that *CHRNA5* expression is predominantly regulated
by the regulatory-SNP rs1979905 rather than the coding-SNP, rs16969968, as all rs1979905 A
allele non-carriers showed similar levels of *CHRNA5* mRNA regardless of the rs16969968 A
allele (Nested one-way ANOVA: F(2) = 229.6; Šidák's post-hoc test for multiple comparisons:
rs1979905 A1 vs. A0 *p* = $8*10^{-6}$, A2 vs. A1 *p* = 0.004, A2 vs A0 *p* = $4*10^{-6}$), suggesting that the
effect of rs1979905 A on *CHRNA5* expression is independent of the rs16969968 SNP genotype.
This was also demonstrated using a conditional eQTL model, where the effect of rs16969968 A
allele on *CHRNA5* expression was lost when co-varying for rs1979905 A allele (rs16969968A: t
= -1.027, p = 0.304; rs1979905A: t = 21.927, *p* = $8.194*10^{-86}$). No association between *CHRNA5*

expression and disease state was detected (**Supplemental Fig. S2**). No trans-eQTL effects were
detected (**Fig. 1E**) between either of these SNPs and the expression of required partner nicotinic
subunit genes, *CHRNA4* and *CHRNB2*.

To assess the α4 and β2 nicotinic subunits required for the formation of α5-containing α4β2α5
receptors, we extended our eQTL analyses to SNPs in *CHRNA4* and *CHRNB2*, focusing on those
associated with altered gene expression or clinical effects(39,40,48). Without exception, eQTL
SNP effects for these genes were weaker than those of rs16969968 and rs1979905 for *CHRNA5*:
the T allele of *CHRNB2* intronic variant rs2072660 (MAF = 0.23) was associated with lower
CHRNB2 expression (t = -4.738, $p = 2.5*10^{-6}$), and a similar association with *CHRNB2*
expression was seen with the T allele of the *CHRNB2* non-coding variant rs4292956 (MAF =
0.07) (t = -4.961, $p = 8.4*10^{-7}$) (**Fig. 1E**). For *CHRNA4,* the G allele of missense variant
rs1044396 (MAF = 0.45) in the coding region of *CHRNA4* was associated with higher *CHRNA4*
expression (t = 3.416, $p = 0.0007$). We also used the Gene Query function of the xQTLServe
online tool (DLPFC of 534 ROS/MAP participants) to identify the T allele of the intronic variant
rs45497800 as associated with decreased *CHRNA4* expression (t = -6.92, $p = 1.32 * 10^{-11}$). We
then replicated this association in our larger cohort of 924 ROS/MAP individuals (MAF = 0.07)
(t = -4.537, $p = 6.48*10^{-6}$) (**Fig. 1E**). No associations were found between these SNPs and the
expression of other high-affinity nicotinic receptor subunit genes (**Fig. 1E**).

CHRNA5 polymorphisms are not associated with smoking status in this largely non-
smoking population

The percentage of participants identified as never, previous, and current smokers (**Supplemental**
Methods) was 68.2, 29.4, and 2.4 respectively. To investigate previously-reported (29,49)
associations between genotype for the *CHRNA5* SNPs and smoking status at baseline
(current/former/never smoked), we used a Chi-squared test and found no relationship
(rs1979905_A : $\chi2(4) = 1.575$, p = 0.813; rs16969968_A: $\chi2(4) = 1.317$, p = 0.858) in this
largely non-smoking population. Smoking status was not used in further analysis, unless
specifically indicated.

A *CHRNA5* polymorphism is negatively associated with brain β-amyloid levels

To address interrelationships among nicotinic subunit expression, nicotinic SNPs, and
neuropathological and cognitive phenotypes, we used clinical and post-mortem data in a multi-
step model, the inclusion and exclusion criteria for this model can be found in **Supplemental**
Fig. S1. As illustrated in **Fig. 1F**, both β-amyloid and tau pathology were negatively associated
with global cognitive performance proximal to death (tau: t = -18.87 $p = 5.77 *10^{-70}$; amyloid: t =
-7.88, $p = -6.99*10^{-15}$) and positively associated with each other (t = 12.59, $p = 2.14*10^{-34}$). Of
the SNPs examined, only the SNP increasing *CHRNA5* expression had a significant association
with AD neuropathology, with the A allele of the regulatory-SNP rs1979905 significantly
negatively associated with β-amyloid load after false discovery rate (FDR) correction for multiple
comparisons (t = -2.81, $p_{FDR} = 0.015$, corrected for 36 models per (45)). Intriguingly, expression
of the *CHRNA5* gene itself showed a direct negative association with β-amyloid load prior to
FDR correction (t = -2.23, $p = 0.026$), whereas the expression of the major nicotinic subunit
genes *CHRNA4* and *CHRNB2* showed significant positive associations with the last global
cognition score, which remained significant after FDR correction (*CHRNA4*: t = 2.98, $p_{FDR} =$
0.013; *CHRNB2*: t = 3.43, $p_{FDR} = 0.003$). Conversely, the rs2072660 T allele, associated with

lower *CHRNB2* expression, was negatively associated with the last global cognition score prior to
FDR correction (t = -1.99 *p* = 0.047). A similar negative association with the last global cognition
score (t = -2.3, *p* = 0.022) was found for the T allele of the *CHRNB2* SNP rs4292956.

CHRNA5 expression does not tightly correlate with other components of the cholinergic
system
To further investigate the interrelationships among *CHRNA5* and the major nicotinic subunits as
well as other components of the cholinergic system, we performed a separate series of gene
expression correlation analyses. Most of the major components of the cholinergic system which
were detected in bulk DLPFC data of the ROS/MAP individuals (*CHRNA2*, *CHRNA4*, *CHRNA7*,
CHRNB2, *CHRM3*, *CHRM1* and *ACHE*) showed significant positive correlation with each other
(**Fig. 1G**). By contrast, *CHRNA5* stood out as showing no positive correlation with any of the
other major cholinergic genes and only weak negative correlations with the expression of
CHRNB2 and *CHRM1* (**Table 1**). Considering that the expression of *CHRNA5*, *CHRNB2,* and
CHRNA4 are required for the assembly of the high-affinity α4β2α5 nicotinic receptor, it was
surprising to see that *CHRNA5* expression was not positively correlated with either *CHRNA4* or
CHRNB2 (**Fig. 1G** and **Table 1**). Therefore, we next investigated whether this lack of correlation
may arise from differences in the cell-type specific expression of *CHRNA5* compared to the
major nicotinic receptor subunits, *CHRNA4* and *CHRNB2,* which are more broadly expressed.
349
CHRNA5 shows stronger expression in chandelier interneurons than most other cell classes

To assess the cell-type specificity of *CHRNA5* expression in the ROS/MAP cohort, we calculated
the average *CHRNA5* expression per cell type per individual using the genotype-matched single-
nucleus RNAseq data available from the DLPFC in a subset of 22 ROS/MAP participants (35)(2
individuals lacked genotyping data). In this small dataset, *CHRNA5* was expressed at a low level
across a number of different excitatory, inhibitory, and nonneuronal cell types (**Fig. 2A**), with
significantly higher expression in inhibitory PVALB+ chandelier cells (as identified by Cain et
al. 2020). Chandelier cells had significantly-higher expression of *CHRNA5* compared to most
other cell types (One-way ANOVA: F(21) = 3.439, *p* = 6.1*10^{-7}; Tukey's post-hoc t-test
Chandelier cells vs. 18 out of 19 other cell types: *p* < 0.05) (**Fig. 2A**). By contrast, chandelier cell
expression of *CHRNA4* and *CHRNB2* were at a similar level in chandelier cells to their
expression levels in many other cell types (**Fig. 2B**). To confirm the identity of the chandelier as
defined by Cain et al 2020 (35), we assessed the expression of *SCUBE3*, a specific marker of
chandelier cells (44), across the different cell type found in the ROS/MAP snRNAseq dataset,
and found the highest expression of this gene in the chandelier cells (**Sup. Fig. S3**).

To confirm the novel finding that chandelier cells show stronger *CHRNA5* expression compared
to other classes of neurons, we probed publicly available cell-type specific gene expression
databases of human brain tissue. Using the Allen Institute SMART-seq single-cell
transcriptomics data from multiple cortical areas https://celltypes.brain-
map.org/rnaseq/human_ctx_smart-seq we found *CHRNA5* expression to be highest in a
PVALB+/SCUBE3+ inhibitory cell type (0.06 trimmed mean *CHRNA5* expression) likely
representing chandelier cells (44), and in a co-clustering PVALB+/MFI+ cell type (0.06 trimmed
mean *CHRNA5* expression). Highest expression of *CHRNA5* in chandelier cells compared to all
other cell types is also replicated in the Seattle Alzheimer's Disease Brain Cell Atlas
(https://knowledge.brain-
map.org/data/5IU4U8BP711TR6KZ843/2CD0HDC5PS6A58T0P6E/compare?cellType=Whole

Taxonomy&geneOption=CHRNA5&metadata=Cognitive Status&comparison=dotplot.) In the
Human Protein Atlas database (brain single cell tissue)
https://www.proteinatlas.org/ENSG00000169684-CHRNA5/single+cell+type/brain, *CHRNA5*
showed highest expression in a PVALB+ inhibitory cell type (c-41) which also showed highest
expression of *SCUBE3* (Inhibitory neurons c-41, 15.1 normalized *CHRNA5* transcripts per
million), likely also representing chandelier cells.

To investigate the cell type-specificity of the *CHRNA5* eQTL effects of the regulatory-SNP
rs1979905 in the single nucleus data from ROS/MAP, we stratified the averaged *CHRNA5*
expression by genotype for the rs1979905 A allele. We found that higher allelic dosage of the
rs1979905 A allele was associated with greater *CHRNA5* expression (**Fig. 2C**), and that this
pattern was most pronounced in subtypes of layer 5 (L5 RORB IT: t = 2.460, p = 0.0249) and
layer 6 (L6 IT THEMIS: t = 2.402, p = 0.028) excitatory neurons (**Fig. 2D**). In the stronger
CHRNA5-expressing PVALB+ chandelier cells, however, the association did not reach
significance. Other neuronal cell types appeared to diverge completely from the typical eQTL
pattern of rs1979905 (**Fig. 2C,D**). This analysis suggests that only a subset of cell-types
contribute to the stepwise expression pattern observed in the prefrontal cortex by rs1979905
genotype.

Cell type specific data from ROS/MAP and the Allen database supports the hypothesis that
CHRNA5 possesses a distinctive expression pattern with enrichment in chandelier cell
interneurons, compared to its more abundant and widely-expressed subunit partners.

Chandelier cells are significantly enriched for genes interacting with β-amyloid

To investigate the potential contribution of chandelier cells to β-amyloid processing, especially
one that might be driven by nicotinic α5-containing receptors, we assessed the molecular function
of genes which are significantly upregulated in chandelier cells compared to PVALB+ non-
chandelier cells (basket cells). For this analysis we took advantage of a list of 222 such genes
previously generated by Bakken and colleagues (34), who investigated the cellular identities of
chandelier neurons in the human cortex. Gene ontology analysis revealed this gene list to be
significant enriched for genes defined as "amyloid-beta binding" (fold enrichment = 7.79, p_{FDR} =
0.01) (**Fig. 3A**), including *SORL1,* an endocytic receptor that directs the amyloid precursor
protein away from the amyloidogenic pathway (50–52) and *EPHA4,* a receptor tyrosine kinase
involved in amyloid regulation (53). Since α5-containing nicotinic receptors are highly
permeable to calcium ions (26,27), we note that the gene ontology analysis also showed
chandelier neurons to be significant enriched for genes with a "calcium ion binding" molecular
function (fold enrichment = 3.77, p_{FDR} = $4.12*10^{-6}$) (**Fig. 3B**), including genes potentially
protective against amyloid pathology such as *MME and SPOCK1*. Two genes are at the
intersection of these functional pathways in chandelier cells, *PRNP and CLSTN2*. The former has
been implicated in nicotinic receptor-mediated regulation of amyloid (54–57) and the latter has
been implicated in cortical GABAergic neurotransmission (58) These findings underscore the
importance of investigating the relationship between the functional status of nicotinic α5 subunits
and chandelier neuron vulnerability to AD neuropathology.
416
A genotype-specific reduction in proportion of chandelier cells with increasing brain β-
amyloid levels

To determine whether impaired function/trafficking α5-containing nicotinic receptors might
promote chandelier neuron vulnerability to neurodegeneration, we examined the interaction of
rs16969968 genotype and AD neuropathology on estimated proportions of chandelier neurons in
the bulk RNAseq dataset. This investigation was based on cell type proportion estimates for
chandelier cells and several other interneuron subclasses, from sets of single-cell-informed
marker genes. The cell type estimates were derived from the bulk DLPFC RNAseq data of a
subset of 640 ROS/MAP participants. Overall, β-amyloid levels were negatively associated with
the proportion of chandelier cells (t = -4, $p = 7.26*10^{-5}$). However, the missense rs16969968 A
allele homozygotes showed significantly lower chandelier cell proportions with increasing β-
amyloid load, compared to rs16969968 A allele non-carriers (interaction term t = -2.842, $p =$
0.005) (**Fig. 4A,B,C**). In a secondary analysis, there is a suggestion of an opposite relationship of
rs1979905 A allele and β-amyloid levels with chandelier cells but this does not reach statistical
significance (interaction term t = 1.81, $p = 0.071$) (**Fig. 4D,E,F**). The observed relationships
between chandelier cell proportions, *CHRNA5* genotypes and amyloid were not altered by the
inclusion of smoking status as a covariate in the analysis. Chandelier cell proportions did not
correlate with tau pathology (t = -0.4, $p = 0.687$), nor was there an interaction of *CHRNA5*
genotype and tau pathology with chandelier cell proportions (data not shown).
To assess whether the interaction between β-amyloid levels and *CHRNA5* SNPs was driven by
the effects of these SNPs on *CHRNA5* expression, we assessed the effect of the interaction of
CHRNA5 levels and β-amyloid load on chandelier cell proportion but found no significant effect
(interaction term, $t = 0.5$, $p = 0.619$). This suggests that the genotype-specific association
between amyloid load and chandelier cell proportion is more likely driven by changes in nicotinic
α5 protein structure and/or trafficking (59) as a consequence of having two copies of the
missense SNP in *CHRNA5*, rather than by the altered *CHRNA5* expression associated with the
rs1979905 SNP genotype.
444
445   The schematic in **Fig. 5** illustrates a working model of the impact of the rs16969968 A allele
446   homozygosity for chandelier cell vulnerability, as well as example mechanisms enriched in
447   chandelier cells and known to alter β-amyloid processing. **Supplemental Table S2** summarizes
448   the impacts of the regulatory and functional SNPs of *CHRNA5*.
449
<u>Discussion</u>
We examined human prefrontal cortical *CHRNA5* expression in aging, probing the connections
between SNPs affecting the expression and function/trafficking of the nicotinic α5 subunit gene
and AD-related neuropathology. The aging prefrontal cortex demonstrates strong eQTL effects of
the common regulatory-SNP rs1979905, and we show that the A allele of rs1979905 is associated
with lower levels of brain β-amyloid. Single-nucleus RNAseq data revealed that chandelier cells
have the greatest abundance of *CHRNA5* in human prefrontal cortex. These neurons are
significantly enriched in amyloid-binding proteins, including some that may be activated via
nicotinic receptors. Lastly, we examined the impact of AD neuropathology on the proportions of
chandelier neurons and determined that the α5-altering common coding-SNP rs16969968 renders
this interneuron population vulnerable to β-amyloid levels. Our findings, summarized in the
working model in **Fig 5**, as well as **Supplemental Table S2**, suggest a potential cell-type specific
neuroprotective role for C*HRNA5* to reduce β-amyloid levels and toxicity.
463
Inhibitory signalling is disrupted in Alzheimer's disease
The disruption of the excitatory/inhibitory balance in the cortex is a hallmark of AD pathology
and is associated with cognitive AD symptoms (3,4). Previous studies have shown disruption of

cortical inhibitory signalling in AD stems from the alteration of the activity of inhibitory neurons
and inhibitory cell loss (60–62). The susceptibility of inhibitory neurons to AD pathology is not
uniform, however. While studies have shown significant drops in somatostatin-positive
interneurons in the cortex of AD patients (63,64), the numbers of parvalbumin-positive cortical
interneurons, including the PVALB+ chandelier cells, appear comparatively more resilient to AD
pathology (64,65). Our study expands on these findings, demonstrating that the preservation of
chandelier cells in AD pathology depends on the genotype of SNP in a nicotinic receptor subunit.
The rs16969968 SNP is a missense mutation found to functionally alter the α5-containing
nicotinic receptors in cell systems and *in vivo*, through altered channel biosynthesis, trafficking,
properties and/or modulation (26,46,59,66,67). Recent optophysiological work suggests that the
nicotinic α5 subunit accelerates the endogenous cholinergic response in prefrontal cortex and
protects it against desensitization (21), but little is known about endogenous cholinergic
modulation of chandelier neurons.
480
Chandelier cells are vulnerable to β-amyloid pathology
Chandelier cells are a specialized subtype of PVALB+ interneurons. They differ anatomically
from the PVALB+ basket cells by their large number of vertically oriented axonal cartridges,
which specifically innervate the axon initial segments of pyramidal neurons (68,69). New
evidence suggests chandelier cells regulate excitation dynamics of neuronal networks (70).
Impairment of these neurons has been implicated in diseases involving pathological excitation in
the cortex, such as epilepsy (71,72) and AD (3,4,65). Our RNASeq findings are in agreement
with previous work showing that the inhibitory output of chandelier cells is sensitive to β-
amyloid pathology (73) but unaffected by tau pathology (74). Chandelier cell axons near β-
amyloid plaques have been found to show deformations, and pyramidal neurons proximal to
plaques show loss of inhibitory input onto their axon initial segments (73). Our findings suggest
that in people homozygous for A allele of the missense *CHRNA5* SNP rs16969968 (11% of the
ROS/MAP participants), this vulnerability of chandelier cells to β-amyloid pathology may be
exacerbated, possibly leading to cell death.
495
Potential mechanisms for a neuroprotective effect of α5-containing nicotinic receptors
Our results suggest that polymorphisms affecting *CHRNA5* expression and function, may alter
both the total β-amyloid levels in the brain, and alter the susceptibility of specific *CHRNA5*-
expressing cell types, such as the chandelier cells, to β-amyloid-mediated toxicity. One possible
explanation for these observations may be the lowered binding of β-amyloid to the α4β2α5
nicotinic receptors (13) expressed by these cells. This protection against β-amyloid binding and
inhibition of the nicotinic response (12) could promote resilience of nicotinic signalling in the
chandelier cells, potentially leading to improved cell survival (14) in AD pathology. Furthermore,
since α4β2α5 nicotinic receptors support higher conductance of calcium ions into the cell (26,27),
another putative neuroprotective mechanism of the α5 subunit may be through driving possible
calcium-dependent neuroprotective pathways in the neurons which express the α4β2α5 nicotinic
receptors (75,76). Such calcium-regulated pathways may include *MME*, *SORL1*, *SPOCK1* or
PRNP, which are specifically enriched in PVALB+ chandelier cells compared to PVALB+ non-
chandelier cells (34), and which have been previously suggested to alter β-amyloid production
and clearance (51,54,55,77).
511
Caveats and opportunities for additional investigation
While the ROS/MAP database offered an opportunity to assess the impact of *CHRNA5*
expression and *CHRNA5*-related SNPs on AD pathology in a large sample, some caveats exist.

As *CHRNA5* expression has previously been shown to be important for animal performance in
demanding attentional tasks (23,28), one of the critical limitations of our study is that a robust
attention assessment of the ROS/MAP individuals was not part of the study design, potentially
explaining the lack of any association between *CHRNA5* expression or polymorphisms and a
cognitive readout. Furthermore, there is limited knowledge on the brain functions regulated by
the inhibitory chandelier cells, which have the highest *CHRNA5* expression of the cell types
assessed in our study. One possibility is that due to their strong effects on the regulation of the
action potential generation at the axonal initial segment of pyramidal cells (68,69), chandelier
cells may be involved in seizure pathology (71,72). However, due to the lack of robust epilepsy
or seizure-related measurements in the ROS/MAP cohort, hypotheses related to this could not be
tested. A previous study in rodents (66) has suggested a link between the rs16969968 SNP and
certain features of psychosis, namely hypofrontality of schizophrenia patients (78), which is
marked by decreased blood flow to the prefrontal cortex, suggesting a decrease in the activity in
this area. However, hypotheses relating to this potential link could not be assessed in our current
study due to a lack of data on the diagnoses of the ROS/MAP participants for schizophrenia or
related disorders.
531
While the ROS/MAP dataset presented an opportunity to study the effects of rs16969968 on a
background of an unusually-low smoking prevalence (79), future work would benefit from more
robust assessment of smoking history. Although most prevalent in the prefrontal cortex, the
α4β2α5 receptor is not the only type of α5-containing nicotinic receptor. Another type of interest
is the α3β4α5 receptor, which is expressed primarily in the habenula (27), and intriguingly has all
its subunits within the same locus (80). Unfortunately, expression data for the relevant *CHRNA3*
and *CHRNB4* genes were not included in the prefrontal bulk RNAseq dataset, complicating
investigation of hypotheses pertaining to the *CHRNA3, CHRNA5, CHRNB4* locus. β-amyloid
pathology affects other types of nicotinic receptors besides the α4β2* subtype, including the
abundant and widely-expressed low-affinity homomeric α7 receptor (10). However, since the
activity and β-amyloid-sensitivity of the neuronal α4β2* receptor can be further modified by the
inclusion of the auxiliary subunit α5 (13), we focus on the high-affinity nicotinic receptor as a
potentially rewarding target of study in the context of altered nicotinic signalling in AD.

A limitation of the single-nucleus RNA sequencing data (35) was its relatively low number of
individuals, limiting the robustness of comparing the effects of rs1979905 on *CHRNA5*
expression across the different cell types, and preventing a similar examination of cell-type
specific effects of rs16969968 on *CHRNA5* expression in the ROS/MAP dataset. This limitation
should be considered when interpreting the findings of our study. Furthermore, the low numbers
of individuals in the single-nucleus cohort also prevented a comparison between our findings
from the estimated cell-type proportions in the bulk RNAseq data and the actual proportions of
different cell types present in the single-nucleus data. Future studies involving more subjects with
single-nucleus RNAseq data could extend our findings on the interaction association of the
rs16969968 genotype and β-amyloid with chandelier cell proportions.

Since cortical cell-type proportions were estimated using patterns of marker gene expression (35),
the decreased chandelier cell proportions may instead reflect reductions of chandelier cell
cartridges in AD (65). While the disruption of cortical E/I balance would remain similar, a
different interpretation of our data would be that the rs16969968 A allele homozygous genotype
exacerbates chandelier cartridge loss and results in lower expression of chandelier-cell-specific
marker genes in individuals with elevated β-amyloid.

Finally, while SNP exploration provides novel insight into the relationship between *CHRNA5* and neuropathology in aging, this work is limited by being correlational. Moreover, gene expression does not necessarily denote protein levels in AD brains (81) and thus differences in nicotinic receptor gene expression may not fully predict receptor levels or binding (82). Many questions remain about the mechanisms by which nicotinic receptors in chandelier cells regulate amyloid processing, as well as the consequences for this cell population when these mechanisms are disrupted. Work in model systems and larger snRNAseq datasets will be necessary to test specific hypotheses raised in this work.

Summary and implications

A growing body of work suggests that cortical excitability is perturbed early in Alzheimer's disease through impairment of inhibitory interneurons (60,83). *CHRNA5* is positioned to modulate the overall excitability of the prefrontal cortex in two ways: through the excitation of a population of deep layer cortical pyramidal neurons (21,84,85) that send projections throughout prefrontal cortex and, as ascertained from our findings in this study, through the excitation of specific subsets of cortical interneurons, the chandelier cells. Our findings suggest that *CHRNA5* is involved in Alzheimer's disease neuropathology. The A allele of the *CHRNA5* regulatory-SNP rs1979905 is associated with higher expression of *CHRNA5* and with reduced β-amyloid load in the brain. In parallel, the A allele of the missense SNP, rs16969968, is associated with fewer chandelier cells in individuals with high β-amyloid levels, suggesting that differences in the protein structure of *CHRNA5* contributes cellular resiliency to β-amyloid pathology. This combination suggests neuroprotective roles of *CHRNA5* in β-amyloid pathology and makes *CHRNA5* a target for therapies aiming to improve neuron survival in Alzheimer's disease.

Acknowledgements: This work was supported by the Canadian Institutes of Health Research (CIHR; EKL; PJT-153101, MOP89825; EKL; NGN-171423, ST; PJT-428404, DF), Krembil Foundation (DF, ST), Koerner Family Foundation (DF), CAMH Discovery Fund (DF, ST), Kavli Foundation (ST), McLaughlin Foundation (ST), Natural Sciences and Engineering Research Council of Canada (RGPIN-2020-05834 and DGECR-2020-00048; ST). JM is supported by the Canadian Research Chairs program, as a Tier 1 Chair. ROS/MAP is supported by P30AG10161, P30AG72975, R01AG15819, R01AG17917. U01AG46152, U01AG61356. ROS/MAP resources can be requested at https://www.radc.rush.edu

Conflicts of Interest: None

References

1. Hampel H, Mesulam M-M, Cuello AC, Farlow MR, Giacobini E, Grossberg GT, et al. The cholinergic system in the pathophysiology and treatment of Alzheimer's disease. Brain [Internet]. 2018 Jul 1;141(7):1917–33. Available from: https://pubmed.ncbi.nlm.nih.gov/29850777

2. Busche MA, Hyman BT. Synergy between amyloid-β and tau in Alzheimer's disease. Nat Neurosci [Internet]. 2020;23(10):1183–93. Available from: https://doi.org/10.1038/s41593-020-0687-6

3. Lauterborn JC, Scaduto P, Cox CD, Schulmann A, Lynch G, Gall CM, et al. Increased excitatory to inhibitory synaptic ratio in parietal cortex samples from individuals with Alzheimer's disease. Nat Commun [Internet]. 2021;12(1):2603. Available from: https://doi.org/10.1038/s41467-021-22742-8

4. Vico Varela E, Etter G, Williams S. Excitatory-inhibitory imbalance in Alzheimer's disease and therapeutic significance. Neurobiol Dis [Internet]. 2019;127:605–15. Available from: https://www.sciencedirect.com/science/article/pii/S0969996118307599

5. Pafundo DE, Miyamae T, Lewis DA, Gonzalez-Burgos G. Cholinergic modulation of neuronal excitability and recurrent excitation-inhibition in prefrontal cortex circuits: implications for gamma oscillations. J Physiol [Internet]. 2013/07/01. 2013 Oct 1;591(19):4725–48. Available from: https://pubmed.ncbi.nlm.nih.gov/23818693

6. Obermayer J, Luchicchi A, Heistek TS, de Kloet SF, Terra H, Bruinsma B, et al. Prefrontal cortical ChAT-VIP interneurons provide local excitation by cholinergic synaptic transmission and control attention. Nat Commun [Internet]. 2019;10(1):5280. Available from: https://doi.org/10.1038/s41467-019-13244-9

7. Bartus RT, Dean RL, Beer B, Lippa AS. The Cholinergic Hypothesis of Geriatric Memory Dysfunction. Science (80-) [Internet]. 1982 Jul 30;217(4558):408–14. Available from: https://doi.org/10.1126/science.7046051

8. Court J, Martin-Ruiz C, Piggott M, Spurden D, Griffiths M, Perry E. Nicotinic receptor abnormalities in Alzheimer's disease. Biol Psychiatry [Internet]. 2001;49(3):175–84. Available from: https://www.sciencedirect.com/science/article/pii/S0006322300011161

9. Nordberg A, Winblad B. Reduced number of [3H]nicotine and [3H]acetylcholine binding sites in the frontal cortex of Alzheimer brains. Neurosci Lett. 1986 Dec;72(1):115–9.

10. Lasala M, Fabiani C, Corradi J, Antollini S, Bouzat C. Molecular Modulation of Human α7 Nicotinic Receptor by Amyloid-β Peptides. Front Cell Neurosci [Internet]. 2019;13. Available from: https://www.frontiersin.org/article/10.3389/fncel.2019.00037

11. Buckingham SD, Jones AK, Brown LA, Sattelle DB. Nicotinic acetylcholine receptor signalling: roles in Alzheimer's disease and amyloid neuroprotection. Pharmacol Rev. 2009 Mar;61(1):39–61.

12. Wu J, Kuo Y-P, George AA, Xu L, Hu J, Lukas RJ. beta-Amyloid directly inhibits human alpha4beta2-nicotinic acetylcholine receptors heterologously expressed in human SH-EP1 cells. J Biol Chem. 2004 Sep;279(36):37842–51.

13. Lamb PW, Melton MA, Yakel JL. Inhibition of neuronal nicotinic acetylcholine receptor channels expressed in Xenopus oocytes by beta-amyloid1-42 peptide. J Mol Neurosci. 2005;27(1):13–21.

14. Kihara T, Shimohama S, Sawada H, Kimura J, Kume T, Kochiyama H, et al. Nicotinic receptor stimulation protects neurons against beta-amyloid toxicity. Ann Neurol. 1997 Aug;42(2):159–63.

15. He N, Wang Z, Wang Y, Shen H, Yin M. ZY-1, A Novel Nicotinic Analog, Promotes

Proliferation and Migration of Adult Hippocampal Neural Stem/Progenitor Cells. Cell Mol
Neurobiol [Internet]. 2013;33(8):1149–57. Available from:
https://doi.org/10.1007/s10571-013-9981-0

16. Nie H, Wang Z, Zhao W, Lu J, Zhang C, Lok K, et al. New nicotinic analogue ZY-1
enhances cognitive functions in a transgenic mice model of Alzheimer's disease. Neurosci
Lett [Internet]. 2013;537:29–34. Available from:
https://www.sciencedirect.com/science/article/pii/S0304394013000153

17. Deardorff WJ, Feen E, Grossberg GT. The Use of Cholinesterase Inhibitors Across All
Stages of Alzheimer's Disease. Drugs Aging [Internet]. 2015;32(7):537–47. Available
from: https://doi.org/10.1007/s40266-015-0273-x

18. Albuquerque EX, Pereira EFR, Alkondon M, Rogers SW. Mammalian Nicotinic
Acetylcholine Receptors: From Structure to Function. Physiol Rev [Internet].
2009;89(1):73–120. Available from: https://doi.org/10.1152/physrev.00015.2008

19. Venkatesan S, Jeoung H-S, Chen T, Power SK, Liu Y, Lambe EK. Endogenous
Acetylcholine and Its Modulation of Cortical Microcircuits to Enhance Cognition. Curr
Top Behav Neurosci. 2020;45:47–69.

20. Power SK, Venkatesan S, Lambe EK. Xanomeline restores endogenous nicotinic
acetylcholine receptor signaling in mouse prefrontal cortex. Neuropsychopharmacol Off
Publ Am Coll Neuropsychopharmacol. 2023 Mar;48(4):671–82.

21. Venkatesan S, Lambe EK. Chrna5 is essential for a rapid and protected response to
optogenetic release of endogenous acetylcholine in prefrontal cortex. J Neurosci.
2020;40(38).

22. Venkatesan S, Chen T, Liu Y, Turner EE, Tripathy SJ, Lambe EK. Chrna5 and lynx
prototoxins identify acetylcholine super-responder subplate neurons. iScience. 2023
Feb;26(2):105992.

23. Bailey CDC, De Biasi M, Fletcher PJ, Lambe EK. The Nicotinic Acetylcholine Receptor
α5 Subunit Plays a Key Role in Attention Circuitry and Accuracy. J Neurosci [Internet].
2010 Jul 7;30(27):9241 LP – 9252. Available from:
http://www.jneurosci.org/content/30/27/9241.abstract

24. Tian MK, Bailey CDC, De Biasi M, Picciotto MR, Lambe EK. Plasticity of Prefrontal
Attention Circuitry: Upregulated Muscarinic Excitability in Response to Decreased
Nicotinic Signaling Following Deletion of α5 or β2 Subunits. J Neurosci [Internet]. 2011
Nov 9;31(45):16458 LP – 16463. Available from:
http://www.jneurosci.org/content/31/45/16458.abstract

25. Ramirez-Latorre J, Yu CR, Qu X, Perin F, Karlin A, Role L. Functional contributions of
α5 subunit to neuronal acetylcholine receptor channels. Nature [Internet].
1996;380(6572):347–51. Available from: https://doi.org/10.1038/380347a0

26. Sciaccaluga M, Monconi C, Martinello K, Catalano M, Bermudez I, Stitzel JA, et al.
Crucial role of nicotinic α5 subunit variants for Ca2+ fluxes in ventral midbrain neurons.
FASEB J [Internet]. 2015;29(8):3389–98. Available from:
https://faseb.onlinelibrary.wiley.com/doi/abs/10.1096/fj.14-268102

27. Scholze P, Huck S. The α5 Nicotinic Acetylcholine Receptor Subunit Differentially
Modulates α4β2* and α3β4* Receptors. Front Synaptic Neurosci [Internet]. 2020;12.
Available from: https://www.frontiersin.org/article/10.3389/fnsyn.2020.607959

28. Howe WM, Brooks JL, Tierney PL, Pang J, Rossi A, Young D, et al. α5 nAChR
modulation of the prefrontal cortex makes attention resilient. Brain Struct Funct. 2018
Mar;223(2):1035–47.

29. Schuch JB, Polina ER, Rovaris DL, Kappel DB, Mota NR, Cupertino RB, et al. Pleiotropic

695   effects of Chr15q25 nicotinic gene cluster and the relationship between smoking,
696   cognition and ADHD. J Psychiatr Res. 2016 Sep;80:73–8.

697 30.   Han W, Zhang T, Ni T, Zhu L, Liu D, Chen G, et al. Relationship of common variants in
698   CHRNA5 with early-onset schizophrenia and executive function. Schizophr Res. 2019
699   Apr;206:407–12.

700 31.   Jensen KP, DeVito EE, Herman AI, Valentine GW, Gelernter J, Sofuoglu M. A CHRNA5
701   Smoking Risk Variant Decreases the Aversive Effects of Nicotine in Humans.
702   Neuropsychopharmacology [Internet]. 2015/05/07. 2015 Nov;40(12):2813–21. Available
703   from: https://pubmed.ncbi.nlm.nih.gov/25948103

704 32.   Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia
705   prevention, intervention, and care: 2020 report of the Lancet Commission.
706   Lancet [Internet]. 2020 Aug 8;396(10248):413–46. Available from:
707   https://doi.org/10.1016/S0140-6736(20)30367-6

708 33.   Bennett DA, Buchman AS, Boyle PA, Barnes LL, Wilson RS, Schneider JA. Religious
709   Orders Study and Rush Memory and Aging Project. J Alzheimers Dis. 2018;64(s1):S161–
710   89.

711 34.   Bakken TE, Jorstad NL, Hu Q, Lake BB, Tian W, Kalmbach BE, et al. Comparative
712   cellular analysis of motor cortex in human, marmoset and mouse. Nature [Internet].
713   2021;598(7879):111–9. Available from: https://doi.org/10.1038/s41586-021-03465-8

714 35.   Cain A, Taga M, McCabe C, Hekselman I, White CC, Green G, et al. Multi-cellular
715   communities are perturbed in the aging human brain and with alzheimer's disease. bioRxiv
716   [Internet]. 2020; Available from:
717   https://www.biorxiv.org/content/early/2020/12/23/2020.12.22.424084

718 36.   Smith RM, Alachkar H, Papp AC, Wang D, Mash DC, Wang J-C, et al. Nicotinic α5
719   receptor subunit mRNA expression is associated with distant 5′ upstream polymorphisms.
720   Eur J Hum Genet [Internet]. 2011;19(1):76–83. Available from:
721   https://doi.org/10.1038/ejhg.2010.120

722 37.   Ji X, Gui J, Han Y, Brennan P, Li Y, McKay J, et al. The role of haplotype in 15q25.1
723   locus in lung cancer risk: results of scanning chromosome 15. Carcinogenesis. 2015
724   Nov;36(11):1275–83.

725 38.   Wessel J, McDonald SM, Hinds DA, Stokowski RP, Javitz HS, Kennemer M, et al.
726   Resequencing of nicotinic acetylcholine receptor genes and association of common and
727   rare variants with the Fagerström test for nicotine dependence. Neuropsychopharmacology
728   [Internet]. 2010/08/25. 2010 Nov;35(12):2392–402. Available from:
729   https://pubmed.ncbi.nlm.nih.gov/20736995

730 39.   Swan GE, Javitz HS, Jack LM, Wessel J, Michel M, Hinds DA, et al. Varenicline for
731   smoking cessation: nausea severity and variation in nicotinic receptor genes.
732   Pharmacogenomics J [Internet]. 2012;12(4):349–58. Available from:
733   https://doi.org/10.1038/tpj.2011.19

734 40.   Sadaghiani S, Ng B, Altmann A, Poline J-B, Banaschewski T, Bokde ALW, et al.
735   Overdominant Effect of a CHRNA4 Polymorphism on Cingulo-Opercular Network
736   Activity and Cognitive Control. J Neurosci [Internet]. 2017;37(40):9657–66. Available
737   from: https://www.jneurosci.org/content/37/40/9657

738 41.   De Jager PL, Ma Y, McCabe C, Xu J, Vardarajan BN, Felsky D, et al. A multi-omic atlas
739   of the human frontal cortex for aging and Alzheimer's disease research. Sci Data
740   [Internet]. 2018;5(1):180142. Available from: https://doi.org/10.1038/sdata.2018.142

741 42.   Bennett DA, Schneider JA, Aggarwal NT, Arvanitakis Z, Shah RC, Kelly JF, et al.
742   Decision Rules Guiding the Clinical Diagnosis of Alzheimer's Disease in Two

Community-Based Cohort Studies Compared to Standard Practice in a Clinic-Based Cohort Study. Neuroepidemiology [Internet]. 2006;27(3):169–76. Available from: https://www.karger.com/DOI/10.1159/000096129

43. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell gene expression data. Nat Biotechnol [Internet]. 2015;33(5):495–502. Available from: https://doi.org/10.1038/nbt.3192

44. Hodge RD, Bakken TE, Miller JA, Smith KA, Barkan ER, Graybuck LT, et al. Conserved cell types with divergent features in human versus mouse cortex. Nature. 2019 Sep;573(7772):61–8.

45. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. J R Stat Soc Ser B [Internet]. 1995 Jan 26;57(1):289–300. Available from: http://www.jstor.org/stable/2346101

46. Kuryatov A, Berrettini W, Lindstrom J. Acetylcholine receptor (AChR) α5 subunit variant associated with risk for nicotine dependence and lung cancer reduces (α4β2)□α5 AChR function. Mol Pharmacol. 2011 Jan;79(1):119–25.

47. Wang J-C, Spiegel N, Bertelsen S, Le N, McKenna N, Budde JP, et al. Cis-Regulatory Variants Affect CHRNA5 mRNA Expression in Populations of African and European Ancestry. PLoS One [Internet]. 2013 Nov 26;8(11):e80204. Available from: https://doi.org/10.1371/journal.pone.0080204

48. Hoft NR, Stitzel JA, Hutchison KE, Ehringer MA. CHRNB2 promoter region: association with subjective effects to nicotine and gene expression differences. Genes Brain Behav [Internet]. 2010/11/04. 2011 Mar;10(2):176–85. Available from: https://pubmed.ncbi.nlm.nih.gov/20854418

49. Jensen KP, Devito EE, Herman AI, Valentine GW, Gelernter J, Sofuoglu M. A CHRNA5 smoking risk variant decreases the aversive effects of nicotine in humans. Neuropsychopharmacology. 2015;40(12):2813–21.

50. Andersen OM, Reiche J, Schmidt V, Gotthardt M, Spoelgen R, Behlke J, et al. Neuronal sorting protein-related receptor sorLA/LR11 regulates processing of the amyloid precursor protein. Proc Natl Acad Sci [Internet]. 2005;102(38):13461–6. Available from: https://www.pnas.org/doi/abs/10.1073/pnas.0503689102

51. Caglayan S, Takagi-Niidome S, Liao F, Carlo A-S, Schmidt V, Burgert T, et al. Lysosomal Sorting of Amyloid-β by the SORLA Receptor Is Impaired by a Familial Alzheimer's Disease Mutation. Sci Transl Med [Internet]. 2014;6(223):223ra20-223ra20. Available from: https://www.science.org/doi/abs/10.1126/scitranslmed.3007747

52. Hung C, Tuck E, Stubbs V, van der Lee SJ, Aalfs C, van Spaendonk R, et al. SORL1 deficiency in human excitatory neurons causes APP-dependent defects in the endolysosome-autophagy network. Cell Rep [Internet]. 2021;35(11):109259. Available from: https://www.sciencedirect.com/science/article/pii/S2211124721006239

53. Tamura K, Chiu Y-W, Shiohara A, Hori Y, Tomita T. EphA4 regulates Aβ production via BACE1 expression in neurons. FASEB J Off Publ Fed Am Soc Exp Biol. 2020 Dec;34(12):16383–96.

54. Griffiths HH, Whitehouse IJ, Hooper NM. Regulation of amyloid-β production by the prion protein. Prion [Internet]. 2012/07/01. 2012 Jul 1;6(3):217–22. Available from: https://pubmed.ncbi.nlm.nih.gov/22449984

55. Nie H, Li Z, Lukas RJ, Shen Y, Song L, Wang X, et al. Construction of SH-EP1-alpha4beta2-hAPP695 cell line and effects of nicotinic agonists on beta-amyloid in the cells. Cell Mol Neurobiol. 2008 Jan;28(1):103–12.

56. Beraldo FH, Arantes CP, Santos TG, Queiroz NGT, Young K, Rylett RJ, et al. Role of

alpha7 nicotinic acetylcholine receptor in calcium signaling induced by prion protein
interaction with stress-inducible protein 1. J Biol Chem. 2010 Nov;285(47):36542–50.

57. Nygaard HB, Strittmatter SM. Cellular Prion Protein Mediates the Toxicity of β-Amyloid
Oligomers: Implications for Alzheimer Disease. Arch Neurol [Internet].
2009;66(11):1325–8. Available from: https://doi.org/10.1001/archneurol.2009.223

58. Lipina T V, Prasad T, Yokomaku D, Luo L, Connor SA, Kawabe H, et al. Cognitive
Deficits in Calsyntenin-2-deficient Mice Associated with Reduced GABAergic
Transmission. Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol. 2016
Feb;41(3):802–10.

59. Maskos U. The nicotinic receptor alpha5 coding polymorphism rs16969968 as a major
target in disease: Functional dissection and remaining challenges. J Neurochem [Internet].
2020;154(3):241–50. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/jnc.14989

60. Hijazi S, Heistek TS, Scheltens P, Neumann U, Shimshek DR, Mansvelder HD, et al.
Early restoration of parvalbumin interneuron activity prevents memory loss and network
hyperexcitability in a mouse model of Alzheimer's disease. Mol Psychiatry [Internet].
2020;25(12):3380–98. Available from: https://doi.org/10.1038/s41380-019-0483-4

61. Wright AL, Zinn R, Hohensinn B, Konen LM, Beynon SB, Tan RP, et al.
Neuroinflammation and neuronal loss precede Aβ plaque deposition in the hAPP-J20
mouse model of Alzheimer's disease. PLoS One. 2013;8(4):e59586.

62. Zheng J, Li H-L, Tian N, Liu F, Wang L, Yin Y, et al. Interneuron Accumulation of
Phosphorylated tau Impairs Adult Hippocampal Neurogenesis by Suppressing GABAergic
Transmission. Cell Stem Cell [Internet]. 2020 Mar 5;26(3):331-345.e6. Available from:
https://doi.org/10.1016/j.stem.2019.12.015

63. Beal MF, Mazurek MF, Svendsen CN, Bird ED, Martin JB. Widespread reduction of
somatostatin-like immunoreactivity in the cerebral cortex in Alzheimer's disease. Ann
Neurol. 1986 Oct;20(4):489–95.

64. Waller R, Mandeya M, Viney E, Simpson JE, Wharton SB. Histological characterization
of interneurons in Alzheimer's disease reveals a loss of somatostatin interneurons in the
temporal cortex. Neuropathology. 2020 Aug;40(4):336–46.

65. Fonseca M, Soriano E, Ferrer I, Martinez A, Tuñon T. Chandelier cell axons identified by
parvalbumin-immunoreactivity in the normal human temporal cortex and in Alzheimer's
disease. Neuroscience [Internet]. 1993;55(4):1107–16. Available from:
https://www.sciencedirect.com/science/article/pii/0306452293903249

66. Koukouli F, Rooy M, Tziotis D, Sailor KA, O'Neill HC, Levenga J, et al. Nicotine
reverses hypofrontality in animal models of addiction and schizophrenia. Nat Med
[Internet]. 2017;23(3):347–54. Available from: https://doi.org/10.1038/nm.4274

67. Forget B, Scholze P, Langa F, Morel C, Pons S, Mondoloni S, et al. A Human
Polymorphism in CHRNA5 Is Linked to Relapse to Nicotine Seeking in Transgenic Rats.
Curr Biol [Internet]. 2018;28(20):3244-3253.e7. Available from:
https://www.sciencedirect.com/science/article/pii/S096098221831128X

68. Fairén A, Valverde F. A specialized type of neuron in the visual cortex of cat: a Golgi and
electron microscope study of chandelier cells. J Comp Neurol. 1980 Dec;194(4):761–79.

69. Somogyi P. A specific "axo-axonal" interneuron in the visual cortex of the rat. Brain Res.
1977 Nov;136(2):345–50.

70. Schneider-Mizell CM, Bodor AL, Collman F, Brittain D, Bleckert A, Dorkenwald S, et al.
Structure and function of axo-axonic inhibition. Calabrese RL, Callaway E, Huang ZJ,
Oberlaender M, editors. Elife [Internet]. 2021;10:e73783. Available from:

839   https://doi.org/10.7554/eLife.73783

840 71. Ribak CE. Axon terminals of GABAergic chandelier cells are lost at epileptic foci. Brain
841   Res [Internet]. 1985;326(2):251–60. Available from:
842   https://www.sciencedirect.com/science/article/pii/0006899385900344

843 72. Zhu Y, Stornetta RL, Zhu JJ. Chandelier Cells Control Excessive Cortical Excitation:
844   Characteristics of Whisker-Evoked Synaptic Responses of Layer 2/3 Nonpyramidal and
845   Pyramidal Neurons. J Neurosci [Internet]. 2004 Jun 2;24(22):5101 LP – 5108. Available
846   from: http://www.jneurosci.org/content/24/22/5101.abstract

847 73. León-Espinosa G, DeFelipe J, Muñoz A. Effects of amyloid-β plaque proximity on the
848   axon initial segment of pyramidal cells. J Alzheimers Dis. 2012;29(4):841–52.

849 74. Blazquez-Llorca L, Garcia-Marin V, Defelipe J. Pericellular innervation of neurons
850   expressing abnormally hyperphosphorylated tau in the hippocampal formation of
851   Alzheimer's disease patients. Front Neuroanat. 2010;4:20.

852 75. Ueda M, Iida Y, Kitamura Y, Kawashima H, Ogawa M, Magata Y, et al. 5-Iodo-A-85380,
853   a specific ligand for alpha 4 beta 2 nicotinic acetylcholine receptors, prevents glutamate
854   neurotoxicity in rat cortical cultured neurons. Brain Res. 2008 Mar;1199:46–52.

855 76. Cingir Koker S, Jahja E, Shehwana H, Keskus AG, Konu O. Cholinergic Receptor
856   Nicotinic Alpha 5 (CHRNA5) RNAi is associated with cell cycle inhibition, apoptosis,
857   DNA damage response and drug sensitity in breast cancer. PLoS One.
858   2018;13(12):e0208982.

859 77. Barrera-Ocampo A, Arlt S, Matschke J, Hartmann U, Puig B, Ferrer I, et al. Amyloid-β
860   Precursor Protein Modulates the Sorting of Testican-1 and Contributes to Its Accumulation
861   in Brain Tissue and Cerebrospinal Fluid from Patients with Alzheimer Disease. J
862   Neuropathol Exp Neurol [Internet]. 2016 Sep 1;75(9):903–16. Available from:
863   https://doi.org/10.1093/jnen/nlw065

864 78. Barch DM, Carter CS, Braver TS, Sabb FW, MacDonald A 3rd, Noll DC, et al. Selective
865   deficits in prefrontal cortex function in medication-naive patients with schizophrenia.
866   Arch Gen Psychiatry. 2001 Mar;58(3):280–8.

867 79. Cornelius ME, Loretan CG, Wang TW, Jamal A, Moma DM. Tobacco Product Use
868   Among Adults — United States, 2020. MMWR Morb Mortal Wkly Rep. 2022;71:397–
869   405.

870 80. Bierut LJ, Stitzel JA, Wang JC, Hinrichs AL, Grucza RA, Xuei X, et al. Variants in
871   nicotinic receptors and risk for nicotine dependence. Am J Psychiatry [Internet].
872   2008/06/02. 2008 Sep;165(9):1163–71. Available from:
873   https://pubmed.ncbi.nlm.nih.gov/18519524

874 81. Johnson ECB, Carter EK, Dammer EB, Duong DM, Gerasimov ES, Liu Y, et al. Large-
875   scale deep multi-layer analysis of Alzheimer's disease brain reveals strong proteomic
876   disease-related changes not observed at the RNA level. Nat Neurosci [Internet].
877   2022;25(2):213–25. Available from: https://doi.org/10.1038/s41593-021-00999-y

878 82. Hansen JY, Markello RD, Tuominen L, Nørgaard M, Kuzmin E, Palomero-Gallagher N, et
879   al. Correspondence between gene expression and neurotransmitter receptor and transporter
880   density in the human brain. Neuroimage [Internet]. 2022;264:119671. Available from:
881   https://www.sciencedirect.com/science/article/pii/S1053811922007923

882 83. Petrache AL, Rajulawalla A, Shi A, Wetzel A, Saito T, Saido TC, et al. Aberrant
883   Excitatory-Inhibitory Synaptic Mechanisms in Entorhinal Cortex Microcircuits During the
884   Pathogenesis of Alzheimer's Disease. Cereb Cortex [Internet]. 2019 Apr 1;29(4):1834–50.
885   Available from: https://pubmed.ncbi.nlm.nih.gov/30766992

886 84. Wada E, McKinnon D, Heinemann S, Patrick J, Swanson LW. The distribution of mRNA

encoded by a new member of the neuronal nicotinic acetylcholine receptor gene family
(alpha 5) in the rat central nervous system. Brain Res. 1990 Aug;526(1):45–53.

85.   Winzer-Serhan UH, Leslie FM. Expression of alpha5 nicotinic acetylcholine receptor
subunit mRNA during hippocampal and cortical development. J Comp Neurol. 2005
Jan;481(1):19–30.

86.   Tai Y, Janas JA, Wang C-L, Van Aelst L. Regulation of chandelier cell cartridge and
bouton development via DOCK7-mediated ErbB4 activation. Cell Rep. 2014
Jan;6(2):254–63.

895
896

Figure legends
Figure 1. **SNPs affecting expression of α4β2* nicotinic receptor subunit genes highlight a**
link between *CHRNA5* and amyloid pathology. *A,* Schematics illustrating different subunit
compositions of prefrontal α4β2* nicotinic receptors, with and without the α5 subunit. *B,*
Localization of the rs16969968 and rs1979905 SNPs in relation to the *CHRNA5* locus. *C,* The A
allele of the missense SNP rs16969968 (left) in the coding region of *CHRNA5* is associated with
lowered *CHRNA5* expression, while the A allele of the rs1979905 SNP (middle) upstream of the
CHRNA5 gene is associated with enhanced *CHRNA5* expression. Data from the DLPFC of
ROS/MAP individuals. *D, CHRNA5* expression appears to be controlled by the zygosity of the
rs1979905 A allele (colors) instead of the rs16969968 A allele (x-axis). Data shown as *CHRNA5*
expression for each subject with means indicated. *E,* eQTL effects of SNPs in nicotinic subunit
genes on their respective gene expression. Shown as β-coefficient with significance (p) in
brackets. **F,** Network plot depicting the relationships between SNPs (black), gene expression
(red), neuropathology (blue), and last global cognition score (green). Solid and dashed lines
indicate whether association was significant after correction for FDR or not, respectively. **G,**
Network plot depicting the correlations present between the expression of select cholinergic
genes in the DLPFC. Colour of lines indicates direction of correlation (negative or positive)
while the thickness indicates correlation strength. All correlations shown are significant after
adjustment for FDR.
916
Figure 2. *CHRNA5* **expression is elevated in chandelier cells and is affected by genotype for**
the rs1979905 A allele. *A,* Expression of *CHRNA5* averaged per cell type per individual, original
gene count values were normalized for each cell by total expression. F-test significance of
ANOVA shown on graph, with red asterisk denoting post-hoc tests demonstrating *CHRNA5*
expression is stronger in chandelier cells compared to all but one other cell type. Mean
expression of *CHRNA5* in chandelier cells displayed (red line). *B,* Expression of *CHRNA4* (left)
and *CHRNB2* (right) across different cell types in the ROS/MAP DLPFC snRNAseq dataset.
Mean expression of *CHRNA4* or *CHRNB2* in chandelier cells displayed (red line)*.* Data shown as
mean + SEM of the data averaged per cell-type per individual. *C,* Expression of *CHRNA5* across
cell types in the PFC in individuals split by genotype for the rs1979905 A allele, expression
averaged per cell type per individual. Number of individuals per rs1979905 A allele genotype: 0,
n = 5; 1, n = 13; 2, n = 4. *D,* Effect of genotype for rs1979905 A allele on expression of *CHRNA5*
in selected cell types, data shown as *CHRNA5* expression averaged per cell type per individual. A
pattern of increasing *CHRNA5* expression with increasing rs1979905 A zygosity is present in
some cell types. Significance shown for linear regression models for L5 and L6 excitatory
neurons. Displayed as mean + SEM. Number of cells per subtype indicated (n_{cell}).

Figure 3. **Chandelier cells are significantly enriched for genes interacting with amyloid.** *A,*
Ontology (molecular function) of gene set upregulated in cortical PVALB+ chandelier cells
versus PVALB+ non-chandelier cells (basket cells). Only molecular functions with significant
fold enrichment after FDR are displayed. "Calcium ion-binding" and "amyloid-beta binding"
functions are highlighted. **B,** Venn diagram displaying genes from *A* with either a "calcium ion-
binding" or a "amyloid-beta binding" molecular function, and their overlap.

Figure 4. **Association of chandelier cell proportions with β-amyloid load is dependent on the**
rs16969968 A allele genotype. Cell type proportion data for interneuron populations is available
for almost a third of the deceased ROS/MAP subjects, allowing the assessment of the interaction
among chandelier cell proportion, brain amyloid load, and the *CHRNA5* SNP haplotype. *A,B*

Stratifying by rs16969968 A allele reveals a significant interaction effect between rs16969968 A
allele and amyloid load on chandelier cell proportions with rs16969968 A allele homozygotes
showing a more negative association between brain amyloid load and chandelier cell proportions
compared to rs16969968 A allele non-carriers (interaction term t = -2.842, $p = 0.005$). Scatter
plots show 95% confidence intervals of linear model predictions, β-coefficients and p values of
individual linear regression models are displayed. **C,** Overlay of the linear models from A,B,
showing 95% confidence intervals. ***D,E,*** Stratifying by rs1979905 A allele shows a suggestion of
an opposite interaction between rs1979905 A allele and amyloid load on chandelier cell
proportions (interaction term t = 1.81, $p = 0.071$). Scatter plots show 95% confidence intervals of
linear model predictions, β-coefficients and p values of individual linear regression models are
displayed. ***F***, Overlay of the linear models from D,E, showing 95% confidence intervals.
954
<u>**Figure 5.**</u> **A working model of the potential role of chandelier cells in β-amyloid processing,**
and of the impact of *CHRNA5* genotype on chandelier cell (ChC) resilience and
vulnerability. Top: Chandelier cells are significantly enriched for multiple genes involved in β-
amyloid processing and degradation including for example neprilysin (NEP), a potentially
CHRNA5-regulated degrader of β-amyloid, and SORL1, a vital component of the APP-recycling
pathway. Bottom: In coding-SNP rs16969968 non-carriers, the α4β2α5 nicotinic receptor is
resistant to inhibition by β-amyloid, preserving nicotinic signalling even at high β-amyloid levels
that inhibit the α4β2 receptors. In coding-SNP rs16969968 homozygous individuals, the
disruption of the α5 subunit may reduce its representation in the receptors or block its protective
function against β-amyloid, leading to disrupted nicotinic signalling at higher β-amyloid levels,
possibly triggering a cytotoxic response in the chandelier cells. Chandelier cell structure is based
on images from Tai et al. 2014 (86).
967

Table 1. Expression correlation of selected cholinergic genes. Results of Pearson's correlation analysis of the expression of selected cholinergic genes using the bulk tissue RNAseq data from the DLPFC of ROS/MAP subjects.

Gene 1	Gene 2	R	p	FDR
CHRNA4	ACHE	0.556	<0.001	<0.001
	CHRM1	0.542	<0.001	<0.001
	CHRM3	0.242	<0.001	0.002
	CHRNA2	0.307	<0.001	<0.001
	CHRNA7	0.131	0.039	0.464
	CHRNB2	0.457	<0.001	<0.001
CHRNA5	ACHE	0.063	0.324	1.000
	CHRM1	-0.309	<0.001	<0.001
	CHRM3	-0.095	0.134	1.000
	CHRNA2	-0.045	0.477	1.000
	CHRNA4	-0.130	0.041	0.464
	CHRNA7	-0.061	0.342	1.000
	CHRNB2	-0.248	<0.001	0.001
CHRNB2	ACHE	0.472	<0.001	<0.001
	CHRM1	0.846	<0.001	<0.001
	CHRM3	0.645	<0.001	<0.001
	CHRNA2	0.283	<0.001	<0.001
	CHRNA4	0.457	<0.001	<0.001
	CHRNA7	0.359	<0.001	<0.001

A

(α4)2(α5)(β2)2   (α4)2(β2)3   (α4)3(β2)2

B

■ - UTR ■ - Exon

CHRNA5

rs1979905   rs16969968

C

CHRNA5 expression vs rs16969968_A ****

CHRNA5 expression vs rs1979905_A ****

D

CHRNA5 Expression vs rs16969968_A **** **** **

rs1979905_A (0)
rs1979905_A (1)
rs1979905_A (2)

E

SNP/Gene	*CHRNA5*	*CHRNA4*	*CHRNB2*
rs1979905_A	**1.08 (5.36*10⁻¹²¹)**	-0.011 (0.604)	-0.010 (0.639)
rs16969968_A	**-0.703 (1.63*10⁻³⁸)**	0.012 (0.581)	0.0002 (0.994)
rs1044396_G	-0.074 (0.164)	**0.071 (0.0007)**	0.023 (0.293)
rs45497800_T	0.039 (0.7)	**-0.181 (6.48*10⁻⁶)**	-0.029 (0.497)
rs4292956_T	0.061 (0.534)	0.025 (0.514)	**-0.209 (8.4*10⁻⁷)**
rs2072660_T	0.09 (0.148)	-0.023 (0.347)	**-0.120 (2.5*10⁻⁶)**

F

Association significant after FDR?
—— TRUE
---- FALSE

Association direction
—— Positive
—— Negative

G

Correlation strength
0.4
0.6
0.8

Correlation direction
—— Positive
—— Negative