Single cell transcriptomic profiling of tauopathy in a novel 3D neuron-astrocyte coculture model

Hannah Drew Rickner[1#], Lulu Jiang[2#], Rui Hong[3], Nicholas K O'Neill[3], Maria Medalla[4,6], Lushuang Zhang[2], Dipan Shaw[7], Benjamin Wolozin[2,5,6]*, Christine S. Cheng[1,3,7,8]*

Author Affiliations

1. Department of Biology, Boston University, Boston, MA, 02215

   Hannah Drew Rickner and Christine S. Cheng

2. Department of Pharmacology and Experimental Therapeutics, Boston University School of Medicine, Boston, MA, 02118

   Lulu Jiang, Lushuang Zhang, and Benjamin Wolozin

3. Program in Bioinformatics, Boston University, Boston, MA, 02215

   Rui Hong and Nicholas K O'Neill and Christine S. Cheng

4. Department of Anatomy & Neurobiology, Boston University School of Medicine, Boston, MA, 02118, USA.

   Maria Medalla

5. Department of Neurology, Boston University School of Medicine

   Benjamin Wolozin

6. Center for Systems Neuroscience, Boston University

   Maria Medalla and Benjamin Wolozin

7. Informatics Group, J. Craig Venter Institute, La Jolla, CA, 92037

   Dipan Shaw and Christine S. Cheng

8. Department of Psychiatry, University of California San Diego, La Jolla, CA, 92093

   Christine S. Cheng

# These authors contributed equally.

*Corresponding authors: bwolozin@bu.edu or csc003@ucsd.edu

ABSTRACT

The use of iPSC derived brain organoid models to study neurodegenerative disease has been hampered by a lack of systems that accurately and expeditiously recapitulate pathogenesis in the context of neuron-glial interactions. Here we report development of a system, termed AstTau, which propagates toxic human tau oligomers in iPSC derived neuron-astrocyte spheroids. The AstTau system develops much of the neuronal and astrocytic pathology observed in tauopathies including misfolded, phosphorylated, oligomeric, and fibrillar tau, strong neurodegeneration, and reactive astrogliosis. Single cell transcriptomic profiling combined with immunochemistry characterizes a model system that can more closely recapitulate late-stage changes in adult neurodegeneration. The transcriptomic studies demonstrate striking changes in neuroinflammatory and heat shock protein (HSP) chaperone systems in the disease process. Treatment with the HSP90 inhibitor PU-H71 was used to address the putative dysfunctional HSP epichaperome and produced a strong reduction of pathology and neurodegeneration, highlighting the potential of AstTau as a rapid and reproducible tool for drug discovery.

KEYWORDS

Pluripotent stem cells; 3D culture; disease modeling; neurodegenerative diseases; tauopathy; phospho-tau, neurons; astrocytes; reactive astrocytosis, cell death, heat shock protein chaperone; single cell transcriptomics

INTRODUCTION

Tauopathies are characterized by the accumulation of misfolded and hyperphosphorylated tau which form neurofibrillary tangles and precipitate neurodegeneration. These intracellular factors such as aggregation of tau and impaired proteostasis contribute to the tauopathy degenerative process, typically in combination with inflammation and vascular dysfunction [1-7]. The resulting neuronal damage and tau pathology is closely associated with cognitive decline in tauopathies [8,9]. This suggest that broadening our understanding of tau pathogenesis will enable the development of effective therapeutics against tauopathies.

Many models for tauopathies have been created, but few if any accurately reflect the human milieu. Animal models develop tangles but lack a human background, often producing results that don't translate successfully to human disease modifying strategies [10,11]. Increasingly, investigators are turning to human induced pluripotent stem cells (hiPSCs) to overcome such challenges, but such 2D models typically exhibit limited arborization, neuron-neuron, neuron-glial interactions, and the extensive tau pathology and degeneration present in tauopathies [12-25]. Three-dimensional (3D) organoid models of tauopathies show great promise, but most require maturation over an extended time frame of months or years, lack an explicit ratio of neuron and glial compartments, and develop limited pathology and neurodegeneration [26-35]. This difficulty in producing an expedient human organoid model combined with the inadequacies of current animal models for tauopathies has stymied the development of effective research and therapeutics.

To address this challenge, we present a model in which hiPSC-derived neurons are seeded with human tau oligomers, combined with hiPSC derived astrocytes and grown in spheroidal culture to create a 3D model termed 'asteroids' [36,37]. The resulting neurons and astrocyte disease model rapidly recapitulates physiologically relevant cellular interactions and exhibits advanced maturation over other cerebral organoid or spheroid models [36-38]. Increasing evidence indicates that tau propagation is an important contributor to multiple human neurodegenerative diseases [39-42]. Accumulating studies demonstrate that exogenous tau oligomers can be internalized, trafficked, and propagated in animal and cell models, spreading disease pathology in a prion-like manner [43-53]. We show that challenging the asteroid model with toxic human oligomeric tau (oTau) yields robust tau pathology, neurodegeneration and reactive astrogliosis over just 21 days of 3D culture (DIV3D). We will refer to this system as AstTau. A key to the success of this model lies in seeding

with a small amount of tau oligomers, and subsequent expansion of the pathology through propagation to other neurons. Most studies of tau propagation have focused on tau fibrils, which propagate pathology but elicit little if any neurodegeneration [43-52]. In contrast, we and others have shown that exposure to tau oligomers propagates tau pathology and induces robust pathology with advanced neurodegeneration [54,55,56]. The tauopathy asteroid system AstTau incorporates this propagation of toxic tau oligomers to create a highly reproducible human 3D model of tauopathy in just three weeks. The AstTau system enables analysis of neurodegenerative processes while also enabling concomitant analysis of the disease responses of astrocytes, which are increasingly noted to play an important yet understudied role in the development of tauopathies [37,57-62]. Additionally, the single cell RNA-sequencing (scRNA-seq) presented below provides detailed transcriptomic profiling of cell specific responses to the pathological progression of tauopathy and enables hypothesis testing using the AstTau system.

RESULTS

3D asteroids rapidly recapitulate neuronal and astrocytic signatures

To establish a 3D spheroidal system of neuron and astrocyte coculture for modeling neurodegenerative disease hiPSC derived neural progenitor cells (NPCs) were directed into neuronal and astrocytic lineages (**Fig. 1a**). The neuronal lineage was created by NGN2 overexpression to produce hiPSC derived neuronal cells (hiNCs), while the astrocytic lineage was generated via small molecule-directed differentiation to produce hiPSC derived astrocytes (hiACs) (**Fig. 1a, Suppl. Fig. 1a**). The resulting hiNCs and hiACs were then combined into 3D spheroidal culture termed asteroids as previously described [37] (~2000 cells each) and allowed to develop over the course of three weeks (**Fig. 1a-b, Suppl. Fig. 1b-c**). Pre-differentiation of hiACs and hiNCs allowed for the development of a mature culture system that can rapidly reproduce physiological interactions when neurons and astrocytes are combined in a 3D culture system (**Suppl. Fig. 1b, Suppl. Vid. 1**). 2D mixed cultures containing neuron and astrocytes do not show the extensive arborization and intimate connections observed with 3D culture (**Suppl. Fig. 1a-b, Suppl. Vid. 1**). Incorporation of astrocytes was necessary for asteroid survival, with 3D cultures of neurons alone dying within 7 days (**Suppl. Fig. 1b**). Both neurons and astrocytes exhibited processes that were striking for their length and complexity, as well as for the intimate association (**Suppl. Fig. 1b, Suppl. Vid. 1**).

Asteroid cellular phenotypes were characterized by acquiring scRNA-seq profiles for 26,789 single cells with an average of 2000 genes per cell across a time course of three weeks (**Fig. 1c-f**). Visualization by uniform manifold approximation and projection (UMAP) demonstrated the separation of cells into trajectory guided clusters (**Fig. 1c**). These clusters were identified as the cycling neuronal progenitors (NEU_P), astrocytes precursors (ASC_P), astrocytes (ASC), inhibitory neurons (IN_NEU) and excitatory neurons (EX_NEU) **(Fig. 1c)** based on the expression of established cell type specific genes including TOP2A and NES for progenitors, SLC1A3 (EAAT1), VIM, and PEA15 for astrocytes, and SNAP25, GAP43, and STMN2 for neurons (**Fig. 1c-d, Suppl. Fig. 3**). Neuronal clusters expressed synaptic markers transcripts including DLG4 and SYP and were further separated by inhibitory neuronal markers including GAD1 and SLC32A1 and excitatory neuronal markers SLC38A1 and GRIA4 **(Fig.1c-d, i, Suppl. Fig. 3).** Manual cell typing was further supported by the automated cell typing platform SingleR (**Suppl. Fig. 3e**). Notably, over 21 days of 3D coculture (21 DIV3D) populations of progenitors decreased while the populations of mature cell types showed a concomitant increase (**Fig. 1e-h**). Quantification of progenitor markers SOX2 and VIM and mature neuronal marker MAP2 decreased and increased respectively over the 21 DIV3D time course by both immunolabeling and scRNA-seq gene expression **(Fig. 1g-h).** Furthermore, analysis of immunolabeled cell type markers showed robust differentiation of neurons that were positive for MAP2, NeuN, and Tuj1 as well as astrocytes that were positive for S100β and GFAP, providing support for robust differentiation using protein-based methods (**Suppl. Fig. 1d**). To compare the maturity of neurons and astrocytes in our asteroid culture system to traditional cerebral organoid models we integrated our asteroid scRNA-seq data with an existing published scRNA-seq data set of forebrain cerebral organoids generated using three previously published protocols with different levels of directed differentiation [38]. The asteroid model (red) was overrepresented in more mature cell type clusters compared to the cerebral organoid model (blue) **(Fig 1j, Suppl. Fig. 3f).**

oTau seeding induces rapid tau misfolding, aggregation, and neurodegeneration in the 3D neuron and astrocyte coculture AstTau model

We proceeded to model tauopathy in asteroid cultures (AstTau) **(Fig. 2a).** Human oligomeric tau (oTau) was isolated from 9-month-old PS19 P301S tau mice by centrifugation using a well-validated method, as described previously [55,63,64]. The PS19 P301S tau mice model overexpresses

human 4R1N P301S Tau and post-extraction these human tau fractions were identified by native gel western blot with human-specific tau antibody Tau13 (**Suppl. Fig. 4a**).

The hiNCs were exposed to human oTau (0.04 mg/mL) for 24 hrs after which they were washed and combined with hiACs to generate the self-aggregating AstTau (**Fig.2a**). Analysis of the AstTau cultures over 21 DIV3D showed rapid development of tau pathology, including tau hyperphosphorylation, misfolding, oligomerization and fibrilization (**Fig. 2,3**). Statistically significant increases were observed for misfolded tau (**Fig. 2b-e**). Misfolding and oligomerization of tau were observed using the TOMA2 and MC1 antibodies respectively, with each showing similar patterns of evolution (**Fig. 2b,c, Fig. 2d,e**). Fibrillar tau pathology also evolved, but at a slower rate as seen in clinical tauopathy progression [65]. Fibrillar tau reactive with the dye thioflavin S (ThioS) became evident only at 21 DIV3D (**Fig. 2f,g**). Additionally, hyperphosphorylated tau exhibiting corresponding increases at positions S202/5 and S262 (**Fig. 3a,b, Suppl. Fig. 5**). This increase was validated by AT8 immunoblotting of AstTau at 21 DIV3D **(Fig. 3c).**

The striking propagated neuropathology was also associated with neurodegeneration by 21 DIV3D. Prominent neuronal injury at this late timepoint was evident by Fluoro Jade B labeling (**Fig. 3e,g**), reduced immunolabeling of MAP2 (**Suppl. Fig. 6a,b**) and increased LDH release (**Suppl. Fig. 6c**). Additionally, we observed reciprocal changes in BAX, a cell death regulator. Immunoblotting demonstrated decreases in full length inactive BAX and corresponding increases in activated cleaved BAX with induction of pathology in the AstTau model at 21 DIV3D, indicative of activated cell death pathways **(Fig 3h-j).**

Astrocytes also showed striking pathophysiological responses reminiscent of that observed in the human brain [57,58]. The astrocytic marker S100β and reactive astrogliosis marker GFAP showed statistically significant increases in reactivity beginning at 14 DIV3D and becoming prominent at 21 DIV3D (**Fig. 3a,d,e,f**). Astrocytic processes in the AstTau cultures showed greater area and more intimate localization with regions of neuronal damage than in the control asteroid cultures (**Fig. Suppl. Vid. 1**). Taken together, these results indicate that over 21 DIV3D, the AstTau system develops a range of responses analogous to observed in the brains of human subjects with tau-mediated neurodegeneration [5,6,57,58].

A requirement for oTau in the seeding process was demonstrated by immunodepleting oTau, which prevented any subsequent tau pathology or neurodegeneration (**Suppl. Fig. 4d**). The role of

seeded oTau was further explored by exposing hiNCs to FITC labeled oTau. FITC labeled oTau was present in >75% of hiNCs 24 hrs after seeding, however the vast majority of the seeded tau was degraded by 5 DIV3D (**Suppl. Fig. 4b-d**), and remaining FITC showing no correlation with accumulating tau pathology (**Suppl. Fig. 4d**). These results indicate that tau pathology developing after 5 DIV3D was generated largely from endogenous tau sources. Additionally, propagation of oTau seeded pathology was demonstrated by merging unlabeled (GFP-) spheroids that had been seeded with oTau (at the 2D stage) with GFP+ labeled spheroids that had not been exposed to oTau. In the combined mixed spheroids misfolded and oligomerized tau was found in GFP- and GFP+ neurons, indicating propagation of pathology from oTau seeded neurons to unexposed neurons **(Fig. 3k)**. Additionally, quantification of the accumulation of immunolabeled MC1 showed the same level of misfolded tau pathology developed in pure oTau seeded cultures and mixed cultures compared to control cultures **(Fig. 3l,m)**.

Further control studies demonstrated that the observed pathology in AstTau was selective for oTau by examining the development of pathology in spheroids exposed to preformed α-synuclein fibrils (PFF-αSYN) (**Suppl. Fig 7, 8**). While PFF-αSYN exposure (1 ug/mL) resulted in the accumulation of phosphorylation α-synuclein **(Suppl. Fig. 8d,e)**, it did not elicit oTau specific pathology, such as the accumulation of toxic tau as marked by TOMA2 and MC1 (**Suppl. Fig. 8a-b,d,f**), hyperphosphorylated tau a p202/AT8 (**Suppl. Fig. 8i,j**), fibrillization or reactive astrogliosis (**Suppl. Fig. 8a,c,d,g**), or neurodegeneration (**Suppl. Fig. 8i,k**). Together these data support a direct role for the human toxic tau oligomers in the development of oTau specific pathology in the AstTau system that recapitulates the pathological progression seen in human subjects with tau-mediated neurodegeneration in just 21 DIV3D.

scRNA-seq of AstTau reveals dynamic transcriptional response to oTau induced pathology

The transcriptome of AstTau was characterized by acquiring robust scRNA-seq profiles for 130,605 single cells from four replicate culture batches with an average of 2000 genes per cell from experimental conditions across a time course of three weeks (**Suppl. Fig. 2a,b**). A comparative analysis of control and AstTau scRNA-seq profiles across 21 DIV3D revealed shared and cell type specific transcriptional responses to oTau induced pathology (**Fig. 4**). UMAP projection demonstrates that all scRNA-seq experimental condition datasets are integrated, and highlights changes in the relative proportion of cell types between control and AstTau (**Fig. 4a-b**).

At 21 DIV3D there is an expansion of the astrocytic compartment (ASC) in AstTau that correlates with the observed immunolabeling (**Fig. 4b**). Notably, there is a decrease in the EX_NEU cluster and a correlated increase in the IN_NEU cluster in AstTau consistent with the observation of glutamatergic excitatory neuronal loss in human subjects with tau-mediated neurodegeneration (**Fig. 4b**).

Differential gene expression (DEG) analysis between the control and AstTau conditions within cell type clusters identified striking transcriptional changes over the 21 DIV3D time course (**Fig. 4c, Suppl. Fig. 9**). Notably, concurrent transcriptional changes were evident in both the ASC and NEU groups by 7 DIV3D, suggesting a high degree of interaction between the cell types as only NEU populations were exposed to toxic human tau (**Fig. 4c**). The AstTau system exhibited regulation of ribosomal transcripts, HSP chaperone transcripts, and TNF/MAPK associated intermediate early response (IER) transcripts (termed neuroinflammatory)[66] (**Fig. 4c**). Unexpectedly, an increase in ribosomal transcripts including RPS26 and RPS28 (green) dominated the early response at 7 DIV3D, suggesting a robust translational stress response (**Fig. 4c**). In contrast, there was a suppression of HSP (red) and neuroinflammatory transcripts (purple) at 7 DIV3D (**Fig. 4c**). This trend was reversed with increasing oTau pathology and by 21 DIV3D there was an increase in stress response transcripts including transcripts associated with neuroinflammation (JUN, DUSP1, and RHOB), and HSP chaperones (HSPA1B, HSPH1, DNAJB1, HSPB1, HSPD1, and HSP90AA1) (**Fig. 4c**). Interestingly, EX_NEU demonstrated the most robust upregulation of neuroinflammatory transcripts, while ASC presented with a stronger HSP response (**Fig. 4c**).

These responses were further elucidated by performing statistical analyses on the average expression of all genes in select curated MsigDB [67] or literature derived gene sets per cell to perform a comparative signature analysis (see Methods) (**Fig. 4d,e**). This demonstrated the observed transcriptional changes at 21 DIV3D in AstTau ASC align with transcriptional signatures upregulated in classical reactive astrocyte including A1 and A2 reactive astrocyte signatures [68] as well as Alzheimer's Disease (AD) associated reactive astrocytes specifically identified in human subjects with AD neurodegeneration by single nuclei RNA-seq (snRNA-seq) [61] (**Fig.4d**). Notably, there was a more significant enrichment ($p < 0.0001$) of the A1 toxic astrocyte signature than the A2 protective astrocyte signature ($p < 0.05$) in AstTau (**Fig. 4d**). The

ASC population also show a significant upregulation of the Reactome HSF1 Activation pathway associated with HSP chaperone production **(Fig. 4d).**

Comparative signature analysis in the AstTau neuronal populations demonstrated a concurrent transcriptional upregulation with neuronal signatures found in human subjects with AD neurodegeneration by snRNA-seq [69] **(Fig. 4e).** However, there was divergent regulation of neuroinflammation demonstrated by the HALLMARK TNFα signaling via NFKB signature between IN_NEU and EX_NEU. There was a significant upregulation of neuroinflammation in the EX_NEU population, while the signature was actually suppressed in the IN_NEU population, demonstrating notable consistency with EX_NEU cell population loss at 21 DIV3D **(Fig. 4e)**. Additionally, both EX_NEU and IN_NEU presented a significant downregulation of the GO cytosolic ribosomal pathway at 21 DIV3D potentially indicative of the translational repression that has been associated with tauopathies [70] **(Fig. 4e).**

Functional gene set enrichment analysis further highlights a trajectory of pathological responses among the AstTau cell types. There was a strong ribosomal pathway enrichment at 7 DIV3D that weakened through 14 and 21 DIV3D in agreement with previous analyses **(Fig. 5a-b, Suppl. Fig. 10)**. Pathways associated with neuroinflammation, neuron stress and senescence, and HSP chaperones were enriched among EX_NEU upregulated genes at 21 DIV3D **(Fig. 5a-b, Suppl. Fig. 10)**. While IN_NEU shared the HSP chaperone pathway enrichment, it did not share the neuroinflammatory pathway enrichment, further supporting cell type specific differential responses to pathology progression **(Fig. 5a, Suppl. Fig. 10)**. ASC demonstrated the most robust HSP chaperone pathway enrichment, as well as cell activation, glial development, and stress response pathways enrichment suggestive of reactive astrogliosis **(Fig. 5a-b, Suppl. Fig. 10)**. Together, the functional gene set pathway enrichment of upregulated transcripts across the 21 DIV3D time course in AstTau reveals a progressive stress response including translational, respirasome, and neuroinflammatory profiles that result in the observed neurodegeneration.

 Importantly, functional gene set enrichment analysis also reveals concordant responses between AstTau and single-nuclei transcriptomic profiles of postmortem AD brain, in particular the HSP chaperone response, indicating activation of similar pathophysiological cascades that are not shared by other neuropathologies [61,69,71] **(Fig. 5c)**. These signatures are not shared by comparative controls including single-nuclei transcriptomic profiles of autism spectrum disorder (ASD) brain

tissue or scRNA-seq of PFF-αSYN exposed asteroid cultures (AstαSYN) (**Fig. 5c**), which highlights the specificity of oTau in the development of the pathological and transcriptional evolution observed in AstTau.

Amelioration of AstTau pathology with the epichaperome HSP90 inhibitor PU-H71

The scRNA-seq data profile suggested that the HSP chaperone response plays a pivotal role in neuronal loss and astrocyte activation in our AstTau model. We applied epichaperome inhibitor PU-H71 in our system in order to validate the involvement of HSP epichaperome dysfunction in tau pathogenesis and explore the potential protective effect of correction. Chaperones are proteins that control neuronal function by regulating protein folding, thereby determining the functional levels and activities of many proteins [72,73]. Accumulating intracellular protein aggregates, such as misfolded tau, are thought to compete with functional endogenous proteins for binding to chaperones, creating dysfunctional, high molecular weight tau-chaperone complexes containing HSc70 and 90 [74]. The formation of this dysfunctional epichaperome (high molecular weight complexes composed of HSP90 and HSc70 chaperones) renders neurons prone to degeneration [74]. AstTau exhibits accumulation of HSc70, as shown by native gel immunoblotting (**Fig. 6a**). The AstTau cultures were treated with the epichaperome specific HSP90 inhibitor PU-H71 (1 µM, 3 days), which has been shown to disrupt the epichaperome complex (**Fig. 6b**) [74]. Subsequent analysis of pathology in AstTau showed a striking reduction in tau pathology and a corresponding reduction of neurodegeneration at 21 DIV3D (**Fig. 6c-j**). We observed a significant reduction in tau phosphorylation (pS202/5, pS262), misfolding (MC1 antibody), oligomerization (TOMA2 antibody) and tangle formation (ThioS) (**Fig. 6c-f**). Fluoro Jade B reactivity and LDH release also decreased, indicating a decrease in neuronal cell death (**Fig. 6h-j**). This reduction in neuronal stress and death was also accompanied by a decrease in reactive astrogliosis, marked by decreased S100β and GFAP staining as compared to oTau+/PU-H71- cultures (**Fig. 6 e, g**).

Analysis of the transcriptional responses by scRNA-seq supports the putative benefit observed for PU-H17, with the transcriptional profile shifting to that observed early in the disease process, perhaps prior to a point where the system was overwhelmed by oTau (**Fig. 7**). PU-H71 scRNA-seq cell transcriptomes were integrated with vehicle treated cell transcriptomes and presented the same cell type clustering and trajectory (**Fig. 7a**). We observed a reduction in transcriptional changes associated with neuronal cell stress and death, including suppression of the

neuroinflammatory transcripts in the EX_NEU population and the HSP chaperone transcripts in IN_NEU **(Fig. 7b)**. Interestingly, PU-H71 initiated a striking re-expression of ribosomal transcripts at 21 DIV3D similar to that observed at 7 DIV3D in all cell types **(Fig. 7b)**. The ASC population also demonstrated a reduction in the upregulation of the A1 toxic astrocytic signature and an increase in the upregulation of the A2 protective astrocytic signature **(Fig. 7c).** Neurons no longer demonstrated an upregulation of AD associated neuronal signatures in the PU-H71 treated AstTau spheroids **(Fig. 7d)**. Additionally, PU-H71 reversed the upregulation of neuroinflammatory signatures in EX_NEU suggesting a mechanism of protection **(Fig. 7d)**. In these analyses the oTau-/PU-H71+ condition is a useful comparison to assess the cellular response to oTau exposure in the presence of PU-H71 versus the transcriptional alterations caused by PU-H71 itself. The ASC population demonstrated a slight decrease in intensity of heat shock and gliosis reactions with PU-H71 treatment at 21 DIV3D corresponding with the decreased GFAP immunolabeling, together suggesting an altered reactive astrocyte phenotype **(Fig. 7e)**. Notably, the reversal of late state upregulated signatures was more pronounced in the EX_NEU population than either the ASC or IN_NEU populations, indicating a potentially cell type specific targeted treatment effect **(Fig. 7e,f, Suppl. Fig. 11).** EnrichmentMap presentation of the 7 DIV3D and 21 DIV3D +/- PU-H71 upregulated functional gene set enrichment pathways in EX_NEU demonstrates the pathway enrichment overlap between 7 DIV3D and 21 DIV3D + PU-H71 that is not shared by the cell stress enriched pathways in the 21 DIV3D – PU-H71 condition **(Fig. 7f).** Together these data suggest that targeting the dysfunctional epichaperome with PU-H71 significantly reduces oTau induced pathology and neurodegeneration in AstTau, accentuating the involvement of HSPs in tauopathies.

DISCUSSION

We have presented a manipulatable and accessible method for modeling tau induced degenerative pathology in a human iPSC derived coculture of neurons and astrocytes in a rapid and reproducible manner. Human oTau seeding in hiNCs before combination with hiACs allows for a discrete analysis of how neuronal tau pathology develops and interacts with neighboring astrocytes in a physiologically relevant spatial context. This targeted and controlled methodology allows for a deeper level of study of tau induced phenotypes *in vitro* beyond that previously shown [28-30,32,33,75,76]. Previous studies with microfluidic and organoid models have developed some

phenotypes of interest such as tau hyperphosphorylation and neurodegeneration in glutamate toxicity challenged MAPT mutant iPSCs [75],[29]. Models such as these and others including 2D models develop tau hyper-phosphorylation but no neurodegeneration [12-20], and represent partial models of tauopathy. By focusing on the oligomeric tau fraction we were able to induce robust pathology and neurodegeneration over a time course that extends well beyond any acute treatment; this approach also highlights an emerging consensus that oTau is more toxic than fibrillar tau [55,56]. The result is a broader and more clinically relevant model of tau induced neurodegeneration from oTau seeding and propagation through fibrillar tau formation and neurodegeneration in just three weeks.

The AstTau system allows for a detailed analysis of astrocytic phenotypes that are associated with tau pathology in a manner not easily studied with *in vivo* models that take 6-8 months to develop glial populations [29,77]. The incorporation of hiACs in the AstTau model is necessary for neuronal survival in 3D culture, while also enabling study of the evolution of the interaction over the disease course. AstTau exhibits remarkable astrocytic stress responses resulting in expansion of the astrocytic compartment, selective transcriptional responses, and changes in morphology consistent with reactive astrogliosis, including increased GFAP expression in AstTau. This early coordinated stress response indicates close crosstalk between neurons exposed to tau oligomers and the otherwise healthy astrocytes. These results highlight the developing importance of glial cell types such as astrocytes to the pathogenesis of neurodegenerative diseases, where they have been found to play roles in inflammation and cell death mediation [57,58]. The reactive astrogliosis in AstTau resembles astrocytic responses observed in AD brain tissue, highlighting the ability of the AstTau system to reproduce physiologically relevant cell type responses [58].

Importantly, we have demonstrated how the observed pathologies are selective for oTau as shown by the absence of such pathology observed in immunodepleted oTau and PFF-αSYN treated controls. Neither tau pathology nor neurodegeneration developed upon treatment with immunodepleted oTau, nor did it occur in neurons or astrocytes treated with PFF-αSYN. Despite the lack of toxic response, PFF-αSYN did induce wide-spread phosphorylated α-synuclein that was evident by immunolabeling. Together these data suggest that the particular degenerative tauopathy observed in AstTau is selective for the type of protein aggregate.

This study provides the first single cell transcriptomic analysis of a 3D coculture based seeded tauopathy model. Informatic analysis identified astrocytic and neuronal cell populations in the AstTau model, and we were able to define inhibitory GABAergic and excitatory glutamatergic neuronal sub-populations produced by a NGN2 overexpression differentiation system [78]. These neuronal populations showed distinct responses to toxic oTau exposure, recapitulating phenotypes observed in postmortem AD brain tissue [60,69,71,79]. There was a progressive loss of glutamatergic neurons in AstTau, consistent with reports that show excitatory neurons are more susceptible to neurodegeneration in tauopathies [80]. Differential gene expression revealed that the glutamatergic neurons mounted a more robust late stage neuroinflammatory response than GABAergic neurons, offering insight to the mechanism of cell type specific degeneration. Additionally, the astrocytic transcriptomic response to neuronal injury was surprisingly rapid and synchronized with the neuronal responses. Astrocytes and neurons both mounted early robust ribosomal responses, consistent with an adaptation to stress [81-86]. Astrocytes were notable for mounting a more robust late stage HSP response as well as presenting with upregulation of autophagic and astrogliosis pathways, perhaps contributing to their underlying resilience and enabling regulation of neuronal response [87,88]. These AstTau specific reactive astrocytes showed remarkable transcriptomic similarity to AD specific disease associated astrocytes (DAAs) identified by Habib et al.,2020 [61]. Additionally, the transcriptomic signature correlates more significantly with the neurotoxic A1 astrocyte signature than the A2 neuroprotective signature, as defined by Zamanian et al., 2012 [68], supporting a potential detrimental role for reactive astrocytes in tauopathy.

Our studies suggest that disruption of chaperone function mediated by tau oligomers contributes to the pathophysiology of tauopathies. Previous work has noted the potential importance of the HSP chaperone system in neurodegeneration[73,89-95]. The HSP chaperone response in AstTau was shared with recent single nuclei postmortem late-stage AD transcriptomic studies [69,71] . Based on this, we hypothesized that the accumulating tau pathology might overwhelm the HSP chaperone system leading chaperone dysfunction. This hypothesis is supported by recent work from Chiosis and colleagues who identified the 'epichaperome', which is a dysfunctional accumulation of HSP chaperone in disease [74,96,97]. They have shown that treatment with PU-H71 disrupts this dysfunctional epichaperome, producing more functional molecular chaperones (HSc70 and HSP90) in 2D iPSC and mouse models of tauopathy [74]. We have now shown striking protection by PU-H71 treatment in the iPSC AstTau system. The accumulation of oTau was associated with

formation of high molecular weight epichaperome complexes while dispersion of these complexes by treatment with PU-H71 rescued neurons from degeneration [74]. We saw a corresponding normalization of neuroinflammatory transcriptional signatures, as well as a reactivation of the ribosomal stress response, suggesting that this response can be beneficial. It is notable that this was a short treatment with PU-H71 late in the disease process provided benefit, suggesting that epichaperome inhibition could be a feasible option for treating the development of tauopathies at varying stages of progression. This provides strong support of the putative benefit of PU-H71 in ongoing clinical trials. It also demonstrates the utility of the AstTau system as a translational tool for drug development.

AstTau was developed to begin to address the need for modeling the contribution of toxic tau oligomers to the pathogenesis of tauopathies in a physiologically relevant human context. The use of iPSC derived neurons approaches this goal, however as with all iPSC neurons the cells express mostly 3R tau. This challenge remains to be solved. Additionally, the model is currently limited in the range of glial cells represented, but future studies will continue to expand the cell type composition and interactions by including microglia, oligodendrocytes, and potentially vascular cell types. Additionally, while this current study focuses on the specific impact of toxic tau oligomers in tauopathy development in healthy iPSC derived cells, the AstTau model is well suited to be used with a range of iPSC backgrounds including those with mutations that have been shown to contribute to disease progression. Future studies with AstTau could model dementia subtypes by propagating varying oligomeric tau conformers in neurons or astrocytes carrying specific disease-associated genetic mutations. These strengths of the rapid and reproducible AstTau model evidence a powerful and accessible new tool for the study of tauopathies.

ACKNOWLEDGEMENTS

We thank Dr. Todd A. Blute for his technical support at BU's Proteomics and Imaging Core Facility. We thank Rakez Kayed (UT Galveston) for provision of the TOMA2 antibody, Nicholas M. Kanaan (Michigan State University) for provision of the Tau13 antibody (originally created by Lester I Binder, Northwestern University), and Peter Davies (Northwell/Hofstra, deceased) for provision of CP13 antibody. We would like to thank the following funding agencies for their support: B.W. & C.S.C : Kilachand Fund of the BU Center for Life Sciences Engineering, B.W.:

AUTHOR CONTRIBUTIONS

H.D.R. designed and performed the experiments and bio-informatics analysis, interpreted the results, and wrote the paper. L.J. designed, performed, and analyzed the experiments, interpreted the results, and wrote the paper. R.H. helped design statistical tests for scRNA-seq analysis. N.K.O'N. helped design statistical tests for scRNA-seq analysis. M.M. helped perform experiments. L.Z. helped analyze the experiments. .D.S. helped design scRNA-seq analysis. B.W. conceived and oversaw the project, provided guidance, interpreted the results, and wrote the paper. C.S.C. conceived and oversaw the project, provided guidance, interpreted the results, and wrote the paper.

DECLARATION OF INTERESTS

The authors declare the following competing interests:

B.W. is co-founder and Chief Scientific Officer for Aquinnah Pharmaceuticals Inc.

B.W., C.S.C, L.J, and H.D.R submitted a patent application through Boston University entitled "Modeling Alzheimer's disease using 3D iPSC cultures."

FIGURES

Figure 1

FIGURE 1: 3D iPSC derived neuron and astrocyte asteroid coculture rapidly recapitulates neuronal and astrocytic interactions and signatures.

a. Experimental schematic diagraming 2D culture of hiPSC derived neurons and astrocytes, 3D asteroid coculture, and immunolabeling and single cell transcriptomic analysis at indicated data collection timepoints.

b. Representative immunolabeling of 21 DIV3D asteroids with cell type markers, including astrocytic precursor markers (VIM), astrocytic markers (EAAT1, GFAP), neuronal precursor markers (SOX2, NES), neuronal markers (MAP2, NEUN), and synaptic markers (DLG4/PSD95, SYP).

c. UMAP of 26,789 single cell transcriptomic profiles by scRNA-seq after filtration and cell type identification, including all cells from 7,14, and 21 DIV3D (see Methods). Five main cell types are identified, excitatory neurons (EX_NEU, dark green), inhibitory neurons (IN_NEU, light green), neuron precursors (NEU_P, teal), astrocytes (ASC, dark red), and astrocyte precursors (ASC_P, light red).

d. Dot plot of key cell type marker transcript expression including cycling, astrocytic, pan-neuronal, GABAergic inhibitory neuronal, glutamatergic excitatory neuronal, and synaptic markers in cell type clusters.

e. UMAP projection colored by timepoint across the 7-21 DIV3D time course (blue gradient) highlighting the model trajectory.

f. Percent composition of cell type clusters across the 7-21 DIV3D time course, indicating a decrease in precursor cell proportions and a correlated increase in mature NEU and ASC proportions.

g. Quantification of select immunolabeled cell type markers across the 7-21 DIV3D time course (blue gradient), indicating a decrease in precursor markers (SOX2, VIM) and an increase in mature neuronal marker (MAP2). Data obtained from 5 independent asteroids. (ANOVA, ns : $p > 0.05$, *: $p <= 0.05$, **: $p <= 0.01$).

h. Violin plot representation of the same select cell type marker transcript expression (SOX2, MAP2, VIM) across the 7-21 DIV3D time course (blue gradient) demonstrating a correlated gene expression trajectory by scRNA-seq (ANOVA).

i. UMAP feature plot indicating robust gene expression of neuronal synaptic markers DLG5 (PSD95) and SYP.

j. UMAP integration of the asteroid scRNA-seq dataset (Asteroid, red) with a published cerebral organoid scRNA-seq dataset (Cerebral Organoid, blue [38])highlighting over representation of the asteroid cells in more mature cell clusters (see Methods).

Figure 2

FIGURE 2. oTau seeding induces rapid tau misfolding and aggregation in the 3D neuron and astrocyte coculture AstTau model.

a. Experimental schematic diagraming 2D culture of hiNCs and hiACs, selective exposure of hiNCs to oTau, 3D asteroid coculture, and immunolabeling and single cell transcriptomic analysis at indicated data collection timepoints.

b. Representative images showing the progressive accumulation of toxic tau oligomers in oTau seeded asteroids at 7, 14 and 21 DIV3D. The tau oligomers are labeled by TOMA2 (green). Neurons were labeled by Tuj1 (βIII tubulin, violet) and astrocyte were labeled by S100β (red). Scale bars = 50μm.

c. Quantification of oTau by fluorescence intensity of TOMA2, normalized to the corresponding DAPI intensity presented as % of WK-1 control. Data obtained from 10 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, **$p<0.01$, ****$p<0.001$ comparisons to vehicle control at each time point.

d. Representative images showing tau misfolding in oTau seeded asteroids at 21 DIV3D. The misfolded tau is labeled by MC1 (misfolded tau, green). Neurons were labeled by Tuj1 (βIII tubulin, violet) and astrocyte were labeled by S100β (red). The 63x bottom row shows high magnification of the highlighted regions for the oTau panels. Scale bars= 10 mm and 50μm.

e. Quantification of MC1 labeled fluorescence intensity normalized by corresponding DAPI intensity. Data obtained from 5 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, **$p<0.01$ comparisons to vehicle control at each time point. ##$p< 0.01$ comparison to 7 DIV3D.

f. Representative images showing ThioS labeling at 21 DIV3D. Neurons were labeled by Tuj1 (βIII tubulin, violet) and astrocyte were labeled by S100β (red). Scale bars =50μm.

g. Quantification of ThioS labeled fluorescence intensity normalized to corresponding DAPI intensity. Data obtained from 5 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, **$p<0.01$ comparisons to vehicle control at each time point. ##$p< 0.01$ comparison to 7 DIV3D.

Figure 3

FIGURE 3. oTau seeding induces rapid accumulation of toxic tau and neurodegeneration in the 3D neuron and astrocyte coculture AstTau model.

a. Representative images showing tau hyperphosphorylation and reactive astrogliosis in oTau seeded asteroids at 14 DIV3D. Hyperphosphorylated tau is labeled with AT8 (pS202/205 tau, red) and activated astrocyte were labeled with GFAP (green). Neurons were labeled by Tuj1 (βIII tubulin, violet). Scale bars = 50μm.

b. Quantification of tau hyperphosphorylation labeled by AT8 fluorescence intensity normalized by corresponding DAPI intensity at 14 DIV3D. Data obtained from 5 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, ***$p < 0.01$.

c. Immunoblotting and quantification of hyper-phosphorylated tau in AstTau. The hyper - phosphorylated tau was detected by AT8 antibody (detecting phosphorylated Serine 202/205) and GAPDH was used as internal control. AT8 band intensity was normalized to internal control. Data obtained from 3 independent asteroid pools. Error bars = SEM. Unpaired t test was performed, *$p < 0.05$.

d. Quantification of reactive astrogliosis labeled by GFAP fluorescence intensity normalized by corresponding DAPI intensity at 14 DIV3D. Data obtained from 5 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, ****$p < 0.001$.

e. Representative images quantifying neurodegeneration with Fluoro Jade Blue labeling at 21 DIV3D. Neurons were labeled with Tuj1 (βIII tubulin, violet) and astrocyte were labeled with S100β (red). Scale bars = 50μm.

f. Quantification of astrocytic S100β labeled fluorescence intensity, normalized to the corresponding DAPI intensity. Data was collected from 5 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, **$p < 0.01$ comparisons to vehicle control at each time point. [##]$p < 0.01$ comparison to 7 DIV3D.

g. Quantification of Fluoro Jade Blue labeled fluorescence intensity normalized by corresponding DAPI intensity. Data obtained from 5 independent asteroids. Error bars = SEM. Two-way

ANOVA with Tukey's multiple comparisons test was performed, **$p<0.01$ comparisons to vehicle control at each time point. ##$p< 0.01$ comparison to 7 DIV3D.

h. Immunoblotting analysis of cell death pathway activation as marked by BAX and cleaved BAX protein level in AstTau. BAX and cleaved BAX was detected by BAX antibody and GAPDH was used as internal control.

i. Quantification of BAX band intensity normalized to the internal control in the immunoblotting analysis indicated in *h*. Data obtained from 3 independent asteroid pools. Error bars = SEM. Unpaired t test was performed, **$p<0.01$.

j. Quantification of cleaved BAX band intensity normalized to the internal control in the immunoblotting analysis indicated in *h*. Data obtained from 3 independent asteroid pools. Error bars = SEM. Unpaired t test was performed, *$p<0.05$.

k. Representative image of a mixed (NGN2_GFP+/oTau - and NGN2_GFP-/oTau +) culture indicating tau propagation from the NGN2_GFP-/oTau + neurons to the NGN2_GFP+/oTau - neurons as evidenced by NGN2_GFP+/MC1+ regions. The misfolded tau is labeled with MC1 (misfolded tau, red). Neurons are labeled with Tuj1 (βIII tubulin, violet) and select neurons are NGN2_GFP+. Closed arrows indicate regions of propagated misfolded tau. Scale bars = 50μm.

l. Representative images of propagated tau misfolding as marked by MC1 (red) in three conditions, control, oTau, and control/oTau mixed cultures. Scale bars= 50μm.

m. Quantification of MC1 labeled fluorescence intensity normalized by corresponding DAPI intensity as shown in *l*. Data obtained from 7 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, *$p<0.05$.

Figure 4

FIGURE 4: scRNA-seq of AstTau reveals dynamic transcriptional response to oTau induced pathology.

a. UMAP projection of 130,605 single cell transcriptomes colored by conditional timepoints across the 7-21 DIV3D control and AstTau time course (blue and orange gradients respectively) highlighting the model integration and trajectory.

b. Percent composition of cell type clusters in control and AstTau across the 7-21 DIV3D time course indicating loss of EX_NEU and expansion of the ASC compartment with oTau treatment.

c. Heatmap representation of the log2FC of the top 25 up and downregulated DEGs in AstTau across the 7-21 DIV3D time course in the three main cell types, EX_NEU, IN_NEU, and ASC. Transcripts associated with the GO Cytosolic Ribosomal, HALLMARK TNFα Signaling via NFKB, and the REACTOME HSF1 Activation pathways are highlighted in green, purple, and red respectively.

d. Single cell average gene expression in the control (blue) and AstTau (orange) ASC populations at 21 DIV3D highlighting the correlation of AstTau ASC transcriptomic signature with published astrocytic signatures including A1 toxic and A2 protective astrocytes (Zamanian et al., 2012 [68]), AD associated astrocytes (Habib et al., 2020 [61]), and the Reactome HSF1 pathway regulating the HSP chaperone system (see Supplemental SourceTable 1. Genesets). (t.test, ns : p > 0.05, *: p <= 0.05, **: p<= 0.01, *** : p<=0.001, ****: p<= 0.0001). Inset box plots show the median, lower and upper hinges that correspond to the first quartile (25th percentile) and third quartile (75th percentile), and the upper and lower whiskers extend from the smallest and largest hinges at most 1.5 times the interquartile range. Mean values numerically presented.

e. Single cell average gene expression in the control (blue) and AstTau (orange) EX_NEU and IN_NEU populations at 21 DIV3D highlighting the correlation of AstTau neurons with published single nuclei AD neuronal signatures (Grubman et al., 2019 [69]), and the cell type differential response in the HALLMARK TNFα signaling via NFKB neuroinflammatory response and concurrent response in the GO Cytosolic Ribosomal pathway (see Supplemental SourceTable 1. Genesets). (t.test, ns : p > 0.05, *: p <= 0.05, **: p<= 0.01, *** : p<=0.001, ****: p<= 0.0001). Inset box plots show the median, lower and upper hinges that correspond to the first quartile (25th percentile) and third quartile (75th percentile), and the upper and lower whiskers extend from the

smallest and largest hinges at most 1.5 times the interquartile range. Mean values numerically presented.

Figure 5

FIGURE 5: Functional gene set enrichment of scRNA-seq DEGs reveals oTau induced pathway perturbations in AstTau.

a. Top 25 functional gene set enrichment pathways by p-value of significant upregulated DEGs (see Methods, p <0.05, log2foldchange > 0.25) between AstTau and control for each timepoint in EX_NEU, IN_NEU, and ASC cell populations presented as –log10 p-value highlighting trends of cell type specific pathway perturbations across the 7-21 DIV3D time course. Manual annotation of shared gene set features presented for clarity. (See Suppl. Fig. 10, Suppl. SourceTable2.Fig5a7a for full annotations).

b. EnrichmentMap clustering presentation of functional gene set enrichment results (see Methods, p < 0.05) with manual annotations across the 7 (light orange), 14 (orange), and 21 DIV3D (dark orange) time course for significantly upregulated DEGs in EX_NEU and ASC, identifying a wide range of cell type specific upregulated responses in AstTau. An HSF1 associated cluster is highlighted in red.

c. Top 10 functional gene set enrichment pathways from the 21 DIV3D AstTau model and two published AD snRNA-seq datasets (Grubman et al., 2019 [69] and Mathys et al., 2019 [71]) presented as -log10 p-value alongside functional gene set enrichment results from a published autism snRNA-seq dataset (Velmeshev et al., 2019 [98]) and the 21 DIV3D AstαSYN model highlighting the enrichment similarity between AstTau and AD brain transcriptomes and the disparity between AstTau and AstαSYN and ASD brain transcriptomes (see Methods).

Figure 6

FIGURE 6: PU-H71 treatment reduces oTau induced pathology and neurodegeneration.

a. Representative blot image showing the native page gel analysis of heat-shock protein HSC-70 in AstTau. GAPDH was used as internal control. Quantification of HSC-70 band intensity in the native page gel showed the significant increase of HSC-70 in AstTau. Data obtained from 4 independent experiments. Error bars = SEM. Unpaired t test was performed, **$p<0.01$.

b. Experimental schematic including 2D culture, oligomer exposure, 3D culture, PU-H71 treatment, and analysis with indicated data collection timepoints.

c. Representative images showing reduction in tau misfolding in oTau seeded asteroids upon exposure to PU-H71 for 3 days, from 19 – 21 DIV3D. The misfolded tau is labeled by MC1 (misfolded tau, green). Neurons were labeled by Tuj1 (βIII tubulin, violet) and astrocyte were labeled by S100β (red). Scale bars = 20 μm.

d. Quantification of MC1 labeled fluorescence intensity normalized to the corresponding DAPI intensity. Data obtained from 5 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, ***$p<0.001$ comparisons to vehicle control at each time point.

e. Representative images showing reduction in ThioS, indicating fibrillar tau, in oTau seeded asteroids upon exposure to PU-H71 for 3 days, from 19 – 21 DIV3D. The misfolded tau is labeled with MC1 (green). Neurons were labeled by Tuj1 (βIII tubulin, violet) and astrocyte were labeled with S100□ (red) and GFAP (blue). Scale bars =50 μm.

f, g. Quantification of ThioS (E) and GFAP (E) labeled fluorescence intensity normalized to corresponding DAPI intensity. Data obtained from 5 independent asteroids. Error bars = SEM. Two-way ANOVA with Tukey's multiple comparisons test was performed, ***$p<0.001$, **$p<0.001$ comparisons to vehicle control at each time point.

h. Representative images showing Fluoro Jade B labeled fluorescence in AstTau upon exposure to PU-H71 for 3 days, from 19 – 21 DIV3D. The damaged neurons are labeled by Fluoro Jade B (green). Neurons were labeled with Tuj1 (βIII tubulin, violet) and astrocyte were labeled with S100β (red) and GFAP (blue). Scale bars = 50 μm.

i. Quantification of Fluoro Jade B labeled fluorescence intensity normalized to the corresponding DAPI intensity. Data obtained from 5 independent asteroids. Error bars = SEM. Two-way

ANOVA with Tukey's multiple comparisons test was performed, ***p<0.001 comparisons to vehicle control at each time point.

j. LDH detection in the conditioned medium collected from 21 DIV3D asteroids. Data obtained from 25 independent asteroids. Error bars = SEM. One-way ANOVA with Tukey's multiple comparisons test was performed, **p<0.01.

Figure 7

FIGURE 7: scRNA-seq transcriptomic modulation supports PU-H71 reversal of the pathological molecular phenotype in AstTau.

a. UMAP projection of 130,605 cells colored by conditional timepoints across the 7-21 DIV3D control and AstTau time course with vehicle treatment (blue and orange gradients respectively) and control and AstTau time course with PU-H71 treatment (green and purple gradients respectively) highlighting the model integration and trajectory.

b. Heatmap representation of the log2FC of the top 25 up and downregulated DEGs in AstTau at 7 DIV3D and 21 DIV3D +/- PU-H71 treatment in the three main cell types EX_NEU, IN_NEU, and ASC highlighting the similarity between 7 DIV3D and 21 DIV3D with PU-H71 treatment. Transcripts associated with the GO Cytosolic Ribosomal, HALLMARK TNFα Signaling via NFKB, and the REACTOME HSF1 Activation pathways are highlighted in green, purple, and red respectively.

c. Single cell average gene expression in the oTau-/PU-H71- (blue), oTau+/PU-H71-(orange), oTau-/PU-H71+ (green) and oTau+/PU-H71+ (purple) ASC populations at 21 DIV3D highlighting the impact of PU-H71 treatment on the A1 toxic and A2 protective astrocytic signature (Zamanian et al., 2012) [68], see Supplemental SourceTable 1. Genesets). (t.test, ns : $p > 0.05$, *: $p <= 0.05$, **: $p<= 0.01$, *** : $p<=0.001$, ****: $p<= 0.0001$). Inset box plots show the median, lower and upper hinges that correspond to the first quartile (25th percentile) and third quartile (75th percentile), and the upper and lower whiskers extend from the smallest and largest hinges at most 1.5 times the interquartile range. Mean values numerically presented.

d. Single cell average gene expression in the oTau-/PU-H71- (blue), oTau+/PU-H71-(orange), oTau-/PU-H71+ (green) and oTau+/PU-H71+ (purple) ASC populations at 21 DIV3D highlighting the impact of PU-H71 treatment on the neuronal signatures including published single nuclei AD neuronal signatures (Grubman et al., 2019 [69]), the HALLMARK TNFα signaling via NFKB neuroinflammatory response, and the GO Cytosolic Ribosomal pathway (see Supplemental Source Table 1. Genesets). (t.test, ns : $p > 0.05$, *: $p <= 0.05$, **: $p<= 0.01$, *** : $p<=0.001$, ****: $p<= 0.0001$). Inset box plots show the median, lower and upper hinges that correspond to the first quartile (25th percentile) and third quartile (75th percentile), and the upper and lower whiskers extend from the smallest and largest hinges at most 1.5 times the interquartile range. Mean values numerically presented.

e. Top 25 functional gene set enrichment pathways by p-value of significant upregulated DEGs (see Methods, p <0.05, log2foldchange > 0.25) between AstTau and control for each timepoint in EX_NEU, IN_NEU, and ASC cell populations +/- PU-H71 presented as –log10 p-value highlighting cell type specific pathway perturbations that are ameliorated by PU-H71 treatment. Manual annotation of shared gene set features presented for clarity. (See Suppl. Fig. 10, Suppl. SourceTable2.Fig5a7a for full annotations).

f. EnrichmentMap clustering presentation of functional gene set enrichment results (see Methods, p < 0.05) with manual annotations across the 7 DIV3D -vehicle (light orange), 21 DIV3D +PU-H71 (purple), and 21 DIV3D-PU-H71 (dark orange) for significantly upregulated DEGs in EX_NEU, identifying a range of upregulated responses in AstTau that are ameliorated with PU-H71 treatment. An HSF1 associated cluster is highlighted in red.

MATERIALS AND METHODS

Oligomer Processing

Generation of S1p oTau fraction: Frozen hippocampus and cortical tissues of 9-month-old PS19 mice were weighed (100mg-250mg) and put in Beckman Centrifuge Tubes, polycarbonate thick wall (cat # 362305). A 10 × volume of homogenization buffer, Hsaio TBS buffer (50 mM Tris, pH 8.0, 274 mM NaCl, 5 mM KCl) supplemented with protease and phosphatase inhibitor cocktails (Roche, cat#05892791001 and cat#04906837001), was used to homogenize brain tissue as described previously [55,63]. Briefly, the homogenate was centrifuged at 48,300 g for 20 min at 4 °C. The supernatant was then centrifuged a second time at 186,340 g at 4°C for 40 min. The TBS-extractable pellet (S1p) fraction was resuspended in a 4x volume of TE buffer relative to the starting weight of the tissue homogenate, aliquoted and frozen at -80°C.

S1p oTau fraction quantification: The molecular weight of tau in the S1p fractions was documented by native page gel electrophoresis, and the concentration of total tau were measured by immunoblot using 3-12% reducing SDS-PAGE gel by comparison to a gradient concentrations of recombinant tau ladders, using the human specific tau13 antibody (detecting total human tau) by immunoblot as described previously (**Suppl. Fig. 4**)[55] All the fractions were then normalized and divided into fractions of 20µg/ml tau for storage and future use.

Immunodepletion of tau from S1p oTau fraction: Tau aggregates in S1p oTau fractions were eliminated by a direct immuno-precipitation kit (Pierce, cat# 26148). Briefly, the tau-5 antibody was coupled to AminoLink plus Coupling Resin, and the fractions were pre-cleared using the Control Agarose Resin with all the materials provided by the kit. The sample was added to the antibody-coupled resin in the spin column and incubated in the column for overnight at 4°C on a gentle rotator. The column was centrifuged, and the flow-through saved for further experimentation. After 3 washes with IP buffer, the spin column was placed into a new collection tube, and tau plus antibodies were eluted from the resin. The eluate was analyzed for presence of tau.

FITC labeling of S1p oTau fraction: The DyLight® 488 Conjugation Kit / DyLight® 488 Labeling Kit (Abcam, cat#ab201799) was used for a simple and quick process for DyLight® 488 labeling / conjugation of S1p oTau fraction. Detailed steps for conjugation of the S1p oTau fraction were followed to the manufacture's instruction. Briefly, modifier was added to S1p oTau fraction aliquot and incubated for 15 mins followed by adding quencher and incubation for 5 mins. The DyLight® 488 conjugated oTau fraction can be used immediately in cell culture treatment.

Generation and quantification of PFF-αSYN: Preformed fibrillar α-synuclein (PFF-αSYN) was produced following standardized methodology [99]. Human α-synuclein monomers (Proteos cat# RP-003) were thawed and centrifuged at 15,000 x g for 10 mins at 4 °C. The supernatant was transferred to a 1.5 mL microcentrifuge tube and the protein concentration was measured by 280 nM absorbance spectrophotometry. Protein concentration was then adjusted to 5 mg/mL with 1X dPBS and monomers were agitated in an orbital thermomixer with heated lid at 1,000 RPM for 7 days at 37 °C. Fibrils were quantified by a Thioflavin T assay and transmission electron microscopy.

Cell Culture and Treatment

All cell cultures were maintained at 37 °C with 5% CO_2. All cell counts were performed in quadruplicate using the Cellometer K2 with AOPI viability dye (Nexcelom) or the Countess3 with 0.4% trypan blue dye (Thermo Fisher).

Neural Progenitor Cell (NPC) Culture: Human iPSC (XCL-1) derived neural progenitor cells (NPCs, Stem Cell Tech cat#70901) were maintained in serum-free STEMdiff™ Neural Progenitor Medium 2 (Stem Cell Tech cat#08560) on Corning® Matrigel® hESC-qualified Matrix (Corning

cat#354277) coated tissue culture plates. NPCs were plated at 50,000 cells/cm2 and passaged at 90% confluency by Accutase^{TM} (Stem Cell Tech cat#07920) dissociation as necessary. A full media change was performed every other day. Low passage (passage <3) NPCs were cryopreserved in STEMdiff™ Neural Progenitor Medium 2 with 10% DMSO, and all NPCs used for experimentation were maintained at passage <6.

iPSC Neuronal Cell (hiNC) Differentiation: NPCs were passaged and plated at 50,000 cells/cm2 in STEMdiff™ Forebrain Neuron Differentiation Media (Stem Cell Tech cat#08600) on Corning® Matrigel® coated tissue culture treated plates and transduced with a NEUROG2 lentivirus (GeneCopoeia cat#LPP-T7381-Lv105-A00-S) at MOI 3 to induce iPSC derived neuronal cells (hiNC). After 24 hours of transduction, a full media change was performed.

iPSC Astrocytic Cell (hiAC) Differentiation: iPSC derived astrocytic cells (hiAC) were differentiated from NPCs by small molecule differentiation in STEMdiff™ Astrocyte Differentiation Media (Stem Cell Tech #cat100-0013) on Corning® Matrigel® coated tissue culture treated plates. A full media change was performed daily for four days, and NPC cultures were passaged at 90% confluence by Accutase^{TM}. NPCs were reseeded at 150,000 cell/cm2 and culture was continued in STEMdiff™ Astrocyte Differentiation Media with a full media change every other day for 14 days, passaging as necessary with Accutase^{TM}. At this stage the differentiated Astrocyte Precursor Cells (APCs) were cryopreserved in STEMdiff™ Astrocyte Differentiation Media with 10% DMSO. At time of experimentation APCs were thawed and plated at 150,000 cell/cm2 in STEMdiff™ Astrocyte Maturation Media (Stem Cell Tech cat#100-0016) on Corning® Matrigel® coated tissue culture treated plate. A full media change was performed every other day for 6 days, with one passage by Accutase^{TM} at 90% confluence as necessary.

Asteroid Generation and Maintenance: A single cell suspension of hiNCs and hiACs was prepared by Accutase^{TM} dissociation and washed once with DMEM/F12 (Stem Cell Tech cat#36254) to remove debris. hiNCs and hiACs were combined at a 1:1 ratio in Asteroid Media (DMEM/F12 (Stem Cell Tech cat#36254) , 1% Glutamax (Thermo Scientific cat#35050061), 1% Sodium Pyruvate (Thermo Scientific cat#11360070), 1% N-2 Supplement (Thermo Scientific cat#17502-048), 1% B-27 Supplement (Thermo Scientific cat#17504044), 10 uM Y-27632 (EMD Millipore cat#SCM075) 1% PenStrep (Thermo Scientific cat#15140148), 1 mg/mL Heparin (Sigma-Aldrich cat#H3149-250KU)) and plated in AggreWell^{TM}800 microwells (Stem Cell Tech cat#34815)

coated with Anti-Adherence Rinsing Solution (Stem Cell Tech cat#07010). The AggreWell plate was immediately centrifuged at 100xg for 3 minutes to capture the cells in the microwells and incubated for 24 hours. A half media change was performed at 24 hours and then every other day for one week. At one week when the spheroids displayed a smooth, bright edge under the cell culture microscope cultures were transferred to ultra-low attachment round bottom 96 well plates (Fisher Scientific cat#07-201-680) and maintained in 100-200 uL asteroid media rotating at 85 rpm. A half media change was performed every other day for up to three weeks. Cultures were monitored and live images captured using the AmScope or EVOS m7000 platforms.

hiNC oTau Treatment: hiNCs were selectively exposed to 0.04 mg/mL oTau by direct administration in cell culture media for 24 hours before incorporation into asteroid culture.

hiNC PFF-αSYN Treatment: Sonication of prepared fibrillar α-synuclein was performed to produce preformed fibrillar α-synuclein (PFF-αSYN) for use in experimentation. Briefly, fibrils were diluted in 1X dPBS to 4 ug/uL and sonicated with a 3.2 mm diameter probe for1 min of 60 pulses at an amplitude of 30%. hiNCs were selectively exposed to 1 ug/mL PFF-αSYN by direct administration in cell culture media for 24 hours before incorporation into asteroid culture.

PU-H71 Treatment: Asteroids were treated with 1 uM PU-H71 by direct administration in cell culture media for 72 hours before timepoint collection.

Mixed Culture: To determine the extent of seeded oTau propagation in the AstTau model the following experiment was performed. Two sets of asteroid cultures were established, NGN2_GFP+/oTau - and NGN2_GFP-/oTau +. NGN2_GFP (NEUROG2) lentivirus was transduced as previously described at MOI 3 (GeneCopoeia cat#LPP-T7381-Lv103-A00-S). At the one-week time point NGN2_GFP+/oTau - and NGN2_GFP-/oTau + asteroids were mixed and placed in 96-well culture as previously described. The asteroids merged together and were cultured to 21 DIV3D.

Sample Collection

Asteroid Fixation: At time of collection asteroids were transferred to a 1.5 mL Protein LoBind Eppendorf (Eppendorf cat#022-43-108-1) and allowed to settle. The supernatant was discarded, and asteroids were fixed in 4 °C 4% PFA in 1X PBS for 15 minutes, rotating at room temperature.

After fixation, asteroids were washed 3x for 10 minutes each with 4°C 1X PBS, rotating at room temperature. Samples were stored in 1X PBS at 4 °C.

Protein Lysate: At the time of collection 50 pooled asteroids were transferred to a 1.5 mL Protein LoBind Eppendorf (Eppendorfcat#022-43-108-1) and allowed to settle. The supernatant was discarded, and samples were flash frozen and stored at -80 °C. Samples were subsequently lysed by mechanical homogenization in 50 uL RIPA lysis buffer (150 mM NaCl, 50 mM Tris-HCL pH 7.4, 1 mM EDTA pH 8.0, 1% NP-40, 0.25% sodium deoxycholate) with Halt™ Protease Inhibitor Cocktail (Thermo Scientificcat#78430), cOmplete™ Protease Inhibitor Cocktail (Roche cat#11697498001), and PhosSTOP™ (Sigma-Aldrich cat#4906845001) for 40 seconds on ice. Lysate concentration was quantified by the Pierce BCA assay according to manufacturer's protocol (Thermo Fisher cat#23225).

Conditioned Media Collection: 50 uL of conditioned cell culture media from replicate asteroids were collected in flat bottomed 96 well plates and frozen at -20 °C.

Sample Processing

Immuno-fluorescence labeling: For immuno-labeling, selected asteroids from each condition were washed in 150µl PBS for 10 mins in a U-bottom 96-well plate and then permeabilized in 150µl PBS/0.01% Triton X-100 (PBST). The asteroids were then blocked in PBST supplemented with 5% BSA and 5% normal donkey serum for 1.5-2 hrs at room temperature (RT). After blocking, asteroids were incubated in primary antibodies diluted in 5% BSA/PBST overnight at 4°C. On the second day, the asteroids were washed 3 times in PBST, 15 min each, before they were transferred into 2° antibodies dilute (1:700 of Dylight-/Alexa-conjugated antibodies made in donkey purchased from Thermo Fisher Scientific in 5% BSA/PBST) for 2 hrs at RT. For DAPI nuclei stain, DAPI (1:10,000) was diluted in PBST and incubated with asteroids for 15 min followed by 2x washes with PBST then 1x with PBS, 10 min each. The asteroids were then mounted onto microscope glass slides in Prolong gold antifade reagent (Thermo Fisher cat#P36930) and stored in the dark until imaging. Primary antibodies used for asteroid labeling were as follows: Tuj1/beta3-Tubulin (chicken, SYSY, cat# 302 306, 1: 300), MAP-2 (rabbit, Millipore, cat# AB5622, 1: 1000), Rabbit monoclonal anti-S100β (Abcam, Cat# ab52642, 1:400); GFAP Monoclonal Antibody (Thermo Fisher Scientific, Cat#13-0300, 1:400); Mouse monoclonal anti-TOMA2 (provided by Dr. Rakez Kayed), 1:300; MC1 (provided by Dr. Peter Davies, Northwell),

1:300; CP-13 (provided by Peter Davies, Northwell), 1:300; AT-8 (Thermo Fisher, cat# MN1020), 1:300.

Thioflavin S staining: Fresh Thioflavin S (ThioS) solution was prepared by dissolving 1g of ThioS (Millipore Sigma cat#T1892) in 100 ml 80% ethanol, and stirring overnight at 4 °C, and filtering for final use. The asteroids to be stained were washed sequentially in 70% and 80% ethanol, 1 min each, prior to incubating in ThioS/80% ethanol solution for 15 min. Asteroids were then sequentially washed in 80% and 70% ethanol, 1 min each, followed by two rinses in PBS. Asteroids were mounted in Prolong Gold antifade reagent (Thermo Fisher cat#P36930) and stored in the dark until imaging.

Fluoro Jade B staining: The Fluoro Jade B reagent was purchased from EMD Millipore (Cat# AG310-30MG) and the staining protocol was followed as instructed by the manufacture. Briefly, the staining solution was prepared from a 0.01% stock solution for Fluoro-Jade B that was made by adding 10 mg of the dye powder to 100 mL of distilled water. To make up 100 mL of staining solution, 4 mL of the stock solution was added to 96 mL of 0.1% acetic acid vehicle. This results in a final dye concentration of 0.0004%. The stock solution, when stored in the refrigerator was stable for months, whereas the staining solution was typically prepared within 10 minutes of use and was not reused. Before staining, the asteroids were rinsed in distilled water and were then treated with 0.06% $KMnO_4$ solution for 15 min. Then the asteroids were stained with Fluoro Jade B working solution for 30 min followed by being washed with PBS twice for 5 min each . Asteroids were mounted in Prolong Gold antifade reagent (Thermo Fisher cat#P36930) and stored in the dark until imaging.

Immunoblotting: 1 ug total protein lysates were analyzed on the ProteinSimple SimpleWestern Jess platform with the 12-230 kDA microplate kit (ProteinSimple cat#SM-W004) according to manufacturer's instructions. Primary antibodies were detected using multiplexed chemiluminescent and fluorescent detection modules according to manufacturer's instructions (ProteinSimple cat#sDM-001, DM-002, DM-007, DM-008, DM-009, DM-010). Primary antibodies used for immunolabeling were as follows: GAPDH (mouse, ProteinTech, 60040-I-Ig, 1: 500), BAX (mouse, Santa Cruz, sc-7480, 1: 25), AT8 p202 Tau (mouse, Thermo Fisher, MN1020, 1:50). The HSC-70 in native cell lysate were analyzed with the NuPAGE™ 3 to 8% Tris-Acetate gel. Novex Tris-Glycine Native Sample Buffer and Novex Tris-Glycine Native

Running Buffer were used for protein lysis and separation in native format. For transfer of proteins to a membrane, the NuPAGE Transfer Buffer was used. The HSC-70 antibody was purchased from Thermo Scientific (cat# PA5-27337).

LDH Cytotoxicity Assay: The CytoTox 96 Non-Radioactive Cytotoxicity Assay or LDH-Glo Cytotoxicity Assay was performed as per manufacturer's instructions using 50 uL conditioned media replicates to measure lactate dehydrogenase (LDH) release (Promega cat# G1780, J2380). For the CytoTox 96 assay 490 nm absorbance readings were taken on a SpectraMaxM5plate reader with SoftMax Pro 7.1 software and data is presented as 490 nm absorbance. For the LDH-Glo assay luminescent readings were taken on a SpectraMaxM5plate reader with SoftMax Pro 7.1 software and data is presented as LDH concentration (μg/mL) calculated by standard curve.

Single Cell RNA Sequencing Sample Preparation and Sequencing: 30 asteroids per condition were pooled in a 1.5 mL Protein LoBind Eppendorf (Eppendorf cat#022-43-108-1) and allowed to settle. The supernatant was carefully discarded, and a single cell suspension was produced by incubation in 500 μL digestion buffer (Accutase^{TM} with 80 U/mL Protector RNase Inhibitor (Sigma-Aldrich cat#03335402001) for 1 hour at 37 °C with gentle pipette mixing every 10 minutes. At the end of the incubation the single cell suspension was washed with 500 μL wash buffer (0.02% BSA in 1X PBS with 80 U/mL Protector RNase Inhibitor) and passed through a 20 μM filter (MACS, Miltenyi Biotec cat#130-101-812) to a fresh 2 mL Protein LoBind Eppendorf. An additional 1 mL of wash buffer was then passed through the same filter for a total single cell suspension of 2 mL. The samples were centrifuged at 300 g for 5 minutes at 4 °C followed by another wash in 1 mL wash buffer. After another centrifugation the supernatant was discarded, and the single cell pellet gently resuspended in 50 μL wash buffer. Cells were counted in quadruplicate on the Cellometer K2 with AOPI and processed through the single cell RNA-sequencing pipeline from 10X Genomics, 3' Version 3 (10X Genomic Chromium).

Briefly, the single cell suspension was mixed with RT reaction mix to target an 8000-cell recovery and 75 μl was loaded onto a chromium microfluidics chip with 40 μL of barcoded beads and 280 μL of partitioning oil. The chip was run on the chromium controller, encapsulating a single cell and barcoded bead within individual oil droplets. Reverse transcription was performed within these individual oil droplets to produce barcoded cDNA. cDNA was then isolated by Silane DynaBeads (Thermo Fisher Scientific, DynaBeads MyONE Silane, cat# 37002D) before PCR

amplification. Amplified cDNA cleanup and size selection was performed using SPRIselect beads (Beckman-Coulter, SPRIselect, cat# B23317) and cDNA quality was assessed by the High-Sensitivity DNA assay on the Agilent 2100 BioAnalyzer (Agilent, High-Sensitivity DNA Kit, cat# 5067-4626). Sequencing libraries were then prepared according to 10X specifications, including fragmentation, sequencing adaptor ligation, and sample index PCR. Between each of these steps, library cleanup and size selection were performed by SPRIselect beads. Final cDNA library quality was assessed by the Agilent BioAnalyzer High-Sensitivity DNA assay and the Qubit High-Sensitivity DNA assay and quality-confirmed libraries were sequenced on Illumina's NextSeq 500 or NOVA-seq platform to a depth of 200 million paired-end reads. Single cell RNA-sequencing data was produced from four independent experimental replicates.

Data Analysis

Image Analysis: Images were captured by Carl Zeiss confocal LSM700. The immuno-fluorescence stained DAPI-positive cells in each image of asteroids were quantified by Image J by automated cell counting. The staining intensity in immuno-fluorescence labeled asteroids were measured by ImageJ. The intensity of MC1, TOMA2, CP13, AT8, ThioS and Fluoro Jade B were normalized by DAPI numbers. Schematics were created with BioRender.com.

GraphPad Prism statistical analysis: Statistical analyses and figures artwork were performed using GraphPad Prism version 9.00 for Windows with two sided α of 0.05. All group data are expressed as mean ± SEM. Column means were compared using one-way ANOVA with treatment as the independent variable. Group means were compared using two-way ANOVA with factors on oTau treatment and time points, respectively. When ANOVA showed a significant difference, pair wise comparisons between group means were examined by Tukey's, Dunnett or uncorrected Fisher's LSD multiple comparison test. Significance was defined when $p < 0.05$. LDH assay data analysis was performed with a paired t-test.

Single Cell RNA Sequencing Data Analysis

CellRanger Pipeline: CellRanger version 3.1.0 (10X Genomics) was used to combine and process the raw Illumina NextSeq 500 RNA and NOVA-seq sequencing files. First each sequencing library was demultiplexed by sample index to generate FASTQ files for paired-end reads using the CellRanger mkfastq pipeline. FASTQ files were then passed to the CellRanger count pipeline, which used STAR aligner [100] to align reads to the human reference genome (GRCh38). The

CellRanger aggr pipeline was then used to equalize the aligned molecule_info.h5 sample libraries across sequencing depths (by each sample cell being down-sampled to have the same confidently mapped reads per cell) and aggregated together to generate the gene-cell barcode matrix.

Seurat Object Filtration: Subsequent filtering, normalization, and scaling of data was performed using Seurat version 3.2.2 [101,102]. The Seurat object was created with a min.cells of 2 and a min.features of 200. Cells with less than 200 and greater than 3000 detected genes or greater than 12% mitochondrial counts were filtered out to discard potential doublets and low-quality captures Gene counts for each cell were normalized by total expression, multiplied by a scale factor of 10,000 and transformed to log scale. PCA based on the highly variable genes detected (dispersion of 2) was performed for dimension reduction and the top 30 principal components (PCs) were selected based on a heuristic elbow plot and jack straw plot analysis. We clustered cells based on graph-based methods (KNN and Louvain community detection method) implemented in Seurat. Clusters were visualized using uniform manifold approximation and projection (UMAP) [103].

Cluster Cell Type Identification: To identify neuronal cell type subpopulations, we performed differential expression analysis using the Wilcoxon rank-sum test implemented in Seurat between previously defined clusters with a min.pct of 0.25 and a logfc.threshold of 0.25. This identified top expressing genes for each cluster, which were then considered alongside the feature expression of canonical gene cell type markers to manually conclude cell type cluster identification. Additionally, the automated cell typing R package SingleR was used to confirm cell type identification using integrated assay with the hpca.se$label.fine from the HumanPrimaryCellAtlasData [104].

DE analysis: Differential expression analysis was performed within each cell type between control and AstTau samples using the Wilcoxon rank-sum test implemented in Seurat with a min.pct of 0.1, a logfc.threshold of 0.25, and a pseudocount of 1E4. A multiple comparison correction was performed using the Benjamin & Hochberg FDR method to produce an adjusted p-value [105]. Differentially expressed genes were evaluated according to their log fold change (greater than $\log_2(.25)$) and adjusted *p* values (<0.05). Figures were generated using the ggplot2 R package[106].

Functional enrichment analysis: Functional gene set enrichment analysis of the significant differentially expressed genes between control and AstTau samples was performed using the R implemented GProfiler2 [107]. The enrichment analysis was run as an ordered query (ordered by

log2FC) using a threshold of 0.05 and using Benjamin & Hochberg FDR for multiple testing correction [105]. Only genes in the Seurat dataset were considered by using a custom domain scope. A custom source GMT, gp__zSEF_sD9Q_d1M, was used. It includes all Hallmark gene sets, curated gene sets, and ontology gene sets from the Molecular Signatures Database (MsigDB) v7.2 [67,108]. The enrichment analysis was assessed and visualized by a heatmap of significance ($-\log10(p$ value)) of the top 25 enriched pathways per sample comparison. Figures were generated using the ComplexHeatmap R package [109].

Comparative functional gene set enrichment analysis between AstTau, AstαSYN and published datasets was performed with the same gprofiler2 settings, but as a non-ordered query. Gene input was as follows: From our AstTau dataset, common genes upregulated across IN_NEU, EX_NEU, and ASC at 21 DIV3D (n=11, p <0.05, log fold change > 0.25). From our AstαSYN dataset, common genes upregulated across IN_NEU, EX_NEU, and ASC at 21 DIV3D (n=491, p <0.05, log fold change > 0.25). From Grubman et al. 2020 [69], common genes upregulated across neurons and astrocytes by snRNA-seq of post-mortem AD brain tissue (DEG 5 and 7, n = 109, log fold change > 1 and FDR < 0.01). From Mathys et al. 2020 [71], common genes upregulated by snRNA-seq of post-mortem AD brain tissue across AST, IN, and EX cell types in Figure 2C, late versus early pathology (n=16, log fold change > 1 and FDR < 0.01)). From Velmeshev et al. 2019 [98], all highlighted genes upregulated by snRNA-seq of post-mortem autism spectrum disorder (ASD) brain tissue across neuronal and non-neuronal cell types in Figure 2 a, b (n=22, p <0.05, log fold change > 0.14). The enrichment analysis was assessed and visualized by a heatmap of significance ($-\log10(p$ value)) of the top 10 unique enriched pathways from AstTau, Grubman, and Mathys. All heatmaps were generated using the ComplexHeatmap R package and color scale generated using dependent R package Circlize [109]. Additional visualization of significant (p< 0.05) enriched pathways was performed using Cytoscape EnrichmentMap [110,111] with an edge cutoff of 0.375. Gene sets in EnrichmentMap cluster by similarity and annotations of shared gene set features were added manually using Cytoscape implemented AutoAnnotate.

Signature analysis: The scaled expression per cell of literature curated and MsigDB ontology gene sets (see Source Table 1. Genesets) was compared between control and AstTau samples by computing the mean expression using colMeans and performing an ANOVA or t.test across comparison pairs using stat_compare_means. Figures were generated using the ggplot2 R package [106].

Liger Integration: To compare cell types of our data with a different dataset [38] we merged the two datasets using Liger [112], which uses integrative Non-Negative Matrix Factorization (iNMF) to simultaneously reduce data dimensionality as well as correct the technical factors. We used Seurat V4.0.4 [113] as previously described for pre-processing before running Liger. Each dataset is normalized to account for differences in total gene-level counts across cells using the NormalizeData function. Furthermore, we used the ScaleData function with parameter do.center = FALSE, to scale normalized datasets without centering by the mean, giving the nonnegative input data required by iNMF. To run Liger, we tuned the two most important parameter k (determines the number of matrix factors in the factorized data) and lambda (the degree of dataset integration) using the suggestK and the suggestLambda function, respectively. Finally, we used k=20 and lambda=5 to integrate the datasets. The resulting integrated dataset was then analyzed using UMAP visualizations and percent cluster cell count calculations as previously described.

Data Availability: Raw and processed scRNA-seq data are available from GEO under accession GSE186356. Processed scRNA-seq datasets are available on Single Cell Portal, including the cell barcodes, UMAP coordinates, and other available characteristics. This processed scRNA-seq data is available at https://singlecell.broadinstitute.org/single_cell/study/SCP1621. The source data underlying Figs. 1,4,5,7and Suppl. Fig.2,3,9,10,11 are provided in the Source Data files as follows: Source data of differential gene expression and celltype_markers are available in supplemental file asteroid1_sourcefile_deg and source data of gprofiler2 functional gene set enrichment, Source Table 1, Source Table 2, and Source Table 3 are available in supplemental file asteroid1_sourcefile_gpro.

Code Availability: The original R scripts for Seurat processing are available on GitHub (https://github.com/satijalab/seurat). All custom code to reproduce the analyses and figures reported in this paper are available on GitHub (https://github.com/ChristineLab/Asteroid1_2021).

REFERENCES

1.   Šimić, G.*, et al.* Tau Protein Hyperphosphorylation and Aggregation in Alzheimer's Disease and Other Tauopathies, and Possible Neuroprotective Strategies. *Biomolecules* **6**, 6 (2016).
2.   Morawe, T., Hiebel, C., Kern, A. & Behl, C. Protein homeostasis, aging and Alzheimer's disease. *Mol Neurobiol* **46**, 41-54 (2012).
3.   Jellinger, K.A. & Attems, J. Neurofibrillary tangle-predominant dementia: comparison with classical Alzheimer disease. *Acta Neuropathologica* **113**, 107-117 (2007).

4.   Iaccarino, L*., et al.* Local and distant relationships between amyloid, tau and neurodegeneration in Alzheimer's Disease. *Neuroimage Clin* **17**, 452-464 (2018).

5.   Dickson, D.W. Neuropathology of non-Alzheimer degenerative disorders. *Int J Clin Exp Pathol* **3**, 1-23 (2009).

6.   Iqbal, K*., et al.* Tau pathology in Alzheimer disease and other tauopathies. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1739**, 198-210 (2005).

7.   Young, J.J., Lavakumar, M., Tampi, D., Balachandran, S. & Tampi, R.R. Frontotemporal dementia: latest evidence and clinical implications. *Ther Adv Psychopharmacol* **8**, 33-48 (2018).

8.   Arriagada, P.V., Growdon, J.H., Hedley-Whyte, E.T. & Hyman, B.T. Neurofibrillary tangles but not senile plaques parallel duration and severity of Alzheimer's disease. *Neurology* **42**, 631-631 (1992).

9.   Nelson, P.T*., et al.* Correlation of Alzheimer Disease Neuropathologic Changes With Cognitive Status: A Review of the Literature. *Journal of Neuropathology & Experimental Neurology* **71**, 362-381 (2012).

10.   Laurijssens, B., Aujard, F. & Rahman, A. Animal models of Alzheimer's disease and drug development. *Drug Discovery Today: Technologies* **10**, e319-e327 (2013).

11.   Duyckaerts, C., Potier, M.-C. & Delatour, B. Alzheimer disease models and human neuropathology: similarities and differences. *Acta neuropathologica* **115**, 5-38 (2008).

12.   Yagi, T*., et al.* Modeling familial Alzheimer's disease with induced pluripotent stem cells. *Human molecular genetics* **20**, 4530-4539 (2011).

13.   Rowland, H.A., Hooper, N.M. & Kellett, K.A.B. Modelling Sporadic Alzheimer's Disease Using Induced Pluripotent Stem Cells. *Neurochem Res* **43**, 2179-2198 (2018).

14.   Ochalek, A*., et al.* Neurons derived from sporadic Alzheimer's disease iPSCs reveal elevated TAU hyperphosphorylation, increased amyloid levels, and GSK3B activation. *Alzheimers Res Ther* **9**, 90 (2017).

15.   Muratore, C.R*., et al.* The familial Alzheimer's disease APPV717I mutation alters APP processing and Tau expression in iPSC-derived neurons. *Human molecular genetics* **23**, 3523-3536 (2014).

16.   Verheyen, A*., et al.* Using Human iPSC-Derived Neurons to Model TAU Aggregation. *PloS one* **10**, e0146127-e0146127 (2015).

17.   Nakamura, M*., et al.* Pathological Progression Induced by the Frontotemporal Dementia-Associated R406W Tau Mutation in Patient-Derived iPSCs. *Stem cell reports* **13**, 684-699 (2019).

18.   Kondo, T*., et al.* Modeling Alzheimer's disease with iPSCs reveals stress phenotypes associated with intracellular Abeta and differential drug responsiveness. *Cell Stem Cell* **12**, 487-496 (2013).

19.   Ghatak, S*., et al.* Mechanisms of hyperexcitability in Alzheimer's disease hiPSC-derived neurons and cerebral organoids vs. isogenic control. *Elife* **8**(2019).

20.   García-León, J.A*., et al.* Generation of a human induced pluripotent stem cell–based model for tauopathies combining three microtubule-associated protein TAU mutations which displays several phenotypes linked to neurodegeneration. *Alzheimer's & Dementia* **14**, 1261-1280 (2018).

21.   Verheyen, A*., et al.* Genetically Engineered iPSC-Derived FTDP-17 MAPT Neurons Display Mutation-Specific Neurodegenerative and Neurodevelopmental Phenotypes. *Stem cell reports* **11**, 363-379 (2018).

22.   Usenovic, M*., et al.* Internalized Tau Oligomers Cause Neurodegeneration by Inducing Accumulation of Pathogenic Tau in Human Neurons Derived from Induced Pluripotent Stem Cells. *The Journal of Neuroscience* **35**, 14234 (2015).

23.   Israel, M.A*., et al.* Probing sporadic and familial Alzheimer's disease using induced pluripotent stem cells. *Nature* **482**, 216-220 (2012).

24.   Hallmann, A.-L*., et al.* Astrocyte pathology in a human neural stem cell model of frontotemporal dementia caused by mutant TAU protein. *Scientific Reports* **7**, 42991 (2017).

25.   Lin, Y.-T.*, et al.* APOE4 Causes Widespread Molecular and Cellular Alterations Associated with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell Types. *Neuron* **98**, 1141-1154.e1147 (2018).

26.   Venkataraman, L., Fair, S.R., McElroy, C.A., Hester, M.E. & Fu, H. Modeling neurodegenerative diseases with cerebral organoids and other three-dimensional culture systems: focus on Alzheimer's disease. *Stem Cell Reviews and Reports* (2020).

27.   Choi, S.H., Kim, Y.H., Quinti, L., Tanzi, R.E. & Kim, D.Y. 3D culture models of Alzheimer's disease: a road map to a "cure-in-a-dish". *Molecular neurodegeneration* **11**, 75-75 (2016).

28.   Park, J.*, et al.* A 3D human triculture system modeling neurodegeneration and neuroinflammation in Alzheimer's disease. *Nat Neurosci* **21**, 941-951 (2018).

29.   Bowles, K.R.*, et al.* Glutamatergic dysfunction precedes neuron loss in cerebral organoids with MAPT mutation. *bioRxiv*, 2021.2002.2003.429623 (2021).

30.   Kim, Y.H.*, et al.* A 3D human neural cell culture system for modeling Alzheimer's disease. *Nat Protoc* **10**, 985-1006 (2015).

31.   Raja, W.*, et al.* Self-Organizing 3D Human Neural Tissue Derived from Induced Pluripotent Stem Cells Recapitulate Alzheimer's Disease Phenotypes. *PloS one* **11**, e0161969 (2016).

32.   Zhang, D.*, et al.* A 3D Alzheimer's disease culture model and the induction of P21-activated kinase mediated sensing in iPSC derived neurons. *Biomaterials* **35**, 1420-1428 (2014).

33.   Lee, H.K.*, et al.* Three Dimensional Human Neuro-Spheroid Model of Alzheimer's Disease Based on Differentiated Induced Pluripotent Stem Cells. *PloS one* **11**, e0163072 (2016).

34.   Jorfi, M., D'Avanzo, C., Tanzi, R.E., Kim, D.Y. & Irimia, D. Human Neurospheroid Arrays for In Vitro Studies of Alzheimer's Disease. *Scientific Reports* **8**, 2450-2450 (2018).

35.   Chen, M.*, et al.* Common proteomic profiles of induced pluripotent stem cell-derived three-dimensional neurons and brain tissue from Alzheimer patients. *Journal of Proteomics* **182**, 21-33 (2018).

36.   Cvetkovic, C., Basu, N. & Krencik, R. Synaptic Microcircuit Modeling with 3D Cocultures of Astrocytes and Neurons from Human Pluripotent Stem Cells. *Journal of Visualized Experiments* (2018).

37.   Krencik, R.*, et al.* Systematic Three-Dimensional Coculture Rapidly Recapitulates Interactions between Human Neurons and Astrocytes. *Stem cell reports* **9**, 1745-1753 (2017).

38.   Bhaduri, A.*, et al.* Cell stress in cortical organoids impairs molecular subtype specification. *Nature* **578**, 142-148 (2020).

39.   Goedert, M., Eisenberg, D.S. & Crowther, R.A. Propagation of Tau Aggregates and Neurodegeneration. *Annu Rev Neurosci* **40**, 189-210 (2017).

40.   Vogels, T.*, et al.* Propagation of tau pathology: integrating insights from post mortem and in vivo studies. *Biological Psychiatry* (2019).

41.   DeVos, S.L.*, et al.* Synaptic Tau Seeding Precedes Tau Pathology in Human Alzheimer's Disease Brain. *Front Neurosci* **12**(2018).

42.   Jucker, M. & Walker, L.C. Self-propagation of pathogenic protein aggregates in neurodegenerative diseases. *Nature* **501**, 45-51 (2013).

43.   Wu, J.W.*, et al.* Small misfolded Tau species are internalized via bulk endocytosis and anterogradely and retrogradely transported in neurons. *J Biol Chem* **288**, 1856-1870 (2013).

44.   Frost, B., Jacks, R.L. & Diamond, M.I. Propagation of tau misfolding from the outside to the inside of a cell. *J Biol Chem* **284**, 12845-12852 (2009).

45.   Guo, J.L. & Lee, V.M. Seeding of normal Tau by pathological Tau conformers drives pathogenesis of Alzheimer-like tangles. *J Biol Chem* **286**, 15317-15331 (2011).

46.   Guo, J.L. & Lee, V.M. Neurofibrillary tangle-like tau pathology induced by synthetic tau fibrils in primary neurons over-expressing mutant tau. *FEBS Lett* **587**, 717-723 (2013).

47. Clavaguera, F., Grueninger, F. & Tolnay, M. Intercellular transfer of tau aggregates and spreading of tau pathology: Implications for therapeutic strategies. *Neuropharmacology* **76**, 9-15 (2014).
48. Clavaguera, F.*, et al.* Transmission and spreading of tauopathy in transgenic mouse brain. *Nat Cell Biol* **11**, 909-913 (2009).
49. Lasagna-Reeves, C.*, et al.* Alzheimer brain-derived Tau oligomers propagate pathology from endogenous Tau. *Scientific Reports* **2**(2012).
50. Clavaguera, F.*, et al.* Brain homogenates from human tauopathies induce tau inclusions in mouse brain. *Proceedings of the National Academy of Sciences* **110**, 9535-9540 (2013).
51. Kfoury, N., Holmes, B.B., Jiang, H., Holtzman, D.M. & Diamond, M.I. Trans-cellular Propagation of Tau Aggregation by Fibrillar Species. *Journal of Biological Chemistry* **287**, 19440-19451 (2012).
52. Simón, D., García-García, E., Royo, F., Falcón-Pérez, J.M. & Avila, J. Proteostasis of tau. Tau overexpression results in its secretion via membrane vesicles. *FEBS Letters* **586**, 47-54 (2012).
53. Brettschneider, J., Del Tredici, K., Lee, V. & Trojanowski, J. Spreading of pathology in neurodegenerative diseases: A focus on human studies. *Nat. Rev. Neurosci.* **16**, 109-120 (2015).
54. Lasagna-Reeves, C.A.*, et al.* Alzheimer brain-derived tau oligomers propagate pathology from endogenous tau. *Sci Rep* **2**, 700 (2012).
55. Jiang, L.*, et al.* TIA1 regulates the generation and response to toxic tau oligomers. *Acta Neuropathologica*, 1-19 (2018).
56. Ashe, P.E.A.*, et al.* TIA1 potentiates tau phase separation and promotes generation of toxic oligomeric tau. *PNAS* **(in press)**(2020).
57. Ben Haim, L., Carrillo-de Sauvage, M.-A., Ceyzériat, K. & Escartin, C. Elusive roles for reactive astrocytes in neurodegenerative diseases. *Frontiers in cellular neuroscience* **9**(2015).
58. Chun, H. & Lee, C.J. Reactive astrocytes in Alzheimer's disease: A double-edged sword. *Neurosci Res* **126**, 44-52 (2018).
59. Leyns, C.E.G. & Holtzman, D.M. Glial contributions to neurodegeneration in tauopathies. *Molecular Neurodegeneration* **12**, 50-50 (2017).
60. Lau, S.-F., Cao, H., Fu, A.K.Y. & Ip, N.Y. Single-nucleus transcriptome analysis reveals dysregulation of angiogenic endothelial cells and neuroprotective glia in Alzheimer's disease. *Proceedings of the National Academy of Sciences* **117**, 25800 (2020).
61. Habib, N.*, et al.* Disease-associated astrocytes in Alzheimer's disease and aging. *Nat Neurosci* **23**, 701-706 (2020).
62. Mostafavi, S.*, et al.* A molecular network of the aging human brain provides insights into the pathology and cognitive decline of Alzheimer's disease. *Nat Neurosci* **21**, 811-819 (2018).
63. Apicco, D.J.*, et al.* Reducing the RNA binding protein TIA1 protects against tau-mediated neurodegeneration in vivo. *Nat Neurosci* **21**, 72-80 (2018).
64. Yoshiyama, Y.*, et al.* Synapse loss and microglial activation precede tangles in a P301S tauopathy mouse model. *Neuron* **53**, 337-351 (2007).
65. Orr, M.E., Sullivan, A.C. & Frost, B. A Brief Overview of Tauopathy: Causes, Consequences, and Therapeutic Strategies. *Trends in pharmacological sciences* **38**, 637-648 (2017).
66. Shih, R.-H., Wang, C.-Y. & Yang, C.-M. NF-kappaB Signaling Pathways in Neurological Inflammation: A Mini Review. *Front Mol Neurosci* **8**, 77-77 (2015).
67. Liberzon, A.*, et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417-425 (2015).
68. Zamanian, J.L.*, et al.* Genomic analysis of reactive astrogliosis. *J. Neurosci.* **32**, 6391-6410 (2012).
69. Grubman, A.*, et al.* A single-cell atlas of entorhinal cortex from individuals with Alzheimer's disease reveals cell-type-specific gene expression regulation. *Nat Neurosci* **22**, 2087-2097 (2019).

70. Radford, H., Moreno, J.A., Verity, N., Halliday, M. & Mallucci, G.R. PERK inhibition prevents tau-mediated neurodegeneration in a mouse model of frontotemporal dementia. *Acta neuropathologica* **130**, 633-642 (2015).

71. Mathys, H*., et al.* Single-cell transcriptomic analysis of Alzheimer's disease. *Nature* **570**, 332-337 (2019).

72. Lanneau, D., Wettstein, G., Bonniaud, P. & Garrido, C. Heat shock proteins: cell protection through protein triage. *ScientificWorldJournal* **10**, 1543-1552 (2010).

73. DeTure, M., Hicks, C. & Petrucelli, L. Targeting Heat Shock Proteins in Tauopathies. *Current Alzheimer Research* **7**, 677-684 (2010).

74. Inda, M.C*., et al.* The epichaperome is a mediator of toxic hippocampal stress and leads to protein connectivity-based dysfunction. *Nature Communications* **11**, 319 (2020).

75. Choi, S.H*., et al.* A three-dimensional human neural cell culture model of Alzheimer's disease. *Nature* **515**, 274-278 (2014).

76. Raja, W.K*., et al.* Self-Organizing 3D Human Neural Tissue Derived from Induced Pluripotent Stem Cells Recapitulate Alzheimer's Disease Phenotypes. *PloS one* **11**, e0161969-e0161969 (2016).

77. Renner, M*., et al.* Self-organized developmental patterning and differentiation in cerebral organoids. *EMBO J* **36**, 1316-1329 (2017).

78. Lin, H.C*., et al.* NGN2 induces diverse neuron types from human pluripotency. *Stem cell reports* **16**, 2118-2127 (2021).

79. Morabito, S*., et al.* Single-nucleus chromatin accessibility and transcriptomic characterization of Alzheimer's disease. *Nature genetics* **53**, 1143-1155 (2021).

80. Leng, K*., et al.* Molecular characterization of selectively vulnerable neurons in Alzheimer's Disease. (bioRxiv, 2020).

81. Wolozin, B. & Ivanov, P. Stress granules and neurodegeneration. *Nat Rev Neurosci* **20**, 649-666 (2019).

82. Meier, S*., et al.* Identification of Novel Tau Interactions with Endoplasmic Reticulum Proteins in Alzheimer's Disease Brain. *Journal of Alzheimer's disease : JAD* **48**, 687-702 (2015).

83. Meier, S*., et al.* Pathological Tau Promotes Neuronal Damage by Impairing Ribosomal Function and Decreasing Protein Synthesis. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **36**, 1001-1007 (2016).

84. Ding, Q., Markesbery, W.R., Chen, Q., Li, F. & Keller, J.N. Ribosome Dysfunction Is an Early Event in Alzheimer's Disease. *The Journal of Neuroscience* **25**, 9171 (2005).

85. Albert, B*., et al.* A ribosome assembly stress response regulates transcription to maintain proteome homeostasis. *eLife* **8**, e45002 (2019).

86. Li, J.-Q., Yu, J.-T., Jiang, T. & Tan, L. Endoplasmic Reticulum Dysfunction in Alzheimer's Disease. *Mol Neurobiol* **51**, 383-395 (2015).

87. Zhao, T., Hong, Y., Yin, P., Li, S. & Li, X.-J. Differential HspBP1 expression accounts for the greater vulnerability of neurons than astrocytes to misfolded proteins. *Proceedings of the National Academy of Sciences*, 201710549 (2017).

88. Taylor, A.R., Robinson, M.B., Gifondorwa, D.J., Tytell, M. & Milligan, C.E. Regulation of heat shock protein 70 release in astrocytes: role of signaling kinases. *Dev Neurobiol* **67**, 1815-1829 (2007).

89. San Gil, R., Ooi, L., Yerbury, J.J. & Ecroyd, H. The heat shock response in neurons and astroglia and its role in neurodegenerative diseases. *Molecular Neurodegeneration* **12**, 65 (2017).

90. Campanella, C*., et al.* Heat Shock Proteins in Alzheimer's Disease: Role and Targeting. *International journal of molecular sciences* **19**, 2603 (2018).

91. Salminen, A., Ojala, J., Kaarniranta, K., Hiltunen, M. & Soininen, H. Hsp90 regulates tau pathology through co-chaperone complexes in Alzheimer's disease. *Progress in Neurobiology* **93**, 99-110 (2011).

92. Nachman, E*., et al.* Disassembly of Tau fibrils by the human Hsp70 disaggregation machinery generates small seeding-competent species. *bioRxiv*, 2019.2012.2016.876888 (2019).

93. Mok, S.A*., et al.* Mapping interactions with the chaperone network reveals factors that protect against tau aggregation. *Nat Struct Mol Biol* **25**, 384-393 (2018).

94. Luo, W*., et al.* Roles of heat-shock protein 90 in maintaining and facilitating the neurodegenerative phenotype in tauopathies. *Proceedings of the National Academy of Sciences* **104**, 9511-9516 (2007).

95. Carnemolla, A., Lazell, H., Moussaoui, S. & Bates, G.P. In Vivo Profiling Reveals a Competent Heat Shock Response in Adult Neurons: Implications for Neurodegenerative Disorders. *PloS one* **10**, e0131985-e0131985 (2015).

96. Inda, M.C*., et al.* The epichaperome is a mediator of toxic hippocampal stress and leads to protein connectivity-based dysfunction. *Nat Commun* **11**, 319 (2020).

97. Rodina, A*., et al.* The epichaperome is an integrated chaperome network that facilitates tumour survival. *Nature* **538**, 397-401 (2016).

98. Velmeshev, D*., et al.* Single-cell genomics identifies cell type–specific molecular changes in autism. *Science* **364**, 685-689 (2019).

99. Volpicelli-Daley, L.A., Luk, K.C. & Lee, V.M. Addition of exogenous alpha-synuclein preformed fibrils to primary neuronal cultures to seed recruitment of endogenous alpha-synuclein to Lewy body and Lewy neurite-like aggregates. *Nat Protoc* **9**, 2135-2146 (2014).

100. Dobin, A*., et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

101. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420 (2018).

102. Macosko, E.Z*., et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).

103. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. arXiv:1802.03426 (2018).

104. Aran, D*., et al.* Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nature Immunology* **20**, 163-172 (2019).

105. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 289-300 (1995).

106. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*, (Springer Science & Business Media 2009).

107. Raudvere, U*., et al.* g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Research* **47**, W191-W198 (2019).

108. Subramanian, A*., et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550 (2005).

109. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847-2849 (2016).

110. Merico, D., Isserlin, R., Stueker, O., Emili, A. & Bader, G.D. Enrichment map: a network-based method for gene-set enrichment visualization and interpretation. *PloS one* **5**, e13984 (2010).

111. Shannon, P*., et al.* Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* **13**, 2498-2504 (2003).

112. Welch, J.D.*, et al.* Single-Cell Multi-omic Integration Compares and Contrasts Features of Brain Cell Identity. *Cell* **177**, 1873-1887.e1817 (2019).
113. Hao, Y.*, et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e3529 (2021).