Thermal cycling hyperthermia sensitizes non-small cell lung cancer A549 cells to EGFR-tyrosine kinase inhibitor Erlotinib

Guan-Bo Lin[1,2,*], Wei-Ting Chen[1,2,*], Yu-Yi Kuo[1,2], Hsu-Hsiang Liu[2,3], You-Ming Chen[2,3], Shr-Jeng Leu[4], Chih-Yu Chao[1,2,3]

[1]Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 106319, Taiwan

[2]Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 100233, Taiwan

[3]Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 106319, Taiwan

[4]Department of Biotechnology and Laboratory Science in Medicine, National Yang Ming Chiao Tung University, Taipei 112304, Taiwan

[*]Equal contribution

Correspondence to: Chih-Yu Chao; **email:** cychao@phys.ntu.edu.tw

Keywords: combination treatment, sensitizer, hyperthermia, non-small cell lung cancer, erlotinib

ABSTRACT

Molecular targeted therapy has emerged as a mainstream treatment for non-small

cell lung cancer (NSCLC), the most common lung cancer, which has been the leading

cause of cancer death in both genders. Erlotinib (Erl), a targeted therapy inhibiting

epidermal growth factor receptor (EGFR) pathways, has been proved to have

noticeable response rate for NSCLC cells. However, limited efficacy arises due to

intrinsic and acquired resistance among most NSCLC patients. Therefore, sensitizers

are required to potentiate the efficacy of Erl in NSCLC treatment. The study

proposed a novel thermal therapy, thermal cycling hyperthermia (TC-HT), as a

supplement to amplify the effects of Erl. We found that TC-HT remarkably reduced

the half-maximal inhibitory concentration (IC50) of Erl to as little as 0.5 μM and

demonstrated that TC-HT could sensitize A549 NSCLC cells to Erl via the

downstream of EGFR signaling cascades. Furthermore, combination treatment

induced G2/M cell cycle arrest, and inhibition of cell proliferation and migration.

Besides, via raising the high temperature of TC-HT slightly, TC-HT treatment alone

can produce excellent antineoplastic effect without hurting the normal cells. The

method is expected to be applicable to other combination therapies and may be a

starter for more sophisticated, side-effect-free anticancer treatments.

INTRODUTION

As one of the most common type of cancer, lung cancer was the largest source of cancer-related deaths in 2020, according to World Health Organization (WHO). In particular, Non-small cell lung cancer (NSCLC) accounts for 85% of lung cancer deaths and there has yet to be effective treatment for it [1]. Available therapies for NSCLC treatment include surgery, chemotherapy, radiation, and targeted therapy. As chemotherapy alone is not effective, due to low response rate, there is urgent need for combination treatment or new agents targeting one or more molecular pathways.

Targeted treatment, meant to inhibit cancer malignancies with focus on critical molecules, has been regarded as a promising therapy to treat NSCLC patients [2]. Among various targeted therapies, anti-epidermal growth factor receptor (EGFR) has been the one with most extensive application in clinical treatment of NSCLC [3]. EGFR is a receptor tyrosine kinase of the ErbB family which plays a key role in cell proliferation, survival, differentiation, and migration [4]. When activated by ligand binding, EGFR will autophosphorylate and trigger downstream signaling cascades, including mitogen-activated protein kinase (MAPK) and phosphatidylinositol 3-kinase (PI3K)-Akt pathways [5], leading to cell cycle progression and inhibiting pro-apoptotic proteins. Therefore, several EGFR-inhibiting agents, such as monoclonal antibodies and tyrosine kinase inhibitors (TKIs) have been developed for anticancer treatments [6]. Erlotinib (Erl), a first generation EGFR-TKI, has exhibited significantly better response rate and benefits in NSCLC patients than conventional platinum-based chemotherapy [7], capable of prolonging the survival of patients with advanced NSCLC after chemotherapy. Regrettably, most tumors will develop acquired resistance to Erl over time [8], plus side effects such as rash and diarrhea for long-term administration of high-dose Erl [9]. Therefore, it is worthwhile to evaluate the effect of combination treatment in potentiating the efficacy of Erl at low dose, in

place of chemotherapy drugs, such as cisplatin and pemetrexed, which have been

applied as sensitizers to enhance the efficacy of Erl [10, 11], with side effects such as

hypertension and severe diarrhea [12].

In this work, we proposed a novel thermal therapy, thermal cycling hyperthermia

(TC-HT), as a substitute for chemotherapy drugs or agents, to investigate whether

such physical cue, TC-HT, may be an effective sensitizer to amplify the anticancer

effect of Erl. Given the safety and effectiveness of TC-HT reported in our previous

studies [13, 14], we aimed to probe the effect of combining Erl and TC-HT in A549

NSCLC cell line. The results showed that TC-HT significantly enhanced the

anticancer effect of Erl by sensitizing A549 cells to Erl. As a consequence, the Erl

dosage required to achieve significant anticancer effect was greatly reduced. The

approach is expected to solve the problems of low response rate and acquired

resistance for cancer drugs in NSCLC patients, while reducing side effects

significantly. The enhanced anticancer effect was achieved in part via EGFR and

downstream pathways as shown by Western blot analysis and flow cytometry. The

results show that Erl and TC-HT synergistically weaken the phosphorylation of EGFR

and modulate the phosphorylated c-Jun N-terminal kinases (p-JNK) and Akt protein

expressions, thereby inducing the mitochondria-dependent apoptosis. Besides,

combination treatment of Erl and TC-HT regulated phosphorylated p38 (p-p38) and

cell division cycle protein 2 (Cdc2), thereby inducing G2/M cell cycle arrest.

Meanwhile, the results of the colony formation assay and wound healing assay

indicated that the combination treatment has significant inhibitory effect on cell

growth and migration. Noteworthily, TC-HT alone, to our surprise, can exhibit potent

anticancer effect by raising the high temperature setting slightly, which is comparable

to the in vitro efficacy of chemotherapy drugs. Moreover, normal lung cell line

IMR-90 was almost unharmed under the same TC-HT treatment condition.

In this paper, the study demonstrated for the first time that TC-HT can potentiate the

anticancer efficacy of Erl on A549 cells, thereby reducing their needed dosage in

treatment. Moreover, by raising the high temperature of TC-HT slightly, TC-HT

treatment could hinder growth of A549 cancer cells, without drug administration.

These findings suggest that combination of TC-HT and Erl is a promising approach in

anticancer treatment, which sheds light on combining TC-HT with other anticancer

drugs for potential treatment.

RESULTS

TC-HT for in vitro application

The thermal cycling treatment was applied with a modified PCR machine as described

previously [13, 14]. The schematic temperature and duration settings are shown in

Figure 1A, where the temperature was increased to the desired high temperature

setting and maintained for 3 min, followed by a natural cooling period for 30 sec. In

practice, the heating device was switched off in the cooling process, and the low

temperature setting was chosen to be 37˚C to mimic human body temperature. A single

cycle containing a high temperature and a low temperature period was repeated for 10

times in this study. Figure 1B showed the actual temperature in the culture well

measured by a thermocouple at moderate temperature TC-HT setting. As can be seen in

Figure 1B, the cycling temperature comes to an equilibrium state after the third heating

period, thereafter the temperature cycling between 41.5~43˚C for moderate

temperature TC-HT. In this work, we adopted the moderate temperature TC-HT

treatment (41.5~43˚C) in the following experiments to study the synergistic anticancer

effect of Erl and TC-HT in A549 cancer cells.

Figure 1. In vitro-applied TC-HT. (A) Schematic representation of the high and low

temperatures and their duration settings during TC-HT treatment. (B) Actual

temperature with moderate temperature TC-HT setting in the culture well measured

every 15 sec by a needle thermocouple.

TC-HT potentiates the anticancer efficacy of Erl in A549 NSCLC cells

To determine the effect of combination Erl and TC-HT treatment on human NSCLC

cell viability, A549 cells were treated with different concentrations of Erl with or

without TC-HT. At 72 h after the treatments, cell viability was assessed by

3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. As

shown in Figure 2A, Erl exhibited antineoplastic effect in a concentration-dependent

manner. The half-maximal inhibitory concentration (IC50) of Erl was about 10 µM in

A549 cells after 72 h treatment. Compared to the treatment with Erl alone, TC-HT

therapy sensitizes A549 cells to Erl and synergistically reduces the cell viability to

about 30% of the control group (Figure 2A). It is noteworthy that the combination of

Erl and TC-HT demonstrated synergistic ability as compared with Erl monotherapy,

which remarkably reduced the IC50 to as little as 0.5 µM (up to 20-fold decrease). To

verify the effect of this combined treatment in normal cells, the normal lung cell line

IMR-90 was used and applied with the same treatment condition. As shown in Figure

2B, the cell viability of IMR-90 normal lung cell line was almost unaffected by Erl

and/or TC-HT regardless of the treatment procedure. Additionally, as shown in Figure

2C, Erl and TC-HT combination treatment resulted in notable morphological changes

in A549 cells, including shrinkage and fragmentation of cells as well as a decreased

number of cells. The result implies that TC-HT is not only a riskless sensitizer to

natural compounds [13, 14], but also an effective and safe sensitizer to increase the

anticancer effect of targeted therapy drug.

Figure 2. Effect of Erl or TC-HT or in combination on the cell viability and

morphological changes of A549 cells. (A) MTT viability assay of A549 NSCLC

cells treated with different concentrations of Erl or in combination with moderate

temperature TC-HT treatment (41.5~43˚C). (B) MTT viability assay of IMR-90

normal lung cells treated with 5 and 10 μM Erl or in combination with moderate

temperature TC-HT treatment (41.5~43˚C). (C) Representative images of

morphological changes of A549 cells under the light microscope after treatment with

10 μM Erl, moderate temperature TC-HT treatment (41.5~43˚C) or in combination.

Scale bar = 100 μm. Data represent the mean ± standard deviation (n=3). Statistical

significance was determined by one-way analysis of variance (ANOVA) followed by

Tukey's post-hoc test (***$P < 0.001$).

Effect of Erl combined with TC-HT on EGFR protein expression in A549 cells

Here we examined the signaling pathways involved in the anticancer mechanism of

Erl and TC-HT combination treatment. With NSCLC cells frequently exhibiting

EGFR overexpression [15], targeted therapy drugs were applied to inhibit the activity.

Meant specifically for cancers with EGFR mutation, Erl inhibits cancer cells by

blocking the intracellular phosphorylation of tyrosine kinase with such mutation [7].

However, NSCLC may develop acquired resistance to Erl over time, according to

some studies [8]. In the study, as shown in Figure 3, Erl alone caused a slight decrease

in EGFR phosphorylation compared to the control cells. Moreover, the study finds

that while TC-HT alone can significantly inhibit phosphorylated EGFR (p-EGFR)

expression, the combination of Erl and TC-HT can induce more pronounced

inhibition effect on p-EGFR expression. It suggests that TC-HT may modulate the

resistance of A549 cells to Erl and alter the activity of EGFR downstream pathways.

Figure 3. Effect of Erl combined with TC-HT on p-EGFR protein expression in

A549 cells. Western blot analysis of p-EGFR protein expression in A549 cells treated

with 10 µM Erl, moderate temperature TC-HT, and the combination treatment. The

expression levels of p-EGFR were normalized to glyceraldehyde-3-phosphate

dehydrogenase (GAPDH). Each relative expression level was compared with control

and represented as fold relative to control. Data represent the mean ± standard

deviation (n=3). Differences of statistical significance were determined by one-way

ANOVA followed by Tukey's post-hoc test (**$P < 0.01$ and *$P < 0.05$).

Effect of Erl combined with TC-HT on EGFR signaling pathways in A549 cells

In the study, we then examined the expressions of JNK and Akt proteins [16] in A549

cells. JNK protein, an EGFR downstream protein belonging to the MAPK family, is

deemed to be responsive to stress stimuli and heat shock, with its phosphorylation

capable of altering the activities of many proteins that reside in mitochondria or act in

the nucleus and trigger apoptosis [17]. As shown in Figure 4A, TC-HT treatment

significantly increased the expression of p-JNK protein, compared with the control or

Erl-treated A549 cells. The result showed that TC-HT alone can effectively promote

apoptosis. Furthermore, we found that the treatment combining Erl and TC-HT is

much more effective in upregulating the p-JNK expression than TC-HT alone, as it

can boost apoptosis even more. On the other hand, another EGFR downstream protein

Akt is involved in cellular survival pathways by inhibiting apoptotic process [18].

Once activated by phosphorylation, Akt actively regulates a number of transcription

factors facilitating the expression of survival proteins. Figure 4B showed that the

expression of phosphorylated Akt (p-Akt) in Erl-treated A549 cells is lower than the

untreated cells, while the inhibitory effect on phosphorylation of Akt increases further

in TC-HT treatment and is even higher in the combination Erl and TC-HT treatment.

Meanwhile, total Akt (t-Akt) protein exhibits a trend similar to that of p-Akt protein,

suggesting that decrease in the phosphorylation of Akt in these treatments is caused

by the reduction in the amount of Akt, leading to a weakened survival signal.

Moreover, it is known that both JNK and Akt signaling pathways are important

regulators in influencing mitochondrial function. In addition, the

mitochondria-dependent apoptosis is associated with a loss in mitochondrial

membrane potential (MMP), causing the release of cytochrome c and other apoptotic

factors [19]. To confirm that the apoptosis caused by Erl and TC-HT could be related

to mitochondrial disruption, we assessed MMP with DiOC6(3) fluorescence staining

via flow cytometry. As shown in Figure S1, we confirmed that Erl and TC-HT

individually had no effect on MMP level compared to control, but the combination

treatment caused significant MMP depolarization in A549 cells, indicating

mitochondrial dysfunction in their apoptosis mechanism.

The injured mitochondria releases cytochrome c into the cytoplasm, cleaving caspase

9 and thus activating caspase 3 in the downstream [20], which enters further the

nucleus and cleaved poly ADP-ribose polymerase (PARP). It should be noted that

PARP plays an important role in mitochondria-mediated apoptosis, in addition to

being a key enzyme for DNA repair. In apoptosis, PARP protein is typically cleaved

and inactivated, thereby suppressing DNA repair and causing programmed cell death

[21]. To understand the apoptosis mechanism in A549 cells triggered by the

combination Erl and TC-HT treatment, the study evaluated the expression of

apoptosis-related proteins with Western blot analysis. Figure 4C showed that the ratio

of cleaved PARP to full length PARP in TC-HT-treated cells was much higher than

that in Erl-treated cells. Moreover, we find that the ratio of cleaved PARP to full

length PARP in combination Erl and TC-HT treatment was much stronger than that in

TC-HT treatment alone, indicating the combination treatment enhances apoptosis of

A549 cells via mitochondrial pathway. In the same vein, MutT homolog 1 (MTH1)

protein has received increasing attention in cancer treatment, due to its key role in

DNA repair [22]. High expression of MTH1 is deemed to be a sign for the metastasis

of NSCLC [23]. Therefore, the study examined MTH1 protein expression via Western

blot analysis. As shown in Figure 4D, MTH1 expression was found to decrease in

Erl-treated cells, while the inhibitive effect on MTH1 protein expression was much

higher in TC-HT treatment, which increased further in the combination Erl and

TC-HT treatment. The result reveals that the combination of Erl and TC-HT may

prevent the MTH1-related DNA repair and induce killing of cancer cells via

apoptosis.

Figure 4. Effect of Erl combined with TC-HT treatment on survival- and

apoptosis-related protein expressions in A549 cells. The anticancer experiments

were conducted in A549 cancer cells treated with 10 μM Erl, moderate temperature

TC-HT, and the combination treatment. Western blot analysis of p-JNK (A), p-Akt

and t-Akt (B), cleaved PARP and full length PARP (C), MTH1 (D) protein

expressions. The expression levels of p-JNK, p-Akt, t-Akt, and MTH1 were

normalized to GAPDH and cleaved PARP was normalized to full length PARP. Each

relative expression level was compared with control and represented as fold relative to

control. Data represent the mean ± standard deviation (n=3). Differences of statistical

significance were determined by one-way ANOVA followed by Tukey's post-hoc test

(***$P < 0.001$ and **$P < 0.01$).

The combination treatment of Erl and TC-HT causes G2/M cell cycle arrest in

A549 cells.

To further evaluate the anticancer effects of combination Erl and TC-HT treatment on

human NSCLC, the cell cycle progression in the A549 cells was examined by flow

cytometric analysis. As shown in Figures 5A and 5B, treatment with moderate TC-HT

led to a higher extent (22.95±0.56%) on the cell cycle arrest in the G2/M phase

compared to the group treated with Erl (13.06±1.16%) and the untreated cells (12.66±

1.18%). Meanwhile, the combination of Erl and moderate TC-HT caused significant

accumulation of cells in the G2/M phase (33.77±2.31%) with a concomitant reduction

of cells in the G0/G1 phase. Besides, no significant differences in the S phase were

observed among the various treatments. In the following, we examined the relevant

proteins involved in the G2/M progression to investigate the mechanism of the

combined treatment effect on cell cycle distribution. Cdc2 protein is a core regulator

in cell cycle, and Cdc2 binds to cyclin B complex guiding G2/M cell cycle transition

[24, 25]. It was reported that the inhibition of Cdc2 expression resulted in G2/M cell

cycle arrest [26-28]. In the study, as shown in Figure 5C, the expression of Cdc2

reduced significantly in combination Erl and TC-HT treatment. Meanwhile, as a

well-known MAPK member, p38 plays an important role in cell cycle regulation, and

many studies have shown that activation of p38 pathway can induce G2/M cell cycle

arrest via Cdc2 suppression in human NSCLC cells [29-31]. Consistently, we found

that the expression of p-p38 was markedly enhanced under the combination treatment

of Erl and moderate TC-HT (Figure 5D). Taken together, these results suggested that

the combined treatment could synergistically induce cell cycle arrest in the G2/M

phase by increasing the activation of p38 while reducing Cdc2 expression in cells,

ultimately inhibiting the proliferation of human lung cancer A549 cells.

Figure 5. Effect of Erl or TC-HT or in combination on the G2/M cell cycle

arrest and protein expressions of Cdc2 and p-p38 in A549 cells. The

experiments of cell cycle and protein analysis were conducted in A549 cancer cells

treated with 10 μM Erl, moderate temperature TC-HT (41.5~43˚C), and the

combination treatment. (A) Flow cytometry analysis on cellular DNA content profiles

of each group of A549 cells. (B) The proportion of A549 cells in G0/G1, S and G2/M

phases of the indicated group. Western bolt analysis of Cdc2 (C) and p-p38 (D)

protein expressions. The expression levels of p-p38 and Cdc2 were normalized to

GAPDH. Each relative expression level was compared with control and represented

as fold relative to control. Data represent the mean ± standard deviation (n=3).

Statistical significance was determined by one-way ANOVA followed by Tukey's

post-hoc test (***$P < 0.001$, **$P < 0.01$, and *$P < 0.05$).

Anti-proliferation and anti-migration effects of the combination treatment (Erl +

moderate temperature TC-HT) in A549 NSCLC cells.

Cancer mortality is associated with cancer recurrence and metastasis, both of which

are common clinical problems in NSCLC [32, 33]. Therefore, it is very important to

reduce the growth and migration ability of cancer cells. To further confirm the

anti-proliferative and anti-migration activities of Erl and TC-HT combination

treatment in NSCLC, we performed wound healing assay and colony formation assay

on A549 cells. As shown in Figures 6A and 6C, Erl-treated cells migrated over a

smaller area than control cells, while cells treated with TC-HT alone exhibited more

reduction in migration. Furthermore, the Erl and TC-HT combination treatment

caused more significant suppression of A549 cell migration compared to those of the

single treatments. In comparison to the control group, the wound closure rate in the

Erl-treated group and the TC-HT-treated group was 83.8% and 59.2%, respectively,

while the combined treatment's wound closure rate was only 35.9%. The results

indicated that the migration ability of A549 cells was significantly suppressed after

the application of Erl and TC-HT combination treatment. On the other hand, we found

that colony formation in A549 cells treated with Erl or TC-HT was decreased,

whereas the colony formation in cells treated with combination treatment was further

and greatly reduced (Figures 6B and 6D). Overall, the combination treatment has the

ability to impede the A549 cell growth and migration, indicating the potential

anticancer efficacy of Erl and TC-HT combination treatment.

Figure 6. Effect of Erl or TC-HT or in combination on the inhibition of A549 cell

colony formation and migration. (A) Wound healing assay to determine the effect of

Erl, TC-HT or combination treatment on the migration ability of A549 cells. After

scratch gaps were made, A549 cells were treated with 10 µM Erl, moderate

temperature TC-HT, and the combination treatment, respectively. (B) The results of

colony formation assay for A549 cells treated with 10 µM Erl, moderate temperature

TC-HT, and the combination treatment, respectively. (C) Wound closure rate for A549

cells was determined as the percentage of the area closed after 24 h from the initial

wound area, and each group was normalized to control group and expressed as a

fraction of 100. The areas of cell-free gaps were measured and quantified using

ImageJ. (D) Colony formation rate for A549 cells at 10 days after treatment with 10

µM Erl, moderate temperature TC-HT, and the combination treatment, respectively.

Each group was normalized to control group and represented in the form of fractions

of 100. The colony counting was performed using ImageJ. Differences of statistical

significance were determined by one-way ANOVA followed by Tukey's post-hoc test

(***$P < 0.001$, **$P < 0.01$, and *$P < 0.05$).

DISCUSSION

The study demonstrated the synergistic anticancer effect of the combination of Erl (an

EGFR-tyrosine kinase inhibitor) and TC-HT on A549 NSCLC cells. EGFR is a

receptor tyrosine kinase critical for the initiation and development of malignant

tumors via MAPK and PI3K/Akt pathways. Although the EGFR-TKIs such as Erl are

meant to target and inhibit the EGFR pathways, their clinical benefit has been limited,

due to the intrinsic and acquired resistance to TKIs among most NSCLC patients.

Amid the efforts to combat the resistance of EGFR-TKIs [34], combination treatment

has been deemed the most promising approach to overcome the resistance problem

[35]. Several studies have investigated the effect of combining Erl and other

anticancer agents such as cisplatin, monoclonal antibodies, aspirin, and capsaicin [10,

36-38], aiming to enhance the therapeutic effect. However, the combined use of

multiple drugs in clinical applications is often hampered by unpredictable drug

interactions or side effects. Therefore, we look for alternate way to enhance the

anticancer effect by combining physical stimulation with the use of drug. The study

demonstrated for the first time the effect of TC-HT physical stimulation in boosting

the anticancer effect of Erl. In comparison to Erl or TC-HT mono treatment, the

combination treatment produced a higher inhibitory effect on A549 NSCLC cells,

without hurting the normal lung cells (Figure 2), thereby circumventing drug

interaction problem and significantly reducing the needed dosage of Erl, which could

lessen side effects. To elucidate the molecular mechanisms underlying the anticancer

effect of the combination treatment on A549 NSCLC cells, the expressions of proteins

in the apoptotic pathway were evaluated. We observed that TC-HT significantly

potentiated the efficacy of the targeted therapy drug Erl in inhibiting the

phosphorylation of EGFR (Figure 3), which amplified the suppressing effect of Erl on

A549 NSCLC cells. Subsequently, we investigated the JNK and Akt proteins in the

EGFR downstream pathway. The activation of JNK is recognized to induce apoptosis

[17], while the activation of Akt serves to inhibit apoptosis [18]. Additionally, both

JNK and Akt proteins play a crucial role in the regulation of mitochondrial function

[19], and thus investigating the activation status of these proteins contributes to a

better understanding of cellular apoptotic tendency. As illustrated in Figures 4A and

4B, our results showed that the combination of TC-HT and Erl increased p-JNK

expression while decreasing p-Akt expression. Concurrently, the combination

treatment of Erl and TC-HT significantly increased MMP depolarization in A549 cells

(Figure S1). In addition, the cleavage of PARP protein and the downregulation of

MTH1 expression are both noteworthy targets for contributing apoptosis in cancer

cells [21-22]. In this study, we find that a significant increase in the expression of

cleaved PARP is accompanied by a substantial decrease in MTH1 expression (Figures

4C and 4D). These findings suggest that the combination treatment of Erl and TC-HT

effectively enhances the apoptosis of A549 NSCLC cells.

The cell cycle dysregulation, a common feature of cancer, could be ameliorated

through the modulation of cell cycle regulatory proteins, offering the potential to

attenuate the uncontrolled proliferation of cancer cells. In this study, we found that the

combination treatment of Erl and TC-HT resulted in G2/M phase arrest (Figure 5A

and 5B). Cdc2 protein is considered to be a core regulator involved in the G2/M cell

cycle transition [24, 25]. As shown in Figure 5C, significant downregulation of Cdc2

expression was found in the combination treatment group of Erl and TC-HT.

Furthermore, the p38 protein was worth investigating, as previous studies in A549

NSCLC cells demonstrated that increased p38 phosphorylation resulted in reducing

the level of Cdc2 expression, eventually leading to G2/M cell cycle arrest [29-31]. In

this study, we also found that the combined treatment of Erl and TC-HT significantly

increased the expression of p-p38. These results indicated that the combination

treatment of Erl and TC-HT effectively reduced the occurrence of mitosis in A549

NSCLC cells, representing the inhibition of cancer cell proliferation. Previous studies

have suggested that Erl alone has only a mild or even negligible effect on G2/M arrest,

while combining Erl with other approaches holds promise for improving its efficacy

[39, 40]. Similarly, our investigation also found a weak influence of Erl on G2/M

arrest when Erl was administered alone, whereas a notable enhancement of G2/M

arrest was observed when Erl was combined with TC-HT. It is worth noting that

TC-HT, as a physical stimulation, offers a new way to avoid potential drug-drug

interactions, meriting further investigation in medical applications.

Inhibiting cancer cell proliferation, recurrence, and migration has long been

considered an important part of cancer treatment, and high rates of recurrence and

metastasis are also prevalent concerns in NSCLC [32, 33]. In this study, we observed

that the combined treatment of Erl and TC-HT effectively reduced the cell viability of

A549 NSCLC cells (Figure 2), and also induced the G2/M cell cycle arrest (Figure 5).

Additionally, we investigated the proliferation and migration of A549 cancer cells

under the combined treatment of Erl and TC-HT. The results in Figures 6A and 6C

highlight significant findings concerning the migratory behavior of A549 cells under

different treatments. We find that single Erl or TC-HT treatment has a reduced

migration area in comparison to control cells. Notably, the combination treatment of

Erl and TC-HT is more effective in inhibiting A549 NSCLC cells migration than the

single Erl or TC-HT treatment. Moreover, A549 NSCLC cells treated with Erl or

TC-HT displayed reduced colony formation, whereas the combination treatment of

Erl and TC-HT exerted a more profound inhibitory effect on A549 cells (Figures 6B

and 6D) than the single treatments. Collectively, these findings illuminate the

combined treatment of Erl and TC-HT as a promising anticancer strategy,

significantly impeding both A549 NSCLC cell migration and colony formation,

suggesting its potential as a prospective anticancer treatment.

Hyperthermia has long been a promising cancer treatment method, to be applied alone

or in combination with other conventional therapies [41]. The treatment, however,

often causes cell damage, due to overdose, a practical problem which can now be

overcome by TC-HT, thanks to its ability to control thermal dosage, as its intermittent

cooling process can avoid excessive thermal dosage accumulation and subsequent

cytotoxic cell damage. Excessive thermal dosage may induce mitochondrial damage

and oxidative stress not only in cancer cells but also in normal cells [42]. However,

studies have shown that cancer cells have higher ROS levels and thus they are more

sensitive to thermal stress [43]. In this study, the results show that moderate

temperature TC-HT treatment (41.5~43°C) alone inhibited the cell viability of A549

NSCLC cells slightly, reducing their viability to 80% of the control group, without

harming IMR-90 normal lung cells (Figures 2A and 2B). To further examine the effect

of TC-HT alone on cancer cells at higher temperature, the temperature setting in

TC-HT application was raised to the range of high temperature TC-HT treatment

(42.5~45.6°C), as shown in 7A. We found that the high temperature TC-HT treatment

alone resulted in significant inhibition of the cell viability of A549 NSCLC cells, with

their viability dropping to only 17.9% of that of the control group, as shown in Figure

7B. It is noteworthy that under high temperature TC-HT treatment (42.5~45.6°C), the

additional incorporation of Erl in the range of 0-10 μM did not cause a significant

additional decrease in cell viability. Moreover, the TC-HT treatment at high

temperature (42.5~45.6°C) alone did not have much influence on the viability of

IMR-90 normal lung cells, which remained at 80% of that of the control group, even in

the case of combined treatment with 10 μM Erl (Figure 7C). The results suggest that

TC-HT is a potential anticancer treatment for NSCLC and perhaps other cancer cells,

capable of reducing the reliance on drugs, and further research to explore its efficacy is

under progress.

Figure 7. Effect of high temperature TC-HT or Erl or in combination on the cell

viability. (A) Actual temperature with high temperature TC-HT setting in the culture

well measured every 15 sec by a needle thermocouple. (B) MTT viability assay of

A549 NSCLC cells treated with different concentrations of Erl or in combination with

high temperature TC-HT treatment (42.5~45.6˚C). (C) MTT viability assay of IMR-90

normal lung cells treated with various concentrations of Erl or in combination with high

temperature TC-HT treatment (42.5~45.6˚C). Data represent the mean ± standard

deviation (n=3). Statistical significance was determined by one-way ANOVA followed

by Tukey's post-hoc test (***$P < 0.001$).

In conclusion, the study demonstrates for the first time that TC-HT, a novel thermal

treatment, is capable of sensitizing A549 cells to EGFR-TKI Erl, via the downstream

of EGFR signaling cascades, including the MAPK and PI3K-Akt pathways. Our

results show that TC-HT can enhance the anticancer efficacy of Erl on A549 cells,

thereby reducing Erl drug dosage and side effects significantly. Furthermore, TC-HT

may become a drug-free cancer treatment method via slightly raising the high

temperature of TC-HT in its application. These findings point out the potential for

TC-HT in combination therapy with other chemotherapy or targeted therapy drugs,

and further studies are needed to examine specific TC-HT parameters in treating

different types of cancers.

MATERIALS AND METHODS

Cell culture

The human NSCLC cell line A549 and normal lung cell line IMR-90 were purchased

from the Bioresource Collection and Research Center of the Food Industry Research

and Development Institute (BCRC, Hsinchu, Taiwan). Both cell lines were seeded on

cell culture flasks and cultured in DMEM (HyClone; Cytiva, Marlborough, MA, USA)

supplemented with 10% fetal bovine serum (HyClone; Cytiva) and 1%

penicillin-streptomycin (Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA)

in a humidified incubator with 5% CO_2 at 37˚C.

Drug treatment and TC-HT exposure

Erl, purchased from MedChemExpress (MCE, Monmouth Junction, NJ, USA), was

dissolved in dimethyl sulfoxide (DMSO) (Sigma-Aldrich; Merck KGaA, Darmstadt,

Germany) to a concentration of 5 mM as the stock solution and stored at -20˚C. A549

cells were seeded in 96-well plates overnight before treated with various

concentrations of Erl. The TC-HT groups were exposed to a thermal cycling treatment

with a high temperature period for 3 min and a cooling period for 30 sec, and this

protocol was repeated continuously for ten cycles, as shown in Figure 1A. The actual

temperatures sensed by the cells at the bottom of the well were measured by a needle

thermocouple. For the combination treatment, TC-HT was applied for 1 h before Erl

treatment. After the single or combination treatment, the cells were maintained in the

cell culture incubator for an additional 72 h for further analyses.

Cell viability assay and morphology

The viability of A549 cells was assessed by MTT (Sigma-Aldrich; Merck KGaA)

assay. After drug or TC-HT treatment, the medium was replaced by DMEM

containing 0.5 mg/mL MTT, and incubated for 4 h at 37˚C. The formazan crystals

were dissolved by 10% sodium dodecyl sulfate (SDS) (Bioman Scientific Co., Ltd.,

Taipei, Taiwan) overnight. Thereafter, the optical density of formazan solution was

measured at 570 nm using Multiskan GO spectrophotometer (Thermo Scientific,

Hudson, NH). Background absorbance at 690 nm was subtracted, and the final

absorbance value was expressed as a percentage of the untreated controls to represent

the cell viability. Cell morphology of A549 cells under different treatment was

observed and photographed by the Zyla 5.5 sCMOS camera (Andor, Belfast, UK).

Western blot analysis

After treatment with Erl or TC-HT or in combination, cells were washed with ice-cold

phosphate buffered saline (PBS) and lysed in RIPA lysis buffer (EMD Millipore,

Billerica, MA, USA) on ice for 30 min. Cell lysates were clarified by centrifugation at

$23,000 \times g$ for 30 min at 4˚C, and then the protein concentration in the supernatant

fraction was measured by the Bradford protein assay (Bioshop, Burlington, Canada).

Proteins (20-40 μg) were subjected to 10% SDS-PAGE and electrotransferred to

polyvinylidene fluoride (PVDF) membranes (EMD Millipore). After incubation in a

blocking buffer (5% bovine serum albumin/TBST) for 1 h at room temperature, the

membranes were immunoblotted with diluted primary antibodies at 4˚C overnight.

The specific primary antibodies against p-EGFR, PARP, p-JNK, phosphorylated Akt

(p-Akt), total Akt (t-Akt), MTH1 (Cell Signaling Technology, Inc., Danvers, MA,

USA), p-p38, Cdc2 and GAPDH (GeneTex, Inc., Irvine, CA, USA) were used. After

being washed with TBST, the membranes were incubated with horseradish

peroxidase-conjugated secondary antibodies (Jackson ImmunoResearch Laboratories,

Inc., West Grove, PA, USA). Images were visualized with an enhanced

chemiluminescence substrate (Advansta, Inc., Menlo Park, CA, USA) and quantified

by Amersham Imager 600 imaging system (AI600, GE Healthcare Life Sciences,

Chicago, IL, USA).

Mitochondrial membrane potential measurement

After treatment with Erl or TC-HT or in combination, cells were washed and

resuspended with PBS followed by staining with 20 nM 3,3’-dihexyloxacarbocyanine

iodide (DiOC$_6$(3)) (Enzo Life Sciences International Inc., Farmingdale, NY, USA) for

30 min at 37˚C in the dark. The fraction of mitochondrial depolarization in cells is

indicated by the decrease of fluorescence intensity measured by a flow cytometer

(FACSVerse; BD Biosciences, San Jose, CA, USA).

Wound healing assay

A549 cells were seeded and cultured as a confluent monolayer in 35 mm diameter

Petri dishes. In vitro, wounds were made by scratching straight lines across the cell

monolayer using a pipette tip (10 µL). Non-adherent cells and debris were removed

by gently rinsing the cells with PBS (HyClone; Cytiva). After PBS was discarded and

replaced with fresh medium, the cells were treated with Erl or TC-HT or in

combination. Each wound was observed and photographed by the Zyla 5.5 sCMOS

camera (Andor) at 0 h and 24 h after treatments. The distances between wound edges

were measured and analyzed using ImageJ software.

Colony formation assay

A549 cells were seeded at 300 cells/dish in 35 mm Petri dishes overnight for cell

adherence. Subsequently, cells were treated with Erl or TC-HT or in combination.

Cell medium was replaced 24 h after the treatment, and the cells were cultured for

additional 10 days, with fresh medium replaced every 3 days during the culture period.

At last, the cells were fixed with 4% paraformaldehyde (PFA) (Sigma-Aldrich; Merck

KGaA) for 10 min and stained with 0.1% crystal violet (Sigma-Aldrich; Merck KGaA)

to make them visible and counted. The number of colonies in each group was

normalized to control group.

Cell cycle analysis

After 24 h treatment with 10 μM Erl or TC-HT or in combination, the cells were

harvested, rinsed with PBS, and fixed with 70% ethanol at 4℃ for 30 min before

staining. The cells were then stained for 30 min in the dark with propidium iodide (PI)

and ribonuclease A (RNase A) (Gibco; Thermo Fisher Scientific, Inc.). Subsequently,

the stained cells were subject to the cell cycle analysis by using a flow cytometer

(FACSVerse; BD Biosciences), and data were analyzed with FlowJo software.

Statistical analysis

The statistical results were expressed as the average from at least three independent

experiments and the error bars are ± standard deviation. Statistical analyses were

performed by using OriginPro 2015 software (OriginLab). Differences of statistical

significance were determined by one-way ANOVA followed by Tukey's post-hoc test.

$P < 0.05$ was considered to indicate a statistically significant difference.

Abbreviations

ANOVA: analysis of variance; Cdc2: cell division cycle protein 2; DMSO: dimethyl

sulfoxide; EGFR: epidermal growth factor receptor; Erl: erlotinib; GAPDH:

glyceraldehyde-3-phosphate dehydrogenase; IC50: half-maximal inhibitory

concentration; MAPK: mitogen-activated protein kinase; MMP: mitochondrial

membrane potential; MTH1: MutT homolog 1; MTT:

3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; NSCLC: non-small

cell lung cancer; p-Akt: phosphorylated Akt; PARP: poly ADP-ribose polymerase;

PBS: phosphate buffered saline; p-EGFR: phosphorylated EGFR; PFA:

paraformaldehyde; PI: propidium iodide; PI3K: phosphatidylinositol 3-kinase; p-JNK:

phosphorylated c-Jun N-terminal kinases; p-p38: phosphorylated p38; PVDF:

polyvinylidene fluoride; RNase A: ribonuclease A; SDS: sodium dodecyl sulfate;

t-Akt: total Akt; TC-HT: thermal cycling hyperthermia; TKIs: tyrosine kinase inhibitors; WHO: World Health Organization;

AUTHOR CONTRIBUTIONS

C.Y.C. conceived and supervised the study. G.B.L., W.T.C., and C.Y.C. designed the study. G.B.L. and W.T.C., performed experiments and collected data. All authors contributed to data analysis. G.B.L., W.T.C., and C.Y.C. wrote the manuscript. All authors contributed to the article and approved the final manuscript.

ACKNOWLEDGMENTS

The authors would like to acknowledge the service provided by the Research Core Facilities 3 Laboratory of the Department of Medical Research at National Taiwan University hospital for use of flow cytometry system.

CONFLICTS OF INTEREST

The authors have declared that no competing interests exist.

FUNDING

This work was supported by research grants from Ministry of Science and Technology (NSTC 112-2112-M-002-033, MOST 110-2112-M-002-004, and MOST 109-2112-M-002-004 to C.Y.C.) of the Republic of China. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

REFERENCES

1.   Belani C, Goss G, Blumenschein Jr G. Recent clinical developments and
rationale for combining targeted agents in non-small cell lung cancer (NSCLC).
Cancer Treat Rev. 2012; 38:174-84.
https://doi.org/10.1016/j.ctrv.2011.05.009
PMID: 21715100

2.   Imyanitov EN, Iyevleva AG, Levchenko EV. Molecular testing and targeted
therapy for non-small cell lung cancer: current status and perspectives. Crit Rev
Oncol Hemat. 2021; 157:103194.
https://doi.org/10.1016/j.critrevonc.2020.103194
PMID: 33316418

3.   Wieduwilt MJ, Moasser MM. The epidermal growth factor receptor family:
biology driving targeted therapeutics.   Cell Mol Life Sci. 2008; 65:1566-84.
https://doi.org/10.1007/s00018-008-7440-8
PMID: 18259690

4.   Yewale C, Baradia D, Vhora I, Patil S, Misra A. Epidermal growth factor
receptor targeting in cancer: a review of trends and strategies. Biomaterials. 2013;
34:8690-707.
https://doi.org/10.1016/j.biomaterials.2013.07.100
PMID: 23953842

5.   Wee P, Wang Z. Epidermal growth factor receptor cell proliferation signaling
pathways. Cancers. 2017; 9:52.
https://doi.org/10.3390/cancers9050052
PMID: 28513565

6.   Ciardiello F, De Vita F, Orditura M, Tortora G. The role of EGFR inhibitors in
nonsmall cell lung cancer. Curr Opin Oncol. 2004; 16:130-5.
https://doi.org/10.1097/00001622-200403000-00008
PMID: 15075904

7.   Gridelli C, Maione P, Bareschino MA, Schettino C, Sacco PC, Ambrosio R, et al.
Erlotinib in the treatment of non-small cell lung cancer: current status and future
developments. Anticancer Res. 2010; 30:1301-10.
https://ar.iiarjournals.org/content/30/4/1301
PMID: 20530444

8.   Lin Y, Wang X, Jin H. EGFR-TKI resistance in NSCLC patients: mechanisms
and strategies. Am J Cancer Res. 2014; 4:411-35.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4163608/#
PMID: 25232485

9.   Melosky B. Supportive care treatments for toxicities of anti-EGFR and other
targeted agents. Curr Oncol. 2012; 19:59-63.

https://doi.org/10.3747/co.19.1054

PMID: 22787412

10. Raimbourg J, Joalland MP, Cabart M, de Plater L, Bouquet F, Savina A,

Decaudin D, Bennouna J, Vallette FM, Lalier L. Sensitization of EGFR

wild-type non–small cell lung cancer cells to EGFR-tyrosine kinase inhibitor

erlotinib. Mol Cancer Ther. 2017; 16:1634-44.

https://doi.org/10.1158/1535-7163.MCT-17-0075

PMID: 28522592

11. Li T, Ling YH, Goldman ID, Perez-Soler R. Schedule-dependent cytotoxic

synergism of pemetrexed and erlotinib in human non-small cell lung cancer cells.

Clin Cancer Res. 2007; 13:3413-22.

https://doi.org/10.1158/1078-0432.CCR-06-2923

PMID: 17545550

12. Almanric K, Marceau N, Cantin A, Bertin É. Risk factors for nephrotoxicity

associated with cisplatin. Can J Hosp Pharm. 2017; 70:99-106.

https://doi.org/10.4212/cjhp.v70i2.1641

PMID: 28487576

13. Chen WT, Sun YK, Lu CH, Chao CY. Thermal cycling as a novel thermal

therapy to synergistically enhance the anticancer effect of propolis on PANC-1

cells. Int J Oncol. 2019; 55:617-28.

https://doi.org/10.3892/ijo.2019.4844

PMID: 31322205

14. Lu CH, Chen WT, Hsieh CH, Kuo YY, Chao CY. Thermal cycling-hyperthermia

in combination with polyphenols, epigallocatechin gallate and chlorogenic acid,

exerts synergistic anticancer effect against human pancreatic cancer PANC-1

cells. PLoS One. 2019; 14:e0217676.

https://doi.org/10.1371/journal.pone.0217676

PMID: 31150487

15. Gaber R, Watermann I, Kugler C, Reinmuth N, Huber RM, Schnabel PA,

Vollmer E, Reck M, Goldmann T. Correlation of EGFR expression, gene copy

number and clinicopathological status in NSCLC. Diagn Pathol. 2014; 9:165.

https://doi.org/10.1186/s13000-014-0165-0

PMID: 25227424

16. Atalay G, Cardoso F, Awada A, Piccart MJ. Novel therapeutic strategies targeting

the epidermal growth factor receptor (EGFR) family and its downstream

effectors in breast cancer. Ann Oncol. 2003; 14:1346-63.

https://doi.org/10.1093/annonc/mdg365

PMID: 12954573

17.   Takeuchi K, Ito F. EGF receptor in relation to tumor development: molecular

basis of responsiveness of cancer cells to EGFR-targeting tyrosine kinase

inhibitors. FEBS J. 2010; 277:316-26.

https://doi.org/10.1111/j.1742-4658.2009.07450.x

PMID: 19922467

18.   Akca H, Tani M, Hishida T, Matsumoto S, Yokota J. Activation of the AKT and

STAT3 pathways and prolonged survival by a mutant EGFR in human lung

cancer cells. Lung Cancer. 2006; 54:25-33.

https://doi.org/10.1016/j.lungcan.2006.06.007

PMID: 16872715

19.   Matsuyama S, Reed JC. Mitochondria-dependent apoptosis and cellular pH

regulation. Cell Death Differ. 2000; 7:1155-65.

https://doi.org/10.1038/sj.cdd.4400779

PMID: 11175252

20.   Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E, Boise LH.

Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis.

BMC Cell Biol. 2013; 14:32.

https://doi.org/10.1186/1471-2121-14-32

PMID: 23834359

21.   Yi M, Dong B, Qin S, Chu Q, Wu K, Luo S. Advances and perspectives of PARP

inhibitors. Exp Hematol Oncol 2019; 8:44573.

https://doi.org/10.1186/s40164-019-0154-9

PMID: 31737426

22.   Gad H, Koolmeister T, Jemth AS, Eshtad S, Jacques SA, Ström CE, Svensson

LM, Schultz N, Lundbäck T, Einarsdottir BO, Saleh A, Göktürk C, Baranczewski

P, et al. MTH1 inhibition eradicates cancer by preventing sanitation of the dNTP

pool. Nature. 2014; 508:215-21.

https://doi.org/10.1038/nature13181

PMID: 24695224

23.   Li DN, Yang CC, Li J, Ou Yang QG, Zeng LT, Fan GQ, Liu TH, Tian XY, Wang

JJ, Zhang H, Dai DP, Cui J, Cai JP. The high expression of MTH1 and NUDT5

promotes tumor metastasis and indicates a poor prognosis in patients with

non-small-cell lung cancer. Biochim Biophys Acta-Mol Cell Res. 2021;

1868:118895.

https://doi.org/10.1016/j.bbamcr.2020.118895

PMID: 33096144

24.   Dorée M, Hunt T. From Cdc2 to Cdk1: when did the cell cycle kinase join its

cyclin partner? J Cell Sci. 2002; 115:2461-4.

https://doi.org/10.1242/jcs.115.12.2461

PMID: 12045216

25. Lim S, Kaldis P. Cdks, cyclins and CKIs: roles beyond cell cycle regulation.

Development. 2013; 140:3079-93.

https://doi.org/10.1242/dev.091744

PMID: 23861057

26. Chang CC, Heller JD, Kuo J, Huang RC. Tetra-O-methyl nordihydroguaiaretic

acid induces growth arrest and cellular apoptosis by inhibiting Cdc2 and survivin

expression. Proc Natl Acad Sci U S A. 2004; 101:13239-44.

https://doi.org/10.1073/pnas.040540710

PMID: 15329416

27. Senju M, Sueoka N, Sato A, Iwanaga K, Sakao Y, Tomimitsu S, Tominaga M,

Irie K, Hayashi S, Sueoka E. Hsp90 inhibitors cause G2/M arrest associated with

the reduction of Cdc25C and Cdc2 in lung cancer cell lines. J Cancer Res Clin

Oncol. 2006; 132:150-8.

https://doi.org/10.1007/s00432-005-0047-7

PMID: 16283383

28. Yoshida M, Matsui Y, Iizuka A, Ikarashi Y. G2-phase arrest through p21(WAF1 /

Cip1) induction and cdc2 repression by gnidimacrin in human hepatoma HLE

cells. Anticancer Res. 2009; 29:1349-54.

https://ar.iiarjournals.org/content/29/4/1349.long

PMID: 19414386

29. Su JC, Lin KL, Chien CM, Lu CM, Chen YL, Chang LS, Lin SR. Novel

indoloquinoline derivative, IQDMA, induces G(2)/M phase arrest and apoptosis

in A549 cells through JNK/p38 MAPK signaling activation. Life Sci. 2009;

85:505-16.

https://doi.org/10.1016/j.lfs.2009.08.006

PMID: 19699753

30. Pai JT, Hsu MW, Leu YL, Chang KT, Weng MS. Induction of G2/M Cell Cycle

Arrest via p38/p21Waf1/Cip1-Dependent Signaling Pathway Activation by

Bavachinin in Non-Small-Cell Lung Cancer Cells. Molecules. 2021; 26:5161.

https://doi.org/10.3390/molecules26175161

PMID: 34500594

31. Luo YH, Wang C, Xu WT, Zhang Y, Zhang T, Xue H, Li YN, Fu ZR, Wang Y,

Jin CH. 18β-Glycyrrhetinic Acid Has Anti-Cancer Effects via Inducing

Apoptosis and G2/M Cell Cycle Arrest, and Inhibiting Migration of A549 Lung

Cancer Cells. Onco Targets Ther. 2021; 14:5131-44.

https://doi.org/10.2147/OTT.S322852

PMID: 34712051

32. Tamura T, Kurishima K, Nakazawa K, Kagohashi K, Ishikawa H, Satoh H,
Hizawa N. Specific organ metastases and survival in metastatic non-small-cell
lung cancer. Mol Clin Oncol. 2015; 3:217-21.
https://doi.org/10.3892/mco.2014.410
PMID: 25469298

33. Uramoto H, Tanaka F. Recurrence after surgery in patients with NSCLC. Transl
Lung Cancer Res 2014; 4:242.
https://doi.org/10.3978/j.issn.2218-6751.2013.12.05
PMID: 25806307

34. Passaro A, Jänne PA, Mok T, Peters S. Overcoming therapy resistance in
EGFR-mutant lung cancer. Nat Cancer. 2021; 2:377-91.
https://doi.org/10.1038/s43018-021-00195-8
PMID: 35122001

35. Tong CW, Wu WK, Loong HH, Cho WC, To KK. Drug combination approach to
overcome resistance to EGFR tyrosine kinase inhibitors in lung cancer. Cancer
Lett. 2017; 405:100-10.
https://doi.org/10.1016/j.canlet.2017.07.023
PMID: 28774798

36. Cavazzoni A, Alfieri RR, Cretella D, Saccani F, Ampollini L, Galetti M, Quaini F,
Graiani G, Madeddu D, Mozzoni P, Galvani E, La Monica S, Bonelli M, et al.
Combined use of anti-ErbB monoclonal antibodies and erlotinib enhances
antibody-dependent cellular cytotoxicity of wild-type erlotinib-sensitive NSCLC
cell lines. Mol Cancer. 2012; 11:91.
https://doi.org/10.1186/1476-4598-11-91
PMID: 23234355

37. Hu X, Wu LW, Weng X, Lin NM, Zhang C. Synergistic antitumor activity of
aspirin and erlotinib: Inhibition of p38 enhanced aspirin plus erlotinib-induced
suppression of metastasis and promoted cancer cell apoptosis. Oncol Lett. 2018;
16:2715-24.
https://doi.org/10.3892/ol.2018.8956
PMID: 30013667

38. Chen JC, Ko JC, Yen TC, Chen TY, Lin YC, Ma PF, Lin YW. Capsaicin
enhances erlotinib-induced cytotoxicity via AKT inactivation and excision repair
cross-complementary 1 (ERCC1) down-regulation in human lung cancer cells.
Toxicol Res 2019; 8:459-70.
https://doi.org/10.1039/c8tx00346g
PMID: 31160978

39. Steen NV, Potze L, Giovannetti E, Cavazzoni A, Ruijtenbeek R, Rolfo C, et al.
Molecular mechanism underlying the pharmacological interactions of the protein
kinase C-β inhibitor enzastaurin and erlotinib in non-small cell lung cancer cells.
Am J Cancer Res. 2017; 7:816-30.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5411790/
PMID: 28469955
40. Cheng F, Peng X, Meng G, Pu Y, Luo K, He B. Poly(ester-thioether)
microspheres co-loaded with erlotinib and α-tocopheryl succinate for
combinational therapy of non-small cell lung cancer. J Mater Chem B. 2020;
8:1728-38.
https://doi.org/10.1039/c9tb02840d
PMID: 32022097
41. Wust P, Hildebrandt B, Sreenivasa G, Rau B, Gellermann J, Riess H, Felix R,
Schlag PM. Hyperthermia in combined treatment of cancer. Lancet Oncol. 2002;
3:487-97.
https://doi.org/10.1016/S1470-2045(02)00818-5
PMID: 12147435
42. Belhadj Slimen I, Najar T, Ghram A, Dabbebi H, Ben Mrad M, Abdrabbah M.
Reactive oxygen species, heat stress and oxidative-induced mitochondrial
damage. A review.   Int J Hyperthermia. 2014; 30:513-23.
https://doi.org/10.3109/02656736.2014.971446
PMID: 25354680
43. Panieri E, Santoro M. ROS homeostasis and metabolism: a dangerous liason in
cancer cells. Cell Death Dis. 2016; 7:e2253.
https://doi.org/10.1038/cddis.2016.105
PMID: 27277675
803
804

805 **SUPPLEMENTARY MATERIALS**
806
807 **Supplementary Figure**
808

Supplementary Figure S1. Flow cytometric analysis of MMP in A549 cells

treated with Erl and/or TC-HT. A549 cells were treated with 10 μM Erl, moderate

temperature TC-HT, and the combination of both treatments. (A) MMP was evaluated

by the $DiOC_6(3)$ fluorescent dye and detected by flow cytometer. (B) Quantification

of the cells with decreased MMP was measured. Data represent the mean ± standard

deviation (n=3). Statistical significance was determined by one-way ANOVA

followed by Tukey's post-hoc test (**$P < 0.01$)

Figure 1

A

B

Figure 2

A

A549

B

IMR-90

C

Ctrl

Erlotinib

TC-HT

Erlotinib + TC-HT

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1