An actomyosin-polarized membrane reservoir mediates the unequal divisions of *Drosophila* neural stem cells

Bryce LaFoya and Kenneth E. Prehoda

Institute of Molecular Biology
Department of Chemistry and Biochemistry
1229 University of Oregon
Eugene, OR 97403

Corresponding author: prehoda@uoregon.edu

Summary

The asymmetric divisions of *Drosophila* neural stem cells (NSCs) produce unequally sized

siblings, with most volume directed into the sibling that retains the NSC fate. Sibling size

asymmetry results from preferential expansion of the NSC sibling surface during division. Here

we show that a polarized membrane reservoir constructed by the NSC in early mitosis provides

the source for expansion. The reservoir is formed from membrane domains that contain folds

and microvilli that become polarized by apically-directed cortical flows of actomyosin early in

mitosis. When furrow ingression begins and internal pressure increases, the stores of

membrane within the apical reservoir are rapidly consumed. Expansion is significantly

diminished in NSCs that lack a reservoir, and membrane expansion equalizes when the

reservoir is not polarized. Our results suggest that the cortical flows that remodel the plasma

membrane during asymmetric cell division function to satisfy the dynamic surface area

requirements of unequally dividing cells.

Introduction

Cellular resources can be partitioned unequally during division to facilitate differences in sibling

cell fate and proliferation potential. For example, dividing *Drosophila* neural stem cells (NSCs)

direct most cell volume into the sibling that retains the NSC fate and remains highly

proliferative. In contrast, the neural precursor (NP) sibling that is destined towards

differentiation and a postmitotic state receives a small fraction of the total volume (Gallaud et

al., 2017; Homem and Knoblich, 2012). While asymmetric spindle positioning can contribute to

unequal cell division, the size asymmetry of the NSC and NP cells results from preferential

expansion of the NSC sibling during anaphase (Cabernard et al., 2010; Connell et al., 2011;

Pham et al., 2019; Roubinet et al., 2017). Biased expansion is characterized by a rapid increase

in surface area near the NSC pole while the area near the other pole remains relatively constant

(Connell et al., 2011). Expansion occurs when intracellular hydrostatic pressure increases

following the initiation of furrow ingression (Pham et al., 2019). While a biased increase in

surface area is a key aspect of the unequal NSC division, little is known about how the plasma

membrane accommodates asymmetric polar expansion. Here we examine the plasma

membrane dynamics that support the geometric changes that accompany the unequal NSC

division.

Cell shape changes can significantly alter surface area, placing significant demands on the

plasma membrane. Many animal cells, including the *Drosophila* NSC, round up during division

to assume a geometry that uses surface area most efficiently for a given volume. The surface

area can increase when the NSC deviates from sphericity as it elongates and expands, like a

pinched balloon. However, the plasma membrane is relatively inelastic and can only undergo

limited stretching to accommodate increases in surface area. One mechanism that cells use to

buffer changes in surface area is to store membrane in cell surface reservoirs (Figard et al.,

2013; Figard and Sokac, 2014; Kabeche et al., 2015). Membrane reservoirs use folds and

microvilli to store membrane that is unnecessary for encapsulating the cell (i.e. excess

membrane). When shape changes decrease surface area, excess membrane can be rapidly

stored in these structures. Conversely, increases in surface area can potentially be rapidly

accommodated by tapping into the reservoir, reducing the amount of excess membrane.

The NSC is not known to store membrane in reservoirs. However, the NSC membrane

undergoes a polarity cycle that resembles its well-characterized protein polarity cycle (LaFoya

and Prehoda, 2021). During the protein cycle, polarity proteins and fate determinants become

polarized to the NSC's apical and basal hemispheres during early mitosis (Gallaud et al., 2017;

Homem and Knoblich, 2012; Knoblich, 2010, 2008). The apical proteins are polarized in part by

apically-directed cortical flows of actomyosin. These cortical movements lead to the formation

of an organized cap near the apical pole that remains until it depolarizes shortly before division

(Oon and Prehoda, 2021, 2019). The plasma membrane undergoes a similar cycle in which

membrane features that are initially dispersed over the surface of the cell move apically, in

synchrony with polarity proteins, before appearing to spread over the surface by moving

basally during anaphase. The dynamic NSC membrane features can be detected with diverse

membrane sensors and some are clearly microvilli suggesting that they could store excess

membrane. Here we examine whether the NSC membrane contains a reservoir and if so, how

membrane dynamics might contribute to the biased expansion that mediates unequal NSC

division.

Results

The apparent surface area of the apical NSC sibling increases during division

To estimate the potential demand for plasma membrane during NSC division, we measured the

expansion of cellular surface area from metaphase to the completion of cytokinesis in

Drosophila larval brain NSCs (Figure 1A,B). We compared the *apparent* total surface area at

both time points, where the apparent surface area refers to that of the main cell body exclusive

of surface area contained in folds or microvilli (Figure 1C). We measured an average surface

area increase of 63 ± 22 µm^2 (n = 10), which was approximately equal to half the apparent

surface area of the NP cell on average (118 ± 12 µm^2; n = 10) (Figure 1D). Expansion during

NSC division is biased, such that most expansion occurs near the apical pole (Connell et al.,

2011; Pham et al., 2019; Roubinet et al., 2017). We conclude that the NSC division creates a

demand for new surface area that is apparently satisfied by a biased expansion process that

directs the increase in surface area towards the apical pole.

Neural stem cell membrane domains store excess membrane

We sought to understand how the plasma membrane supports the increase in apparent

surface area of the NSC's apical region during division. The NSC membrane is heterogeneous,

with distinct domains characterized by localized areas of relatively high membrane sensor

intensity (Figure 1E) (LaFoya and Prehoda, 2021). NSC membrane domains are distributed

across the cell surface in interphase and early mitosis and can be detected with diverse

membrane sensors. The higher sensor intensity within NSC membrane domains could arise

from an increased density of specific membrane components in a single bilayer (Figure 1F-

scheme 1). Alternately, however, increased amounts of plasma membrane in the form of folds

or microvilli could also lead to higher sensor intensity (Figure 1F-scheme 2), and some microvilli

are known to be present (LaFoya and Prehoda, 2021). We term the fine structure of the

membrane in folds and microvilli, "excess membrane", as it extends away from the cell surface

and therefore appears to be unnecessary for encapsulating the NSC. We examined the

prevalence of excess membrane by measuring the thickness of domains in early mitosis when

the NSC is rounded up using super resolution imaging (Azuma and Kei, 2015). NSC membrane

domains were nearly exclusively associated with increased thickness relative to surrounding

areas, which had a relatively uniform thickness approximately equal to the diffraction limited

width (Figure 1G). We conclude that excess membrane is stored on the NSC surface in

discrete regions that are thicker than surrounding regions and often contain microvilli. The

presence of excess membrane makes the total NSC membrane surface area greater than the

apparent surface area of the cell body, a phenomenon that has been compared to the

"coastline paradox" in which the size of an interface is influenced by the presence of structure

at different length scales (Sokac, 2017).

Excess plasma membrane moves apically to form a polarized membrane reservoir

NSC membrane domains that are dispersed over the cell surface in early mitosis become

concentrated around the apical pole by metaphase (Figure 2A and Video S1) (LaFoya and

Prehoda, 2021). Given our finding that membrane domains store excess membrane, we

hypothesized that the polarization process causes excess membrane to accumulate near the

apical pole. Consistent with this hypothesis, we measured a significant increase in apical

membrane thickness at metaphase compared to prophase (Figure 2B,C). We did not detect

any basally-directed movement of excess membrane during this time period. We also

observed deformation of domains along the apical-basal axis as they moved towards the

apical pole (c.f. Figure 2A inset, Video S1), suggesting the presence of tension-induced

stretching. We conclude that the polarization of the NSC membrane occurs at least in part from

accumulation of excess membrane around the apical pole due to its directional transport from

around the cell surface.

Cortical actomyosin flows drive apical plasma membrane accumulation

The apically-directed membrane domain movements that construct the polarized membrane

reservoir do not occur in Latrunculin A-treated NSCs (LaFoya and Prehoda, 2021), indicating

that they require F-actin. The requirement of F-actin, along with the recent discovery of

apically-directed cortical flows of actomyosin in prophase NSCs (Oon and Prehoda, 2021),

suggest that cortical flows could drive the formation of the apical membrane reservoir. We

examined whether cortical cytoskeletal flows are correlated with membrane polarization

dynamics using simultaneous imaging of the membrane and F-actin. As previously reported

(LaFoya and Prehoda, 2021), we observed enrichment of F-actin at sites of excess membrane

(Figure 3 and Video S2). Here we find that the movements that polarize membrane domains are

highly correlated with cortical F-actin dynamics (Figure 3B and Video S2). Cortical F-actin has

been reported to become enriched near the apical pole at metaphase (Oon and Prehoda, 2021;

Roubinet et al., 2017), and our results suggest that F-actin enrichment is the result of its

presence in the apically-polarized membrane domains. The correlated motions of actin and

membrane, along with the requirement of F-actin for these movements, suggest that cortical

flows of actin are responsible for constructing the apical membrane reservoir.

The apical membrane reservoir is consumed during NSC division

Our data indicate that the NSC constructs a polarized membrane reservoir in early mitosis by

moving excess membrane to the apical pole using actomyosin driven cortical flows. We

examined whether the apical membrane reservoir is consumed during anaphase, when the

apical region's apparent surface area increases (Connell et al., 2011; Pham et al., 2019). The

internal pressure of the NSC changes dramatically during this phase of the cell cycle, peaking

shortly after furrow ingression followed by a rapid decrease that coincides with membrane

expansion (Pham et al., 2019). We followed the dynamics of the apical membrane reservoir

during anaphase using super resolution imaging with high temporal resolution. We observed

thinning of the apical membrane that coincided with the broad equatorial narrowing associated

with the start of cytokinesis and the increase in apical surface area (Figure 4 and Video S3, S4;

reservoir formation and consumption are shown in Video S4). Microvilli that extended over a

micron from the cell surface were reabsorbed into the cell as furrowing continued (Figure 4A-

inset and Video S3, S4). The reservoir was consumed locally in smaller domains rather than

uniformly across the apical surface, suggesting that mechanical failure points may exist that

begin to unfold once the pressure reaches a certain threshold. As membrane reservoir

consumption occurred, the apical surface expanded and the remaining domains of excess

membrane appeared to move basally, as previously reported (LaFoya and Prehoda, 2021). As

apical expansion occurred, F-actin formed puncta that spread over the surface (Figure 5 and

Video S5). Thus, NSCs construct a polarized membrane reservoir in early metaphase that is

consumed during furrow ingression as the apical membrane expands.

The membrane reservoir is required for expansion and completion of cytokinesis

Our data indicate that the apical membrane reservoir is consumed during anaphase,

suggesting that the reservoir mediates preferential expansion at the apical pole. To investigate

how the NSC division process proceeds without a reservoir, we examined the divisions of

cyclodextrin-treated NSCs. Cyclodextrin treatment can ablate membrane domains in diverse

cell types (Barman and Nayak, 2007; Pang et al., 2007), perhaps because it extracts

cholesterol, significantly decreasing membrane rigidity (Chakraborty et al., 2020). In previous

work, we found that cyclodextrin treatment inhibits formation of NSC membrane features

(LaFoya and Prehoda, 2021). Consistent with previous observations, we found that the

membranes of cyclodextrin-treated NSCs were relatively featureless such that the plasma

membrane smoothly followed the contours of the cell (Figure 6A), apparently lacking any

excess membrane. In the absence of a membrane reservoir, very little polar expansion

occurred, and furrowing was incomplete and unstable (Figure 6B,C and Video S6). These

results suggest that expansion requires a membrane reservoir. Furthermore, reservoir-

mediated expansion may provide a "pressure relief valve" that prevents the intracellular

pressure from continuing to increase as the furrow ingresses. The increase in intracellular

hydrostatic pressure that coincides with the onset of furrow ingression (Pham et al., 2019), may

be released by membrane reservoir-mediated expansion and without a reservoir, intracellular

pressure may increase to the point where the furrow is no longer able to ingress.

Membrane reservoir polarization is required for biased expansion

Our data suggest that membrane reservoir polarization mediates biased expansion during NSC

division. We examined how failure to polarize the membrane reservoir influences expansion

using NSCs expressing RNAi directed against the heterotrimeric G-protein Gγ1 (hereafter Gγ).

Gγ is known to be required for asymmetric NSC division (Izumi et al., 2004), and we found that

NSCs expressing Gγ RNAi do not polarize their membranes in early mitosis (Figure 7 and Video

S7). Like wild-type NSCs, cells expressing Gγ RNAi had excess membrane distributed over the

cell surface in early mitosis. However, the apically-directed movements that polarize the

membrane reservoir of wild-type NSCs failed to occur in Gγ RNAi-expressing cells. Without

these polarizing movements, these cells failed to form a membrane reservoir at the apical pole

as excess membrane remained distributed across the cell surface as furrow ingression

commenced (Figure 7B,C). In the absence of a polarized membrane reservoir, the expansion

that occurred following ingression was equally distributed to both poles and daughter cell size

was approximately symmetric. We conclude that the biased expansion of the NSC membrane

is mediated by polarization of the membrane reservoir before division.

Discussion

The morphological changes that accompany cell division can place significant demands on the

plasma membrane by rapidly increasing total surface area. During the unequal divisions of

Drosophila NSCs, the surface area increase is biased, with the nascent NSC surface expanding

as the furrow ingresses while the NP surface remains relatively constant (Connell et al., 2011;

Pham et al., 2019). In general, the demand for membrane can be buffered by reservoirs of

excess membrane stored in folds and microvilli, or eisosomes in yeast (Figard et al., 2016;

Figard and Sokac, 2014; Goudarzi et al., 2017; Kabeche et al., 2015). We discovered a

membrane reservoir that mediates the biased expansion that occurs during NSC asymmetric

division (Figure 7D). The reservoir is constructed from localized regions of excess membrane

that are dynamically collected at the apical pole by apically-directed cortical flows of

actomyosin that begin and end in early mitosis (Oon and Prehoda, 2021). When expansion

begins following the initiation of cleavage furrow ingression, the excess membrane in the

reservoir is consumed, coinciding with a rapid drop in internal hydrostatic pressure that occurs

during anaphase (Pham et al., 2019). In addition to explaining the source of membrane for

biased NSC expansion, these results have implications for the function of the complex cortical
actomyosin dynamic phases that occur during NSC asymmetric division. Furthermore, they
raise questions about how cortical actomyosin dynamics are coupled to the organization of the
NSC membrane, and how intracellular pressure and actomyosin dynamics are coupled to
expansion-induced tapping of the reservoir. Polarized membrane structures have also been
reported in the asymmetrically-dividing *C. elegans* zygote (Hirani et al., 2019; Scholze et al.,
2018), suggesting that they may play a more general role in asymmetric division.

Examination of NSC membrane dynamics revealed that the cell goes to great lengths to
prepare for biased expansion, generating directional cortical flows to collect excess membrane
at the apical pole (Figure 2). Directional cortical flows of actomyosin play a role in the
polarization of both the plasma membrane and proteins during the NSC division (LaFoya and
Prehoda, 2021; Oon and Prehoda, 2021, 2019), and their role in cortical protein polarity of the
C. elegans zygote has been extensively studied (Illukkumbura et al., 2019; Lang and Munro,
2017; Munro et al., 2004). While cortical movements of polarity proteins such as aPKC are
driven by actomyosin cortical flows in the NSC, F-actin is not strictly required for its polarity.
For example, in Latrunculin A treated NSCs that lack an actin cytoskeleton, aPKC polarizes but
ultimately diffuses into the basal hemisphere (Oon and Prehoda, 2019). Thus, currently
available data suggest that actomyosin cortical flows in the NSC function primarily to organize
the membrane and are not essential for establishing protein polarity, even while the cortical
cytoskeleton maintains protein polarity by preventing spreading into the basal hemisphere
(LaFoya and Prehoda, 2021; Oon and Prehoda, 2019).

Localized regions of excess membrane move to the apical pole during prophase to generate
the NSC membrane reservoir. How might actomyosin-generated cortical flows be coupled to
the directional movement of excess membrane domains? In cultured mammalian epithelial
cells, microvilli move rapidly across the cell surface to form the brush border (Meenderink et
al., 2019). In this context, microvillar motion has been proposed to be driven by treadmilling of
an interior actin bundle. In the NSC, apically-directed cortical flows of actomyosin not only
move excess membrane but deform the surrounding tissue (LaFoya and Prehoda, 2021; Oon
and Prehoda, 2021), suggesting that the flows are coupled to a broad range of membrane-
associated structures, including adhesions. The deformations of excess membrane domains
along the apical-basal axis suggest that tension in the membrane is generated as the domains
are gathered at the apical pole.

The apical reservoir is tapped shortly after its creation in metaphase (Figure 4), coinciding with

the onset of furrow ingression and the peak of internal hydrostatic pressure (Pham et al., 2019).

Simultaneously with these processes, myosin II redistributes from the apical pole towards the

cleavage furrow (Oon and Prehoda, 2021; Pham et al., 2019; Roubinet et al., 2017). Reservoir

consumption could be caused by myosin relocalization or the initiation of furrow ingression, or

a combination of both processes. Alternately redistribution of myosin could be a mechanical

consequence of reservoir tapping. Future work will be directed at understanding the physical

mechanisms that couple cortical actomyosin dynamics to membrane movements in the NSC,

and the role these dynamics play in coupling reservoir consumption and cell surface

expansion.

Acknowledgments

This work was supported by NIH grants GM127092 and F32GM134705.

Star Methods

Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed to the Lead

Contact, Kenneth Prehoda (prehoda@uoregon.edu).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

The raw data supporting the current study have not been deposited in a public repository

because of their large file size, but are available from the corresponding author on request.

Experimental Model and Subject Details

Fly Strains

For tissue specific expression of UAS controlled transgenes in NSCs, a Worniu-Gal4 driver line

was used. For live imaging of membrane dynamics, two different classes of membrane-bound

fusion proteins were used, both of which nearly exclusively label the plasma membrane and

not internal membranes in NSCs. Farnesyl-GFP expresses the C-terminal region of human K-

Ras tagged with GFP which becomes farnesylated and plasma membrane-anchored in cells.

PH-GFP and PH-mCherry express the pleckstrin homology domain of human PLCδ tagged

with GFP or mCherry, which binds to the plasma membrane lipid phosphoinositide PI(4,5)P2.

Live imaging of actin dynamics was achieved using the F-actin marker Lifeact-mRuby

expressed in NSCs. Progression through mitosis was tracked by expressing the microtubule

marker Zeus-mCherry, which labels the mitotic spindle in NSCs. Symmetrical cell division in

NSCs was induced through the expression of dsRNAi of G protein γ1 (Gγ1, FBgn0004921)

under UAS control. Miranda-GFP expresses GFP tagged Miranda at the endogenous loci

(Ramat et al., 2017).

Method Details

Live Imaging

Intact *Drosophila* central nervous systems were dissected from third instar larvae in a bath of

Schneider's Insect Media. These larval brain explants were then mounted dorsal side down on

sterile poly-D-lysine coated 35mm glass bottom dishes (ibidi Cat#81156) containing modified

minimal hemolymph-like solution (HL3.1). Explants were imaged on a Nikon Eclipse Ti-2 (60x

H_2O objective) equipped with a Yokogawa CSU-W1 SoRa spinning disk head and dual

Photometrics Prime BSI sCMOS cameras. GFP tagged proteins were illuminated with 488nm.

Proteins tagged with mCherry and mRuby were illuminated with 561nm laser light. For

standard resolution time lapse imaging a z-stack of 41-61 optical sections with a step size of

0.5 μm was acquired every 10 seconds. For super resolution imaging, individual NSCs were

imaged using SoRa optics which achieve super resolution through optical photon

reassignment. For time lapse super resolution imaging, images were acquired every 2-60

seconds. For z-stacks acquired during super resolution imaging, a step size of 0.3 μm was

used. Where time relative to anaphase onset is noted, anaphase onset was identified as the

frame in which the apical membrane begins its expansion which coincides with the separation

of the sister chromatids (LaFoya and Prehoda, 2021). To examine the effects of cyclodextrin on

membrane dynamics, explants were treated with 15 mM methyl-β-cyclodextrin (solubilized in

HL3.1) for 4 hours prior to imaging. To study cyclodextrin's effects on cytokinesis (Figure 6b),

NSCs expressing PH-GFP and Zeus-mCherry and incubated with methyl-β-

cyclodextrin (n=10) or left untreated (n=10) were imaged throughout mitosis and assessed on

their ability to complete cytokinesis or not.

Image Processing and Analysis

Images were analyzed using ImageJ (FIJI package) and Imaris (Bitplane) software. NSCs were

identified through their location in the brain and large size, as well as the presence of

transgenes expressed from the NSC specific promoter Worniu. 3D projections were created by

z-stack processing in ImageJ, using the maximum intensity projection method for standard

resolution imaging and standard deviation projection method for super resolution imaging.

Only optical slices from the front hemisphere of the NSC were used to create 3D projections.

We define a medial optical slice as a single optical slice through the center of the cell along the

apical-basal axis. Photobleaching during time-lapse imaging was corrected for using ImageJ

bleach correction tool in histogram matching mode. When image deconvolution was applied,

deconvolution was performed using Nikon Elements standard 2D deconvolution mode. Image

deconvolution was applied to Video S3 NSCs #3, 7, & 8. Kymographs (Fig. 3b) were created

from time-lapse video using a line along the apical-basal axis using ImageJ multi kymograph

tool. To reduce image noise, Gaussian blur was applied to the Zeus-mCherry channel in Figure

6c using ImageJ.

For the cell surface area quantifications (Figure 1d), the surface area of cells marked with the

basal marker Miranda was determined using the following method. In ImageJ, a medial optical

slice was used to measure the diameter of the cell at five different angles. From the average

diameter of the cell, the surface area was calculated using the equation $SA = 4\pi r^2$, where SA =

surface area and r = radius. The surface area of the NSC at metaphase was compared to the

combined surface area of the NSC and its progeny cell just after they became separated

during division. The overall increase in apparent surface area that accompanies NSC division

was plotted for n=10 NSCs.

To quantify the thickness of membrane domains (Figure 1g), the thickness of the plasma

membrane within membrane domains and outside of membrane domains was determined

using the following method. NSCs were marked with two different membrane markers, Farnysl-

GFP and PH-mCherry. A membrane domain was identified as a region of the plasma

membrane which had high signal intensity in both channels, relative to other regions of the

plasma membrane. Medial optical slices from seven different NSCs were used to measure the
thickness of the plasma membrane within four distinct membrane domains and also at four
regions outside of membrane domains for each cell (totaling n=28 for each). Gardner-Altman
estimation plots for two group datasets and 95% confidence intervals of membrane thickness
were prepared using the DABEST package (Ho et al., 2019).
325
To quantify membrane thickness at the apical and basal poles (Figures 2b, 2c, 4c, 4d, 7b, and
7c) medial optical slices of NSCs were acquired using super resolution imaging. Images were
loaded into ImageJ, and a line which traversed the cell membrane at (or near to) either the
apical or basal pole was drawn. A plot profile was generated and the signal intensity peak
correlating to the cell membrane was identified based on its position and high intensity over
background signal. The length of the base of the peak was measured to determine the
thickness of the cell membrane. Measurements were taken at various stages of the cell cycle
and compared. When quantifying the effects of Gγ1 RNAi on membrane thickness the same
protocol was used, but only NSCs which divided symmetrically were quantified. Gardner-
Altman estimation plots for paired datasets and 95% confidence intervals of membrane
thickness were prepared using the DABEST package (Ho et al., 2019) for n=4 NSCs.
337
338 ***Quantification and Statistical Analysis***
339 Gardner-Altman estimation plots and 95% confidence intervals of datasets were prepared
340 using the DABEST package (Ho et al., 2019). Statistical details can be found in the relevant
341 methods section and figure legend.
342
343 **Key Resources Table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, Peptides, and Recombinant Proteins		
Methyl-β-cyclodextrin	Caymen Chemical	Cat#21633, CAS# 128446-36-6
Experimental Models: Organisms/Strains		
Farnesyl-GFP	Bloomington *Drosophila* Stock Center	BDSC Cat#80052, RRID:BDSC_80052
Lifeact-mRuby	Bloomington *Drosophila* Stock Center	BDSC Cat#35545, RRID:BDSC_35545

Miranda-GFP	Jens Januschke Lab	
PH-GFP	Bloomington *Drosophila* Stock Center	BDSC Cat#39693, RRID:BDSC_39693
PH-mCherry	Bloomington *Drosophila* Stock Center	BDSC Cat#51658, RRID:BDSC_51658
Worniu-Gal4	Chris Q. Doe Lab	
Zeus-mCherry	Chris Q. Doe Lab	
Gγ1 RNAi	Bloomington *Drosophila* Stock Center	BDSC Cat#34372, RRID:BDSC_34372
Software and Algorithms		
ImageJ (FIJI package)	National Institutes of Health	
Imaris	Bitplane	
Elements	Nikon	

Video Legends

Video S1: NSC membrane dynamics in early mitosis that lead to reservoir formation

Super resolution videos of PLCδ-PH-GFP (worniu-GAL4>UAS- PLCδ-PH-GFP; "PH-GFP")
expressing NSCs in early mitosis shown in standard deviation projections and in medial
sections.

Video S2: Correlated NSC membrane and cortical actin dynamics in early mitosis

Standard resolution videos of PLCδ-PH-GFP (worniu-GAL4>UAS- PLCδ-PH-GFP; "PH-GFP")
and cortical F-actin (worniu-GAL4>UAS-Lifeact-mRuby; "Lifeact") expressing NSCs in early
mitosis shown in maximum intensity projections and in medial sections.

Video S3: NSC membrane dynamics during anaphase expansion

Super resolution videos of PLCδ-PH-GFP ("PH-GFP") expressing NSCs during anaphase
shown in standard deviation projections and in medial sections. Image deconvolution was
applied to NSCs #3, 7, & 8.

Video S4: NSC membrane dynamics during mitosis

Super resolution videos of PLCδ-PH-GFP (worniu-GAL4>UAS- PLCδ-PH-GFP; "PH-GFP")
expressing NSCs showing complete mitosis in standard deviation projections and in medial
sections.

Video S5: Correlated NSC membrane and cortical actin dynamics during anaphase expansion

Super resolution videos of PLCδ-PH-GFP (worniu-GAL4>UAS- PLCδ-PH-GFP; "PH-GFP") and
cortical F-actin (worniu-GAL4>UAS-Lifeact-mRuby; "Lifeact") expressing NSCs in early mitosis
shown in standard deviation projections and in medial sections.

Video S6: Membrane dynamics in cyclodextrin-treated NSCs

Super resolution videos of PLCδ-PH-GFP (worniu-GAL4>UAS- PLCδ-PH-GFP; "PH-GFP") and
microtubules (worniu-GAL4>UAS-Zeus-mCherry; "Zeus-mCherry") expressing NSCs treated
with cyclodextrin (NSCs #1-4) or untreated (NSC #5) shown in medial sections.

Video S7: Membrane dynamics in Gγ RNAi expressing NSCs

Super resolution (NSC #1) and standard resolution (NSCs #2 & 3) videos of PLCδ-PH-GFP
(worniu-GAL4>UAS- PLCδ-PH-GFP; "PH-GFP") and Gγ RNAi (worniu-GAL4>UAS- Gγ RNAi)
expressing NSCs shown in 3D projections and in medial sections.

References

Azuma T, Kei T. 2015. Super-resolution spinning-disk confocal microscopy using optical

photon reassignment. *Opt Express* **23**:15003–15011. doi:10.1364/OE.23.015003

Barman S, Nayak DP. 2007. Lipid raft disruption by cholesterol depletion enhances influenza A

virus budding from MDCK cells. *J Virol* **81**:12169–12178. doi:10.1128/JVI.00835-07

Cabernard C, Prehoda KE, Doe CQ. 2010. A spindle-independent cleavage furrow positioning

pathway. *Nature* **467**:91–94. doi:10.1038/nature09334

Chakraborty S, Doktorova M, Molugu TR, Heberle FA, Scott HL, Dzikovski B, Nagao M,

Stingaciu L-R, Standaert RF, Barrera FN, Katsaras J, Khelashvili G, Brown MF, Ashkar

R. 2020. How cholesterol stiffens unsaturated lipid membranes. *Proc Natl Acad Sci U S*

A **117**:21896–21905. doi:10.1073/pnas.2004807117

Connell M, Cabernard C, Ricketson D, Doe CQ, Prehoda KE. 2011. Asymmetric cortical

extension shifts cleavage furrow position in Drosophila neuroblasts. *Mol Biol Cell*

22:4220–4226. doi:10.1091/mbc.E11-02-0173

Figard L, Sokac AM. 2014. A membrane reservoir at the cell surface: unfolding the plasma

membrane to fuel cell shape change. *Bioarchitecture* **4**:39–46. doi:10.4161/bioa.29069

Figard L, Wang M, Zheng L, Golding I, Sokac AM. 2016. Membrane Supply and Demand

Regulates F-Actin in a Cell Surface Reservoir. *Dev Cell* **37**:267–278.

doi:10.1016/j.devcel.2016.04.010

Figard L, Xu H, Garcia HG, Golding I, Sokac AM. 2013. The plasma membrane flattens out to

fuel cell-surface growth during Drosophila cellularization. *Dev Cell* **27**:648–655.

doi:10.1016/j.devcel.2013.11.006

Gallaud E, Pham T, Cabernard C. 2017. Drosophila melanogaster Neuroblasts: A Model for

Asymmetric Stem Cell Divisions. *Results Probl Cell Differ* **61**:183–210. doi:10.1007/978-

3-319-53150-2_8

Goudarzi M, Tarbashevich K, Mildner K, Begemann I, Garcia J, Paksa A, Reichman-Fried M,

Mahabaleshwar H, Blaser H, Hartwig J, Zeuschner D, Galic M, Bagnat M, Betz T, Raz E.

2017. Bleb Expansion in Migrating Cells Depends on Supply of Membrane from Cell

Surface Invaginations. *Dev Cell* **43**:577-587.e5. doi:10.1016/j.devcel.2017.10.030

Hirani N, Illukkumbura R, Bland T, Mathonnet G, Suhner D, Reymann A-C, Goehring NW. 2019.

Anterior-enriched filopodia create the appearance of asymmetric membrane

microdomains in polarizing C. elegans zygotes. *J Cell Sci* **132**. doi:10.1242/jcs.230714

Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. 2019. Moving beyond P values: data
analysis with estimation graphics. *Nat Methods* **16**:565–566. doi:10.1038/s41592-019-
0470-3

Homem CCF, Knoblich JA. 2012. Drosophila neuroblasts: a model for stem cell biology
139:4297–4310. doi:10.1242/dev.080515

Illukkumbura R, Bland T, Goehring NW. 2019. Patterning and polarization of cells by
intracellular flows. *Curr Opin Cell Biol* **62**:123–134. doi:10.1016/j.ceb.2019.10.005

Izumi Y, Ohta N, Itoh-Furuya A, Fuse N, Matsuzaki F. 2004. Differential functions of G protein
and Baz-aPKC signaling pathways in Drosophila neuroblast asymmetric division. *J Cell*
Biol **164**:729–738. doi:10.1083/jcb.200309162

Kabeche R, Howard L, Moseley JB. 2015. Eisosomes provide membrane reservoirs for rapid
expansion of the yeast plasma membrane. *J Cell Sci* **128**:4057–4062.
doi:10.1242/jcs.176867

Knoblich JA. 2010. Asymmetric cell division: recent developments and their implications for
tumour biology. *Nat Rev Mol Cell Biol* **11**:849–860. doi:10.1038/nrm3010

Knoblich JA. 2008. Mechanisms of asymmetric stem cell division. *Cell* **132**:583–597.
doi:10.1016/j.cell.2008.02.007

LaFoya B, Prehoda KE. 2021. Actin-dependent membrane polarization reveals the mechanical
nature of the neuroblast polarity cycle. *Cell Rep* **35**:109146.
doi:10.1016/j.celrep.2021.109146

Lang CF, Munro E. 2017. The PAR proteins: from molecular circuits to dynamic self-stabilizing
cell polarity. *Dev Camb Engl* **144**:3405–3416. doi:10.1242/dev.139063

Meenderink LM, Gaeta IM, Postema MM, Cencer CS, Chinowsky CR, Krystofiak ES, Millis BA,
Tyska MJ. 2019. Actin Dynamics Drive Microvillar Motility and Clustering during Brush
Border Assembly. *Dev Cell* **50**:545-556.e4. doi:10.1016/j.devcel.2019.07.008

Munro E, Nance J, Priess JR. 2004. Cortical flows powered by asymmetrical contraction
transport PAR proteins to establish and maintain anterior-posterior polarity in the early
C. elegans embryo. *Dev Cell* **7**:413–424. doi:10.1016/j.devcel.2004.08.001

Oon CH, Prehoda KE. 2021. Phases of cortical actomyosin dynamics coupled to the
neuroblast polarity cycle. *eLife* **10**:e66574. doi:10.7554/eLife.66574

Oon CH, Prehoda KE. 2019. Asymmetric recruitment and actin-dependent cortical flows drive
the neuroblast polarity cycle. *eLife* **8**. doi:10.7554/eLife.45815

Pang DJ, Hayday AC, Bijlmakers M-J. 2007. CD8 Raft Localization Is Induced by Its Assembly

440   into CD8αβ Heterodimers, Not CD8αα Homodimers. *J Biol Chem* **282**:13884–13894.

441   doi:10.1074/jbc.M701027200

442   Pham TT, Monnard A, Helenius J, Lund E, Lee N, Müller DJ, Cabernard C. 2019.

443   Spatiotemporally Controlled Myosin Relocalization and Internal Pressure Generate

444   Sibling Cell Size Asymmetry. *iScience* **13**:9–19. doi:10.1016/j.isci.2019.02.002

445   Ramat A, Hannaford M, Januschke J. 2017. Maintenance of Miranda Localization in Drosophila

446   Neuroblasts Involves Interaction with the Cognate mRNA. *Curr Biol CB* **27**:2101-

447   2111.e5. doi:10.1016/j.cub.2017.06.016

448   Roubinet C, Tsankova A, Pham TT, Monnard A, Caussinus E, Affolter M, Cabernard C. 2017.

449   Spatio-temporally separated cortical flows and spindle geometry establish physical

450   asymmetry in fly neural stem cells. *Nat Commun* **8**:1383. doi:10.1038/s41467-017-

451   01391-w

452   Scholze MJ, Barbieux KS, De Simone A, Boumasmoud M, Süess CCN, Wang R, Gönczy P.

453   2018. PI(4,5)P2 forms dynamic cortical structures and directs actin distribution as well

454   as polarity in Caenorhabditis elegans embryos. *Dev Camb Engl* **145**.

455   doi:10.1242/dev.164988

456   Sokac AM. 2017. Seeing a Coastline Paradox in Membrane Reservoirs. *Dev Cell* **43**:541–542.

457   doi:10.1016/j.devcel.2017.11.013

Figure 1 Neural stem cell membrane domains store excess membrane

(**A**) *Drosophila* larval central nervous system explants used in this study. Larval brain lobe neural stem cells were imaged from explants dissected from third instar larvae (vnc: ventral nerve cord).

(**B**) Model for the unequal division of neural stem cells. Preferential expansion of the apical hemisphere that occurs when the internal pressure increases during anaphase contributes to the formation of a larger apical sibling. NSC = neural stem cell. NP = neural progenitor.

(**C**) The apparent surface area of the cell follows the geometry of the main cell body and does not include the surface area contributed by the fine structure of the plasma membrane (e.g. folds and microvilli).

(**D**) Overall increase in apparent surface area that accompanies NSC division. The difference in apparent surface area at the completion of cytokinesis compared to metaphase is shown for divisions of 11 different NSCs.

(**E**) Examples of NSC membrane structure associated with localized regions of increased membrane sensor (farnesyl-GFP) intensity. A maximum intensity projection (3D projection) and medial section are shown.

(**F**) Schemes for localized enrichment of membrane sensor: 1) a configuration that yields a localized increase in sensor concentration by increased density on a single plasma membrane. 2) a configuration that yields a localized increase in intensity with uniform sensor concentration but increased membrane density.

(**G**) Variation of NSC membrane thickness associated with localized regions of increased membrane sensor intensity (membrane domains). The thickness of the membrane is shown inside and outside the domain as shown in the inset. The outside thickness is approximately equal to the diffraction limited minimum thickness.

Figure 2 Apically-directed membrane domain movements during prophase form a meta-phase membrane reservoir

(**A**) Frames from Video S1 showing the apically-directed movement of excess membrane during prophase that leads to the accumulation of excess membrane (wor-niu-GAL4>UAS-PLCd-PH-GFP; "PH-GFP") near the apical pole at metaphase. A standard deviation projection ("3D projection") is shown along with a medial section. Inset shows the basal and apical extent of one excess membrane domain as it moves apically. Time is shown relative to anaphase onset (mm:ss).

(**B**) Membrane thickness increases at the apical pole from prophase to metaphase. A paired Gardner-Altman estimation plot is shown for apical membrane thickness before apical movement commenced and metaphase. The 95% confidence interval is shown along with the bootstrap estimation distribution.

(**C**) Membrane thickness does not significantly increase at the basal pole between prophase and metaphase. Data shown as in (B).

Figure 3 Apically-directed flows of actomyosin underlie the movement of excess membrane towards the apical pole

(**A**) Selected frames from Video S2 showing coordinated movement of excess membrane domains (worniu-GAL4>UAS-PLCd-PH-GFP; "PH-GFP") and cortical F-actin (worniu-GAL4>UAS-Lifeact-mRuby; "Lifeact") during prophase. A maximum intensity projection ("3D projection") is shown along with a medial section. Arrows show correlated movement of an excess membrane domain and the corresponding localized cortical actin enrichment. Time is shown relative to anaphase onset (mm:ss).

(**B**) Kymograph constructed from Video S2 of a section along the apical-basal axis showing apically-directed movements of excess membrane domains and cortical F-actin. Time is shown relative to anaphase onset (mm:ss).

Figure 4 Consumption of the apical membrane reservoir during anaphase expansion
(**A**) Frames from Video S3 showing the consumption of the apical membrane reservoir through anaphase, furrow ingression, and late cytokinesis. Box shows inset area in row 3. A standard deviation projection ("3D projection") is shown of the membrane (wor-niu-GAL4>UAS-PLCd-PH-GFP; "PH-GFP") along with a medial section. Arrows show microvilli as it is absorbed into the cell body. Time is shown relative to anaphase onset (mm:ss).
(**B**) Straightened membrane sections from Video S3 showing the apical pole to the furrow at times relative to anaphase onset.
(**C**) Membrane thickness at the apical pole at metaphase and the completion of cytokinesis. A paired Gardner-Altman estimation plot is shown for apical membrane thickness at both time points. The 95% confidence interval is shown along with the bootstrap estimation distribution.
(**D**) Membrane thickness at the basal pole at metaphase and the completion of cytokinesis. Data is shown as in (C) but for the membrane near the basal pole.

Figure 5 Cortical actin dynamics during membrane reservoir consumption
Selected frames from Video S5 showing correlated F-actin and membrane dynamics (wor-niu-GAL4>UAS-PLCd-PH-GFP; "PH-GFP") and cortical F-actin (worniu-GAL4>UAS-LifeAct-mRu-by; "Lifeact") during anaphase. Standard deviation projection view of the apical hemisphere (apical pole is marked with a yellow circle). Time is shown relative to anaphase onset (mm:ss).

Figure 6 Cyclodextrin treatment removes excess membrane
(**A**) Standard deviation projection ("3D projection") and medial section showing the effect of cyclodextrin treatment on the NSC membrane (worniu-GAL4>UAS-PLCd-PH-GFP; "PH-GFP").
(**B**) Effect of cyclodextrin treatment on cytokinesis. "Completion" denotes cells that completed furrowing whereas "failure" denotes cells whose furrow failed to completely close.
(**C**) Selected frames from Video S6 showing the effect of cyclodextrin treatment on the NSC membrane and NSC cytokinesis. Arrows mark furrow that initially ingresses and ultimately retracts. The Zeus-mCherry microtubule binding protein is shown in the second row and merge. Time is shown relative to anaphase onset (mm:ss).

Figure 7 Membrane reservoir asymmetry is required for biased expansion
(**A**) Selected frames from Video S7 of an NSC expressing RNAi directed against Gγi. A standard deviation projection ("3D projection") is shown of the membrane (worniu-GAL4>UAS-PLCd-PH-GFP; "PH-GFP") along with a medial section.
(**B**) Membrane thickness at the apical pole at metaphase and the completion of cytokinesis for Gγi-expressing NSCs. A paired Gardner-Altman estimation plot is shown for apical membrane thickness at both time points. The 95% confidence interval is shown along with the bootstrap estimation distribution.
(**C**) Membrane thickness at the basal pole at metaphase and the completion of cytokinesis. Data is shown as in (B) but for the membrane near the basal pole.