POPULATION GENOMICS OF STONE AGE EURASIA

Morten E. Allentoft[1,2]*§, Martin Sikora[1]*§, Alba Refoyo-Martínez[1]§, Evan K. Irving-Pease[1]§, Anders Fischer[3,4]§, William Barrie[5], Andrés Ingason[6,1]§, Jesper Stenderup[1], Karl-Göran Sjögren[3], Alice Pearson[7], Barbara Mota[8,9], Bettina Schulz Paulsson[3], Alma Halgren[10], Ruairidh Macleod[1,5,11], Marie Louise Schjellerup Jørkov[12], Fabrice Demeter[1,13], Maria Novosolov[1], Lasse Sørensen[14], Poul-Otto Nielsen[14], Rasmus H.A. Henriksen[1], Tharsika Vimala[1], Hugh McColl[1], Ashot Margaryan[15,16], Melissa Ilardo[17], Andrew Vaughn[18], Morten Fischer Mortensen[14], Anne Birgitte Nielsen[19], Mikkel Ulfeldt Hede[20], Peter Rasmussen[14], Lasse Vinner[1], Gabriel Renaud[21], Aaron Stern[18], Theis Zetner Trolle Jensen[15], Niels Nørkjær Johannsen[22], Gabriele Scorrano[1], Hannes Schroeder[15], Per Lysdahl[23], Abigail Daisy Ramsøe[1], Andrei Skorobogatov[24], Andrew Joseph Schork[6,25], Anders Rosengren[6,1], Anthony Ruter[1], Alan Outram[26], Aleksey A. Timoshenko[27], Alexandra Buzhilova[28], Alfredo Coppa[29], Alisa Zubova[30], Ana Maria Silva[31,59], Anders J. Hansen[1], Andrey Gromov[30], Andrey Logvin[32], Anne Birgitte Gotfredsen[1], Bjarne Henning Nielsen[33], Borja González-Rabanal[34], Carles Lalueza-Fox[35], Catriona J. McKenzie[26], Charleen Gaunitz[1], Concepción Blasco[36], Corina Liesau[36], Cristina Martinez-Labarga[37], Dmitri V. Pozdnyakov[27], David Cuenca-Solana[38,39], David O. Lordkipanidze[40,41], Dmitri En'shin[42], Domingo C. Salazar-García[43,44], T. Douglas Price[45], Dušan Borić[29,46], Elena Kostyleva[47], Elizaveta V. Veselovskaya[48], Emma R. Usmanova[49,50], Enrico Cappellini[15], Erik Brinch Petersen[51], Esben Kannegaard[52], Francesca Radina[53], Fulya Eylem Yediay[1], Henri Duday[54], Igor Gutiérrez-Zugasti[38], Inna Potekhina[55,56], Irina Shevnina[32], Isin Altinkaya[1], Jean Guilaine[57], Jesper Hansen[58], Joan Emili Aura Tortosa[43], João Zilhão[59,60], Jorge Vega[61], Kristoffer Buck Pedersen[62], Krzysztof Tunia[63], Lei Zhao[1], Liudmila N. Mylnikova[27], Lars Larsson[64], Laure Metz[65], Levon Yeppiskoposyan[66,93], Lisbeth Pedersen[67], Lucia Sarti[68], Ludovic Orlando[69], Ludovic Slimak[65], Lutz Klassen[52], Malou Blank[3], Manuel González-Morales[38], Mara Silvestrini[70], Maria Vretemark[71], Marina S. Nesterova[27], Marina Rykun[72], Mario Federico Rolfo[73], Marzena Szmyt[74], Marcin Przybyła[75], Mauro Calattini[68], Mikhail Sablin[76], Miluše Dobisíková[77], Morten Meldgaard[78], Morten Johansen[79], Natalia Berezina[28], Nick Card[80], Nikolai A. Saveliev[81], Olga Poshekhonova[42], Olga Rickards[37], Olga V. Lozovskaya[82], Otto Christian Uldum[79], Paola Aurino[83], Pavel Kosintsev[84,85], Patrice Courtaud[54], Patricia Ríos[36], Peder Mortensen[86], Per Lotz[87,88], Per Åke Persson[89], Pernille Bangsgaard[90], Peter de Barros Damgaard[1], Peter Vang Petersen[14], Pilar Prieto Martinez[91], Piotr Włodarczak[63], Roman V. Smolyaninov[92], Rikke Maring[22,52], Roberto Menduiña[61], Ruben Badalyan[91], Rune Iversen[51], Ruslan Turin[24], Sergey Vasilyev[27], Sidsel Wåhlin[23], Svetlana Borutskaya[28], Svetlana Skochina[42], Søren Anker Sørensen[87], Søren H. Andersen[94], Thomas Jørgensen[87], Yuri B. Serikov[95], Vyacheslav I. Molodin[27], Vaclav Smrcka[96], Victor Merz[97], Vivek Appadurai[6], Vyacheslav Moiseyev[30], Yvonne Magnusson[98], Kurt H. Kjær[1], Niels Lynnerup[12], Daniel J. Lawson[99], Peter H. Sudmant[10,18], Simon Rasmussen[100], Thorfinn Korneliussen[1]@, Richard Durbin[7,101]@, Rasmus Nielsen[10,1]@, Olivier Delaneau[8]@, Thomas Werge[1,6,102]@, Fernando Racimo[1]@, Kristian Kristiansen[1,3]@, Eske Willerslev[1,5,101,103]*@

Affiliations

[1]Lundbeck Foundation GeoGenetics Centre, Globe Institute, University of Copenhagen, Copenhagen, Denmark. [2]Trace and Environmental DNA (TrEnD) Laboratory, School of Molecular and Life Sciences, Curtin University, Perth, Australia. [3]Department of Historical Studies, University of Gothenburg, Gothenburg, Sweden. [4]Sealand Archaeology, Gl. Roesnaesvej 27, 4400 Kalundborg, Denmark. [5]GeoGenetics Group, Department of Zoology, University of Cambridge, Cambridge, UK. [6]Institute of Biological Psychiatry, Mental Health Services, Copenhagen University Hospital, Roskilde, Denmark. [7]Department of Genetics, University of Cambridge, Cambridge, UK. [8]Department of Computational Biology, University of Lausanne, Switzerland. [9]Swiss Institute of Bioinformatics, University of Lausanne, Switzerland. [10]Department of Integrative Biology, University of California, Berkeley, USA. [11]Research department of Genetics, Evolution and Environment, University College London, London, UK. [12]Laboratory of Biological Anthropology, Department of Forensic Medicine, University of Copenhagen, Copenhagen, Denmark. [13]Muséum national d'Histoire naturelle, CNRS, Université de Paris, Musée de l'Homme, Paris, France. [14]The National Museum of Denmark, Ny Vestergade 10, Copenhagen, Denmark. [15]Section for Evolutionary Genomics, GLOBE Institute, University of Copenhagen, Copenhagen, Denmark. [16]Centre for Evolutionary Hologenomics, University of Copenhagen, Copenhagen, Denmark. [17]Anthropology Department, University of Utah, USA. [18]Center for Computational Biology, University of California, Berkeley, USA. [19]Department of Geology, Lund University, Lund, Sweden. [20]Tårnby Gymnasium og HF, Kastrup, Denmark. [21]Department of Health Technology, Section of Bioinformatics, Technical University of Denmark, Kongens Lyngby, Denmark. [22]Department of Archaeology and Heritage Studies, Aarhus University, Aarhus, Denmark. [23]Vendsyssel Historiske Museum, DK-9800 Hjørring, Denmark. [24]Terra Ltd., Letchik Zlobin St. 20, Voronezh, 394055, Russian Federation. [25]Neurogenomics Division, The Translational Genomics Research Institute (TGEN), Phoenix, AZ, USA. [26]Department of Archaeology, University of Exeter, Exeter, UK. [27]Institute of Archaeology and Ethnography, Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russian Federation. [28]Research Institute and Museum of Anthropology, Lomonosov Moscow State University, Mokhovaya str. 11, Moscow, Russian Federation. [29]Department of Environmental Biology, Sapienza University of Rome, Rome, Italy. [30]Peter the Great Museum of Anthropology and Ethnography (Kunstkamera), Russian Academy of Sciences, Saint Petersburg, Russian Federation. [31]CIAS, Department of Life Science, University of

Coimbra, Coimbra, Portugal. [32]Kostanay Regional University A. Baitursynov, Kostanay, Kazakhstan.
[33]Vesthimmerlands Museum, Søndergade 44, Aars, Denmark. [34]Grupo EvoAdapta, Departamento de Ciencias Históricas, Universidad de Cantabria, Santander, Spain. [35]Institute of Evolutionary Biology, CSIC-Universitat Pompeu Fabra, Barcelona, Spain. [36]Departamento de Prehistoria y Arqueología Department, Universidad Autónoma de Madrid, Madrid, Spain. [37]Department of Biology, University of Rome "Tor Vergata", Rome, Italy. [38]Instituto Internacional de Investigaciones Prehistóricas de Cantabria, Universidad de Cantabria, Santander, Spain. [39]Centre de Recherche en Archéologie, Archeosciences, Histoire (CReAAH), UMR-6566 CNRS, Rennes, France. [40]Georgian National Museum, Tbilisi, Georgia. [41]Tbilisi State University, Tbilisi, Georgia. [42]IPND, Tyumen Scientific Centre, Siberian Branch of the Russian Academy of Sciences, Tyumen, Russian Federation. [43]Departament de Prehistòria, Arqueologia i Història Antiga, Universitat de València, València, Spain. [44]Department of Geological Sciences, University of Cape Town, Cape Town, South Africa. [45]Laboratory for Archaeological Chemistry, Department of Anthropology, University of Wisconsin-Madison, Madison, USA. [46]Department of Anthropology, New York University, New York, USA. [47]Institute of Humanities, Ivanovo State University, Ivanovo, Russian Federation. [48]Institute of Ethnology and Anthropology, Russian Academy of Sciences, Moscow, Russian Federation. [49]Saryarka Archaeological Institute, Buketov Karaganda University, Karaganda, Kazakhstan. [50]South Ural State University, Chelyabinsk, Russia. [51]The Saxo Institute, University of Copenhagen, Copenhagen, Denmark. [52]Museum Østjylland, Stemannsgade 2, Randers, Denmark. [53]Soprintendenza Archeologia Belle Arti e Paesaggio per la Città Metropolitana di Bari, Via Pier l'Eremita, 25, 70122, Bari, Italy. [54]UMR 5199 PACEA, CNRS, Université de Bordeaux, 33615 Pessac, France. [55]Institute of Archaeology, National Academy of Sciences of Ukraine, Kyiv, Ukraine. [56]National University of Kyiv-Mohyla Academy, Kyiv, Ukraine. [57]Collège de France, 75231 Paris cedex 05, France. [58]Odense City Museums, Overgade 48, Odense, Denmark. [59]UNIARQ, University of Lisbon, Lisbon, Portugal. [60]ICREA, University of Barcelona, Barcelona, Spain. [61]ARGEA Consultores SL, C. de San Crispín, Madrid, Spain. [62]Museum Sydøstdanmark, Algade 97, 4760 Vordingborg, Denmark. [63]Institute of Archaeology and Ethnology, Polish Academy of Sciences, Kraków, Poland. [64]Department of Archaeology and Ancient History, Lund University, Lund, Sweden. [65]CNRS UMR 5608, Toulouse Jean Jaurès University, Maison de la Recherche, 5 Allées Antonio Machado, 31058 Toulouse, Cedex 9, France. [66]Institute of Molecular Biology, National Academy of Sciences, Yerevan, Armenia. [67]HistorieUdvikler, Gl. Roesnaesvej 27, DK-4400 Kalundborg, Denmark. [68]Department of history and cultural heritage, University of Siena, Siena, Italy. [69]Centre d'Anthropobiologie et de Génomique de Toulouse, CNRS UMR 5288, Université Paul Sabatier, Toulouse, France. [70]Soprintendenza per i Beni Archeologici delle Marche, Via Birarelli 18, 60100, Ancona, Italy. [71]Västergötlands museum, Stadsträdgården, Skara, Sweden. [72]Cabinet of Anthropology, Tomsk State University, Tomsk, Russian Federation. [73]Department of History, Humanities and Society, University of Rome "Tor Vergata", Rome, Italy. [74]Institute for Eastern Research, Adam Mickiewicz University in Poznań, Poznań, Poland. [75]Institute of Archaeology, Jagiellonian University, Ul. Gołębia 11, 31-007, Kraków, Poland. [76]Zoological Institute of Russian Academy of Sciences, Universitetskaya nab. 1, 199034, St. Petersburg, Russian Federation. [77]Department of Anthropology, Czech National Museum, Prague, Czech Republic. [78]Department of Health and Nature, University of Greenland, Greenland. [79]The Viking Ship Museum, Vindeboder 12, Roskilde, Denmark. [80]Archaeology Institute, University of Highlands and Islands, Scotland, UK. [81]Scientific Research Center "Baikal region", Irkutsk State University; 1, K. Marx st., Irkutsk, 664003, Russian Federation. [82]Laboratory for Experimental Traceology, Institute for the History of Material Culture of the Russian Academy of Sciences, Dvortsovaya nab., 18, 191186, St. Petersburg, Russian Federation. [83]Soprintendenza Archeologia, Belle Arti e Paesaggio per la provincia di Cosenza, Cosenza, Italy. [84]Paleoecology Laboratory, Institute of Plant and Animal Ecology, Ural Branch of the Russian Academy of Sciences, Ekaterinburg, Russian Federation. [85]Department of History of the Institute of Humanities, Ural Federal University, Ekaterinburg, Russian Federation. [86]Centre for the Study of Early Agricultural Societies, Department of Cross-Cultural and Regional Studies, University of Copenhagen, 2300 Copenhagen, Denmark. [87]Museum Nordsjælland, Frederiksgade 9, 3400 Hillerød. [88]Museum Vestsjælland, Klosterstræde 18, 4300 Holbæk, Denmark. [89]Museum of Cultural History, University of Oslo, P.O. Box 6762. St. Olavs Plass NO-0130 Oslo, Norway. [90]ArchaeoScience, GLOBE Institute, University of Copenhagen, Copenhagen, Denmark. [91]Department of History, University of Santiago de Compostela, Spain. [92]Lipetsk Regional Scientific Public Organisation "Archaeological Research", Lipetsk, Russian Federation. [92]Institute of Archaeology and Ethnography, National Academy of Sciences, Yerevan, Armenia. [93]Russian-Armenian University, Yerevan, Armenia. [94]Moesgaard Museum, Moesgård Allé 15, Højbjerg, Denmark. [95]Nizhny Tagil State Socio-Pedagogical Institute, Nizhny Tagil, Russia. [96]Institute for History of Medicine, First Faculty of Medicine, Charles University, Prague, Czech Republic. [97]Centre for Archaeological Research Toraighyrov University, Pavlodar, Kazakhstan. [98]Malmö Museer, Malmöhusvägen 6, Malmö, Sweden. [99]Institute of Statistical Sciences, School of Mathematics, University of Bristol, Bristol, UK. [100]Novo Nordisk Foundation Centre for Protein Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen N, Denmark. [101]Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge, UK. [102]Department of Clinical Medicine, University of Copenhagen, 2200 Copenhagen N, Denmark. [103]MARUM, University of Bremen, Bremen, Germany.

* Corresponding authors; email: morten.allentoft@curtin.edu.au, martin.sikora@sund.ku.dk, ew482@cam.ac.uk
§ Joint first authors

120 @ Joint last authors
121
122
123 **Summary**
124

The transitions from foraging to farming and later to pastoralism in Stone Age Eurasia (c. 11-
3 thousand years before present, BP) represent some of the most dramatic lifestyle changes in
human evolution. We sequenced 317 genomes of primarily Mesolithic and Neolithic
individuals from across Eurasia combined with radiocarbon dates, stable isotope data, and
pollen records. Genome imputation and co-analysis with previously published shotgun
sequencing data resulted in >1600 complete ancient genome sequences offering fine-grained
resolution into the Stone Age populations. We observe that: 1) Hunter-gatherer groups were
more genetically diverse than previously known, and deeply divergent between western and
eastern Eurasia. 2) We identify hitherto genetically undescribed hunter-gatherers from the
Middle Don region that contributed ancestry to the later Yamnaya steppe pastoralists; 3) The
genetic impact of the Neolithic transition was highly distinct, east and west of a boundary
zone extending from the Black Sea to the Baltic. Large-scale shifts in genetic ancestry
occurred to the west of this "Great Divide", including an almost complete replacement of
hunter-gatherers in Denmark, while no substantial ancestry shifts took place during the same
period to the east. This difference is also reflected in genetic relatedness within the
populations, decreasing substantially in the west but not in the east where it remained high
until c. 4,000 BP; 4) The second major genetic transformation around 5,000 BP happened at a
much faster pace with Steppe-related ancestry reaching most parts of Europe within 1,000-
years. Local Neolithic farmers admixed with incoming pastoralists in eastern, western, and
southern Europe whereas Scandinavia experienced another near-complete population
replacement. Similar dramatic turnover-patterns are evident in western Siberia; 5) Extensive
regional differences in the ancestry components involved in these early events remain visible
to this day, even within countries. Neolithic farmer ancestry is highest in southern and eastern
England while Steppe-related ancestry is highest in the Celtic populations of Scotland, Wales,
and Cornwall (this research has been conducted using the UK Biobank resource); 6) Shifts in
diet, lifestyle and environment introduced new selection pressures involving at least 21
genomic regions. Most such variants were not universally selected across populations but
were only advantageous in particular ancestral backgrounds. Contrary to previous claims, we
find that selection on the FADS regions, associated with fatty acid metabolism, began before
the Neolithisation of Europe. Similarly, the lactase persistence allele started increasing in
frequency before the expansion of Steppe-related groups into Europe and has continued to
increase up to the present. Along the genetic cline separating Mesolithic hunter-gatherers
from Neolithic farmers, we find significant correlations with trait associations related to skin
disorders, diet and lifestyle and mental health status, suggesting marked phenotypic
differences between these groups with very different lifestyles. This work provides new
insights into major transformations in recent human evolution, elucidating the complex
interplay between selection and admixture that shaped patterns of genetic variation in
modern populations.

163
164
165 **Introduction**
166
167 The transition from hunting and gathering to farming represents one of the most dramatic shifts in
     lifestyle and diet in human evolution with lasting effects on the modern world. For millions of years
our ancestors relied on hunting and foraging for survival but c.12,000 years ago in the Fertile

Crescent of the Near East, plant cultivation and animal husbandry were developed[1–3]. This
ultimately resulted in a more sedentary lifestyle accompanied by increasing population sizes and
higher social complexity. Expanding populations and the adoption of herding, carried farming
practices into Europe and parts of SW Asia in the following millennia, and farming was also
developed independently in other parts of the World. Today, 50% of the Earth's habitable land is
used for agriculture and very few hunter-gatherers remain[4,5]. Understanding the changes to the
human gene pool during this shift from hunter-gathering to farming between the Mesolithic and
Neolithic periods is central to understanding ourselves and the events that led to a major
transformation of our planet.
178
While the Neolithisation process has been studied extensively with ancient DNA (aDNA)
technology, several key questions remain unaddressed. Population movements during the Neolithic
can be traced in the gene pools across the European continent as farming was introduced from the
Near East. Several regional studies have testified to varying degrees of reproductive interaction
with local Mesolithic groups, ranging from genetic continuity[6] to gradual population admixture[7–10]
to almost complete replacement[11]. However, our knowledge of the population structure in the
Mesolithic period and how it was formed is limited, partly because of a paucity of data from
skeletons older than 8,000 years, compromising resolution into subsequent demographic transitions.
Moreover, the spatiotemporal mapping of population dynamics east of Europe, including Siberia,
Central- and North Asia during the same time period remains patchy. In these regions the
'Neolithic' typically refers to new forms of lithic material culture, and/or the presence of
ceramics[12]. For instance, the Neolithic cultures of the Central Asian Steppe possessed pottery, but
retained a hunter-gatherer economy alongside stone blade technology similar to the preceding
Mesolithic cultures[13]. The archaeological record testifies to a boundary, ranging from the eastern
Baltic to the Black Sea, east of which hunter-gatherer societies persist for much longer than in
western Europe[14]. The population genomic implications of this "Great Divide" is, however, largely
unknown. Southern Scandinavia represents another enigma in the Neolithisation debate[15]. The
introduction of farming reached a 1,000-year standstill at the doorstep to Southern Scandinavia
before finally progressing into Denmark around 6,000 BP. It is not known what caused this delay
and whether the transition to farming in Denmark, was facilitated by the migration of people (demic
diffusion), similar to the rest of Europe[11,16,17] or mostly involved cultural diffusion[18,19]. Starting at
around 5,000 BP, a new ancestry component emerged on the eastern European plains associated
with Yamnaya Steppe pastoralists culture and swept across Europe mediated through expansion of
the Corded Ware complex (CWC) and related cultures[20,21]. The genetic origin of the Yamnaya and
the fine-scale dynamics of the formation and expansion of the CWC are largely unresolved
questions of central importance to clarify the formation of the present day European gene pool.
205
Rapid dietary changes and expansion into new climate zones represent shifts in environmental
exposure, impacting the evolutionary forces acting on the gene pool. The Neolithisation can
therefore be considered as a series of large-scale selection pressures imposed on humans from
around 12,000 years ago. Moreover, close contact with livestock and higher population densities
have likely enhanced exposure and transmission of infectious diseases, introducing new challenges
to our survival[22,23]. While signatures of selection can be identified from patterns of genetic diversity
in extant populations[24,25], this can be challenging in species such as humans, which show very wide
geographic distributions and have thus been exposed to highly diverse and changing local
environments through space and time. In the complex mosaic of ancestries that constitute a modern
human genome any putative signatures of selection may therefore misrepresent the timing and
magnitude of the actual event unless we can use ancient DNA to chart the individual ancestry
components back into the evolutionary past.
218

To investigate these formative processes in Eurasian prehistory, we conducted the largest ancient
DNA study to date on human Stone Age skeletal material. We sequenced low-coverage genomes of
317 radiocarbon-dated (AMS) primarily Mesolithic and Neolithic individuals, covering major parts
of Eurasia. We combined these with published shotgun-sequenced data to impute a dataset of
>1600 diploid ancient genomes. Genomic data from 100 AMS-dated individuals from Denmark
supported detailed analyses of the Stone Age population dynamics in Southern Scandinavia. When
combined with genetically-predicted phenotypes, proxies for diet (δ^{13}C/δ^{15}N), mobility (^{87}Sr/^{86}Sr)
and vegetation cover (pollen) we could connect this with parallel shifts in phenotype, subsidence
and landscape. To test for traces of divergent selection in health and lifestyle-related genetic
variants, we used the imputed ancient genomes to reconstruct polygenic risk scores for hundreds of
complex traits in the ancient Eurasian populations. Additionally, we used a novel chromosome
painting technique based on tree sequences, in order to model ancestry-specific allele frequency
trajectories through time. This allowed us to identify many new phenotype-associated genetic
variants with hitherto unknown evidence for positive selection in Eurasia throughout the Holocene.
233

Results/Discussion

Samples and data

In this study we present genomic data from 317 ancient individuals (Fig 1, Extended data fig. 2,
Supplement Table I). A total of 272 were radiocarbon dated within the project, while 39 dates were
derived from literature and 15 were dated by archaeological context. Dates were corrected for
marine and freshwater reservoir effects (Supplementary Note 8) and ranged from the Upper
Palaeolithic (UP) c. 25,700 calibrated years before present (cal. BP) to the mediaeval period (c.
1200 cal. BP). However, 97% of the individuals (N=309) span 11,000 cal. BP to 3,000 cal. BP, with
a heavy focus on individuals associated with various Mesolithic and Neolithic cultures.
243

Geographically, the sampled skeletons cover a vast territory across Eurasia, from Lake Baikal to the
Atlantic coast, from Scandinavia to the Middle East, and they derive from a variety of contexts,
including burial mounds, caves, bogs and the seafloor (Supplementary Notes 6-7). Broadly, we can
divide our research area into three large regions: 1) central, western and northern Europe, 2) eastern
Europe including western Russia and Ukraine, and 3) the Urals and western Siberia. Our samples
cover many of the key Mesolithic and Neolithic cultures in Western Eurasia, such as the
Maglemose and Ertebølle cultures in Scandinavia, the Cardial in the Mediterranean, the Körös and
Linear Pottery (LBK) in SE and Central Europe, and many archaeological cultures in Ukraine,
western Russia, and the trans-Ural (e.g. Veretye, Lyalovo, Volosovo, Kitoi). Our sampling was
particularly dense in Denmark from where we present a detailed and continuous sequence of 100
genomes spanning from the early Mesolithic to the Bronze Age. Dense sample sequences were also
obtained from Ukraine, Western Russia, and the trans-Ural, spanning from the Early Mesolithic
through the Neolithic, up to c. 5,000 BP.
257

We extracted ancient DNA from tooth cementum or petrous bone and shotgun sequenced the 317
genomes to a depth of genomic coverage ranging from 0.01X to 7.1X (mean = 0.75X, median =
0.26X), with 81 individuals having >1X coverage. Using a new imputation method designed for
low-coverage sequencing data[26], we performed genotype imputation based on the 1,000 Genomes
phased data as a reference panel. We also imputed >1,300 previously published shotgun-sequenced
ancient genomes. This resulted in a "raw" dataset containing 8.5 million common Single Nucleotide
Polymorphisms (SNPs) (>1% MAF and imputation info score > 0.5) from 1,664 imputed diploid
ancient genomes. This number includes 42 high-coverage ancient genomes (Table S2.1,
Supplementary Note 2) that were down-sampled to values between 0.1X and 4X for validation.

This demonstrated that 1-fold genome coverage provides remarkably high imputation accuracy
(r^2>0.95 at common variants with MAF above 5%) and closely matches what is obtained for
modern samples (Extended Fig. 1A-D). African genomes, however, exhibit lower imputation
accuracy as a result of the poor representation of this ancestry in the reference panel. For European
genomes, this translates into genotyping error rates usually below 5% for the most challenging
genotypes to impute (heterozygous genotypes or with two copies of the non-reference allele;
Supplementary Fig. S2.1-S2.2). Imputation accuracy also depends on minor allele frequency and
genomic coverage (Supplementary Fig. S2.3). We find that coverage values as low as 0.1x and
0.4X are sufficient to obtain r^2 imputation accuracy of 0.8 and 0.9 at common variants
(MAF>=10%), respectively. As further validation, we increased genomic coverage to 27.5X, 18.9X
and 5.4X on a previously published trio (mother, father, son) from the Late Neolithic mass burial at
Koszyce in Poland [27]. This allowed for a validation of imputed genotypes and haplotypes using
Mendel's rules of inheritance. We obtained Mendelian error rates from 0.1% at 4X to 0.55% at
0.1X (Extended Fig. 1E). Similarly, we obtained switch error rates between 2% and 6%. Altogether,
our validation analysis showed that ancient European genomes can be imputed confidently from
coverages above 0.4X and highly valuable data can still be obtained with coverages as low as 0.1X
when using specific QC on the imputed data, although at very low coverage a bias arise towards the
major allele (see Supplementary Note 2). We filtered out samples with poor coverage or variant
sites with low MAF in downstream analyses depending on the specific data quality requirements.
For most analyses we use a subset of 1,492 imputed ancient genomes (213 sequenced in this study)
after filtering individuals with very low coverages (<0.1X) and/or low imputation quality (average
genotype probability < 0.8) and close relatives. This dataset allows us to characterise the ancient
cross-continental gene pools and the demographic transitions with unprecedented resolution.
290
We performed broad-scale characterization of this dataset using principal component analysis
(PCA) and model-based clustering (ADMIXTURE), recapitulating and providing increased
resolution into previously described ancestry clines in ancient Eurasian populations (Fig. 1;
Extended Data Fig. 2; Supplementary Note 3d). Strikingly, inclusion of the imputed ancient
genomes in the inference of the principal components reveals much higher variance among the
ancient groups than previously anticipated using projection onto a PC-space inferred from modern
individuals alone (Extended Data Fig. 2). This is particularly notable in a PCA of West Eurasian
individuals, where genetic variation among all present-day populations is confined within a small
central area of the PCA (Extended Data Fig. 2C, D). These results are consistent with much higher
genetic differentiation between ancient Europeans than present-day populations reflecting lower
effective population sizes and genetic isolation among ancient groups.
302
To obtain a finer-scale characterization of genetic ancestries across space and time, we assigned
imputed ancient individuals to genetic clusters by applying hierarchical community detection on a
network of pairwise identity-by-descent (IBD)-sharing similarities[28] (Extended Data Fig. 3;
Supplementary Note 3c). The obtained clusters capture fine-scale genetic structure corresponding to
shared ancestry within particular spatiotemporal ranges and/or archaeological contexts, and were
used as sources and/or targets in supervised ancestry modelling (Extended Data Fig. 4;
Supplementary Note 3i). We focus our subsequent analyses on three panels of putative source
clusters reflecting different temporal depths: "deep", using a set of deep ancestry source groups
reflecting major ancestry poles; "postNeol", using diverse Neolithic and earlier source groups; and
"postBA", using Late Neolithic and Bronze Age source groups (Extended Data Fig. 4).
313
314
315

316
Fig 1. Sample overview and broad scale genetic structure. (A), (B) Geographic and temporal distribution of the 317
ancient genomes reported here. Age and geographic region of ancient individuals are indicated by plot symbol colour
and shape, respectively. Random jitter was added to geographic coordinates to avoid overplotting. (C), (D) Principal
component analysis of 3,316 modern and ancient individuals from Eurasia, Oceania, and the Americas (C), as well as
restricted to 2,126 individuals from western Eurasia (west of Urals) (D). Principal components were defined using both
modern and imputed ancient genomes passing all filters, with the remaining low-coverage ancient genomes projected.
Ancient genomes sequenced in this study are indicated with black circles (imputed genomes passing all filters, n=213)
or grey diamonds (pseudo-haploid projected genomes, n=104). Genomes of modern individuals are shown in grey, with
population labels corresponding to their median coordinates.
326

Deep population structure of western Eurasians

Our study comprises the largest genomic dataset on European hunter-gatherers to date, including
113 imputed hunter-gatherer genomes of which 79 were sequenced in this study. Among them, we
report a 0.83X genome of an Upper Palaeolithic (UP) skeleton from Kotias Klde Cave in Georgia,
Caucasus (NEO283), directly dated to 26,052 - 25,323 cal BP (95%). In the PCA of all non-African
individuals, it occupies a position distinct from other previously sequenced UP individuals, shifted
towards west Eurasians along PC1 (Supplementary Note 3d). Using admixture graph modelling, we
find that this Caucasus UP lineage derives from a mixture of predominantly West Eurasian UP
hunter-gatherer ancestry (76%) with ~24% contribution from a "basal Eurasian" ghost population,
first observed in West Asian Neolithic individuals[29] (Extended Data Fig. 5A). Models attempting to
reconstruct major post-LGM clusters such as European hunter-gatherers and Anatolian farmers
without contributions from this Caucasus UP lineage provided poor admixture graph fits or were
rejected in *qpAdm* analyses (Extended Data Fig. 5B,C). These results thus suggest a central role of

the descendants related to this Caucasus UP lineage in the formation of later West Eurasian
populations, consistent with recent genetic data from the nearby Dzudzuana Cave, also in
Georgia[30].
343
We performed supervised admixture modelling using a set of twelve possible source clusters
representing Mesolithic hunter-gatherers from the extremes of the HG cline, as well as temporal or
geographical outgroups of deep Eurasian lineages (Fig 2A). We replicate previous results of broad-
scale genetic structure correlated to geography in European hunter-gatherers after the LGM[17], while
also revealing novel insights into their fine-scale structure. Ancestry related to southern European
hunter-gatherers (source: Italy_15000BP_9000 BP) predominates in western Europe. This includes
Denmark, where our 28 sequenced and imputed hunter-gatherer genomes derive almost exclusively
from this cluster, with remarkable homogeneity across a 5,000 year transect (Fig. 3A). In contrast,
hunter-gatherer individuals from the eastern and far northern reaches of Europe show the highest
proportions of Russian hunter-gatherer ancestry (source: RussiaNW_11000BP_8000BP; Fig. 2B,
D), with genetic continuity until ~5,000 BP in Russia. Ancestry related to Mesolithic hunter-
gatherer populations from Ukraine (source: Ukraine_10000BP_4000BP) is carried in highest
proportions in hunter-gatherers from a geographic corridor extending from south-eastern Europe
towards the Baltic and southern Scandinavia. Swedish Mesolithic individuals derive up to 60% of
their ancestry from that source (Fig. 2C). Our results thus indicate northwards migrations of at least
three distinct waves of hunter-gatherer ancestry into Scandinavia: a predominantly southern
European source into Denmark; a source related to Ukrainian and south-eastern European hunter-
gatherers into the Baltic and southern Sweden; and a northwest Russian source into the far north,
before venturing south along the Atlantic coast of Norway[31] (Fig. 2). These movements are likely to
represent post glacial expansions from refugia areas shared with many plant and animal species[32,33].
364
Despite the major role of geography in shaping European hunter-gatherer structure, we also
document more complex local dynamics. On the Iberian Peninsula, the earliest individuals,
including a ~9,200-year-old hunter-gatherer (NEO694) from Santa Maira (eastern Spain),
sequenced in this study, show predominantly southern European hunter-gatherer ancestry with a
minor contribution from UP hunter-gatherer sources (Fig. 3). In contrast, later individuals from
Northern Iberia are more similar to hunter-gatherers from eastern Europe, deriving ~30-40% of
their ancestry from a source related to Ukrainian hunter-gatherers[34,35]. The earliest evidence for this
gene flow is observed in a Mesolithic individual from El Mazo, Spain (NEO646) that was dated,
calibrated and reservoir-corrected to c. 8,200 BP (8365-8182 cal BP, 95%) but context-dated to
slightly older (8550-8330 BP, see[36]). The younger date coincides with some of the oldest Mesolithic
geometric microliths in northern Iberia, appearing around 8,200 BP at this site[36]. In southern
Sweden, we find higher amounts of southern European hunter-gatherer ancestry in late Mesolithic
coastal individuals (NEO260 from Evensås; NEO679 from Skateholm) than in the earlier
Mesolithic individuals from further inland, suggesting either geographic genetic structure in the
Swedish Mesolithic population or a possible eastward expansion of hunter-gatherers from
Denmark, where this ancestry prevailed (Fig. 3). An influx of southern European hunter-gatherer-
related ancestry in Ukrainian individuals after the Mesolithic (Fig. 3) suggests a similar eastwards
expansion in south-eastern Europe[17]. Interestingly, two herein reported ~7,300-year-old imputed
genomes from the Middle Don River region in the Pontic-Caspian steppe (Golubaya Krinitsa,
NEO113 & NEO212) derive ~20-30% of their ancestry from a source cluster of hunter-gatherers
from the Caucasus (Caucasus_13000BP_10000BP) (Fig. 3). Additional lower coverage (non-
imputed) genomes from the same site project in the same PCA space (Fig. 1D), shifted away from
the European hunter-gatherer cline towards Iran and the Caucasus. Our results thus document
genetic contact between populations from the Caucasus and the Steppe region as early as 7,300
years ago, providing documentation of continuous admixture prior to the advent of later nomadic

Steppe cultures, in contrast to recent hypotheses, and also further to the west than previously
reported[17,37].
392

393
Fig 2. Genetic structure of European hunter-gatherers (A) Ancestry proportions in 113 imputed ancient genomes
representing European hunter-gatherer contexts (right) estimated from supervised non-negative least squares analysis
using deep Eurasian source groups (left). Individuals from target groups are grouped by genetic clusters. (B)-(D) Moon
charts showing spatial distribution of ancestry proportions in European hunter-gatherers deriving from three deep
Eurasian source groups; Italy_15000BP_9000BP; Ukraine_10000BP_4000BP; RussiaNW_11000BP_8000BP (source
origins shown with coloured symbol). Estimated ancestry proportions are indicated by both size and amount of fill of
moon symbols.
401
402
403 *Major genetic transitions in Europe*

404 Previous ancient genomics studies have documented multiple episodes of large-scale population
405 turnover in Europe within the last 10,000 years[6,11,14,16,17,20,21,34,38–41]. The 317 genomes reported here
406 fill important knowledge gaps, particularly in northern and eastern Europe, allowing us to track the
407 dynamics of these events at both continental and regional scales.
408
Our analyses reveal profound differences in the spatiotemporal Neolithisation dynamics across
Europe. Supervised admixture modelling ("deep" set) and spatiotemporal kriging[42] document a
broad east-west distinction along a boundary zone running from the Black Sea to the Baltic. On the
western side of this "Great Divide", the Neolithic transition is accompanied by large-scale shifts in
genetic ancestry from local hunter-gatherers to Neolithic farmers with Anatolian-related ancestry
(Boncuklu_10000BP; Fig. 3; Extended Data Fig. 4, 6). The arrival of Anatolian-related ancestry in
different regions spans an extensive time period of over 3,000 years, from its earliest evidence in
the Balkans (Lepenski Vir) at ~8,700 BP[17] to c. 5,900 BP in Denmark. On the eastern side of this
divide, no ancestry shifts can be observed during this period. In the East Baltic region (see also[43]),

Ukraine and Western Russia local hunter-gatherer ancestry prevails until ~5,000 BP without
noticeable input of Neolithic Anatolian-related farmer ancestry (Fig. 3; Extended Data Fig. 4, 6).
This Eastern genetic continuity is in remarkable congruence with the archaeological record showing
persistence of pottery-using hunter-gatherer-fisher groups in this wide region, and delayed
introduction of cultivation and husbandry by several thousand years (Supplementary Note 5).
423
From approximately 5,000 BP, an ancestry component appears on the eastern European plains in
Early Bronze Age Steppe pastoralists associated with the Yamnaya culture and it rapidly spreads
across Europe through the expansion of the Corded Ware complex (CWC) and related cultures[20,21].
We demonstrate that this "steppe" ancestry (Steppe_5000BP_4300BP) can be modelled as a
mixture of ~65% ancestry related to herein reported hunter-gatherer genomes from the Middle Don
River region (MiddleDon_7500BP) and ~35% ancestry related to hunter-gatherers from Caucasus
(Caucasus_13000BP_10000BP) (Extended Data Fig. 4). Thus, Middle Don hunter-gatherers, who
already carry ancestry related to Caucasus hunter-gatherers (Fig. 2), serve as a hitherto unknown
proximal source for the majority ancestry contribution into Yamnaya genomes. The individuals in
question derive from the burial ground Golubaya Krinitsa (Supplementary Note 3). Material culture
and burial practices at this site are similar to the Mariupol-type graves, which are widely found in
neighbouring regions of Ukraine, for instance along the Dnepr River. They belong to the group of
complex pottery-using hunter-gatherers mentioned above, but the genetic composition at Golubaya
Krinitsa is different from the remaining Ukrainian sites (Fig 2A, Extended Data Fig. 4). We find
that the subsequent transition of the Late Neolithic and Early Bronze Age European gene pool
happened at a faster pace than during the Neolithisation, reaching most parts of Europe within a
~1,000-year time period after first appearing in eastern Baltic region ~4,800 BP (Fig. 3). In line
with previous reports we observe that beginning c. 4,200 BP, steppe-related ancestry was already
dominant in samples from France and the Iberian peninsula, while it reached Britain only 400 years
later[11,38,44]. Strikingly, because of the delayed Neolithisation in Southern Scandinavia these
dynamics resulted in two episodes of large-scale genetic turnover in Denmark and southern Sweden
within a 1,000-year period (Fig. 3).
446
447

Fig. 3. Genetic transects of Eurasia. Regional timelines of genetic ancestry compositions within the past 15,000 years in western Eurasia (top) and the Eurasian Steppe belt east of the Urals (bottom). Ancestry proportions in 972 imputed ancient genomes from these regions (covering c. 12,000 BP to 500 BP), inferred using supervised admixture modelling with "deep" hunter-gatherer ancestry source groups. Geographic areas included in timelines are indicated with fill colour (west Eurasia) and grey shading (eastern Steppe region). Excavation locations of the ancient skeletons are indicated with black crosses. Coloured bars within the timelines represent ancestry proportions for temporally consecutive individuals, with the width corresponding to their age difference. Individuals with identical age were offset along the time axis by adding random jitter, ages. We note that the inclusion of only shotgun-sequenced samples may affect the exact timing of events in some regions from where such data are sparse.

We next investigated fine-grained ancestry dynamics underlying these transitions. We replicate previous reports[11,16,17,21,41,45,46] of widespread, but low-level admixture between Neolithic farmers and local hunter-gatherers resulting in a resurgence of HG ancestry in many regions of Europe during the middle and late Neolithic (Extended Data Fig. 7). Estimated hunter-gatherer ancestry proportions among early Neolithic people rarely exceed 10%, with notable exceptions observed in individuals from south-eastern Europe (Iron Gates), Sweden (Pitted Ware Culture) as well as herein reported early Neolithic genomes from Portugal (western Cardial), estimated to harbour 27% – 43% Iberian hunter-gatherer ancestry (Iberia_9000BP_7000BP). The latter result, suggesting extensive first-contact admixture, is in agreement with archaeological inferences derived from modelling the

spread of farming along west Mediterranean Europe[47]. Individuals associated with Neolithic
farming cultures from Denmark show some of the highest overall hunter-gatherer ancestry
proportions (up to ~25%), mostly derived from Western European-related hunter-gatherers
(EuropeW_13500BP_8000BP) supplemented with marginal contribution from local Danish groups
in some individuals (Extended Data Fig. 7D; Supplementary Note 3f). We estimated the timing of
the admixture using the linkage-disequilibrium-based method DATES[48] at ~6,000 BP. Both lines of
evidence thus suggest that a significant part of the hunter-gatherer admixture observed in Danish
Neolithic individuals occurred already before the arrival of the incoming Neolithic people in the
region (Extended Data Fig. 7), and further imply Central Europe as a key region in the resurgence
of HG ancestry. Interestingly, the genomes of two ~5,000-year-old Danish male individuals
(NEO33, NEO898) were entirely composed of Swedish hunter-gatherer ancestry, and formed a
cluster with Pitted Ware Culture (PWC) individuals from Ajvide on the Baltic island of Gotland
(Sweden)[49–51]. Of the two individuals, NEO033 also displays an outlier Sr-signature (Fig. 4),
potentially suggesting a non-local origin matching his unusual ancestry. Overall, our results
demonstrate direct contact across the Kattegat and Öresund during Neolithic times (Extended Data
Fig. 3, 4), in line with archaeological finds from Zealand (east Denmark) showing cultural affinities
to PWC on the Swedish west coast[52–55].
486
Further, we find evidence for regional stratification in early Neolithic farmer ancestries in
subsequent Neolithic groups. Specifically, southern European early farmers appear to have provided
major genetic ancestry to mid- and late Neolithic groups in Western Europe, while central European
early farmer ancestry is mainly observed in subsequent Neolithic groups in eastern Europe and
Scandinavia (Extended Data Fig. 7D-F). These results are consistent with distinct migratory routes
of expanding farmer populations as previously suggested[8]. For example, similarities in material
culture and flint mining activities could suggest that the first farmers in South Scandinavia
originated from or had close social relations with the central European Michelsberg Culture[56].
495
The second continental-wide and CWC-mediated transition from Neolithic farmer ancestry to
Steppe-related ancestry was found to differ markedly between geographic regions. The contribution
of local Neolithic farmer ancestry to the incoming groups was high in eastern, western and southern
Europe, reaching >50% on the Iberian Peninsula ("postNeol" set; Extended Data Fig. 4, 6B, C)[34].
Scandinavia, however, portrays a dramatically different picture, with a near-complete replacement
of the local Neolithic farmer population inferred across all sampled individuals (Extended Data Fig.
7B, C). Following the second transition, Neolithic Anatolian-related farmer ancestry remains in
Scandinavia, but the source is now different. It can be modelled as deriving almost exclusively from
a genetic cluster associated with the Late Neolithic Globular Amphora Culture (GAC)
(Poland_5000BP_4700BP; Extended Data Fig. 4). Strikingly, after the Steppe-related ancestry was
first introduced into Europe (Steppe_5000BP_4300BP), it expanded together with GAC-related
ancestry across all sampled European regions (Extended Data Fig. 7I). This suggests that the spread
of steppe-related ancestry throughout Europe was predominantly mediated through groups that were
already admixed with GAC-related farmer groups of the eastern European plains. This finding has
major implications for understanding the emergence of the CWC. A stylistic connection from GAC
ceramics to CWC ceramics has long been suggested, including the use of amphora-shaped vessels
and the development of cord decoration patterns[57]. Moreover, shortly prior to the emergence of the
earliest CWC groups, eastern GAC and western Yamnaya groups exchanged cultural elements in
the forest-steppe transition zone northwest of the Black Sea, where GAC ceramic amphorae and
flint axes were included in Yamnaya burials, and the typical Yamnaya use of ochre was included in
GAC burials[58], indicating close interaction between the groups. Previous ancient genomic data from
a few individuals suggested that this was limited to cultural influences and not population
admixture[59]. However, in the light of our new genetic evidence it appears that this zone, and

possibly other similar zones of contact between GAC and Yamnaya (or other closely-related
steppe/forest-steppe groups) were key in the formation of the CWC through which steppe-related
ancestry and GAC-related ancestry co-dispersed far towards the west and the north[cf. 60]. This
resulted in regionally diverse situations of interaction and admixture[61,62] but a significant part of the
CWC dispersal happened through corridors of cultural and demic transmission which had been
established by the GAC during the preceding period[63,64].
525
526
Fine-scale structure and multiproxy analysis of Danish transect

We present a detailed and continuous sequence of multiproxy data from Denmark, from the Early
Mesolithic Maglemose, via the Kongemose and Late Mesolithic Ertebølle epochs, the Early and
Middle Neolithic Funnel Beaker Culture and the Single Grave Culture, to Late Neolithic and
Bronze Age individuals (Fig. 4). To integrate multiproxy data from as many skeletons as possible
we made use of non-imputed data for the admixture analyses (Supplementary Note S3d) which
were not restricted to the >0.1X coverage cut-off used elsewhere. This provided genetic profiles
from 100 Danish individuals (Fig 4), spanning c. 7,300 years from the earliest known skeleton in
Denmark (the Mesolithic "Koelbjerg Man" (NEO254, 10,648-10,282 cal. BP, 95% probability
interval) and formerly known as the "Koelbjerg Woman"[65]), to a Bronze Age skeleton from Hove Å
(NEO946) dated to 3322-2967 cal. BP (95%). Two temporal shifts in genomic admixture
proportions confirm the major population genetic turnovers (Fig. 4) that was inferred from imputed
data (Fig. 3). The multiproxy evidence, however, unveils the dramatic concomitant changes in all
investigated phenotypic, environmental and dietary parameters (Fig. 4).
541

543
Fig 4. Environmental, dietary, phenotypic and ancestry shifts in Denmark through time. Two dramatic population
turnovers are evident from chronologically-sorted multiproxy data representing 100 Danish Stone Age and early Bronze
Age skeletons sequenced in this study. The figure shows concomitant changes in several investigated parameters
including (from the top) admixture proportions from non-imputed autosomal genome-wide data, Y-chromosomal and
mitochondrial haplogroups, genetic phenotype predictions (based on imputed data) as well as $^{87}Sr/^{86}Sr$ and $\delta^{13}C$ and
$\delta^{15}N$ isotope data as possible proxies for mobility and diet, respectively. Predicted height values represent differences
(in cm) from the average height of the present-day Danish population, based on genotypes at 310 height-associated loci
(Supplementary Note 4f). Probabilities for the indicated natural eye and hair colours are based on genotypes at 18
pigmentation-associated loci (Supplementary Note 4f) with grey denoting probability of intermediate eye colour
(including grey, green and hazel). Lower panel shows changes in vegetation as predicted from pollen analyses at Lake
Højby in Zealand (Supplementary Note 12). Black vertical lines mark the first presence of Anatolian farmer ancestry
and Steppe-related ancestry, respectively.
556
557
During the Danish Mesolithic, individuals from the Maglemose, Kongemose and Ertebølle cultures
displayed a remarkable genetic homogeneity across a 5,000 year transect deriving their ancestry
almost exclusively from a southern European source (source: Italy_15000BP_9000BP) that later
predominates in western Europe (Fig. 2). These cultural transitions occurred in genetic continuity,
apparent in both autosomal and uniparental markers, which rules out demic diffusion and supports
the long-held assumption of a continuum of culture and population[e.g. 66–68]. Genetic predictions
indicate blue eye pigmentation with high probability in several individuals throughout the duration
of the Mesolithic (Supplementary Note 4f), consistent with previous findings [11,20,45]. In contrast,
none of the analysed Mesolithic individuals displayed high probability of light hair pigmentation.
Height predictions for Mesolithic individuals generally suggest slightly lower or perhaps less
variable genetic values than in the succeeding Neolithic period. However, we caution that the
relatively large genetic distance to modern individuals included in the GWAS panel make these
scores poorly applicable to Mesolithic individuals (Supplementary Note 4c) and are dependent on
the choice of GWAS filters used. Unfortunately, only a fraction of the 100 Danish skeletons
included were suitable for stature estimation by actual measurement, why these values are not
reported.
574
Stable isotope $\delta^{13}C$ values in collagen inform on the proportion of marine versus terrestrial protein,
while $\delta^{15}N$ values reflect the trophic level of protein sources[69,70]. Both the Koelbjerg Man and the
second earliest human known from Denmark, (Tømmerupgårds Mose – not part of the present
study; see[71]) showed more depleted dietary isotopic values, representing a lifestyle of inland hunter-
fisher-gatherers of the early Mesolithic forest. A second group consisted of coastal fisher-hunter-
gatherers dating to the late half of the Maglemose epoch onwards (Supplementary Figs. S10.1 and
S10.2). During this period global sea-level rise gradually changed the landscape of present-day
Denmark from an interior part of the European continent to an archipelago, where all human groups
had ample access to coastal resources within their annual territories. Increased $\delta^{13}C$ and $\delta^{15}N$ values
imply that from the late Maglemose marine foods gradually increased in importance, to form the
major supply of proteins in the final Ertebølle period[71,cf. 72]. Interestingly, rather stable $^{87}Sr/^{86}Sr$
isotope ratios throughout the Mesolithic indicate limited mobility, in agreement with the evidence
for genetic continuity reported here and modelled in previous work[73,74 Fig. 3], and/or dietary sources
from homogeneous environments.
589
590 The arrival of Neolithic farmer-related ancestry at c. 5,900 BP in Denmark resulted in a population
591 replacement with very limited genetic contribution from the local hunter-gatherers. The shift is
592 abrupt and brings changes in all the measured parameters. This is a clear case of demic diffusion,
593 which settles a long-standing debate concerning the neolithisation process in Denmark[15,56,75,76], at
594 least at a broader population level. The continuing use of coastal kitchen middens well into the

Neolithic[77,78] remains, however, an enigma, although this may represent sites where local remnants of Mesolithic groups survived in partly acculturated form, or it could be middens taken over by the newcomers. Concomitant shifts in both autosomal and uniparental genetic markers show that the migration by incoming farmers was not clearly sex-biased but more likely involved nuclear family units. Diet shifted abruptly to terrestrial sources evidenced by δ^{13}C values around -20 ‰ and δ^{15}N values around 10 ‰ in line with archaeological evidence that domesticated crops and animals were now providing the main supply of proteins (Supplementary Note 6). Isotope values remained stable at these levels throughout the following periods, although with somewhat greater variation after c. 4,500 BP. However, five Neolithic and Early Bronze Age individuals have δ^{13}C and δ^{15}N values indicating intake of high trophic marine food. This is most pronouncedly seen for NEO898 (Svinninge Vejle) who was one of the two aforementioned Danish Neolithic individuals displaying typical Swedish PWC hunter-gatherer ancestry. A higher variability in ^{87}Sr/^{86}Sr values can be seen with the start of the Neolithic and this continues in the later periods, which suggests that the Neolithic farmers in Denmark consumed food from more diverse landscapes and/or they were more mobile than the preceding hunter-gatherers (Supplementary Note 11). The Neolithic transition also marks a considerable rise in frequency of major effect alleles associated with light hair pigmentation[79], whereas polygenic score predictions for height are generally low throughout the first millennium of the Neolithic (Funnel Beaker epoch), echoing previous findings based on a smaller set of individuals[45,80].

We do not know how the Mesolithic Ertebølle population disappeared. Some may have been isolated in small geographical pockets of brief existence and/or adapted to a Neolithic lifestyle but without contributing much genetic ancestry to subsequent generations. The most recent individual in our Danish dataset with Mesolithic WHG ancestry is "Dragsholm Man" (NEO962), dated to 5,947-5,664 cal. BP (95%) and archaeologically assigned to the Neolithic Funnel Beaker farming culture based on his grave goods[81,82]. Our data confirms a typical Neolithic diet matching the cultural affinity but contrasting his WHG ancestry. Thus, Dragsholm Man represents a local person of Mesolithic ancestry who lived in the short Mesolithic-Neolithic transition period and adopted a Neolithic culture and diet. A similar case of very late Mesolithic WHG ancestry in Denmark was observed when analysing human DNA obtained from a piece of chewed birch pitch dated to 5,858–5,661 cal. BP (95%)[83].

The earliest example of Anatolian Neolithic ancestry in our Danish dataset is observed in a bog skeleton of a female from Viksø Mose (NEO601) dated to 5,896-5,718 cal. BP (95%) (and hence potentially contemporaneous with Dragsholm Man) whereas the most recent Danish individual showing Anatolian ancestry without any Steppe-related ancestry is NEO943 from Stenderup Hage, dated to 4,818-4,415 cal. BP (95%). Using Bayesian modelling we estimate the duration between the first appearance of Anatolian ancestry to the first appearance of Steppe-related ancestry in Denmark to be between 876 and 1100 years (95% probability interval, Supplementary Note 9) indicating that the typical Neolithic ancestry was dominant for less than 50 generations in Denmark. From this point onwards the steppe-ancestry was introduced, signalling the rise of the late Neolithic Corded Ware derived cultures in Denmark (i.e. Single Grave Culture), followed by the later Neolithic Dagger epoch and Bronze Age cultures. While this introduced a major new component in the Danish gene pool, it was not accompanied by apparent shifts in diet. Our complex trait predictions indicate an increase in "genetic height" occurring concomitant with the introduction of Steppe-related ancestry, which is consistent with Steppe individuals (e.g., Yamnaya) being genetically taller on average[45] and with previous results from other European regions[80,84].

These major population turnovers were accompanied by significant environmental changes, as apparent from high-resolution pollen diagrams from Lake Højby in Northwest Zealand

reconstructed using the Landscape Reconstruction Algorithm (LRA[85] (Supplementary Note 8).
While the LRA has previously been applied at low temporal resolution regional scale [e.g. 86,87], and at
local scale to Iron Age and later pollen diagrams [e.g. 88,89], this is the first time this quantitative
method is applied at local scale to a pollen record spanning the Mesolithic and Neolithic periods in
Denmark. Comparison with existing pollen records show that the land cover changes demonstrated
here reflect the general vegetation development in eastern Denmark, while the vegetation on the
sandier soils of western Jutland maintains a more open character throughout the sequence
(Supplementary Note 12). We find that during the Mesolithic (i.e. before c. 6,000 BP) the
vegetation was dominated by primary forest trees (*Tilia, Ulmus, Quercus, Fraxinus, Alnus* etc.).
The forest composition changed towards more secondary, early successional trees (*Betula and then*
Corylus) in the earliest Neolithic, but only a minor change in the relationship between forest and
open land is recorded. From c. 5,650 BP deforestation intensified, resulting in a very open
grassland-dominated landscape. This open phase was short-lived, and secondary forest expanded
from 5,500 to 5,000 BP, until another episode of forest clearance gave rise to an open landscape
during the last part of the Funnel Beaker epoch. We thus conclude that the agriculture practice was
characterised by repeated clearing of the forest with fire, followed by regrowth. This strategy
changed with the onset of the Single Grave Culture, when the forest increased again, but this time
dominated by primary forest trees, especially *Tilia* and *Ulmus*. This reflects the development of a
more permanent division of the landscape into open grazing areas and forests. In contrast, in
western Jutland this phase was characterised by large-scale opening of the landscape, presumably as
a result of human impact aimed at creating pastureland[90].

666
Finally, we investigated the fine-scale genetic structure in southern Scandinavia after the
introduction of Steppe-related ancestry using a temporal transect of 38 Late Neolithic and Early
Bronze Age Danish and southern Swedish individuals. Although the overall population genomic
signatures suggest genetic stability, patterns of pairwise IBD-sharing and Y-chromosome
haplogroup distributions indicate at least three distinct ancestry phases during a ~1,000-year time
span: i) An early stage between ~4,600 BP and 4,300 BP, where Scandinavians cluster with early
CWC individuals from Eastern Europe, rich in Steppe-related ancestry and males with an R1a Y-
chromosomal haplotype (Extended Data Fig. 8A, B); ii) an intermediate stage until c. 3,800 BP,
where they cluster with central and western Europeans dominated by males with distinct sub-
lineages of R1b-L51 (Extended Data Fig. 8C, D; Supplementary Note 3b) and includes Danish
individuals from Borreby (NEO735, 737) and Madesø (NEO752) with distinct cranial features
(Supplementary Note 6); and iii) a final stage from c. 3,800 BP onwards, where a distinct cluster of
Scandinavian individuals dominated by males with I1 Y-haplogroups appears (Extended Data Fig.
8E). Using individuals associated with this cluster (Scandinavia_4000BP_3000BP) as sources in
supervised ancestry modelling (see "postBA", Extended Data Fig. 4), we find that it forms the
predominant source for later Iron- and Viking Age Scandinavians, as well as ancient European
groups outside Scandinavia who have a documented Scandinavian or Germanic association (e.g.,
Anglo-Saxons, Goths; Extended Data Fig. 4). Y-chromosome haplogroup I1 is one of the dominant
haplogroups in present-day Scandinavians,s, and we document its earliest occurrence in a ~4,000-
year-old individual from Falköping in southern Sweden (NEO220). The rapid expansion of this
haplogroup and associated genome-wide ancestry in the early Nordic Bronze Age indicates a
considerable reproductive advantage of individuals associated with this cluster over the preceding
groups across large parts of Scandinavia.
690
691

693   In contrast to the significant number of ancient hunter-gatherer genomes from western Eurasia
694   studied to date, genomic data from hunter-gatherers east of the Urals remain sparse. These regions
695   are characterised by an early introduction of pottery from areas further east and are inhabited by
696   complex hunter-gatherer-fisher societies with permanent and sometimes fortified settlements
697   (Supplementary Note 5; [91]).
698
Here, we substantially expand the knowledge on ancient Stone Age populations of this region by
reporting new genomic data from 38 individuals, 28 of which date to pottery-associated hunter-
gatherer contexts [e.g. 92] between 8,300-5,000 BP (Supplementary Table II).The majority of these
genomes form a previously only sparsely sampled [48,93] "Neolithic Steppe" cline spanning the
Siberian Forest Steppe zones of the Irtysh, Ishim, Ob, and Yenisei River basins to the Lake Baikal
region (Fig. 1C; Extended Data Fig. 2A, 3E). Supervised admixture modelling (using the "deep" set
of ancestry sources) revealed contributions from three major sources in these hunter gatherers from
east of Urals: early West Siberian hunter-gatherer ancestry (SteppeC_8300BP_7000BP) dominated
in the western Forest Steppe; Northeast Asian hunter-gatherer ancestry (Amur_7500BP) was
highest at Lake Baikal; and Paleosiberian ancestry (SiberiaNE_9800BP) was observed in a cline of
decreasing proportions from northern Lake Baikal westwards across the Forest Steppe (Extended
Data Fig. 4, 9). [93]
711
We used these Neolithic hunter-gatherer clusters ("postNeol" ancestry source set, Extended Data
Fig. 4) as putative source groups in more proximal admixture modelling to investigate the
spatiotemporal dynamics of ancestry compositions across the Steppe and Lake Baikal after the
Neolithic period. We replicate previously reported evidence for a genetic shift towards higher
Forest Steppe hunter-gatherer ancestry (SteppeCE_7000BP_3600BP) in late Neolithic and early
Bronze Age individuals (LNBA) at Lake Baikal [93,94]. However, ancestry related to this cluster is
already observed at ~7,000 BP in herein-reported Neolithic hunter-gatherer individuals both at Lake
Baikal (NEO199, NEO200), and along the Angara river to the north (NEO843). Both male
individuals at Lake Baikal belonged to Y-chromosome haplogroup Q1, characteristic of the later
LNBA groups in the same region. (Extended Data Fig. 3, 6A). Together with an estimated date of
admixture of ~6,000 BP for the LNBA groups, these results suggest gene flow between hunter-
gatherers of Lake Baikal and the south Siberian forest steppe regions already during the early
Neolithic. This is consistent with archaeological interpretations of contact. In this region, bifacially
flaked tools first appeared near Baikal [95] from where the technique spread far to the west. We find
its reminiscences in Late Neolithic archaeological complexes (Shiderty 3, Borly, Sharbakty 1, Ust-
Narym, etc.) in Northern and Eastern Kazakhstan, around 6,500-6,000 BP [96,97]. Our herein-reported
genomes also shed light on the genetic origins of the early Bronze Age Okunevo culture in the
Minusinsk Basin in Southern Siberia. In contrast to previous results, we find no evidence for Lake
Baikal hunter-gatherer ancestry in the Okunevo[93,94], suggesting that they instead originate from a
three-way mixture of two different genetic clusters of Siberian forest steppe hunter-gatherers and
Steppe-related ancestry (Extended Data Fig. 4D). We date the admixture with Steppe-related
ancestry to ~4,600 BP, consistent with gene flow from peoples of the Afanasievo culture that
existed near Altai and Minusinsk Basin during the early eastwards' expansion of Yamnaya-related
groups [20,94].
736
From around 3,700 BP, individuals across the Steppe and Lake Baikal regions display markedly
different ancestry profiles (Fig. 3; Extended Data Fig. 4D, 9). We document a sharp increase in
non-local ancestries, with only limited ancestry contributions from local hunter-gatherers. The early
stages of this transition are characterised by influx of Yamnaya-related ancestry, which decays over

time from its peak of ~70% in the earliest individuals. Similar to the dynamics in western Eurasia,
Yamnaya-related ancestry is here correlated with late Neolithic GAC-related farmer ancestry
(Poland_5000BP_4700BP; Extended Data Fig. 9G), recapitulating the previously documented
eastward expansion of admixed Western Steppe pastoralists from the Sintashta and Andronovo
complexes during the Bronze Age[20,48,98]. However, GAC-related ancestry is notably absent in
individuals of the Okunevo culture, providing further support for two distinct eastward migrations
of Western Steppe pastoralists during the early (Yamnaya) and later (Sintashta, Andronovo) Bronze
Age. The later stages of the transition are characterised by increasing Central Asian
(Turkmenistan_7000 BP_5000BP) and Northeast Asian-related (Amur_7500BP) ancestry
components (Extended Data Fig. 9G). Together, these results show that deeply structured hunter-
gatherer ancestry dominated the eastern Eurasian Steppe substantially longer than in western
Eurasia, before successive waves of population expansions swept across the Steppe within the last
4,000 years, including a large-scale introduction of domesticated horse lineages concomitant with
new equestrian equipment and spoke-wheeled chariotry [20,48,98,99].
755

Genetic legacy of Stone Age Europeans

To investigate the distribution of Stone Age and Early Bronze Age ancestry components in modern
populations, we used ChromoPainter [100] to "paint" the chromosomes of individuals in the UK
Biobank (https://www.ukbiobank.ac.uk) using a panel of 10 ancient donor populations
(Supplementary Note 3h). Painting was done following the pipeline of Margaryan et al. [101] based on
GLOBETROTTER [102], and admixture proportions were estimated using Non-Negative Least
squares. Haplotypes in the modern genomes are assigned to the genetically closest ancient
population as measured by meiosis events, which favours more recent matches in time. Therefore,
ancestry proportions assigned to the oldest groups (e.g. WHG) should be interpreted as an excess of
this ancestry, which cannot be explained by simply travelling through more recent ancient
populations up to present times.
767

768 First, we selected non-British individuals from the UK Biobank if their country of birth was
769 European, African, or Asian. Because many of these individuals are admixed or British, we set up a
770 pipeline (Supplementary Note 3g) to select individuals of a typical ancestral background for each
771 country. This resulted in 24,511 individuals from 126 countries, who were then chromosome
772 painted to assess the average admixture proportions for each ancestry per country.
773

The various hunter-gatherer ancestries are not homogeneously distributed amongst modern
populations (Fig. 5). WHG-related ancestry is highest in present-day individuals from the Baltic
States, Belarus, Poland, and Russia; EHG-related ancestry is highest in Mongolia, Finland, Estonia
and Central Asia; and CHG-related ancestry is maximised in countries east of the Caucasus, in
Pakistan, India, Afghanistan and Iran, in accordance with previous results [103]. The CHG-related
ancestry likely reflects both Caucasus hunter-gatherer and Iranian Neolithic signals, explaining the
relatively high levels in south Asia [104]. Consistent with expectations [105,106], Neolithic Anatolian-
related farmer ancestry is concentrated around the Mediterranean basin, with high levels in southern
Europe, the Near East, and North Africa, including the Horn of Africa, but is less frequent in
Northern Europe. This is in direct contrast to the Steppe-related ancestry, which is found in high
levels in northern Europe, peaking in Ireland, Iceland, Norway, and Sweden, but decreases further
south. There is also evidence for its spread into southern Asia. Overall, these results refine global
patterns of spatial distributions of ancient ancestries amongst modern populations.
787

The availability of a large number of modern genomes (n=408,884) from self-identified "white"
British individuals who share similar PCA backgrounds [107] allowed us to further examine the
distribution of ancient ancestries at high resolution in Britain (Supplementary Note 3h). Although
regional ancestry distributions differ by only a few percent, we find clear evidence of geographical
heterogeneity across the United Kingdom as visualised by assigning individuals to their birth
county and averaging ancestry proportions per county (Fig. 5, inset boxes). The proportion of
Neolithic farmer ancestry is highest in southern and eastern England today and lower in Scotland,
Wales, and Cornwall. Steppe-related ancestry is inversely distributed, peaking in the Outer
Hebrides and Ireland, a pattern only previously described for Scotland [108]. This regional pattern was
already evident in the Pre-Roman Iron Age and persists to the present day even though immigrating
Anglo-Saxons had relatively less Neolithic farmer ancestry than the Iron-Age population of
southwest Briton (Extended Data Fig. 4). Although this Neolithic farmer/steppe-related dichotomy
mirrors the modern 'Anglo-Saxon'/'Celtic' ethnic divide, its origins are older, resulting from
continuous migration from a continental population relatively enhanced in Neolithic farmer
ancestry, starting as early as the Late Bronze Age [109]. By measuring haplotypes from these
ancestries in modern individuals, we are able to show that these patterns differentiate Wales and
Cornwall as well as Scotland from England. We also found higher levels of WHG-related ancestry
in central and Northern England. These results demonstrate clear ancestry differences within an
'ethnic group' (white British) traditionally considered relatively homogenous, which highlights the
need to account for subtle population structure when using resources such as the UK Biobank
genomes.
809

810
811 **Fig 5. The genetic legacy of Stone Age ancestry in modern populations.**
812 From top left clockwise: Neolithic Farmer, Yamnaya, Caucasus hunter-gatherer, Eastern hunter-gatherer, Western
813 hunter-gatherer. Panels show average admixture proportion in modern individuals per country estimated using NNLS
814 (large maps), average per county within the UK (top left insert), and PCA (PC2 vs PC1) of admixture proportions, with
815 the top 10 highest countries by admixture fraction labelled and PCA loadings for that ancestry.
816

Sociocultural insights

We used patterns of pairwise IBD sharing between individuals and runs of homozygosity (ROH)
within individuals (measured as the fraction of the genome within a run of homozygosity f(ROH))
to examine our data for temporal shifts in relatedness within genetic clusters. Both measures show
clear trends of a reduction of within-cluster relatedness over time, in both western and eastern
Eurasia (Fig. 6). This pattern is consistent with a scenario of increasing effective population sizes
during this period [110]. Nevertheless, we observe notable differences in temporal relatedness patterns
between western and eastern Eurasia, mirroring the wider difference in population dynamics
discussed above. In the west, within-group relatedness changes substantially during the Neolithic
transition (~9,000 to ~6,000 BP), where clusters of Neolithic farmer-associated individuals show
overall reduced IBD sharing and f(ROH) compared to clusters of HG-associated individuals (Fig.
6A,C). In the east, genetic relatedness remains high until ~4,000 BP, consistent with a much longer
persistence of smaller localised hunter-gatherer groups (Fig. 6B,D).
830

Next, we examined the data for evidence of recent parental relatedness, by identifying individuals
harbouring a large fraction of their genomes (> 50cM) in long (>20cM) ROH segments [111]. We only
detect 39 such individuals out of a total sample of 1,540 imputed ancient genomes (Fig. 6E), in line
with recent results indicating that close kin mating was not common in human prehistory
[41,103,111,112]. With the exception of eight ancient American individuals from the San Nicolas Islands
in California [113], no obviously discernible spatiotemporal or cultural clustering was observed among
the individuals with recent parental relatedness. Interestingly, an ~1,700-year-old Sarmatian
individual from Temyaysovo (tem003) [114] was found homozygous for almost the entirety of
chromosome 2, but without evidence of ROHs elsewhere in the genome, suggesting an ancient case
of uniparental disomy. Among several noteworthy familial relationships (see Supplementary Fig.
S3c.2), we report a Mesolithic father/son burial at Ertebølle (NEO568/NEO569), as well as a
Mesolithic mother/daughter burial at Dragsholm (NEO732/NEO733).
843
844

845
Fig 6. Patterns of co-ancestry. (A)-(D) Panels show within-cluster genetic relatedness over time, measured either as
the total length of genomic segments shared IBD between individuals (A, B) or the proportion of individual genomes
found in a run of homozygosity f(ROH) (C,D). Results for both measures are shown separately for individuals from
western (A, C) or eastern Eurasia (B, D). Small grey dots indicate estimates for individual pairs (A, B) or individuals
(C, D), with larger coloured symbols indicating median values within genetic clusters. (E) Distribution of ROH lengths
for 39 individuals with evidence for recent parental relatedness (>50 cM total in ROHs > 20 cM).
852
853
854 *Pathogenic structural variants in ancient vs. modern-day humans*
855
Rare, recurrent copy-number variants (CNVs) are known to cause neurodevelopmental disorders
and are associated with a range of psychiatric and physical traits with variable expressivity and
incomplete penetrance[115,116]. To understand the prevalence of pathogenic structural variants over
time we examined 50 genomic regions susceptible to recurrent CNV, known to be the most
prevalent drivers of human developmental pathologies[117]. The analysis included 1442 ancient
imputed genomes passing quality control for CNV analysis (Supplementary Note 4i) and 1093
modern human genomes for comparison [118,119]. We identified CNVs in ancient individuals at ten
loci using a read-depth based approach and digital Comparative Genomic Hybridization [120]
(Supplementary Table S4i.1; Supplementary Figs. S4i.1-S41.20). Although most of the observed
CNVs (including duplications at 15q11.2 and *CHRNA7*, and CNVs spanning parts of the TAR locus
and 22q11.2 distal) have not been unambiguously associated with disease in large studies, the
identified CNVs include deletions and duplications that have been associated with developmental
delay, dysmorphic features, and neuropsychiatric abnormalities such as autism (most notably at
1q21.1, 3q29, 16p12.1 and the DiGeorge/VCFS locus, but also deletions at 15q11.2 and

duplications at 16p13.11). The individual harbouring the 16p13.1 deletion, RISE586 [20], a 4,000 BP
woman aged 20-30 from the Únětice culture (modern day Czech Republic), had almost complete
skeletal remains, which allowed us to test for the presence of various skeletal abnormalities
associated with the 16p13.11 microdeletion [121]. RISE586 exhibited a hypoplastic tooth,
spondylolysis of the L5 vertebrae, incomplete coalescence of the S1 sacral bone, among other
minor skeletal phenotypes. The skeletal phenotypes observed in this individual are relatively
common (~10%) in European populations and are not specific to 16p13.1 thus do not indicate
strong penetrance of this mutation in RISE586 [122–125]. However, these results do highlight our
ability to link putatively pathogenic genotypes to phenotypes in ancient individuals. Overall, the
carrier frequency in the ancient individuals is similar to that reported in the UK Biobank genomes
(1.25% vs 1.6% at 15q11.2 and *CHRNA7* combined, and 0.8% vs 1.1% across the remaining loci
combined) [126]. These results suggest that large, recurrent CNVs that can lead to several pathologies
were present at similar frequencies in the ancient and modern populations included in this study.
883
884   *Ancestry-stratified patterns of natural selection in the last 13,000 years*
885
The Neolithic transition led to a fundamental change in lifestyle, diet and exposure to pathogens
that imposed drastically new selection pressures on human populations. To detect genetic candidate
targets of selection, we used a set of 1,015 imputed ancient genomes from West Eurasia that were
fitted to a four-way admixture model of demographic history in this region (Supplementary Note 3i)
and identified phenotype-associated variants with evidence for directional selection over the last
13,000 years, with a special focus on the Neolithic transition (Supplementary Note 4a). We adapted
CLUES [127] to model time-series data (Supplementary Note 4a) and used it to infer allele frequency
trajectories and selection coefficients for 33,323 quality-controlled phenotype-associated variants
ascertained from the GWAS Catalogue [128]. An equal number of putatively neutral, frequency-paired
variants were used as a control set. To control for possible confounders, we built a causal model to
distinguish direct effects of age on allele frequency from indirect effects mediated by read depth,
read length, and/or error rates (Supplementary Note 4b), and developed a mapping bias test used to
evaluate systematic differences between data from ancient and present-day populations
(Supplementary Note 4a). Because admixture between groups with differing allele frequencies can
confound interpretation of allele frequency changes through time, we also applied a novel
chromosome painting technique, based on inference of a sample's nearest neighbours in the
marginal trees of a tree sequence (Supplementary Note 3i). This allowed us to accurately assign
ancestral path labels to haplotypes found in both ancient and present-day individuals. By
conditioning on these haplotype path labels, we could infer selection trajectories while controlling
for changes in admixture proportions through time (Supplementary Note 4a).
906
Our analysis identified no genome-wide significant (p < 5e-8) selective sweeps when using
genomes from present-day individuals alone (1000 Genomes Project populations GBR, FIN and
TSI), although trait-associated variants were enriched for signatures of selection compared to the
control group (p < 2.2e-16, Wilcoxon signed-rank test). In contrast, when using imputed aDNA
genotype probabilities, we identified 11 genome-wide significant selective sweeps in the GWAS
variants, and none in the control group, consistent with selection acting on trait-associated variants
(Supplementary Note 4a, Supplementary Figs. S4a.4 to S4a.14). However, when conditioned on
one of our four ancestral histories—genomic regions arriving in present day genomes through
Western hunter-gatherers (WHG), Eastern hunter-gatherers (EHG), Caucasus hunter-gatherers
(CHG) or Anatolian farmers (ANA)—we identified 21 genome-wide significant selection peaks
(including the 11 from the pan-ancestry analysis) (Fig. 7). This suggests that admixture between
ancestral populations has masked evidence of selection at many trait associated loci in modern
populations.

Fig 7. Genome-wide selection scan for trait associated variants. A) Manhattan plot of p-values from selection scan
with CLUES, based on a time-series of imputed aDNA genotype probabilities. Twenty-one genome-wide significant
selection peaks highlighted in grey and labelled with the most significant gene within each locus. Within each sweep,
SNPs are positioned on the y-axis and coloured by their most significant marginal ancestry. Outside of the sweeps,
SNPs show p-values from the pan-ancestry analysis and are coloured grey. Red dotted lines indicate genome-wide
significance (p < 5e-8), while the grey dotted line shows the Bonferroni significance threshold, corrected for the number
of tests (p < 1.35e-6). B) Detailed plots for three genome-wide significant sweep loci: (i) MCM6, lactase persistence;
(ii) SLC45A2, skin pigmentation; and (iii) FADS2, lipid metabolism. Rows show results for the pan-ancestry analysis
(ALL) plus the four marginal ancestries: Western hunter-gatherers (WHG), Eastern hunter-gatherers (EHG), Caucasus
hunter-gatherers (CHG) and Anatolian farmers (ANA). The first column of each loci shows zoomed Manhattan plots of
the p-values for each ancestry (significant SNPs sized by their selection coefficients), and column two shows allele

933 trajectories for the top SNPs across all ancestries (grey shading for the marginal ancestries indicates approximate
934 temporal extent of the pre-admixture population).

935
936 *Selection on diet-associated loci*

937
We find strong changes in selection associated with lactose digestion after the introduction of
farming, but prior to the expansion of the Yamnaya pastoralists into Europe around 5,000 years ago
[20,21], settling controversies regarding the timing of this selection [129–132]. The strongest overall signal
of selection in the pan-ancestry analysis is observed at the *MCM6 / LCT* locus (rs4988235; p=9.86e-
31; s=0.020), where the derived allele results in lactase persistence [133,134] (Supplementary Note 4a).
The trajectory inferred from the pan-ancestry analysis indicates that the lactase persistence allele
began increasing in frequency only c. 7,000 years ago, and has continued to increase up to present
times (Fig. 7). Our ancestry-stratified analysis shows, however, that selection at the MCM6/LCT
locus is much more complex than previously thought. In the pan-ancestry analysis, this sweep is led
by the lactase persistence SNP rs4988235, whereas in the ancestry-stratified analysis, this signal is
primarily driven by sweeps in two of the ancestral backgrounds (EHG and CHG), each of which
differ in their most significant SNPs (Fig. 7). Conversely, in the WHG background, we find no
evidence for selection at rs4988235, but strong selection at rs12465802 within the last c. 2,000
years. Overall, our results suggest that there were multiple, asynchronous selective sweeps in this
genomic region in recent human history, and possibly targeting different loci.

953
We also find strong selection in the FADS gene cluster — *FADS1* (rs174546; p=2.65e-10; s=0.013)
and *FADS2* (rs174581; p=1.87e-10; s=0.013) — which are associated with fatty acid metabolism
and known to respond to changes in diet from a more/less vegetarian to a more/less carnivorous diet
[135–140]. In contrast to previous results [138–140], we find that much of the selection associated with a
more vegetarian diet occurred in Neolithic populations before they arrived in Europe, but then
continued during the Neolithic (Fig. 7). The strong signal of selection in this region in the pan-
ancestry analysis is driven primarily by a sweep occurring on the EHG, CHG and ANA haplotypic
backgrounds (Fig. 7). Interestingly, we find no evidence for selection at this locus in the WHG
background, and most of the allele frequency rise in the EHG background occurs after their
admixture with CHG (around 8ka, [141]), within whom the selected alleles were already close to
present-day frequencies. This suggests that the selected alleles may already have existed at
substantial frequencies in early farmer populations in the Middle East and among Caucasus Hunter
gatherers (associated with the ANA and CHG and backgrounds, respectively) and were subject to
continued selection as eastern groups moved northwards and westwards during the late Neolithic
and Bronze Age periods.

969
When specifically comparing selection signatures differentiating ancient hunter-gatherer and farmer
populations [142], we also observe a large number of regions associated with lipid and sugar
metabolism, and various metabolic disorders (Supplementary Note 4e). These include, for example,
a region in chromosome 22 containing *PATZ1*, which regulates the expression of *FADS1*, and
MORC2, which plays an important role in cellular lipid metabolism [143–145]. Another region in
chromosome 3 overlaps with *GPR15,* which is both related to immune tolerance and to intestinal
homeostasis [146–148]. Finally, in chromosome 18, we recover a selection candidate region spanning
SMAD7, which is associated with inflammatory bowel diseases such as Crohn's disease [149–151].
Taken together these results suggest that the transition to agriculture imposed a substantial amount
of selection for humans to adapt to our new diet and that some diseases observed today in modern
societies can likely be understood as a consequence of this selection.

Selection on immunity-associated variants

In addition to diet-related selection, we observe selection in several loci associated with immunity/defence functions and with autoimmune disease (Supplementary Note 4a). Some of these selection events occurred earlier than previously claimed and are likely associated with the transition to agriculture and may help explain the high prevalence of autoimmune diseases today. Most notably, we detect a 33 megabase (Mb) wide selection sweep signal in chromosome 6 (chr6:19.1–50.9 Mb), spanning the human leukocyte antigen (HLA) region (Supplementary Note 4a). The selection trajectories of the variants within this locus support multiple independent sweeps, occurring at different times and with differing intensities. The strongest signal of selection at this locus in the pan-ancestry analysis is at an intergenic variant, located between *HLA-A* and *HLA-W* (rs7747253; p=8.86e-17; s=-0.018), associated with heel bone mineral density [152], the derived allele of which rapidly reduced in frequency, beginning c. 8,000 years ago (Extended Data Fig. 10). In contrast, the signal of selection at *C2* (rs9267677; p= 9.82e-14; s= 0.04463), also found within this sweep, and associated with educational attainment [153], shows a gradual increase in frequency beginning c. 4,000 years ago, before rising more rapidly c. 1,000 years ago. This highlights the complex temporal dynamics of selection at the HLA locus, which not only plays a role in the regulation of the immune system, but also has association with many other non-immune-related phenotypes. The high pleiotropy in this region makes it difficult to determine which selection pressures may have driven these increases in frequencies at different periods of time. However, profound shifts in lifestyle in Eurasian populations during the Holocene, including a change in diet and closer contact with domestic animals, combined with higher mobility and increasing population sizes, are likely drivers for strong selection on loci involved in immune response.

We also identified selection signals at the *SLC22A4* (rs35260072; p=1.15e-10; s=0.018) locus, associated with increased itch intensity from mosquito bites [154], and find that the derived variant has been steadily rising in frequency since c. 9,000 years ago (Extended Data Fig. 11). However, in the same *SLC22A4* candidate region as rs35260072, we find that the frequency of the previously reported SNP rs1050152 plateaued c. 1,500 years ago, contrary to previous reports suggesting a recent rise in frequency [45]. Similarly, we detect selection at the *HECTD4* (rs11066188; p=3.02e-16; s=0.020) and *ATXN2* (rs653178; p=1.92e-15; s=0.019) locus, associated with celiac disease and rheumatoid arthritis [155], which has been rising in frequency for c. 9,000 years (Extended Data Fig. 12), also contrary to previous reports of a more recent rise in frequency [45]. Thus, several disease-associated loci previously thought to be the result of recent adaptation may have been subject to selection for a longer period of time.

Selection on the 17q12.13 locus

We further detect signs of strong selection in a 12 Mb sweep in chromosome 17 (chr17:36.1–48.1 Mb), spanning a locus on 17q21.3 implicated in neurodegenerative and developmental disorders (Supplementary Note 4a). The locus includes an inversion and other structural polymorphisms with indications of a recent positive selection sweep in some human populations [156,157]. Specifically, partial duplications of the *KANSL1* gene likely occurred independently on the inverted (H2) and non-inverted (H1) haplotypes (Fig. 8B) and both are found in high frequencies (15-25%) among current European and Middle Eastern populations but are much rarer in Sub-Saharan African and East Asian populations. We used both SNP genotypes and WGS read depth information to determine inversion (H1/H2) and *KANSL1* duplication (d) status in the ancient individuals studied here (see Supplementary Note 4g).

The H2 haplotype is observed in two of three previously published genomes[158] of Anatolian
aceramic Neolithic individuals (Bon001 and Bon004) from around 10,000 BP, but data were
insufficient to identify *KANSL1* duplications. The oldest evidence for *KANSL1* duplications is
observed in an Iranian early Neolithic individual (AH1 from 9,900 BP[2]) followed by two Georgian
Mesolithic individuals (NEO281 from 9,724 BP and KK1 [6] from 9,720 BP) all of whom are
heterozygous for the inversion and carry the inverted duplication. The *KANSL1* duplications are
also detected in two Russian Neolithic individuals: NEO560 from 7,919 BP (H1d) and NEO212
from 7,390 BP (H2d). With both H1d and H2d having spread to large parts of Europe with
Anatolian Neolithic Farmer ancestry, their frequency seems unchanged in most of Europe as
Steppe-related ancestry becomes dominant in large parts of the subcontinent (Extended Data Fig.
8D). The fact that both H1d and H2d are found in apparently high frequencies in both early
Anatolian Farmers and the earliest Yamnaya/Steppe-related ancestry groups suggests that any
selective sweep acting on the H1d and H2d variants would probably have occurred in populations
ancestral to both.
1045
We note that the strongest signal of selection observed in this locus is at *MAPT* (rs4792897;
p=4.65e-10; s=0.03 (Fig. 8A; Supplementary Note 4a), which codes for the tau protein [159] and is
involved in a number of neurodegenerative disorders, including Alzheimer's disease and
Parkinson's disease [160–164]. However, the region is also enriched for evidence of reference bias in
our imputed dataset—especially around the *KANSL1* gene—due to complex structural
polymorphisms (Supplementary Note 4i).

1052

Fig 8. Selection at the *MAPT* / 17q21.31 inversion locus. A) Results for the pan-ancestry analysis (ALL) plus the four marginal ancestries: Western hunter-gatherers (WHG), Eastern hunter-gatherers (EHG), Caucasus hunter-gatherers (CHG) and Anatolian farmers (ANA). Grey shading for the marginal ancestries indicates approximate temporal extent of the pre-admixture population. B) Haplotypes of the 17q21.31 locus: the ancestral (non-inverted) H1 17q21.31 and the inverted H2 haplotype. Duplications of the *KANSL1* gene have occurred independently on both lineages yielding H1D and H2D haplotypes. C) Frequency of the 17q21.31 inversion and duplication haplotypes across modern-day global populations (Human Genome Diversity Project [119]). D) Change in the frequency of the 17q21.31 inversion haplotype through time.

Selection on pigmentation-associated variants

Our results identify strong selection for lighter skin pigmentation in groups moving northwards and westwards, in agreement with the hypothesis that selection is caused by reduced UV exposure and resulting vitamin D deficiency. We find that the most strongly selected alleles reached near-fixation several thousand years ago, suggesting that this was not associated with recent sexual selection as proposed [165,166] (Supplementary Note 4a).

In the pan-ancestry analysis we detect strong selection at the *SLC45A2* locus (rs35395; p=4.13e-23; s=0.022) locus [167,168], with the selected allele (responsible for lighter skin), increasing in frequency from c. 13,000 years ago, until plateauing c. 2,000 years ago (Fig. 7). The dominating hypothesis is that high melanin levels in the skin are important in equatorial regions owing to its protection

1074 against UV radiation, whereas lighter skin has been selected for at higher latitudes (where UV
1075 radiation is less intense) because some UV penetration is required for cutaneous synthesis of
1076 vitamin D [169,170]. Our findings confirm pigmentation alleles as major targets of selection during the
1077 Holocene [45,171,172] particularly on a small proportion of loci with large effect sizes [168].
1078
Additionally, our results provide unprecedentedly detailed information about the duration and
geographic spread of these processes (Fig. 7) suggesting that an allele associated with lighter skin
was selected for repeatedly, probably as a consequence of similar environmental pressures
occurring at different times in different regions. In the ancestry-stratified analysis, all marginal
ancestries show broad agreement at the *SLC45A2* locus (Fig. 7) but differ in the timing of their
frequency shifts. The ANA ancestry background shows the earliest evidence for selection, followed
by EHG and WHG around c. 10,000 years ago, and CHG c. 2,000 years later. In all ancestry
backgrounds except WHG, the selected haplotypes reach near fixation by c. 3,000 years ago, whilst
the WHG haplotype background contains the majority of ancestral alleles still segregating in
present-day Europeans. This finding suggests that selection on this allele was much weaker in
ancient western hunter-gatherer groups during the Holocene compared to elsewhere. We also detect
strong selection at the *SLC24A5* (rs1426654; p=6.45e-09; s=0.019) which is also associated with
skin pigmentation [167,173]. At this locus, the selected allele increased in frequency even earlier than
SLC45A2 and reached near fixation c. 3,500 years ago (Supplementary Note 4a). Selection on this
locus thus seems to have occurred early on in groups that were moving northwards and westwards,
and only later in the Western hunter-gatherer background after these groups encountered and
admixed with the incoming populations.
1096
1097 *Selection among major axes of ancient population variation*
1098
Beyond patterns of genetic change at the Mesolithic-Neolithic transition, much genetic variability
observed today reflects high genetic differentiation in the hunter-gatherer groups that eventually
contributed to modern European genetic diversity [142]. Indeed, a substantial number of loci
associated with cardiovascular disease, metabolism and lifestyle diseases trace their genetic
variability prior to the Neolithic transition, to ancient differential selection in ancestry groups
occupying different parts of the Eurasian continent (Supplementary Note 4d). These may represent
selection episodes that preceded the admixture events described above, and led to differentiation
between ancient hunter-gatherer groups in the late Pleistocene and early Holocene. One of these
overlaps with the *SLC24A3* gene which is a salt sensitivity gene significantly expressed in obese
individuals [174,175]. Another spans *ROPN1* and *KALRN*, two genes involved in vascular disorders [176–
178]. A further region contains *SLC35F3*, which codes for a thiamine transport and has been
associated with hypertension in a Han Chinese cohort [179,180]. Finally, there is a candidate region
containing several genes (*CH25H, FAS*) associated with obesity and lipid metabolism [181–183] and
another peak with several genes (*ASXL2, RAB10, HADHA, GPR113*) involved in glucose
homeostasis and fatty acid metabolism [184–193]. These loci predominantly reflect ancient patterns of
extreme differentiation between Eastern and Western Eurasian genomes, and may be candidates for
selection after the separation of the Pleistocene populations that occupied different environments
across the continent (roughly 45,000 years ago [103]).
1117
1118 *Genetic trait reconstruction and the phenotypic legacy of ancient Europeans*
1119
1120 When comparing modern European genomes in the UK Biobank to ancient Europeans, we find
1121 strong differentiation at certain sets of trait-associated variants, and differential contribution of
1122 different ancestry groups to various traits. We reconstructed polygenic scores for phenotypes in
1123 ancient individuals, using effect size estimates obtained from GWASs performed using the

>400,000 UK Biobank genomes [107] (http://www.nealelab.is/uk-biobank) and looked for overdispersion among these scores across ancient populations, beyond what would be expected under a null model of genetic drift [194] (Supplementary Note 4c). We stress that polygenic scores and Q_X statistic may both be affected by population stratification, so these results should be interpreted with caution [195–198]. The most significantly overdispersed scores are for variants associated with pigmentation, anthropometric differences and disorders related to diet and sugar levels, including diabetes (Fig. 9). We also find psychological trait scores with evidence for overdispersion related to mood instability and irritability, with Western Hunter-gatherers generally showing smaller genetic scores for these traits than Neolithic Farmers. Intriguingly, we find highly inconsistent predictions of height based on polygenic scores in western hunter-gatherer and Siberian groups computed using effect sizes estimated from two different - yet largely overlapping - GWAS cohorts (Supplementary Note 4c), highlighting how sensitive polygenic score predictions are to the choice of cohort, particularly when ancient populations are genetically divergent from the reference GWAS cohort [198]. Taking this into account, we do observe that the Eastern hunter-gatherer and individuals associated with the Yamnaya culture have consistently high genetic values for height, which in turn contribute to stature increases in Bronze Age Europe, relative to the earlier Neolithic populations [45,80,199].

We performed an additional analysis to examine the data for strong alignments between axes of trait-association [200] and ancestry gradients, rather than relying on particular choices for population clusters (Supplementary Note 4e). Along the population structure axis separating ancient East Asian and Siberian genomes from Steppe and Western European genomes (Fig. 1), we find significant correlations with trait-association components related to impedance, body measurements, blood measurements, eye measurement and skin disorders. Along the axis separating Mesolithic hunter-gatherers from Anatolian and Neolithic farmer individuals, we find significant correlations with trait-association components related to skin disorders, diet and lifestyle traits, mental health status, and spirometry-related traits (Fig. 9). Our findings show that these phenotypes were genetically different among ancient groups with very different lifestyles. However, we note that the realised value of these traits is highly dependent on environmental factors and gene-environment interactions, which we do not model in this analysis.

1155

Fig. 9 A. First two principal components of a PCA of ancient Western Eurasian samples, coloured by height polygenic scores. B. Top 6 UK Biobank trait-association components from DeGAs that have the highest correlation (P < 5e-3) with the principal component separating ancient farmers from hunter-gatherer populations (PC1 in Panel A). C. Top UK Biobank traits with highest overdispersion in polygenic scores among ancient populations, out of a total of 320 sets of trait-associated SNPs tested ($Q_X > 79.4$, $P < 0.05/320$).

In addition to the above reconstructions of genetic traits among the ancient individuals, we also estimated the contribution from different ancestral populations (EHG, CHG, WHG, Yamnaya and Anatolian farmer) to variation in polygenic phenotypes in present-day individuals, leveraging the exceptional resolution offered by the UK Biobank genomes [107] to investigate this. We calculated ancestry-specific polygenic risk scores based on the chromosome painting of the >400,000 UKB genomes (Supplementary Note 4h); this allowed us to identify if any of the ancient ancestry components were over-represented in modern UK populations at loci significantly associated with a given trait, and also avoids exporting risk scores over space and time. Working with large numbers of imputed ancient genomes provides high statistical power to use ancient populations as "ancestral sources". We focused on phenotypes whose polygenic scores were significantly over-dispersed in the ancient populations (Supplementary Note 4c), as well as a single high effect variant, ApoE4, known to be a significant risk factor in Alzheimer's Disease ([201,202]). We emphasise that this approach makes no reference to ancient phenotypes but describes how these ancestries contributed to the modern genetic landscape. In light of the ancestry gradients within the British Isles and Eurasia (Fig. 5), these results support the hypothesis that ancestry-mediated geographic variation in disease risks and phenotypes is commonplace. It points to a way forward for disentangling how ancestry contributed to differences in risk of genetic disease – including metabolic and mental health disorders – between present-day populations.

1180
Taken together, these analyses help to settle the famous discussion of selection in Europe relating to
height [45,80,203]. The finding that steppe individuals have consistently high genetic values for height
(Supplementary Note 4c), is mirrored by the UK Biobank results, which find that the 'Steppe'
ancestral components (Yamnaya/EHG) contributed to increased height in present-day populations
(Supplementary Note 4h). This shows that the height differences in Europe between north and south
may not be due to selection, as claimed in many previous studies, but may be a consequence of
differential ancestry.
1188
Likewise, European hunter gatherers are genetically predicted to have dark skin pigmentation and
dark brown hair [11,20,21,79,83,168,204,205], and indeed we see that the WHG, EHG and CHG components
contributed to these phenotypes in present-day individuals whereas the Yamnaya and Anatolian
farmer ancestry contributed to light brown/blonde hair pigmentation (Supplementary Note 4h).
Interestingly, loci associated with overdispersed mood-related polygenic phenotypes recorded
among the UK Biobank individuals (like increased anxiety, guilty feelings, and irritability) showed
an overrepresentation of the Anatolian farmer ancestry component; and the WHG component
showed a strikingly high contribution to traits related to diabetes. We also found that the ApoE4
effect allele is preferentially found on a WHG/EHG haplotypic background, suggesting it likely was
brought to western Europe by early hunter-gatherers (Supplementary Note 4h). This is in line with
the present-day European distribution of this allele, which is highest in north-eastern Europe, where
the proportion of these ancestries are higher than in other regions of the continent [206].
1201
1202 *Conclusion*
1203
Our study has provided fundamental new insights into one of the most transformative periods of
human biological and cultural evolution. We have demonstrated that a clear east-west division
known from Stone Age material culture, extending from the Black Sea to the Baltic and persisting
across several millennia, was genetically deeply rooted in populations with different ancestries. We
showed that the genetic impact of the Neolithic transition was highly distinct, east and west of this
boundary. We have identified a hitherto unknown source of ancestry in hunter-gatherers from the
Middle Don region contributing ancestry to the Yamnaya pastoralists, and we have documented
how the later spread of steppe-related ancestry into Europe was very rapid and mediated through
admixture with people from the Globular Amphora Culture. Additionally, we have observed two
near-complete population replacements in Denmark within just 1,000 years, concomitantly with
major changes in material culture, which rules out cultural diffusion as a main driver and settles
generation-long archaeological debates. Our analyses revealed that the ability to detect signatures of
natural selection in modern human genomes is drastically limited by conflicting selection pressures
in different ancestral populations masking the signals. Developing methods to trace selection in
individual ancestry components allowed us to effectively double the number of significant selection
peaks, which helped clarify the trajectories of a number of traits related to diet and lifestyle. Our
results emphasise how the interplay between major ancient selection and admixture events
occurring across Europe and Asia in the Stone and Bronze Ages have profoundly shaped patterns of
genetic variation in modern human populations.

1228

Extended Data Figures

1230
Extended Data Fig.1. Imputation accuracy of aDNA. Panel A shows imputation accuracy across the 42 high-
coverage ancient genomes when downsampled to lower depth of coverage values. Panels B-D show imputation
accuracy for 1X depth of coverage across 21 high-coverage ancient European genomes. In panels A-D, imputation
accuracy is shown as the squared Pearson correlation between imputed and true genotype dosages as a function of
minor allele frequency of the target variant sites. Panel E shows imputation accuracy as Mendelian error rates for a re-
sequenced Neolithic trio [27] downsampled to a range of low coverage values.

1237
1238

1239

Extended Data Fig. 2. Genetic structure of the 317 herein-reported ancient genomes. (A)-(D) Principal component
analysis of 3,316 modern and ancient individuals from Eurasia, Oceania and the Americas (A, B), as well as restricted
to 2,126 individuals from western Eurasia (west of Urals) (C, D). Shown are analyses with principal components
inferred either using both modern and imputed ancient genomes passing all filters, and projecting low coverage ancient
genomes (A, C); or only modern genomes and projecting all ancient genomes (B, D). Ancient genomes sequenced in
this study are indicated either with black circles (imputed genomes) or grey diamonds (projected genomes). (E) Model-
based clustering results using ADMIXTURE for 284 newly reported genomes (excluding close relatives and individuals
flagged for possible contamination). Results shown are based on ADMIXTURE runs from K=2 to K=15 on 1,584
ancient individuals. Low-coverage individuals represented by pseudo-haploid genotypes are indicated with alpha
transparency.
1250

1251

Extended Data Fig. 3. Genetic clustering of ancient individuals. Genetic clusters inferred from pairwise identity-by-descent (IBD) sharing of 1,401 ancient Eurasian individuals, indicated using colored symbols throughout (A) Network graph of pairwise IBD sharing between 596 ancient Eurasians predating 3,000 BP, highlighting within- and between-cluster relationships. Each node represents an individual, and the width of edges connecting nodes indicates the fraction of the genome shared IBD between the respective pair of individuals. Network edges were restricted to the 10 highest sharing connections for each individual, and the layout was computed using the force-directed Fruchterman-Reingold algorithm. (B) Temporal distribution of clustered individuals, grouped by broad ancestry cluster. (C), (D) Geographical distribution of clustered individuals, shown for individuals predating 3,000 BP (C) and after 3,000 BP (D). (E)-(H) Fine-scale population structure among genetic clusters. Modern individuals are shown in gray, with population labels corresponding to their median coordinates. (E), (F) PCA of 3,119 Eurasian (E) or 2,126 west Eurasian (F) individuals. (G), (H) t-distributed stochastic neighbour embedding (t-SNE) using the first 12 principal components of the all Eurasian panel (E). Shown are embeddings with two different exaggeration factors ρ, emphasising local (G, ρ=1) or global (H, ρ=30) structure.

1266

1277

1278

Extended Data Fig. 5. Deep Eurasian population structure. (A) Admixture graph fit relating deep Eurasian lineages
predating the Last Glacial Maximum (LGM) to later West Eurasian ancestry clusters (worst |Z| = 3.65). (B) Rotating
outgroup *qpAdm* analysis showing fit results for modelling post-LGM target groups as mixtures of all possible
combinations involving one to five source groups. Colours of the individual matrix cells indicate the fit for a particular
model, either rejected at p<0.01 (grey), 0.01≤ p<0.05 (light blue) or p ≥ 0.05 (dark blue). Cells with crosses indicate
infeasible models involving negative admixture proportions. (C) Estimated ancestry proportions from *qpAdm* for post-
LGM target groups inferred from the model fitting with least number of source groups.

1286
1287

1288　**Extended Data Fig 6**. Spatiotemporal kriging of four major ancestry clusters over the last 12,000 years of human history.
1289　LVN = ancestry maximized in Anatolian farmer populations. WHG = ancestry maximized in western European hunter-
1290　gatherers. EHG = ancestry maximized in eastern European hunter-gatherers. IRN = ancestry maximized in Iranian
1291　Neolithic individuals and Caucasus hunter-gatherers.

1292
Extended Data Fig. 7. Genetic transitions in Europe. (A)-(C) Ancestry proportions contributed from preceding local
groups to later individuals during the two major western Eurasian genetic transitions. (A) contribution to individuals

with farmer ancestry from preceding local hunter-gatherer groups; (B,C) contribution to individuals with Steppe-related
ancestry from preceding local farmer groups. Coloured areas in all maps indicate the geographic extent of individuals
included in respective regions. (D) Composition of hunter-gatherer ancestry proportions from different source groups in
individuals with farmer ancestry, shown as heatmap (top) and barplots (bottom). Grey bars represent contributions from
local hunter-gatherers (E)-(G) Moon charts showing spatial distribution of estimated ancestry proportions of mid- to
late Neolithic farmer individuals from three clusters of early Neolithic European farmers (locations indicated with
coloured symbols). Estimated ancestry proportions are indicated by size and amount of fill of moon symbols. (H, I)
Estimated time of admixture between (H) local hunter-gatherer groups and farmers and (I) eastern European farmers
with GAC-related ancestry and Steppe pastoralist groups. Black diamonds and error bars represent point estimate and
standard errors of admixture time, coloured bars show temporal range of included target individuals. The time to
admixture was adjusted backwards by the average age of individuals for each region. (J) Correlation between estimated
proportions of Steppe-related and GAC farmer-related ancestries, across west Eurasian target individuals. Symbol shape
and colour indicate the genetic cluster of respective individuals.
1308

1309
Extended Data Fig. 8. Fine-scale structure in Late Neolithic Scandinavians. (A)-(E) Geographic locations and PCA
based on pairwise IBD sharing (middle) of 148 European individuals predating 3,000 BP. Geographic locations are
shown for 65 individuals belonging to the five genetic clusters observed in 38 ancient Scandinavians (temporal
sequence shown in timeline in centre of plot). Individual assignments and frequency distribution of major Y
chromosome haplogroups are indicated in maps and timeline. Plot symbols with black circles indicate the 38
Scandinavian individuals in the PCA panels. Ancestry proportions for the 38 Scandinavian individuals estimated using
proximal source groups from outside Scandinavia ("postNeolScand" source set) are shown on the right of the respective
cluster results.

1318

Extended Data Fig. 9. Genetic transformations across the Eurasian Steppe. (A)-(C) Principal component analysis of modern and ancient individuals from Eurasia, Oceania and the Americas, highlighting estimated ancestry proportions from "deep" Siberian ancestry sources (individuals highlighted with dashed line). Present-day individuals are shown in gray, with population labels corresponding to their median coordinates. (D)-(E) Moon charts showing spatial distribution of estimated ancestry proportions of Siberian hunter-gatherers before 5,000 BP from "deep" Siberian ancestry sources (names and locations indicated with coloured symbols). Estimated ancestry proportions are indicated by size and amount of fill of moon symbols. (G) Timelines of ancestry proportions from "postNeol" sources in Central and North Asian ancient individuals after 5,000 BP. Symbol shape and colour indicate the genetic cluster of each individual.

1332
Extended Data Fig. 10. Selection at the HLA locus. Results for the pan-ancestry analysis (ALL) plus the four
marginal ancestries: Western hunter-gatherers (WHG), Eastern hunter-gatherers (EHG), Caucasus hunter-gatherers
(CHG) and Anatolian farmers (ANA). Row one shows zoomed Manhattan plots of the p-values for each ancestry
(significant SNPs sized by their selection coefficients), and row two shows allele trajectories for the top SNPs across all
ancestries (grey shading for the marginal ancestries indicates approximate temporal extent of the pre-admixture
population).
1339
1340

1341
Extended Data Fig. 11. Selective sweep at the *SLC22A4* locus. Results for the pan-ancestry analysis (ALL) plus the
four marginal ancestries: Western hunter-gatherers (WHG), Eastern hunter-gatherers (EHG), Caucasus hunter-gatherers
(CHG) and Anatolian farmers (ANA). Row one shows zoomed Manhattan plots of the p-values for each ancestry
(significant SNPs sized by their selection coefficients), and row two shows allele trajectories for the top SNPs across all
ancestries (grey shading for the marginal ancestries indicates approximate temporal extent of the pre-admixture
population).
1348
1349
1350

Extended Data Fig. 12. Selective sweep at the *HECTD4* locus. Results for the pan-ancestry analysis (ALL) plus the four marginal ancestries: Western hunter-gatherers (WHG), Eastern hunter-gatherers (EHG), Caucasus hunter-gatherers (CHG) and Anatolian farmers (ANA). Row one shows zoomed Manhattan plots of the p-values for each ancestry (significant SNPs sized by their selection coefficients), and row two shows allele trajectories for the top SNPs across all ancestries (grey shading for the marginal ancestries indicates approximate temporal extent of the pre-admixture population).

Data availability

All collapsed and paired-end sequence data for novel samples sequenced in this study will be made publicly available on the European Nucleotide Archive, together with trimmed sequence alignment map files, aligned using human build GRCh37. Previously published ancient genomic data used in this study is detailed in Supplementary Table VII, and are all already publicly available. Bioarchaeological data (including Accelerator Mass Spectrometry results) are included in the online supplementary materials of this submission.

Code availability

The modified version of CLUES used in this study is available from https://github.com/standard-aaron/clues. The pipeline and conda environment necessary to replicate the analysis of allele frequency trajectories of trait-associated variants in Supplementary Note 4a are available on Github at https://github.com/ekirving/mesoneo_paper. The pipeline to replicate the analyses for Supplementary Note 4c-4e can be found at https://github.com/albarema/neo. All other analyses relied upon available software which has been fully referenced in the manuscript and detailed in the relevant supplementary notes.

References

1.   Riehl, S., Zeidi, M. & Conard, N. J. Emergence of agriculture in the foothills of the Zagros Mountains of Iran.

     Science **341**, 65–67 (2013).

2.   Broushaki, F. *et al.* Early Neolithic genomes from the eastern Fertile Crescent. *Science* **353**, 499–503 (2016).

3.   Shennan, S. *The First Farmers of Europe: An Evolutionary Perspective*. (Cambridge University Press, 2018).

4.   Ellis, E. C., Klein Goldewijk, K., Siebert, S., Lightman, D. & Ramankutty, N. Anthropogenic transformation of the

biomes, 1700 to 2000. *Glob. Ecol. Biogeogr.* no–no (2010).

5.   Burger, J. R. & Fristoe, T. S. Hunter-gatherer populations inform modern ecology. *Proceedings of the National*

Academy of Sciences of the United States of America vol. 115 1137–1139 (2018).

6.   Jones, E. R. *et al.* The Neolithic Transition in the Baltic Was Not Driven by Admixture with Early European

Farmers. *Curr. Biol.* **27**, 576–582 (2017).

7.   Nikitin, A. G. *et al.* Interactions between earliest Linearbandkeramik farmers and central European hunter

gatherers at the dawn of European Neolithization. *Sci. Rep.* **9**, 19544 (2019).

8.   Betti, L. *et al.* Climate shaped how Neolithic farmers and European hunter-gatherers interacted after a major

slowdown from 6,100 BCE to 4,500 BCE. *Nat Hum Behav* **4**, 1004–1010 (2020).

9.   Furholt, M. Mobility and Social Change: Understanding the European Neolithic Period after the Archaeogenetic

Revolution. *Journal of Archaeological Research* **29**, 481–535 (2021).

10. Immel, A. *et al.* Genome-wide study of a Neolithic Wartberg grave community reveals distinct HLA variation and

hunter-gatherer ancestry. *Commun Biol* **4**, 113 (2021).

11. Brace, S. *et al.* Ancient genomes indicate population replacement in Early Neolithic Britain. *Nat Ecol Evol* **3**, 765–

771 (2019).

12. Fowler, C., Harding, J. & Hofmann, D. Introduction. in *The Oxford Handbook of Neolithic Europe* (eds. Fowler,

C., Harding, J. & Hofmann, D.) (Oxford University Press, 2015).

13. Kislenko, A. & Tatarintseva, N. The eastern Ural steppe at the end of the Stone Age. *Late prehistoric exploitation*

of the Eurasian steppe 183–216 (1999).

14. Mittnik, A. *et al.* The genetic prehistory of the Baltic Sea region. *Nat. Commun.* **9**, 442 (2018).

15. Fischer, A. & Kristiansen, K. *The Neolithisation of Denmark. 150 years of debate*. (J.R. Collis Publications.

Sheffield, 2002).

16. Lipson, M. *et al.* Parallel palaeogenomic transects reveal complex genetic history of early European farmers.

Nature **551**, 368 (2017).

17. Mathieson, I. *et al.* The genomic history of southeastern Europe. *Nature* **555**, 197–203 (2018).

18. Fort, J. Demic and cultural diffusion propagated the Neolithic transition across different regions of Europe. *J. R.*

Soc. Interface **12**, (2015).

19. Fort, J., Mercè Pareta, M. & Sørensen, L. Estimating the relative importance of demic and cultural diffusion in the

spread of the Neolithic in Scandinavia. *J. R. Soc. Interface* **15**, 20180597 (2018).

20. Allentoft, M. E. *et al.* Population genomics of Bronze Age Eurasia. *Nature* **522**, 167–172 (2015).

21. Haak, W. *et al.* Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature*

522, 207–211 (2015).

22. Page, A. E. *et al.* Reproductive trade-offs in extant hunter-gatherers suggest adaptive mechanism for the Neolithic

expansion. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 4694–4699 (2016).

23. Marciniak, S., Bergey, C., Silva, A. M. & Hałuszko, A. An integrative skeletal and paleogenomic analysis of

prehistoric stature variation suggests relatively reduced health for early European farmers. *bioRxiv* (2021).

24. Nielsen, R. Molecular signatures of natural selection. *Annu. Rev. Genet.* **39**, 197–218 (2005).

25. Vitti, J. J., Grossman, S. R. & Sabeti, P. C. Detecting natural selection in genomic data. *Annu. Rev. Genet.* **47**, 97–

120 (2013).

26. Rubinacci, S., Ribeiro, D. M., Hofmeister, R. J. & Delaneau, O. Efficient phasing and imputation of low-coverage

sequencing data using large reference panels. *Nat. Genet.* **53**, 412 (2021).

27. Schroeder, H. *et al.* Unraveling ancestry, kinship, and violence in a Late Neolithic mass grave. *Proc. Natl. Acad.*

Sci. U. S. A. **116**, 10705–10710 (2019).

28. Greenbaum, G., Rubin, A., Templeton, A. R. & Rosenberg, N. A. Network-based hierarchical population structure

analysis for large genomic data sets. *Genome Res.* **29**, 2020–2033 (2019).

29. Lazaridis, I. *et al.* Genomic insights into the origin of farming in the ancient Near East. *Nature* **536**, 419–424

(2016).

30. Lazaridis, I., Belfer-Cohen, A., Mallick, S. & Patterson, N. Paleolithic DNA from the Caucasus reveals core of

West Eurasian ancestry. *BioRxiv* (2018).

31. Günther, T. *et al.* Population genomics of Mesolithic Scandinavia: Investigating early postglacial migration routes

and high-latitude adaptation. *PLoS Biol.* **16**, e2003703 (2018).

32. Klütsch, C. F. C., Manseau, M. & Wilson, P. J. Phylogeographical analysis of mtDNA data indicates postglacial

expansion from multiple glacial refugia in woodland caribou (Rangifer tarandus caribou). *PLoS One* **7**, e52661

(2012).

33. Horreo, J. L. & Fitze, P. S. Postglacial Colonization of Northern Europe by Reptiles. in *Origin and Evolution of*

Biodiversity (ed. Pontarotti, P.) 197–214 (Springer International Publishing, 2018).

34. Olalde, I. *et al.* The genomic history of the Iberian Peninsula over the past 8000 years. *Science* **363**, 1230–1234

(2019).

35. Villalba-Mouco, V. *et al.* Survival of Late Pleistocene Hunter-Gatherer Ancestry in the Iberian Peninsula. *Curr.*

Biol. **29**, 1169–1177.e7 (2019).

36. García-Escárzaga, A. *et al.* Human forager response to abrupt climate change at 8.2 ka on the Atlantic coast of

Europe. *Sci. Rep.* **12**, 6481 (2022).

37. Wang, C.-C. *et al.* Ancient human genome-wide data from a 3000-year interval in the Caucasus corresponds with

eco-geographic regions. *Nat. Commun.* **10**, 590 (2019).

38. Olalde, I. *et al.* The Beaker phenomenon and the genomic transformation of northwest Europe. *Nature* **555**, 190–

196 (2018).

39. Antonio, M. L. *et al.* Ancient Rome: A genetic crossroads of Europe and the Mediterranean. *Science* **366**, 708–714

(2019).

40. Brunel, S. *et al.* Ancient genomes from present-day France unveil 7,000 years of its demographic history. *Proc.*

Natl. Acad. Sci. U. S. A. **117**, 12791–12798 (2020).

41. Cassidy, L. M. *et al.* A dynastic elite in monumental Neolithic society. *Nature* **582**, 384–388 (2020).

42. Racimo, F. *et al.* The spatiotemporal spread of human migrations during the European Holocene. *Proc. Natl. Acad.*

Sci. U. S. A. **117**, 8989–9000 (2020).

43. Saag, L. *et al.* Extensive Farming in Estonia Started through a Sex-Biased Migration from the Steppe. *Curr. Biol.*

27, 2185–2193.e6 (2017).

44. Seguin-Orlando, A. *et al.* Heterogeneous Hunter-Gatherer and Steppe-Related Ancestries in Late Neolithic and

Bell Beaker Genomes from Present-Day France. *Curr. Biol.* **31**, 1072–1083.e10 (2021).

45. Mathieson, I. *et al.* Genome-wide patterns of selection in 230 ancient Eurasians. *Nature* **528**, 499–503 (2015).

46. Martiniano, R. *et al.* The population genomics of archaeological transition in west Iberia: Investigation of ancient

substructure using imputation and haplotype-based methods. *PLoS Genet.* **13**, e1006852 (2017).

47. Isern, N., Zilhão, J., Fort, J. & Ammerman, A. J. Modeling the role of voyaging in the coastal spread of the Early

Neolithic in the West Mediterranean. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 897–902 (2017).

48. Narasimhan, V. M. *et al.* The formation of human populations in South and Central Asia. *Science* **365**, (2019).

49. Skoglund, P. *et al.* Origins and genetic legacy of Neolithic farmers and hunter-gatherers in Europe. *Science* **336**,

466–469 (2012).

50. Malmström, H. *et al.* Ancient mitochondrial DNA from the northern fringe of the Neolithic farming expansion in

Europe sheds light on the dispersion process. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **370**, 20130373 (2015).

51. Coutinho, A. *et al.* The Neolithic Pitted Ware culture foragers were culturally but not genetically influenced by the

Battle Axe culture herders. *Am. J. Phys. Anthropol.* **172**, 638–649 (2020).

52. Becker, C. J. Den grubekeramiske kultur i Danmark. Aarbøger for nordisk Oldkyndighed og Historie. 153–274

(1951).

53. Iversen, R. In a World of Worlds: The Pitted Ware Complex in a large scale perspective. *Acta Archaeol.* **81**, 5–43

(2010).

54. Iversen, R., Philippsen, B. & Persson, P. Reconsidering the Pitted Ware chronology. *Praehistorische Zeitschrift*

96, 44–88 (2021).

55. Klassen, L., Iversen, R. & Rasmussen, L. W. *The Pitted Ware Culture on Djursland Supra-regional: Significance*

and Contacts in the Middle Neolithic of Southern Scandinavia. (Aarhus Universitetsforlag, 2020).

56. Sørensen, L. From hunter to farmer in Northern Europe: Migration and adaptation during the Neolithic and Bronze

Age. *Acta Archaeologica* **85**, (2014).

57. Furholt, M. Die Złota-Gruppe in Kleinpolen: Ein Beispiel für die Transformation eines Zeichensystems?

Germania: Anzeiger der Römisch-Germanischen Kommission des Deutschen Archäologischen Instituts **86**, 1–28

(2008).

58. Szmyt, M. In the Reaches Far of Two Worlds: On the Study of Contacts between the Societies of the Globular

Amphora and Yamnaya Cultures. in *A Turning of Ages: Jubilee Book Dedicated to Professor Jan Machnik on His*

70th Anniversary (ed. Kadrow, S.) 443–466 (Institute of Archaeology and Ethnology, Polish Academy of

Sciences, Cracow Branch, 2000).

59. Tassi, F. *et al.* Genome diversity in the Neolithic Globular Amphorae culture and the spread of Indo-European

languages. *Proc. Biol. Sci.* **284**, (2017).

60. Nordqvist, K. & Heyd, V. The Forgotten Child of the Wider Corded Ware Family: Russian Fatyanovo Culture in

Context. *Proceedings of the Prehistoric Society* **86**, 65–93 (2020).

61. Papac, L. *et al.* Dynamic changes in genomic and social structures in third millennium BCE central Europe. *Sci*

Adv **7**, (2021).

62. Saag, L. *et al.* Genetic ancestry changes in Stone to Bronze Age transition in the East European plain. *Sci Adv* **7**,

(2021).

63. Johannsen, N. & Laursen, S. Routes and Wheeled Transport in Late 4th–Early 3rd Millennium Funerary Customs

of the Jutland Peninsula: Regional Evidence and European Context. **85**, 15–58 (2010).

64. Kristiansen, K. *et al.* Re-theorising mobility and the formation of culture and language among the Corded Ware

Culture in Europe. *Antiquity* **91**, 334–347 (2017).

65. Hansen, J. *et al.* The Maglemosian skeleton from Koelbjerg revisited: Identifying sex and provenance. *Danish*

Journal of Archaeology **6**, 55–66 (2017).

66. Petersen, P. V. Chronological and regional variation in the Late Mesolithic of Eastern Denmark. *Journal of Danish*

Archaeology **3**, 7–18 (1984).

67. Fischer, A. People and the sea – settlement and fishing along the Mesolithic coasts. in *The Danish Storebælt Since*

the Ice Age – Man, Sea and Forest (eds. Pedersen, L., Fischer, A. & Aaby, B.) 63–77 (A/S Storebælt Fixed Link,

1997).

68. Sørensen, S. A. *The Kongemose Culture*. (University Press of Southern Denmark, 2017).

69. Schoeninger, M. J. & Moore, K. Bone stable isotope studies in archaeology. *Journal of World Prehistory* **6**, 247–

296 (1992).

70. Hedges, R. E. M. & Reynard, L. M. Nitrogen isotopes and the trophic level of humans in archaeology. *J. Archaeol.*

Sci. **34**, 1240–1251 (2007).

71. Fischer, A. *et al.* Coast-inland mobility and diet in the Danish Mesolithic and Neolithic: evidence from stable

isotope values of humans and dogs. *J. Archaeol. Sci.* **34**, 2125–2150 (2007).

72. Tauber, H. 13C evidence for dietary habits of prehistoric man in Denmark. *Nature* **292**, 332–333 (1981).

73. Loog, L. *et al.* Estimating mobility using sparse data: Application to human genetic variation. *Proc. Natl. Acad.*

Sci. U. S. A. **114**, 12213–12218 (2017).

74. Racimo, F. *et al.* The Spatiotemporal Spread of Human Migrations during the European Holocene. *Proc. Natl.*

Acad. Sci. U. S. A. **117**, 8989–9000 (2020).

75. Worsaae, J. J. A. Om en ny deling af Steen- og Bronzealderen. Oversigten over det Kongelige Danske

Videnskabernes Selskabs Forhandlinger (reprint). in *The Neolithisation of Denmark. 150 years of debate* (eds.

Fischer, A. & Kristiansen, K.) 47–56 (J.R. Collis Publications, 1859).

76. Becker, C. J. Mosefundne lerkar fra yngre stenalder (Neolithic pottery in Danish bogs), Copenhagen. *Aarbøger for*

nordisk Oldkyndighed og Historie (1947).

77. Andersen, S. H. 'Køkkenmøddinger' (shell middens) in Denmark: A survey. *Proc. Prehist. Soc.* **66**, 361–384

(2000).

78. Gutiérrez-Zugasti, I. *et al.* Shell midden research in Atlantic Europe: State of the art, research problems and

perspectives for the future. *Quat. Int.* **239**, 70–85 (2011).

79. Olalde, I. *et al.* Derived immune and ancestral pigmentation alleles in a 7,000-year-old Mesolithic European.

Nature **507**, 225–228 (2014).

80. Cox, S. L., Ruff, C. B., Maier, R. M. & Mathieson, I. Genetic contributions to variation in human stature in

prehistoric Europe. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 21484–21492 (2019).

81. Brinch Petersen, E. Gravene ved Dragsholm. Fra jægere til bønder for 6000 år siden. *Nationalmuseets arbejdsmark*

1974, 112–120 (1974).

82. Price, T. D. *et al.* New information on the Stone Age graves at Dragsholm, Denmark. *Acta Archaeol.* **78**, 193–219

(2007).

83. Jensen, T. Z. T. *et al.* A 5700 year-old human genome and oral microbiome from chewed birch pitch. *Nat.*

Commun. **10**, 5520 (2019).

84. Grasgruber, P., Sebera, M., Hrazdíra, E., Cacek, J. & Kalina, T. Major correlates of male height: A study of 105

countries. *Econ. Hum. Biol.* **21**, 172–195 (2016).

85. Sugita, S. Theory of quantitative reconstruction of vegetation II: all you need is LOVE. *Holocene* **17**, 243–257

(2007).

86. Nielsen, A. B. *et al.* Quantitative reconstructions of changes in regional openness in north-central Europe reveal

new insights into old questions. *Quat. Sci. Rev.* **47**, 131–147 (2012).

87. Githumbi, E. *et al.* Pollen-based maps of past regional vegetation cover in Europe over twelve millennia -

evaluation and potential. *Frontiers in Ecology and Evolution* **(Provisionally accepted)**, (2022).

88. Nielsen, A. B. & Odgaard, B. V. Quantitative landscape dynamics in Denmark through the last three millennia

based on the Landscape Reconstruction Algorithm approach. *Veg. Hist. Archaeobot.* **19**, 375–387 (2010).

89. Søe, N. E., Odgaard, B. V., Nielsen, A. B., Olsen, J. & Kristiansen, S. M. Late Holocene landscape development

around a Roman Iron Age mass grave, Alken Enge, Denmark. *Veg. Hist. Archaeobot.* **26**, 277–292 (2017).

90. Kristiansen, K., Melheim, L., Bech, J.-H., Mortensen, M. F. & Frei, K. M. Thy at the Crossroads: A Local Bronze

Age Community's Role in a Macro-Economic System. in *Contrasts of the Nordic Bronze Age* 269–282 (Brepols

Publishers, 2020).

91. Borzunov, V. A. The neolithic fortified settlements of the Western Siberia and Trans-Urals. *Rossijskaâ arheologiâ*

20–34 (2013).

92. Piezonka, H. *et al.* The emergence of hunter-gatherer pottery in the Urals and West Siberia: New dating and stable

isotope evidence. *J. Archaeol. Sci.* **116**, 105100 (2020).

93. Damgaard, P. de B. *et al.* The first horse herders and the impact of early Bronze Age steppe expansions into Asia.

Science **360**, (2018).

94. Yu, H. *et al.* Paleolithic to Bronze Age Siberians Reveal Connections with First Americans and across Eurasia.

Cell **181**, 1232–1245.e20 (2020).

95. Okladnikov, A. P. Neolit i bronzovyi vek Pribaikaliya [Neolithic and Bronze Age of the Baikal region]. Materialy i

issledovaniya po arkheologii SSSR; №. 18 [Materials and researches on Archeology of the USSR; Vol. 18].

Moscow, Leningrad, AS USSR Publ., 1950, Part 1 and 2, 411 p. (1950).

96. Merz, V. On the origin of the complexes of the Baikal type in the Eneolithic of Kazakhstan. in *Paleodemography*

and migration processes in Western Siberia in antiquity and the Middle Ages (ed. Kiryushin, Y. F.) 39–42 (Altai

State University, 1994).

97. Merz, V. Periodization of the Holocene complexes of Northern and Central Kazakhstan based on the materials of

the multilayer site Shiderty 3. (Kemerovo State University, 2008).

98. Damgaard, P. de B. *et al.* 137 ancient human genomes from across the Eurasian steppes. *Nature* **557**, 369–374

(2018).

99. Librado, P. *et al.* The origins and spread of domestic horses from the Western Eurasian steppes. *Nature* **598**, 634–

640 (2021).

100. Lawson, D. J., Hellenthal, G., Myers, S. & Falush, D. Inference of population structure using dense haplotype data.

PLoS Genet. **8**, e1002453 (2012).

101. Margaryan, A. *et al.* Population genomics of the Viking world. *Nature* **585**, 390–396 (2020).

102. Hellenthal, G. *et al.* A genetic atlas of human admixture history. *Science* **343**, 747–751 (2014).

103. Sikora, M. *et al.* The population history of northeastern Siberia since the Pleistocene. *Nature* **570**, 182–188 (2019).

104. Shinde, V. *et al.* An Ancient Harappan Genome Lacks Ancestry from Steppe Pastoralists or Iranian Farmers. *Cell*

179, 729–735.e10 (2019).

105. Hofmanová, Z. *et al.* Early farmers from across Europe directly descended from Neolithic Aegeans. *Proc. Natl.*

Acad. Sci. U. S. A. **113**, 6886–6891 (2016).

106. Feldman, M. *et al.* Late Pleistocene human genome suggests a local origin for the first farmers of central Anatolia.

Nat. Commun. **10**, 1218 (2019).

107. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209

(2018).

108. Galinsky, K. J., Loh, P.-R., Mallick, S., Patterson, N. J. & Price, A. L. Population Structure of UK Biobank and

Ancient Eurasians Reveals Adaptation at Genes Influencing Blood Pressure. *Am. J. Hum. Genet.* **99**, 1130–1139

(2016).

109. Patterson, N. *et al.* Large-scale migration into Britain during the Middle to Late Bronze Age. *Nature* (2021)

doi:10.1038/s41586-021-04287-4.

110. Palamara, P. F., Lencz, T., Darvasi, A. & Pe'er, I. Length distributions of identity by descent reveal fine-scale

1593   demographic history. *Am. J. Hum. Genet.* **91**, 809–822 (2012).

1594   111. Ringbauer, H., Novembre, J. & Steinrücken, M. Parental relatedness through time revealed by runs of

1595   homozygosity in ancient DNA. *Nat. Commun.* **12**, 5425 (2021).

1596   112. Sikora, M. *et al.* Ancient genomes show social and reproductive behavior of early Upper Paleolithic foragers.

1597   *Science* **358**, 659–662 (2017).

1598   113. Scheib, C. L. *et al.* Ancient human parallel lineages within North America contributed to a coastal expansion.

1599   *Science* **360**, 1024–1027 (2018).

1600   114. Krzewińska, M. *et al.* Ancient genomes suggest the eastern Pontic-Caspian steppe as the source of western Iron

1601   Age nomads. *Sci Adv* **4**, eaat4457 (2018).

1602   115. Girirajan, S., Campbell, C. D. & Eichler, E. E. Human copy number variation and complex genetic disease. *Annu.*

1603   *Rev. Genet.* **45**, 203–226 (2011).

1604   116. Weise, A. *et al.* Microdeletion and microduplication syndromes. *J. Histochem. Cytochem.* **60**, 346–358 (2012).

1605   117. Girirajan, S. *et al.* Phenotypic heterogeneity of genomic disorders and rare copy-number variants. *N. Engl. J. Med.*

1606   **367**, 1321–1331 (2012).

1607   118. Mallick, S. *et al.* The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature* **538**,

1608   201–206 (2016).

1609   119. Bergström, A. *et al.* Insights into human genetic variation and population history from 929 diverse genomes.

1610   *Science* **367**, (2020).

1611   120. Sudmant, P. H. *et al.* Diversity of human copy number variation and multicopy genes. *Science* **330**, 641–646

1612   (2010).

1613   121. Nagamani, S. C. S. *et al.* Phenotypic manifestations of copy number variation in chromosome 16p13.11. *Eur. J.*

1614   *Hum. Genet.* **19**, 280–286 (2011).

1615   122. Cremin, B., Goodman, H., Spranger, J. & Beighton, P. Wormian bones in osteogenesis imperfecta and other

1616   disorders. *Skeletal Radiol.* **8**, 35–38 (1982).

1617   123. Foreman, P. *et al.* L5 spondylolysis/spondylolisthesis: a comprehensive review with an anatomic focus. *Childs.*

1618   *Nerv. Syst.* **29**, 209–216 (2012).

1619   124. Vleeming, A. *et al.* The sacroiliac joint: an overview of its anatomy, function and potential clinical implications. *J.*

1620   *Anat.* **221**, 537–567 (2012).

125. Salanitri, S. & Seow, W. K. Developmental enamel defects in the primary dentition: aetiology and clinical

management. *Aust. Dent. J.* **58**, 133–40; quiz 266 (2013).

126. Crawford, K. *et al.* Medical consequences of pathogenic CNVs in adults: analysis of the UK Biobank. *J. Med.*

Genet. **56**, 131–138 (2019).

127. Stern, A. J., Wilton, P. R. & Nielsen, R. An approximate full-likelihood method for inferring selection and allele

frequency trajectories from DNA sequence data. *PLoS Genet.* **15**, e1008384 (2019).

128. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays

and summary statistics 2019. *Nucleic Acids Res.* **47**, D1005–D1012 (2019).

129. Enattah, N. S. *et al.* Independent introduction of two lactase-persistence alleles into human populations reflects

different history of adaptation to milk culture. *Am. J. Hum. Genet.* **82**, 57–72 (2008).

130. Itan, Y., Powell, A., Beaumont, M. A., Burger, J. & Thomas, M. G. The origins of lactase persistence in Europe.

PLoS Comput. Biol. **5**, e1000491 (2009).

131. Ségurel, L. & Bon, C. On the Evolution of Lactase Persistence in Humans. *Annu. Rev. Genomics Hum. Genet.* **18**,

297–319 (2017).

132. Segurel, L. *et al.* Why and when was lactase persistence selected for? Insights from Central Asian herders and

ancient DNA. *PLoS Biol.* **18**, e3000742 (2020).

133. Enattah, N. S. *et al.* Identification of a variant associated with adult-type hypolactasia. *Nat. Genet.* **30**, 233–237

(2002).

134. Bersaglieri, T. *et al.* Genetic signatures of strong recent positive selection at the lactase gene. *Am. J. Hum. Genet.*

74, 1111–1120 (2004).

135. Willer, C. J. *et al.* Discovery and refinement of loci associated with lipid levels. *Nat. Genet.* **45**, 1274–1283 (2013).

136. Gallois, A. *et al.* A comprehensive study of metabolite genetics reveals strong pleiotropy and heterogeneity across

time and context. *Nat. Commun.* **10**, 4788 (2019).

137. Ligthart, S. *et al.* Bivariate genome-wide association study identifies novel pleiotropic loci for lipids and

inflammation. *BMC Genomics* **17**, 443 (2016).

138. Buckley, M. T. *et al.* Selection in Europeans on Fatty Acid Desaturases Associated with Dietary Changes. *Mol.*

Biol. Evol. **34**, 1307–1318 (2017).

139. Ye, K., Gao, F., Wang, D., Bar-Yosef, O. & Keinan, A. Dietary adaptation of FADS genes in Europe varied across

time and geography. *Nat Ecol Evol* **1**, 167 (2017).

140. Mathieson, S. & Mathieson, I. FADS1 and the Timing of Human Adaptation to Agriculture. *Mol. Biol. Evol.* **35**,

2957–2970 (2018).

141. Lazaridis, I. The evolutionary history of human populations in Europe. *Curr. Opin. Genet. Dev.* **53**, 21–27 (2018).

142. Luu, K., Bazin, E. & Blum, M. G. B. pcadapt: an R package to perform genome scans for selection based on

principal component analysis. *Mol. Ecol. Resour.* **17**, 67–77 (2017).

143. Fairn, G. D. & McMaster, C. R. Emerging roles of the oxysterol-binding protein family in metabolism, transport,

and signaling. *Cell. Mol. Life Sci.* **65**, 228–236 (2008).

144. Lehto, M. & Olkkonen, V. M. The OSBP-related proteins: a novel protein family involved in vesicle transport,

cellular lipid metabolism, and cell signalling. *Biochim. Biophys. Acta* **1631**, 1–11 (2003).

145. Sánchez-Solana, B., Li, D.-Q. & Kumar, R. Cytosolic functions of MORC2 in lipogenesis and adipogenesis.

Biochim. Biophys. Acta **1843**, 316–326 (2014).

146. Zhong, X. *et al.* The zinc-finger protein ZFYVE1 modulates TLR3-mediated signaling by facilitating TLR3 ligand

binding. *Cell. Mol. Immunol.* **17**, 741–752 (2020).

147. Kim, S. V. *et al.* GPR15-mediated homing controls immune homeostasis in the large intestine mucosa. *Science*

340, 1456–1459 (2013).

148. Nguyen, L. P. *et al.* Role and species-specific expression of colon T cell homing receptor GPR15 in colitis. *Nature*

Immunology vol. 16 207–213 (2015).

149. Monteleone, G., Boirivant, M., Pallone, F. & MacDonald, T. T. TGF-β1 and Smad7 in the regulation of IBD.

Mucosal Immunology vol. 1 S50–S53 (2008).

150. Kennedy, B. W. C. Mongersen, an Oral SMAD7 Antisense Oligonucleotide, and Crohn's Disease. *The New*

England journal of medicine vol. 372 2461 (2015).

151. Garo, L. P. *et al.* Smad7 Controls Immunoregulatory PDL2/1-PD1 Signaling in Intestinal Inflammation and

Autoimmunity. *Cell Rep.* **28**, 3353–3366.e5 (2019).

152. Morris, J. A. *et al.* An atlas of genetic influences on osteoporosis in humans and mice. *Nat. Genet.* **51**, 258–266

(2018).

153. Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide association study of educational

attainment in 1.1 million individuals. *Nat. Genet.* **50**, 1112–1121 (2018).

154. Jones, A. V. *et al.* GWAS of self-reported mosquito bite size, itch intensity and attractiveness to mosquitoes

implicates immune-related predisposition loci. *Hum. Mol. Genet.* **26**, 1391–1406 (2017).

155. Gutierrez-Achury, J. *et al.* Functional implications of disease-specific variants in loci jointly associated with

coeliac disease and rheumatoid arthritis. *Hum. Mol. Genet.* **25**, 180–190 (2016).

156. Stefansson, H. *et al.* A common inversion under selection in Europeans. *Nat. Genet.* **37**, 129–137 (2005).

157. Steinberg, K. M. *et al.* Structural diversity and African origin of the 17q21.31 inversion polymorphism. *Nat.*

Genet. **44**, 872–880 (2012).

158. Kılınç, G. M. *et al.* The Demographic Development of the First Farmers in Anatolia. *Curr. Biol.* **26**, 2659–2666

(2016).

159. Andreadis, A., Brown, W. M. & Kosik, K. S. Structure and novel exons of the human tau gene. *Biochemistry* **31**,

10626–10633 (1992).

160. Alonso, A. C., Grundke-Iqbal, I. & Iqbal, K. Alzheimer's disease hyperphosphorylated tau sequesters normal tau

into tangles of filaments and disassembles microtubules. *Nat. Med.* **2**, 783–787 (1996).

161. Desikan, R. S. *et al.* Genetic overlap between Alzheimer's disease and Parkinson's disease at the MAPT locus.

Mol. Psychiatry **20**, 1588–1595 (2015).

162. Satake, W. *et al.* Genome-wide association study identifies common variants at four loci as genetic risk factors for

Parkinson's disease. *Nat. Genet.* **41**, 1303–1307 (2009).

163. Simón-Sánchez, J. *et al.* Genome-wide association study reveals genetic risk underlying Parkinson's disease. *Nat.*

Genet. **41**, 1308–1312 (2009).

164. Shulman, J. M. & De Jager, P. L. Evidence for a common pathway linking neurodegenerative diseases. *Nature*

genetics vol. 41 1261–1262 (2009).

165. Aoki, K. Sexual selection as a cause of human skin colour variation: Darwin's hypothesis revisited. *Ann. Hum.*

Biol. **29**, 589–608 (2002).

166. Frost, P. The puzzle of European hair, eye, and skin color. *Advances in Anthropology* **2014**, (2014).

167. Lona-Durazo, F. *et al.* Meta-analysis of GWA studies provides new insights on the genetic architecture of skin

pigmentation in recently admixed populations. *BMC Genet.* **20**, 59 (2019).

168. Ju, D. & Mathieson, I. The evolution of skin pigmentation-associated variation in West Eurasia. *Proc. Natl. Acad.*

Sci. U. S. A. **118**, (2021).

169. Jablonski, N. G. & Chaplin, G. The evolution of human skin coloration. *J. Hum. Evol.* **39**, 57–106 (2000).

170. Engelsen, O. The relationship between ultraviolet radiation exposure and vitamin D status. *Nutrients* **2**, 482–495

(2010).

171. Voight, B. F., Kudaravalli, S., Wen, X. & Pritchard, J. K. A Map of Recent Positive Selection in the Human

Genome. *PLoS Biol.* **4**, e72 (2006).

172. Sabeti, P. C. *et al.* Genome-wide detection and characterization of positive selection in human populations. *Nature*

449, 913–918 (2007).

173. Martin, A. R. *et al.* An Unexpectedly Complex Architecture for Skin Pigmentation in Africans. *Cell* **171**, 1340–

1353.e14 (2017).

174. Wu, H. *et al.* Transcriptome Sequencing to Detect the Potential Role of Long Noncoding RNAs in Salt-Sensitive

Hypertensive Rats. *Biomed Res. Int.* **2019**, 2816959 (2019).

175. Logsdon, B. A., Hoffman, G. E. & Mezey, J. G. Mouse obesity network reconstruction with a variational Bayes

algorithm to employ aggressive false positive control. *BMC Bioinformatics* **13**, 53 (2012).

176. Wang, L. *et al.* Peakwide Mapping on Chromosome 3q13 Identifies the Kalirin Gene as a Novel Candidate Gene

for Coronary Artery Disease. *The American Journal of Human Genetics* vol. 80 650–663 (2007).

177. Ikram, M. A., Seshadri, S. & Bis, J. C. Genomewide Association Studies of Stroke. *Journal of Vascular Surgery*

vol. 50 467 (2009).

178. Krug, T. *et al.* Kalirin: a novel genetic risk factor for ischemic stroke. *Hum. Genet.* **127**, 513–523 (2010).

179. Zang, X.-L. *et al.* Association of a SNP in SLC35F3 Gene with the Risk of Hypertension in a Chinese Han

Population. *Frontiers in Genetics* vol. 7 (2016).

180. Zhang, K. *et al.* Genetic implication of a novel thiamine transporter in human hypertension. *J. Am. Coll. Cardiol.*

63, 1542–1555 (2014).

181. Russo, L. *et al.* Cholesterol 25-hydroxylase (CH25H) as a promoter of adipose tissue inflammation in obesity and

diabetes. *Mol Metab* **39**, 100983 (2020).

182. Zhao, J., Chen, J., Li, M., Chen, M. & Sun, C. Multifaceted Functions of CH25H and 25HC to Modulate the Lipid

Metabolism, Immune Responses, and Broadly Antiviral Activities. *Viruses* **12**, (2020).

183. Demir, A., Kahraman, R., Candan, G. & Ergen, A. The role of FAS gene variants in inflammatory bowel disease.

Turk. J. Gastroenterol. **31**, 356–361 (2020).

184. Izawa, T. *et al.* ASXL2 Regulates Glucose, Lipid, and Skeletal Homeostasis. *Cell Rep.* **11**, 1625–1637 (2015).

185. Zou, W. *et al.* Myeloid-specific Asxl2 deletion limits diet-induced obesity by regulating energy expenditure. *J.*

Clin. Invest. **130**, 2644–2656 (2020).

186. Park, U.-H., Yoon, S. K., Park, T., Kim, E.-J. & Um, S.-J. Additional Sex Comb-like (ASXL) Proteins 1 and 2

Play Opposite Roles in Adipogenesis via Reciprocal Regulation of Peroxisome Proliferator-activated Receptor γ.

Journal of Biological Chemistry vol. 286 1354–1363 (2011).

187. Ponsuksili, S. *et al.* Epigenome-wide skeletal muscle DNA methylation profiles at the background of distinct

metabolic types and ryanodine receptor variation in pigs. *BMC Genomics* **20**, 492 (2019).

188. Samad, M. B. *et al.* [6]-Gingerol, from Zingiber officinale, potentiates GLP-1 mediated glucose-stimulated insulin

secretion pathway in pancreatic β-cells and increases RAB8/RAB10-regulated membrane presentation of GLUT4

transporters in skeletal muscle to improve hyperglycemia in Leprdb/db type 2 diabetic mice. *BMC Complementary*

and Alternative Medicine vol. 17 (2017).

189. Vazirani, R. P. *et al.* Disruption of Adipose Rab10-Dependent Insulin Signaling Causes Hepatic Insulin

Resistance. *Diabetes* **65**, 1577–1589 (2016).

190. Hsieh, P. *et al.* Exome Sequencing Provides Evidence of Polygenic Adaptation to a Fat-Rich Animal Diet in

Indigenous Siberian Populations. *Mol. Biol. Evol.* **34**, 2913–2926 (2017).

191. Thapa, D. *et al.* The protein acetylase GCN5L1 modulates hepatic fatty acid oxidation activity via acetylation of

the mitochondrial β-oxidation enzyme HADHA. *J. Biol. Chem.* **293**, 17676–17684 (2018).

192. Baloni, P. *et al.* Genome-scale metabolic model of the rat liver predicts effects of diet restriction. *Sci. Rep.* **9**, 9807

(2019).

193. Ong, H. S. & Yim, H. C. H. Microbial Factors in Inflammatory Diseases and Cancers. *Regulation of Inflammatory*

Signaling in Health and Disease 153–174 (2017) doi:10.1007/978-981-10-5987-2_7.

194. Berg, J. J. & Coop, G. A population genetic signal of polygenic adaptation. *PLoS Genet.* **10**, e1004412 (2014).

195. Berg, J. J. *et al.* Reduced signal for polygenic adaptation of height in UK Biobank. *Elife* **8**, (2019).

196. Sohail, M. *et al.* Polygenic adaptation on height is overestimated due to uncorrected stratification in genome-wide

association studies. *Elife* **8**, (2019).

197. Irving-Pease, E. K., Muktupavela, R., Dannemann, M. & Racimo, F. Quantitative Human Paleogenetics: What can

Ancient DNA Tell us About Complex Trait Evolution? *Front. Genet.* **12**, 703541 (2021).

198. Refoyo-Martínez, A. *et al.* How robust are cross-population signatures of polygenic adaptation in humans? *bioRxiv*

2020.07.13.200030 (2021) doi:10.1101/2020.07.13.200030.

199. Ruff, C. B. *Skeletal Variation and Adaptation in Europeans: Upper Paleolithic to the Twentieth Century.* (John

Wiley & Sons, 2017).

200. Tanigawa, Y. *et al.* Components of genetic associations across 2,138 phenotypes in the UK Biobank highlight

adipocyte biology. *Nat. Commun.* **10**, 4064 (2019).

201. Corder, E. H. *et al.* Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset

families. *Science* **261**, 921–923 (1993).

202. Strittmatter, W. J. *et al.* Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4

allele in late-onset familial Alzheimer disease. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 1977–1981 (1993).

203. Rosenstock, E. *et al.* Human stature in the Near East and Europe ca. 10,000–1000 BC: its spatiotemporal

1772   development in a Bayesian errors-in-variables model. *Archaeol. Anthropol. Sci.* **11**, 5657–5690 (2019).

1773 204. Wilde, S. *et al.* Direct evidence for positive selection of skin, hair, and eye pigmentation in Europeans during the

1774   last 5,000 y. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 4832–4837 (2014).

1775 205. González-Fortes, G. *et al.* Paleogenomic Evidence for Multi-generational Mixing between Neolithic Farmers and

1776   Mesolithic Hunter-Gatherers in the Lower Danube Basin. *Curr. Biol.* **27**, 1801–1810.e10 (2017).

1777 206. Belloy, M. E., Napolioni, V. & Greicius, M. D. A Quarter Century of APOE and Alzheimer's Disease: Progress to

1778   Date and the Path Forward. *Neuron* **101**, 820–838 (2019).

Acknowledgements

This publication, the culmination of a research effort lasting over a decade, is dedicated to the memory of Pia
Bennike, who was part of a small core team that initiated the project. Sadly Pia passed away in 2017 but this
study, and many others, testifies to her tremendous efforts and knowledge on Danish prehistoric skeletal
material.

We are deeply indebted to former and present staff members at the National Museum, the Anthropological
Laboratory, the regional museums and citizen scientists of Denmark, who for many generations carefully
collected, recorded and curated the prehistoric skeletal remains that form a key component of this study. We
are equally thankful to curators of the many other institutions across major parts of Eurasia, who to the
benefit of following generations curated human skeletal remains and gave us access and permission to
sample this precious material. We also thank all the former and current staff at the Lundbeck Foundation
GeoGenetics Centre and the GeoGenetics Sequencing Core, and to colleagues across the many institutions
detailed below. We are particularly grateful to Line Olsen as project manager for the Lundbeck Foundation
GeoGenetics Centre project, and to Pernille Selmer Olsen for assisting with sample processing. We thank
UK Biobank Ltd. for access to the UK Biobank genomic resource. We are thankful to Illumina Inc. for
collaboration and to L. Speidel for assistance in running *Relate*. EW thanks St. John's College, Cambridge,
for providing a stimulating environment of discussion and learning.

The Lundbeck Foundation GeoGenetics Centre is supported by the the Lundbeck Foundation (R302-2018-
2155, R155-2013-16338), the Novo Nordisk Foundation (NNF18SA0035006), the Wellcome Trust
(UNS69906), Carlsberg Foundation (CF18-0024), the Danish National Research Foundation (44113220) and
the University of Copenhagen (KU2016 programme). This research has been conducted using the UK
Biobank Resource and the iPSYCH Initiative, funded by the Lundbeck Foundation (R102-A9118 and R155-
2014-1724). This work was further supported by the Swedish Foundation for Humanities and Social
Sciences grant (Riksbankens Jubileumsfond M16-0455:1) to KK. M.E.A. was supported by Marie
Skłodowska-Curie Actions of the EU (grant no. 300554), The Villum Foundation (grant no. 10120) and
Independent Research Fund Denmark (grant no. 7027-00147B). W.B. is supported by the Hanne and Torkel
Weis-Fogh Fund (Department of Zoology, University of Cambridge); AP is funded by Wellcome grant
WT214300, B.S.d.M and O.D. by the Swiss National Science Foundation (SFNS PP00P3_176977) and
European Research Council (ERC 679330); M.N. by the Human Frontier Science Program Postdoctoral
Fellowship (LT000143/2019-L4); R. Macleod by an SSHRC doctoral studentship grant (G101449:
'Individual Life Histories in Long-Term Cultural Change'); G.R. by a Novo Nordisk Foundation Fellowship
(gNNF20OC0062491); N.N.J. by Aarhus University Research Foundation; H.S. by a Carlsberg Foundation
Fellowship (CF19-0601); G.S. by Marie Skłodowska-Curie Individual Fellowship 'PALAEO-ENEO' (grant

agreement number 751349); A.J. Schork by a Lundbeckfonden Fellowship (R335-2019-2318) and the
National Institute on Aging (NIH award numbers U19AG023122, U24AG051129, and UH2AG064706);
A.V.L. and I.V.S. by the Science Committee, Ministry of Education and Science of the Republic of
Kazakhstan (AP08856317); B.G.R. and MGM by the Spanish Ministry of Science and Innovation (Project
HAR2016-75605-R); C.M.-L. and O.R. by the Italian Ministry for the Universities (grants '2010-11
prot.2010EL8TXP_001 Biological and cultural heritage of the central-southern Italian population trough 30
thousand Years' and '2008 prot. 2008B4J2HS_001 Origin and diffusion of farming in central-southern Italy:
a molecular approach'); D.C.-S. and I.G.Z. by the Spanish Ministry of Science and Innovation (Project
HAR2017-86262-P). D.C.S.G. acknowledges funding from the Generalitat Valenciana
(CIDEGENT/2019/061) and the Spanish Government (EUR2020-112213); D.B. was supported by the
NOMIS Foundation and Marie Skłodowska-Curie Global Fellowship 'CUSP' (grant no. 846856); E.R.U. by
the Science Committee, Ministry of Education and Science of the Republic of Kazakhstan (AP09261083:
"Transcultural Communications in the Late Bronze Age (Western Siberia - Kazakhstan - Central Asia)");
E.C. by Villum Fonden (17649); J.E.A.T. by the Spanish Ministry of Economy and Competitiveness,
(HAR2013-46861-R) and Generalitat Valenciana (Aico/ 2018/125 and Aico 2020/97). L.Y. acknowledges
funding by the Science Committee of the Armenian Ministry of Education and Science (Project 21AG-
1F025), L.V. by ERC Consolidator Grant 'PEGASUS' (agreement no. 681605); M. Sablin by the Russian
Ministry of Science and Higher Education (075-15-2021-1069); N.C. by Historic Environment Scotland;
S.V. by the Russian Ministry of Science and Higher Education (075-15-2020-910); V.M. by the Science
Committee, Ministry of Education and Science of the Republic of Kazakhstan (AR08856925). V.A. is
supported by a Lundbeckfonden Fellowship (R335-2019-2318); P.H.S. by the National Institute of General
Medical Sciences (R35GM142916); S.R. by the Novo Nordisk Foundation (NNF14CC0001); R.D. by the
Wellcome Trust (WT214300); R.N. by the National Institute of General Medical Sciences (NIH grant
R01GM138634); F. Racimo by a Villum Fonden Young Investigator Grant (no. 00025300). T.W. and V.A.
are supported by the Lundbeck Foundation iPSYCH initiative (R248-2017-2003).
1839

1840   Author Information

1841   These authors contributed equally: Morten E. Allentoft, Martin Sikora, Alba Refoyo-Martínez, Evan K.
1842   Irving-Pease, Anders Fischer, William Barrie & Andrés Ingason
1843

1844   These authors equally supervised research: Thorfinn Korneliussen, Richard Durbin, Rasmus Nielsen, Olivier
1845   Delaneau, Thomas Werge, Fernando Racimo, Kristian Kristiansen & Eske Willerslev
1846

1847   <u>Affiliations</u>
1848

Lundbeck Foundation GeoGenetics Centre, GLOBE Institute, University of Copenhagen,
Copenhagen, Denmark
Morten E. Allentoft, Martin Sikora, Alba Refoyo-Martínez, Evan K. Irving-Pease, Andrés Ingason, Jesper
Stenderup, Fabrice Demeter, Maria Novosolov, Ruairidh Macleod, Rasmus A. Henriksen, Tharsika Vimala,
Hugh McColl, Lasse Vinner, Gabriele Scorrano, Abigail Ramsøe, Anders Rosengren, Anthony Ruter, Anne
Birgitte Gotfredsen, Charleen Gaunitz, Fulya Eylem Yediay, Isin Altinkaya, Lei Zhao, Peter de Barros
Damgaard, Kurt H. Kjær, Thorfinn Korneliussen, Rasmus Nielsen, Thomas Werge, Fernando Racimo,
Kristian Kristiansen & Eske Willerslev
1857

1858   **Trace and Environmental DNA (TrEnD) Laboratory, School of Molecular and Life Sciences, Curtin**
1859   **University, Perth, Australia**

1860   Morten E. Allentoft
1861

1862   **GeoGenetics Group, Department of Zoology, University of Cambridge, Cambridge, UK**
1863   William Barrie, Alice Pearson, Ruairidh Macleod & Eske Willerslev
1864

1865   **Department of Historical Studies, University of Gothenburg, Gothenburg, Sweden**
1866   Anders Fischer, Karl-Göran Sjögren, Bettina Schulz Paulsson, Malou Blank & Kristian Kristiansen
1867

1868   **Institute of Biological Psychiatry, Mental Health Services, Copenhagen University Hospital, Roskilde,**
1869   **Denmark**
1870   Andrés Ingason, Vivek Appadurai,
1871

1872   **The National Museum of Denmark, Ny Vestergade 10, Copenhagen, Denmark**
1873   Lasse Sørensen, Poul Otto Nielsen, Morten Fischer Mortensen, Peter Rasmussen, Peter Vang Petersen,
1874

1875   **Sealand Archaeology, Gl. Roesnaesvej 27, 4400 Kalundborg, Denmark**
1876   Anders Fischer
1877

1878   **Department of Genetics, University of Cambridge, Cambridge, UK**
1879   Alice Pearson & Richard Durbin
1880

1881   **Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge, UK**
1882   Richard Durbin & Eske Willerslev
1883

1884   **Department of Computational Biology, University of Lausanne, Switzerland**
1885   Bárbara Sousa da Mota & Olivier Delaneau
1886

1887   **Swiss Institute of Bioinformatics, University of Lausanne, Switzerland**
1888   Bárbara Sousa da Mota
1889

1890   **Department of Integrative Biology, University of California, Berkeley, USA**
1891   Alma S. Halgren, Peter H. Sudmant & Rasmus Nielsen
1892

1893   **Center for Computational Biology, University of California, Berkeley, USA**
1894   Andrew Vaughn, Aaron J. Stern, Peter H. Sudmant & Rasmus Nielsen
1895

1896   **Laboratory of Biological Anthropology, Department of Forensic Medicine, University of Copenhagen,**
1897   **Copenhagen, Denmark**
1898   Marie Louise Schjellerup Jørkov & Niels Lynnerup
1899

1900   **Muséum National d'Histoire Naturelle, Paris, France**
1901   Fabrice Demeter
1902

1903   **Research department of Genetics, Evolution and Environment, University College London, London,**
1904   **UK**
1905   Ruairidh Macleod
1906

1907 **Section for Evolutionary Genomics, GLOBE Institute, University of Copenhagen, Copenhagen,**
1908 **Denmark**
1909 Ashot Margaryan, Theis Zetner Trolle Jensen, Hannes Schroeder, Enrico Cappellini
1910

1911 **Centre for Evolutionary Hologenomics, University of Copenhagen, Copenhagen, Denmark**
1912 Ashot Margaryan
1913

1914 **Anthropology Department, University of Utah, USA**
1915 Melissa Illardo
1916

1917 **Department of Geology, Lund University, Lund, Sweden**
1918 Anne Birgitte Nielsen
1919

1920 **Department of Archaeology and Ancient History, Lund University, Lund, Sweden**
1921 Lars Larsson
1922

1923 **Tårnby Gymnasium og HF, Kastrup, Denmark**
1924 Mikkel Ulfeldt Hede
1925

1926 **Department of Health Technology, Section of Bioinformatics, Technical University of Denmark,**
1927 **Kongens Lyngby, Denmark**
1928 Gabriel Renaud
1929

1930 **Department of Archaeology and Heritage Studies, Aarhus University, Aarhus, Denmark**
1931 Niels N. Johannsen, Rikke Maring
1932

1933 **Institute of Biological Psychiatry, Mental Health Centre Sct Hans, Copenhagen University Hospital,**
1934 **Denmark**
1935 Andrew J. Schork, Anders Rosengren & Thomas Werge
1936

1937 **Department of Clinical Medicine, University of Copenhagen, 2200 Copenhagen N, Denmark**
1938 Thomas Werge
1939

1940 **Neurogenomics Division, The Translational Genomics Research Institute (TGEN), Phoenix, AZ, USA**
1941 Andrew J. Schork
1942

1943 **Terra Ltd., Letchik Zlobin St. 20, Voronezh, 394055, Russian Federation**
1944 Andrey Skorobogatov, Ruslan Turin
1945

1946 **Department of Archaeology, University of Exeter, Exeter, UK**
1947 Alan K. Outram & Catriona J. McKenzie
1948

1949 **Institute of Archaeology and Ethnography, Siberian Branch of the Russian Academy of Sciences,**
1950 **Novosibirsk, Russian Federation**
1951 Aleksey A. Timoshchenko, Dmitri V. Pozdnyakov, Liudmila N. Mylnikova, Marina S. Nesterova &
1952 Vyacheslav I. Molodin
1953

1954 **Institute of Ethnology and Anthropology, Russian Academy of Sciences, Moscow, Russian Federation**
1955 Elizaveta V. Veselovskaya, Sergey Vasilyev
1956

1957 **Centre for Egyptological studies, Russian Academy of Sciences, Moscow, Russia**
1958 Sergey Vasilyev
1959

1960 **Research Institute and Museum of Anthropology, Lomonosov Moscow State University, Mokhovaya**
1961 **str. 11, Moscow, Russian Federation**
1962 Alexandra Buzhilova, Natalia Berezina, Svetlana Borutskaya
1963

1964 **Department of Environmental Biology, Sapienza University of Rome, Rome, Italy**
1965 Alfredo Coppa, Dušan Borić
1966

1967 **Peter the Great Museum of Anthropology and Ethnography (Kunstkamera), Russian Academy of**
1968 **Sciences, Saint Petersburg, Russian Federation**
1969 Alisa Zubova & Vyacheslav Moiseyev
1970

1971 **CIAS, Department of Life Science, University of Coimbra, Coimbra, Portugal**
1972 Ana Maria Silva
1973

1974 **UNIARQ, University of Lisbon, Lisbon, Portugal**
1975 Ana Maria Silva & João Zilhão
1976

1977 **Kostanay Regional University A. Baitursynov, Kostanay, Kazakhstan**
1978 Andrey V. Logvin, Irina V. Shevnina
1979

1980 **Vesthimmerlands Museum, Søndergade 44, Aars, Denmark**
1981 Bjarne Henning Nielsen
1982

1983 **Museum Nordsjælland, Frederiksgade 9, 3400 Hillerød**
1984 Per Lotz, Søren Anker Sørensen, Thomas Jørgensen
1985

1986 **Museum Vestsjælland, Klosterstræde 18, 4300 Holbæk, Denmark**
1987 Per Lotz
1988

1989 **Vendsyssel Historiske Museum, DK-9800 Hjørring, Denmark**
1990 Per Lysdahl, Sidsel Wåhlin
1991

1992 **Moesgaard Museum, Moesgård Allé 15, Højbjerg, Denmark**
1993 Søren H. Andersen
1994

1995 **Grupo EvoAdapta, Departamento de Ciencias Históricas, Universidad de Cantabria, Santander, Spain**
1996 Borja González-Rabanal
1997

1998 **Museu de Ciències Naturals de Barcelona, Barcelona, Spain**
1999 Carles Lalueza-Fox
2000

2001 **Institute of Evolutionary Biology, CSIC-Universitat Pompeu Fabra, Barcelona, Spain**
2002 Carles Lalueza-Fox
2003
2004 **ICREA, University of Barcelona, Barcelona, Spain**
2005 João Zilhão
2006
2007 **Departamento de Prehistoria y Arqueología Department, Universidad Autónoma de Madrid,**
2008 **Madrid, Spain**
2009 Concepción Blasco, Corina Liesau, Patricia Ríos
2010
2011 **Department of Biology, University of Rome "Tor Vergata", Rome, Italy**
2012 Cristina Martinez-Labarga, Olga Rickards
2013
2014 **Department of History, Humanities and Society, University of Rome "Tor Vergata", Rome, Italy**
2015 Mario Federico Rolfo
2016
2017 **Instituto Internacional de Investigaciones Prehistóricas de Cantabria, Universidad de Cantabria,**
2018 **Santander, Spain**
2019 David Cuenca-Solana, Igor Gutiérrez Zugasti & Manuel González-Morales
2020
2021 **Centre de Recherche en Archéologie, Archeosciences, Histoire (CReAAH), UMR-6566 CNRS, Rennes,**
2022 **France**
2023 David Cuenca-Solana
2024
2025 **Georgian National Museum, Tbilisi, Georgia**
2026 David O. Lordkipanidze
2027
2028 **Tbilisi State University, Tbilisi, Georgia**
2029 David O. Lordkipanidze
2030
2031 **IPND, Tyumen Scientific Centre, Siberian Branch of the Russian Academy of Sciences, Tyumen,**
2032 **Russian Federation**
2033 Dmitri Enshin, Olga Poshekhonova, Svetlana N. Skochina
2034
2035 **Departament de Prehistòria, Arqueologia i Història Antiga, Universitat de València, València, Spain**
2036 Domingo C. Salazar-García, J.Emili Aura Tortosa
2037
2038 **Department of Geological Sciences, University of Cape Town, Cape Town, South Africa**
2039 Domingo C. Salazar-García
2040
2041 **Department of Anthropology, New York University, New York, USA.**
2042 Dušan Borić
2043
2044 **Institute of Humanities, Ivanovo State University, Ivanovo, Russian Federation**
2045 Elena Kostyleva
2046
2047 **Saryarka Archaeological Institute, Buketov Karaganda University, Karaganda, Kazakhstan**

2048 Emma R. Usmanova
2049
2050 **South Ural State University, Chelyabinsk, Russia**
2051 Emma R. Usmanova
2052
2053 **The Saxo Institute, University of Copenhagen, Copenhagen, Denmark**
2054 Erik Brinch Petersen, Rune Iversen
2055
2056 **Museum Østjylland, Stemannsgade 2, Randers, Denmark**
2057 Esben Kannegaard, Lutz Klassen
2058
2059 **Soprintendenza Archeologia Belle Arti e Paesaggio per la Città Metropolitana di Bari, Via Pier**
2060 **l'Eremita, 25, 70122, Bari, Italy**
2061 Francesca Radina
2062
2063 **Soprintendenza per i Beni Archeologici delle Marche, Via Birarelli 18, 60100, Ancona, Italy**
2064 Mara Silvestrini
2065
2066 **Soprintendenza Archeologia, Belle Arti e Paesaggio per la provincia di Cosenza, Cosenza, Italy**
2067 Paola Aurino
2068
2069 **UMR 5199 PACEA, CNRS, Université de Bordeaux, 33615 Pessac, France**
2070 Henri Duday, Patrice Courtaud
2071
2072 **Institute of Archaeology, National Academy of Sciences of Ukraine, Kyiv, Ukraine**
2073 Inna Potekhina
2074
2075 **National University of Kyiv-Mohyla Academy, Kyiv, Ukraine**
2076 Inna Potekhina
2077
2078 **Collège de France, 75231 Paris cedex 05, France**
2079 Jean Guilaine
2080
2081 **Odense City Museums, Overgade 48, Odense, Denmark**
2082 Jesper Hansen
2083
2084 **Museum Sydøstdanmark, Algade 97, 4760 Vordingborg, Denmark**
2085 Kristoffer Buck Pedersen
2086
2087 **Institute of Archaeology and Ethnology, Polish Academy of Sciences, Kraków, Poland**
2088 Krzysztof Tunia, Piotr Włodarczak
2089
2090 **CNRS UMR 5608, Toulouse Jean Jaurès University, Maison de la Recherche, 5 Allées Antonio**
2091 **Machado, 31058 Toulouse, Cedex 9, France**
2092 Laure Metz & Ludovic Slimak
2093
2094 **ARGEA Consultores SL, C. de San Crispín, Madrid, Spain**

2095   Jorge Vega, Roberto Menduiña

2096

2097   **Institute of Molecular Biology, National Academy of Sciences, Yerevan, Armenia**
2098   Levon Yepiskoposyan

2099

2100   **Russian-Armenian University, Yerevan, Armenia**
2101   Levon Yepiskoposyan

2102

2103   **Institute of Archaeology and Ethnography, National Academy of Sciences, Yerevan, Armenia**
2104   Ruben Badalyan

2105

2106   **HistorieUdvikler, Gl. Roesnaesvej 27, DK-4400 Kalundborg, Denmark**
2107   Lisbeth Pedersen

2108

2109   **Department of history and cultural heritage, University of Siena, Siena, Italy**
2110   Lucia Sarti, Mauro Calattini

2111

2112   **Centre d'Anthropobiologie et de Génomique de Toulouse, CNRS UMR 5288, Université Paul Sabatier,**
2113   **Toulouse, France**
2114   Ludovic Orlando

2115

2116   **Västergötlands museum, Stadsträdgården, Skara, Sweden**
2117   Maria Vretemark

2118

2119   **Cabinet of Anthropology, Tomsk State University, Tomsk, Russian Federation**
2120   Marina P. Rykun

2121

2122   **Institute for Eastern Research, Adam Mickiewicz University in Poznań, Poznań, Poland**
2123   Marzena H. Szmyt

2124

2125   **Institute of Archaeology, Jagiellonian University, Ul. Gołębia 11, 31-007, Kraków, Poland**
2126   Marcin Przybyła

2127

2128   **Zoological Institute of Russian Academy of Sciences, Universitetskaya nab. 1, 199034, St. Petersburg,**
2129   **Russian Federation**
2130   Mikhail Sablin

2131

2132   **Department of Anthropology, Czech National Museum, Prague, Czech Republic**
2133   Miluše Dobisíková

2134

2135   **Department of Health and Nature, University of Greenland, Greenland**
2136   Morten Meldgaard

2137

2138   **The Viking Ship Museum, Vindeboder 12, Roskilde, Denmark**
2139   Morten Johansen, Otto Christian Uldum

2140

2141   **Archaeology Institute, University of Highlands and Islands, Scotland, UK**

2142   Nick Card
2143
2144   **Laboratory for Experimental Traceology, Institute for the History of Material Culture of the Russian**
2145   **Academy of Sciences, Dvortsovaya nab., 18, 191186, St. Petersburg, Russian Federation**
2146   Olga V. Lozovskaya
2147
2148   **Scientific Research Center "Baikal region", Irkutsk State University; 1, K. Marx st., Irkutsk, 664003,**
2149   **Russian Federation**
2150   Nikolai A. Saveliev
2151
2152   **Paleoecology Laboratory, Institute of Plant and Animal Ecology, Ural Branch of the Russian Academy**
2153   **of Sciences, Ekaterinburg, Russian Federation**
2154   Pavel Kosintsev
2155
2156   **Department of History of the Institute of Humanities, Ural Federal University, Ekaterinburg, Russian**
2157   **Federation**
2158   Pavel Kosintsev
2159
2160   **Centre for the Study of Early Agricultural Societies, Department of Cross-Cultural and Regional**
2161   **Studies, University of Copenhagen, 2300 Copenhagen, Denmark**
2162   Peder Mortensen
2163
2164   **Museum of Cultural History, University of Oslo, P.O. Box 6762. St. Olavs Plass NO-0130 Oslo,**
2165   **Norway**
2166   Per Persson
2167
2168   **ArchaeoScience, GLOBE Institute, University of Copenhagen, Copenhagen, Denmark**
2169   Pernille Bangsgaard
2170
2171   **Department of History, University of Santiago de Compostela, Spain**
2172   Pilar Prieto Martinez
2173
2174   **Lipetsk Regional Scientific Public Organisation "Archaeological Research", Lipetsk, Russian**
2175   **Federation**
2176   Roman V. Smolyaninov
2177
2178   **Laboratory for Archaeological Chemistry, Department of Anthropology, University of Wisconsin-**
2179   **Madison, Madison, USA**
2180   T. Douglas Price
2181
2182   **Nizhny Tagil State Socio-Pedagogical Institute, Nizhny Tagil, Russia**
2183   Yuri B. Serikov
2184
2185   **Institute for History of Medicine, First Faculty of Medicine, Charles University, Prague, Czech**
2186   **Republic**
2187   Vaclav Smrcka
2188

Centre for Archaeological Research Toraighyrov University, Pavlodar, Kazakhstan
Victor Merz

Malmö Museer, Malmöhusvägen 6, Malmö, Sweden
Yvonne Magnusson

Institute of Statistical Sciences, School of Mathematics, University of Bristol, Bristol, UK
Daniel J. Lawson

Novo Nordisk Foundation Centre for Protein Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen N, Denmark
Simon Rasmussen

MARUM, University of Bremen, Bremen, Germany
Eske Willerslev

Contributions
M.E.A., M.S., A.R.-M., E.K.I.-P., A.F., W.B., and A.I. contributed equally to this work. M.E.A., M.S., T.S.K., R.D., R.N., O.D., T.W., F. Racimo, K.K. and E.W. led the study. M.E.A., M.S., A.F., C.L.-F., R.N., T.W., K.K. and E.W. conceptualised the study. M.E.A., M.S., H.S., L.O., T.S.K., R.D., R.N., O.D., T.W., F. Racimo, K.K. and E.W. supervised the research. M.E.A., L.O., R.D., R.N., T.W., K.K. and E.W. acquired funding for research. A.F., J.S., K.G.S., M.L.S.J., M.U.H., A.A.T., A.C., A.Z., A.M.S., A.J.H, A.G., A.V.L., B.H.N., B.G.R, C.B., C.L., C.M-L., D.V., D.C.-S., D.L., D.N., D.C.S.-G., D.B., E.K., E.V.V., E.R.U., E. Kannegaard, F. Radina, H.D., I.G.Z., I.P., I.V.S., J.G., J.H., J.E.A.T., J.Z., J.V., K.B.P., K.T., L.N., L.L., L.M., L.Y., L.P., L. Sarti, L. Slimak, L.K., M.G.M., M. Silvestrini, M.V., M.S.N., M.P.R., M.H.S., M.P., M.C., M. Sablin, N.C., O.P., O.R., O.V.L., P.A., P.K., P.C., P. Ríos, P. Lotz, P. Lysdahl, P.P., P.B., P.d.B.D., P.V.P., P.P.M., P.W., R.V.S., R. Maring, R. Menduiña, R.B., R.T., S.V., S.W., S.B., S.N.S., S.A.S., S.H.A., T.D.P., T.J., Y.B.S., V.I.M., V.S., V.M, Y.M. and N.L. were involved in sample collection. M.E.A., M.S., A.R.-M., E.K.I.-P., W.B., A.I., J.S., A.P., B.S.d.M., M.I., L.V., A.J. Stern, C.G., F.E.Y, D.J.L., T.S.K., R.D., R.N., O.D., F. Racimo, K.K. and E.W. were involved in developing and applying methodology. J.S., C.G. and L.V. led the DNA laboratory work research component. K.G.S. led bioarchaeological data curation. M.E.A., M.S., A.R.-M., E.K.I.-P., W.B., A.I., A.P., B.S.d.M., B.S.P., A.S.H., R.A.H., T.V., H.M., A.M., A.V., A.B.N., P. Rasmussen, G.R., A. Ramsøe, A.S., A.J. Schork, A. Rosengren, C.J.M., I.A., L.Z., R.Maring, V.S., V.A., P.H.S, S.R., T.S.K., O.D. and F. Racimo undertook formal analyses of data. M.E.A., M.S., A.R.-M., E.K.I.-P., A.F., W.B., A.I., K.G.S., D.J.L., P.H.S., T.S.K., and F. Racimo drafted the main text (M.E.A. and M.S. led this). M.E.A., M.S., A.R.-M., E.K.I.-P., A.F., W.B., A.I., K.G.S., A.P., B.S.d.M., B.S.P, A.S.H., R. Macleod, R.A.H., T.V., M.F.M., A.B.N., M.U.H., P. Rasmussen, A.J. Stern, N.N.J., H.S., G.S., A. Ramsøe, A.S., A. Rosengren, A.K.O., A.B., A.C., A.G., A.V.L., A.B.G., C.J.M., D.C.S.-G., E. Kostyleva, E.R.U., E. Kannegaard, I.G.Z., I.P., I.V.S., J.G., J.H., J.E.A.T., L.Z, L.Y., L.P., L.K., M.B., M.G.M., M.V., M.P.R., M.J., N.B., O.V.L., O.C.U., P.K., P. Lysdahl, P.B., P.W., R.V.S., R. Maring, R.B., R.I., S.V., S.W., S.B., S.H.A., T.J., V.S., D.J.L., P.H.S., S.R., T.S.K., O.D. and F. Racimo drafted supplementary notes and materials. M.E.A., M.S., A.R.-M., E.K.I.-P., A.F., W.B., A.I., G.G.S., A.S.H., M.L.S.J., F.D., R. Macleod, L. Sørensen, P.O.N., R.A.H., T.V., H.M., A.M., N.N.J., H.S., A. Ramsøe, A.S., A.J. Schork, A. Ruter, A.K.O., B.H.N., B.G.R., D.C.-S., D.C.S.-G., I.G.Z., I.P., J.G., J.E.A.T., L.Z., L.O., L.K., M.G.M., P.d.B.D., R.I., S.A.S., D.J.L., P.H.S., T.S.K., R.D., R.N., O.D., T.W., F. Racimo, K.K. and E.W. were involved in reviewing drafts and editing (M.E.A., A.F., K.G.S., F.D., R. Macleod, H.M. and T.V. led this). All co-authors read, commented on, and agreed upon the submitted manuscript.

2236
2237 <u>Corresponding authors</u>
2238
2239 Correspondence to Morten E. Allentoft (morten.allentoft@curtin.edu.au), Martin Sikora
2240 (martin.sikora@sund.ku.dk), Eske Willerslev (ew482@cam.ac.uk).
2241

2242 **Ethics declarations**

2243
2244 <u>Competing interests</u>
2245
2246 The authors declare no competing interests.