Microbial population dynamics decouple nutrient affinity from environmental concentration

Justus Wilhelm Fink*

Institute of Integrative Biology, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland

Noelle A. Held

Institute of Biogeochemistry and Pollutant Dynamics,
Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland and
Department of Environmental Microbiology, Swiss Federal Institute
of Aquatic Science and Technology (Eawag), Dübendorf, Switzerland

Michael Manhart†

Institute of Integrative Biology, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland
Department of Environmental Microbiology, Swiss Federal Institute of
Aquatic Science and Technology (Eawag), Dübendorf, Switzerland and
Center for Advanced Biotechnology and Medicine and Department of Biochemistry and Molecular Biology,
Robert Wood Johnson Medical School, Rutgers University, Piscataway, NJ, USA
(Dated: October 3, 2022)

How the growth rate of a microbial population depends on the environmental availability of chemical nutrients and other resources is a fundamental question in microbiology. Models of this dependence, such as the widely-used Monod model, are generally characterized by a threshold concentration (or affinity) of the resource, above which the population can grow near its maximum rate. What values should we expect for these thresholds, and how should they depend on the environmental concentration of the resource? We perform a survey of resource threshold data across a wide range of organisms and resources. We find that the thresholds vary across orders of magnitude, even for the same organism and resource. To explain this variation, we develop an evolutionary model to show that demographic fluctuations (genetic drift) can constrain the adaptation of resource thresholds. We find that this effect fundamentally differs depending on the type of population dynamics: populations undergoing periodic bottlenecks of fixed size will adapt their thresholds in proportion to the environmental resource concentration, but populations undergoing periodic dilutions of fixed size will evolve thresholds that are largely decoupled from the environmental concentration. Our model not only provides testable predictions for laboratory evolution experiments, but it also reveals how an evolved resource threshold may not reflect the organism's environment. In particular, this explains how organisms in resource-rich environments can still evolve fast growth at low resource concentrations. Altogether our results demonstrate the critical role of population dynamics in shaping fundamental ecological traits.

Keywords: Microbial evolution | Monod model | resource competition | nutrient affinity | selection-drift balance

INTRODUCTION

Microbial populations rely on a wide range of resources, including chemical nutrients such as sugars, minerals, and metals, as well as space, light, and prey [1]. These resources vary in abundance or concentration across time and environments, which typically elicits differences in growth rates [2–4]. A significant literature discusses how natural populations can be classified as oligotrophs or copiotrophs [4–6], that differ, among other things, in their growth rate response to resource concentration. The most widely-used quantitative model of the relationship between growth rate and resource concentra-

tion is attributed to Jacques Monod [7]. In the Monod model, growth rate increases linearly with resource concentration below a threshold concentration, and then saturates at high concentrations of the resource. The threshold concentration, also sometimes known as the resource affinity, therefore plays a key role in determining the ability of the population to grow on scarce resources. This suggests that lower resource concentrations in the environment may drive populations to evolve commensurately lower thresholds [8, 9], one of the main predictions of resource-ratio theory [10–12]. Quantitative models and data for the dependence of growth rate on resource concentration are important both for predicting the behavior of a population under different environmental conditions [13–15], as well as for inferring the natural environmental niche from evolved traits of the population. This inverse approach has been used, for example, to infer separate niches for ammonia-oxidizing archaea and bacteria in the global nitrogen cycle based on kinetic parameters

---

* To whom correspondence should be addressed. Email: justus.fink@env.ethz.ch
† To whom correspondence should be addressed. Email: mmanhart@rutgers.edu

for resource consumption [16–19].

Even though these concepts have been central elements of microbiology and ecology for decades, there is limited experimental evidence that directly demonstrates the evolution of growth rate dependence on resources. Continuous culture for 200–300 generations led to improved growth rate at low glucose concentrations for *Escherichia coli* [20, 21] and *Saccharomyces cerevisiae* [22], but these changes were not clearly attributable to genetic (rather than physiological) adaptation. The Long-Term Evolution Experiment (LTEE) of *E. coli* found that the glucose threshold actually increased over the first 2000 generations, although the growth rate at much higher glucose concentrations significantly increased as well [23]. More recently, Bernhardt et al. [12] observed adaptation in the phosphorus threshold of *Chlamydomonas reinhardtii* when limited for phosphorus, but they did not obtain consistent outcomes for nitrogen and light. Perhaps the most explicit evidence so far is from Hart et al. [24], who found that a synthetic auxotroph strain of *S. cerevisiae* significantly reduced its threshold for lysine through genetic adaptations.

While laboratory experiments can test the basic principle, mathematical models are better suited to exploring the wide range of environments necessary to establish the link between environment and evolved traits. Previous modeling studies on this topic have focused on how tradeoffs in growth rate at low versus high resource concentrations define an optimum strategy for a single strain [13] or can facilitate coexistence of multiple strains or species when resource concentrations fluctuate [25, 26]. More recent work has shown how this coexistence can spontaneously evolve if such tradeoffs constrain the effects of mutations [27, 28]. However, the evidence for these tradeoffs, especially on spontaneous mutations, is limited [27–30]. Thus their importance for explaining the evolved variation in growth rate dependence, especially the resource threshold, is unclear.

Here we address this problem using both empirical and modeling approaches. We first perform a survey of data for the growth rate dependence on resource concentration across a wide range of organisms and resources. We find that the measured resource thresholds vary over orders of magnitude, even within some single species on the same resource, such as *E. coli* strains on glucose. We also find no evidence for tradeoffs between growth rate at low versus high resource concentrations. To better understand the potential causes of this variation, we model evolution for populations with a single limiting resource under feast-and-famine conditions (batch dynamics with fixed biomass or fixed dilution factor) and steady-state growth (chemostat dynamics). We show how demographic fluctuations, known as genetic drift, inhibit selection on lower thresholds, which leads to a general relationship between the evolved resource threshold, environmental resource concentration, and the effective population size. Using this result, we determine that populations with fixed-bottleneck batch dynamics will evolve resource thresh-olds that are proportional to the environmental resource concentration, but populations with fixed-dilution batch dynamics evolve thresholds that are practically independent of the environment. Besides providing a testable theory for laboratory evolution experiments, our results help to explain how species evolving under high concentrations can maintain fast growth at low concentrations and why evolved resource thresholds may not reflect the environment of origin.

RESULTS

The Monod model quantifies growth rate dependence on resource concentration

Consider a population of microbes consuming a resource; we will generally focus on chemical nutrients such as carbon or nitrogen sources, but some aspects of the model apply to other types of resources as well (e.g., prey or light). While microbes consume many different resources simultaneously [31, 32], for simplicity here we assume only a single resource limits growth (Supplementary Information Sec. S1). The best-known dependence of population growth rate g on resource concentration R is the Monod model [7]:

$$g(R) = g^{\mathrm{max}} \cdot \frac{R}{R + K}, \qquad (1)$$

where g^{max} is the maximum growth rate — achieved when the resource is unlimited — and K is a threshold concentration for the resource at which growth rate is slowed to half its maximum (Fig. 1). Decreasing the threshold K therefore allows the population to grow faster at lower resource concentrations. The resource threshold K is not to be confused with a related but distinct concept of R^* from resource-ratio theory [10, 12]. Note that the Monod model of Eq. (1) is used to describe both steady state [12] and non-steady-state [25, 28] relationships between growth rate and environmental resource concentration. While there are many alternative models of how growth rate depends on resource concentration (Supplementary Information Sec. S2, Table S1), we focus on the Monod model due to its wider usage and available data.

The parameter K is sometimes labeled as the affinity for the resource [33], but this is potentially misleading as K is inversely proportional to ability to grow on the resource. We instead use the term *specific affinity* to refer to the parameter combination g^{max}/K, which measures how much the growth rate increases per unit change in resource concentration, starting from a low concentration [34]. The specific affinity is therefore a common measure for oligotrophic growth ability [9, 16, 19, 33]. Note that both K and g^{max} are required to fully characterize the growth rate dependence; for example, the specific affinity g^{max}/K alone does not suffice because while it describes the growth rate response at low concentrations,

FIG. 1. **Monod model of growth rate dependence on resource concentration.** The population growth rate $g(R)$ as a function of the external resource concentration R for two hypothetical strains: a wild-type (green) and a derived mutant strain (orange), with equal maximum growth rates ($g^{\max} = 1$) but different resource thresholds ($K_{\mathrm{wt}} = 5$, $K_{\mathrm{mut}} = 3$). The inset shows a magnified view at low concentrations near K_{wt} and K_{mut} (dotted vertical lines). Note that the growth rates do not fully overlap at the highest concentration shown, but eventually converge to the same value $g^{\max}$ outside the range of this plot.

it does not define the threshold for a low concentration (which is determined separately by K). Since we are primarily interested in how these traits evolve in relation to the environmental concentration R, we focus primarily on the threshold K since, being a concentration as well, one can directly compare it to R.

One can derive the Monod model of Eq. (1) by modeling biomass growth as a two-step process, in which uptake of the external resource into the cell occurs at a rate proportional to the external concentration R [35]. However, the dependence of growth rate on resource concentration expressed by Eq. (1) is surprisingly robust to additional model complexities [36, 37], albeit with the resulting traits $g^{\max}$ and K being emergent properties of whole cells or populations. In particular, the threshold K is not equivalent to the Michaelis-Menten constant for resource uptake kinetics [36, 38, 39], despite the mathematical similarity between the Michaelis-Menten and Monod models (Eq. 1); this is because the Monod model describes the whole process of producing new biomass, of which uptake is just one step.

Resource thresholds vary widely across resources and organisms

To explore the diversity of microbial growth strategies, we have compiled 247 measurements of resource thresholds K from previously-published studies (Methods; Dataset S1; Fig. S1), substantially extending previ-

ous surveys [40–43]. Figure 2A shows an overview of this data, sorted by resource. The data includes a wide range of resources, with phosphate, glucose, and nitrate having the largest number of measurements due to their emphasis in marine and laboratory systems. Organisms include prokaryotes and eukaryotes as well as autotrophs and heterotrophs (marked by different symbols in Fig. 2A).

Measured values of the threshold K vary over several orders of magnitude, ranging from below 10^{-6} µM (for thiamine and vitamin B12) to above 10^4 µM (for one glucose measurement). This variation is not attributable to measurement uncertainties, which never exceeded 20% in the studies that reported them. It also is not an artifact of technical aspects of the measurements (Fig. S2) such as temperature (linear regression, $R^2 \approx 0.089$, $p \approx 1.2 \times 10^{-5}$) or experimental method (linear regression, $R^2 \approx 0.160$, $p \approx 1.3 \times 10^{-3}$), nor does the variation appear to be systematically biased by experimental design such as the degree of pre-acclimation to the growth medium (Fig. S3). We furthermore find no evidence for a major bias from simultaneous limitation for other resources besides the focal resource (colimitation; Supplementary Information Sec. S1).

Instead, most variation of thresholds K corresponds to variation in the identity of the organisms and resources themselves (Fig. S2A). Figure 2B shows a subset of measurements on glucose, which have systematic differences in K between taxa. For example, measurements of *S. cerevisiae* and *Streptococcus* almost all have K values higher than those of *E. coli* (Mann-Whitney U test, $p \approx 1.40 \times 10^{-6}$). Phosphate and silicate similarly show significant variation between species (Fig. 2C,D), as do nitrate and ammonium (Fig. S4). Even within some taxa, there is large variation of K; glucose K in *E. coli* varies over three orders of magnitude (Fig. 2B). This variation within a single resource and taxon does not appear to be explained by technical covariates of the measurements (Fig. S2B), but rather corresponds to genetically-distinct strains of *E. coli* (Fig. S5), suggesting that even subspecies-level genetic variation can lead to significant differences in the threshold K. Indeed, Ferenci [45] reported single target genes, like the membrane-associated lamB or the stress-factor rpoS, that affect the resource threshold of *E. coli* on glucose when mutated. The genetic differences in our dataset are unknown, but we grouped *E. coli* measurements by strain labels to find reproducible glucose thresholds within strains (e.g., ML30, see Fig. S5A).

How can we explain this wide variation in resource thresholds? Intuitively, we expect evolution to reduce K, since mutations that reduce K increase growth rate (Eq. (1)). For example, Fig. 1 shows the growth rate dependence for a hypothetical wild-type strain (green line) and a mutant (orange) with lower threshold K. Since the mutant has a greater relative growth rate advantage at low resource concentrations, there could be stronger selection pressure to reduce K at those low concentrations. This is hinted by some patterns in the data: for

FIG. 2. **Survey of measured resource thresholds.** (A) Complete set of resource thresholds K in our survey, grouped by resource (in decreasing order of number of data points). Each point represents a different measurement; color indicates whether the organism is a prokaryote (green) or eukaryote (orange), and shape indicates whether the organism can grow as an autotroph (square) or only as a heterotroph (circle). Dashed lines mark concentrations of one molecule per cell for approximate prokaryotic and eukaryotic cell volumes [44]. (B) Subset of K measurements from panel A for glucose, grouped by taxon (only those with at least two measurements). We use the taxonomic identity given in the original publications, where an ending in *sp.* means the isolate is a representative of the genus but was not identified at the species level. Symbols are the same as in panel A. (C) Subset of K measurements from panel A for phosphate, grouped by taxon (with at least three measurements). (D) Subset for silicate, grouped by taxon (with at least two measurements). See the Supplemental Information for additional plots with K measurements for nitrate (Fig. S4A) and ammonium (Fig. S4B).

example, *E. coli* often grows in mammalian large intestines where there are few simple sugars such as glucose, while *S. cerevisiae* and *Streptococcus* often grow in high-sugar environments (fruit and the oral microbiome, respectively) [46, 47], which could explain their large difference in glucose thresholds.

Variation in specific affinity has trends similar to those of the resource threshold

Since K alone does not define the growth rate at low resource concentrations, it is essential to consider the maximum growth rate g^{max} or specific affinity g^{max}/K as well. We show the variation in maximum growth rate g^{max} across resources in Fig. 3A (reported for 97.5% of all entries for thresholds K; Dataset S1). The most striking feature of this data is that while maximum growth rates g^{max} vary less between resources than do thresholds K (compare Figs. 3A and 2A), there is a clear bimodal-

ity between fast-growing heterotrophs (circles) and slow-growing autotrophs (squares). This suggests that the maximum growth rate depends primarily on the background medium and organism identity rather than on the focal resource, which makes sense given that g^{max} is a property of the population when the focal resource is in excess. Indeed, a closer look at the covariation between g^{max} and K in autotrophs (Fig. 3B) reveals that resources have comparable distributions of g^{max} but stratify in terms of thresholds, with the lowest values for phosphate. In particular, the distributions for phosphate and nitrate are indistinguishable in terms of maximum growth rate (Mann-Whitney U test, $p = 0.0801$), but clearly different in terms of resource threshold (Mann-Whitney U test, $p = 1.28 \times 10^{-12}$). Also, the species differences in maximum growth rate on glucose and phosphate are less pronounced (Fig. S6) and more of the variation can be explained by experiment temperature (Fig. S7) compared to variation in K.

We can also compute the specific affinity g^{max}/K for each data point. Figure S8 shows that the variation in specific affinity is similar to variation in K: the variation spans orders of magnitude, even for single species, and there are systematic differences between taxa (e.g., $E.\ coli$ compared to $S.\ cerevisiae$ and $Streptococcus$; Mann-Whitney U test, $p \approx 1.20 \times 10^{-6}$; Fig. S8B). The similarity in patterns of variation between the threshold and specific affinity is because variation in g^{max}/K is dominated by variation in K (Fig. S7B); on a logarithmic scale, g^{max}/K depends on additive contributions from g^{max} and K, and variation in K is much larger than variation in g^{max} (compare Figs. 2A and 3A).

There is no evidence for a tradeoff between resource threshold and maximum growth rate

Many previous studies have considered the possibility of tradeoffs between g^{max} and K (positive correlation), such that genotypes growing faster with abundant resources will grow slower when resources are scarce [13, 25–28]. If this were true, evolution at high resource concentrations may select for increasing maximum growth rate g^{max} at the expense of the resource threshold K, leading to high values of K. If we consider all organisms and resources in our data set, we do find a significant positive correlation between g^{max} and K (Spearman $\rho \approx 0.39$, $p \approx 5.7 \times 10^{-10}$; Fig. 3B). However, this correlation is an artifact of the biased sampling of organism-resource pairs, which are dominated by fast-growing heterotrophs on glucose (which tend to have higher K thresholds) and slow-growing autotrophs on other resources (which tend to have lower K thresholds compared to glucose); the correlation disappears when we separate heterotrophs (Fig. S9A,B) from autotrophs (Fig. S9C,D). If we further separate individual resources, we see no significant correlations for phosphate, nitrate, ammonium, or glucose across organisms

(Figs. 3C,D and S9E–H), while there is actually a negative correlation (opposite of a tradeoff) for silicate g^{max} and K (Spearman $\rho \approx -0.56$, $p \approx 0.0025$; Fig. 3E). In Fig. 3F we test the covariation of g^{max} with K for two individual species ($E.\ coli$ and $S.\ cerevisiae$) for a single resource (glucose). The $E.\ coli$ data shows a positive correlation indicative of a tradeoff, but it has modest magnitude and low statistical significance (Spearman $\rho \approx 0.26$, $p \approx 0.26$). $Saccharomyces\ cerevisiae$, on the other hand, shows a positive correlation between the two traits (Spearman $\rho \approx -0.75$, $p \approx 0.008$). The lack of tradeoff appears irrespective of experimental method (i.e., batch or chemostat; Fig. S3B) and also holds when comparing the maximum growth rate g^{max} to the specific affinity g^{max}/K (Fig. S10).

Much of the previous literature arguing for tradeoffs in these traits based their evidence on measurements for resource uptake kinetics [27, 28, 30, 48] rather than on population growth as we consider here. However, we find little to no correspondence between traits of uptake kinetics with traits of population growth in data points where we have measurements for both (Fig. S11) [43], consistent with previous analyses [36, 38]. It is therefore not surprising that the observed tradeoffs in uptake do not translate to tradeoffs in growth. For example, Litchman et al. [30] reported a tradeoff between uptake traits for nitrate, but we see no correlation in growth traits for nitrate (Spearman $\rho \approx 0.03$, $p \approx 0.84$; Figs. 3D and S10C). Altogether the absence of evidence for a systematic correlation between K and g^{max} suggests that selection for g^{max} does not explain the evolved variation in K.

Models of population dynamics with mutations to resource threshold

To test how the environmental resource concentration shapes the evolution of the resource threshold K, we turn to a model of population dynamics with mutations altering traits of the Monod growth rate dependence (Methods; Supplementary Information Secs. S3–S5; Table S2). We consider a microbial population consisting of a wild-type and a mutant, with biomasses $N_{\mathrm{wt}}(t)$ and $N_{\mathrm{mut}}(t)$ that vary over time t. They grow at rates depending on the resource concentration R according to the Monod model (Eq. (1)), but with potentially different values of the traits g^{max} and K depending on the effect of the mutation [25, 28]. The relative rate at which the mutant increases or decreases in frequency compared to the wild-type is given by the selection coefficient s (Supplementary Information Sec. S6) [49, 50]. We show that s decomposes into two additive terms

$$s \approx s_{\mathrm{high}} + s_{\mathrm{low}}, \tag{2}$$

where s_{high} measures selection on growth at high resource concentrations, and is therefore proportional to variation in the maximum growth rate g^{max}, while s_{low} measures

FIG. 3. **Survey of maximum growth rates and trait correlations.** (A) Empirical maximum growth rates g^{max} for the microbial isolates in our survey. There are slightly fewer data points for maximum growth rate compared to thresholds in Fig. 2A, since some publications only reported the resource threshold. Markers indicate whether the organisms can grow as an autotroph (square) or only as a heterotroph (circle); colors indicate if the isolate is prokaryotic (green) or eukaryotic (orange). Dashed lines mark reference doubling times. (B) Covariation of maximum growth rate g^{max} and resource threshold K across the entire set of isolates from panel A. Here colors indicate the limiting resource, with the number of measurements n given in brackets. Marker shapes (squares are autotrophs, circles are heterotrophs) are the same as in panel A. We compute the Spearman rank correlation ρ and p-value across the pooled set of isolates. (C) Subset of measurements from panel B for phosphate (only autotroph isolates shown). (D) Subset of measurements from panel B for nitrate. (E) Subset of measurements from panel B for silicate. (F) Covariation between maximum growth rate g^{max} and glucose threshold K for measurements of *E. coli* (green) and *S. cerevisiae* (orange), with Spearman rank correlations ρ and p-values by species.

selection on growth at low resource concentrations, and is therefore proportional to variation in the threshold K (Figs. S12–S15; Supplementary Information Secs. S7–S9).

We consider selection in three prototypical regimes of population dynamics. In the first case, the population grows as a batch culture with serial transfers (Supplementary Secs. S3). That is, there is a fixed initial concentration R_0 of the resource, and the population grows until the resource is exhausted. Figure 4A shows these dynamics for the hypothetical wild-type and mutant strains of Fig. 1. Although the mutant has the same maximum growth rate $g^{\max}$ as the wild-type, its lower value of K allows it to continue growing fast at lower concentrations of the resource, decelerating more abruptly at the end of growth (see inset of Fig. 4A for more dramatic examples). Then a fixed amount of biomass N_0 — sampled from the whole culture, so that the relative frequencies of the mutant and wild-type are preserved on average — is transferred to a new environment with the same initial concentration R_0 of the resource as before, and the cycle repeats (Fig. 4B, top panel). This dilution step represents a form of mortality for the population. We refer to this regime as *fixed-bottleneck batch dynamics*, since the bottleneck of biomass between transfers is held fixed. Boom-bust dynamics such as these are believed to be common in some natural environments [51, 52], with a fixed bottleneck size being plausible for populations that serially colonize new environments [53] or reset to a fixed density by culling [4] between cycles of growth.

The second regime is the same as the first, except instead of transferring a fixed amount of biomass to the next cycle, we transfer a fixed fraction $1/D$, where D is the dilution factor (Fig. 4B, bottom panel); we therefore refer to this regime as *fixed-dilution batch dynamics*. Note that the dilution factor D and the bottleneck biomass N_0 are related according to $D = R_0 Y / N_0 + 1$, where Y is the yield (biomass produced per unit resource; Supplementary Information Sec. S3). These dynamics are plausible for populations that experience a constant death rate between growth cycles or are regularly purged by the environment, as believed to occur in the human gut microbiome [54]. This case is also the most common protocol in laboratory evolution experiments owing to its simplicity [55]. While these two regimes of batch dynamics may appear to be only subtly different (comparing the two panels of Fig. 4B), we will show later that these two dilution protocols have different dependences on the resource concentration, which lead to different evolutionary outcomes.

Finally, we consider the regime of *chemostat dynamics*, where the population grows as a continuous culture with a constant supply of the resource and a constant dilution rate d (Supplementary Information Sec. S5). Chemostats are used as a device for experimental evolution [12, 22] and the same dynamics are often applied to describe natural populations in the ocean [13, 56].

Selection quantifies variation in growth traits between isolates at different resource concentrations

We previously observed wide variation in resource thresholds K (Fig. 2A) and maximum growth rates $g^{\max}$ (Fig. 3A) across isolates, but the significance of this variation is difficult to assess by itself. For example, glucose K for *E. coli* varied across four orders of magnitude, but how significant is this variation for evolution? Our model of selection under different population dynamics gives us precisely the metric to quantify this variation. We demonstrate this in Fig. 4C by calculating the two components of selection (Eq. (2)) between all pairs of *E. coli* isolates measured on glucose. We do this for batch dynamics starting at different initial concentrations R_0 of glucose. While selection on variation in $g^{\max}$ (s_{high}) always increases with higher R_0, selection on variation in K (s_{low}) depends non-monotonically on the concentration R_0, such that selection is maximized at some intermediate concentration (Fig. S16, Supplementary Information Sec. S10). Intuitively, this optimal concentration approximately equals the threshold K itself (Fig. S16C). On the other hand, if the resource concentration R_0 also increases the initial population size N_0 (i.e., transfer from a pre-growth cycle with fixed dilution factor), selection on variation in K depends monotonically on R_0 and is maximized at the lowest concentration (Fig. S17).

We calculate selection between *E. coli* isolates at 10 µM glucose, which is in the middle of the range of observed thresholds K, as well as at two higher concentrations corresponding to the conditions of the *E. coli* LTEE (139 µM) [57] and a common laboratory concentration (11000 µM ≈ 0.2% w/v). Figure 4C indeed shows that variation in the thresholds K is highly significant for evolution at concentrations around the threshold values, whereas at the highest concentration, selection on the variation in K is small compared to the selection in $g^{\max}$.

The resource threshold evolves downward over successive mutations

With our model of population dynamics, we can predict how the traits of the Monod growth rate dependence (Eq. (1)) will evolve over long times. For simplicity, we focus on the "strong-selection weak-mutation" (SSWM) regime of evolutionary dynamics, where each new mutation either fixes or goes extinct before the next mutation arises (Fig. S18; Supplementary Information Sec. S11) [58].

We first simulate a population growing under fixed-bottleneck batch dynamics, with an initial threshold K that is higher than the external resource concentration R_0; the population therefore decelerates gradually into starvation over each growth cycle (Fig. 5A, left inset). Mutations then regularly arise and alter the threshold K with a random effect size (Fig. S18; Supplementary Infor-

FIG. 4. **Selection on variation in resource thresholds over batch population dynamics.** (A) Simulated growth of wild-type (green) and mutant (orange) strains competing under batch dynamics, with the transient resource concentration (gray) on the right vertical axis (Supplementary Information Sec. S3). The strain pair is the same as in Fig. 1; the initial resource concentration is $R_0 = 25$, with strains at equal initial frequencies and equal yields. (B) The same strain competition from panel A continued over multiple growth cycles under fixed-bottleneck batch dynamics (top panel, $N_0 = 0.01$) and fixed-dilution batch dynamics (bottom panel, $D = 100$). (C) Each point represents the predicted selection coefficients $|s_{high}|$ and $|s_{low}|$ (Eq. (2); Supplementary Information Sec. S8) for pairs of *E. coli* isolates with measured growth traits on glucose (from Fig. 2D). The three colors represent different glucose concentrations. We assume the isolates in each pair start competing at equal initial frequencies, set the initial cell density to $N_0 = 4.6 \times 10^5$ cells/mL, and use a biomass yield of $Y = 3.3 \times 10^8$ cells/µmol glucose measured by a previous study [23].

mation Sec. S11). Each mutation stochastically fixes or goes extinct according to a fixation probability, which depends on the mutation's selection coefficient. Over time these beneficial mutations accumulate and the threshold K systematically decreases. By the end of the simulation, the threshold K is 1000 times smaller than the resource concentration R_0, leading to growth curves that grow much faster and abruptly decelerate into starvation (Fig. 5A, right inset).

Such an abrupt arrest is, for example, realized by *E. coli* in glucose-limited batch culture through a dynamic surge in gene expression late in the growth cycle [59], often involving the use of separate transporters with lower Michaelis-Menten constants [60]. The presence of these transporter systems has been raised as evidence for evolutionary adaptation of the species at micromolar glucose concentrations [8, 60, 61]. But our model shows that a feast-and-famine environment dominated by concentrations orders of magnitude higher would still allow *E. coli* to evolve the low resource threshold K observed in extant strains.

Adaptation in the resource threshold stalls when it reaches selection-drift balance

The threshold K does not evolve downward forever; in Fig. 5A adaptation slows down and the threshold levels off after a few tens of thousands of mutations, even though there is no change in the supply of beneficial

mutations. This occurs because selection on beneficial mutations is inhibited by random demographic fluctuations in the population, known as genetic drift [62]. The strength of genetic drift is measured by $1/N_e$, where N_e is the effective population size [63]; smaller populations experience greater fluctuations. In the simplest cases, N_e is proportional to the actual ("census") population size, but in more complex systems N_e may depend on other aspects of demography (such as spatial dynamics [64] or age structure [65]) as well as additional sources of noise in the population dynamics [66].

Beneficial mutations will therefore no longer fix with high probability if their selection equals genetic drift, a condition known as selection-drift balance [67–69]:

$$ s = \frac{1}{N_e}. \tag{3} $$

Since there are different concepts of effective population size for different phenomena [70], we note that N_e here specifically refers to the effective population size for the variance in mutant frequency change per unit time. Selection-drift balance occurs in our model under batch dynamics because the growth deceleration phase becomes shorter as K decreases over evolution (insets of Figs. 4A and 5A), which means there is weaker selection to reduce it further. Once the threshold K becomes sufficiently small, selection is no longer strong enough to overcome genetic drift (Supplementary Information Sec. S12; Fig. S19).

FIG. 5. **Evolution of the resource threshold.** (A) The resource threshold K evolving under fixed-bottleneck batch dynamics. Each gray line is one of 10 independent stochastic simulations using an effective population size $N_e = 1000$ and mutation effects κ drawn from a uniform distribution (Supplementary Information Sec. S11). The insets show the growth curve in a single batch cycle before adaptation (left inset) and at the final state (right inset). The green dashed line marks our prediction K_{evo} at selection-drift balance. (B) Evolved resource threshold K_{evo} as a function of the effective population size N_e. In the gray region, the effective population size is too small and all evolution is neutral. If N_e is sufficiently large (white region), the evolved resource threshold K_{evo} has selection-drift balance along the green line. Parameters are $|\kappa_{max}| = 0.001$, $g^{max} = 1$, $N_0 = 0.01$, and $Y = 1$ for both strains. (C) The evolved glucose threshold K_{evo} as a function of initial glucose concentration R_0 for two regimes of batch dynamics: fixed-bottleneck dynamics (blue line) or fixed-dilution dynamics (orange line). We use parameters based on the LTEE: $N_0 = 4.6 \times 10^5$ cells/mL (for fixed-bottleneck case), $D = 100$ (for fixed-dilution case), $N_e = V N_0$ where $V = 10$ mL, $g^{max} = 0.888$/h, and $Y = 3.3 \times 10^8$ cells/μmol [23]. We also set $\kappa_{max} = 6 \times 10^{-6}$ (Fig. S26). On the right axis is a histogram of glucose threshold K data for *E. coli* isolates (from Fig. 2B).

By combining Eqs. (2) and (3), we can calculate the value of the evolved threshold at which selection-drift balance occurs (Fig. S20). Under typical limits of the parameters, the evolved threshold is approximately (Supplementary Information Sec. S13)

$$K_{evo} \approx \frac{R_0}{N_e |\kappa_{max}| \log(N_e |\kappa_{max}| R_0 Y / N_0)}, \qquad (4)$$

where κ_{max} is the maximum effect size of a beneficial mutation reducing K. We calculate an example of K_{evo} in Fig. 5A (dashed green line), which corresponds well with the simulations. This result is robust to a wide range of effective population sizes and frequency-dependent effects (Fig. S21; Supplementary Information Sec. S11). We also observe an equivalent result for the adaptation of the specific affinity g^{max}/K (Fig. S22; Supplementary Information Sec. S14) instead of the threshold K alone.

One salient feature of Eq. (4) is that the evolved threshold K_{evo} scales inversely with the effective population size N_e, as shown in Fig. 5B. That is, larger populations or those with lower genetic drift can evolve proportionally lower thresholds K_{evo} that are orders of magnitude lower than the environmental resource concentration R_0. This suggests why we observe such low thresholds K for many organisms and resources (Fig. 2); this also explains why these thresholds are difficult to measure from time-series data, since low thresholds produce extremely abrupt deceleration at the end of growth (insets of Figs. 4A and 5A and Fig. S23; Supplementary Information Sec. S15). Hints of the influence of N_e are found in ammonia-oxidizing archaea and bacteria from marine environments, which tend to have lower resource

thresholds than lineages isolated from soil [18]. Assuming that these measurements more or less reflect cell growth rate, our scaling relationship Eq. (4) suggests that this ordering can arise from the smaller effective population size N_e for spatially-structured environments like soil.

The other important feature of Eq. 4 is the dependence of the evolved threshold K_{evo} on the resource concentration R_0. For fixed effective population size N_e, there is an optimal value of R_0 that minimizes the evolved threshold K_{evo} (left insets of Fig. S20A,B), just as we observed for selection on individual mutations (Fig. S16). We note that for sufficiently low values of the effective population size N_e, genetic drift is stronger than selection on any mutation κ (Fig. S19A), and so the threshold K evolves neutrally (gray region in Fig. 5B).

In contrast to batch dynamics, selection under chemostat dynamics does not depend on the threshold K itself (Supplementary Information Sec. S9). Intuitively, this is because reductions in K cause the environmental resource concentration to decrease proportionally (Supplementary Information Sec. S5), such that the growth rate remains constant. Not only does this keep a constant strength of selection on new mutations, but the effective population size will actually increase as K evolves lower, making beneficial mutations even easier to fix. Therefore selection-drift balance never occurs for K under chemostat dynamics; the threshold K will continue to evolve downward until adaptation is limited by the supply of mutations or other factors (Discussion). Note that selection-drift balance also does not occur for mutations to the maximum growth rate g^{max} under either batch or chemostat dynamics, since selection does not depend on the magnitude of growth rate (Supplementary

Information Secs. S8 and S9).

Population dynamics can decouple the evolved threshold from the resource concentration

In general the effective population size N_e that controls genetic drift may be shaped by a variety of demographic factors besides the census population size [63]. However, in well-mixed batch cultures, N_e is primarily determined by the number of cells at the bottleneck of each transfer [67, 71]; we assume that other sources of stochasticity (such as individual cell division events) are much smaller than the sampling noise of these transfers. Therefore the effective population size N_e is proportional to the bottleneck biomass N_0 (assuming constant biomass per cell).

Under fixed-bottleneck batch dynamics, the effective population size N_e is thus an independent parameter of the population, so that the strength of genetic drift does not depend on the resource concentration (Fig. S24A). In this case, the evolved threshold K_{evo} is in approximately linear proportion to the resource concentration R_0 (Eq. (4); Figs. 5C and S25A), making the evolved threshold a biomarker of the resource's environmental concentration. This is consistent with our original speculation about the systematic differences in glucose K between *E. coli* and *S. cerevisiae*, owing to the different glucose availability in their different environments.

However, for fixed-dilution batch dynamics, the bottleneck biomass N_0, and therefore the effective population size, are coupled to the resource concentration R_0 because the dilution factor D is fixed: $N_e \propto N_0 = R_0 Y/(D-1)$ (Supplementary Information Sec. S3). This is because increasing the resource concentration increases the biomass at the end of each growth cycle, but then the bottleneck biomass must also increase proportionally to hold the dilution factor fixed. The scaling of N_e with R_0, though, cancels out the scaling of K_{evo} with R_0 in Eq. (4), leading to an evolved threshold K_{evo} that is approximately independent of the environmental concentration R_0 (Figs. 5C and S25B). Conceptually, this occurs because increasing the resource R_0 decreases selection on the mutation but also decreases genetic drift, since the effective population size N_e increases (Fig. S24B). Thus, the population dynamics decouple the evolved threshold from the environmental concentration.

This has major consequences for interpreting empirical variation. We predict the evolved threshold K_{evo} for *E. coli* on glucose as a function of glucose concentration R_0 in Fig. 5C, using parameters estimated from the LTEE (Fig. S26). On the same plot, we show a histogram of all measured glucose thresholds for *E. coli* (from Fig. 2B) on the right vertical axis. We see that, under fixed-bottleneck batch dynamics, we would expect *E. coli* to have evolved in glucose concentrations above 100 μM to account for the observed glucose thresholds. However, under fixed-dilution batch dynamics, the evolved glucose threshold depends so weakly on the environmental concentration that almost any concentration of glucose is possible to explain the data.

DISCUSSION

Modeling insights to interpret threshold data

Since it is often difficult to measure resource concentrations and population dynamics in natural environments, can we use the evolved threshold K as a biomarker to infer them? This logic is often implicit in environmental studies, which attempt to draw conclusions about the environmental conditions of an isolate based on its abilities to grow at different resource concentrations [16–19]. However, our model shows that it is not as simple as assuming the threshold K for a resource is proportional to its concentration in the environment, since that proportionality is altered by the population dynamics, at least through the effective population size N_e (Eq. (4)). In particular, this proportionality is confounded in the case of fixed-dilution batch dynamics, where the evolved threshold K is largely independent of the resource concentration R_0 (Fig. 5C).

Under fixed-bottleneck batch dynamics, though, the linear scaling of K with R_0 does approximately hold. In this case, one can compare two populations with unknown, but identical effective population sizes N_e and mutation effects κ; for example, two isogenic populations located at different points along a resource gradient. In this case, one can calculate the ratio of evolved resource thresholds K_{evo} for the two populations to estimate the ratio of resource concentrations. But in many scenarios, one might not even know the type of bottlenecks the population is experiencing. To classify the population dynamics as fixed-bottleneck or fixed-dilution, one could correlate a set of evolved thresholds K_{evo} with their different resource concentrations R_0; a strong linear correlation would support fixed-bottleneck batch dynamics, while little to no correlation would indicate fixed-dilution batch or chemostat dynamics.

Role of the mutation supply in shaping evolved resource thresholds

We have focused on the role of selection-drift balance as a null model for the evolved variation in resource thresholds, since the competition between selection and genetic drift is a universal feature of all evolving populations. In doing so we have assumed the supply of mutations on K is constant, but real populations will at some point run out of beneficial mutations on the threshold K, potentially reaching this mutation-selection balance before selection-drift balance [68]. Many mutations will also be pleiotropic, affecting both the resource threshold K and the maximum growth rate g^{max} (as well as possibly other traits) simultaneously. The correlation be-

tween pleiotropic effects on both traits is important: if pleiotropy is synergistic, so that mutations that decrease K also tend to increase g^{max}, then the population might evolve lower K than otherwise expected since its selection is enhanced by additional selection on g^{max}. On other other hand, if there is a tradeoff between K and g^{max}, the population might evolve higher K if its selection is outweighed by selection for higher g^{max}. Indeed, this is what appears to happen in the LTEE, where K for glucose actually increased over the first 2000 generations, but that was offset by a stronger improvement in the maximum growth rate g^{max} [23].

Such a tradeoff between K and g^{max} is interesting both for its consequences on the stochiometric composition of community biomass [48, 72] as well as from an evolutionary point of view, since the population can then diversify into stably-coexisting lineages. While there is significant theoretical work on this hypothesis [25–28], it has limited empirical evidence. Some of these previous studies claiming tradeoffs found them only in parameters for the Michaelis-Menten model of resource uptake [27, 28, 30, 48, 73], which we and others have shown are not equivalent to parameters of the Monod model of growth (Fig. S11) [36, 38]. In the larger set of data we have collected in this work (Fig. 2F), we find no compelling evidence of a correlation; _E. coli_ shows a weak but insignificant tradeoff, while _S. cerevisiae_ shows a slight synergy [74].

Interpretation of this tradeoff (or lack thereof) is also complicated by the sample of strains and environmental conditions being considered. For the tradeoff to affect the evolved threshold as we have discussed, the tradeoff must exist across the spectrum of spontaneous mutations available to an organism (i.e., there is an underlying physiological constraint). This has also been the underlying assumption of previous models on this topic [25–28]. Testing this would require a distribution of K and g^{max} measurements over a large mutant library in a single environment, which has not been done to our knowledge. However, even in the absence of an underlying correlation in mutation effects, such a tradeoff could still emerge across clones within a rapidly evolving population, at least transiently [75, 76]. Further systematic measurements of these traits within and between populations will be necessary to resolve the issue of a tradeoff in the future.

Other factors shaping evolved resource thresholds

Besides mutation supply, there are other phenomena that may lead to different evolved outcomes for the threshold K. One important assumption in our model is that we only consider a single resource, whereas real populations are dependent on several resources [77], including those from biotic sources such as cross-feeding and predation. Some of these resources may be rarely or never limiting, and therefore their thresholds K will evolve

only as byproducts of selection on mutations for other traits. In this sense many observed resource thresholds may actually be spandrels, an evolutionary term (defined in analogy with the architectural structure) for traits that evolve for reasons other than direct selection [78]. Selection for other traits may occur simply because competition in natural environments is likely more complex and could include lag phases [50] and other strategies for low-resource survival [5, 79–81]. On the other hand, multiple resources could also be simultaneously colimiting [31, 32]. While we have shown how colimitation under measurement conditions affects estimates of g^{max} and K (Supplementary Information Sec. S1), the effect of colimitation, as well as more complex sources of nutrients such as cross-feeding and predation, on the evolution of these traits remains an important problem for future work.

We can predict the consequences of relaxing other assumptions in our model as well. For example, simultaneous competition of multiple mutations (clonal interference) generally reduces the efficacy of selection [82, 83], which would make it more likely to evolve higher thresholds than what we predict from SSWM dynamics. Another assumption in our model is that the population under batch dynamics always grows until complete exhaustion of the resources during each cycle, but earlier transfers could reduce the amount of growth occurring during deceleration, which would reduce selection on the threshold K. However, the population may adapt its maximum growth rate to simply saturate earlier and restore selection on its deceleration phase. Finally, populations may also have higher than expected K thresholds if they simply have not had enough time to reach selection-drift balance, which takes a timescale of order N_{e} generations (Fig. S21) [84].

Population dynamics are essential for understanding microbial ecology

Broadly speaking, our results provide a valuable example of how ecological traits are influenced by factors other than abiotic environmental features. In particular, we have shown how population dynamics can confound our naive expectations for the evolutionary fate of such traits. While here we have focused on the role of genetic drift, other potentially important factors include mutation supply, pleiotropy, recombination, and spatial structure. Altogether our results mean that the threshold K may not be a reliable biomarker of environmental resource concentrations. This does not mean that K evolves independently of the environment, however. Rather, it is linked to additional environmental processes like the bottleneck between growth cycles. To understand the systematic differences between species, we need to know not only the resource concentrations they have evolved in, but also which type of population dynamics best reflects the time scales of growth, death, and resource supply in natural environments.

Materials and Methods

Literature survey of measured growth rate dependence on resources

We collected 247 measurements of Monod model parameters (K and $g^{\max}$; Eq. (1)) through a targeted literature search that included prior surveys and reviews [40, 42], the phytoplankton trait database (130 data points) by Edwards et al. [43], as well as original research papers. In all but two cases, we traced data from surveys and reviews back to their original papers, which we report in Dataset S1 (sheet 1). We included only experiments that directly measured population growth rates, rather than nutrient uptake rates or respiration. We excluded measurements where the actual limiting resource was unclear, such as measurements in rich medium with added glucose. Where possible we checked the raw data of growth rate over resource concentrations to determine if the focal resource concentration was measured up to saturation and had sufficient sampling of concentrations around K. For a subset of measurements of *E. coli* on glucose, we also checked for the concentration of a nitrogen source to determine the relative impact of colimitation (Supplementary Data Table S1, sheet 2; Supplementary Information Sec. S1). If the original K value was reported as weight per volume, we converted these into units of micromolar (μM) using the calculated molecular weight of the compound's chemical formula. We preserved significant digits from the original studies. See Dataset S1 for more details.

Models of population dynamics

We mathematically model population dynamics using systems of ordinary differential equations for the wild-type and mutant biomasses as well as the extracellular resource concentration (Supplementary Information Secs. S3 and S5). We numerically integrate these equations using standard algorithms in Scipy [85] (Supplementary Information Sec. S4).

ACKNOWLEDGMENTS

JWF and MM were supported by an Ambizione grant from the Swiss National Science Foundation (PZ00P3_180147). NAH was supported by the Principles of Microbial Ecosystems collaboration of the Simons Foundation (grant ID 542379) and an ETH Zurich Career Seed Grant (SEED-26 21-2). We thank Ayush Pathak for a critical reading of the manuscript.

[1] M. Schaechter, J. L. Ingraham, and F. C. Neidhardt. *Microbe*. ASM Press, Washington, DC, 2006.

[2] T. E. Shehata and A. G. Marr. Effect of nutrient concentration on the growth of Escherichia Coli. *Journal of Bacteriology*, 107:210–216, 1971.

[3] Nicolai S. Panikov. *Microbial Growth Kinetics*. London: Chapman & Hall, 1995.

[4] David L. Kirchman. Growth rates of microbes in the oceans. *Annual Review of Marine Science*, 8:285–309, 2016.

[5] Jeanne S. Poindexer. *Oligotropy*, pages 63–89. Springer US, 1981.

[6] Noah Fierer, Mark A. Bradford, and Robert B. Jackson. Toward an ecological classification of soil bacteria. *Ecology*, 88:1354–1364, 2007.

[7] J. Monod. The growth of bacterial cultures. *Annu Rev Microbiol*, 3:371–394, 1949.

[8] Arthur L. Koch. The adaptive responses of escherichia coli to a feast and famine existence. *Advances in Microbial Physiology*, 6:147–217, 1971.

[9] Don K. Button. Kinetics of nutrient-limited transport and microbial growth. *Microbiological Reviews*, 49:270–279, 1985.

[10] D. Tilman. *Resource competition and community structure*. Princeton University Press, Princeton, NJ, 1982.

[11] T. E. Miller, J. H. Burns, P. Munguia, E. L. Walters, J. M. Kneitel, P. M. Richards, N. Mouquet, and H. L. Buckley. A critical review of twenty years' use of the resource-ratio theory. *Am Nat*, 165:439–448, 2005.

[12] Joey R. Bernhardt, Pavel Kratina, Aaron Louis Pereira, Manu Tamminen, Mridul K. Thomas, and Anita Narwani. The evolution of competitive ability for essential resources. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375(1798):20190247, 2020.

[13] James P. Grover. *Resource Competition*. Springer US, 1997.

[14] Emilio Marañón, Pedro Cermeño, María Huete-Ortega, Daffne C. López-Sandoval, Beatriz Mouriño-Carballido, and Tamara Rodríguez-Ramos. Resource Supply Overrides Temperature as a Controlling Factor of Marine Phytoplankton Growth. *PLOS ONE*, 9(6):e99312, 2014.

[15] Erik Askov Mousing, Katherine Richardson, and Marianne Ellegaard. Global patterns in phytoplankton biomass and community size structure in relation to macronutrients in the open ocean. *Limnology and Oceanography*, 63:1298–1312, 2018.

[16] Willm Martens-Habbena, Paul M. Berube, Hidetoshi Urakawa, José R. de la Torre, and David A. Stahl. Ammonia oxidation kinetics determine niche separation of nitrifying archaea and bacteria. *Nature*, 461:976–979, 2009.

[17] James I. Prosser and Nicol W. Graeme. Archaeal and bacterial ammonia-oxidisers in soil: The quest for niche specialisation and differentiation. *Trends in Microbiology*, 20:523–31, 2021.

[18] Dimitri K. Kits, Christopher J. Sedlacek, and Elena V. Lebedeva et al. Kinetic analysis of a complete nitrifier reveals an oligotrophic lifestyle. *Nature*, 549:269–272, 2017.

[19] Man-Young Jung, Christopher J. Sedlacek, K. Dimitri Kits, Anna J. Mueller, Sung-Keun Rhee, Linda Hink, Graeme W. Nicol, and et al. Ammonia-oxidizing archaea possess a wide range of cellular ammonia affinities. *The ISME Journal*, 16:272–283, 2022.

[20] Daniel Dykhuizen and Daniel Hartl. Evolution of competitive ability in escherichia coli. *Evolution*, 35:581, 1981.

[21] Karin Kovárŏvá. *Growth Kinetics of Escherichia Coli: Effect of Temperature, Mixed Substrate Utilization and Adaptation to Carbon-Limited Growth*. PhD thesis, ETH Zurich, Zurich, Switzerland, 1996.

[22] Julian Adams, Charlotte Paquin, Paul W. Oeller, and Lester W. Lee. Physiological characterization of adaptive clones in evolving populations of the yeast, saccharomyces cerevisiae. *Genetics*, 110:173–185, 1985.

[23] Farida Vasi, Michael Travisano, and Richard E. Lenski. Long-Term Experimental Evolution in Escherichia coli. II. Changes in Life-History Traits During Adaptation to a Seasonal Environment. *The American Naturalist*, 144(3):432–456, 1994.

[24] Samuel Frederick Mock Hart, Chi-Chun Chen, and Wenying Shou. Pleiotropic mutations can rapidly evolve to directly benefit self and cooperative partner despite unfavorable conditions. *eLife*, 10:e57838, 2021.

[25] Frank M. Stewart and Bruce R. Levin. Partitioning of Resources and the Outcome of Interspecific Competition: A Model and Some General Considerations. *The American Naturalist*, 107(954):171–198, 1973.

[26] Antony M Dean. Protecting Haploid Polymorphisms in Temporally Variable Environments. *Genetics*, 169(2):1147–1156, 2005.

[27] Robert E. Beardmore, Ivana Gudelj, David A. Lipson, and Laurence D. Hurst. Metabolic trade-offs and the maintenance of the fittest and the flattest. *Nature*, 472(7343):342–346, 2011.

[28] Meike T. Wortel. Evolutionary coexistence in a fluctuating environment by specialization on resource level. *BioRxiv*, preprint:2021.05.18.444718, 2021.

[29] Kai W. Wirtz. A generic model for changes in microbial kinetic coefficients. *Journal of Biotechnology*, 97(2):147–162, 2002.

[30] Elena Litchman, Christopher A. Klausmeier, Oscar M. Schofield, and Paul G. Falkowski. The role of functional traits and trade-offs in structuring phytoplankton communities: scaling from cellular to ecosystem level. *Ecology Letters*, 10(12):1170–1181, 2007.

[31] Mak A. Saito, Tyler J. Goepfert, and Jason T. Ritt. Some thoughts on the concept of colimitation: Three definitions and the importance of bioavailability. *Limnology and Oceanography*, 53(1):276–290, 2008.

[32] S. W. Harpole, J. T. Ngai, E. E. Cleland, E. W. Seabloom, E. T. Borer, M. E. S. Bracken, J. J. Elser, D. S. Gruner, H. Hillebrand, J. B. Shurin, and J. E. Smith. Nutrient co-limitation of primary producer communities. *Ecol Lett*, 14:852–862, 2011.

[33] Don K. Button. Affinity of organisms for substrate. *Limnology and Oceanography*, 31:435–456, 1986.

[34] F P Healey. Slope of the Monod equation as an indicator of advantage in nutrient competition. *Microbial ecology*, 5(4):281–286, 1980.

[35] R. V. O'Neill, D. L. DeAngelis, J. J. Pastor, B. J. Jackson, and W. M. Post. Multiple nutrient limitations in ecological models. *Ecol Modelling*, 46:147–163, 1989.

[36] J. L. Snoep, M. Mrwebi, J. M. Schuurmans, J. M. Rohwer, and M. J.YR 2009 Teixeira de Mattos. Control of specific growth rate in Saccharomyces cerevisiae. *Microbiology*, 155(5):1699–1707, 2009.

[37] S. Sharma and R. Steuer. Modelling microbial communities using biochemical resource allocation analysis. *J R Soc Inter*, 16:20190474, 2019.

[38] David Tilman and Susan Soltau Kilham. Phosphate and Silicate Growth and Uptake Kinetics of the Diatoms Asterionella Formosa and Cyclola Meneghiniana in Batch and Semicontinuous Culture1. *Journal of Phycology*, 12(4):375–383, 1976.

[39] François M. M. Morel. Kinetics of Nutrient Uptake and Growth in Phytoplankton. *Journal of Phycology*, 23(1):137–150, 1987.

[40] J.D. Owens and J.D. Legan. Determination of the Monod substrate saturation constant for microbial growth. *FEMS Microbiology Letters*, 46(4):419–432, 1987.

[41] H W Jannasch. Growth characteristics of heterotrophic bacteria in seawater. *Journal of Bacteriology*, 95(2):722–723, 1968.

[42] Karin Kovárová-Kovar and Thomas Egli. Growth kinetics of suspended microbial cells: From single-substrate-controlled growth to mixed-substrate kinetics. *Microbiology and Molecular Biology Reviews*, 62:646–666, 1998.

[43] Kyle F. Edwards, Christopher A. Klausmeier, and Elena Litchman. Nutrient utilization traits of phytoplankton. *Ecology*, 96(8):2311–2311, 2015.

[44] Ron Milo and Rob Phillips. *Cell Biology by the Numbers*. Garland Science, Taylor Francis Group, 2016.

[45] Thomas Ferenci. 'growth of bacterial cultures' 50 years on: Towards an uncertainty principle instead of constants in bacterial growth kinetics. *Research in Microbiology*, 150:431–38, 1999.

[46] Huai-Feng Liu, Ben-Hong Wu, Pei-Ge Fan, Shao-Hua Li, , and Lian-Sheng Li. Sugar and acid concentrations in 98 grape cultivars analyzed by principal component analysis. *Journal of the Science of Food and Agriculture*, 86:1526—36, 2006.

[47] Zerihun T.Dame, Farid Aziat, Rupasri Mandal, Ram Krishnamurthy, and Souhaila Bouatra et al. The human saliva metabolome. *Metabolomics*, 11:1864–83, 2015.

[48] Elena Litchman, Kyle F. Edwards, and Christopher A. Klausmeier. Microbial resource utilization traits and trade-offs: implications for community structure, functioning, and biogeochemical impacts at present and in the future. *Frontiers in Microbiology*, 6, 2015.

[49] Chevin Luis-Miguel. On measuring selection in experimental evolution. *Biology Letters*, 7(2):210–213, 2011.

[50] Michael Manhart, Bharat V. Adkar, and Eugene I. Shakhnovich. Trade-offs between microbial growth phases lead to frequency-dependent and non-transitive selection. *Proceedings of the Royal Society B: Biological Sciences*, 285(1872), 2018.

[51] Michael J. Behrenfeld, Yongxiang Hu, Robert T. O'Malley, Emmanuel S. Boss, Chris A. Hostetler, David A. Siegel, Jorge L. Sarmiento, Jennifer Schulien, Johnathan W. Hair, Xiaomei Lu, Sharon Rodier, and Amy Jo Scarino. Annual boom–bust cycles of polar phytoplankton biomass revealed by space-based lidar. *Nature Geoscience*, 10(2):118–122, 2017.

[52] David M. Needham, Cheryl-Emiliane T. Chow, Jacob A. Cram, Rohan Sachdeva, Alma Parada, and Jed A. Fuhrman. Short-term observations of marine bacterial

and viral communities: patterns, connections and resilience. *The ISME Journal*, 7(7):1274–1285, 2013.

[53] Tim N. Enke, Manoshi S. Datta, Julia Schwartzman, nathan Cermak, Désirée Schmitz, Julien Barrere, Alberto Pascual-García, and Otto X. Cordero. Modular assembly of polysaccharide-degrading marine microbial communities. *Current Biology*, 29, 2019.

[54] Jonas Cremer, Markus Arnoldini, and Terence Hwa. Effect of water flow and chemical environment on microbiota growth and composition in the human colon. *Proceedings of the National Academy of Sciences*, 114:6438—-43, 2017.

[55] J. E. Barrick and R. E. Lenski. Genome dynamics during experimental evolution. *Nat Rev Genet*, 14:827–839, 2013.

[56] S. Dutkiewicz, M. J. Follows, and J. G. Bragg. Modeling the coupling of ocean ecology and biogeochemistry. *Global Biogeochemical Cycles*, 23(4), 2009.

[57] Richard E. Lenski, Michael R. Rose, Suzanne C. Simpson, and Scott C. Tadler. Long-Term Experimental Evolution in Escherichia coli. I. Adaptation and Divergence During 2,000 Generations. *The American Naturalist*, 138(6):1315–1341, 1991.

[58] J. H. Gillespie. Molecular evolution over the mutational landscape. *Evolution*, 38:1116–1129, 1984.

[59] Anat Bren, Yuval Hart, Erez Dekel, Daniel Koster, and Uri Alon. The last generation of bacterial growth in limiting nutrient. *BMC Systems Biology*, 7(1):27, 2013.

[60] Thomas Ferenci. Adaptation to life at micromolar nutrient levels: The regulation of Escherichia Coli glucose transport by endoinduction and camp. *FEMS Microbiology Reviews*, 18:301–317, 1996.

[61] J. W. Lengeler. Carbohydrate transport in bacteria under environmental conditions, a black box? *Antonie van Leeuwenhoek*, 63, 1993.

[62] J. F. Crow and M. Kimura. *An Introduction to Population Genetics Theory*. Harper and Row, New York, 1970.

[63] B. Charlesworth. Effective population size and patterns of molecular evolution and variation. *Nat Rev Genet*, 10:195–205, 2009.

[64] O. Hallatschek, P. Hersen, S. Ramanathan, and D. R. Nelson. Genetic drift at expanding frontiers promotes gene segregation. *Proc Natl Acad Sci USA*, 104:19926–19930, 2007.

[65] M. Shpak. Selection against demographic stochasticity in age-structured populations. *Genetics*, 177:2181–2194, 2007.

[66] J. H. Gillespie. Natural selection for within-generation variance in offspring number II. Discrete haploid models. *Genetics*, 81:403–413, 1975.

[67] M. Kimura. *The Neutral Theory of Molecular Evolution*. Cambridge University Press, Cambridge, 1983.

[68] A. W. R. Serohijos and E. I. Shakhnovich. Merging molecular mechanism and evolution: Theory and computation at the interface of biophysics and evolutionary population genetics. *Curr Opin Struct Biol*, 26:84–91, 2014.

[69] M. Lynch and K. Hagner. Evolutionary meandering of intermolecular interactions along the drift barrier. *Proc Natl Acad Sci USA*, 112:E30–38, 2015.

[70] W. J. Ewens. *Mathematical Population Genetics*. Springer-Verlag, New York, 2004.

[71] R. E. Lenski, M. R. Rose, S. C. Simpson, and S. C. Tadler. Long-term experimental evolution in Escherichia coli. I. Adaptation and divergence during 2,000 generations. *Am Nat*, 138:1315–1341, 1991.

[72] Kevin J. Flynn, John A. Raven, T. Alwyn V. Rees, Zoe Finkel, Antonietta Quigg, and John Beardall. Is the Growth Rate Hypothesis Applicable to Microalgae?1. *Journal of Phycology*, 46(1):1–12, 2010.

[73] Karin Elbing, Christer Larsson, Roslyn M. Bill, Eva Albers, Jacky L. Snoep, Eckhard Boles, Stefan Hohmann, and Lena Gustafsson. Role of Hexose Transport in Control of Glycolytic Flux in Saccharomyces cerevisiae. *Applied and Environmental Microbiology*, 70(9):5323–5330, 2004.

[74] Naomi Ziv, Mark L. Siegal, and David Gresham. Genetic and Nongenetic Determinants of Cell Growth Variation Assessed by High-Throughput Microscopy. *Molecular Biology and Evolution*, 30(12):2568–2578, 2013.

[75] K. Gomez, J. Bertram, and J. Masel. Directional selection rather than functional constraints can shape the G matrix in rapidly adapting asexuals. *Genetics*, 211:715–729, 2019.

[76] Jie Lin, Michael Manhart, and Ariel Amir. Evolution of Microbial Growth Traits Under Serial Dilution. *Genetics*, 215(3), 2020.

[77] M. Kaspari and J. S. Powers. Biogeochemistry and geographical ecology: embracing all twenty-five elements required to build organisms. *Am Nat*, 188:S62–S73, 2016.

[78] S. J. Gould and R. C. Lewontin. The spandrels of San Marco and the Panglossian paradigm: A critique of the adaptationist programme. *Proc R Soc B*, 205:581–598, 1979.

[79] Arthur L. Koch. Oligotrophs versus copiotrophs. *BioEssays*, 23(7):657–661, 2001.

[80] R. Cavicchioli, M. Ostrowski, F. Fegatella, A. Goodchild, and N. Guixa-Boixereu. Life under Nutrient Limitation in Oligotrophic Marine Environments: An Eco/Physiological Perspective of Sphingopyxix alaskensis (Formerly Sphingomonas alaskensis). *Microbial Ecology*, 45(3):203–217, 2003.

[81] Jana Grote, J. Cameron Thrash, Megan J. Huggett, Zachary C. Landry, Paul Carini, Stephen J. Giovannoni, and Michael S. Rappé. Streamlining and Core Genome Conservation among Highly Divergent Members of the SAR11 Clade. *mBio*, 3(5):e00252–12, 2012.

[82] P. J. Gerrish and R. E. Lenski. The fate of competing beneficial mutations in an asexual population. *Genetica*, 102/103:127–144, 1998.

[83] S. Schiffels, G. J. Szöllösi, V. Mustonen, and M. Lässig. Emergent neutrality in adaptive asexual evolution. *Genetics*, 189:1361–1375, 2011.

[84] M. Kimura and T. Ohta. The average number of generations until fixation of a mutant gene in a finite population. *Genetics*, 61:763–771, 1969.

[85] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Com-

puting in Python. *Nature Methods*, 17:261–272, 2020.

[86] Jacques Monod and A Audureau. Mutation et adaptation enzymatique chez escherichia coli-mutabile. In *Annales de l'Institut Pasteur*, volume 72, pages 868–878. Masson Editeur 120 Blvd Saint-Germain, 75280 Paris 06, France, 1946.

FIG. S1. **Historical trends of measured resource thresholds.** The number of measured resource thresholds K published in peer-reviewed journals aggregated by year, based on our literature survey (Dataset S1). Colors indicate the number of measurements for individual resources.

FIG. S2. **Comparison of technical covariates for the resource threshold.** (A) Linear regressions of technical covariates against thresholds $\log_{10} K$ from the complete dataset (Fig. 2A), with degrees of freedom (df), number of data points (n), and p-values indicated. Each bar represents a separate regression fit, where R^2 measures the variation explained by a single variable as predictor for the resource threshold. (C) Linear regressions of technical covariates against glucose thresholds $\log_{10} K$ for all *E. coli* measurements (shown in Fig. 2B).

FIG. S3.   **Comparison of resource threshold measurements with and without acclimation.** (A) Empirical resource thresholds for glucose, grouped by taxon (only those with at least two measurements). The data shown here are identical to Fig. 2B, but colors indicate which measurements included a phase of acclimation (red). We infer acclimation from the type of experiments used to measure the resource threshold: For *batch solid culture*, growth rate is inferred from the area increase of single cell colonies on agar plates. For *batch experiments*, the growth rate is observed from exponential phase of a liquid culture with varying initial resource concentration. For *chemostat experiments*, the residual resource concentration is observed in steady state with varying growth rate by tuning the rate of liquid outflow. For *serial transfer experiments*, the growth is only measured in exponential phase after multiple transfers. We consider measurements to be acclimated if they derive from chemostat or serial transfer experiments. (B) Covariation between maximum growth rate g^{max} and glucose threshold K for isolates of *E. coli*. The data shown here are identical to *E. coli* datapoints in Fig. 3F. We calculate the Spearman rank correlation ρ and p-value across all isolates with acclimation (red dots). (C) Pairwise comparison of resource threshold measurements before and after acclimation. We identify a subset of publications in our database (see legend) which have explicitly tested the effect of acclimation. Each publication has two measurements for the organism's resource threshold K which we report together with full citations in our database (Supplementary Data Table 1). A black diagonal line indicates exact match between measurements with and without acclimation, with diagonal lines in dashes ($y = 10x$) and dots ($y = 0.1x$) as visual guides for the eye.

FIG. S4.   **Survey of resource thresholds and specific affinities for nitrate and ammonium in our survey.**
(A) Subset of K measurements from Fig. 2A for nitrate, grouped by taxon (only those with at least two measurements).
Symbols are the same as in Fig. 2A: Color indicates whether the organism is a prokaryote (green) or eukaryote (orange), and
shape indicates whether the organism can grow as an autotroph (square) or only as a heterotroph (circle). We use the taxonomic
identity given in the original publications, where an ending in *sp.* means the isolate is a representative of the genus but was
not identified at the species level. (B) Subset of K measurements from Fig. 2A for ammonium, grouped by taxon (with at least
two measurements). (C) Subset of $g^{\max}/K$ measurements from Fig. S8A for nitrate, grouped by taxon. (D) Subset of $g^{\max}/K$
measurements from Fig. S8A for ammonium, grouped by taxon.

FIG. S5. **Variation in glucose thresholds by experiment type and substrain label.** Subset of data from Fig. 2B for *E. coli* on glucose, with different strains separated. The strains ML 30 and ML308 were derived from a natural isolate in human feces by Jacques Monod in 1946 and differ in their genes for lactose utilization [86]: the lacI repressor is non-functional in ML308. We only show substrains with two or more measurements from the data. The three panels show the same data but are colored according to (A) publication, (B) temperature, and (C) experimental method (batch or chemostat).

FIG. S6. **Survey of maximum growth rates in our survey grouped by resource and taxon.** (A) Subset of g^{max} measurements from Fig. 3A for glucose, grouped by taxon (only those with at least two measurements). Symbols are the same as in Fig. 3A: Color indicates whether the organism is a prokaryote (green) or eukaryote (orange), and shape indicates whether the organism can grow as an autotroph (square) or only as a heterotroph (circle). We use the taxonomic identity given in the original publications, where an ending in *sp.* means the isolate is a representative of the genus but was not identified at the species level. (B) Subset of g^{max} measurements from Fig. 3A for phosphate, grouped by taxon (with at least three measurements). (C) Subset of g^{max} measurements from Fig. 3A for silicate, grouped by taxon (with at least two measurements). (D) Subset of g^{max} measurements from Fig. 3A for nitrate, grouped by taxon. (E) Subset of g^{max} measurements from Fig. 3A for ammonium, grouped by taxon.

FIG. S7.   **Comparison of technical covariates for maximum growth rate and specific affinity.** (A) Linear regression of technical covariates against maximum growth rate on glucose $g^{\max}$ for all *E. coli* measurements, with degrees of freedom (df), number of data points (n), and p-values indicated. We follow the same analysis as in Fig. S2B, but using $g^{\max}$ as the target variable for regression (no log transform). (B) Linear regression of technical covariates against the specific affinity $\log_{10}(g^{\max}/K)$ on glucose for all *E. coli*. The set of underlying isolates is identical to panel A. Here we use the log-transformed maximum growth rate $\log_{10} g^{\max}$ as a predictor, to compare the contributions of variation in $\log_{10} g^{\max}$ and variation in $\log_{10} K$ to the total variation in $\log_{10}(g^{\max}/K)$. The fraction of variation R^2 explained by $\log_{10} g^{\max}$ is too small to be visible.

FIG. S8.   **Survey of specific affinities computed from our survey.** (A) Variation in specific affinity $a = g^{\max}/K$ for the microbial isolates in our survey. For each isolate, we compute the trait value from the maximum growth rate $g^{\max}$ (Fig. 3A) and resource threshold K (Fig. 2A). Each point represents a different measurement; color indicates whether the organism is a prokaryote (green) or eukaryote (orange), and shape indicates whether the organism can grow as an autotroph (square) or only as a heterotroph (circle). The set of isolates shown here is fewer than in the total dataset, since some publications only reported the resource threshold K and not the maximum growth rate $g^{\max}$. (B) Subset of K measurements from panel A for glucose, grouped by taxon (only those with at least two measurements). We use the taxonomic identity given in the original publications, where an ending in *sp.* means the isolate is a representative of the genus but was not identified at the species level. Symbols are the same as in panel A. (C) Subset of K measurements from panel A for phosphate, grouped by taxon (with at least three measurements). (D) Subset for silicate, grouped by taxon (with at least two measurements). Compare also additional plots with $g^{\max}/K$ measurements for nitrate (Fig. S4C) and ammonium (Fig. S4D).

FIG. S9. **Covariation of Monod growth traits for autotroph and heterotroph isolates.** (A) Covariation of resource threshold K with maximum growth rate $g^{\max}$ for all heterotrophs (subset of points from Fig. 3B). We compute the Spearman rank correlation ρ and p-value across all resources. Colors indicate the limiting resource, with the number of measurements n in brackets. (B) Covariation of specific affinity $g^{\max}/K$ with $g^{\max}$ for all heterotrophs (subset from Fig. S10A). (C) Covariation of resource threshold with maximum growth rate for all autotrophs (subset from Fig. 3B). (D) Covariation of specific affinity with maximum growth rate for all autotrophs (subset from Fig. S10A). (E) Covariation of resource threshold with maximum growth rate for ammonia only (subset from panel C). See Fig. 3C–E for phosphate, nitrate, and silicate. (F) Covariation of specific affinity with maximum growth rate for ammonia only (subset from panel D). See Fig. S10B–D for phosphate, nitrate, and silicate. (G) Covariation of resource threshold with maximum growth rate for glucose only (subset from panel A). See Fig. 3F for covariation within species. (H) Covariation of specific affinity with maximum growth rate for glucose only (subset from panel B). See Fig. S10D for covariation within species.

FIG. S10.   **Covariation between maximum growth rate and specific affinity by resource.** (A) Covariation of maximum growth rate $g^{\max}$ and specific affinity $g^{\max}/K$ across all resources and isolates (from Fig. S8A). Marker shapes distinguish autotrophs (squares) from heterotrophs (circles); colors indicates the limiting resource, with the number of measurements n given in brackets. We compute the Spearman rank correlation ρ and p-value across the pooled set of isolates. (B) Subset of measurements from panel A for phosphate (only autotroph isolates shown). (C) Subset of measurements from panel A for nitrate. (D) Subset of measurements from panel A for silicate. (E) Covariation between maximum growth rate $g^{\max}$ and glucose specific affinity $g^{\max}/K$ for measurements of *E. coli* (green) and *S. cerevisiae* (orange), with Spearman rank correlations ρ and p-values by species.

FIG. S11. **Covariation between uptake and growth rate parameters for phosphate based on the phytoplankton trait database by Edwards et al. [43]** (A) Covariation between the half-saturation constant k for uptake rate and the resource threshold K, also known has the half-saturation constant for growth rate in the Monod model (Eq. (1)). The dashed diagonal line indicates perfect agreement ($x = y$), and we calculate the Spearman rank correlation ρ with p-value. We show all datapoints from Edwards et al. [43] which included half-saturation constants for uptake and growth rate. These data points are for phosphate as the limiting resource. (B) Covariation between maximum uptake rate v^{max} in the Michaelis-Menten model and the maximum growth rate g^{max} in the Monod model, with Spearman rank correlation ρ and p-value. The data shown here corresponds to the same measurements as in panel A but with one fewer data point, since one isolate lacked the measurement for maximum growth rate. Color and marker shape are equivalent to Fig. 2A and indicate that the subset of data shown here includes only eukaryotic organisms (orange fill) capable of autotrophy (square shape).

FIG. S12. **Selection within a batch growth cycle.** (A) The fold-change spectrum (thick lines) throughout the growth cycle for high and low initial concentration R_0 (dotted lines). Curves are computed from the weight term in Eq. (S42) with effective biomass yield $\bar{Y} = 1$ and $N_0 = 0.01$. (B) The transient resource level for increasing initial concentration of resource (gray lines). The lowest line ($R_0 = 1$) corresponds to the resource trajectory for the selection scenario used for the phase diagram in Fig. S14. The transient resources are converted into generations using the equations for resource consumption, assuming an identical biomass yield $Y = 1$ for both strains and initial biomass $N_0 = 0.01$.

FIG. S13. **Test of the selection coefficient approximation.** The predicted selection coefficient across a sample of wild-type and mutant strains, compared to the selection coefficient from simulation of the differential equations. The black diagonal line indicates perfect agreement between simulation and prediction. We draw the wild-type traits over four orders of magnitude and sample relative mutant effects on maximum growth rate, resource threshold, and biomass yield from a rectangular region in trait space: $[-0.5, 0.5]^3$. Each strain pair is systematically evaluated at different frequencies $x = 0.01, 0.5, 0.99$ using the general Eq. (S47) and contributes three data points. Without loss of generality, we fix the initial biomass to $N_0 = 0.01$ and initial resource concentration to $R_0 = 1$. The trait values for the resource threshold span two orders of magnitude around this concentration such that we cover both limiting scenarios with dominant selection on maximum growth rate ($R_0 \gg K_1$) and resource threshold ($R_0 \ll K_1$).

FIG. S14. **Diagram of selection across mutation effects under batch growth.** The space of mutation effects on maximum growth rate $\gamma = (g_{\mathrm{mut}}^{\max} - g_{\mathrm{wt}}^{\max})/g_{\mathrm{wt}}^{\max}$ and resource threshold $\kappa = (K_{\mathrm{mut}} - K_{\mathrm{wt}})/K_{\mathrm{wt}}$ relative to a wild-type strain (central dot), with green marking the space of mutations that are overall beneficial ($s > 0$) and red marking mutations that are overall deleterious ($s < 0$) according to Eq. (S47).

FIG. S15. **The gleaner and opportunist strategies in the Monod growth model.** (A) The growth rate $g(R)$ as a function of the external resource concentration R for two strains with a tradeoff. The opportunist strain (green) has a higher maximum growth rate $g^{\max} = 1.5$ compared to the gleaner strain (orange) with $g^{\max} = 1$. But the gleaner has the growth rate advantage at low concentrations due to a smaller resource threshold $K = 0.5$ (orange dotted line) relative to the opportunist with $K = 10$ (green dotted line). (B) A single growth cycle for the gleaner and opportunist strain pair from panel A. We simulate the population dynamics according to Eq. (S11), starting from an initial mutant frequency $x = 0.5$ and total initial biomass $N_0 = 0.01$. On a separate axis, the transient resource concentration R (gray line, initial value $R_0 = 24$) and in the panel title, the components of selection as computed from Eq. (S47).

FIG. S16. **Selection dependence on resource concentration under fixed-bottleneck batch dynamics.** (A) Dependence on external resource concentration R_0 of the total selection coefficient (purple) and its two components at high (s_{high}, red) and low (s_{low}, blue) resource concentrations under a fixed bottleneck (Eq. (S47)). (B) Selection s_{low} on growth at low resource concentrations (top panel) decomposed into two constituent factors (bottom panel) at fixed bottleneck biomass ($N_0 = 10^{-3}$). The top panel is the same as panel A for the approximate selection coefficient s_{low}; in the bottom panel, the two factors that constitute s_{low} with the trait factor from Eq. (S60) in green and the log term from Eq. (S61) in gray. The log term is related but not identical to the number of generations in the growth cycle. Panels A and B are based on an example mutation with relative effects $\gamma = 0.01$ on maximum growth rate and $\kappa = -0.04$ on resource threshold over the wild-type traits $g_{wt} = 1$ and $K_{wt} = 1$. (C) Selection on the threshold K as a function of resource concentration R_0 for three different values of K (different shades of blue). The inset shows a numerical calculation (orange points) of the optimal resource concentration R_{opt} that maximizes selection on K as a function of the wild-type threshold K_{wt}; the gray line is the identity. Parameters are the same as in panels A and B, but we include two alternative wild-type resource thresholds $K_{wt} = 10$ (lightest blue) and $K_{wt} = 0.1$ (darkest blue). (D) The distribution of beneficial selection coefficients (purple) as a function of initial resource concentration R_0, with the variance (which is proportional to the speed of adaptation) shown as the dashed gray line and plotted against the right axis. The inset shows the underlying sample of mutations according to their relative effects on maximum growth rate γ and resource threshold κ. We sample the effects of mutations from independent Gaussian distributions for γ (mean $\mu = 0$, s.d. $\sigma = 0.01$) and κ (mean $\mu = 0$, s.d. $\sigma = 0.02$). All panels assume initial population biomass $N_0 = 0.001$, initial mutant frequency $x = 0.01$, and equal yield for mutant and wild-type.

FIG. S17. **Selection dependence on resource concentration under fixed-dilution batch dynamics.** Same as Fig. S16A, but for fixed-dilution batch dynamics with $D = 1000$.

FIG. S18. **Schematic of evolutionary dynamics in the strong-selection weak-mutation (SSWM) regime.** Top panel shows a schematic of the population biomass undergoing cycles of batch dynamics with serial transfers. The middle panel shows the genetic composition of the population. The population begins with a resource threshold K_1. Then a mutation arises with a different threshold K_2 (blue), which increases in frequency until it fixes. Then another mutation with a threshold K_3 arises (magenta), and the process continues. The bottom panel shows a simplified algorithm for this process that we use in our simulations (Supplementary Information Sec. S11), where mutations are determined to fix or go extinct one at a time based on their selection coefficients, without explicitly simulating their intermediate frequency dynamics.

FIG. S19. **Selection-drift balance under batch dynamics.** (A) Selection s_{low} on the resource threshold K as a function of the current wild-type trait value in the population (blue line, Eq. (S47)), with mutation effect $\kappa = -10^{-2}$. The horizontal black line marks the strength of genetic drift; its intersection with the selection coefficient defines the value of K_{evo} at which selection-drift balance occurs (vertical dotted line; Eq. (3)). Above this point, selection is stronger than genetic drift, and so the resource threshold will adapt downward until it reaches that point. (B) Selection on the specific affinity $a = g^{\text{max}}/K$ as a function of the current wild-type trait value in the population (orange line, Eq. (S52c)) assuming a relative mutation effect $\alpha = 10^{-2}$ that acts directly on the specific affinity instead of on the resource threshold. For the specific affinity, adaptation means the trait value increases. Similar panel A, the intersection of the selection coefficient with the black line (strength of genetic drift) defines the evolved trait value a_{evo} at selection-drift balance. Parameters are identical in both panels with $R_0 = 1$, $N_0 = 0.01$, $N_{\text{e}} = 10^3$, and $x = 0.001$. We set mutant and wild-type to equal maximum growth rate and equal yield. This plot is based on fixed bottleneck biomass N_0, but we observe similar dependences for fixed dilution factor D. In that case, we rewrite Eq. (S47) (resp. Eq. (S52c)) in terms of D (replacing N_0 using Eq. (S21)) and see that the selection coefficient depends on the wild-type trait K with the same functional form.

FIG. S20. **The evolved resource threshold as a function of experimental parameters with independent genetic drift.** (A) We numerically solve for the evolved resource threshold K_{evo} under selection-drift balance (using Eqs. (2) and (3)) as a function of the fixed-bottleneck biomass N_0 and initial resource concentration R_0, where the effective population size $N_{\text{e}} = 10^5$ is an independent parameter. Where the selection-drift balance condition is infeasible (gray area), the resource threshold evolves neutrally without steady state. The insets show cross-sections along initial resource concentration R_0 (solid line) and bottleneck biomass N_0 (dashed line). (B) Same as panel A, but for a larger effective population size $N_{\text{e}} = 10^7$. (C) Same as panel A but for fixed-dilution batch dynamics, with varying D instead of N_0. (D) Same as panel C, but for a larger effective population size $N_{\text{e}} = 10^7$. All panels use identical growth rate $g^{\text{max}} = 1$ and biomass yield $Y = 1$ for wild-type and mutant strain with a fixed mutation effect $\kappa = 0.01$ on the resource threshold. The initial mutant frequency $x = 1/N_{\text{e}}$ is adjusted to the effective population size N_{e}.

FIG. S21. **Simulated evolutionary trajectories of the resource threshold under batch dynamics.** We simulate the time-course of resource threshold evolution in the SSWM regime (Supplementary Information Sec. S11) for different strengths of genetic drift ($1/N_e$). Each line corresponds to a separate run of the stochastic evolution process. Here mutations are sampled with random effect $\kappa = (K_{\mathrm{mut}} - K_{\mathrm{wt}})/K_{\mathrm{wt}}$ from a uniform distribution in $[-0.1, 0.1]$ and accepted or rejected according to their probability of fixation (Supplementary Information Sec. S11). (A)–(C) In the top row, we use the approximate fixation probability Eq. (S63) which is based on the selection coefficient at the initial mutant frequency $x = 1/N_e$. (D)–(F) In the bottom row, we use the integrated form of the fixation probability from Eq. (S62) that takes into account the frequency-dependence of the mutant selection coefficient. For each panel, we numerically calculate the resource threshold K_{evo} at selection-drift balance (dashed line) using Eqs. (2) and (3). To guide the eye, we also mark the resource threshold $K = R_0$ that matches the environmental concentration (gray line). All panels are based on identical maximum growth rate $g^{\mathrm{max}} = 1$ and biomass yield $Y = 1$ for the mutant and wild-type strain such that only the resource threshold evolves. The length of the growth cycle is constant with ≈ 6.6 generations at initial resource concentration $R_0 = 1$ and fixed-bottleneck biomass $N_0 = 0.01$.

effective population size

FIG. S22.   **Simulated evolutionary trajectories for the specific affinity under batch dynamics.** We simulate the time-course of evolution in the SSWM regime (Supplementary Information Sec. S11) similar to Fig. S21, but assuming that mutations directly affect the specific affinity $a = g^{\max}/K$ instead of the threshold K alone. Here mutations are sampled with random effect $\alpha = (a_{\mathrm{mut}} - a_{\mathrm{wt}})/a_{\mathrm{wt}}$ from a uniform distribution in $[-0.1, 0.1]$ and accepted or rejected according to their probability of fixation (compare also Sec. S14). Here we use the approximate fixation probability Eq. (S63) which is based on the selection coefficient at the initial mutant frequency $x = 1/N_{\mathrm{e}}$. The panels (A)–(C) only differ in the effective population size N_{e} used for the simulation. For each panel, we numerically calculate the resource threshold a_{evo} at selection-drift balance (dashed line) using Eqs. (S52c) and (3). All panels are based on identical maximum growth rate $g^{\max} = 1$ and biomass yield $Y = 1$ for the mutant and wild-type strain such that only the specific affinity evolves. The length of the growth cycle is constant with ≈ 6.6 generations at initial resource concentration $R_0 = 1$ and fixed-bottleneck biomass $N_0 = 0.01$.

FIG. S23.   **Detecting the resource threshold K from time-series data.** We use an initial resource concentration $R_0 = 10$ close to the resource threshold of the wild-type strain ($K_{\mathrm{wt}} = 5$, see Fig. 1) to simulate a monoculture growth curve from Eq. (S11) (Supplementary Information Sec. S3). The population leaves steady exponential growth phase (gray area) to enter deceleration (white area). To fit the resource threshold K, the time-series must include multiple data points in the deceleration phase (orange dots). On the right axis, a bracket marks the fold-change from the onset of deceleration at biomass to the saturation.

FIG. S24. **Environmental dependence of selection and genetic drift under batch dynamics.** The top panel shows selection s_{low} on the threshold K (blue solid line, left axis) and the strength of genetic drift $1/N_e$ (dashed black line, right axis) as functions of the resource concentration R_0 under fixed-bottleneck batch dynamics. In this case, the effective population size is independent of the resource concentration. We use parameters based on the LTEE (same as in Fig. 5C): $N_0 = 4.6 \times 10^5$ cells/mL and $N_e = V N_0$ using culture volume $V = 10$ mL, $g^{max} = 0.888$/h, and $Y = 3.3 \times 10^8$ cells/µmol [23]. The bottom panel shows the same but for fixed-dilution batch dynamics, with $D = 100$; in this case the effective population size is proportional to the resource concentration, and thus the strength of genetic drift decreases with R_0.

FIG. S25. **The evolved resource threshold as a function of experimental parameters under coupled genetic drift.** (A) Same as Fig. S20A, but where the effective population size N_e is proportional to the biomass bottleneck N_0 as in well-mixed laboratory experiments with fixed-bottleneck batch dynamics. We set $N_e = N_0 V$, where $V = 10^5$ is the culture volume such that a bottleneck biomass of $N_0 = 0.01$ corresponds to an effective population size of $N_e = 10^3$ cells. (B) Same as panel A but for fixed-dilution batch dynamics, where the effective population size is $N_e = N_0 V = V R_0 Y/(D-1)$ (Eq. (S21)).

FIG. S26. **Inferred mutation effect for the Long-Term Evolution Experiment.** Evolved glucose threshold K_{evo} as a function of the dilution factor D under fixed-dilution batch dynamics. If we assume the glucose threshold for *E. coli* in the LTEE is under selection-drift balance, then we can use this dependence to infer the value of the mutation effect κ that would be consistent with the other known parameters of the system. We numerically solve for selection-drift balance using Eqs. (2) and (3) with dilution factor $D = 100$, initial glucose concentration $R_0 = 139 \,\mu\text{M}$, and evolved resource threshold $K = 0.489 \,\mu\text{M}$ (red dot). We obtain an estimate of $\kappa = 6 \times 10^{-6}$.

Supplementary Information:
Microbial population dynamics decouple nutrient affinity from environmental concentration

Justus Wilhelm Fink*

Institute of Integrative Biology, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland

Noelle A. Held

Institute of Biogeochemistry and Pollutant Dynamics,
Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland and
Department of Environmental Microbiology, Swiss Federal Institute
of Aquatic Science and Technology (Eawag), Dübendorf, Switzerland

Michael Manhart†

Institute of Integrative Biology, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland
Department of Environmental Microbiology, Swiss Federal Institute of
Aquatic Science and Technology (Eawag), Dübendorf, Switzerland and
Center for Advanced Biotechnology and Medicine and Department of Biochemistry and Molecular Biology,
Robert Wood Johnson Medical School, Rutgers University, Piscataway, NJ, USA
(Dated: October 3, 2022)

S1. EFFECT OF COLIMITATION ON ESTIMATES OF MONOD GROWTH TRAITS

The Monod model (Eq. (1)) assumes there is only a single limiting resource whose concentration affects growth rate. However, microbes rely on multiple resources to grow, and therefore their growth rate may depend on the concentrations of all these resources simultaneously. Here we address how these other resources would affect the estimation of Monod growth traits for a focal resource. For simplicity, we consider the case of two essential, independent resources, where resource 1 is the focal resource (e.g., glucose) that we vary over a range of concentrations to measure its Monod parameters $g_1^{\max}$ and K_1, and resource 2 is another resource (e.g., ammonium) that is fixed in the background medium. While there is no consensus on the best model for this behavior, we consider three of the most widely-used models:

Liebig model [1–3]:

$$g(R_1, R_2) = \min\left(\frac{g_1^{\max} R_1}{R_1 + K_1}, \frac{g_2^{\max} R_2}{R_2 + K_2} \right), \quad \text{(S1)}$$

Additive model [1, 4]:

$$g(R_1, R_2) = g^{\max} \frac{R_1 R_2}{K_2 R_1 + R_1 R_2 + K_1 R_2}, \quad \text{(S2)}$$

Multiplicative model [2, 4]:

$$g(R_1, R_2) = g^{\max} \left(\frac{R_1}{R_1 + K_1} \right) \left(\frac{R_2}{R_2 + K_2} \right). \quad \text{(S3)}$$

Assuming one of these models is the true description of how growth rate depends on resource concentrations, we imagine fitting an apparent Monod model $g_{\mathrm{app}}(R_1)$ for resource 1 to data generated by the true model, with fixed R_2:

$$g_{\mathrm{app}}(R_1) = g_{1,\mathrm{app}}^{\max}(R_2) \frac{R_1}{R_1 + K_{1,\mathrm{app}}(R_2)}, \quad \text{(S4)}$$

where $g_{1,\mathrm{app}}^{\max}$ is the apparent maximum growth rate for resource 1 and $K_{1,\mathrm{app}}$ is its apparent threshold, both of which may depend on the concentration R_2 of resource 2. All of the true models correspond exactly to the apparent Monod model — with apparent parameters equaling the true ones, $g_{1,\mathrm{app}}^{\max} = g_1^{\max}$ and $K_{1,\mathrm{app}} = K_1$ — if the concentration R_2 is much larger than its threshold K_2, since the growth rate no longer depends on resource 2 once its concentration is saturating. Therefore $R_2 \gg K_2$ is the general condition on the background resource which determines whether we are in the desired regime of limitation only for resource 1.

If the concentration R_2 is smaller or not much larger than its threshold K_2, we can then use the models to determine how colimitation with resource 2 affects estimates of Monod parameters for resource 1. For all of the true models, the apparent maximum growth rate $g_{1,\mathrm{app}}^{\max}$ is an underestimate of the true maximum growth rate $g_1^{\max}$. Specifically,

---

* To whom correspondence should be addressed. Email: justus.fink@env.ethz.ch

† To whom correspondence should be addressed. Email: mmanhart@rutgers.edu

Liebig model:

$$g_{1,\text{app}}^{\max}(R_2) \approx \min\left(g_1^{\max}, \frac{g_2^{\max}R_2}{R_2+K_2}\right), \quad (S5)$$

Additive model: $g_{1,\text{app}}^{\max}(R_2) = g^{\max}\frac{R_2}{R_2+K_2}, \quad (S6)$

Multiplicative model:

$$g_{1,\text{app}}^{\max}(R_2) = g^{\max}\frac{R_2}{R_2+K_2}. \quad (S7)$$

That is, the apparent maximum growth rate for resource 1 is a Monod-type function of resource 2, meaning it is very close to the true $g_1^{\max}$ for large R_2 as expected, but becomes a significant underestimate when R_2 is below its threshold K_2. Note that the apparent parameters for the Liebig model are only approximate because the Liebig model will not exactly fit the Monod model for a single resource; this is because at some concentration there is a sharp transition in limitation between resources, owing to the minimum function.

The apparent threshold $K_{1,\text{app}}$ is also an underestimate of the true K_1 for the Liebig and additive models, but equals the true value for the multiplicative model:

Liebig model:

$$K_{1,\text{app}}(R_2) \approx \frac{K_1}{2\frac{g_1^{\max}}{\min\left(g_1^{\max},\frac{g_2^{\max}R_2}{R_2+K_2}\right)}-1}, \quad (S8)$$

Additive model: $K_{1,\text{app}}(R_2) = K_1\frac{R_2}{R_2+K_2}, \quad (S9)$

Multiplicative model: $K_{1,\text{app}}(R_2) = K_1. \quad (S10)$

Note also that this means the apparent specific affinity $g_{1,\text{app}}^{\max}/K_{1,\text{app}}$ is always correct for the additive model, since the dependence on R_2 cancels out between the apparent maximum growth rate and threshold, while it is biased for the Liebig and multiplicative models.

To what extent might these biases affect our data? We can test this condition in a subset of measurements for *E. coli* on glucose where the nitrogen source is ammonium and has a reported concentration. The measured K for ammonium in *E. coli* is 2.6 μM (Dataset S1, sheet 1). In the experiments that measure K for glucose, the ammonium concentrations are all orders of magnitude higher (0.16 mM to 18.7 mM; Dataset S1, sheet 2). This indicates that ammonium was not colimiting with glucose in these experiments. Indeed, for almost all the resources included in our data, the K thresholds are much lower than typical laboratory concentrations. This is not surprising in light of our evolutionary model that predicts K will often evolve to be much lower than the environmental concentration of the resource (Eq. (4)), and presumably why colimitation of essential independent resources has been rarely observed empirically [3].

S2. ALTERNATIVE MODELS OF GROWTH RATE DEPENDENCE ON RESOURCE CONCENTRATIONS

Table S1 lists several common models for growth rate dependence on resource concentration R. Some of these models are mathematically equivalent; for example, Holling [11] proposed a classification scheme for growth models (commonly referred to as Type I, II, and III) for the response of predator growth rate on prey density, which exactly corresponds to other models of growth in Table S1. Some of these models are also equivalent in certain limits. At high resource concentrations $R/K \gg 1$, all of the models are approximately equivalent to the constant growth model, since the assumption is that resources are saturating and growth is limited by other processes. On the other hand, at low concentrations $R/K \ll 1$, the Monod, Blackman, and Bertalanffy models are approximately equivalent to the linear model.

There are also some important differences between models. The Blackman, Monod, Bertalanffy, and Hill models all saturate at high resource concentrations, but the nature of that saturation qualitatively differs. That is, the Monod model converges most slowly due its power law dependence on R. The Hill model also converges as a power law, but assuming $n > 1$, it does so more quickly than Monod. The Bertalanffy model converges even more rapidly due to its exponential dependence on R. Finally, the Blackman model converges to a constant immediately at the threshold concentration $R = K$.

The model most significantly different from the rest is the Droop model, since it depends not on the external resource concentration directly, but only on the resource concentration internal to the cell. Therefore this requires inclusion of a separate resource uptake process to be included in our framework. Under steady-state (chemostat) growth, this will also be equivalent to the Monod model under a shift in the resource concentration parameter $Q - Q_0 \to R$, but under non-steady state conditions (e.g., batch dynamics), the Droop model can differ [16].

S3. MODEL OF BATCH POPULATION DYNAMICS

For batch culture we describe the dynamics of the wild-type and mutant biomasses $N_{\text{wt}}(t)$ and $N_{\text{mut}}(t)$ and the extracellular resource concentration $R(t)$ using the following differential equations [20, 21]:

model	definition	references
constant	$g(R) = g^{\max}\Theta(R)$	[5–7]
linear	$g(R) = g^L \cdot \frac{R}{K}$	[8]
Blackman or Holling Type I	$g(R) = g^{\max} \cdot \left(1 + \left(\frac{R}{K} - 1\right)\Theta(K-R)\right)$	[9–11]
Monod or Holling Type II	$g(R) = g^{\max} \cdot \frac{R}{R+K}$	[11, 12]
Droop (depends on internal concentration Q)	$g(Q) = g^{\max} \cdot \frac{Q-Q_0}{Q}$	[13–16]
Bertalanffy	$g(R) = g^{\max}\left(1 - e^{-R/K}\right)$	[4, 17]
Hill, Moser, or Holling Type III	$g(R) = g^{\max} \cdot \frac{R^n}{R^n + K^n}$	[11, 18, 19]

TABLE S1. **Overview of models for microbial population growth rate.** For each entry, the column "references" lists works that establish or build on the model and have been cited elsewhere in this text. The symbol Θ denotes the Heaviside step function which is 1 for a positive argument and zero otherwise.

definition		definition	
biomass concentrations	$N_{\text{wt}}(t), N_{\text{wt}}(t)$	effective growth rate	$\bar{g}(R) = \frac{1-x}{Y_{\text{wt}}/\bar{Y}} \cdot g_{\text{wt}}(R) + \frac{x}{Y_{\text{mut}}/\bar{Y}} g_{\text{mut}}(R)$
initial mutant frequency	x		
extracellular resource conc.	$R(t)$	effective yield	$\bar{Y} = \left[\frac{1-x}{Y_{\text{wt}}} + \frac{x}{Y_{\text{mut}}}\right]^{-1}$
initial biomass concentration	N_0		
initial resource concentration	R_0	effective max. growth rate	$\bar{g}^{\max} = \frac{1-x}{Y_{\text{wt}}/\bar{Y}} \cdot g_{\text{wt}}^{\max} + \frac{x}{Y_{\text{mut}}/\bar{Y}} \cdot g_{\text{mut}}^{\max}$
population growth rates	$g_{\text{wt}}(R), g_{\text{mut}}(R)$		
biomass yields	$Y_{\text{wt}}, Y_{\text{mut}}$	critical concentration	$Z = K_{\text{wt}} K_{\text{mut}} \left[\frac{g_{\text{wt}}^{\max}/\bar{g}^{\max}}{Y_{\text{wt}}/\bar{Y}} \frac{1-x}{K_{\text{wt}}} + \frac{g_{\text{mut}}^{\max}/\bar{g}^{\max}}{Y_{\text{mut}}/\bar{Y}} \frac{x}{K_{\text{mut}}}\right]$
max. growth rates	$g_{\text{wt}}^{\max}, g_{\text{mut}}^{\max}$		
resource thresholds	$K_{\text{wt}}, K_{\text{mut}}$		
specific affinity	$a = g^{\max}/K$		

TABLE S2. **Key notation and definitions used in the model.** The subscripts "wt" and "mut" correspond to *wild-type* and *mutant*. Sometimes we drop the subscript "wt" and use a plain letter (K or $g^{\max}$ or a) for the wild-type trait (for example, in the main text).

$$\frac{1}{N_{\text{wt}}} \frac{dN_{\text{wt}}}{dt} = g_{\text{wt}}(R), \qquad N_{\text{wt}}(0) = (1-x)N_0,$$

$$\frac{1}{N_{\text{mut}}} \frac{dN_{\text{mut}}}{dt} = g_{\text{mut}}(R), \quad N_{\text{mut}}(0) = xN_0,$$

$$\frac{dR}{dt} = -\frac{1}{Y_{\text{wt}}} \frac{dN_{\text{wt}}}{dt} - \frac{1}{Y_{\text{mut}}} \frac{dN_{\text{mut}}}{dt}, \qquad R(0) = R_0.$$
$$\text{(S11)}$$

See Table S2 for a summary of the main notation and
definitions used throughout this article. Growth begins
with an external resource concentration R_0 and total
biomass N_0, a fraction x of which is the mutant strain.
The strains then grow with per-capita rates $g_{\text{wt}}(R)$ and
$g_{\text{mut}}(R)$, which depend on the extracellular resource con-
centration $R(t)$; here we neglect other growth dynam-
ics such as lag [5, 6] and death [22] for simplicity, but
they are straightforward to add within this framework.
The resource concentration $R(t)$ declines in proportion to
growth of biomass, where the yields Y_{mut} and Y_{wt} for each
strain set the amount of new biomass per unit resource.
Here we neglect resource consumption due to mainte-

nance of existing biomass [23], since we expect consump-
tion for maintenance to be much less than consumption
for growth during rapid growth. Growth continues until
the resource is depleted or the growth rates reach zero.
While it is difficult to analytically solve these dynamics
in general, it is straightforward to numerically solve the
model for a given set of parameters (Sec. S4).
We note that for the Monod model in the limit of low
resource concentration R, or any model of growth rate
that depends approximately linearly on R (Table S1),
the batch dynamics of Eq. S11 are equivalent to a lo-
gistic growth model. We can integrate the equation for
resource consumption dR/dt in Eq. S11 to express the
current resource concentration $R(t)$ as a function of the
biomasses of wild-type N_{wt} and mutant strain N_{mut}:

$$R = R_0 + \frac{N_0}{\bar{Y}} - \frac{N_{\text{wt}}}{Y_{\text{wt}}} - \frac{N_{\text{mut}}}{Y_{\text{mut}}}, \qquad \text{(S12)}$$

167 where $\bar{Y}$ is the effective population yield (Table S2). Sub-
168 stituting R from Eq. (S12) into the equations for dN_{wt}/dt
169 and dN_{mut}/dt from Eq. S11 with linear growth rate de-
170 pendence ($g(R) \approx g^{\max} R/K$), we obtain

$$\frac{1}{N_{\text{wt}}}\frac{dN_{\text{wt}}}{dt} = \frac{g_{\text{wt}}^{\max}}{K_{\text{wt}}}\left(R_0 + \frac{N_0}{\bar{Y}} - \frac{N_{\text{wt}}}{Y_{\text{wt}}} - \frac{N_{\text{mut}}}{Y_{\text{mut}}}\right),$$
(S13a)

$$\frac{1}{N_{\text{mut}}}\frac{dN_{\text{mut}}}{dt} = \frac{g_{\text{mut}}^{\max}}{K_{\text{mut}}}\left(R_0 + \frac{N_0}{\bar{Y}} - \frac{N_{\text{wt}}}{Y_{\text{wt}}} - \frac{N_{\text{mut}}}{Y_{\text{mut}}}\right).$$
(S13b)

This is equivalent to logistic growth for both species or competitive Lotka-Volterra dynamics.

Once the resource R is depleted during a single cycle of batch growth, we transfer a fraction $1/D$ of the population to an environment with a new supply of resources at the original concentration R_0, after which the population resumes growth in the new environment according to Eq. (S11). The factor D is known as the dilution factor and is the ratio of the total biomass at the end of the previous growth cycle and the total biomass at the beginning of the next growth cycle [7].

In principle the dilution factor D and the bottleneck biomass concentration N_0 can vary over each growth cycle, depending on how we perform the transfers. Let superscript n refer to the dynamics during the nth batch growth cycle over a series of dilutions and transfers. The biomass at the beginning of the $(n+1)$th cycle, $N_0^{(n+1)}$, equals the biomass at the end of the previous cycle n divided by the dilution factor $D^{(n)}$ for that cycle:

$$N_0^{(n+1)} = \frac{1}{D^{(n)}}\left(N_{\text{wt}}^{(n)}(t_{\text{sat}}) + N_{\text{mut}}^{(n)}(t_{\text{sat}})\right),$$
(S14)

where t_{sat} is the saturation time of the growth cycle. To determine the relationship with the bottleneck size of the previous growth cycle, we use the relationship between resource and biomass concentrations (Eq. (S12)) to show that at the end of the growth cycle, $R(t_{\text{sat}}) = 0$, and so

$$R^{(n)}(t_{\text{sat}}) = 0$$
$$= R_0 + \frac{N_0^{(n)}}{\bar{Y}^{(n)}} - \frac{N_{\text{wt}}^{(n)}(t_{\text{sat}})}{Y_{\text{wt}}} - \frac{N_{\text{mut}}^{(n)}(t_{\text{sat}})}{Y_{\text{mut}}}.$$
(S15)

Using this, we can insert the identity to obtain

$$N_0^{(n+1)} = \frac{1}{D^{(n)}}\left(N_{\text{wt}}^{(n)}(t_{\text{sat}}) + N_{\text{mut}}^{(n)}(t_{\text{sat}})\right)$$
$$= \frac{1}{D^{(n)}}\left(\frac{R_0 + \frac{N_0^{(n)}}{\bar{Y}^{(n)}}}{\frac{N_{\text{wt}}^{(n)}(t_{\text{sat}})}{Y_{\text{wt}}} + \frac{N_{\text{mut}}^{(n)}(t_{\text{sat}})}{Y_{\text{mut}}}}\right)$$
$$\cdot\left(N_{\text{wt}}^{(n)}(t_{\text{sat}}) + N_{\text{mut}}^{(n)}(t_{\text{sat}})\right)$$
$$= \frac{1}{D^{(n)}}\left(R_0 + \frac{N_0^{(n)}}{\bar{Y}^{(n)}}\right)\left(\frac{1 - x^{(n+1)}}{Y_{\text{wt}}} + \frac{x^{(n+1)}}{Y_{\text{mut}}}\right)^{-1}$$
(S16)

where we have used the fact that the frequencies of each strain at the end of the nth cycle equal their frequencies at the beginning of the $(n+1)$th cycle:

$$\frac{N_{\text{mut}}^{(n)}(t_{\text{sat}})}{N_{\text{wt}}^{(n)}(t_{\text{sat}}) + N_{\text{mut}}^{(n)}(t_{\text{sat}})} = x^{(n+1)}.$$
(S17)

Using the equation for the effective yield (Table S2), we obtain

$$N_0^{(n+1)} = \frac{1}{D^{(n)}}\left(R_0 + \frac{N_0^{(n)}}{\bar{Y}^{(n)}}\right)\bar{Y}^{(n+1)}.$$
(S18)

This establishes the general relationship between the bottleneck size and the dilution factor.

Under fixed-bottleneck batch dynamics (Fig. 4B, top panel), $N_0^{(n)}$ is a constant value N_0, and so we can rearrange Eq. (S18) to determine how the dilution factor varies at each cycle:

$$D^{(n)} = \frac{R_0 \bar{Y}^{(n+1)}}{N_0} + \frac{\bar{Y}^{(n+1)}}{\bar{Y}^{(n)}}.$$
(S19)

This shows that the dilution factor changes only if the strains have different yields, such that the effective yields $\bar{Y}^{(n)}$ change over cycles as the strain frequencies change. On the other hand, under fixed-dilution batch dynamics (Fig. 4B, bottom panel), $D^{(n)}$ is a constant D, and Eq. (S18) simplifies to

$$N_0^{(n+1)} = \frac{1}{D}\left(R_0 + \frac{N_0^{(n)}}{\bar{Y}^{(n)}}\right)\bar{Y}^{(n+1)}.$$
(S20)

Under both serial transfer regimes, the steady state occurs when $D^{(n+1)} = D^{(n)}$, $N_0^{(n+1)} = N_0^{(n)}$, and $\bar{Y}^{(n+1)} = \bar{Y}^{(n)}$, which implies

$$D = \frac{R_0 \bar{Y}}{N_0} + 1.$$
(S21)

This steady state occurs if 1) all strains have the same yields, such that the effective yield is constant; 2) one strain goes extinct; or 3) the two strains stably coexist.

S4. NUMERICAL METHODS FOR BATCH DYNAMICS

It is not possible to analytically solve the ordinary differential equations for batch dynamics (Eq. S11). To obtain explicit solutions to this model, we therefore numerically integrate the equations using the Scipy package [24]. We use the default Runge-Kutta algorithm "RK45" in the function `solve_ivp`. This interpolates the differential

equation in fourth-order expansion over a short step size δt. The step size is automatically adjusted by `solve_ivp` to keep the error of integration below a threshold fixed by the user through the parameters `atol` and `rtol`. Our choices of `atol` $= 10^{-12}$ and `rtol` $= 10^{-8}$ are more restrictive than the default setting and ensure low errors on the state variables N_{wt}, N_{mut}, and R.

The population dynamics in Eq. S11 reach the final equilibrium when all resources have been converted into biomass. This final equilibrium is the only attractor since the resources are finite and biomass is strictly increasing (no cell death within a batch growth cycle); in particular, this system does not allow for limit cycles. However, the time to reach this equilibrium is infinite for all growth models in Table S1 (including the Monod model) except for the constant growth rate model. This is because the smooth decline of growth rate prevents full depletion of resources and allows populations to grow indefinitely at infinitesimal but strictly positive growth rates. (The constant growth rate model allows for the same growth rate at arbitrarily low resource concentrations, which means the resources deplete to zero in finite time [5–7].)

For numerical calculations we must therefore set a finite saturation time t_{sat} such that the population dynamics are sufficiently close to their equilibrium state. We choose this time using the selection coefficient, which quantifies the relative change in the strain frequencies. Define the cumulative selection coefficient up to time t for a batch growth cycle as

$$s_t = \log\left(\frac{N_{\mathrm{mut}}(t)}{N_{\mathrm{wt}}(t)}\right) - \log\left(\frac{N_{\mathrm{mut}}(0)}{N_{\mathrm{wt}}(0)}\right). \tag{S22}$$

(We further motivate this definition of selection in Sec. S6.) The total selection coefficient for the batch cycle is the cumulative selection coefficient in the limit of infinite time:

$$s = \lim_{t \to \infty} s_t. \tag{S23}$$

We want to define the saturation time t_{sat} as the time where the difference between the cumulative selection up to that time and the total selection is less than some tolerance. We can do this by determining an upper bound on the difference between total selection s and the selection at finite time t. As the population continues to grow after time t, the change in frequencies is bounded by the remaining available resources $R(t)$. The two possible extremes are if all remaining resources go to the wild-type, in which case the biomass of the wild-type increases by $R(t)Y_{\mathrm{wt}}$ and the mutant biomass remains constant, or if all remaining resources go to the mutant, in which case the biomass of the mutant increases by $R(t)Y_{\mathrm{mut}}$ and the wild-type remains constant. Therefore the largest possible change in selection occurs in one of these two scenarios, and so the deviation in selection at time t from its equilibrium value is bounded by

$$|s_t - s| \leq \max\left\{ \log\left(1 + \frac{R(t)Y_{\mathrm{mut}}}{N_{\mathrm{mut}}(t)}\right), \right.$$
$$\left. \log\left(1 + \frac{R(t)Y_{\mathrm{wt}}}{N_{\mathrm{wt}}(t)}\right) \right\}. \tag{S24}$$

We define the saturation time t_{sat} as the shortest time (infimum) such that the difference between the cumulative selection at that time and the total selection is smaller than a given error tolerance $\epsilon > 0$:

$$t_{\mathrm{sat}} = \inf\left\{ t > 0 : |s_t - s| < \epsilon \right\}. \tag{S25}$$

We implement this algorithmically by evaluating the simulation up to an initial time t, then evaluating the maximum future error on the selection coefficient from the right hand side of Eq. (S24), and then extending the simulation to $t + 10$ if the error exceeds a defined tolerance $\epsilon = 10^{-8}$. We iterate this process until the error is less than the threshold.

S5. MODEL OF CHEMOSTAT POPULATION DYNAMICS

Similar to the batch model of Eq. (S11), the dynamics of biomass and resource concentrations under continuous culture (chemostat) are

$$\frac{1}{N_{\mathrm{wt}}}\frac{dN_{\mathrm{wt}}}{dt} = g_{\mathrm{wt}}(R) - d, \qquad N_{\mathrm{wt}}(0) = (1-x)N_0,$$
$$\frac{1}{N_{\mathrm{mut}}}\frac{dN_{\mathrm{mut}}}{dt} = g_{\mathrm{mut}}(R) - d, \quad N_{\mathrm{mut}}(0) = xN_0,$$

$$\frac{dR}{dt} = -g_{\mathrm{wt}}(R)\frac{N_{\mathrm{wt}}(t)}{Y_{\mathrm{wt}}} - g_{\mathrm{mut}}(R)\frac{N_{\mathrm{mut}}(t)}{Y_{\mathrm{mut}}}$$
$$+ d(R_{\mathrm{source}} - R(t)),$$
$$R(0) = R_0, \tag{S26}$$

where R_{source} is the concentration of the resource in the source media fed into the culture. In a laboratory chemostat, the dilution rate is $d = \omega/V$, where ω is the outflow rate (volume per time) and V is the volume of the culture vessel [25].

In the SSWM regime where mutations arise only rarely (Sec. S11), we can assume that the mutant arises on the background of the wild-type at steady-state growth. Let N_{wt}^* be the steady-state concentration of wild-type biomass and R^* be the steady-state concentration of the resource. Note that R^* here is the chemostat-specific realization of the ecological concept of a minimum resource concentration required for positive net growth, as used in resource-ratio theory [26, 27]. Since $dN_{\mathrm{wt}}/dt = 0$ in steady state, the resource concentration R^* must satisfy

$$g_{\mathrm{wt}}(R^*) = d. \tag{S27}$$

For the Monod model, we can solve this explicitly for R^* to obtain

$$R^* = K_{\mathrm{wt}} \frac{d}{g_{\mathrm{wt}}^{\max} - d}. \tag{S28}$$

Note that this concentration R^* is independent of the source concentration R_{source}. Using the steady-state condition for the resource $dR/dt = 0$, we can then obtain the steady-state biomass concentration

$$N_{\mathrm{wt}}^* = (R_{\mathrm{source}} - R^*)Y_{\mathrm{wt}}. \tag{S29}$$

This establishes a feasibility condition for steady state: the dilution rate d must be less than the growth rate at the source concentration $g_{\mathrm{wt}}(R_{\mathrm{source}})$, which is the maximum that the culture can realize for the given resource supply. This criterion has been used by Jannasch [28, 29] to define a *minimum resource threshold* required for population growth at a given dilution factor d. This minimum resource threshold corresponds to the steady-state concentration R^*, which is related to the parameter K but also depends on d.

S6. DEFINITION OF SELECTION COEFFICIENT

The instantaneous selection coefficient $\sigma(t)$ measures the rate of change in the logarithm of relative mutant frequency:

$$\sigma(t) = \frac{d}{dt} \log\left(\frac{N_{\mathrm{mut}}(t)}{N_{\mathrm{wt}}(t)}\right). \tag{S30}$$

This is a sufficient statistic for frequency change in the sense that knowledge of the instantaneous selection coefficient and the current mutant frequency is sufficient to predict the future mutant frequency over a short time horizon.

For population growth under batch dynamics, the repeated bottlenecks between growth cycles introduce randomness in the frequency trajectory of a mutant. We assume that this stochastic sampling at transfer dominates over the random fluctuations in individual birth rates within the growth cycle. Thus, the genetic drift in our model of serial transfer evolution occurs at the timescale of one growth cycle. To compare the strength of drift and selection on the same timescale, we integrate the instantaneous selection coefficient (Eq. (S30)) over time

$$s = \frac{1}{\Delta t} \int_0^{\Delta t} \sigma(t) \ dt, \tag{S31}$$

where Δt is the length of the growth cycle. Note that the selection coefficient s is still defined as a rate per unit time and in the limit $\Delta t \to 0$ exactly matches the instantaneous selection coefficient (Eq. (S30)).

For batch dynamics the selection coefficient s determines the change of frequency over multiple growth cycles. At the beginning of the nth cycle, the initial mutant frequency $x^{(n)}$ is given by

$$x^{(n)} = \frac{N_{\mathrm{mut}}^{(n)}(0)}{N_{\mathrm{mut}}^{(n)}(0) + N_{\mathrm{wt}}^{(n)}(0)}, \tag{S32}$$

where $N_{\mathrm{wt}}^{(n)}$ and $N_{\mathrm{mut}}^{(n)}$ refer to the biomass of wild-type and mutant strains. The population grows to saturation and possibly experiences some frequency change, which sets the mutant frequency $x^{(n+1)}$ of the next cycle. This change is summarized by the selection coefficient

$$s^{(n)} = \log\left(\frac{x^{(n+1)}}{1 - x^{(n+1)}}\right) - \log\left(\frac{x^{(n)}}{1 - x^{(n)}}\right), \tag{S33}$$

which we compute from the integral definition (Eq. (S31)) using a timescale of $\Delta t = 1$ (per growth cycle). Knowledge of $s^{(n)}$ is sufficient to predict the initial mutant frequency in the next cycle

$$x^{(n+1)} = \frac{x^{(n)} \exp(s^{(n)})}{1 + x^{(n)}\left[\exp(s^{(n)}) - 1\right]}, \tag{S34}$$

neglecting the stochastic effects of the dilution. Thus, given the starting mutant frequency $x^{(1)}$ and the selection coefficients for each growth cycle $s^{(n)}$, the recursion in Eq. (S34) allows us to predict the mutant frequency trajectory without simulating the population dynamics within each growth cycle.

S7. DERIVATION OF THE SELECTION COEFFICIENT FOR BATCH DYNAMICS

For populations growing in batch culture, the selection coefficient reduces to the cumulative difference of growth rates:

$$s = \int_0^\infty \left[g_{\mathrm{mut}}(R(t)) - g_{\mathrm{wt}}(R(t))\right] \ dt, \tag{S35}$$

where we have inserted the equations for mutant and wild-type growth from our model of populations dynamics (Eq. S11) into the definition of s from Eq. (S31). The integral extends to infinite time for the growth dynamics to reach equilibrium (Sec. S4) so we therefore do not normalize by the time scale as in Eq. (S31); rather we leave it as understood that the selection coefficient is defined per growth cycle. We can change variables of the integral in Eq. (S35) from time t to resource concentration R:

$$s = \int_{R_0}^0 \left[g_{\mathrm{mut}}(R) - g_{\mathrm{wt}}(R)\right] \cdot \frac{1}{dR/ \ dt} \ dR, \tag{S36}$$

where we have used the fact that R ranges from R_0 at the beginning of the growth cycle to 0 at the end of the growth cycle, and that R depends monotonically on t so that $dt/dR = (dR/dt)^{-1}$.

To compute this integral, we need to express the transient resource consumption rate dR/dt as an explicit function of current resource concentration R. As a first step, we rewrite the differential equation for resources (Eq. S11) into the product form

$$\frac{d}{dt}R(t) = -\frac{N_{\text{wt}}(t) + N_{\text{mut}}(t)}{\bar{Y}} \quad (S37)$$
$$\cdot \left[\frac{1-x(t)}{Y_{\text{wt}}/\bar{Y}} g_{\text{wt}}(R) + \frac{x(t)}{Y_{\text{mut}}/\bar{Y}} g_{\text{mut}}(R) \right],$$

where we use the shorthand $\bar{Y}$ for the effective biomass yield (Table S2). This product separates into the joint biomass $N_{\text{wt}}(t) + N_{\text{mut}}(t)$ and a new parameter, that we term the *effective growth rate*:

$$\bar{g}(t,R) = \frac{1-x(t)}{Y_{\text{wt}}/\bar{Y}} g_{\text{wt}}(R) + \frac{x(t)}{Y_{\text{mut}}/\bar{Y}} g_{\text{mut}}(R). \quad (S38)$$

Equation (S37) suggests that this mean of wild-type and mutant growth rates acts as the effective growth rate of the joint population $N_{\text{wt}}(t) + N_{\text{mut}}(t)$. This effective growth rate is time-dependent due to the underlying frequency change $x(t)$. Using this equation for the joint biomass (derived from Eq. (S12))

$$N_{\text{wt}}(t) + N_{\text{mut}}(t) = \left(R_0 - R(t) + \frac{N_0}{\bar{Y}} \right) \bar{Y}(t), \quad (S39)$$

we insert this and the equation for mean growth rate (Eq. (S38)) into Eq. (S36) for the selection coefficient:

$$s = \int_0^{R_0} \left(\frac{g_{\text{mut}}(R) - g_{\text{wt}}(R)}{\bar{g}(t(R), R)} \right) \cdot \left(\frac{1}{N_0/\bar{Y} + R_0 - R} \right) \, dR. \quad (S40)$$

Equation (S40) is an exact expression but requires full knowledge of the resource trajectory $R(t)$ and its inverse $t(R)$ to calculate the mean growth rate $\bar{g}(t(R), R)$ in the denominator. For a constant growth rate model (Table S1), this exact expression can be computed [5, 6]. However, for general growth models $g(R)$ and the Monod model in particular, the integral Eq. (S40) can only be solved under an approximation. Previous work invoked the assumption of small initial mutant frequency $x \ll 1$ to replace mean growth rate and effective biomass yield by the wild-type traits [20, 21], but here we introduce a novel approximation that holds for all initial mutant frequencies.

We assume that the frequency change over the growth cycle is small, such that the mean growth rate only depends on the resource concentration

$$\bar{g}(R) \approx \frac{1-x}{Y_{\text{wt}}/\bar{Y}} g_{\text{wt}}(R) + \frac{x}{Y_{\text{mut}}/\bar{Y}} g_{\text{mut}}(R), \quad (S41)$$

but not otherwise on time t. That is, we neglect the time dependence of the mutant frequency $x(t)$. Thus, we get the explicit integral formula for the selection coefficient:

$$s \approx \int_0^{R_0} \left(\frac{g_{\text{mut}}(R) - g_{\text{wt}}(R)}{\bar{g}(R)} \right) \cdot \left(\frac{1}{N_0/\bar{Y} + R_0 - R} \right) \, dR. \quad (S42)$$

This equation neglects the frequency change $x(t)$ within the growth cycle but still includes dependence on the initial mutant frequency x. One can show that the approximate integral in Eq. (S42) corresponds to a first-order expansion of the exact integral (Eq. (S40) in terms of transient selection coefficients inside the growth cycle, meaning that it is equivalent to a weak-selection approximation. We numerically evaluate the accuracy of this approximation in the case of the Monod model in the next section (Sec. S8).

The exact selection coefficient in its integral form (Eq. (S40)) reveals generic properties of batch-culture competition. First, there is no direct selection for cell yield. A mutant with higher efficiency Y_{mut} but equal growth response is neutral. Thus, with an uncorrelated mutation supply, we expect cell yield to evolve neutrally [7, 30]. Second, the selection on the growth rate function $g(R)$ is distributed unequally across concentrations. In the integrand of Eq. (S40), the difference in growth rates at each resource concentration R is weighted by the *fold-change spectrum* $1/(N_0/\bar{Y} + R_0 - R)$. This weight peaks at the initial resource concentration R_0 (see Fig. S12A) and is independent of the growth rate model $g(R)$. For growth cycles with large fold-change ($R_0 Y/N_0 \gg 1$), the selection coefficient s roughly corresponds to the growth rate difference at initial concentrations because most generations occur at near-constant concentrations close to R_0 (compare Fig. S12B).

A third important property holds only approximately in Eq. (S42), where we see that selection only acts on ratios of growth rates, since the growth rates appear in both the numerator and denominator of the integrand. The dependence on growth rate ratios means that alternative growth models can still lead to equivalent selection on traits. For example, if we take any growth rate model from Table S1 where the mutant and wild-type differ only in their maximum growth rates g^{max} (but not other parameters such as K), then their selection coefficients will depend only on the ratio $g_{\text{mut}}^{\text{max}}/g_{\text{mut}}^{\text{max}}$ and not other details of the specific model.

S8. CALCULATION OF THE SELECTION COEFFICIENT FOR THE MONOD MODEL

In this section, we apply the integral formula Eq. (S42) to calculate the selection coefficient for a wild-type and mutant strain competing under the Monod model. Let

$$\Delta g^{\max} = g_{\mathrm{mut}}^{\max} - g_{\mathrm{wt}}^{\max}, \quad \Delta K = K_{\mathrm{mut}} - K_{\mathrm{wt}} \quad \text{(S43)}$$

denote the absolute trait differences in maximum growth rate and resource threshold between the two strains. First, we rewrite the relative growth rate difference

$$\frac{g_{\mathrm{mut}}(R) - g_{\mathrm{wt}}(R)}{\bar{g}(R)} = \frac{\Delta g^{\max}}{\bar{g}^{\max}} \quad \text{(S44)}$$
$$- \frac{\Delta K}{R + Z} \cdot \frac{g_{\mathrm{wt}}^{\max} g_{\mathrm{mut}}^{\max}}{\bar{g}^{\max} \bar{g}^{\max}},$$

using the effective maximum growth rate

$$\bar{g}^{\max} = \frac{1-x}{Y_{\mathrm{wt}}/\bar{Y}} \cdot g_{\mathrm{wt}}^{\max} + \frac{x}{Y_{\mathrm{mut}}/\bar{Y}} \cdot g_{\mathrm{mut}}^{\max} \quad \text{(S45)}$$

and the critical resource concentration

$$Z = K_{\mathrm{wt}} K_{\mathrm{mut}} \cdot \left[\frac{g_{\mathrm{wt}}^{\max}/\bar{g}^{\max}}{Y_{\mathrm{wt}}/\bar{Y}} \cdot \frac{1-x}{K_{\mathrm{wt}}} \right. \quad \text{(S46)}$$
$$\left. + \frac{g_{\mathrm{mut}}^{\max}/\bar{g}^{\max}}{Y_{\mathrm{mut}}/\bar{Y}} \cdot \frac{x}{K_{\mathrm{mut}}} \right]$$

as effective traits of the joint population to simplify the notation (Table S2). Equation (S44) consists of two terms, one proportional to the difference in maximum growth rates $\Delta g^{\max}$ and the other proportional to the difference in resource thresholds ΔK. Therefore after substituting this expression into Eq. (S42) and carrying out the integral over R, we obtain a selection coefficient consisting of two distinct components:

$$s \approx s_{\mathrm{high}} + s_{\mathrm{low}} \quad \text{(S47a)}$$

where

$$s_{\mathrm{high}} = \frac{\Delta g^{\max}}{\bar{g}^{\max}} \log\left(1 + \frac{R_0 \bar{Y}}{N_0}\right) \quad \text{(S47b)}$$

$$s_{\mathrm{low}} = -\frac{\Delta K}{R_0 + N_0/\bar{Y} + Z} \left(\frac{g_{\mathrm{wt}}^{\max}}{\bar{g}^{\max}} \frac{g_{\mathrm{mut}}^{\max}}{\bar{g}^{\max}} \right) \quad \text{(S47c)}$$
$$\cdot \log\left(\left(1 + \frac{R_0 \bar{Y}}{N_0}\right) \left(1 + \frac{R_0}{Z}\right) \right).$$

This is the basis for Eq. (2) in the main text under batch dynamics.

The formula for the selection coefficient in Eq. (S42) is based on an approximation of small frequency change. In Fig. S13 we compare the approximate selection coefficient against the exact selection coefficient obtained from numerically solving the differential equations for batch dynamics (Eq. S11). The simulations show that the approximate selection coefficient is accurate up to large values of order $s \approx 1$. This means that, while we mainly consider the scenario of weak selection ($|s| < 1$), the approximation is excellent even when selection is strong. Intuitively, the approximation should break down because of wrongly estimating the mean resource consumption rate, which we expect to occur when the yields and realized growth rates differ strongly between the two strains. In Fig. S14 we also show a phase diagram of this selection coefficient as a function of the mutant's traits $g^{\max}$ and K relative to their wild-type values.

The decomposition in Eq. (S47) is useful because the terms correspond to components of selection on distinct phases of growth. The first component, s_{high}, measures selection on growth at high resource concentrations, and is therefore proportional to the mutational change $\Delta g^{\max}$ in the trait $g^{\max}$. This mutational effect is weighed by the logarithm of the total fold-change of growth, which equals the dilution factor $D = R_0 Y/N_0 + 1$ (Eq. (S21)). An important feature of selection s_{high} is that it depends on the nominal maximum growth rate $g^{\max}$, which is always greater than the realized maximum growth rate $g(R_0)$ that actually occurs at the beginning of growth. Therefore the calculation of selection from actual growth data requires an inference of these nominal rates, since the realized rates measured at the beginning of growth curves could produce misleading results if growth begins at low resource concentrations [31].

The second component of selection, s_{low}, corresponds to growth at low resource concentrations, and is proportional to the mutant's change ΔK of the threshold K. There is a negative sign in s_{low} since selection is positive for mutations that decrease K ($\Delta K < 0$). For the hypothetical mutant and wild-type in Figs. 1 and 4A,B, $s_{\mathrm{high}} = 0$ since the mutant does not change $g^{\max}$, while $s_{\mathrm{low}} \approx 0.516$, since the mutant has a significantly lower threshold K. In Fig. S15 we show a more complex pair of strains with a gleaner-opportunist tradeoff (one strain has higher $g^{\max}$ but also higher K), where both components of selection are nonzero [20, 21, 27, 32].

We also briefly discuss an interpretation for the parameter Z. The instantaneous selection coefficient (Eq. (S30)) within the batch culture growth cycle

$$\sigma(t) = g_{\mathrm{mut}}(R(t)) - g_{\mathrm{wt}}(R(t)) \quad \text{(S48)}$$

can be decomposed into two components

$$\sigma = \sigma_{\max} + \sigma_{\mathrm{lin}}, \quad \text{(S49a)}$$

where

$$\sigma_{\max} = \Delta g^{\max} \frac{R}{R + K_{\mathrm{wt}}} \frac{R}{R + K_{\mathrm{mut}}}, \quad \text{(S49b)}$$

$$\sigma_{\mathrm{lin}} = \Delta a \cdot R \frac{K_{\mathrm{wt}}}{R + K_{\mathrm{wt}}} \frac{K_{\mathrm{mut}}}{R + K_{\mathrm{mut}}}. \quad \text{(S49c)}$$

Here $a = g^{\max}/K$ is the specific affinity, and Δa is the difference in specific affinities between the mutant and the wild-type. The first component $\sigma_{\max}$ quantifies

growth rate difference at excess conditions, where both strains grow close to their maximum growth rates. The second component σ_{lin} measures growth rate differences in the opposite regime, where both strains grow below their resource thresholds. The relative size of the two components varies shifts with resource concentration and also depends on the mutation effect on maximum growth rate and specific affinity.

The effective parameter Z acts as an intrinsic scale in the resource dependence. Normalizing for different relative mutation effects, both components contribute equally to growth rate difference exactly at external concentration $R = Z$ such that

$$\frac{\sigma_{\text{max}}(Z)}{\Delta g^{\text{max}}/\bar{g}^{\text{max}}} = \frac{\sigma_{\text{lin}}(Z)}{\Delta a/\bar{a}}, \qquad (S50)$$

where the effective specific affinity $\bar{a}$ is defined (in analogy with the effective maximum growth rate defined in Eq. (S45)) as

$$\bar{a} = \frac{1-x}{Y_{\text{wt}}/\bar{Y}} \cdot \frac{g_{\text{wt}}^{\text{max}}}{K_{\text{wt}}} + \frac{x}{Y_{\text{mut}}/\bar{Y}} \cdot \frac{g_{\text{mut}}^{\text{max}}}{K_{\text{mut}}}. \qquad (S51)$$

This means, at concentration Z both components of the growth rate difference in Eq. (S49) receive equal selection pressure.

The decomposition in Eq. (S49) more generally suggests an alternative parametrization of the Monod model and its selection coefficient. We can replace the resource threshold K by the specific affinity $a = g^{\text{max}}/K$. This alternative trait corresponds to the growth rate in the limit of low resource concentrations where the Monod model behaves linearly (see Sec. S2). The selection coefficient in Eq. (S47) can be rewritten as

$$s \approx s_{\text{max}} + s_{\text{lin}}, \qquad (S52a)$$

where

$$s_{\text{max}} = \frac{\Delta g^{\text{max}}}{\bar{g}^{\text{max}}} \qquad (S52b)$$
$$\cdot \left[\frac{R_0 + N_0/\bar{Y}}{R_0 + N_0/\bar{Y} + Z} \cdot \log\left(1 + \frac{R_0 \bar{Y}}{N_0}\right) \right.$$
$$\left. - \frac{Z}{R_0 + N_0/\bar{Y} + Z} \cdot \log\left(1 + \frac{R_0}{Z}\right) \right],$$

$$s_{\text{lin}} = \frac{\Delta a}{\bar{a}} \cdot \left[\frac{Z}{R_0 + N_0/\bar{Y} + Z} \right. \qquad (S52c)$$
$$\left. \cdot \log\left(1 + \frac{R_0 \bar{Y}}{N_0}\right)\left(1 + \frac{R_0}{Z}\right) \right].$$

The selection coefficient maps the life-history traits to relative fitness, and the parametrization in a is well-suited to study the structure of this map under environmental variation. In the limit of high nutrient concentrations, the total resources are large compared to the

critical concentration Z. The selection coefficient then reduces to the component of maximum growth:

$$s \approx s_{\text{max}} \qquad \text{as } R_0 \to \infty. \qquad (S53)$$

In the opposite limit, the selection coefficient only acts on the growth rate at low concentrations. In this sense, the selection coefficient recovers the limiting behaviour of the underlying growth response:

$$s \approx s_{\text{lin}} \qquad \text{as } R_0 \to 0. \qquad (S54)$$

This means that g and a are the marginal traits that exclusively control growth in the limiting environments. The selection coefficient reduces to one component or the other. For the parametrization based on K given below, this is not true — both g and K contribute at low concentrations.

S9. DERIVATION OF THE SELECTION COEFFICIENT FOR CHEMOSTAT DYNAMICS

For a population in chemostat conditions (Eq. (S26)), the instantaneous selection coefficient $\sigma(t)$ (Eq. (S30)) only depends on the difference in growth rates. At a given resource concentration $R(t)$, this growth rate difference can be decomposed in to two trait components

$$\sigma \approx \sigma_{\text{high}} + \sigma_{\text{low}}, \qquad (S55a)$$

where

$$\sigma_{\text{high}} = \frac{\Delta g^{\text{max}}}{\bar{g}^{\text{max}}} \bar{g}(R), \qquad (S55b)$$

$$\sigma_{\text{low}} = -\frac{\Delta K}{R + Z} \frac{g_{\text{wt}}^{\text{max}}}{\bar{g}^{\text{max}}} \frac{g_{\text{mut}}^{\text{max}}}{\bar{g}^{\text{max}}} \bar{g}(R). \qquad (S55c)$$

We can derive this by multiplying Eq. (S44) with the mean growth rate for the Monod model

$$\bar{g}(R) = \bar{g}^{\text{max}} \frac{R}{R + K_{\text{wt}}} \frac{R}{R + K_{\text{mut}}} \qquad (S56)$$
$$+ \bar{a}R \frac{K_{\text{wt}}}{R + K_{\text{wt}}} \frac{K_{\text{mut}}}{R + K_{\text{mut}}}.$$

The two components σ_{high} and σ_{low} are consistent with our results for batch culture conditions (Eq. (S47)). By integrating the instantaneous component σ_{high} over the growth cycle, we recover the component s_{high} for batch-culture growth.

We assume a specific scenario for selection in chemostat populations, where mutants arise at small frequency x on top of a wild-type population. This is plausible if mutations occur not too frequently, such that the chemostat population is replaced by a mutant and reaches the new steady state before the next mutation arises. The

wild-type population under steady-state chemostat conditions has a resource concentration given by (Eq. S28)

$$R^* = K_{\text{wt}} \frac{d}{g_{\text{wt}}^{\max} - d}, \tag{S57}$$

where growth rate matches the dilution factor $g_{\text{wt}}(R^*) = d$ (Eq. S27). After the mutant appears, the resource concentration $R(t) \approx R^*$ remains constant over a short timespan while the mutant still has low frequency $x \ll 1$. In this time window, the mean growth rate Eq. (S56) is set by the wild-type only and thus equals the dilution rate:

$$\bar{g}(R^*) \approx d. \tag{S58}$$

We insert Eq. (S57) and Eq. (S58) into Eq. (S55) to calculate the selection coefficient at invasion with small mutant frequency $x \ll 1$:

$$\sigma_{\text{high}} = \frac{\Delta g^{\max}}{g_{\text{wt}}^{\max}} d \tag{S59a}$$

$$\sigma_{\text{low}} = -\frac{-\Delta K}{-d\Delta K + K_{\text{mut}} g_{\text{wt}}^{\max}} \frac{g_{\text{mut}}^{\max}}{g_{\text{wt}}^{\max}} \tag{S59b}$$
$$\cdot (g_{\text{wt}}^{\max} - d)d.$$

Note that if we express this selection coefficient in terms of the relative mutation effect $\Delta K / K_{\text{wt}}$, then the selection coefficient is independent of the wild-type trait K_{wt} (compare to Fig. S19 for batch culture, where the selection coefficient increases with K_{wt} for fixed relative mutation effect). This has been observed independently in calculations by Dykhuizen et al. [33], who similarly decompose the growth rate difference in chemostats. As in the case of batch dynamics, the chemostat selection coefficient in Eq. (S59) can also be rewritten in terms of the specific affinity $a = g^{\max}/K$ instead of the resource threshold K.

S10. DEPENDENCE OF SELECTION ON RESOURCE CONCENTRATION

In this section, we use the explicit formula for s in batch culture (Eq. (S47)) to describe how selection varies with the initial resource concentration R_0 of the growth cycle. For fixed initial biomass N_0, there is an optimum concentration that maximizes selection on the resource threshold K. Figure S16A shows non-monotonic behavior of s_{low} with initial resource concentration R_0 for an example mutation with beneficial effects on both the maximum growth rate $g^{\max}$ and the resource threshold K. In particular, this optimum does not rely on a tradeoff between the two traits. Instead, Fig. S16B demonstrates that s_{low} is the product of two opposing forces: the overall budget for selection in the growth cycle (equivalent to number of generations) increases with

R_0, but the relative selection pressure on the resource threshold decreases. We can identify these two factors from Eq. (S47c) for s_{low} on the resource threshold: the selection coefficient is the product of a trait term

$$s_{\text{low}} \propto -\frac{\Delta K}{R_0 + N_0/\bar{Y} + Z} \frac{g_{\text{wt}}^{\max}}{\bar{g}^{\max}} \frac{g_{\text{mut}}^{\max}}{\bar{g}^{\max}}, \tag{S60}$$

which decreases (in magnitude) with R_0, and a logarithmic term

$$s_{\text{low}} \propto \log\left(\left(1 + \frac{R_0 \bar{Y}}{N_0}\right)\left(1 + \frac{R_0}{Z}\right)\right), \tag{S61}$$

which increases with R_0 via the number of generations in the growth cycle. The optimum concentration, in general, is determined by the wild-type resource threshold (compare Fig. S16C). Figure S16D shows how this causes the distribution of fitness effects to vary in width non-monotonically with the resource concentration as well; the width of this distribution is generally proportional to the speed of adaptation [34], which thus also displays a local maximum and minimum over resource concentrations.

These effects are not observed in batch dynamics with fixed-dilution factor, where selection s_{low} decreases strictly monotonically with resource concentration. The same example mutation in Fig. S17 reaches peak selection at the lowest nutrient concentration R_0. Intuitively, the fixed dilution factor D means the total budget for selection (number of generations) is independent of the initial concentration R_0 and low concentrations mean a larger fraction of time spent in deceleration, but not fewer generations.

S11. MODEL OF EVOLUTIONARY DYNAMICS UNDER STRONG-SELECTION WEAK-MUTATION

We can map the dynamics of the mutant frequency over batch growth cycles to the Wright-Fisher model of population genetics, where each batch growth cycle corresponds to a discrete time step [5, 35]. First, we assume the mutation arises only at the beginning of the growth cycle at frequency $1/N_0$, where N_0 is the bottleneck population size measured in number of cells. Let $s(x)$ be the selection coefficient for the mutant over a whole batch growth cycle, with explicit dependence on the frequency x of the mutant at the beginning of the cycle. In the limit of large population size ($N_0 \gg 1$) and weak selection ($|s(x)| \ll 1$), the fixation probability for the mutant is [36]

$$p(s) = \frac{\int_0^{1/N_0} \exp\left(-2N_0 \int_0^x s(y)\,\mathrm{d}y\right)\,\mathrm{d}x}{\int_0^1 \exp\left(-2N_0 \int_0^x s(y)\,\mathrm{d}y\right)\,\mathrm{d}x}. \tag{S62}$$

However, if the selection coefficient $s(x)$ is approximately constant over mutant frequencies x, we can simplify this to

$$p(s) = \frac{1 - e^{-2s}}{1 - e^{-2N_0 s}}. \tag{S63}$$

We briefly describe the scheme for simulating trait evolution. In general, a mutation can change both growth traits

$$g_{\text{mut}}^{\text{max}} = (1 + \gamma) \cdot g^{\text{max}}, \tag{S64}$$
$$K_{\text{mut}} = (1 + \kappa) \cdot K, \tag{S65}$$

where γ is the mutation effect on the wild-type maximum growth rate g^{max} and κ is the relative effect on the resource threshold K. Given the absence of correlation between g^{max} and K for autotrophs on phosphate, nitrate and ammonium (Figs. 3C–D, S9E) and for heterotrophs on glucose (Figs. 3F, S9G), we assume that mutations affect K independently of maximum growth rate ($\gamma = 0$). We simulate evolutionary trajectories of the resource threshold K by first randomly sampling a mutation effect κ from a uniform distribution on the interval $(-0.1, 0.1)$. We then calculate the selection coefficient of this mutation using Eq. (S47) and the fixation probability according to Eq. (S63). We randomly accept or reject the mutation according to this probability, and then the cycle repeats with a new mutation (Fig. S18). We also test the effect of frequency-dependence selection using the fixation probability of Eq. (S62), but Fig. S21D-F shows that it does not noticeably affect evolution of the resource threshold.

S12.   DERIVATION OF SELECTION-DRIFT BALANCE CONDITION

In the limit of weak selection ($s \ll 1$), we can expand Eq. (S63) to leading order in s:

$$p(s) \approx \frac{1}{N_0} + \left(1 - \frac{1}{N_0}\right) s, \tag{S66}$$

where the first term captures the probability of fixation due purely to demographic fluctuations (genetic drift), while the second term captures the correction due to selection. The balance between selection and drift therefore occurs when these two contributions are approximately equal, which gives us $s \approx 1/N_0$ (Eq. (3) from the main text) under the additional assumption that N_0 is large.

Now we consider the effect of a mutation arising at some intermediate time t during a growth cycle. Since at this time there are $N_{\text{wt}}(t)$ wild-type cells, the initial frequency of the mutant is $1/N_{\text{wt}}(t)$, and the amount of remaining resources is $R(t) = R_0 - (N_{\text{wt}}(t) - N_{\text{wt}}(0))/Y_{\text{wt}}$. Therefore the frequency of the mutant at the end of this cycle is

$$x(t) = \frac{e^{s(t)}}{e^{s(t)} + N_{\text{wt}}(t) - 1}, \tag{S67}$$

where $s(t)$ is the selection coefficient for this mutant arising at time t, assuming a growth cycle that starts when the mutation arises (so we use $R(t)$ as the initial amount of resources and $N_{\text{wt}}(t)$ as the initial population size).

Let $p(t)$ be the probability that this mutant ultimately fixes. This is the probability that n mutant cells survive the transfer, multiplied by the probability those mutants fix, averaged over all possible n:

$$\begin{aligned}
p(t) &= \sum_{n=0}^{N_0} \binom{N_0}{n} (x(t))^n (1 - x(t))^{N_0 - n} \left(\frac{1 - e^{-2ns(0)}}{1 - e^{-2N_0 s(0)}}\right) \\
&\approx \sum_{n=0}^{N_0} \binom{N_0}{n} (x(t))^n (1 - x(t))^{N_0 - n} \\
&\quad \cdot \left(\frac{n}{N_0} + n\left(1 - \frac{n}{N_0}\right) s(0)\right) \\
&= x(t)\left[1 + (N_0 - 1)s(0)(1 - x(t))\right] \\
&\approx \frac{1}{N_{\text{wt}}(t)} + \left(\frac{N_0 - 1}{N_{\text{wt}}(t)}\right)\left(\frac{N_{\text{wt}}(t) - 1}{N_{\text{wt}}(t)}\right) s(0) \\
&\quad + \left(\frac{N_{\text{wt}}(t) - 1}{[N_{\text{wt}}(t)]^2}\right) s(t),
\end{aligned} \tag{S68}$$

where we have invoked the weak-selection approximation to the fixation probability (Eq. (S66)) on the second line, evaluated moments of the binomial distribution on the third line, and then expanded the frequency $x(t)$ (Eq. (S67)) to leading order in $s(t)$ on the last line. By neglecting terms that are higher-order in $1/N_{\text{wt}}(t)$ and $s(t)$, we obtain

$$p(t) \approx \frac{1}{N_{\text{wt}}(t)} + \left(\frac{N_0 - 1}{N_{\text{wt}}(t)}\right) s(0). \tag{S69}$$

Note that this only depends on the selection coefficient of the mutant starting at the beginning of the cycle; to leading order there is no dependence on the selection coefficient during that first cycle $s(t)$. If we calculate the condition for selection-drift balance as before, we obtain $s(0) \approx 1/N_0$ as before. That is, the dependence on the wild-type population size at which the mutant first arises $N_{\text{wt}}(t)$ is irrelevant to the selection-drift balance. Therefore mutations arising during growth cycles have no effect on the selection-drift balance condition to leading order.

S13.   THE EVOLVED RESOURCE THRESHOLD AT SELECTION-DRIFT BALANCE

In this section, we calculate the evolved resource threshold K_{evo} as a function of environmental concen-

tration R_0 and effective population size N_{e}. We assume mutations have a maximum relative effect $|\kappa_{\max}| = |\Delta K / K_{\mathrm{wt}}|$ on the resource threshold, but no effect on maximum growth rate or biomass yield. Therefore the maximum possible selection coefficient for any mutant on the background of a wild-type trait K is thus

$$
s_{\mathrm{low}} = |\kappa_{\max}| \left(\frac{\frac{K}{R_0}}{(1 + \kappa_{\max})\frac{K}{R_0} + 1 + \frac{N_0}{R_0 Y}} \right) \tag{S70}
$$
$$
\cdot \log \left[\left(1 + \frac{R_0 Y}{N_0} \right) \left(1 + \frac{1}{(1 + \kappa_{\max})\frac{K}{R_0}} \right) \right],
$$

where we have rewritten the selection coefficient
(Eq. (S47)) in terms of the ratio K/R_0 between the
wild-type threshold and the initial resource concentra-
tion. Note that we write Y for the wild-type biomass
yield, which remains unchanged throughout evolution.

To simplify Eq. (S70), we assume that the maximum
mutation effect is small ($|\kappa_{\max}| \ll 1$), the value of
the threshold K relative the initial resource concentra-
tion is small ($K/R_0 \ll 1$), and the fold-change over
the growth cycle is large ($R_0 Y / N_0 \gg 1$). This is true
for growth cycles in typical laboratory evolution experi-
ments, with typical dilution factors between $D = 100$ [37]
and $D = 1500$ [38]. We therefore approximate the selec-
tion coefficient in Eq. (S70) by keeping only leading-order
terms in these parameters:

$$
s_{\mathrm{low}} \approx |\kappa_{\max}| \frac{K}{R_0} \log \left(\frac{R_0 Y / N_0}{K / R_0} \right). \tag{S71}
$$

The evolved threshold K_{evo} is defined as the value of
the threshold K such that the selection coefficient for
a mutation on this threshold equals the fixation prob-
ability of a neutral mutation. We must therefore also
assume that the maximum strength of selection, which
occurs for large K, is greater than the neutral fixa-
tion probability (Fig. S19A). In the limit of small $|\kappa_{\max}|$
and large $R_0 Y / N_0$, the maximum selection coefficient is
$|\kappa_{\max}| \log(R_0 Y / N_0)$, and so this must be greater than
$1/N_{\mathrm{e}}$. To solve for K_{evo}, we then set the selection coeffi-
cient in Eq. (S71) equal to $1/N_{\mathrm{e}}$ (using Eq. (3)) and solve
to obtain

$$
K_{\mathrm{evo}} \approx - \frac{R_0}{N_{\mathrm{e}}|\kappa_{\max}| W_{-1} \left(- \frac{1}{N_{\mathrm{e}}|\kappa_{\max}| R_0 Y / N_0} \right)}, \tag{S72}
$$

704 where $W_{-1}(z)$ is the -1 branch of the Lambert W func-
705 tion, defined as the solution of the equation $y e^y = z$ for
706 $-e^{-1} \le z < 0$ [39]. The latter condition is met since the
707 argument of the W function, $-1/(N_{\mathrm{e}}|\kappa_{\max}| R_0 Y / N_0)$ is
708 certainly less than zero, but also

$$
- \frac{1}{N_{\mathrm{e}}|\kappa_{\max}| R_0 Y / N_0} \ge - \frac{1}{N_{\mathrm{e}}|\kappa_{\max}| e \log (R_0 Y / N_0)}
$$
$$
> - \frac{1}{e}, \tag{S73}
$$

where on the first line we have used the fact that
$e \log(R_0 Y / N_0) \le R_0 Y / N_0$ and on the second line we have
used $N_{\mathrm{e}}|\kappa_{\max}| \log(R_0 Y / N_0) > 1$ from our previous as-
sumption that the maximum strength of selection is big-
ger than genetic drift. We can further simplify Eq. (S72)
using the approximation $W_{-1}(z) \approx \log(-z)$ for $|z| \ll 1$,
which gives us Eq. (4) in the main text.

We note that this calculation does not work for the
chemostat selection coefficient (Eq. (S59)) since it does
not depend on the wild-type trait K_{wt} outside of the
relative mutation effect $\Delta K / K_{\mathrm{wt}}$. Therefore the selec-
tion coefficient does not decrease as K evolves lower, and
there is no selection-drift balance.

S14. EVOLUTION TO SELECTION-DRIFT BALANCE FOR THE SPECIFIC AFFINITY

In this section we repeat our evolutionary analysis us-
ing the specific affinity $a = g^{\max}/K$, instead of the re-
source threshold K, as the focal trait for mutation and se-
lection. First we simulate evolution in the SSWM regime,
then we predict the evolved trait from a selection-drift
balance condition and derive a scaling relationship with
resource concentration R_0 and effective population size
N_{e}. In combination with the maximum growth rate $g^{\max}$,
the specific affinity a gives an alternative parametrization
of the Monod model of growth. Equation (S52) decom-
poses the total selection coefficient s in batch culture,
where the component s_{lin} captures the trait differences
in the specific affinity $a = g^{\max}/K$.

We assume mutations have a relative effect α on the specific affinity

$$
a_{\mathrm{mut}} = a \cdot (1 + \alpha), \tag{S74}
$$

but leave the maximum growth rate $g^{\max}$ and biomass
yield Y unchanged. The effect size α is sampled at ran-
dom from a uniform distribution, with maximum value
$\alpha_{\max} > 0$. This means a single mutation can increase
the specific affinity at most by a fixed percentage $\alpha_{\max}$.
This set of assumptions mirrors the evolutionary sim-
ulations carried out for the resource threshold K. We
simulate the trait evolution over long times, where each
new mutation either fixes or goes extinct before the next
mutation arises.

Figure S22 shows that evolution of the specific affin-
ity $a = g^{\max}/K$ leads to behavior that is analogous to
when mutations target the resource threshold K: the
specific affinity a evolves upwards over successive muta-
tions, improving the growth rate at low concentration,

but eventually the trait a stalls in adaptation around an upper limit. The limiting value depends on the effective population size N_e between transfers (compare panels in Fig. S22). Following the same reasoning as in Sec. S13, we define the evolved trait a_{evo} as the trait value where selection-drift balance is achieved:

$$s_{lin} = \frac{1}{N_e}. \qquad (S75)$$

747 Figure S22 shows that the simulated trajectories are pre-
748 dicted well by Eq. (S75), which we solve numerically for
749 specific affinity a_{evo} at selection-drift balance.

We follow the same steps as in Sec. S13 to derive a similar scaling relationship for a_{evo} as a function of the resource concentration R_0 and the effective population size N_e. The maximum possible selection coefficient for any mutation on the background of a wild-type trait a is

$$s_{lin} = \alpha_{max} \left(\frac{\frac{g^{max}}{aR_0}}{(1 + \alpha_{max})\left(1 + \frac{N_0}{R_0 Y}\right) + \frac{g^{max}}{aR_0}} \right) \qquad (S76)$$
$$\cdot \log\left[\left(1 + \frac{R_0 Y}{N_0}\right)\left(1 + \frac{aR_0}{g^{max}}(1 + \alpha_{max})\right)\right],$$

where we have rewritten the selection component (Eq. (S52c)) in terms of the ratio $g^{max}/(aR_0) = K/R_0$ between the wild-type traits and the initial resource concentration. To simplify Eq. (S76), we assume that the maximum mutation effect is small ($\alpha_{max} \ll 1$), the fold-change over the growth cycle is large ($R_0 Y/N_0 \gg 1$), and the evolved value of the specific affinity a is large relative to the initial resource concentration ($g^{max}/(aR_0) \ll 1$). This last assumption is equivalent to assuming a highly-adapted resource threshold ($K/R_0 \ll 1$), just as we did in Sec. S13. We thus approximate the selection coefficient in Eq. (S76) by keeping only the leading-order terms in these parameters:

$$s_{lin} \approx \alpha_{max} \frac{g^{max}}{aR_0} \log\left(\frac{R_0 Y/N_0}{g^{max}/(aR_0)}\right). \qquad (S77)$$

The evolved specific affinity a_{evo} is defined as the value of the specific affinity such that the selection coefficient for a mutation on this trait value equals the fixation probability of a neutral mutation. Again, we must assume that the maximum strength of selection, which occurs for small a, is greater than the neutral fixation probability (Fig. S19B). In the limit of small α_{max} and large $R_0 Y/N_0$, the maximum selection coefficient is $\alpha_{max} \log(R_0 Y/N_0)$ so this must be greater than $1/N_e$. To calculate a_{evo}, we then set the selection coefficient in Eq. (S77) equal to $1/N_e$ and solve to obtain

$$a_{evo} \approx -g^{max} \frac{N_e \alpha_{max}}{R_0}$$
$$\cdot W_{-1}\left(-\frac{1}{N_e \alpha_{max} R_0 Y/N_0}\right), \qquad (S78)$$

where $W_{-1}(z)$ is the -1 branch of the Lambert W function, introduced above in Eq. (S72). Just as before, we confirm that the evolved trait a_{evo} is confined to this solution branch and use the approximation $W_{-1}(z) \approx \log(-z)$ to arrive at the final scaling relationship

$$a_{evo} \approx g^{max} \frac{N_e \alpha_{max}}{R_0} \log\left(N_e \alpha_{max} \frac{R_0 Y}{N_0}\right), \qquad (S79)$$

which is the analogous result to Eq. (4) in the main text.
How does the evolved specific affinity a_{evo} (Eq. (S78))
compare to the evolved resource threshold K_{evo}
(Eq. (S72))? They are mathematically equivalent if the
mutation effects sizes α_{max} and $|\kappa_{max}|$ are equal, which
holds in the limit where they are both small. That is, if
we express the relation $a_{mut} = a(1 + \alpha_{max})$ for the muta-
tion effect on a as $g_{mut}^{max}/K_{mut} = (g_{wt}^{max}/K_{wt})(1 + \alpha_{max})$,
and then use the fact that g^{max} is unchanged by the mu-
tation ($g_{mut}^{max} = g_{wt}^{max}$), we then get

$$K_{mut} = \frac{K_{wt}}{1 + \alpha_{max}} \qquad (S80)$$
$$\approx K_{wt}(1 - \alpha_{max}),$$

which, compared with the definition of $\kappa = (K_{mut} -$
$K_{wt})/K_{wt}$, shows that $\alpha_{max} = |\kappa_{max}|$ when both are
small.

Altogether this shows that focusing on specific affinity
a leads to equivalent evolutionary outcomes as focusing
on the threshold K, including the dependence on the re-
source concentration R_0 and the mode of population dy-
namics (fixed-bottleneck or fixed-dilution batch dynam-
ics, or chemostat dynamics). This makes sense since mu-
tations that affect a but leave g^{max} constant must there-
fore only affect K, and thus the only difference between
these approaches is the choice of mathematical param-
eterization. We can also speculate what would happen
if mutations affect both the maximum growth rate g^{max}
and the specific affinity a simultaneously (but assuming
no correlation in effects). We expect that the maximum
growth rate will evolve to the highest physiologically-
feasible value, which will serve as the effective maximum
growth rate to convert between a and K. Intuitively, this
would still lead to identical selection-drift balance for the
resource threshold K and the specific affinity a.

S15. EFFECT OF EVOLVED RESOURCE
THRESHOLD VALUES ON MEASUREMENT
APPROACHES

In the main text we present a survey of empirical val-
ues for the resource threshold K, as well as an evolution-
ary model suggesting that K should generally be much
smaller than the concentration of the corresponding re-
source in the evolutionary environment. Here we explore
what these values of K mean for three approaches to
measuring K under laboratory conditions.

A. Inferring thresholds under chemostat growth

Arguably the most direct approach to measuring K is to use a chemostat (Sec. S5). This setup takes an inverse approach to the Monod model relation in Eq. (1): instead of varying the resource concentration R and measuring the growth rate g, as suggested by the functional form of the model, we vary the growth rate (by controlling the dilution rate d, which must equal the growth rate g in steady state) and measure the resource concentration R. We first identify the maximum growth rate $g^{\max}$ by gradually increasing the dilution rate d until the population collapses; the maximum dilution rate that the population can sustain equals the maximum growth rate $g^{\max}$. Then we set the dilution rate to half the maximum growth rate ($d = g^{\max}/2$) and measure the resource concentration at this state, which by definition of the Monod model (Eq. (1)) must equal the threshold concentration K.

In light of what we know about typical values of the threshold K, what challenges does this pose for such measurements? We must either directly measure resource concentrations in the medium around the value K (which may be difficult depending on the sensitivity of such a measurement), or infer the resource concentration from the biomass concentration $N^* = (R_{\text{source}} - K)Y$ (Eq. (S29)) In the latter case, we would also need to know the source concentration R_{source} we are supplying to the culture as well as the yield Y. However, we are not limited by low biomass concentrations in the chemostat, as we can arbitrarily increase the biomass concentration by increasing the source concentration R_{source}. For example, for *E. coli* on glucose, the threshold is $K \sim 10\,\mu\text{M}$ (Fig. 2B), the yield is $Y = 3.3 \times 10^8$ cells/μmol [30], and a typical laboratory concentration of glucose to provide could be $R_{\text{source}} = 11000\,\mu\text{M}$ (0.2% w/v). In this case the concentration of *E. coli* would be 3.6×10^9 cells/mL, which is high enough to easily measure through different standard techniques. For example, this cell density corresponds to an optical density (OD) of approximately 3.6 (using 1 OD = 10^9 cells/mL, for wavelengths of 600 nm and a path length of 1 cm), which is easily measured in a standard spectrophotometer.

B. Inferring thresholds under batch growth using the initial growth rate

A second approach uses cultures under batch growth. This takes a direct approach to the Monod model compared to the chemostat: we vary the initial concentration of the resource over some range around the concentration K and measure the initial growth rate of the biomass as a function of these concentrations. We then fit this data to the Monod model (Eq. (1)) and infer the concentration K. Note that this assumes that the population can rapidly adjust its growth rate to the external resource concentration, so that the measurement is not biased by the previous state of the culture (e.g., under starvation).

Therefore we need to perform this experiment with initial resource concentrations R_0 that are around the value of K. The total biomass concentration at the end of such a batch growth cycle would be $KY + N_0$, where N_0 is the initial biomass concentration. Using the previous example of *E. coli* on glucose, the biomass concentration KY is approximately 3.3×10^6 cells/mL, which corresponds to an OD of 3.3×10^{-3}. However, to measure growth, we must start at a concentration at least 10–100 times lower than this to have a sufficiently large dynamic range of the biomass to accurately measure the growth rate. This range of concentrations is too low to be detected on typical spectrophotometers, which usually have a lower limit of 10^{-3}–10^{-2} OD, so only methods with greater sensitivity to low concentrations (e.g., colony counting on plates or flow cytometry) would be suitable. In this case, note that the difficulty with measuring K this way is not due to its magnitude relative to a typical glucose concentration R, but that the biomass produced by this resource concentration (KY) is low compared to the lower limit of typical detection methods.

C. Inferring thresholds under batch dynamics using the deceleration into starvation

The third approach also uses batch cultures, but instead of considering how the initial growth rate varies with initial resource concentration, we use a fixed initial resource concentration R_0 and infer K from how growth rate spontaneously decelerates into starvation at the end of the growth cycle. Equation S11 defines the ODEs for batch growth with a wild-type and mutant strain. If we simplify this to a single strain, insert the Monod model for growth rate (Eq. (1)), and integrate the resource consumption equation (to express resource $R(t)$ in terms of biomass $N(t)$, as in Eq. (S12)), we obtain a single nonlinear ODE for the biomass concentration:

$$\frac{d}{dt}N(t) = g^{\max}\frac{R_0 - N(t)/Y}{R_0 - N(t)/Y + K}N(t). \qquad (S81)$$

In principle we can fit this ODE to time-series data for the biomass concentration $N(t)$ (the growth curve) and infer the threshold K.

Intuitively, though, this only works if the growth curve has enough data during the deceleration phase of growth where the threshold K is relevant; see Fig. S23 for a schematic example. Previous work has studied this as a problem of statistical estimation, calculating parameter sensitivities to identify the optimum measurement concentration and discussing variable transformations to simplify the regression (see Robinson [40] for an overview). The basic conclusion is that the initial resource concentration R_0 must be near the value of the threshold K itself for the fit to work robustly.

We can justify the intuition for this conclusion as follows. If the initial resource concentration R_0 is instead much greater than the threshold K, then the fold-change during deceleration will be too small to provide sufficient dynamic range for a fit. That is, deceleration approximately begins at the time t_{decel} when $R(t_{\text{decel}}) = K$, so that the biomass concentration is $N(t_{\text{decel}}) = N_0 + (R_0 - K)Y$. Since the final biomass concentration at saturation is $N(t_{\text{sat}}) = N_0 + R_0 Y$, the fold-change during deceleration is therefore

$$\frac{N(t_{\text{sat}})}{N(t_{\text{decel}})} = \frac{N_0 + R_0 Y}{N_0 + (R_0 - K)Y}$$
$$= 1 + \frac{\frac{K}{R_0}}{1 - \frac{K}{R_0} + \frac{N_0}{R_0 Y}}. \qquad (S82)$$

However, if R_0 is much larger than K, then this fold-change is approximately $1 + K/R_0$, meaning that it is very close to 1. Visually, this appears as a growth curve with an abrupt transition from the maximum growth rate g^{max} to zero growth (inset of Fig. 4). Since typical concentrations of many resources (such as glucose) used in the laboratory are indeed much larger than the K thresholds, this is why these growth curves usually do not contain useful data on the threshold K.

On the other hand, if R_0 is much less than K, then the growth dynamics are approximately logistic:

$$\frac{d}{dt}N(t) \approx g^{\text{max}} \frac{R_0}{K} N(t) \left(1 - \frac{N(t)}{R_0 Y}\right), \qquad (S83)$$

which we obtain similarly with Eq. (S81) but in the limit $R_0 \ll K$. In this case, one can only infer the combined parameter $g^{\text{max}} R_0/K$ from the growth curve and not the threshold K by itself. Therefore the threshold K can only be inferred from the growth curve if the initial concentration R_0 is around the value of K itself. However, this is the same parameter regime as needed for the previous method of inferring K from the initial growth rates, and thus it poses the same practical challenges, such as sensitivity to very low biomass concentrations.

[1] R. V. O'Neill, D. L. DeAngelis, J. J. Pastor, B. J. Jackson, and W. M. Post. Multiple nutrient limitations in ecological models. *Ecol Modelling*, 46:147–163, 1989.

[2] M. Zinn, B. Witholt, and T. Egli. Dual nutrient limited growth: models, experimental observations, and applications. *J Biotechnol*, 113:263–279, 2004.

[3] Mak A. Saito, Tyler J. Goepfert, and Jason T. Ritt. Some thoughts on the concept of colimitation: Three definitions and the importance of bioavailability. *Limnology and Oceanography*, 53(1):276–290, 2008.

[4] E. Sperfeld, D. Martin-Creuzburg, , and A. Wacker. Multiple resource limitation theory applied to herbivorous consumers: Liebig's minimum rule vs. interactive colimitation. *Ecol Lett*, 15:142–150, 2012.

[5] Michael Manhart, Bharat V. Adkar, and Eugene I. Shakhnovich. Trade-offs between microbial growth phases lead to frequency-dependent and non-transitive selection. *Proceedings of the Royal Society B: Biological Sciences*, 285(1872), 2018.

[6] M. Manhart and E. I. Shakhnovich. Growth tradeoffs produce complex microbial communities on a single limiting resource. *Nat Commun*, 9:3214, 2018.

[7] Jie Lin, Michael Manhart, and Ariel Amir. Evolution of Microbial Growth Traits Under Serial Dilution. *Genetics*, 215(3), 2020.

[8] R. H. MacArthur. Species packing and competitive equilibria for many species. *Theor Pop Biol*, 1:1–11, 1970.

[9] F. F. Blackman. Optima and limiting factors. *Annals of Botany*, 19:281–295, 1905.

[10] J. R. Casey and M. J. Follows. A steady-state model of microbial acclimation to substrate limitation. *PLoS Comput Biol*, 16:e1008140, 2020.

[11] C. S. Holling. Some characteristics of simple types of predation and parasitism i. *The Canadian Entomologist*, 91:358–369, 1959.

[12] J. Monod. The growth of bacterial cultures. *Annu Rev Microbiol*, 3:371–394, 1949.

[13] M. R. Droop. Some thoughts on nutrient limitation in algae. *Journal of Phycology*, 9:264–272, 1973.

[14] U. Sommer. A comparison of the droop and the monod models of nutrient limited growth applied to natural populations of phytoplankton. *Funct Ecol*, 5:535–544, 1991.

[15] W. G. Sunda, K. W. Shertzer, and D. R. Hardison. Ammonium uptake and growth models in marine diatoms: Monod and Droop revisited. *Mar Ecol Prog Ser*, 386:29–41, 2009.

[16] H. Wang, P. V. Garcia, S. Ahmed, and C. M. Heggerud. Mathematical comparison and empirical review of the Monod and Droop forms for resource-based population dynamics. *Ecol Model*, 466:109887, 2022.

[17] L. Von Bertalanffy. Quantitative laws in metabolism and growth. *Q Rev Biol*, 32:217–231, 1957.

[18] H. Moser. *The Dynamics of Bacterial Populations Maintained in the Chemostat*. Carnegie Institution of Washington, 1958.

[19] S. F. M. Hart, D. Skelding, A. J. Waite, J. C. Burton, and W. Shou. High-throughput quantification of microbial birth and death dynamics using fluorescence microscopy. *Quant Biol*, 7:69–81, 2019.

[20] Frank M. Stewart and Bruce R. Levin. Partitioning of Resources and the Outcome of Interspecific Competition: A Model and Some General Considerations. *The American Naturalist*, 107(954):171–198, 1973.

[21] Meike T. Wortel. Evolutionary coexistence in a fluctuating environment by specialization on resource level. *BioRxiv*, preprint:2021.05.18.444718, 2021.

[22] S. J. Schink, E. Biselli, C. Ammar, and U. Gerland. Death rate of E. coli during starvation is set by maintenance cost and biomass recycling. *Cell Syst*, 9:64–73, 2019.

[23] P. van Bodegom. Microbial maintenance: A critical review on its quantification. *Microb Ecol*, 53:513–523, 2007.

[24] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020.

[25] Nicolai S. Panikov. *Microbial Growth Kinetics*. London: Chapman & Hall, 1995.

[26] D. Tilman. *Resource competition and community structure*. Princeton University Press, Princeton, NJ, 1982.

[27] Joey R. Bernhardt, Pavel Kratina, Aaron Louis Pereira, Manu Tamminen, Mridul K. Thomas, and Anita Narwani. The evolution of competitive ability for essential resources. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375(1798):20190247, 2020.

[28] H W Jannasch. Growth characteristics of heterotrophic bacteria in seawater. *Journal of Bacteriology*, 95(2):722–723, 1968.

[29] Jeanne S. Poindexer. *Oligotropy*, pages 63–89. Springer US, 1981.

[30] Farida Vasi, Michael Travisano, and Richard E. Lenski. Long-Term Experimental Evolution in Escherichia coli. II. Changes in Life-History Traits During Adaptation to a Seasonal Environment. *The American Naturalist*, 144(3):432–456, 1994.

[31] E. Atolia, S. Cesar, H. A. Arjes, M. Rajendram, H. Shi, B. D. Knapp, S. Khare, A. Aranda-Díaz, R. E. Lenski, and K. C. Huang. Environmental and physiological factors affecting high-throughput measurements of bacterial growth. *mBio*, 11:e01378–20, 2020.

[32] S. Sharma and R. Steuer. Modelling microbial communities using biochemical resource allocation analysis. *J R Soc Inter*, 16:20190474, 2019.

[33] Daniel Dykhuizen and Daniel Hartl. Evolution of competitive ability in escherichia coli. *Evolution*, 35:581, 1981.

[34] M. M. Desai, D. S. Fisher, and A. W. Murray. The speed of evolution and maintenance of variation in asexual populations. *Curr Biol*, 17:385–394, 2007.

[35] J. F. Crow and M. Kimura. *An Introduction to Population Genetics Theory*. Harper and Row, New York, 1970.

[36] Motoo Kimura. On the Probability of Fixation of Mutant Genes in a Population. *Genetics*, 47(6):713–719, 1962.

[37] R. E. Lenski, M. R. Rose, S. C. Simpson, and S. C. Tadler. Long-term experimental evolution in Escherichia coli. I. Adaptation and divergence during 2,000 generations. *Am Nat*, 138:1315–1341, 1991.

[38] Nittay Meroz, Nesli Tovi, Yael Sorokin, and Jonathan Friedman. Community composition of microbial microcosms follows simple assembly rules at evolutionary timescales. *Nature Communications*, 12, 2021.

[39] E. W. Weisstein. Lambert W-function. From MathWorld–A Wolfram Web Resource. `https://mathworld.wolfram.com/LambertW-Function.html`, 2022. Accessed 2022-09-03.

[40] Joseph A.Robinson. Determining microbial kinetic parameters using nonlinear regression analysis. *Advances in Microbial Ecology*, 8:61–114, 1985.