Nucleus accumbens D1- and D2-expressing neurons control the balance between feeding and activity-mediated energy expenditure

Roman Walle [1,*], Anna Petitbon [1], Giulia R. Fois [2], Christophe Varin[3], Enrica Montalban [1], Lola Hardt [1], Andrea Contini [1], Maria Florencia Angelo [1], Mylène Potier [1,4], Rodrigue Ortole [1], Asma Oummadi [1], Véronique De Smedt-Peyrusse [1], Roger A Adan [5], Bruno Giros [6,7], Francis Chaouloff [8], Guillaume Ferreira [1], Alban de Kerchove d'Exaerde[3], Fabien Ducrocq [1], François Georges [2] and Pierre Trifilieff [1,*]

[1] Université de Bordeaux, INRAE, Bordeaux INP, NutriNeuro, 33000, Bordeaux, France
[2] Univ. Bordeaux, CNRS, IMN, UMR5293 F-33000 Bordeaux
[3] Laboratory of Neurophysiology, ULB Neuroscience Institute, WELBIO, Université Libre de Bruxelles (ULB), Brussels, Belgium
[4] Bordeaux Sciences Agro, F-33175 Gradignan, France
[5] Department of Translational Neuroscience, UMC Utrecht Brain Center, University Medical Center Utrecht, Heidelberglaan 100, 3584CG Utrechtthe Netherlands; Altrecht Eating Disorders Rintveld, Zeist, the Netherlands.
[6] Department of Psychiatry, Douglas Hospital, McGill University; Montreal, QC, Canada.
[7] Université de Paris Cité, INCC UMR 8002, CNRS ; F-75006 Paris, France.
[8] Endocannabinoids and NeuroAdaptation, NeuroCentre INSERM U1215, 33077 Bordeaux, France; Université de Bordeaux, 33077 Bordeaux, France
*Correspondence to Pierre Trifilieff (pierre.trifilieff@inrae.fr) or Roman Walle (roman.walle@live.fr)

Contact information: Pierre Trifilieff (Lead contact) or Roman Walle
INRAE, UMR 1286, Laboratoire NutriNeuro, Université de Bordeaux, Batiment UFR Pharmacie - 2eme Tranche - 2eme Etage – CC34, 146 rue Léo Saignat, 33076 BORDEAUX Cedex - France.
pierre.trifilieff@inrae.fr; roman.walle@live.fr
Phone: +33557571248

ABSTRACT

Accumulating evidence points to dysregulations of common brain systems in eating disorders (ED), the Nucleus Accumbens (NAc) in particular. However, to which extent alterations in NAc activity contribute to ED-like dimensions remains unclear. Using chemogenetic manipulations of dopamine D1 receptor-expressing and D2-expressing neuronal subpopulations of the NAc in male mice, we assessed their respective roles in incentive processes, food intake and the balance between feeding and exercise-mediated energy expenditure. We found that D1-neurons facilitate effort for a food reward but decreased food intake, while D2-neurons have opposite effects. The anorexigenic influence of D1-neurons was accompanied by increased voluntary exercise while the orexigenic effect of D2-neurons was concomitant with decreased running, congruent with D2-neurons being more active than D1-neurons during feeding while it is the opposite during running. Chronic manipulations of each subpopulations had limited effects on energy balance. However, repeated activation of D1-neurons combined with inhibition of D2-neurons biased behavior toward activity-related energy expenditure leading to weight/fat loss, whilst the opposite manipulations favored energy intake and hence weight/fat gain. Strikingly, concomitant activation of D1-neurons and inhibition of D2-neurons precipitated weight loss in anorexia models. These results suggest that dysregulations of NAc dopaminoceptive neurons might be at the core of EDs.

Keywords: Eating disorders, energy balance, motivation, nucleus accumbens, chemogenetics, calcium imaging

INTRODUCTION

Overeating/obesity-related pathologies and anorexia nervosa (AN) are multifactorial diseases resulting from an interaction between genetic and environmental factors. These eating disorders (ED) are characterized by psychiatric symptoms coupled to metabolic alterations, making them increasingly viewed as "metabo-psychiatric" diseases[1,2]. Genome-wide association studies (GWAS) reveal an inverse genetic correlation between AN and obesity[3], as well as overlapping environmental factors[4]. Dimensional approaches through the Research Domain Criteria (RDoC) framework to EDs reveals alterations in common domains such as negative (stress, anxiety) and positive (motivation, reward processing) valence systems[5]. In particular, beyond the opposite feeding patterns, which might be related to inverse alterations of food-related reward valuation[5], reward processing tends to be dysregulated in opposite manners in overeating and AN. In particular, an increased drive to exercise due to enhanced reinforcing value of physical activity is a typical feature of AN that persists after remission and might even precede the development of the disease[6–8]. On the other hand, obesity is associated with physical hypoactivity even after recovery[9]. These differential alterations of physical activity in EDs have been proposed to be linked to the rewarding impact of exercise and/or the motivation to engage in physical activity[10,11]. Such overlapping symptomatic dimensions in EDs suggest opposite dysfunctions within common brain networks, especially the reward system.

In regard to its well-established role in reward processing, motivation and locomotor activity, the nucleus accumbens (NAc) has received particular attention as it stands as a hub within the reward system, computing sensory, cognitive and reward information to modulate goal-directed behaviors[12]. Functional imaging reveals that NAc activity is regulated in opposite manner in AN and obese subjects in response to food cues[13]. Hence, this suggests that the reward system is hyperresponsive in AN while hyporesponsive in obese subjects[14], at least in relation to food-related stimuli. In the same vein, studies aiming at addressing the integrity of dopamine transmission found decreased and increased striatal dopaminergic D2/3 receptor binding in obese subjects[15,16] and in AN individuals[17], respectively, even though these latter imaging data in obesity and AN are controversial[18–20].

Altogether, these findings point to dopamine-dependent transmission of the NAc playing a key role in the pathophysiology of EDs. In the NAc, most neurons are accounted for by dopaminoceptive medium spiny neurons (MSNs) which are equally divided in two subpopulations expressing either D1 (D1-MSNs) or D2 (D2-MSNs) receptors. The respective roles of D1- and D2-MSNs of the NAc have been the subject of extensive investigation for their implication in food-related reward processing and motivational processes[21–27]. Recent studies

also highlight a direct role of both MSN subpopulations of the NAc in the regulation of food intake. In particular, D1-MSNs of the NAc shell have been shown to negatively regulate feeding[28–31], although chronic blockade of their synaptic outputs prevented the development of obesity[32]. On the other hand, striatal overexpression of the D2 receptor restricted to the developmental period facilitates weight gain due to decreased energy expenditure[33], while downregulation of striatal D2 receptor at adulthood leads to compulsion towards palatable food[34]. Conversely, its overexpression in the NAc induces rapid weight loss in female mice exposed to the so-called Activity-Based Anorexia (ABA) model[35]. Interestingly, such effects of manipulations of D2 receptor expression mainly result from physical activity-related energy expenditure rather than from a direct effect on food intake[35–37]. However, how MSN subpopulations act in concert to regulate the balance between energy intake and activity-mediated expenditure and to which extent dysregulation of MSN network could be sufficient to mimic specific obesity- or AN-related dimensions remain unknown.

To solve this issue, we used imaging to characterize the signature of activity of D1- and D2-neurons during feeding and voluntary running. We assessed the impact of acute or chronic manipulation of the activity of D1- and D2-neurons on incentive processes as well as the balance between feeding and voluntary activity-mediated energy expenditure using a chemogenetic approach. The rational for using such long-lasting manipulation was to mimic to some extent the changes in resting state functional connectivity of the NAc described in obesity[38,39] and AN[40,41]. We provide evidence that the balance between D1- and D2-neurons of the NAc differentially determines the decision to engage in food seeking, feeding or voluntary exercise. Our data point to the NAc being at the crossroad between reward processing and energy balance, further supporting that dysfunction of that network could be a main player in ED etiology.

RESULTS

D1-neurons and D2-neurons of the NAc respectively facilitate and diminish incentive processes towards palatable food

Alterations in food reward valuation are main features of EDs[5]. Because the respective roles of NAc D1- and D2-MSNs in food-oriented incentive processes was recently reconsidered[23–27], we aimed at assessing the consequence of long-lasting manipulation of either D1- or D2-expressing neurons – as a proxy for pathological dysregulation of these neuronal subpopulations - in motivation to exert effort to obtain a food reward. Activating or inhibitory DREADDs were expressed in either D1- or D2-expressing neurons of the NAc core (see supplementary material) through stereotaxic injections in D1- or D2-cre mice (Figure 1A). Of note, as previously described[22,42], we found very little viral-mediated expression of cre-

dependent DREADD, if any, in ChAT-positive neurons (SF1A), supporting that cholinergic
interneurons were mainly spared by DREADD manipulations. In accordance with the literature,
D2-neurons exclusively project to the ventral pallidum (VP), while D1-neurons of the NAc core
project to both the VP and the SNc/VTA (Figure 1 B,G).

We found that chemogenetic inhibition or activation of D2-neurons increased and decreased
performance in a progressive ratio task, respectively (Figure 1C-E; SF1B). There was no effect
on licking patterns during consumption of a palatable sweetened milk solution (SF1C) - a proxy
for hedonic reactivity -, nor on motoric abilities (Figure 2F, SF1D). However, locomotor activity
was increased and decreased by D2-neurons inhibition and activation, respectively (SF1E).
Regarding D1-neurons, both chemogenetic inhibition and activation decreased performance
in the progressive ratio task (SF1F-G). This was accompanied by an inability to increase press
rate as a function of ratio requirement in the PRx2 task for the DREADD Gq group, together
with increased locomotion, but no change in hedonic reactivity (SF1G, I, J, K). Decreasing the
dose of CNO recapitulated the effects in PRx2, though sparing the ability to increased press
rate as a function of ratio requirement (Figure 2H-K). Use of the DREADD ligand JHU37160
(J60) led to the same findings, confirming that this was not related to a non-specific CNO effect
(SF2H).

Considering the previously described satietogenic effect of NAc shell D1-neurons activation,
we hypothesized that the diminution of lever pressing could be related to a decreased hunger
drive rather than blunted motivational performance *per se*.

158 NAc D1-neurons and D2-neurons exert opposite effects on food consumption

We therefore tested the effect of chemogenetic manipulations of NAc neuronal subpopulations
on food intake. In the absence of effort required – i.e. when the reward was delivered in a
pavlovian conditioning setting (Figure 2A) –, D2-neurons inhibition in food-deprived animals
decreased consumption of the sweetened milk reward, while their activation had the opposite
effect (Figure 2B and SF2A). The mirroring pattern was observed under manipulations of D1-
neurons (Figure 2B and SF2A). To confirm the effect on food intake *per se*, we assessed the
effects of modulating NAc subpopulations activity on *ad libitum* food consumption (SF2B).
Inhibition of D2-neurons acutely decreased chow consumption, hence resulting in lower
cumulative food intake over 48 hours (SF2B). D2-neurons activation had the opposite effect,
though the increased chow consumption was delayed (SF2B). We interpret this latter finding
as the result of an initial overshadowing of the orexigenic effect of D2-neurons activation by
the blunted motivational drive, decreasing the willingness to exert effort to retrieve the food
pellets despite the increased hunger drive. This is supported by the observation that in the
case of easily-accessible liquid food (milk), under D2-neurons activation, food consumption

immediately increases as animals mostly stay at the dipper to consume (Figure 2B and SF2A).
Such effects on food intake resulted in a significant weight loss for D2-neurons inhibition and
weight gain for D2-neurons activation, these being maintained for 48 hours (SF2B). Of note,
these effects were not related to differences in peripheral glucose tolerance (SF2D).
Manipulation of D1-neurons activity resulted in opposite findings with their inhibition leading to
acute increased food intake whilst their activation strongly decreased it (SF2B), with no effect
on glucose metabolism (SF2D). However, these acute effects on food intake were not
accompanied by significant changes in weight (SF2B).

As food intake and willingness to work for a food reward were regulated in opposite manners
by D1- and D2-neuron manipulations, we next aimed at determining whether an increase in
the motivational valence of food might lead to different outcomes on food intake. We therefore
challenged mice in a fasting/refeeding protocol wherein animals were allowed access to
palatable food (high-fat) after an overnight fasting (SF2C). We found effects comparable to
those measured during *ad libitum* chow intake, i.e. D2-neurons inhibition decreased food
intake while their activation had a delayed orexigenic effect (SF2C). Manipulation of D1-
neurons triggered the opposite effects (SF2C). Both high (2 mg/kg) and low (0.5 mg/kg) doses
of CNO resulted in similar outcomes, with milder effects at 0.5 mg/kg. We therefore kept CNO
at the dose of 2 mg/kg for the following experiments.

These opposite effects of D1 and D2-neurons manipulations on food intake made us
hypothesize that each neuronal subpopulation might display opposite pattern of activity during
food consumption. Using GCaMP-mediated calcium imaging coupled with fiber photometry in
a pavlovian conditioning paradigm (Figure 2A), we found that, in trained animals, while the
relative bulk activity of D1- and D2-neurons was comparable in response to the reward-
predicting cues (Figure 2C), D2-neurons were significantly more active than D1-neurons – that
even tended to be inhibited – when animals engaged in reward consumption (Figure 2D).
Together with our chemogenetic experiments, these findings support that the balance towards
higher activity of D2-neurons over D1-neurons promotes feeding. However, reward prediction
seems to be comparably encoded by these two neuronal subpopulations, suggesting that
alterations in feeding as a result of chemogenetic manipulations of D1- and D2-neurons do not
originate from a perturbation of associative processes between the reward and predicting cues.
To further explore this hypothesis, we assessed the effect of chemogenetic activation of D1-
or D2-neurons on mesoaccumbens dopamine dynamics given the role of this pathway in
associative learning. We first confirmed through electrophysiological recordings that activation
of D1- and D2-neurons exert opposite effect on the firing of dopamine cells in the VTA, namely
inhibition and activation, respectively (SF2E). We next assessed the impact on
mesoaccumbens dopamine dynamics in a pavlovian conditioning paradigm, through

expression of the dopamine sensor dLight1.3 coupled with fiber photometry (Figure 2A). In trained animals, activation of either D1- or D2-neurons had no effect on dopamine response to food cues (Figure 2E, G). During reward consumption, activation of D1-neurons did not affect dopamine dynamics (Figure 2F), while activation of D2-neurons enhanced dopamine response (Figure 2H). Altogether, these latter findings further support that chemogenetic manipulations of NAc neuronal subpopulations directly impact feeding response rather than associative processes and that these effects could be partially mediated by an influence on dopamine transmission.

Opposite balance between NAc D1- and D2-neurons activities during food consumption and voluntary physical activity

As mentioned above, alterations in voluntary physical activity are typical symptoms of EDs that go in opposite directions with food consumption. We hypothesized that running and feeding would be accompanied by an opposite balance in the activity of D1- and D2-neurons. We therefore measured the relative bulk activity pattern of D1- and D2-neurons of the NAc during food consumption and wheel running through recording of calcium dynamics coupled with fiber photometry (Figure 3A-B, SF3A-C). In food- or wheel running-deprived animals, bulk activity of both D1- and D2-neuron were comparably increased when animals engaged towards either food consumption or the running wheel (Figure 3A-B, SF3A-B). Strikingly, the activity of both D1- and D2-neurons dropped during food reward consumption, with the relative activity of D2-neurons remaining significantly higher than that of D1-neurons (Figure 3A, SF3A-B). In opposite, during running, the relative activity of D1-neurons remained significantly higher than that of D2-neurons (Figure 3B, SF3A-B). Interestingly, when animals stopped running, D2-neurons activity remained higher than that of D1-neurons (SF3C). Altogether, these findings support that the balance of activity between D1- and D2-neurons of the NAc is biased towards D2-neurons during food consumption while voluntary wheel running correlates with higher activity of D1-neurons over D2-neurons.

The effects of acute chemogenetic NAc neuron manipulations on food intake occur at the expense of voluntary physical activity

Based on the aforementioned findings we then hypothesized that altering the balance between D1- and D2-neurons could bias choice towards eating or running. We performed a choice protocol in which fasted mice had free access to food and a running wheel (Figure 3C, SF3D-G). The acute satietogenic effect of D2-neurons inhibition was accompanied by increased wheel running distance, while activation of D2-neurons resulted in a strong decrease in physical activity with no significant effect on food intake (Figure 3D, SF3D-E), in accordance with the delayed effect of such manipulation on feeding (SF2B-C). In opposite, inhibition of D1-

244 neurons increased and decreased food intake and wheel running, respectively, while the
245 satietogenic effect of D1-neurons activation was accompanied by increased wheel running
246 (Figure 3E ; SF3F-G).

Chronic activation of NAc D2-neurons leads to increased fat mass related to decreased voluntary physical activity

Altogether, our findings suggest that the opposite effects of D1- and D2-neurons manipulation
on food intake do not reflect alterations of reward processing *per se*, but could rather
correspond to an imbalance between energy supply and physical activity-mediated
expenditure, which is a typical feature of EDs[43]. Therefore, we next asked whether repeatedly
manipulating D1- and D2-neurons activity – as a proxy for pathological dysregulation of these
neuronal subpopulation - could be sufficient for the development of some obesity- or AN-like
phenotypes. To do so, mice were housed individually with free access to high fat diet and a
running wheel in order to evaluate the relative preference between the two rewards. CNO was
administered every day at the onset of the dark period for 3 weeks (Figure 4A).

Chronic D2-neurons activation (D2 Gq) transiently increased food consumption as compared
to WT littermate controls treated with CNO (Figure 4B). This was accompanied by decreased
voluntary wheel running throughout the entire period of CNO activation compared to controls
(Figure 4C, SF4A). These resulted in a significantly higher gain of fat mass measured by TD-
NMR at the end of the chronic CNO period (Figure 4D, SF4B-E). Importantly, cessation of
CNO administration led to an immediate reversion of voluntary physical activity, together with
a significant decrease in food consumption (Figure 4B-C, SF4A), resulting in significant weight
and fat mass loss compared to controls (Figure 4D, SF4B-E). Despite the acute increase in
wheel running and decrease in food consumption, as previously found (Figure 3), animals
subjected to chronic inhibition of D2-neurons (D2 Gi) did not significantly differ from controls
for most of the main variables measured (Figure 4B-D, SF4A-E) but displayed significant
weight gain that might be due to a trend of higher food consumption throughout the chronic
manipulation (Figure 4B).

Enhanced voluntary wheel running under chronic activation of D1-neurons is compensated by increased food consumption

273 Despite the strong satietogenic effect of D1-neurons activation (D1 Gq) the very first day of
274 CNO administration, the higher voluntary physical activity that was maintained across chronic
275 D1-neurons activation was accompanied by higher food consumption during the entire period
276 of CNO administration compared to controls (Figure 4E-F, SF4F). As a consequence, animals
277 with chronic activation of D1-neurons did not differ from controls in terms of body weight, fat

and lean mass variation (Figure 4G, SF4G-J), suggesting that energy expenditure resulting from increased physical activity was compensated by higher energetic supply. Cessation of CNO administration progressively normalized both wheel running and food intake relative to controls, leading to significant gain in fat mass in the D1 Gq group (Figure 4G, SF4G-J).

Chronic inhibition of D1-neurons (D1 Gi) blunted wheel running, with no difference in food consumption compared to controls (Figure 4E-F, SF4F). This resulted in significant weight gain and increased fat mass as compared to the D1 Gq group (Figure 4G, SF4G-J). Interestingly, lower voluntary physical activity was maintained despite cessation of CNO administration (Figure 4E-F) with no change in body weight or fat mass (Figure 4G, SF4G-J).

Concomitant chronic manipulation of both NAc subpopulations bypasses the equilibrium between activity-related energy expenditure and energy intake

Altogether, our findings suggest that decreased energy expenditure due to lower voluntary wheel running, as observed under activation of D2-neurons or inhibition of D1-neurons, does not lead to compensatory decrease in food intake. Such an alteration of energy balance leads to fat gain, i.e. an obesity-like phenotype. This is in line with previous observation consecutive to decreased expression of the D2 receptor[36,37] – which disinhibits D2-neurons. However, enhanced running-induced energy expenditure, as we found under D1-neurons activation led to compensatory increase in food intake, thus maintaining energy balance, as reflected by the maintenance of weight and fat mass. We therefore asked whether potentiating the disequilibrium between D1- and D2-neuron activities by manipulating both subpopulations concomitantly in opposite manner, could potentiate the aforementioned phenotypes, eventually leading to bidirectional disruptions of the balance between food intake and activity-induced energy expenditure. We took advantage of a unique mouse model expressing the flippase recombinase in D1-expressing neurons[44] and crossed this mouse line with the D2-cre line, thereby allowing the concomitant manipulation of both NAc subpopulations (Figure 5A). Simultaneous activation of D1-neurons together with inhibition of D2-neurons strongly increased wheel running. However, in this case, after an acute decrease the first day of CNO administration, food consumption remained equivalent to control animals (Figure 5B-C, SF5A), leading to weight and fat loss (Figure 5D, SF5B-E). In opposite, the concomitant inhibition of D1-neurons together with activation of D2-neurons resulted in significantly lower wheel running compared to controls, together with an initial increase in food consumption over the first 3 days, that returned to control levels thereafter (Figure 5B-C, SF5A). Such a behavioral pattern resulted in weight gain and increased fat mass (Figure 5D, SF5B-E). Cessation of CNO administration normalized voluntary physical activity in both groups (Figure 5C, SF5A). Interestingly, this was accompanied by an increase of food intake (Figure 5B) and fat gain in

animals in which D1- and D2-neurons were previously activated and inhibited, respectively (Figure 5D, SF5B-E). Conversely, food intake (Figure 5B) and body weight (Figure 5D, SF5B-E) were decreased in animals in which D1- and D2-neurons were previously simultaneously inhibited and activated, respectively.

Chronic imbalance of D1- and D2-neuron activities in favor of D1-neurons leads to anorexia-like phenotype

The latter findings show that the effect of chronic concomitant manipulation of D1- and D2-neurons on voluntary activity can overcome metabolic signals related to energy intake. In particular, imbalance of activity between D1- and D2-neurons in favor of D1-neurons of the NAc lead to weight loss due to disproportional voluntary exercise with regard to food intake, a main feature of AN. Therefore, we next asked to which extend such manipulation of NAc subpopulation could lead to pathological weight loss (up to 20%). We first exposed mice to a gradual food deprivation in the presence of the running wheel. Under such conditions, all groups displayed a decrease in weight that was accelerated under concomitant activation of D1-neurons and inhibition of D2-neurons due to higher voluntary wheel running despite food restriction (Figure 6 A-B). Strikingly, in these animals, serum levels of the anorexigenic adipokine leptin where decreases while the orexigenic gut peptide ghrelin was elevated (Figure 6C), which is a hallmark of AN[45]. Nonetheless, in the absence of a running wheel, concomitant manipulations of NAc subpopulations did not prevent the ability of leptin and ghrelin to exert their respective anorexigenic and orexigenic influence despite that the effect of chemogenetic manipulations of NAc subpopulations was dominant on food intake (SF6A-B). These findings support that imbalanced activity of NAc subpopulations in favor of D1-neurons can overcome peripheral feeding signals.

To further support the translational validity of our findings, we subjected animals to the Activity-Based Anorexia (ABA) model[46]. Concomitant activation of D1-neurons together with inhibition of D2-neurons precipitated weight loss due to higher voluntary activity (Figure 6D-F, SF6C-D), leading to lower probability of "survival" (20% weight loss, Figure 6G). Concomitant inhibition of D1-neurons together with activation of D2-neurons was "protective" due to blunted voluntary running (Figure 6D-G, SF 6C-D).

DISCUSSION

Beyond the well-documented role of NAc in the modulation of incentive processes, the present study highlights an underestimated differential implication of D1- and D2-neuronal subpopulations in the balance between food intake and physical activity-induced energy expenditure. As an illustration, the use of the above-mentioned double transgenic mice reveals

that repeated (i) activation of D2-neurons together with (ii) inhibition of D1-neurons is sufficient to elicit an obesity-like phenotype as assessed by a decrease in energy expenditure. In opposite, repeated hyperactivity of D1-neurons together with hypoactivity of D2-neurons directs the energetic balance towards physical activity at the expense of energy intake, which are typical features of AN.

Even though dimensional approaches to EDs is just starting to emerge, increasing evidence supports alterations in common domains across EDs, especially those linked to reward-related positive valence system[5]. In particular, reward valuation is elevated in individuals with obesity-related binge eating syndromes, but excessively diminished in patients with AN[47,48]. Similarly, increased willingness to work for disorder-specific reward, i.e. exercise and food for AN and bulimia, respectively, are increased[49,50]. Such reward-related symptoms correlate with opposite brain activity patterns, especially in cortico-striatal circuits[13], hence reinforcing the hypothesis of shared neurobiological substrates, and therefore potential continuum in EDs. Our data suggest that opposite alterations in the activity of NAc subcircuits could underlie a subset of symptomatic dimensions of ED phenotypes, in particular the balance between calorie intake and activity-related energy expenditure.

We found that the bulk activity balance between D1- and D2-neuron subpopulations was in favor of D2-neurons during food intake while towards D1-neurons during voluntary physical exercise. In accordance, acute manipulations of the two subpopulations led to opposite consequences on food intake and voluntary physical activity in opposite manner, highlighting the implication of these neuronal subpopulations in the regulation of energy balance. However, it is important to stress that, under chronic manipulations, the homeostatic system – likely comprising hypothalamic network - seems to be able to overcome alteration of D1- and D2-neurons' activity. Thus, under chronic activation of D1-neurons, despite an acute decrease in food intake, enhanced energy expenditure through physical activity is accompanied by increased food intake, allowing maintenance of fat mass and body weight. These findings are mostly consistent with a previous study by the group of DiLeone[31]. However, our findings reveal that it is only when both NAc subpopulations are concomitantly manipulated, i.e. activation and inhibition of D1- and D2-neurons, respectively, that such an homeostatic system does not fully compensate for energy expenditure, hence leading to weight loss. Nonetheless, animals still eat comparable amount of food than controls, suggesting that the energetic imbalance mainly results from exaggerated physical activity-mediated energy expenditure rather than reduced food consumption. Similarly, when D1-neurons or D2-neurons are respectively subjected to chronic inhibition and activation, weight/fat gain mostly originates from decreased physical activity since animals consume comparable amount of food than controls. Finally, under food restriction (i.e. progressive food restriction or ABA model), we found that protection from, or

precipitation of, weight loss depending on the manipulations of MSN subpopulations, mainly resulted from alterations of voluntary activity. Yet, avoidance and compulsion towards exercise are typical features of obesity-related disorders and AN, respectively, that have been proposed to be one main pathophysiological mechanism leading to, and even preceding, the development of EDs[6,51]. Even though speculative, overall, our findings suggest that chronic dysfunction in the balance of activities of D1- and D2-neuron of the NAc could participate to the etiology of EDs by altering, positively or negatively, the willingness to engage in, and/or the rewarding value of, physical activity. These findings expand the importance of the balance of activities between striatal D1- and D2-neurons to control behavior[52,53].

Of importance is the observation that wheel-running, rather than reflecting non-specific locomotor responses, is a goal-directed behavior accounted for by the reinforcing value of exercise. Thus, as for other natural rewards such as food, rats/mice are willing to exert efforts, i.e. lever presses or nose pokes, to access a wheel[54,55]. In accordance, our findings using chemogenetic inhibition demonstrate that D1-neurons facilitate both wheel-running and willingness to exert effort to obtain a food reward, while D2-neurons rather play an opposite role. These data further suggest that alterations in voluntary physical activity under D1- and D2-neurons manipulation reflect changes in reward/incentive processes and reveal the antagonistic role of the two NAc subpopulations on this dimension. However, such an opposite effect of D1- and D2-neurons on motivational processes has been recently challenged. Similarly to our findings, the chemogenetic inhibition of D2-neurons of the NAc enhances performance in a motivational task[21,22], while optogenetic manipulations of D2-neurons of the NAc rather support a "pro-motivational" role, similarly to D1-neurons[23–25]. However, recent data clarified this discrepancy, showing that optogenetic stimulation of D2-neurons during reward-predicting cue enhances operant responding, while the very same manipulation during reward consumption has the opposite effect[27]. Such a dissociated role of D2-neurons on distinct phases of the motivational dimension remains unclear and might involve reward prediction error mechanisms through dynamic feedback modulation of VTA dopamine neurons[27]. Nonetheless, through the use of long-lasting chemogenetic approach, our results support that chronic increase or decrease in D2-neurons excitability, as putatively found in some pathological conditions, will likely lead to enhanced and blunted motivational processes, respectively.

In addition to their antagonistic effect on the "ability to exert effort", our results reveal that D1-neurons inhibit, whereas D2-neurons of the NAc enhance food intake. To the best of our knowledge, such a role of D2-neurons has not been reported before. However, even though a recent study showed that activation of D1- and D2-neurons of the NAc enhance and terminate sugar intake, respectively[56], the acute anorexigenic potency of D1-neurons has been

previously described for the NAc shell, through projections onto the lateral hypothalamus and the midbrain[28–30]. These latter findings raise the question of whether these two pathways are redundant to control food intake or modulate distinct – though complementary – dimensions of feeding behavior. It has been proposed that the NAc-VTA pathway modulate feeding through a direct control onto motor behaviors that comprise a feeding response[28,57]. Regarding NAc-LH projections that mostly originate from the shell of the NAc, GABA signalling – notably originating from NAc D1-expressing neurons – has been shown to be increased during food intake, consequently terminating feeding, notably through decrease of orexin production[58]. These findings also highlight that, beyond the core and shell subdivisions, subgroups of NAc D1-neurons likely control discrete behavioral dimensions, related to their location in subterritories of the NAc as well as their input and output projections. Finally, a main output structure of D1-expressing neurons of the NAc is the ventral pallidum (VP). Lesion of the VP – similarly to lesions of the LH - has been shown to counteract the orexigenic effect of NAc inhibition[59], suggesting a "pro-feeding" role of the VP. Accordingly, arkypallidal neurons of the VP potentiate reward consumption[60]. Even though, to the best of our knowledge, a direct implication of D1-neuron-to-VP projections in the regulation of food intake remains to be demonstrated, all these findings support that D1- neurons of the NAc can decrease food intake through several output pathways involving the LH, the VTA and the VP. Of note, even though we cannot exclude some effects related to spread of viral infections to the shell of the NAc, our chemogenetic manipulations were mostly restricted to the core subdivision, that sends projections to the VP and VTA but not the LH. The circuits by which D2-neurons of the NAc regulate feeding remain unclear. As mentioned, we cannot rule out that some of the effects we observed in this study might be related to an alteration in the activity of cholinergic interneurons (CIN) due the use of the D2-cre line - even though likely targeting at most a subset of ChAT-positive interneurons[22]. However, increasing CINs activity reduces palatable food consumption while their inhibition had the opposite effect[61], in contrary to our findings with manipulations of D2-neurons. This suggests that most of the effects we observed were due to alterations of D2-MSNs. The sole output of this neuronal subpopulation is the VP. This raises the intriguing ideas that, if the VP is involved in the effects on feeding that we observed under manipulation of NAc subpopulations, it is likely that D1- and D2-neurons should project onto distinct VP neuronal subpopulations, though, to the best of our knowledge, this is an important aspect that remains to be demonstrated. Finally, the lateral inhibition between NAc subpopulations has been largely overlooked and is likely to participate to their differential roles, including feeding behavior.

Together with the current study, these observations suggest that, under physiological conditions, the recruitment of the NAc network in favor of D1-neurons would bias behavior towards effort exertion – including food-seeking – at the expense of food consumption. In

opposite, a predominant activity of D2-neurons would blunt voluntary activity whilst promoting food intake. Such a pivotal role of the NAc neuronal network is in accordance with the implication of the NAc as a "sensory sentinel", allowing rapid control over food consumption in response to motivational or sensory signals[28,29,57,62]. It remains to be determined whether these behavioral dimensions bidirectionally modulated by D1- and D2-neurons depend on distinct subpopulations in the NAc and/or their projecting areas. Nonetheless, the current work suggests that a chronic imbalance in the NAc neuronal network could be sufficient to alter homeostatic balance, eventually leading to EDs.

MATERIALS AND METHODS

Experimental models and subject details

Adult male D1-Cre, D2-Cre, D1-Flp and D1Flp/D2cre (homemade breeding from D1Flp and D2cre mice) mice were used in this study (see table 1 for details). Mice (12 to 28 weeks old) were group housed (4-7 animals), unless otherwise specified, in standard polypropylene cages and maintained in a temperature and humidity-controlled environment under a 12:12 light-dark cycle (07:30 on) with *ad libitum* access to water and food. For food consumption experiments, mice were housed in standard polypropylene individual cages. All behavioral procedures were carried out during the light phase unless otherwise specified. Mice were habituated to handling prior to behavioral procedures, and to injection proceeding before test day. All animal care and experimental procedures were in accordance with the INRAE Quality Reference System and to French legislations (Directive87/148, Ministère de l'Agriculture et de la Pêche) and European (Directive 86/609/EEC). They followed ethical protocols approved by the Region Aquitaine Veterinary Services (Direction Départementale de la Protection des Animaux, approval ID: B33-063-920) and by the animal ethic committee of Bordeaux CEEA50. Every effort was made to minimize suffering and reduce the number of animals used.

Stereotaxic viral injections and implantation of optic fibers

Mice analgesia was achieved with the subcutaneous injection of buprenorphine (0.05 mg/kg). The animals were then anesthetized using isoflurane (4% induction, 1.5% maintenance) and placed in a stereotaxic frame onto a heating pad. The AAV8-hSyn-DIO-hM4D-mCherry virus (titer of 4.5×10^{12} vg/ml I) was used for the Cre-dependent expression of hM4D(Gi) receptor, the AAV8-hSyn-DIO-hM3D-mCherry virus (titer of 5.2×10^{12} vg/ml) was used for the Cre-dependent expression of hM3D(Gq) receptors and the AAV8-hSyn-DIO-mCherry virus (1×10^{13} vg/ml) was used for the cre dependent expression of mCherry. The AAV8-hSyn1-dFRT-mCherry(rev)-dFRT-WPRE-hGHp(A) and AAV8-hSyn1-dlox-EGFP(rev)-dlox-WPRE-hGHp(A) were used for the labelling of D1-Flp and D2-cre expressing neurons (titer of $4-5 \times 10^{12}$ vg/ml).

The AAV8-hSyn1-dFRT-hM4D(Gi)-mCherry(rev)-dFRT-WPRE-hGHp(A) and AAV8-hSyn1-dFRT-hM3D(Gq)-mCherry(rev)-dFRT-WPRE-hGHp(A), were used for the flp-dependent expression of hM3D(Gq) and hM4D(Gi)-receptors (titer of 4 x 10^{12} vg/ml). A total of 0.5µl of virus solution was injected bilaterally in the nucleus accumbens using a 10µl Hamilton syringe at a rate of 100nl/mn at the following stereotaxic coordinates relative to Bregma (in mm) : AP= +1,7 ; ML= +/- 1,1 ; DV= -4 from the surface of the brain. The syringe was left in place for 5 min before being slowly removed. For measures of calcium dynamics by fiber photometry, the AAV5-hSyn1-dlox-jGCaMP8m(rev)-dlox-WPRE-SV40p(A) viral vector was used for the cre-dependent expression of GCaMP8m (8.0 x 10^{12} vg/ml). For the measure of dopamine dynamics coupled to chemogenetic activation of D1- or D2-neurons, the AAV9-hSyn1-chl-dLight1.3b-WPRE-bGHp(A) vector was co-injected in the NAc of D1- or D2-cre mice together with the AAV8-hSyn-DIO-hM3D-mCherry vector. An optic fiber (RWD Fiber Optic Cannulae 0,5 NA, 400µm flat tip, L:6mm) was implanted just above the viral vector infusion site (DV= -3,75) and immobilized with an opaque dental cement. The skin incision was sutured, postoperative analgesia was prolonged with subcutaneous injections, and mice were place on a heating pad until full recovery from the anesthesia. Behavioral experiments were started 3 to 4 weeks after the surgeries.

Ventral tegmental area recordings

Stereotaxic surgery for electrophysiology experiments was performed under isoflurane anesthesia as previously described [63]. Recording pipettes were inserted into the VTA with the skull flat, at the following coordinates: -3.16mm from bregma; 0.5mm from midline. A glass micropipette (tip diameter=2-3µm, 4-6Mohm) filled with a 2% pontamine sky blue solution in 0.5M sodium acetate was lowered into the VTA. DA neurons were identified according to well-established electrophysiological features [64]. These included: 1) an half action potential width ≥1.1ms; 2) slow spontaneous firing rate (<10Hz); 3) single and burst spontaneous firing patterns. Through these electrodes, the extracellular potential was recorded with an Axoclamp2B amplifier in the bridge mode. The extracellular potential amplified 10 times by the Axoclamp2B amplifier was further amplified 100 times and filtered (low-pass filter at 300 Hz and high-pass filter at 0.5 kHz) via a differential AC amplifier (model 1700; A-M Systems, Carlsborg, WA). Single neuron spikes were discriminated and digital pulses were collected on-line using a laboratory interface and software (CED 1401, SPIKE 2, Cambridge Electronic Design). Four parameters of VTA DA neuron impulse activity were computed over 200-second epochs after a 5-minute stable baseline period: 1) the basal firing rate, 2) the basal bursting rate, 3) the mean number of spikes per bursts and 4) the number of spontaneously active cells per track. The onset of a burst was defined as the occurrence of two spikes with an interspike

525 interval <80msec[64]. VTA DA neurons were recorded under saline or CNO injection (2mg/kg)
526 in D1- or D2-cre mice expressing the cre-dependent hM3D(Gq) DREADDs in the NAc.

Behavioral Procedures

For most behavioral procedures, two experimenters were involved, one in charge of injections
(CNO/saline), the other performing the behavioral tasks. The latter was blind of animals'
genotype, viral vector and injection. For all behavioral experiments, mice received CNO (0.5
or 2mg/kg dissolved in sterile 0.9% saline as indicated in the text; see table 2 for details),
JHU37160 (1 mg/kg dissolved in sterile 0.9% saline), or 0,9% saline solution intraperitoneal
(IP) injections 30 min before the beginning of the test. For the experiments requiring training
sessions, there were no injections during these steps. CNO and JHU37160 solutions were
prepared daily and protected from light. For acute CNO treatment experiments, each mouse
received either CNO or saline in a counterbalanced manner for within subject analysis, across
2 sessions of each behavioral test at minimum. The same procedure was applied for
experiments using JHU37160. For experiments using WT littermate controls, all animals were
treated with CNO when indicated. Experimental designs were randomized regarding genotype,
virus type, home-cage and testing chamber. Different cohorts were used for each experiment,
except for the measure of licking microstructures that were performed in the same animals as
the ones used for operant conditioning experiments.

Operant conditioning

Mice were placed under food deprivation and maintained at 85-90% of their baseline *ad libitum*
weight. Fourteen operant chambers (30x40x36 cm, Imetronic Bordeaux), each located into a
soundproof and light-attenuating cabinet were used. They consisted of 2 plexiglas panels and
2 PVC panels on top of a metal grid floor, lit by a houselight located at the top of the chamber
during sessions. A trough was located on the center of one PVC panel, 4cm above the metal
grid, and was surrounded by 2 equally distant retractable metal levers (2x4x1cm). A cue light
was located 8.5 cm above each lever. The reinforcer was composed of a 15 µl drop of
sweetened condensed milk (3.25 Kcal/g) diluted at 10% in tap water delivered in the trough.
Both levers were extended at the beginning of each session, except for pavlovian training.
Only one lever was reinforced across all operant procedures and pressing on the non-
reinforced lever had no consequence. For each mouse, all of the operant procedures were
executed in the same operant chamber. Operant chambers were connected to a computer
equipped with POLY Software (Imetronic), which allowed for control of the sessions
parameters and recording of the data (number of presses on both levers, number of licks in
the full or empty trough).

Pavlovian Training preceding operant conditioning

Three to 4 weeks after AAV injections, animals started pavlovian training. Mice were placed in the operant chamber for 30 min sessions, one session per day, during which a reward – a drop of sweetened condensed milk – was delivered in the trough every minute, associated with the emission of a sound (65 db, 3000 Hz, 200ms) and the illumination of the diode located above the future reinforced lever. The reward had to be consumed by the animal in order for a new reward to be delivered. The learning criterion was set at a mean of 20 rewards earned out of 30, across at least 3 daily consecutive sessions.

Fixed Ratio Training (FR)

When the mice reached the pavlovian training criterion, animals started operant conditioning sessions during which they had to press on the reinforced lever to obtain the reward. Reinforced lever presses were signaled by the emission of the sound and the cue light. Operant learning started with daily 1-hour fixed ratio 1 schedule of reinforcement (FR1), during which each reinforced-lever press was rewarded. The learning criterion was set at 100 reward dose earned, across four consecutive sessions.

Random Ratio Training (RR)

Once all the mice completed the operant learning criterion, their ability to press on the reinforced lever in order to receive a reward was increased using random ratio schedules. During the sessions, each lever press on the reinforced lever had a given probability of being reinforced (pRR5=0.2, pRR10=0.1, pRR20=0.05). Animals underwent those 3 different ratio across one-hour sessions, each day, 5 days a week. Moving from one ratio to the next one was achieved once lever response values across two consecutive sessions were stable (less than 10% variation).

Progressive Ratio Task (PRx2)

The motivational component of operant conditioning was evaluated using a progressive ratio task. During the session, the first reward was delivered after two lever presses on the reinforced lever and the following required amount of lever presses doubled after every reward earned. Session was considered terminated when animals failed to lever press for three minutes, thus defining the session duration. PRx2 tasks were intercalated by RR20 sessions to prevent operant responding extinction. Mice underwent one PRx2 task without injections for basal performances. Then 4 PRx2 tasks were carried out, during which each mouse received CNO (2 mg/kg or 0.5mg/kg ; IP) or saline IP injection. Order of Saline/CNO administration was counterbalanced within groups. The 2 sessions of PRx2 under CNO and under saline were

Licking microstructures

A homemade setup was used to record licking microstructures [42,65]. The test was run in
polypropylene cages equipped with a feeding bottle and a metal grid floor connected to a
computer with a dedicated software. The animal's contact between the grid floor and the
metallic sipping tip of the bottle allowed for the precise detection of licks. Mice were habituated
to the testing chamber with bottles filled with water for 2 days through 30 min sessions. On the
test day, the bottle was filled with 10% condensed sweet milk diluted in tap water and mice
were injected with CNO (2 mg/kg) or saline solution 30 min before the beginning of the 30-min
testing session (within subject repeated-measures). The system allowed for the detection of
the total number of licks, total number of bursts and number of licks per burst. Bursts were
defined as sequences of high frequency licks, in which the interlick interval was less than
250ms. Relation between number of licks per burst and hedonic reactivity have been
demonstrated in[42,66,67].

Motor coordination in rotarod

A rotarod apparatus (Bioseb) was used to assess motor coordination under CNO (2mg/kg) and
saline (within subject repeated-measures). Mice were placed on the rod rotating at 4rpm.
During the test, the rotation of the rod increased from 4rpm to 40rpm in 300 sec. The latency
to fall was recorded.

Measure of 48-hour ad libitum food intake

Single-housed mice were food deprived for 2 hours and were injected with CNO (2 mg/kg ; IP)
30mn before the onset of the dark phase. Chow pellets (A04, SAFE) were placed in a dish into
the cage. Food dishes were weighed after 1 ; 2 ; 4 ; 12 ; 24 and 48 hours. The same procedure
was performed with animal receiving simultaneous CNO (2mg/kg; IP; within subject repeated-
measures) and leptin infusions (5mg/kg; IP) 3 hours before dark phase, or CNO (2mg/kg; IP)
and ghrelin infusions (3mg/kg; IP) at the beginning of the light phase (9.30 AM). In this latter
condition, chow pellets were placed in a dish into the cage 30 minutes after injections (10.00
AM).

Fasting Refeeding procedure

Single-housed mice previously exposed to high fat diet (Research Diet ref: D12451), were food
deprived for 12 hours, then injected with CNO (2 mg/kg or 0.5mg/kg ; IP) 30mn before the

onset of the dark phase. Following high fat diet access, food was weighed at 1 ; 2 ; 4 ; 12 ; 24 and 48 hours.

Pavlovian conditioning for the measure of food intake coupled with fiber photometry recordings

Food-deprived mice were placed in the operant chamber for 20 min sessions, during which a drop of sweetened condensed milk was delivered 3 seconds after the emission of a sound (65 db, 3000 Hz, 7s) and the illumination of a diode lasting 10 seconds (ITI : 45s after reward consumption). Animal were first trained during 10 sessions. For the measure of food intake, during the last two sessions, animals were received i.p. saline or CNO (2mg/kg) injections in a counterbalanced manner, 30 minutes before the trials. Licking patterns were analyzed as a proxy for reward consumption. The same procedure was used for the recording of calcium transients in animals expressing GCaMP8m in either D1- or D2-expressing neurons of the NAc. Implanted animals were connected to A RWD (R820 Fiber Photometry) fiber photometry setup. A 470nm LED light stimulation coupled to a 405nm isobestic stimulation was delivered through a fiber-optic patch cord (O.D. 1.25mm, Ceramic Ferrule, 400µm core diameter, 0.5 NA ; R-FC-L-N5-400-L1, RWD). Intensities were set to reach a 100µW power for 470nm LED and 40µW for 405nm isobestic LED. Activity was recorded at 40Hz. Mice were subjected to 7 sessions of 15 minutes. For recording of dopamine transients, the same behavioral paradigm and recording settings were used. The effect of CNO was assessed after extensive pavlovian training (10 sessions) over 2 sessions, intercalated with one session with saline injections (3 total with one session before and one after the CNO injections). The z-score around each behavioral event was then calculated using a baseline of 1.5sec (from -2sec to -0.5sec before the event) as $\frac{\Delta F/F - (means\ of\ values\ in\ baseline\ period)}{standard\ deviation\ of\ values\ in\ baseline\ period}$. Area Under the Curve (AUC) were calculated in MATLAB using the trapz function.

In vivo fiber photometry recording of calcium transients during voluntary wheel running or free-feeding

Animals expressing GCaMP8m in either D1- or D2-neurons of the NAc (see above) were singled-housed with *ad libitum* access to a running wheel and *ad libitum* food during 2 weeks. Two days before testing, either the running wheel was removed or animals were food restricted. The behavioral apparatus consisted in a four-chamber maze composed of two waiting areas each leading to distinct compartments containing either palatable grain-based food pellets (Dustless Precision Pellets, Bio-Serv) or a running wheel. A session consisted in recording baseline activity in the waiting area for 60 sec, then the door was opened giving access to either the running wheel or 6 food pellets distributed every 30 sec. Animals were

recorded during 3 min. 2-3 sessions per day were performed (for either the running wheel or
food consumption) during 6 consecutive days. Animals' behavior was video-recorded
synchronized with the fiber photometry recording (TTL), and a post-analysis of behavioral
epochs identified through the BORIS Software, allowing the alignment with fiber photometry
signal and a peri-event analysis.

Fiber photometry data were acquired with Doric Neuroscience Studio software through a fiber
photometry console controlling excitation lights, demodulated fluorescence signals and
received timestamps of behavioral events (Doric Lenses). Fiber photometry uses the same
fiber to both excite and record from animal via optical fiber patch cord (RWD Optic Fiber 0,5
NA, 400µm flat tip, L:1m). Blue excitation light (460 nm Doric Lenses LED) and isosbestic light
(405 nm Doric Lenses LED) passed through a filter cube (iFMC4; Doric Lenses). Light intensity
at the tip of the fiber was measured before every recording session and kept at 40 µW for
isosbestic light and at 100 µW for blue excitation light. During recording, both light stimulations
were alternatively triggered at 10hz. Then fluorescence signals were collected through a
Fluorescence Detector Amplifier (DC x1 ,Doric Lenses). Signals were sampled at 12048.19Hz.

Fiber photometry signals were analyzed using custom MATLAB (MathWorks) scripts. Signals
were demodulated according to the 460 nm and the 405 nm light emission and resampled at
10 Hz. The isosbestic signal (405 nm signal) was scaled using a least-squares linear
regression to best fit the 460 nm signal. This scaled isosbestic signal was subtracted from the
calcium-sensitive signal at 460 nm to compute a movement corrected, and bleaching-corrected
change in fluorescence (ΔF), then the ΔF/F was calculated by dividing the change in
fluorescence by the scaled isosbestic signal. To detect calcium transients, the ΔF/F signal was
bandpass filtered (4th-order Butterworth filter: lower cutoff frequency 0.1 Hz, higher cutoff
frequency of 2 Hz), and calcium activity peaks were detected when the signal exceeded a
threshold defined by the median value + 2 median absolute deviation (MAD) with a prominence
of at least one MAD and a distance between peaks of at least 0.1 s. For each behavioral event,
the photometry signals were z-scored using a pre-event baseline ranging from -40 s to -30 s
for 'feeding room entrance', 'feeding period', 'running room entrance', 'running period' events,
or from -8 s to -5 s for 'feeding', 'stop running', and 'locked wheel' events. Resulting photometry
signals were then analyzed to first determine significant event-related transients around
behavioral events in either D1- or D2-neurons. For each event, using the compilation of peri-
event z-scored photometry signals, a bootstrapping procedure was used to calculate
confidence intervals (95% CI, 2000 random resampling with replacement). A 95% CI was
calculated for each time point using the 2.5 and 97.5 percentiles of the bootstrap distribution
and was then expanded by a factor of $\sqrt{(n/(n-1)}$ to account for narrowness bias [68,69].
Significant transients were defined during periods for which the 95% CI does not contain 0

(corresponding to the pre-event baseline) for at least 1 s. Second, we compared photometry signals recorded from D1-neurons and D2-neurons around each behavioral event using permutation tests (2000 random partitions) [70]. Signals were defined as significantly different if the exact permutation-based p-value stays below the significance threshold set at 0.05 for at least 1 s. In addition, the area under curve (AUC), the average z-score, the peak z-score, and the average transient rate were calculated respectively as the sum of the z-scored photometry signal, the averaged value of the z-scored signal, the maximum value of the z-scored signal, and the average frequency of detected calcium transients. For each behavioral event, these values are calculated during a pre and a post periods as indicated in the figures.

Effect of acute manipulation of neurons on wheel-running/free-feeding

The setup consisted of a magnetic fast-track running wheel mounted onto a mouse igloo shelter (BioServ) and connected to a magnet odometer. Single-housed mice had free access to regular chow and the running wheel for 2 days for habituation. On the test day, mice were fasted for 12 hours then placed in the cage with the running wheel and *ad libitum* chow placed inside the cage for 2 hours for baseline consumption and wheel activity. CNO (2 mg/kg) or saline (within-subject repeated measures) was then injected and food intake and wheel activity were recorded every hour for 3 hours. For each condition (pre- or post-infusion), data are expressed as the difference between CNO and saline condition, divided by the value under saline condition. Animals with at least one outlier ratio value (Grubb's test) were excluded from analysis.

Effect of chronic manipulation of D1- and D2-neurons on wheel-running/free-feeding

Single-housed mice were exposed to a running wheel and standard diet for 1 week prior to the beginning of the test. 24-hour wheel running activity, food consumption and weight variation were recorded and defined as a baseline. On the first day of the test animals' food was switched to HFD and CNO (2mg/kg) was administered every day before the onset of the dark period for 23 days, during which variation of percentage of the aforementioned parameters were evaluated. Measurements were pursued for 10 days with no CNO administration. Subsequently, mice were subjected to progressive food restriction over 7 days (decrease of 0.5 g per day, starting at 3g) with a daily CNO infusion and free access to the running wheel.

Effect of concomitant manipulation of D1- and D2-neurons in the ABA model

Mice were individually housed and given free access to a running wheel, food and water for 7 days (day −7 to day 0). On day 0, 3 h into the dark phase, food was removed and mice were placed on scheduled feeding of 3 h per day starting at dark onset. CNO was injected 30 minutes before addition of the food.

MRI for body composition

The last day of chronic chemogenetic manipulation experiments, whole body composition of was measured (basal condition, after CNO treatment period and after CNO withdrawal period) using EchoMRI, minispec LF90II (Bruker). It acquires and analyzes TD-NMR signals from all protons in the entire sample volume and can provide 3 components of interest: Fat, Free Body Fluid and Lean Tissue values. Total mass and fat mass variation between each period were calculated.

Glucose Tolerance Test

Single-housed mice were food deprived 5 hours prior to the test at 09:00. Glycemia was measured using single-use test strips and glycemia meter (ACCU-CHEK Performa, Roche). The tip of the tail of the animal was cut for the first measurement, in order to obtain 1-2 drops of blood. For the following blood tests, the tail was massaged from the base to the tip so that a small volume of blood comes out. The first drop was wiped away and the second drop was used to measure glycemia. Mice were injected with CNO (2mg/kg ; IP) or saline 20 minutes prior to the beginning of the test. Glucose (2g/kg ; IP) was injected to the mice at t=0mn. Nine blood samples were collected throughout the test (-30mn before CNO/Saline injection; 0mn before glucose injection; +15mn ; +30mn ; +45mn ; +60mn ; +90mn ; +120mn ; +180mn) and mice were left quiet in a closed soundproof room in between samples.

Plasma dosage of ghrelin and leptin

At the end of the progressive food-restriction procedure, CNO was injected 90 minutes before blood samples collection in EDTA pre-coated tubes through cardiac puncture. Plasma was obtained by centrifugation (4500 rpm, 5 min at 4 °C) and then stored at –80 °C until further analysis. Plasma concentrations of acylated ghrelin and leptin were determined by commercial ELISA kits according to the manufacturers' instructions (Spi Bio #A05117.96 and Millipore #EZML-82K, respectively).

Immunohistochemistry

Mice were deeply anesthetized using a lethal IP dose of sodic pentobarbital (Exagon, 300mg/kg) together with lidocaine (Lurocaine 30m g/kg), and transcardially perfused with an ice cold 4% paraformaldehyde solution in PBS 1X. Brains were removed and immersed in 4% PFA overnight at 4°C, then transferred in cryoprotectant solution (30% glycerol ; 30% ethylene glycol in PBS-1X pH 7.4), and stored at -20°C until further use. Brains were rinsed in PBS 1X for 30 min before slicing. 40μm coronal sections were collected using a vibratome (Leica) into cryoprotectant solution and stored at -20°C until further use. Brain slices were mounted on

slides and coverslipped with VectaShield containing DAPI mounting medium (Vector Labs). mCherry expression was visualized with a Hamamatsu NANOZOOMER 2.0HT microscope (Bordeaux Imaging Center facilities). For mCherry signal amplification, brain sections were rinsed in PBS-1X, treated with blocking buffer (10% Fetal Bovine Serum in TBS: Tris 0.1M, NaCl 0.9%, pH 7.4) for 2 h at room temperature (RT) and incubated overnight at 4°C with primary antibody against mCherry (rabbit; 1:1000). For GFP signal amplification primary antibody against EGFP (mouse; 1:1000). Sections were then incubated with the secondary antibodies (donkey anti-rabbit; 1:1000 and goat anti-mouse; 1:1000) conjugated to a fluorophore for 2 hours RT and mounted on slides and coverslipped with VectaShield containing DAPI mounting medium and the signal was visualized with a Hamamatsu NANOZOOMER 2.0HT microscope. To evaluate viral expression in cholinergic interneurons, the procedure was repeated with a primary antibody against ChAT (Choline acetyl transferase; goat ; 1:500) and a secondary antibody (donkey anti-goat ; 1:1000).

Statistical analysis

Data are expressed as the mean ±SEM and statistical analyses were conducted using Prism software with a threshold for statistically significant differences set at $p \leq 0.05$. Each figure represents behavioral tests conducted on different male animal cohorts. The results were analyzed using adequate statistical and post hoc tests with repeated measurement when appropriate (see table 2 for details). Outlier values were removed with Grubbs' method when appropriate (see table 2 for details).

Data Availability: The dataset generated during and/or analyzed during the current study are available from the corresponding authors on reasonable request.

Code Availability: MatLab codes used in this study are available here: https://github.com/christophe-varin/FiberPhoto

REFERENCES

1. Watson, H. J. *et al.* Genome-wide association study identifies eight risk loci and implicates metabo-psychiatric origins for anorexia nervosa. *Nat Genet* **51**, 1207–1214 (2019).

2. Weiss, F. *et al.* Psychiatric Aspects of Obesity: A Narrative Review of Pathophysiology and Psychopathology. *J Clin Med* **9**, (2020).

3. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits. *Nat Genet* **47**, 1236–1241 (2015).

4. Krug, I. *et al.* Eating-related environmental factors in underweight eating disorders and obesity: are there common vulnerabilities during childhood and early adolescence? *Eur Eat Disord Rev* **21**, 202–208 (2013).

5. Wildes, J. E. & Marcus, M. D. Application of the Research Domain Criteria (RDoC) framework to eating disorders: emerging concepts and research. *Curr Psychiatry Rep* **17**, 30 (2015).

6. Casper, R. C. Not the Function of Eating, but Spontaneous Activity and Energy Expenditure, Reflected in 'Restlessness' and a 'Drive for Activity' Appear to Be Dysregulated in Anorexia Nervosa: Treatment Implications. *Front Psychol* **9**, 2303 (2018).

7. Shroff, H. *et al.* Features associated with excessive exercise in women with eating disorders. *Int J Eat Disord* **39**, 454–461 (2006).

8. Taranis, L. & Meyer, C. Associations between specific components of compulsive exercise and eating-disordered cognitions and behaviors among young women. *Int J Eat Disord* **44**, 452–458 (2011).

9. Matikainen-Ankney, B. A. & Kravitz, A. V. Persistent effects of obesity: a neuroplasticity hypothesis. *Ann N Y Acad Sci* **1428**, 221–239 (2018).

10. Rizk, M. *et al.* Physical Activity in Eating Disorders: A Systematic Review. *Nutrients* **12**, (2020).

11. Ruegsegger, G. N. & Booth, F. W. Running from Disease: Molecular Mechanisms Associating Dopamine and Leptin Signaling in the Brain with Physical Inactivity, Obesity, and Type 2 Diabetes. *Front Endocrinol (Lausanne)* **8**, 109 (2017).

12. Floresco, S. B. The Nucleus Accumbens: An Interface Between Cognition, Emotion, and Action. *Annual Review of Psychology* **66**, 25–52 (2015).

13. Frank, G. K. W. *et al.* Anorexia nervosa and obesity are associated with opposite brain reward response. *Neuropsychopharmacology* **37**, 2031–2046 (2012).

14. Kaye, W. H. *et al.* Does a shared neurobiology for foods and drugs of abuse contribute to extremes of food ingestion in anorexia and bulimia nervosa? *Biol Psychiatry* **73**, 836–842 (2013).

15. Volkow, N. D. *et al.* Low dopamine striatal D2 receptors are associated with prefrontal metabolism in obese subjects: possible contributing factors. *Neuroimage* **42**, 1537–1543 (2008).

16. Wang, G. J. *et al.* Brain dopamine and obesity. *Lancet* **357**, 354–357 (2001).

17. Bailer, U. F. *et al.* Interaction between serotonin transporter and dopamine D2/D3 receptor radioligand measures is associated with harm avoidant symptoms in anorexia and bulimia nervosa. *Psychiatry Res* **211**, 160–168 (2013).

18. Broft, A. *et al.* Striatal dopamine type 2 receptor availability in anorexia nervosa. *Psychiatry Res* **233**, 380–387 (2015).

19. Dang, L. C. *et al.* Associations between dopamine D2 receptor availability and BMI depend on age. *NeuroImage* **138**, 176–183 (2016).

20. Gaiser, E. C. *et al.* Elevated Dopamine D2/3 Receptor Availability in Obese Individuals: A PET Imaging Study with [11C](+)PHNO. *Neuropsychopharmacology* **41**, 3042–3050 (2016).

21. Carvalho Poyraz, F. *et al.* Decreasing Striatopallidal Pathway Function Enhances

Motivation by Energizing the Initiation of Goal-Directed Action. *Journal of Neuroscience* **36**, 5988–6001 (2016).

22. Gallo, E. F. *et al.* Accumbens dopamine D2 receptors increase motivation by decreasing inhibitory transmission to the ventral pallidum. *Nature Communications* **9**, (2018).

23. Natsubori, A. *et al.* Ventrolateral Striatal Medium Spiny Neurons Positively Regulate Food-Incentive, Goal-Directed Behavior Independently of D1 and D2 Selectivity. *The Journal of Neuroscience* **37**, 2723–2733 (2017).

24. Soares-Cunha, C. *et al.* Activation of D2 dopamine receptor-expressing neurons in the nucleus accumbens increases motivation. *Nat Commun* **7**, 11829 (2016).

25. Soares-Cunha, C. *et al.* Nucleus Accumbens Microcircuit Underlying D2-MSN-Driven Increase in Motivation. *eneuro* **5**, ENEURO.0386-18.2018 (2018).

26. Soares-Cunha, C. *et al.* Nucleus accumbens medium spiny neurons subtypes signal both reward and aversion. *Mol Psychiatry* **25**, 3241–3255 (2020).

27. Soares-Cunha, C. *et al.* Distinct role of nucleus accumbens D2-MSN projections to ventral pallidum in different phases of motivated behavior. *Cell Rep* **38**, 110380 (2022).

28. Bond, C. W. *et al.* Medial Nucleus Accumbens Projections to the Ventral Tegmental Area Control Food Consumption. *J Neurosci* **40**, 4727–4738 (2020).

29. O'Connor, E. C. *et al.* Accumbal D1R Neurons Projecting to Lateral Hypothalamus Authorize Feeding. *Neuron* **88**, 553–564 (2015).

30. Thoeni, S., Loureiro, M., O'Connor, E. C. & Lüscher, C. Depression of Accumbal to Lateral Hypothalamic Synapses Gates Overeating. *Neuron* **107**, 158-172.e4 (2020).

31. Zhu, X., Ottenheimer, D. & DiLeone, R. J. Activity of D1/2 Receptor Expressing Neurons in the Nucleus Accumbens Regulates Running, Locomotion, and Food Intake. *Front Behav Neurosci* **10**, (2016).

32. Matikainen-Ankney, B. A. *et al.* Nucleus Accumbens D1 Receptor–Expressing Spiny Projection Neurons Control Food Motivation and Obesity. *Biological Psychiatry* **93**, 512–523 (2023).

33. Labouesse, M. A. *et al.* Striatal dopamine 2 receptor upregulation during development predisposes to diet-induced obesity by reducing energy output in mice. *Proc Natl Acad Sci U S A* **115**, 10493–10498 (2018).

34. Johnson, P. M. & Kenny, P. J. Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nature Neuroscience* **13**, 635–641 (2010).

35. Welch, A. C. *et al.* Dopamine D2 receptor overexpression in the nucleus accumbens core induces robust weight loss during scheduled fasting selectively in female mice. *Mol. Psychiatry* (2019) doi:10.1038/s41380-019-0633-8.

36. Beeler, J. A., Faust, R. P., Turkson, S., Ye, H. & Zhuang, X. Low Dopamine D2 Receptor Increases Vulnerability to Obesity Via Reduced Physical Activity, Not Increased Appetitive Motivation. *Biological Psychiatry* **79**, 887–897 (2016).

37. Friend, D. M. *et al.* Basal Ganglia Dysfunction Contributes to Physical Inactivity in Obesity. *Cell Metabolism* **25**, 312–321 (2017).

38. Donofry, S. D., Stillman, C. M. & Erickson, K. I. A review of the relationship between eating behavior, obesity and functional brain network organization. *Soc Cogn Affect Neurosci* **15**, 1157–1181 (2020).

39. Pujol, J. *et al.* Dysfunctional Brain Reward System in Child Obesity. *Cereb Cortex* **31**, 4376–4385 (2021).

40. Haynos, A. F. *et al.* Resting state functional connectivity of networks associated with reward and habit in anorexia nervosa. *Hum Brain Mapp* **40**, 652–662 (2018).

41. Uniacke, B. *et al.* Resting-state connectivity within and across neural circuits in anorexia nervosa. *Brain Behav* **9**, e01205 (2018).

42. Ducrocq, F. *et al.* Causal Link between n-3 Polyunsaturated Fatty Acid Deficiency and

Motivation Deficits. *Cell Metab.* **31**, 755-772.e7 (2020).

43. Lowe, M. R. Weight suppression is a risk factor for eating disorders: Implications for
etiology, maintenance, and treatment. *Am J Clin Nutr* **112**, 907–908 (2020).

44. Bonnavion, P. *et al.* Unexpected inhibition of motor function by dopamine activation of
D1/D2 co-expressing striatal neurons. 2022.04.05.487163 Preprint at
https://doi.org/10.1101/2022.04.05.487163 (2022).

45. Schorr, M. & Miller, K. K. The endocrine manifestations of anorexia nervosa: mechanisms
and management. *Nat Rev Endocrinol* **13**, 174–186 (2017).

46. Adan, R. a. H. *et al.* Neurobiology driving hyperactivity in activity-based anorexia. *Curr*
Top Behav Neurosci **6**, 229–250 (2011).

47. Steinglass, J. E. *et al.* Increased capacity to delay reward in anorexia nervosa. *J Int*
Neuropsychol Soc **18**, 773–780 (2012).

48. Wierenga, C. E. *et al.* Hunger does not motivate reward in women remitted from anorexia
nervosa. *Biol Psychiatry* **77**, 642–652 (2015).

49. Klein, D. A. *et al.* Behavioral assessment of the reinforcing effect of exercise in women
with anorexia nervosa: further paradigm development and data. *Int J Eat Disord* **43**, 611–
618 (2010).

50. Schebendach, J., Broft, A., Foltin, R. W. & Walsh, B. T. Can the reinforcing value of food
be measured in bulimia nervosa? *Appetite* **62**, 70–75 (2013).

51. Ekkekakis, P., Vazou, S., Bixby, W. R. & Georgiadis, E. The mysterious case of the public
health guideline that is (almost) entirely ignored: call for a research agenda on the causes
of the extreme avoidance of physical activity in obesity. *Obes Rev* **17**, 313–329 (2016).

52. Le Merrer, J. *et al.* Balance Between Projecting Neuronal Populations of the Nucleus
Accumbens Controls Social Behavior in Mice. *Biol Psychiatry* **95**, 123–135 (2024).

53. Varin, C., Cornil, A., Houtteman, D., Bonnavion, P. & de Kerchove d'Exaerde, A. The
respective activation and silencing of striatal direct and indirect pathway neurons support
behavior encoding. *Nat Commun* **14**, 4982 (2023).

54. Iversen, I. H. Techniques for establishing schedules with wheel running as reinforcement
in rats. *J Exp Anal Behav* **60**, 219–238 (1993).

55. Hurel, I. *et al.* Beyond the Activity-Based Anorexia Model: Reinforcing Values of Exercise
and Feeding Examined in Stressed Adolescent Male and Female Mice. *Front Pharmacol*
10, 587 (2019).

56. Sandoval-Rodríguez, R. *et al.* D1 and D2 neurons in the nucleus accumbens enable
positive and negative control over sugar intake in mice. *Cell Reports* **42**, 112190 (2023).

57. Kelley, A. E., Baldo, B. A., Pratt, W. E. & Will, M. J. Corticostriatal-hypothalamic circuitry
and food motivation: integration of energy, action and reward. *Physiol Behav* **86**, 773–
795 (2005).

58. Yonemochi, N., Ardianto, C., Ueda, D., Kamei, J. & Ikeda, H. GABAergic function in the
lateral hypothalamus regulates feeding behavior: Possible mediation via orexin.
Neuropsychopharmacol Rep **39**, 289–296 (2019).

59. Stratford, T. R. & Wirtshafter, D. Evidence that the nucleus accumbens shell, ventral
pallidum, and lateral hypothalamus are components of a lateralized feeding circuit. *Behav*
Brain Res **226**, 548–554 (2012).

60. Vachez, Y. M. *et al.* Ventral arkypallidal neurons inhibit accumbal firing to promote reward
consumption. *Nat Neurosci* **24**, 379–390 (2021).

61. Aitta-Aho, T. *et al.* Accumbal Cholinergic Interneurons Differentially Influence Motivation
Related to Satiety Signaling. *eNeuro* **4**, ENEURO.0328-16.2017 (2017).

62. Baldo, B. A. & Kelley, A. E. Discrete neurochemical coding of distinguishable motivational
processes: insights from nucleus accumbens control of feeding. *Psychopharmacology*
(Berl) **191**, 439–459 (2007).

63. Georges, F. & Aston-Jones, G. Activation of ventral tegmental area cells by the bed

nucleus of the stria terminalis: a novel excitatory amino acid input to midbrain dopamine neurons. *J Neurosci* **22**, 5173–5187 (2002).

64. Grace, A. A. In vivo and in vitro intracellular recordings from rat midbrain dopamine neurons. *Ann N Y Acad Sci* **537**, 51–76 (1988).

65. Ducrocq, F. *et al.* Decrease in Operant Responding Under Obesogenic Diet Exposure is not Related to Deficits in Incentive or Hedonic Processes. *Obesity* **27**, 255–263 (2019).

66. Johnson, A. W. Characterizing ingestive behavior through licking microstructure: Underlying neurobiology and its use in the study of obesity in animal models. *International Journal of Developmental Neuroscience* **64**, 38–47 (2018).

67. Ward, R. D., Kellendonk, C., Kandel, E. R. & Balsam, P. D. Timing as a window on cognition in schizophrenia. *Neuropharmacology* **62**, 1175–81 (2012).

68. Jean-Richard-Dit-Bressel, P., Clifford, C. W. G. & McNally, G. P. Analyzing Event-Related Transients: Confidence Intervals, Permutation Tests, and Consecutive Thresholds. *Front Mol Neurosci* **13**, 14 (2020).

69. Liu, Y. *et al.* The Mesolimbic Dopamine Activity Signatures of Relapse to Alcohol-Seeking. *J Neurosci* **40**, 6409–6427 (2020).

70. Maris, E. & Oostenveld, R. Nonparametric statistical testing of EEG- and MEG-data. *J Neurosci Methods* **164**, 177–190 (2007).

Acknowledgments: We thank the Bordeaux Imaging Center funded by ANR-10-INBS-04 for imaging, Motac Neuroscience for lending the rotarod apparatus, Céline Ducroix-Crépy and Jean-Christophe Helbling (INRAE UMR 1286) for genotyping and the staff from the animal facility of INRAE UMR 1286 for animal care.

Funding: This study was supported by INRAE and University of Bordeaux; Idex Bordeaux "chaire d'installation" (ANR-10-IDEX-03-02) (PT), NARSAD Young Investigator Grants from the Brain and Behavior Foundation (PT), ANR "SynLip" (ANR-16-CE16-0022) (PT), ANR "FrontoFat" (ANR-20-CE14-0020) (PT), ANR "StriaPOM" (ANR-21-CE14-0018) (PT), Institut de Recherche en Santé publique (IReSP) Aviesan APP-addiction 2019 (PT), Labex "BRAIN" (PT and RW), Region Nouvelle Aquitaine 2014-1R30301-00003023 (PT); Fondation pour la Recherche Médicale (FRM "Environnement et Santé") (GF and PT), ANR "HA-CTion" (ANR-19-HBPR-0002) (GF and PT); PRESTIGE-2017-2-0031 (AC), "Fondation pour la Recherche Médicale" SPF201809007095 (AC); Canadian Institute for Health Research (201309OG-312343-PT; 201803PJT-399980-PT) (BG); FD and RW are recipients of PhD fellowships from the French Ministry of Research. AP is a recipient of a PhD fellowship from the "Ecole Universitaire de Recherche" (EUR Neuro, Bordeaux Neurocampus); AKE is Research Director of FRS-FNRS and WELBIO investigator. AKE was supported by grants from from FRS-FNRS (#23587797, #33659288, #33659296), Fondation Simone et Pierre Clerdent (Prize 2018) and Fondation ULB.

Authors contributions: RW, GF, FD, FG and PT designed research, RW, AP, GRF, EM, AC, MFA, MP, RO, AO, VDP and FD performed research, RW, FD, FG and PT supervised

research, RW, AP, GRF, CV, EM, LH, AC, MFA, MP, FD, FG and PT analyzed data, RAA, AKE, BG and FC provided expertise, reagents and material, RW, FC, GF, FD, FG and PT wrote the manuscript. All authors edited and approved the manuscript.

Competing interests: The authors declare no competing interests.

FIGURES LEGENDS

Figure 1: Effect of chemogenetic manipulations of D2- and D1-neurons of the NAc on the motivational dimension of feeding behavior.

(A) Experimental timeline. **(B)** Representative coronal sections for mCherry expression after injection of a DREADD-mCherry-carrying viral vector in the NAc core of D2-cre mice (left), highlighting projections in the ventral pallidum (VP, middle) but not the midbrain (right). **(C-F)** Effects of chemogenetic-mediated D2-neurons activation (DREADD Gq, n=7) and inhibition (DREADD Gi, n=9) under administration of CNO at 2 mg/kg on the number of reinforced lever presses (C), breakpoint (D) the percentage of mice that keep on responding as a function of time (E) and the ratio requirement plotted as the rate of lever presses as a function of PR ratios (F). **(G)** Representative coronal sections for mCherry expression after injection DREADD-mCherry-carrying viral vector in the NAc core of D1-cre mice (left), highlighting projections in the ventral pallidum (VP, middle) and the midbrain (right). **(H-K)** Effects of chemogenetic-mediated D1-neurons activation (DREADD Gq, n=8) and inhibition (DREADD Gi, n=9) under administration of CNO at 0.5 mg/kg on the number of reinforced lever presses (H), breakpoint (I) the percentage of mice that keep on responding as a function of time (J) and the ratio requirement plotted as the rate of lever presses as a function of PR ratios (K). Data are averages of 2 PR sessions under CNO or saline administrations in a within-subject, counterbalanced manner. AcbC: Accumbens core; AcbSh: Accumbens shell; Cpu: Caudate Putamen; DG: Dentate Gyrus; LV: Lateral ventricle; SNC: *Substantia nigra pars compacta*; SNR: *Substantia nigra pars reticulata*; Vp: Ventral Pallidum; VTA: Ventral tegmental area. * : $p < 0.05$; ** : $p < 0.01$; *** : $p < 0.001$; Error bars = s.e.m. See Table 2 for statistical analyses.

Figure 2: D1- and D2-neurons of the NAc are differentially activated during reward consumption and oppositely regulate reward consumption in a pavlovian conditioning paradigm

(A) Schematic representation of the procedure. **(B)** Effect of chemogenetic inhibition (DREADD Gi, D2-neurons n=7, D1-neurons n=6) and activation (DREADD Gq, n=8, D1-

neurons n=9) of either D2-neurons or D1-neurons on the number of licks on the trough that delivers the reward. **(C-D)** Comparison of bulk calcium dynamics in D1- (n=5 mice) and D2-neurons (n=5mice) aligned to the onset of the conditioned stimuli (CS) (C), or to the onset of feeding (lick) and quantification of the AUC and average z-score (D), measured through expression of the calcium sensor GcAMP in the NAc, in a pavlovian conditioning paradigm. **(E-F)** Effect of chemogenetic activation of D1-neurons (D1 Gq, n=4 mice) on dopamine dynamics and quantification of the AUC and average z-score, measured through the dopamine sensor dLight coupled with fiber photometry in the NAc, aligned with CS onset (E) or lick (F). **(G-H)** Effect of chemogenetic activation of D2-neurons (D2 Gq, n=3 mice) on calcium dynamics measured through the dopamine sensor dLight coupled with fiber photometry in the NAc, aligned with CS onset (G) or lick (H). $p < 0.05$; ** : $p < 0.01$; *** : $p < 0.001$; Error bars = s.e.m. See Table 2 for statistical analyses.

Figure 3: D1- and D2-neurons of the NAc are differentially activated during feeding and running and oppositely regulate the balance between feeding and voluntary running

(A) (Top) Experimental timeline for recording of calcium dynamics during feeding. (Bottom) Normalized calcium activity of D1 and D2-neurons during feeding trials (Z score ; 12 trials / animal) with quantification of the AUC, average z-score, Peak z-score and transient rates. **(B)** (Top) Experimental timeline for recording of calcium dynamics during running. (Bottom) Normalized calcium activity of D1 and D2-neurons during running trials (Z score ; 12 trials / animal) with quantification of the AUC, average z-score, Peak z-score and transient rates. For both A and B, D2-cre: n=5 ; D1-cre: n=5; Blue and red bars represent statistical difference to baseline, black bar significant difference between the relative activity of both subpopulations. **(C-E)** (C) Experimental design. Normalized chow consumption and wheel running comparing 1-hour pre-CNO and 1-hour post-CNO injection for chemogenetic activation and inhibition of D2-neurons (D) and D1-neurons (E). (D2-cre: DREADD Gi n=7, DREADD Gq n=7 ; D1-cre: DREADD Gi n=8, DREADD Gq n=8. * : $p < 0.05$; ** : $p < 0.01$; *** : $p < 0.001$; Error bars = s.e.m. See Table 2 for statistical analyses.

Figure 4: Effect of chronic chemogenetic manipulations of D2-neurons and D1-neurons of the NAc in a wheel-running/high-fat free-feeding choice procedure.

(A) Experimental design. **(B-D)** High-fat consumption over time (B), variation in wheel running (C) and variation in total and fat mass (D) were measured under chronic D2-neurons inhibition (D2 Gi) and activation (D2Gq) compared to WT controls, as well as after CNO cessation. (E-G) The same design was followed for chronic D1-neurons manipulations. D2-WT n=9; D2-Gi n=8; D2-Gq n=9; D1-WT n=12; D1-Gi n=11; D1-Gq n=12. \$: $0.05 < p < 0.1$; *: $p < 0.05$; **: $p < 0.01$; Error bars = s.e.m. See Table 2 for statistical analyses.

Figure 5: Effect of chronic and simultaneous chemogenetic manipulations of D2-neurons and D1-neurons of the NAc in a wheel-running/high-fat free-feeding choice procedure.

(A) Schematic design for the transgenic and viral strategy for concomitant chemogenetic manipulation of D1- and D2-neuroins of the NAc (left) and representative pattern of expression of cre-dependent GFP and flp-dependent mCherry in double transgenic D1-FlpxD2-cre mice (right). **(B-D)** High-fat consumption over time (B), variation in wheel running (C) and variation in total and fat mass (D) were measured under chronic manipulations of both subpopulations (D1Gq-D2Gi: concomitant activation of D1-neurons and inhibition of D2-neurons; D1Gi-D2Gq: concomitant inhibition of D1-neurons and activation of D2-neurons) compared to WT controls, as well as after CNO cessation. D1-D2 WT n=9 ; D1Gi-D2Gq n=9, D1Gq-D2Gi n=8. $: 0.05<p<0.1 * : p<0.05 ; ** : p<0.01; *** : p<0.001 Error bars = s.e.m. See Table 2 for statistical analyses.

Figure 6: Concomitant activation of D1- and inhibition of D2-neurons precipitates weight loss in rodent models of anorexia nervosa

(A-C) Measures of weight variation (A) and wheel-running (B) in a procedure of progressive food restriction. (C) Dosage of plasma leptin and ghrelin levels and weight following the food-restriction procedure. D1-D2 WT n=9 ; D1Gi-D2Gq n=9, D1Gq-D2Gi n=8. $: 0.05<p<0.1 * : p<0.05 ; ** : p<0.01; *** : p<0.001 Error bars = s.e.m. See Table 2 for statistical analyses. **(D-G)** Measures of weight variation (D), wheel running (E) and chow consumption (F) in an Activity-Based Anorexia (ABA) model. (G) Probability of survival (i.e. % of animals displaying less than 20% weight loss) over the ABA procedure. D1-D2 WT n=8 ; D1Gi-D2Gq n=9, D1Gq-D2Gi n=8. $: 0.05<p<0.1 * : p<0.05 ; ** : p<0.01; *** : p<0.001 Error bars = s.e.m. See Table 2 for statistical analyses.

 TABLES

 Table 1: Key resources

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Primary antibody anti-DS red (rabbit)	Takara Bio	632496
Primary antibody anti-GFP (mouse)	Clontech	632280
Primary antibody anti-Choline acetyl transferase (goat)	Merck Millipore	AB144P
Secondary antibody anti-rabbit alexa 568 (donkey)	ThermoFisher	A-10042
Secondary antibody anti-rabbit alexa 488 (donkey)	ThermoFisher	A-21206
Secondary antibody anti-goat alexa 568 (donkey)	ThermoFisher	A-11057
Secondary antibody anti-goat alexa 488 (donkey)	ThermoFisher	A-11055
Bacterial and Virus Strains		
AAV8-hSyn1-DIO-EGFP	University of Zurich	v115
AAV8-hSyn1-dFRT-mCherry	University of Zurich	v188
AAV8-hSyn-DIO-hM4D(Gi)-mCherry	Addgene	44362-AAV8
AAV8-hSyn-DIO-hM4D(Gi)-mCherry	University of Zurich	V84
AAV8-hSyn-DIO-hM3D(Gq)-mCherry	Addgene	44361-AAV8
AAV8-hSyn-DIO-hM3D(Gq)-mCherry	University of Zurich	V89
AAV8-hSyn1-dFRT-hM4D(Gi)-mCherry(rev)-dFRT-WPRE-hGHp(A)	University of Zurich	V189
AAV8-hSyn1-dFRT-hM3D(Gq)-mCherry(rev)-dFRT-WPRE-hGHp(A)	University of Zurich	V190
AAV8-hSyn1-dlox-jGCaMP8m(rev)-dlox-WPRE-SV40p(A)	University of Zurich	V628
AAV9-hSyn1-chl-dLight1.3b-WPRE-bGHp(A)	University of Zurich	v565
Chemicals, Peptides, and Recombinant Proteins		
Clozapine-N-Oxide (CNO)	ENZO	BML-NS105
Recombinant Mouse Leptin Protein, CF	Bio-Techne	498-OB-05M
Ghrelin	Bachem	H-4862.1000
JHU37160 dihydrochloride (J60 DREADD ligand)	hellobio	HB6261
Critical Commercial Assays		
ACCU-CHEK Performa	Roche	N/A
Fiber Optic Cannulae 0,5 NA, 400µm flat tip, L:6mm	RWD	N/A
Acylated Ghrelin (mouse, rat) Express ELISA kit	Bertin Bioreagent	A05117
Mouse Leptin ELISA Kit - 96-Well Plate	Sigma Aldrich - Merck	EZML-82K
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	Janvier Labs	N/A
Mouse: Tg(Drd1-cre)FK150Gsat/Mmucd	GENSAT	RRID: MMRRC_029178-UCD
Mouse: Tg(Drd2-cre)ER44Gsat/Mmucd	GENSAT	RRID: MMRRC_017263-UCD
Mouse D1Flp (D1 expressing flippase and YFP)	Produced by the Laboratory of B. Giros	N/A

 Table 2: Summary of statistical analyses

Data	Statistical test	n	Outcome measure	P value
Figure 1				
Figure 1C	Two-way RM ANOVA	D2 cre mice: Dreadd Gi = 9 Dreadd Gq= 7	Lever presses	Virus effect $F_{(1, 14)}=6.604$ p=0.0222 CNO effect $F_{(1, 14)}=0.3902$ p=0.5423 Interaction $F_{(1, 14)}=19.29$ p=0.0006

				Bonferroni's multiple comparisons test: DREADD Gi: Sal vs CNO p=0.0258 DREADD Gq: Sal vs CNO p=0.0096
Figure 1D	Two-way RM ANOVA	D2 cre mice: Dreadd Gi = 9 Dreadd Gq= 7	Break point	Virus effect F(1, 14)=6.773 p=0.0209 CNO effect F(1, 14)=0.04977 p=0.8267 Interaction F(1, 14)=14.92 p=0.0017 Bonferroni's multiple comparisons test: DREADD Gi: Sal vs CNO p=0.0310 DREADD Gq: Sal vs CNO p=0.0327
Figure 1E	Log-rank (Mantel-Cox) test	D2 cre mice: Dreadd Gi = 9	Survival Curve	χ2=4.511 ; p value=0.0337
Figure 1E	Gehan-Breslow-Wilcoxon test	D2 cre mice: Dreadd Gq= 7	Survival Curve	χ2=6.338 ; p value=0.0118
Figure 1F	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre mice: Dreadd Gi = 9	Rate (presses/s)	Ratio effect F(3.138, 25.10)=20.67 p<0.0001 (ε correction: 0.3922) CNO effect F(1.000, 8.000)=0.1515 p=0.7073 (ε correction: 1.000) Interaction F(3.379, 27.03)=1.024 p=0.4040 (ε correction: 0.4224)
Figure 1F	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre mice: Dreadd Gq= 7	Rate (presses/s)	Ratio effect F(1.245, 7.469)=4.953 p=0.0539 (ε correction: 0.1556) CNO effect F(1.000, 6.000)=2.183 p=0.1900 (ε correction: 1.000) Interaction F(2.468, 12.34)=0.7316 p=0.5278 (ε correction: 0.3085)
Figure 1H	Two-way RM ANOVA	D1 cre mice: Dreadd Gi = 9 Dreadd Gq=8	Lever presses	Virus effect F(1, 15)=1.992 p=0.1785 CNO effect F(1, 15)=28.22 p<0.0001 Interaction F(1, 15)=1.658 p=0.2174
Figure 1I	Two-way RM ANOVA	D1 cre mice: Dreadd Gi = 9 Dreadd Gq=8	Break point	Virus effect F(1, 15)=0.06963 p=0.7955 CNO effect F(1, 15)=25.29 p=0.0001 Interaction F(1, 15)=0.2090 p=0.6541
Figure 1J	Log-rank (Mantel-Cox) test	D1 cre mice: Dreadd Gi = 9	Survival Curve	χ2=8.910 ; p value=0.0028
Figure 1J	Log-rank (Mantel-Cox) test	D1 cre mice: Dreadd Gq= 8	Survival Curve	χ2=3.652 ; p value=0.0560
Figure 1K	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 cre mice: Dreadd Gi = 9	Rate (presses/s)	Ratio effect F(2.171, 17.37)=17.20 p<0.0001 (ε correction: 0.2713) CNO effect F(1, 8)=1.422 p=0.2673 (ε correction: 1.000) Interaction F(2.167, 14.90)=1.135 p=0.3517 (ε correction: 0.2709)
Figure 1K	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 cre mice: Dreadd Gq=8	Rate (presses/s)	Ratio effect F(1.361, 9.525)=9.687 p<0.0001 (ε correction: 0.1944) CNO effect F(1, 7)=0.3425 p=0.5768 (ε correction: 1.000) Interaction F(2.055, 13.80)=1.131 p=0.3605 (ε correction: 0.2936)
Figure 2				

Figure 2B	Paired T-test	D2 cre mice Dreadd Gi = 7	Number of licks	t=3.232, df=6, p=0.0179
Figure 2B	Paired T-test	D2 cre mice Dreadd Gq = 8	Number of licks	t=2.761, df=7, p=0.0281
Figure 2B	Paired T-test	D1 cre mice Dreadd Gi = 6	Number of licks	t=3.798, df=5, p=0.0127
Figure 2B	Paired T-test	D1 cre mice Dreadd Gq = 9	Number of licks	t=5.236, df=8, p=0.0008
Figure 2C	Two-way ANOVA	D1 cre mice = 5 (event =528) D2 cre mice = 5 (event =591)	GCaMP CS Area under the curve	Time effect F(1, 2234)=35.21 p<0.0001 Group effect F(1, 2234)=2.404 p=0.1211 Interaction F(1, 2234)=1.450 p=0.2286
Figure 2C	Two-way ANOVA	D1 cre mice = 5 (event =528) D2 cre mice = 5 (event =591)	GCaMP CS Average Zscore	Time effect F(1, 2234)=35.21 p<0.0001 Group effect F(1, 2234)=2.404 p=0.1211 Interaction F(1, 2234)=1.450 p=0.2286
Figure 2D	Two-way ANOVA	D1 cre mice = 5 (event =593) D2 cre mice = 5 (event =660)	GCaMP lick Area under the curve	Time effect F(1, 2502)=302.5 p<0.0001 Group effect F(1, 2502)=4.301 p=0.0382 Interaction F(1, 2502)=5.141 p=0.0235 Bonferroni's multiple comparisons test: -2-0s: D1 vs D2 p=0.0043 0-2s: D1 vs D2 p=0.9999
Figure 2D	Two-way ANOVA	D1 cre mice = 5 (event =593) D2 cre mice = 5 (event =660)	GCaMP lick Average Zscore	Time effect F(1, 2502)=302.5 p<0.0001 Group effect F(1, 2502)=4.301 p=0.0382 Interaction F(1, 2502)=5.141 p=0.0235 Bonferroni's multiple comparisons test: -2-0s: D1 vs D2 p=0.0043 0-2s: D1 vs D2 p=0.9882
Figure 2E	Two-way RM ANOVA	D1 cre mice = 4 event Saline = 252 event CNO = 168	Dlight CS Area under the curve	Time effect F(1, 418)=308.5 p<0.0001 CNO effect F(1, 418)=1.019 p=0.3134 Interaction F (1, 418) = 0.5410 p=0.4624
Figure 2E	Two-way RM ANOVA	D1 cre mice = 4 event Saline = 252 event CNO = 168	Dlight CS Average Zscore	Time effect F(1, 418)=311.3 p<0.0001 CNO effect F(1, 418)=0.9687 p=0.3256 Interaction F(1, 418)=0.4511 p=0.5022
Figure 2F	Two-way RM ANOVA	D1 cre mice = 4 event Saline = 271 event CNO = 186	Dlight LICK Area under the curve	Time effect F(1, 455)=4.083 p=0.0439 CNO effect F(1, 455)=4.010 p=0.0458 Interaction F(1, 455)=2.429 p=0.1198
Figure 2F	Two-way RM ANOVA	D1 cre mice = 4 event Saline = 271 event CNO = 186	Dlight LICK Average Zscore	Time effect F(1, 455)=2.457 p=0.1177 CNO effect F(1, 455)=3.798 p=0.0519 Interaction F(1, 455)=2.241 p=0.1351
Figure 2G	Two-way RM ANOVA	D2 cre mice = 3 event Saline = 170 event CNO = 105	Dlight CS Area under the curve	Time effect F(1, 273)=200.2 p<0.0001 CNO effect F(1, 273)=1.849 p=0.1751 Interaction F(1, 273)=2.151 p=0.1436
Figure 2G	Two-way RM ANOVA	D2 cre mice = 3 event Saline = 170 event CNO = 105	Dlight CS Average Zscore	Time effect F(1, 273)=203.5 p<0.0001 CNO effect F(1, 273)=1.887 p=0.1706 Interaction F(1, 273)=2.355 p=0.1260
Figure 2H	Two-way RM ANOVA	D2 cre mice = 3 event Saline = 192 event CNO = 112	Dlight LICK Area under the curve	Time effect F(1, 302)=82.15 p<0.0001 CNO effect F(1, 302)=31.40 p<0.0001 Interaction F(1, 302)=35.76 p<0.0001 Bonferroni's multiple comparisons test:

| | | | | -2 0s: Sal vs CNO p=0.9999 |
				0 2s: Sal vs CNO p=0.0001
Figure 2H	Two-way RMANOVA	D2 cre mice = 3 event Saline = 192 event CNO = 112	Dlight LICK Average Zscore	Time effect F(1, 302)=79.18 p<0.0001 CNO effect F(1, 302)=32.06 p<0.0001 Interaction F(1, 302)=36.33 p<0.0001 Bonferroni's multiple comparisons test: -2 0s: Sal vs CNO p=0.9999 0 2s: Sal vs CNO p=0.0001
Figure 3				
Figure 3A	Mixed-effect model (Satterthwaite correction for degrees of freedom)	D1 Cre = 5 D2 Crep = 5 12 trials per animals	GCamp Calcium activity during feeding trials (Zscore; D1 vs D2)	Intercept effect F(1, 12.499)=197.66 p=4.97e-9 Genotype effect F(1, 12.499)=0.1867 p=0.6731 Time effect F(1, 37.133)=4.1772 p=0.0481 Genotype x Time F(1, 37.133)=10.665 p=0.0024 post hoc-permutation test between genotypes
Figure 3A	Two-way RM ANOVA	D1 Cre = 5 event = 59 D2 Cre = 5 event = 60	GCamp Calcium activity during feeding trials Area under the curve	Interaction F(1, 117)=19.88 p<0.0001 Time effect F(1, 117)=56.85 p<0.0001 Group effect F(1, 117)=7.376 p=0.0076 -40-0s: D1 vs D2 p=0.9999 0-60s: D1 vs D2 p=0.0001
Figure 3A	Two-way RM ANOVA	D1 Cre = 5 event = 59 D2 Cre = 5 event = 60	GCamp Calcium activity during feeding trials Average Zscore	Interaction F(1, 117)=22.11 p<0.0001 Time effect F(1, 117)=18.37 p<0.0001 Group effect F(1, 117)=5.489 p=0.0208 -40-0s: D1 vs D2 p=0.9999 0-60s: D1 vs D2 p=0.0001
Figure 3A	Two-way RM ANOVA	D1 Cre = 5 event = 59 D2 Cre = 5 event = 60	GCamp Calcium activity during feeding trials Peak Zscore	Interaction F(1, 117)=31.07 p<0.0001 Time effect F(1, 117)=186.1 p<0.0001 Group effect F(1, 117)=0.07555 p=0.7839 -40 0s: D1 vs D2 p=0.0099 0 60s: D1 vs D2 p=0.0391
Figure 3A	Two-way RM ANOVA	D1 Cre = 5 event = 59 D2 Cre = 5 event = 60	GCamp Calcium activity during feeding trials Transient rate	Interaction F(1, 117)=12.67 p=0.0005 Time effect F(1, 117)=47.33 p<0.0001 Group effect F(1, 117)=0.2608 p=0.6105 -40 0s: D1 vs D2 p=0.0089 0 60s: D1 vs D2 p=0.0654
Figure 3B	Mixed-effect model (Satterthwaite correction for degrees of freedom)	D1GCamp = 5 D2GCamp = 5 12 trials per animals 5 event per trials	GCamp Calcium activity during wheel running trials (Zscore D1 vs D2)	Intercept effect F(1, 9.998)=138.675 p=6.15e-7 Genotype effect F(1, 9.998)=1.2689 p=0.2879 Time effect F(1, 20.362)=0.6199 P= 0.4402 Genotype x Time F(1, 20.362)=0.4579 p=0.5062 (post hoc-permutation test between genotypes)
Figure 3B	Two-way RM ANOVA	D1 Cre = 5 event = 60 D2 Cre = 5 event = 59	GCamp Calcium activity during wheel running trials Area under the curve	Interaction F(1, 117)=16.73 p<0.0001 Time effect F(1, 117)=96.48 p<0.0001 Group effect F(1, 117)=7.219 p=0.0083 -40-0s: D1 vs D2 p=0.9999

				0-60s: D1 vs D2 p=0.0001
Figure 3B	Two-way RM ANOVA	D1 Cre = 5 event = 60 D2 Cre = 5 event = 59	GCamp Calcium activity during wheel running trials Average Zscore	Interaction F(1, 117)=18.22 p<0.0001 Time effect F(1, 117)=51.69 p<0.0001 Group effect F(1, 117)=5.598 p=0.0196 -40-0s: D1 vs D2 p=0.9999 0-60s: D1 vs D2 p=0.0001
Figure 3B	Two-way RM ANOVA	D1 Cre = 5 event = 60 D2 Cre = 5 event = 59	GCamp Calcium activity during wheel running trials Peak Zscore	Interaction F(1, 117)=1.616 p=0.2062 Time effect F(1, 117)=56.99 p<0.0001 Group effect F(1, 117)=0.1438 p=0.7052
Figure 3B	Two-way RM ANOVA	D1 Cre = 5 event = 60 D2 Cre = 5 event = 59	GCamp Calcium activity during wheel running trials Transient rate	Interaction F(1, 118)=3.456 p=0.0655 Time effect F(1, 118)=60.18 p<0.0001 Group effect F (1, 118) = 0.004899 p=0.9443
Figure 3D	Two-way RM ANOVA	D2Gi = 7 D2Gq = 8 1 D2Gq outlier value was excluded with Grubbs' method (Alpha = 0.01)	Chow consumption (ratio CNO/SAL)	Infusion effect F(1, 12)=8.570 p=0.0127 Virus effect F(1, 12)=0.3412 p=0.5700 Interaction F(1, 12)=0.5111 p=0.4823
Figure 3D	Two-way RM ANOVA	D2Gi = 7 D2Gq = 8 1 D2Gq outlier value was excluded with Grubbs' method (Alpha = 0.01)	Wheel running activity (ratio CNO/SAL)	Infusion effect F(1, 12)=8.410 p=0.0113 Virus effect F(1, 12)=23.88 p=0.0004 Interaction F(1, 12)=46.90 p<0.0001 Bonferroni's multiple comparisons test: D2Gi vs D2Gq: Pre infusion p>0.9999; Post infusion p<0.0001 Pre infusion vs Post infusion : D2Gi p=0.0326 D2Gq p<0.0001
Figure 3E	Two-way RM ANOVA	D1Gi = 9 D1Gq = 9 1 D1Gi and 1 D2Gq outlier value were excluded with Grubbs' method (Alpha = 0.01)	Chow consumption (ratio CNO/SAL)	Infusion effect F(1, 14)=0.06936 p=0.7961 Virus effect F(1, 14)=5.686 p=0.0318 Interaction F(1, 14)=10.97 p=0.0058 Bonferroni's multiple comparisons test: D1Gi vs D1Gq: Pre infusion p>0.9999; Post infusion p=0.0008 Pre infusion vs Post infusion : D1Gi p=0.0980 D1Gq p=0.0483
Figure 3E	Two-way RM ANOVA	D1Gi = 9 D1Gq = 9 1 D1Gi and 1 D2Gq outlier value were excluded with Grubbs' method (Alpha = 0.01)	Wheel running activity (ratio CNO/SAL)	Infusion effect F(1, 14)=0.6277 p=0.4414 Virus effect F(1, 14)=4.852 p=0.0449 Interaction F(1, 14)=18.60 p=0.0007 Bonferroni's multiple comparisons test: D1Gi vs D1Gq: Pre infusion p=0.4164; Post infusion p=0.0002 Pre infusion vs Post infusion : D1Gi p=0.0520 D1Gq p=0.0057
Figure 4				
Figure 4B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 WT = 9 D2 GI = 8 D2 Gq = 9	High fat consumption D1-9; D24-25	Time effect F(3.466, 79.71)=70.40 p<0.0001 (ε correction: 0.3851) Virus effect F(2, 23)=0.6509 p=0.5309 Interaction F(18, 207)=4.489 p<0.0001 Bonferroni's multiple comparisons test:

				D2Gq vs D2WT: Day3 p=0.0091; Day25 p=0.0330
				D2Gq vs D2Gi: Day1 p=0.0256; Day24 p=0.0197; Day25 p=0.0312
Figure 4B	One way ANOVA	D2 WT = 9 D2 GI = 8 D2 Gq = 9	High fat consumption mean D10-D22	F(2, 23)=0.6708 p=0.5210
Figure 4B	One way ANOVA	D2 WT = 9 D2 GI = 8 D2 Gq = 9	High fat consumption mean D26-D33	F(2, 23)=1.567 p=0.2302
Figure 4C	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 WT = 7 D2 GI = 7 D2 Gq = 8 (2 WT, 1 D1-Gi and 1 D1-Gq animals were exposed to a running wheel without recording)	Running wheel variation D1-9; D24-25	Time effect F(3.868, 72.36)=4.052 p=0.0055 (ε correction: 0.2276) Virus effect F(2, 19)=1.659 p=0.2167 Interaction F(34, 318)=3.536 p<0.0001 Bonferroni's multiple comparisons test: D2Gi vs D2WT: Day1 p=0.0272 D2Gq vs D2WT: Day11 p=0.0305; Day13 p=0.0038; Day15 p=0.0208; Day19 p=0.0328 ; Day21 p=0.0145 ; Day23 p=0.0228
Figure 4D	One way ANOVA	D2 WT = 9 D2 GI = 8 D2 Gq = 9	Weight variation following CNO period	F(2, 23)=0.3597 p=0.7017
Figure 4D	Brown-Forsythe ANOVA test	D2 WT = 9 D2 GI = 8 D2 Gq = 9 1 D2 WT outlier value was removed with Grubbs' method (Alpha = 0.01)	Fat mass variation following CNO period	F(2.00, 10.99)=5.966 p=0.0176 Dunnett's T3 multiple comparisons test D1 WT vs. D1 Gi p=0.0026 D1 WT vs. D1 Gq p=0.0447
Figure 4D	One way ANOVA	D2 WT = 9 D2 GI = 8 D2 Gq = 9	Weight variation following CNO removal	F(2, 23)=12.67 p=0.0002 Bonferroni's multiple comparisons test: D2WT vs D2Gq p=0.0002 D2Gi vs D2Gq p=0.0040
Figure 4D	One way ANOVA	D2 WT = 9 D2 GI = 8 D2 Gq = 9	Fat mass variation following CNO removal	F(2, 23)=8.537 p=0.0017 Bonferroni's multiple comparisons test: D2WT vs D2Gq p=0.0020 D2Gi vs D2Gq p=0.0184
Figure 4E	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 WT = 12 D1 GI = 11 D1 Gq = 12 2 outlier values removed with Grubbs' method (Alpha = 0.01)	High fat consumption D1-9; D24-25	Time effect F(3.418, 108.4)=34.19 p<0.0001 (ε correction: 0.4883) Virus effect F(2, 32)=1.470 p=0.2451 Interaction F(14.222)=6.702 p<0.0001 Bonferroni's multiple comparisons test: D1Gq vs D1WT: Day1 p=0.0183; Day3 p=0.0771; Day5 p=0.0612; Day7 p=0.0411 D1Gq vs D1Gi: Day1 p=0.0022; Day5 p=0.0096; Day24 p=0.0082; Day25 p=0.0558

Figure 4E	One way ANOVA	D1 WT = 12 D1 GI = 11 D1 Gq = 12	High fat consumption mean D10-D22	F(2, 32)=6.579 p=0.0040 Bonferroni's multiple comparisons test: D1Gq vs D1WT: p=0.0152 D1Gq vs D1Gi: p=0.0084
Figure 4E	One way ANOVA	D1 WT = 12 D1 GI = 11 D1 Gq = 12	High fat consumption mean D26-D33	F(2, 32)=2.153 p=0.1327
Figure 4F	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 WT = 9 D1 GI = 11 D1 Gq = 11 (3 WT and 1 D1-Gq animals were exposed to a running wheel without recording)	Running wheel variation D1-9; D24-25	Time effect F(4.651, 128.9)=10.12 p<0.0001 (ε correction: 0.2736) Virus effect F(2, 28)=11.07 p=0.0003 Interaction F(34, 471)=3.334 p<0.0001 Bonferroni's multiple comparisons test: D1Gq vs D1WT: Day1 p=0.0531; Day3 p=0.0223; Day5 p=0.0129; Day7 p=0.0218; Day9 p=0.0213; Day11 p=0.0086; Day13 p=0.0290; Day15 p=0.0285 D1Gi vs D1WT: Day1 p=0.0299; Day17 p=0.0187; Day19 p=0.0703; Day21 p=0.0174; Day25 p=0.0223; Day27 p=0.0770; Day29 p=0.0090; Day31 p=0.0325; Day33 p=0.0808
Figure 4G	Brown-Forsythe ANOVA test	D1 WT = 12 D1 GI = 11 D1 Gq = 12	Weight variation following CNO period	F (2.00, 20.08) =3.861 P=0.0381 Dunnett's T3 multiple comparisons test D1 WT vs. D1 Gi p=0.3297 D1 WT vs. D1 Gq p=0.4184 D1 Gi vs. D1 Gq p=0.0826
Figure 4G	One way ANOVA	D1 WT = 12 D1 GI = 11 D1 Gq = 12	Fat mass variation following CNO period	F (2, 32) =3.492 P=0.0425 Bonferroni's multiple comparisons test: D1Gi vs D1Gq p=0.0463
Figure 4G	Brown-Forsythe ANOVA test	D1 WT = 12 D1 GI = 11 D1 Gq = 12 1 D1 Gq and 1D1 GI outlier value were removed with Grubbs' method (Alpha = 0.01)	Weight variation following CNO removal	F (2.00, 17.99) =6.066 P=0.0097 Dunnett's T3 multiple comparisons test D1 WT vs. D1 Gi p=0.9612 D1 WT vs. D1 Gq p=0.0044 D1 Gi vs. D1 Gq p=0.0478
Figure 4G	Kruskal-Wallis test	D1 WT = 12 D1 GI = 11 D1 Gq = 12 1 D1 Gq and 1D1 GI outlier value were removed with Grubbs' method (Alpha = 0.01)	Fat mass variation following CNO removal	Kruskal-Wallis statistic: 9.225; p=0.0099 Dunn's multiple comparisons test D1WT vs D1Gq p=0.0139 D1Gi vs D1Gq p=0.0544
Figure 5				
Figure 5B	Mixed-effects model (Geisser-	D1-D2 WT = 11 D1Gq-D2Gi = 8	High fat consumption	Time effect F(4.272, 115.3)=32.84 p<0.0001 (ε correction: 0.6103)

	Greenhouse's epsilon correction)	D1Gi-D2Gq = 11	D1-9; D24-25	Virus effect $F_{(2, 27)}$=0.3648 p=0.6977 Interaction $F_{(14, 189)}$=7.911 p<0.0001 Bonferroni's multiple comparisons test: WT vs D1GqD2Gi: Day1 p=0.0498; Day24 p=0.0056; Day25 p=0.0769 WT vs D1GiD2Gq: Day3 p=0.0648; Day24 p=0.0967 D1GiD2Gq vs D1GqD2Gi: Day1 p=0.0057; Day3 p=0.0277; Day24 p=0.0004; Day25 p=0.0001
Figure 5B	One way ANOVA	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	High fat consumption mean D10-22	$F_{(2, 27)}$=1.880 p=0.1720
Figure 5B	One way ANOVA	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	High fat consumption mean D26-33	$F_{(2, 27)}$=1.425 p=0.2580
Figure 5C	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1-D2 WT = 9 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9 (2 WT and 2 D1Gq-D2Gi animals were exposed to a running wheel without recording) 6 outlier values removed with Grubbs' method (Alpha = 0.01)	Running wheel variation	Time effect $F_{(2.425, 53.64)}$=2.841 p=0.0573 (ε correction: 0.1427) Virus effect $F_{(2, 23)}$=8.839 p=0.0014 Interaction $F_{(34, 376)}$=3.486 p<0.0001 Bonferroni's multiple comparisons test: WT vs D1Gq-D2Gi: Day1 p=0.0791; Day3 p=0.0122; Day5 p=0.0013; Day7 p=0.0148; Day9 p=0.0105; Day13 p=0.0106; Day15 p=0.0164; Day17 p=0.0059; Day19 p=0.0010 D1Gi-D2Gq vs WT: Day1 p=0.0966; Day3 p=0.0600; Day5 p=0.0005; Day11 p=0.0336; Day13 p=0.0039; Day15 p=0.0331; Day17 p=0.0047; Day19 p=0.0282; Day21 p=0.0568; Day23 p=0.0443
Figure 5D	One way ANOVA	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	Weight variation (D1-D23)	$F_{(2, 27)}$=9.421 p=0.0008 Bonferroni's multiple comparisons test: WT vs D1Gi-D2Gq: p=0.0638 D1Gq-D2Gi vs D1Gi-D2Gq p=0.0006
Figure 5D	Kruskal-Wallis test	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11 1 D1Gq-D2Gi outliers value was removed with Grubbs' method (Alpha = 0.01)	Fat mass variation (D1-D23)	Kruskal-Wallis statistic: 16.74 p=0.0002 Dunn's multiple comparisons test WT vs D1Gq-D2Gq p=0.0408 D1Gq-D2G vs D1Gi-D2Gq p<0.0001
Figure 5D	One way ANOVA	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	Weight variation (D23-D36)	$F_{(2, 27)}$=19.79 p<0.0001 Bonferroni's multiple comparisons test: WT vs D1Gi-D2Gq p=0.0004 D1Gq-D2Gi vs D1Gi-D2Gq p<0.0001
Figure 5D	One way ANOVA	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	Fat mass variation (D23-D36)	$F_{(2, 27)}$=52.97 p<0.0001 Bonferroni's multiple comparisons test: WT vs D1Gi-D2Gq p=0.0171 WT vs D1Gq-D2Gi p<0.0001

				D1Gq-D2Gi vs, D1Gi-D2Gq p<0.0001
Figure 6				
Figure 6A	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	Weight variation (Food restricted procedure)	Time effect F(1.421, 38.36)=348.2 p<0.0001 (ε correction: 0.2368) Virus effect F(2, 27)=4.600 p=0.0191 Interaction F(12, 162)=10.63 p<0.0001 Bonferroni's multiple comparisons test: WT vs D1Gq-D2Gi: 1.5g(Day5) p=0.0467; 1.5g(Day6) p=0.0212
Figure 6B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1-D2 WT = 9 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9 (2 WT and 2 D1Gq-D2Gi animals were exposed to a running wheel without recording) 1 outlier value removed with Grubbs' method (Alpha = 0.01)	Running wheel variation (Food restricted procedure)	Time effect F(3.503, 79.39)=1.644 p=0.1784 (ε correction: 0.5838) Virus effect F(2, 23)=7.484 p=0.0031 Interaction F(12, 136)=5.156 p<0.0001 Bonferroni's multiple comparisons test: WT vs D1Gq-D2Gi: 3g p=0.0027; 2.5g p=0.0085; 2g p=0.0429; 1.5(1) p=0.0318; 1.5(3) p=0.0648
Figure 6C	Brown-Forsythe ANOVA test	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11 1 D1Gi-D2Gq outlier value was removed with Grubbs' method (Alpha = 0.01)	Peripheral Leptin	F(2.00, 16.01)=9.434 p=0.0020 Dunnett's T3 multiple comparisons test WT vs D1Gq-D2Gi: p=0.0057 WT vs D1Gi-D2Gq: p=0.1179 D1Gq-D2Gi vs D1Gi-D2Gq p=0.0240
Figure 6C	One way ANOVA	D1-D2 WT = 9 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9 (2 D1-D2 WT and 2 D1Gi-D2Gq animals samples were not detected by Elisa kit) 1 D1Gq-D2Gi outlier value was removed with Grubbs' method (Alpha = 0.01)	Peripheral Ghrelin	F(2, 22)=4.870 p=0.0177 Bonferroni's multiple comparisons test: WT vs D1Gi-D2Gq p=0.0124
Figure 6C	One way ANOVA	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	Weight	F(2, 27)=4.982 p=0.0144 Bonferroni's multiple comparisons test: WT vs D1Gi-D2Gq p=0.0142
Figure 6D	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1-D2WT= 8 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9	Weight loss during ABA protocol	Time effect F(5, 105)=324.2 p<0.0001 (ε correction: 0.6460) group effect F(2, 22)=17.45 p<0.0001 Interaction F(10, 105)=7.702 p<0.0001 Bonferroni's multiple comparisons test: D1-D2WT vs D1Gq-D2Gi: D2 p=0.0046; D3 p=0.0312; D5 p=0.0220

				D1-D2WT vs D1GI-D2Gq: D2 p=0.0051; D3 p=0.0038; D4 p=0.0006; D5 p=0.0001
Figure 6E	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1-D2WT= 8 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9	Wheel running activity during ABA protocol	Time effect F(2.906, 61.29)=53.66 P<0.0001 (ε correction: 0.2642) group effect F(2, 22)=7.078 P:0.0042 Interaction F(22, 232)=9.935 P<0.0001 Bonferroni's multiple comparisons test: D1-D2WT vs D1Gq-D2Gi: D1 night p=0.0004 D2 night p=0.0051 D1-D2WT vs D1GI-D2Gq: D1 night p=0.0023; D2 night p=0.0001; D3 night p=0.0001; D4 night p=0.0190; D5 night p=0.0381
Figure 6F	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1-D2WT= 8 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9	Chow consumption during ABA protocol	Time effect F(2.492, 52.96)=192.8 p<0.0001 (ε correction: 0.6230) group effect F(2, 22)=2.496 p=0.1055 Interaction F(8, 85)=0.4240 p=0.9037
Figure 6G	Log-rank (Mantel-Cox) test	D1-D2WT= 8 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9	Survival Curve	D1-D2WT vs D1Gq-D2Gi: χ2= 2.358; p=0.1247 D1-D2WT vs D1Gi-D2Gq: χ2=1.125; p=0.2888 D1Gi-D2Gq vs D1Gq-D2Gi: χ2=5.583; p=0.0189
			Supplementary figure	
Supplementary Figure 1				
Figure Supp 1B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 GI = 9 D2 Gq = 7	Lever press (Learning)	Time effect F(1.121, 8.972)=139.2 p<0.0001 (ε correction: 0.3738) lever effect F(1.545, 12.36)=177.2 p<0.0001 (ε correction: 0.5151) Interaction F(1.299, 7.508)=46.80 p<0.0001 (ε correction: 0.1444) Bonferroni's multiple comparisons test: Reinforced Lever (L1): D1 Gi vs D1 Gq p=0.7850 Non reinforced Lever (L2): D1 Gi vs D1 Gq p=0.5598
Figure Supp 1C	Two-way RM ANOVA	D2 Gi = 8 D2 Gq= 7	Lick/burst	Virus effect F(1, 13)=0.04570 p=0.8340 CNO effect F(1, 13)=6.896 p=0.0209 Interaction F(1, 13)=0.03268 P=0.8593
Figure Supp 1D	Two-way RM ANOVA	D2 GI =8 D2 Gq = 7	RPM	Virus effect F(1, 13)=1.871 p=0.1946 CNO effect F(1, 13)=1.976 p=0.1833 Interaction F(1, 13)=0.3563 p=0.5608
Figure Supp 1E	Two-way RM ANOVA	D2 GI =9 D2 Gq = 8	Distance	Virus effect F(1, 15)=11.92 =0.0036 CNO effect F(1, 15)=1.319 p=0.2687 Interaction F(1, 15)=28 p=0.0001 Bonferroni's multiple comparisons test: D2Gi: Sal vs CNO p=0.0006 D2Gq: Sal vs CNO p=0.0245

Figure	Two-way RM ANOVA	D2 GI =9	Mean speed	Virus effect F(1, 15)=11.91 p=0.0036
Supp 1E		D2 Gq = 8		CNO effect F(1, 15)=1.309 p=0.2704
				Interaction F(1, 15)=28.02 p=0.0001
				Bonferroni's multiple comparisons test:
				D2Gi: Sal vs CNO p=0.0006
				D2Gq: Sal vs CNO p=0.0243
Figure	Mixed-effects model	D1 GI = 9	Lever press	Time effect F(1.104, 8.833)=103.7 p<0.0001
Supp 1F	(Geisser-Greenhouse's epsilon correction)	D1 Gq = 8	(Learning)	(ε correction: 0.3680)
				lever effect F(1.555, 12.44)=110.4 p<0.0001
				(ε correction: 0.5184)
				Interaction F(1.925, 13.69)=45.22 <0.0001
				(ε correction: 0.2139)
				Bonferroni's multiple comparisons test:
				Reinforced Lever (L1): D1 Gi vs D1 Gq p=0.3512
				Non reinforced Lever (L2): D1 Gi vs D1 Gq p>0.9999
Figure	Two-way RM ANOVA	D1 GI = 9	Lever presses	Virus effect F(1, 15)=11.59 p=0.0039
Supp 1G		D1 Gq = 8	CNO (2mg/kg)	CNO effect F(1, 15)=13.59 p=0.0022
				Interaction F(1, 15)=0.3135 p=0.5838
Figure	Two-way RM ANOVA	D1 GI = 9	Break point	Virus effect F(1, 15)=9.109 p=0.0086
Supp 1G		D1 Gq = 8	CNO (2mg/kg)	CNO effect F(1, 15)=17.29 p=0.0008
				Interaction F(1, 15)=0.00277 p=0.9587
Figure	Mixed-effects model	D1 Gi = 9	Rate (presses/s)	Ratio effect F(2.634, 21.07)=13.85 p<0.0001
Supp 1G	(Geisser-Greenhouse's epsilon correction)		CNO (2mg/kg)	(ε correction: 0.3292)
				CNO effect F(1.00, 8.00)=0.8356 p=0.3874
				(ε correction: 1.000)
				Interaction F(4.193, 32.49)=1.724 p=0.1664
				(ε correction: 0.5241)
Figure	Mixed-effects model	D1 Gq= 8	Rate (presses/s)	Ratio effect F(2.371, 16.60)=4.209 p=0.0281
Supp 1G	(Geisser-Greenhouse's epsilon correction)		CNO (2mg/kg)	(ε correction: 0.3534)
				CNO effect F(1.000, 7.000)=15.63 p=0.0055
				(ε correction: 1.000)
				Interaction F(2.449, 16.65)=7.832 p=0.0039
				(ε correction: 0.2854)
				Bonferroni's multiple comparisons test:
				Saline vs CNO: ratio 16 p=0.0757; ratio 32 p=0.0165; ratio 64 p=0.0139
Figure	Two-way RM ANOVA	D1 GI = 9	Lever presses	Virus effect F(1, 16)=2.623 p=0.1249
Supp 1H		D1 Gq = 9	J60 (1mg/kg)	J60 effect F(1, 16)=5.102 p=0.0382
				Interaction F(1, 16)=0.2318 p=0.6367
Figure	Two-way RM ANOVA	D1 GI = 9	Break point	Virus effect F(1, 16)=5.789 p=0.0286
Supp 1H		D1 Gq = 9	J60 (1mg/kg)	J60 effect F(1, 16)=5.869 p=0.0277
				Interaction F(1, 16)=0.07715 p=0.7848
Figure	Two-way RM ANOVA	D1 GI =11	Distance	Virus effect F(1, 19)=9.705 p=0.0057
Supp 1I		D1 Gq = 10		CNO effect F(1, 19)=9.692 p=0.0057
				Interaction F(1, 19)=14.98 p=0.0010
				Bonferroni's multiple comparisons test:
				D1Gi: Sal vs CNO p=0.9999
				D1Gq: Sal vs CNO p=0.0002

Figure				
Figure Supp 1I	Two-way RM ANOVA	D1 GI =11 D1 Gq = 10	Mean speed	Virus effect F(1, 19)=9.675 p=0.0058 CNO effect F(1, 19)=9.690 p=0.0057 Interaction F(1, 19)=14.96 p=0.0010 Bonferroni's multiple comparisons test: D1Gi: Sal vs CNO p=0.9999 D1Gq: Sal vs CNO p=0.0002
Figure Supp 1J	Two-way RM ANOVA	D1 Gi = 11 D1 Gq=10	Lick/burst	Virus effect F(1, 19)=3.917 p=0.0625 CNO effect F(1, 19)=2.378 p=0.1395 Interaction F(1, 19)=0.9080 p=0.3526
Figure Supp 1K	Two-way RM ANOVA	D1 Gi = 8 D1 Gq = 8	RPM	Virus effect F(1, 14)=0.04791 p=0.8299 CNO effect F(1, 14)=1.774 p=0.2042 Interaction F(1, 14)=0.05004 p=0.8262
Supplementary Figure 2				
Figure Supp 2B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre mice Dreadd Gi = 7	Cumulative Chow consumption (1h - 4h)	Time effect F(1.672, 10.03)=2065 p<0.0001 (ε correction: 0.3345) CNO effect F(1.000, 6.000)=32.69 p=0.0012 (ε correction: 1.000) Interaction F(2.306, 13.84)=5.830 p=0.0122 (ε correction: 0.4612) Bonferroni's multiple comparisons test: DREADD Gi Sal vs CNO: 4h p=0.0084; 12h p=0.0171 24h p=0.0416; 48h p=0.0455
Figure Supp 2B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre mice Dreadd Gi = 7	Weight variation	Time effect F(1.000, 6.000)=19.95 p=0.0043 (ε correction: 1.000) CNO effect F(1.000, 6.000)=7.204 p=0.0363 (ε correction: 1.000) Interaction F(1.000, 6.000)=13.50 p=0.0104 (ε correction: 1.000) Šídák's multiple comparisons test: DREADD Gi Sal vs CNO: 24h p=0.0500; 48h p=0.2485
Figure Supp 2B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre mice Dreadd Gq = 8	Cumulative Chow consumption (1h - 4h)	Time effect F(1.278, 8.944)=975.3 p<0.0001 (ε correction: 0.2555) CNO effect F(1.000, 7.000)=9.034 p=0.0198 (ε correction: 1.000) Interaction F(2.182, 15.28)=21.06 p<0.0001 (ε correction: 0.4365) Bonferroni's multiple comparisons test: DREADD Gq Sal vs CNO: 48h p=0.0043
Figure Supp 2B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre mice Dreadd Gq = 8	Weight variation	Time effect F(1.0, 7.0)=0.008117 p=0.9307 (ε correction: 1.000) CNO effect F(1.000, 7.000)=26.40 p=0.0013 (ε correction: 1.000) Interaction F(1.000, 7.000)=1.854 p=0.2155 (ε correction: 1.000)
Figure Supp 2B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 cre mice Dreadd Gi = 9	Cumulative Chow consumption (1h - 48h)	Time effect F(1.874, 14.99)=1514 p<0.0001 (ε correction: 0.3747) CNO effect F(1.000, 8.000)=7.473 p=0.0257 (ε correction: 1.000) Interaction F(1.795, 14.36)=0.6985 p=0.4986 (ε correction: 0.3590)

Figure Supp 2B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 cre mice Dreadd Gi = 9	Weight variation	Time effect F(1.00, 8.00)=0.8960 p=0.3716 (ε correction: 1.000) CNO effect F(1.00, 8.00)=0.6924 p=0.4295 (ε correction: 1.000) Interaction F(1.000, 8.000)=3.569 p=0.0955 (ε correction: 1.000)
Figure Supp 2B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 cre mice Dreadd Gq = 9	Cumulative Chow consumption (1h - 48h)	Time effect F(1.994, 15.95)=1729 p<0.0001 (ε correction: 0.3987) CNO effect F(1.000, 8.000)=21.63 p=0.0016 (ε correction: 1.000) Interaction F(1.638, 13.10)=2.277 p=0.1473 (ε correction: 0.3275
Figure 2B	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 cre mice Dreadd Gq = 9	Weight variation	Time effect F(1.00, 8.00)= 42.30 p=0.0002 (ε correction: 1.000) CNO effect F(1.00, 8.00)=2.831 p=0.1310 (ε correction: 1.000) Interaction F(1.00, 8.00)=0.008763 p=0.9771 (ε correction: 1.00)
Figure Supp 2C	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre Dreadd Gi = 7	Fasting refeeding: High fat consumption (1h - 24h) CNO: 2mk/kg	Time effect F(1.652, 9.912)=899.7 p<0.0001 (ε correction: 0.4130) CNO effect F(1.000, 6.000)=15.96 p=0.0072 (ε correction: 1.000) Interaction F(1.969, 11.82)=16.90 p=0.0004 (ε correction: 0.4923) Bonferroni's multiple comparisons test: DREADD GI Sal vs CNO: 1h p=0.0012; 2h p=0.0027; 4h p=0.0057
Figure Supp 2C	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre Dreadd Gi =7	Fasting refeeding: High fat consumption (1h - 24h) CNO: 0.5mk/kg	Time effect F(1.515, 9.089)=484.68 p<0.0001 (ε correction: 0.3787) CNO effect F(1.00, 6.00)=0.1468 p=0.7148 (ε correction: 1.000) Interaction F(1.00, 7.00)=7.520 p=0.0210 Bonferroni's multiple comparisons test: DREADD GI Sal vs CNO: 1h p=0.028; 2h p=0.0099
Figure Supp 2C	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre Dreadd Gq = 8	Fasting refeeding: High fat consumption (1h - 24h) CNO: 2mk/kg	Time effect F(1.251, 8.754)=201.0 p<0.0001 (ε correction: 0.3126) CNO effect F(1.000, 7.000)=17.16 p=0.0043 (ε correction: 1.000) Interaction F(1.152, 8.067)=6.396 p=0.00321 (ε correction: 0.2881) Bonferroni's multiple comparisons test: DREADD Gq Sal vs CNO: 24h p=0.0370
Figure Supp 2C	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre Dreadd Gq = 7 (1 animal excluded)	Fasting refeeding: High fat consumption (1h - 24h) CNO: 0.5mk/kg	Time effect F(1.246, 7.478)=104.9 p<0.0001 (ε correction: 0.3116) CNO effect F(1.000, 6.000)=0.5173 p=0.4990 (ε correction: 1.000) Interaction F(2.218, 13.31)=1.949 p=0.1787 (ε correction: 0.5545)

Figure	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1 cre Dreadd Gi = 9	Fasting refeeding: High fat consumption (1h - 24h) CNO: 2mk/kg	Time effect F(1.181, 9.451)=414.5 p<0.0001 (ε correction: 0.2953) CNO effect F(1.00, 8.00)=0.3249 p=0.5843 (ε correction: 1.000) Interaction F(2.622, 20.32)=3.468 p=0.0400 (ε correction: 0.6556)
Figure Supp 2C	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1 cre Dreadd Gi = 8 1 outlier value removed with Grubbs' method (Alpha = 0.01)	Fasting refeeding: High fat consumption (1h - 24h) CNO: 0.5mk/kg	Time effect F(1.309, 9.163)=813.7 p<0.0001 (ε correction: 0.3272) CNO effect F(1.000, 7.000)=9.101 p=0.0195 (ε correction: 1.000) Interaction F(1.5658, 10.95)=3.844 p=0.0623 (ε correction: 0.3911) Bonferroni's multiple comparisons test: DREADD Gq Sal vs CNO: 1-24h p>0.9999
Figure Supp 2C	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1 cre Dreadd Gq = 8 2 outlier values removed with Grubbs' method (Alpha = 0.01)	Fasting refeeding: High fat consumption (1h - 24h) CNO: 2mk/kg	Time effect F(1.407, 9.851)=195.6 p<0.0001 (ε correction: 0.3518) CNO effect F(1.000, 7.000)=16.21 p=0.0050 (ε correction: 1.000) Interaction F(1.548, 10.06)=2.039 p=0.1836 (ε correction: 0.3870)
Figure Supp 2C	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1 cre Dreadd Gq = 9	Fasting refeeding: High fat consumption (1h - 24h) CNO: 0.5mk/kg	Time effect F(1.938, 15.50)=606.0 p<0.0001 (ε correction: 0.4844) CNO effect F(1.000, 8.000)=14.46 p=0.0052 (ε correction: 1.000) Interaction F(1.642, 13.14)=8.142 p=0.1406 (ε correction: 0.4106) Bonferroni's multiple comparisons test: DREADD Gq Sal vs CNO: 1h p=0.0080; 2h p=0.0111; 4h p=0.0072
Figure Supp 2D	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre mice Dreadd Gi = 7	% Blood glucose change vs baseline	Time effect F(2.576, 15.46)=104.2 p<0.0001 (ε correction: 0.3220) CNO effect F(1.00, 6.00)=1.070 p=0.3409 (ε correction: 1.000) Interaction F(3.077, 18.46)=0.4382 p=0.7330 (ε correction: 0.3846)
Figure Supp 2D	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 cre mice Dreadd Gq = 8	% Blood glucose change vs baseline	Time effect F(1.698, 11.89)=55.82 p<0.0001 (ε correction: 0.2123) CNO effect F(1.00, 7.00)=0.01822 p=0.8964 (ε correction: 1.000) Interaction F(2.704, 18.93)=1.226 p=0.3380 (ε correction: 0.3380)
Figure Supp 2D	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 cre mice Dreadd Gi = 9	% Blood glucose change vs baseline	Time effect F(1.901, 15.21)=74.77 p<0.0001 (ε correction: 0.2376) CNO effect F(1.00, 8.00)=0.1887 p=0.6754 (ε correction: 1.000) Interaction F (2.197, 17.58)=0.3944 p=0.6987 (ε correction: 0.2747)
Figure Supp 2D	Mixed-effects model	D1 cre mice Dreadd Gq = 9	% Blood glucose change vs baseline	Time effect F(2.193, 17.54)=130.8 p<0.0001 (ε correction: 0.2741) CNO effect F(1.00, 8.00) = 2.419 p=0.1585

				(ε correction: 1.000)
				Interaction F(3.290, 26.32) = 1.257 p=0.3601
				(ε correction: 0.4113)
Figure Supp 2E	Mann Whitney test	D2cre mice (n=6), Neurons: Saline:42 / CNO:44	Firing rate (Hz)	U=712, p= 0.0612
Figure Supp 2E	Mann Whitney test	D2cre mice (n=6), Neurons: Saline:42 / CNO:44	Bursting rate (Hz)	U=579.5, p=0.0026
Figure Supp 2E	Mann Whitney test	D2cre mice (n=6), Neurons: Saline:42 / CNO:44	%SIB	U=627, p=0.0097
Figure Supp 2E	Mann Whitney test	D2cre mice (n=6), Neurons: Saline:42 / CNO:44	MSB	U=791, p=0.2369
Figure Supp 2E	Unpaired T-test	D1cre mice (n=4), Neurons: Saline:40 / CNO:39	Firing rate (Hz)	t(77)=3.888, p=0.0002
Figure Supp 2E	Mann Whitney test	D1cre mice (n=4), Neurons : Saline:40 / CNO:39	Bursting rate (Hz)	U=530.5, p=0.0133
Figure Supp 2E	Mann Whitney test	D1cre mice (n=4), Neurons : Saline:40 / CNO:39	%SIB	U=634.5, p= 0.1534
Figure Supp 2E	Unpaired T-test	D1cre mice (n=4), Neurons : Saline:40 / CNO:39	MSB	t(77)=1.953, p=0.0545
Supplementary Figure 3				
Figure Supp 3C	Mixed-effect model (Satterthwaite correction for degrees of freedom)	D1GCamp = 5 D2GCamp = 5 12 trials per animals	Peri-event (Stop Run) (Zscore ; D1 vs D2)	Intercept effect F(1, 8.560)=5.056 p=0.0526 Genotype effect F(1, 8.560)=0.486 p=0.5044 Time effect F(1, 10.113)=2.3046 p=0.1596 Genotype x Time F(1, 10.11)=3.172 p=0.105 post hoc-permutation test between genotypes
Figure Supp 3D	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D2 cre mice Dreadd Gi = 7	Chow consumption	Time effect F(1.782, 26.72)=15.45 p<0.0001 (ε correction: 0.4454) CNO effect F(1.00, 60.00)=0.5417 p=0.4646 (ε correction: 1.000) Interaction F (1.533, 22.99)=4.968 p=0.0227 (ε correction: 0.3632) Bonferroni's multiple comparisons test: DREADD Gi Sal vs CNO: 3h p=0.0006
Figure Supp 3D	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D2 cre mice Dreadd Gq = 8	Chow consumption	Time effect F(1.922, 13.45)=3.763 p=0.0517 (ε correction: 0.4805) CNO effect F(1.000, 7.000)=2.435 p=0.1626 (ε correction: 1.000) Interaction F (1.647, 11.53)=1.052 p=0.3664 (ε correction: 0.4118)
Figure Supp 3E	Mixed-effects model	D2 cre mice Dreadd Gi = 7	Wheel running activity	Time effect F(1.894, 11.36)=11.19 p=0.0023 (ε correction: 0.4735)

				CNO effect F(1.000, 6.000)=17.85 p=0.0055 (ε correction: 1.000)
				Interaction F(1.582, 9.490)=4.728 p=0.0438 (ε correction: 0.3954)
				Bonferroni's multiple comparisons test: DREADD Gi Sal vs CNO: 3h p=0.0871; 5h p=0.0231
Figure Supp 3E	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D2 cre mice Dreadd Gq = 8 2 outlier values removed with Grubbs' method (Alpha = 0.01)	Wheel running activity	Time effect F(2.084, 14.59)=35.25 p<0.0001 (ε correction: 0.5210) CNO effect F(1.00 , 7.00)=14.38 p=0.0068 (ε correction: 1.000) Interaction F(1.762, 11.45)=9.475 p=0.0045 (ε correction: 0.4404) Bonferroni's multiple comparisons test: DREADD Gq Sal vs CNO: 3h p=0.0011; 4h p=0.0726
Figure Supp 3F	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1 cre Dreadd Gi = 9	Chow consumption	Time effect F(1.856, 14.84)=10.76 p=0.0015 (ε correction: 0.4639) CNO effect F(1.00, 8.00)=1.739 p=0.2238 (ε correction: 1.000) Interaction F(2.506, 20.06)=3.931 p=0.0287 (ε correction: 0.6269) Bonferroni's multiple comparisons test: DREADD Gq Sal vs CNO: 1-24h p>0.9999
Figure Supp 3F	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1 cre Dreadd Gq = 9 2 outlier values removed with Grubbs' method (Alpha = 0.01)	Chow consumption	Time effect F(2.569, 20.55)=21.55 p<0.0001 (ε correction: 0.6422) CNO effect F(1.00, 8.00)=6.794 p=0.0313 (ε correction: 1.000) Interaction F(2.150, 16.12)=6.184 p=0.0091 (ε correction: 0.5374) Bonferroni's multiple comparisons test: DREADD Gq Sal vs CNO: 3h p=0.0043
Figure Supp 3G	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1 cre Dreadd Gi = 9 2 outlier values removed with Grubbs' method (Alpha = 0.01)	Wheel running activity	Time effect F(2.393, 19.15)=48.19 p<0.0001 (ε correction: 0.5983) CNO effect F(1.000, 8.000)=2.106 p=0.1848 (ε correction: 1.000) Interaction F(2.518, 18.89)=5.949 p=0.0067 (ε correction: 0.6295) Bonferroni's multiple comparisons test: DREADD Gi Sal vs CNO: 3h p=0.0328
Figure Supp 3G	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1 cre Dreadd Gq = 9	Wheel running activity	Time effect F(1.858, 14.86)=1.387 p=0.2787 (ε correction: 0.4644) CNO effect F(1.000, 8.000)=12.14 p=0.0083 (ε correction: 1.000) Interaction F(1.420, 11.36)=10.31 p=0.0048 (ε correction: 0.3551) Bonferroni's multiple comparisons test: DREADD Gq Sal vs CNO: 3h p=0.0097; 4h p=0.0536; 5h p=0.0450
Supplementary Figure 4				

Figure	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D2 WT = 7 D2 GI = 7 D2 Gq = 8 (2 WT, 1 D1-Gi and 1 D1-Gq animals were exposed to a running wheel without recording)	Running wheel variation D1-33	Time effect $F_{(7.220, 136.5)}=3.247$ p=0.0029 (ϵ correction: 0.2188) Virus effect $F_{(2, 19)}=0.6835$ p=0.5168 Interaction $F_{(66, 624)}=2.884$ p<0.0001 Bonferroni's multiple comparisons test: D2Gi vs D2WT: Day1 p=0.0278;
Figure Supp 4B	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D2 WT = 9 D2 GI = 8 D2 Gq = 9	Total Mass	Time effect $F_{(1.15, 26.42)} = 16.03$ p=0.0003 (ϵ correction: 0.5744) Virus effect $F_{(2, 23)}=0.8456$ p=0.4422 Interaction $F_{(4, 46)}=1.247$ p=0.3047
Figure Supp 4C	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D2 WT = 9 D2 GI = 8 D2 Gq = 9	Fat Mass	Time effect $F_{(1.216, 27.96)}=40.10$ p<0.0001 (ϵ correction: 0.5744) Virus effect $F_{(2, 23)}=0.5713$ p=0.5726 Interaction $F_{(4, 46)}=1.448$ p=0.2336
Figure Supp 4D	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D2 WT = 9 D2 GI = 8 D2 Gq = 9	Lean Mass	Time effect $F_{(1.525, 35.08)}=116.5$ p<0.0001 (ϵ correction: 0.7625) Virus effect $F_{(2, 23)}=0.8900$ p=0.4243 Interaction $F_{(4, 46)}=2.186$ p=0.0853
Figure Supp 4E	One way ANOVA	D2 WT = 8 D2 GI = 8 D2 Gq = 9 1 outlier value removed with Grubbs' method (Alpha = 0.01)	Lean mass variation (mean D1-D23)	$F_{(2, 22)}=7.281$ p=0.0037 Bonferroni's multiple comparisons test: D2 WT vs D2 Gi p=0.0342 D2 WT vs D2 Gq p=0.0039
Figure Supp 4E	One way ANOVA	D2 WT = 9 D2 GI = 8 D2 Gq = 9	Lean mass variation (mean D23-D36)	$F_{(2, 23)}=3.737$ p=0.0393 Bonferroni's multiple comparisons test: D2 WT vs D2 Gq p=0.0825
Figure Supp 4F	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1 WT = 9 D1 GI = 11 D1 Gq = 11 (3 WT and 1 D1-Gq animals were exposed to a running wheel without recording)	Running wheel variation D1-33	Time effect $F_{(8.351, 228.5)}=5.553$ p<0.0001 (ϵ correction: 0.2531) Virus effect $F_{(2, 28)} = 0.0087$ p=0.0087 Interaction $F_{(66, 903)} = 2.061$ p<0.0001 Bonferroni's multiple comparisons test: D1Gq vs D1WT: Day2 p=0.0042; Day3 p=0.0045; Day4 p=0.0006; Day5 p=0.0151; Day6 p=0.0054; Day8 p=0.0197; Day9 p=0.0132; Day10 p=0.0059; Day11 p=0.0273; Day12 p=0.0052;
Figure Supp 4G	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D1 WT = 12 D1 GI = 11 D1 Gq = 12	Total Mass	Virus effect $F_{(2, 32)}=1.123$ p=0.3377 Time effect $F_{(1.534, 49.10)}=52.54$ p<0.0001 (ϵ correction: 0.7672) Interaction $F_{(4, 64)}=1.275$ p=0.2891
Figure Supp 4H	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D1 WT = 12 D1 GI = 11 D1 Gq = 12	Fat Mass	Virus effect $F_{(2, 32)} = 2.059$ p=0.1441 Time effect $F_{(1.366, 43.71)}=223.4$ p<0.0001 (ϵ correction: 0.6830) Interaction $F_{(4, 64)}=3.723$ p=0.0087 Bonferroni's multiple comparisons test:

				D1Gq vs D1WT: D23 p=0.0345;
Figure Supp 4I	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D1 WT = 12 D1 GI = 11 D1 Gq = 12	Lean Mass	Virus effect $F_{(2, 32)}$ = 2.498 p=0.0982 Time effect $F_{(1.262, 40.40)}$=122.9 p<0.0001 (ε correction: 0.6312) Interaction $F_{(4, 64)}$=1.547 p=0.1994
Figure Supp 4J	One way ANOVA	D1 WT = 12 D1 GI = 11 D1 Gq = 12	Lean mass variation (mean D1-D23)	$F_{(2, 32)}$=2.193 p=0.1280
Figure Supp 4J	One way ANOVA	D1 WT = 12 D1 GI = 11 D1 Gq = 12	Lean mass variation (mean D23-D36)	$F_{(2, 32)}$=0.7046 p=0.5018
Supplementary Figure 5				
Figure Supp 5A	Mixed-effects model (Geisser-Greenhouse's epsilon correction)	D1-D2 WT = 9 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9 (2 WT and 2 D1Gq-D2Gi animals were exposed to a running wheel without recording)	Running wheel variation D1-33	Time effect $F_{(5.387, 120.0)}$=3.491 p=0.0045 (ε correction: 0.1633) Virus effect $F_{(2, 23)}$=7.604 p=0.0029 Interaction $F_{(66, 735)}$=3.960 p<0.0001 Bonferroni's multiple comparisons test: D1-D2 WT vs. D1Gq-D2Gi: Day1 p=0.00152; Day3 p=0.0096; Day4 p=0.0410; Day5 p=0.0052; Day6 p=0.0111; Day7 p=0.0121; Day8 p=0.0113; Day9 p=0.0047; Day10 p=0.0379; Day11 p=0.0120; Day13 p=0.0140; Day14 p=0.0221; Day16 p=0.0055; Day17 p=0.0028; Day18 p=0.0002; Day20 p=0.0413; Day21 p=0.0149; Day22 p=0.0380;
Figure Supp 5B	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	Total Mass	Virus effect $F_{(2, 27)}$=5.959 p=0.0072 Time effect $F_{(1.313, 35.45)}$=14.92 p=0.0002 (ε correction: 0.6564) Interaction $F_{(4, 54)}$ = 5.141 p=0.0014 Bonferroni's multiple comparisons test: D1-D2 WT vs. D1Gq-D2Gi: D0 p=0.0010; D23 p=0.0023 D34 p=0.0176
Figure Supp 5C	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	Fat Mass	Virus effect $F_{(2, 27)}$=6.471 p=0.0051 Time effect $F_{(1.279, 33.88)}$=18.28 p=0.0001 (ε correction: 0.6393) Interaction $F_{(4, 53)}$=4.210 p=0.0049 Bonferroni's multiple comparisons test: D1-D2 WT vs. D1Gq-D2Gi: D0 p=0.0022; D23 p=0.0040 D34 p=0.0096 D1-D2 WT vs. D1Gi-D2Gq: D0 p=0.0038;

Figure Supp 5D	Two-way RM ANOVA (Geisser-Greenhouse's epsilon correction)	D1-D2 WT = 11 D1Gq-D2Gi = 8 D1Gi-D2Gq = 11	Lean Mass	Virus effect $F_{(2, 27)}=2.368$ p=0.1129 Time effect $F_{(1.260, 33.39)}=32.66$ p=0.0001 (ε correction: 0.6301) Interaction $F_{(4, 53)}=4.563$ p=0.0031 Bonferroni's multiple comparisons test: D1-D2 WT vs. D1Gq-D2Gi: D23 p=0.0308; D34 p=0.0437
Figure Supp 5E	One way ANOVA	D1-D2WT= 11 D1Gi-D2Gq = 8 D1Gq-D2Gi = 11	Lean mass variation (mean D1-D23)	$F_{(2, 32)}=6.872$ p=0.0039 Bonferroni's multiple comparisons test: D1-D2 WT vs D1Gi-D2Gq p=0.0948 D1Gq-D2Gi vs D1Gi-D2Gq p=0.0034
Figure Supp5E	One way ANOVA	D1-D2WT= 10 D1Gi-D2Gq = 8 D1Gq-D2Gi = 10 2 outlier values removed with Grubbs' method (Alpha = 0.01)	Lean mass variation (mean D23-D36)	$F_{(2, 25)}=15.07$ p<0.0001 Bonferroni's multiple comparisons test: D1-D2 WT vs D1Gi-D2Gq p=0.0104 D1Gq-D2Gi vs D1Gi-D2Gq p<0.0001
Supplementary Figure 6				
Figure Supp 6A	Two-way RM ANOVA	D1-D2WT= 8 D1Gq-D2Gi = 7 D1Gi-D2Gq = 7	Cumulative Chow consumption 2h under leptin or saline	Group effect $F_{(2, 19)}=92.61$ p<0.0001 Leptin effect $F_{(1, 19)}=14.83$ p=0.0011 Interaction $F_{(2, 19)}=0.8090$ p=0.4601
Figure Supp 6A	Two-way RM ANOVA	D1-D2WT= 8 D1Gq-D2Gi = 7 D1Gi-D2Gq = 7	Cumulative Chow consumption 24h under leptin or saline	Group effect $F_{(2, 19)}=19.20$ p=0.0543 Leptin effect $F_{(1, 19)}=54.72$ p<0.0001 Interaction $F_{(2, 19)}=3.410$ p<0.0001
Figure Supp 6B	Mixed-effects model	D1-D2WT= 8 D1Gq-D2Gi = 7 D1Gi-D2Gq = 7 1 outlier value removed with Grubbs' method (Alpha = 0.01)	Cumulative Chow consumption 2h under ghrelin or saline	Group effect $F_{(1, 18)}=30.26$ p<0.0001 Ghrelin effect $F_{(2, 19)}=80.60$ p<0.0001 Interaction $F_{(2, 18)}=16.60$ p<0.0001 Bonferroni's multiple comparisons test: Saline: D1-D2 WT vs D1Gq-D2Gi p=0.0032 D1-D2 WT vs D1Gi-D2Gq p>0.9999 D1Gq-D2Gi vs D1Gi-D2Gq p=0.0547 Ghrelin: D1-D2 WT vs D1Gq-D2Gi p<0.0001 D1-D2 WT vs D1Gi-D2Gq p=0.0025 D1Gq-D2Gi vs D1Gi-D2Gq p<0.0001
Figure Supp 6B	Two-way RM ANOVA	D1-D2WT= 8 D1Gq-D2Gi = 7 D1Gi-D2Gq = 7	Cumulative Chow consumption 24h under ghrelin or saline	Group effect $F_{(2, 19)}=9.446$ P=0.0014 Ghrelin effect $F_{(1, 19)}=6.859$ P=0.0169 Interaction $F_{(2, 19)}=2.327$ P=0.1247
Figure Supp 6C	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1-D2WT= 8 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9	Total Wheel running (m) activity during ABA protocol	Time effect $F_{(3.201, 67.22)}=10.35$ p<0.0001 (ε correction: 0.6402) group effect $F_{(2, 22)}=7.092$ p=0.0042 Interaction $F_{(10, 105)}=7.672$ p<0.0001 Bonferroni's multiple comparisons test:

				D1-D2WT vs D1Gq-D2Gi: D1 night p=0.0196
				D1-D2WT vs D1GI-D2Gq: D2 night p=0,0600; D3 night p=0,0900
Figure Supp 6D	Mixed-effects model (Geisser-Greenhouse's epsilon correction	D1-D2WT= 8 D1Gq-D2Gi = 8 D1Gi-D2Gq = 9	Wheel running variation during ABA protocol	Time effect F(2.856, 59.27)=13.67 p<0.0001 (ε correction: 0.7140) group effect F(2, 22)=13.22 p=0.0002 Interaction F(8, 83)=1.159 p=0.3336

1088

Chronic concommitant modulation of D1- and D2 neurons

A

hSyn-DIO-hm3d (Gq) x hSyn-FRT-hm4d (Gi)
or
hSyn-FRT-hm3d (Gq) x hSyn-DIO-hm4d (Gi)

D2-cre X D1-Flp →

D1-Flp / D2-Cre

Ac
D2-neuron
250 μm
Cre expressing neurons (green)

Ac
D1-neuron
250 μm
Flippase expressing neurons (red)

Ac
D1/D2
Coexpressing neuron
250 μm
Merge

B

CNO

■ D1-D2_WT ■ D1Gq-D2Gi ■ D1Gi-D2Gq

Food consumption (g)

D0 (chow) D1 D3 D5 D7 D9 Mean D10-22 D24 D25 Mean D26-33

C

CNO

Wheel running variation (%)

● D1-D2_WT ● D1Gq-D2Gi ▲ D1Gi-D2Gq

Basal D1 D3 D5 D7 D9 D13 D15 D17 D19 D21 D23 D25 D27 D9 D11 D33

D

CNO period

Total mass variation (%)

D1-D2_WT D1Gq-D2Gi D1Gi-D2Gq

Fat mass variation (%)

D1-D2_WT D1Gq-D2Gi D1Gi-D2Gq

CNO removal

Total mass variation (%)

D1-D2_WT D1Gq-D2Gi D1Gi-D2Gq

Fat mass variation (%)

D1-D2_WT D1Gq-D2Gi D1Gi-D2Gq

Food restricted procedure (Under CNO)

A — Weight variation (%) vs Food restriction (g). Legend: D1-D2 WT, D1Gq-D2Gi, D1Gi-D2Gq

B — Wheel running (m) vs Food restriction (g). Legend: D1-D2 WT, D1Gq-D2Gi, D1Gi-D2Gq

C — Leptin (ng/ml plasma), Ghrelin (pg/ml plasma), Weight (g). Groups: D1-D2 WT, D1Gq-D2Gi, D1Gi-D2Gq

Activity Based Anorexia (CNO infusion 30 min before dark onset)

D — Weight variation (%) vs Basal, D1, D2, D3, D4, D5. Legend: D1Gq-D2Gi, D1-D2 WT, D1Gi-D2Gq

E — Wheel running (m) vs Basal Night, Basal Day, Night 1, Day 1, Night 2, Day 2, Night 3, Day 3, Night 4, Day 4, Night 5, Day 5. Legend: D1-D2 WT, D1Gq-D2Gi, D1Gi-D2Gq

F — Chow consumption (g) vs Basal, D1, D2, D3, D4. Legend: D1-D2 WT, D1Gq-D2Gi, D1Gi-D2Gq

G — Probability of Survival vs Day. Legend: D1Gq-D2Gi, D1-D2 WT, D1Gi-D2Gq