Acoustic cues and season affect mobbing responses in a bird

community

<u>Short title:</u> cues and context affect heterospecific communication

Ambre Salis[1*], Jean-Paul Léna[1], Thierry Lengagne[1]

[1]Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-
69622, Villeurbanne, France

[*]Corresponding author: ambre.salis@univ-lyon1.fr

AS - ORCID : 0000-0003-0753-8981

JPL- ORCID : 0000-0002-2135-820X

TL- ORCID : 0000-0001-7840-6068

Address: Darwin C & Forel, 3-6 Rue Raphaël Dubois, 69622 Villeurbanne

Acknowledgments – Funding. This work was supported by the French Ministry of Research
and Higher Education funding (to A.S. PhD grants 2019-2022). We are grateful to Coline
Guignier and Lou Sayd for useful help with fieldwork. We thank the *Ligue de Protection des*
Oiseaux (LPO AuRA) for the information on the presence of coal and crested tits in the
territory.

Conflict of interest : The authors declare they have no conflict of interest relating to the
content of this article.

Ethics. All the authors follow the ASAB/ABS Guidelines for the Use of Animals in Research
and with The European Code of Conduct for Research Integrity. This work did not require
special permit.

Data statement. Raw data, R code, and Supplementary files are all added in Zenodo.
https://doi.org/10.5281/zenodo.7271850.

ABSTRACT

Heterospecific communication is common for birds when mobbing a predator. However, joining the mob should depend on the number of callers already enrolled, as larger mobs imply lower individual risks for the newcomer. In addition, some 'community informant' species seem more reliable regarding the information transferred in mobbing calls. Birds should therefore rely on both the number of callers and the species identity of the caller(s) when mobbing. In the present study, we tested the potential interaction between two acoustic cues (number of callers and the species identity of the caller). Overall, we found that soundtracks with three callers triggered more mobbing than soundtracks with one caller and that soundtracks with coal tits' calls triggered more mobbing than soundtracks with crested tits' calls. Our results therefore support the hypothesis that birds consider both the species and the number of callers when joining a mobbing chorus in winter. However, when the same experiments were replicated during spring, only soundtracks with three coal tits triggered a mobbing response, illustrating therefore how the seasonal context interacts can strongly affect the results of studies on heterospecific communication.

Keywords: heterospecific communication, mobbing call, community response, cooperation, seasonal effect

INTRODUCTION

Clustering around a predator and actively harassing it instead of fleeing is a widespread phenomenon termed 'mobbing'. Particularly common in birds (Carlson et al. 2018), mobbing encourages the predator to give up hunting and move to another location in both the short and long term (the Move-On Hypothesis, Curio 1978, Flasskamp 1994). Other benefits, such as monitoring the predators and enhancing learning opportunities for offspring, have been proposed (Curio 1978). Costs associated with such behavior are however non-negligible: in addition to the loss of time and energy when responding to an individual calling, the direct confrontation with a predator could result in direct aggression from the predator (Curio and Regelmann 1986, Poian and Yorke 1989, Sordahl 1990). Mobbing efficiency (i.e., the ratio of costs / benefits) can be improved by increasing the number of mobbing individuals (Krams et al. 2010, Wheatcroft and Price 2018). Indeed, larger groups decrease both the individual risk of being targeted by the predator (Hamilton selfish herd or dilution effect, Foster and Treherne 1981), and the overall success of the predator through confusion effect (Carlson et al. 2018). Larger groups also increase the chances of repelling the predator (Hendrichsen et al. 2006). Such an increase of participants can be achieved both with conspecific and heterospecific individuals, and heterospecific mobs are indeed well documented (e.g., Dutour et al. 2017a, Goodale and Kotagama 2005, Hua et al. 2016). Although heterospecific mobbing responses probably emerged as simple by-product mutualism (Kostan 2002), the relationships between species can be complex. Indeed, participation in such mobs is often unequal (Dutour et al. 2017b), with some species risking less by following the group at a distance (Magrath et al. 2015). In opposition, other species

seem particularly active and trustworthy regarding the information conveyed in the calls

(Farine et al. 2015). For such species that are active, reliable, and highly responded to, the

term 'community informant' has been proposed (Carlson et al. 2020).

The rationale to join mobbing birds should therefore depend on two main acoustic

cues: (i) the number of birds already mobbing, as a greater number of birds indicates a lower

risk for new participants, and (ii) the species identity of the caller(s), since some species

convey more reliable and relevant information than others. To test these hypotheses, we built

a set of playback experiments using a factorial design. We broadcast soundtracks of either one

or three coal tits (*Periparus ater*) and one or three crested tits (*Lophophanes cristatus*) to free-

ranging birds of both species, and recorded their behavioral response (calling and

approaching, the most conspicuous signs of mobbing in birds). Following a recent study

(Carlson et al. 2020), coal tits and crested tits contrast in their call reliability (i.e., coal tits

vary their calls when facing different threats) and heterospecific attraction when mobbing a

predator.

Heterospecific communication related to mobbing is prevalent in winter in passerines

communities (Dutour et al. 2019), notably because of an increased tendency to flock with

heterospecifics to increase predator defense and foraging efficiency (Goodale et al. 2015). We

therefore chose to test first and foremost birds during winter. Yet, we also replicated the same

experiment in spring to test whether the same rules apply during the reproductive season for

which interspecific communication is less prominent. Indeed, throughout the year, the

physical and social environment of birds varies greatly, possibly impacting their

communication. In spring, the increased aggressiveness due to territoriality and nest defense

could affect results on mobbing behavior (Betts et al. 2005, Jiang et al. 2020). By replicating this experiment in a different season, we test how environmental parameters such as season can affect our biological conclusions.

Our experiment therefore aims at determining the relative flexibility of heterospecific relationships and stability of acoustic cues throughout birds' seasonal activity. By looking at the mobbing response of both coal and crested tits to each other's calls, as much as the mobbing response of the overall community, we aim at determining how context affects the acoustic cues used by birds when investing in mobbing.

MATERIAL AND METHODS

General organization

We aimed at testing the mobbing response of free-ranging birds to different soundtracks. Playback tests were done in a 10 km² area of coniferous forest in the East of France (46°13'05.0"N 5°41'50.8"E). Densities of coal and crested tits are high in this area, as shown by the long-term ornithological census in the region: both species were detected in 94% of points, spaced at 150 m from each other (participative database Faune-ain.org administered by the LPO AuRA DT Ain). To circumvent inherent spatial variability, we established 100 spots for the playback tests. Each spot was selected close to a tree allowing birds' approach and concealment of experimenters, following existing trails. All spots were separated by ~ 100 m (mean and standard deviation X = 110.9 ± 27.2 m) since this distance is sufficient to degrade bird sounds (Morton 1975). In addition, we performed a complementary

subset of experiments (n = 22 birds tested) to verify that birds do not follow the observer between successive spots. For this purpose, we followed the same methodology than the one used by Salis et al. on great tits (2022). More specifically, both observers were equipped with the acoustic material and binoculars, and after each test, while one observer was launching the playback experiments on a subsequent location, the other was following the birds from the previous location. We found that from one test to the next one, no bird followed us, and no bird moved farther than 50 meters from their original position (see details in Supplementary File 1). While birds can travel large distances in a short period, it is unlikely that we tested the same birds in consecutive tests in the present experiment given the absence of human following and the absence of attraction from the subsequent playbacks.

We created a factorial design in which our four different treatments (different emitter species and number of callers, see paragraph Playbacks for details) were broadcast on each spot. These experiments were first carried out in winter, and then replicated in spring. Each spot consequently received eight playback tests. We avoided spatial and temporal autocorrelation by (i) alternating the four treatments at consecutive spots, and (ii) doing the same number of tests of each treatment, each day. The 400 tests in each season were done in a short period (two weeks) to avoid a potential intra-seasonal effect, and each consecutive test spaced by at least five minutes. We changed the order in which the spots were tested each day (different beginning point each day and different directions in the trails). Post hoc analyses (Sup. File 2) show no effect of order of playback treatment nor of the repeated presentation of playbacks on our results.

Playbacks

We created four treatments: soundtrack with only one calling coal tit (1CO), three coal tits calling simultaneously (3CO), only one calling crested tit (1CR), and lastly, three crested tits (3CR). We did not use a negative control (e.g., heterospecific song or background noise) since we were interested in the difference between our treatments. Moreover, background noise has been used in several studies (Dutour et al. 2019, Salis et al. 2022, Suzuki et al. 2016) and never triggered a response from Parids. To prepare our soundtracks, we elicited mobbing calls from wild crested tit and coal tit by broadcasting a mobbing chorus of various birds (including coal and crested tits, Dutour et al. 2016). Once birds arrived to mob they were recorded with a ME-67 Sennheiser microphone connected to a K6 basis and a Fostex FR2LE recorder. At last, the recordings were then cleared of any other bird call, their amplitude homogenized at 50% on the entire file with AvisoftSasLab (Avisoft Bioacoustics, Glienicke, Germany), and saved as WAV files. We selected recordings with a number of calls around the mean ($\pm$ 1 SD) of previous recordings obtained by our team (coal tit: 82 ± 26 notes per min, $N = 30$, crested tit: 134 ± 44 notes per min, $N = 10$). For the treatments with three birds (trio treatments), we superimposed recordings of three different birds calling to simulate a chorus. As a result, the final duty cycle (i.e., the amount of signal present in the playbacks) was higher for the three-birds treatment (~ 9 seconds) than for the one-bird treatments (~ 6.5 seconds, details in Sup. File. 3). Nevertheless, the calls substantially overlapped, reducing the risk for the focal birds to consider the three-birds treatments as only one bird calling intensely. For each treatment, we built five different soundtracks to circumvent the idiosyncrasy of recorded subjects (Kroodsma 1989).

Test procedure

One test consisted in playing 30 sec of a mobbing call sequence at each spot with a

Bose Soundlink Revolve loudspeaker perched on a tripod (H: 1m), put near a tree and at an

amplitude of 84.01 ± 2.70 dB (calculated at 1 m with Lutron SL-4001, C weighting, slow

settings, re. 20 µPa, Templeton et al., 2016). 30 sec is enough to trigger a mobbing response

from nearby birds (previous recordings were obtained with such a stimulation), who can

approach and call as a response, sometimes with additional aggressive behavior (e.g., wing

flicking, Salis et al. 2021). A stimulation of only 30 seconds also limited the influence of the

first birds to call on the following birds recruited. The two observers positioned themselves at

10 m from the tripod at vantage points before launching the soundtrack with an NW-A45

Sony. Before launching any test, we made sure that no bird was already in the vicinity nor

uttering mobbing calls in a distance. If a bird was detected, we waited only it left the area

(~10m around the loudspeaker). We observed the area with binoculars and all birds either

calling and/or approaching from the beginning of the test to 15 sec after the end of the

soundtrack. One bird was considered as approaching if it came in the 10 m radius around the

tripod (Dutour et al. 2017b). Only birds uttering specific and known mobbing calls (see Sup.

File 4 for spectrograms) were noted as calling. If a bird displayed the complete sequence of

mobbing behavior (i.e., simultaneously calling and approaching the loudspeaker), it was then

considered as giving a mobbing response. The two observers agreed on the lowest number of

birds seen simultaneously by both experimenters.

Statistical analyses

All statistical analyses were done with R studio (R v.3.6.1, R core team 2022).

Since the number of responding birds during the winter cannot be strictly compared to the one observed during the spring, analyses were done separately for each season. We used three response variables: the number of responding birds at the community level, the number of responding coal tits, and the number of responding crested tit. All these three response variables were thus analyzed separately for each season and following the same model framework. More specifically, for each response variable, we constructed an initial Hurdle mixed model. Hurdle models are two stage mixture models enabling to take into account excess of zero for count data (Zuur et al. 2009). This model framework is particularly suitable in our case since it involves a first process that determine the occurrence of an event (here the presence of at least one responding bird, i.e., mobbing occurrence) using a Binomial distribution, and a second one that determine the number of events (here when the number of responding birds is > 0, i.e. mobbing intensity) using a truncated count distribution. Such models are notably used in parasitology to segregate the effects on the host susceptibility and those affecting the infection intensity (e.g., Planade et al. 2009). We used a truncated Negative Binomial distribution to model the second stage (i.e. Hurdle Negative Binomial model, hereafter referred as HNB model) since preliminary analyses revealed that a truncated Poisson distribution performed less well than the HNB models, and the residuals of the latter did not indicate a lack of fit (function *testResiduals* of the package *DHARMa,* v. 0.4.5, Hartig and Hartig 2017). Both the effect of the emitter species, the one of the number of callers and their interactive effect were introduced in both fixed parts of the initial model. Moreover, both

the spot location and the soundtracks' ID were introduced as random effects in the model. The random effects were not added in the models analyzing the response of crested tits because of a general lower response precluding the correct estimation of the random effects. All initial full models were implemented using the package *glmmTMB* (v.1.1.2.3, Brooks et al. 2017), and were further reduced to construct all nested models (i.e., a set of 25 models including the full one for each response variable). Multimodel inference (Anderson and Burnham 2002) was used to select the best supported models for each response variable. For this task, we used the Bayesian Information Criterion (BIC) instead of the AIC since the former is more sensitive to the sample size than the latter (although both criteria gave a similar model selection, results not shown). We calculated the weighted BIC (wBIC) to calculate the evidence ratio between two models (i.e., calculate how much better is the best model compared to the other ones). The delta BIC (difference between two consecutive models) are given (a delta >2 is commonly considered to represent a significant difference between models). Effects sizes of the differences between treatments were calculated with odds ratios (OR).

RESULTS

Mobbing responses in winter

Eleven different species were attracted to our soundtracks (Figure 1A), with a maximum diversity of six species at one test. The two main species, apart from coal and crested tits, were the Goldcrest *(Regulus regulus,* present in 39% of our tests) and the Marsh

tit (*Poecile palustris*, 23%). As indicated by the best supported model (lowest BIC and an evidence ratio of 10, Table 1A), mobbing occurrence (the probability that at least one bird responded the playback), irrespective of the species (i.e., at the community level, Figure 2A) was affected by an additive effect of both the number of callers in the playback and the caller species (Table 1A). Indeed, birds mobbed more often the coal tit soundtracks compared to the crested tit soundtracks, and more to soundtracks with three birds rather than only one bird calling (1CO: 64%, 3CO: 77%, 1CR: 33%, 3CR: 59%). This additive effect was also detected when looking at the mobbing intensity (i.e., the number of mobbing birds when mobbing occurs, Figure 2B). Indeed, the largest mobs were initiated by playbacks with three coal tits (4.22 ± 3.65 birds, mean ± standard deviation) while the smaller mobs were initiated by playbacks with one crested tit (1.88 ± 1.24 birds).

When focusing on the response of coal tits or the one of the crested tits, the best supported model comprised an additive effect of the number of callers and the emitter species (Table 1B and 1C), resulting in a lower response toward singletons of crested tits (8% of points attracted coal tits or crested tits), intermediate scores toward trios of crested tits and singletons of coal tits, and the highest occurrence of response toward soundtracks with three coal tits (46% triggered a response from coal tits and 41% triggered a response from crested tits, Figure 2C and 2E). However, for the crested tit, the model with an interaction between number of callers and emitter species was also well supported (Delta < 2 and evidence ratio of 0.35/0.29 = 1.21, Table 1C). Indeed, the difference between 1CR and 3CR was higher (OR: 4.74, 95%CI: [1.92; 10.40]) than the difference between 1CO and 3CO (OR: 1.35, 95%CI: [0.76; 2.40]). Regarding mobbing intensity, no difference was detected between any of the

four playback treatments for the coal tit's response nor for the crested tit's response (Figure 2D and 2F).

Mobbing responses in spring

In spring, we detected a lower mobbing propensity: the occurrence of response to any treatment did not reach 25%, compared to almost 50% in winter. 15 different species were attracted to our soundtracks (Figure 1B), with a maximum diversity of five species at one test. The two most common species that responded were the Goldcrest (present in 24/% of our tests) and the Common Chaffinch (*Fringilla coelebs*, 18%). The mobbing occurrence at the community level was not impacted by any of the four playback treatments (Table 1D, Figure 3A). However, both the model with an interaction between number of callers and emitter species and the model with the number of callers alone were also well supported (similar BIC and wBIC, Table 1D). As a consequence, there is not strong evidence for any impact of either number of callers nor emitter species in our playbacks on the community response. The effect sizes depict a higher response towards the 3CO treatment than towards any of the three other playbacks (e.g., 3CO vs 3CR: 3.03, 95%CI: [1.70; 5.38]), while the three other playbacks triggered a similar response (e.g., 3CR vs 1CO: 1.14, 95%CI: [0.64; 2.05]). When focusing on the intensity of response (Figure 3B), only the emitter species impacted the size of the mobs, with coal tits' playbacks attracting more birds (respectively 1.69 ± 0.98 birds and 2.23 ± 1.51 for 1CO and 3CO, and 1.27 ± 0.63 birds and 1.44 ± 0.96 birds for 1CR and 3CR).

263   The best supported model regarding the presence of at least one coal tit included the

264 emitter species of the playback (Figure 3C, Table 1E), while the presence of at least one

265 crested tit was more impacted by the number of callers in the playback (Figure 3E, Table 1F).

266 We did not detect a difference in mobbing intensity between the four playbacks (Figure 3D

267 and 3F).

Figure 1. Number of spots (100 per acoustic treatment) in which at least one bird of each species mobbed (i.e., approached and called), when facing each of our four acoustic treatments (3CO: three coal tits, 1CO: one coal tit, 3CR: three crested tits, 1CR: one crested tit). Species taxonomy : Blue tit = *Cyanistes caeruleus,* Carrion crow = *Corvus corone,* Crested tit = *Lophophanes cristatus*, Coal tit = *Periparus ater*, Common blackbird = *Turdus merula,* Common chaffinch = *Fringilla coelebs*, Common chiffchaff = *Phylloscopus collybita*, Eurasian nuthatch = *Sitta europaea*, Eurasian wren = *Troglodytes troglodytes*, European blackcap = *Sylvia atricapilla,* European Robin = *Erithacus rubecula*, Goldcrest = *Regulus regulus*, Great tit = *Parus major*, Long-tailed tit = *Aegithalos caudatus,* Marsh tit = *Poecile palustris*, Mistle thrush = *Turdus viscivorus*, Treecreeper = *Certhia familiaris*, Willow tit = *Poecile montanus.*

Figure 2. Mobbing response of the bird community tested in winter to our four different mobbing soundtracks (1CO: one coal tit, 3CO: three coal tits, 1CR: one crested tit, 3CR: three crested tits). Error bars are 95% confidence intervals. Graphs on the left represent mobbing occurrence: the proportion of spots in which at least one bird mobbed (i.e., approach and called, $N = 100$ per treatment). Graphs on the right represent mobbing intensity: the number of birds that responded when there was a mobbing response (sample sizes are the proportion of the graphs on the left). The upper graphs are the response of the general bird community, middle graphs responses from coal tits, and lower graphs are response from crested tits.

Figure 3. Mobbing response of the bird community tested in a replication of the first experiment, during the reproductive season (spring). Birds' responses are recorded when facing four different mobbing soundtracks (1CO: one coal tit, 3CO: three coal tits, 1CR: one crested tit, 3CR: three crested tits). Error bars are 95% confidence intervals. Graphs on the left represent mobbing occurrence: the proportion of spots in which at least one bird mobbed (i.e., approach and called, $N = 100$ per treatment). Graphs on the right represent mobbing intensity: the number of birds that responded when there was a mobbing response (sample sizes are the proportion of the graphs on the left). The upper graphs are the response of the general bird community, middle graphs responses from coal tits, and lower graphs are response from crested tits.

Table 1. Hurdle models selection tables. A full mixed Hurdle model was constructed for each
response variable (number of responding birds at the community level, number of responding
coal tits, number of responding crested tits), and further reduced to construct the whole set of
reduced models. The full model included both the effect of the emitter species, the one of the
number of callers as well as their interactive effect in both fixed parts of the model (i.e.
mobbing occurrence modeled using a Binomial distribution, and mobbing intensity using a
truncated Negative Binomial distribution). We provide the three best supported models
according to the Bayesian Information Criterion (BIC), and the ones with the lowest BIC are
in bold. The weighted BIC (wBIC) represent the relative likelihood of our models. With
wBIC we can calculate evidence ratio between two models (e.g., the first model is 0.76 /
0.076 = 10 times more supported than the second model). The delta BIC (difference between
two consecutive models) are given (a delta >2 is commonly considered to represent a
significant difference between models).

	Occurrence (Presence of mobbers)	Intensity (Number of mobbers)	BIC	wBIC	Delta
Winter					
A. Community	**Emitter Species + Number of callers**	**Emitter Species + Number of callers**	1471.13	0.76	0
	Emitter Species + Number of callers	Emitter Species	1475.73	0.076	4.6
	Emitter Species * Number of callers	Emitter Species + Number of callers	1476.03	0.065	4.9
B. Coal tit	**Emitter Species + Number of callers**	**ø**	758.13	0.61	0
	Emitter Species + Number of callers	Number of callers	761.04	0.14	2.91
	Emitter Species + Number of callers	Emitter Species	761.56	0.11	3.43

C. Crested tit	**Emitter Species + Number of callers**	ø	732.47	0.35	0
	Emitter Species * Number of callers	ø	732.83	0.29	0.35
	Emitter Species + Number of callers	Emitter Species	735.36	0.08	2.89

Spring					
D. Community	**ø**	**Emitter Species**	962.94	0.18	0
	Number of callers	Emitter Species	963.17	0.17	0.21
	Emitter Species * Number of callers	Emitter Species	963.47	0.14	0.53
E. Coal tit	**Emitter Species**	ø	473.21	0.54	0
	ø	ø	475.71	0.16	2.5
	Emitter Species	Emitter Species	475.83	0.15	2.63
F. Crested tit	**Number of callers**	ø	281.35	0.75	0
	Number of callers	Emitter Species	285.37	0.10	4.03
	Emitter Species * Number of callers	ø	286.37	0.06	5.02

DISCUSSION

In winter, coal tits' soundtracks triggered more mobbing response from conspecifics and heterospecifics than crested tits' soundtracks; and soundtracks with three callers triggered more mobbing response from the bird community than soundtracks with only one caller. However, when replicating the experiment in spring, we found a lower general response but

also difference between playbacks, with increased responses only toward the 3 coal tits'

playbacks. This interaction between context and acoustic cues demonstrates the flexible

nature of heterospecific communication.

In winter, both the number of caller and emitter species influence mobbing responses

The mobbing response towards soundtracks with three individuals calling triggered

higher responses than soundtracks with only one individual calling. These results corroborate

the hypothesis that a greater number of birds mobbing a predator represents a lower risk for a

potential mobber, as it probably represents a lower risk of attack from the predator (confusion

and dilution effect, Sridhar et al. 2009). One previous study found similar results with the

Australian magpie (*Gymnorhina tibicen,* Igic et al. 2019), but with flee calls. Magpies flew

and scanned more in response to soundtracks with more individuals. Birds can consider

several acoustic criteria to assess the number of birds calling. First, birds can consider the

number of spatially different acoustic sources (Bradbury and Vehrencamp 2011). In our

study, we launched the soundtracks with only one loudspeaker whatever the treatment to

suppress this effect. Therefore, in our tests, the acoustic criteria that remain available are the

duty cycle (i.e., the proportion of the calling sequence when the signal is present), and the

count of calling individual through individual signatures. Our experiment does not add any

insights on which criteria was used by birds. Based on the current literature, the duty cycle is

probably one major coding strategy for increased risk in Parids (Landsborough et al. 2020,

Salis et al. 2022), and Parids modify their response to unknown non-Parids calls with different

duty cycles (Dutour et al. 2022). Regarding caller identity, great tits (*Parus major*) increased

their mobbing response toward soundtracks made with calls of several individuals compared to soundtracks with only one individual calling (Dutour et al. 2021). In this latter experiment, the duty cycles of the different treatments were strictly equal. This result was however not replicated when testing the response to different number of heterospecifics (chaffinches *Fringilla coelebs*, Dutour and Randler 2021). In our experiments, we believe that the overlapping of the calls in the three birds treatments avoid the risk of interpreting these treatments as only one bird calling intensely. However, experiments with non-overlapping calls and similar duty cycles could provide insight on this issue.

A mobbing response occurred more often when broadcasting coal tits' mobbing calls compared to crested tits' calls, but also more birds responded to it. Unexpectedly, even crested tits responded more to coal tits' mobbing calls than to calls from their own species. Coal tits therefore appear to be listened to and heavily responded to, leading to larger (and possibly more efficient) mobs. This is in line with the hypothesis that species from the same community show different levels of reliability (Magrath et al. 2015). The notion of "community informant" was developed for Parids in Carlson et al. (2020). They investigated whether the birds possessed a reliable way of encoding predator information, and if several heterospecifics relied on these calls. They showed that the great tit (*Parus major*) best fitted the definition of community informant. The coal tit approached the definition, with only one caveat: the dunnock (*Prunella modularis*) did not respond to it. As the authors suggested, the lack of response from one species does not mean that other species from the community do not respond to it (Carlson et al. 2020). Indeed, in our study, 14 species responded to coal tits' soundtracks. In opposition, the crested tit did not meet any of the criteria set by Carlson and

colleagues. Coal tits appear therefore to be one important species regarding predator

information in the community, and this is congruent with their increased sensibility to

predation by pygmy owls (*Glaucidium passerinum*) in winter (Suhonen et al. 1993).

Replicating the experiment in spring: A lower general response

In winter, Parids living in temperate regions often flock with heterospecifics,

sometimes leading to impressive mobs (up to 20 birds in the present experiment). In

opposition, during the reproductive period (May-July), Parids nest and defend their territory

with intensity (Hinde 1952). Spring territoriality may explain one of our results: we detected a

lower number of birds mobbing to the different soundtracks in spring. We here propose that in

spring, when all birds defend their territory, the number of birds that can respond is restricted

to the neighbors. Moreover, in spring, aggressivity toward conspecifics is high and may

therefore reduce the number of potential birds responding to mobbing calls. This aggressivity

may also explain why not so many birds responded to conspecific mobbing calls in spring

(coal tits to coal tits and crested tits to crested tits). Secondly, our results depict an apparent

decrease in mobbing occurrence in spring compared to winter. This difference must be taken

with cautiousness, as the community and the density of the population may vary with the

seasons: a decrease in mobbing response may simply be related to fewer individuals in the

territory. An order effect due to tests in winter being done before the tests in spring is unlikely

given the absence of order effect in our experiment at a short time scale (see Sup. Mat. For

details). In addition, in spring, we were able to hear coal tits singing at the 100 spots studied.

We are therefore confident that, in spring, each spot could have recorded one coal tit's

mobbing response. This suggests that at least for the coal tit, the response to conspecific and

heterospecific mobbing calls decreases in spring. This result is consistent with Dutour et al.

(2019) who detected in Parids a higher mobbing response toward heterospecific calls in

winter compared to summer. The proximal reasons for such a decrease can be various.

Increased territoriality and aggression in spring may very well limit cooperative

communication, since the mobbing calls may resemble intra-specific aggression/territoriality

calls, leading to a lower relevance for heterospecifics. Other factors such as decreased

predator pressure in spring (Dutour et al. 2017b) could also result in a lower investment in

mobbing in spring. The ratio cost/benefits in responding to distanced mobbing calls is

therefore probably flexible through different times of the year. Given that most of these

factors are intercorrelated, determining which one is responsible for the difference in mobbing

is unfeasible in natural conditions.

Replicating the experiment in spring: Almost no response to crested mobbing

calls

In addition to a general lower mobbing response in spring, the differences between

treatments were also impacted by the season. Indeed, while we selected similar models for the

community, coal tits', and crested tits' response in winter (additive effect of number of caller

and emitter species), we found different models in spring. A general tendency was detected,

with only the playbacks with three coal tits triggering more response than the three other

playbacks. This suggests that the crested tit is not considered as informative in spring, even

when mobbing in groups, and unexpectedly, even to conspecifics. Three explanations can be

proposed. Firstly, a group of three crested tits in spring may be too rare to bear meaning, as

they are in pairs and defending their nest. However, this hypothesis does not stand as this is

also the case for the coal tit, but that the difference between one and three callers still stands

in spring for this species. Alternatively, the contact with crested tits may be reduced in spring

if crested tits densities are lower during this season, hence decreasing learning opportunities

for heterospecifics. However, crested tits stay on the same territory throughout the year

(Ekman 1979) making this hypothesis unlikely despite our experiments do not allow us to

formally rule out it. We rather suggest that this lack of mobbing response may emerge from

reduced reliability of the calls. To be efficient, an acoustic signal needs to be easily

distinguishable from other signals (Bradbury and Vehrencamp 2011). The song and mobbing

calls of the crested tits are extremely similar (Cramp and Perrins 1993, Hailman 1989). As

crested tits produce both songs and mobbing calls in spring, we can hypothesize that the

global vocal production of crested tits therefore becomes less reliable from an external

individual, hence leading to a decreased response to such calls. In contrast, the coal tit appears

to be reliable and responded to in both seasons. This result is in adequacy with Jiang et al.

(2020) who also found that between seasons differences in playback responses did not affect

the nuclear status of some particular species (in their case, David's Fulvetta *Alcippe davidi*).

The difference between the response of the bird community to coal tits' playbacks compared

to crested tits' playbacks may also be due to a higher aggressiveness from crested tits. Crested

tits are known to be more aggressive during spring (Campbell 1958), and crested tits, larger

than coal tits, have higher rank dominance status (Suhonen et al. 1993). We have however

little data on whether the heterospecific aggressivity is higher than coal tits' aggressivity since

dominance status are not necessarily linked to increased aggressiveness (Wilson 1992).

==Finally, difference in nest predation may impact the reliability of the information produced,==

==but to our knowledge, nest predators are very similar between Parid species== (Cramp and

Perrins 1993).

To conclude, individuals from different species rely on several acoustic cues when

responding to conspecific or heterospecific mobbing calls. Both the number of callers and the

species calling are salient to the receivers, and those acoustic cues are not used in the same

way throughout the year. These results emphasize the importance of seasons in studies

investigating the complexity of heterospecific communication.

REFERENCES

Anderson DR, Burnham KP. 2002. Avoiding pitfalls when using information-theoretic

methods. J Wildl Manag. 66:912–918.

Betts MG, Hadley AS, Doran PJ. 2005. Avian Mobbing Response is Restricted by Territory

Boundaries☐: Experimental Evidence from Two Species of Forest Warblers. Ethology.

111:821–835. https://doi.org/10.1111/j.1439-0310.2005.01109.x

Bradbury JW, Vehrencamp SL. 1998. Principles of animal communication. Sunderland, MA:

Sinauer Associates.

Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ,

Maechler M, Bolker BM (2017). glmmTMB Balances Speed and Flexibility Among Packages

for Zero-inflated Generalized Linear Mixed Modeling. The R Journal. 9:378–400.

https://journal.r-project.org/archive/2017/RJ-2017-066/index.html

Broughton RK, Maziarz M, Hinsley SA. 2019. Social structure of Coal Tits *Periparus ater* in

temperate deciduous forest. J Ornithol. 160:117-126. https://doi.org/10.1007/s10336-018-

1594-4

Campbell, B. 1958. The crested tit. Forestry commission leaflet, Her Majesty's stationery

office.

Carlson NV, Healy SD, Templeton CN. 2018. Mobbing. Cur Biol. 28:R1081–R1082.

doi:10.1016/j.cub.2018.06.025

Carlson NV, Healy SD, Templeton CN. 2020. What makes a 'community informant'?

Reliability and anti-predator signal eavesdropping across mixed-species flocks of tits. Anim

Behav Cogn. 7:214–246. https://doi.org/10.26451/abc.07.02.13.2020

Cramp S, Perrins CM, Brooks DJ. 1993. Vol. VII: Flycatchers to shrikes. Oxford, UK:

Oxford University Press.

Cully JF, Ligon JD. 1986. Seasonality of Mobbing Intensity in the Pinyon Jay. Ethology

71:333–339. https://doi.org/10.1111/j.1439-0310.1986.tb00597.x

Curio E. 1978. The adaptive significance of avian mobbing: I Teleonomic hypotheses and

predictions. Z Tierpsychol. 48:175–183. https://doi.org/10.1111/j.1439-0310.1978.tb00254.x

Curio E, Regelmann K. 1986. Predator harassment implies a real deadly risk: A reply to

Hennessy Ethology. 72:75–78. https://doi.org/10.1111/j.1439-0310.1986.tb00607.x

Dutour M, Lena JP, Lengagne T. 2016. Mobbing behaviour varies according to predator

dangerousness and occurrence. Anim Behav. 119:119–124.

https://doi.org/10.1016/j.anbehav.2016.06.024

Dutour M, Léna JP, Lengagne T. 2017a. Mobbing calls: a signal transcending species boundaries. Anim Behav. 131:3–11. https://doi.org/10.1016/j.anbehav.2017.07.004

Dutour M, Lena JP, Lengagne T. 2017b. Mobbing behaviour in a passerine community increases with prevalence in predator diet. Ibis. 159:324–330. https://doi.org/10.1111/ibi.12461

Dutour M, Cordonnier M, Léna JP, Lengagne T. 2019. Seasonal variation in mobbing behaviour of passerine birds. J Ornithol. 160:509–514. https://doi.org/10.1007/s10336-019-01630-5

Dutour M, Fernández GJ, Randler C. 2022. How great tits respond to urgency□based information in allopatric Southern house wren mobbing calls. Ethology. 128: 676-683. https://doi.org/10.1111/eth.13329

Dutour M, Kalb N, Salis A, Randler C. 2021. Number of callers may affect the response to conspecific mobbing calls in great tits (*Parus major*). Behav Ecol Sociobiol. 75:1–8. https://doi.org/10.1007/s00265-021-02969-7

Dutour M, Randler C. 2021. Mobbing responses of great tits (*Parus major*) do not depend on the number of heterospecific callers. Ethology. 127:379-384. https://doi.org/10.1111/eth.13138

Ekman J. 1979. Coherence, composition and territories of winter social groups of the willow tit *Parus montanus* and the crested tit *P cristatus*. Ornis Scand. 10:56–68. https://doi.org/10.2307/3676345

Farine DR, Aplin LM, Sheldon BC, Hoppitt W. 2015. Interspecific social networks promote

information transmission in wild songbirds. Proc Royal Soc B. 282:20142804.

https://doi.org/10.1098/rspb.2014.2804

Flasskamp A. 1994. The adaptive significance of avian mobbing V An experimental test of

the 'move on'hypothesis. Ethology. 96:322–333. https://doi.org/10.1111/j.1439-

0310.1994.tb01020.x

Foster WA, Treherne JE. 1981. Evidence for the dilution effect in the selfish herd from fish

predation on a marine insect. Nature. 293:466–467. https://doi.org/10.1038/293466a0

Goodale E, Kotagama SW. 2005. Testing the roles of species in mixed-species bird flocks of

a Sri Lankan rain forest. J Trop Ecol. 21:669–676.

https://doi.org/10.1017/S0266467405002609

Goodale E, Ratnayake CP, Kotagama SW. 2014. Vocal Mimicry of Alarm-Associated Sounds

by a Drongo Elicits Flee and Mobbing Responses from Other Species that Participate in

Mixed-Species Bird Flocks. Ethology. 120:266–274. https://doi.org/10.1111/eth.12202

Goodale E, Ding P, Liu X, Martínez A, Si X, Walters M, et al. 2015. The structure of mixed-

species bird flocks, and their response to anthropogenic disturbance, with special reference to

East Asia. Avian Res. 6:14. https://doi.org/10.1186/s40657-015-0023-0

Hailman JP. 1989. The organization of major vocalizations in the Paridae. Wil Bull. 101:305–

343.

Hartig F, Hartig MF. 2017. Package 'DHARMa'. Vienna, Austria: R Development Core

Team.

Hendrichsen D, Christiansen PK, Nielsen E, Dabelsteen T, Sunde P. 2006. Exposure affects

the risk of an owl being mobbed–experimental evidence. J Avian Biol. 37:13–18.

https://doi.org/10.1111/j.2005.0908-8857.03658.x

Hinde, RA. 1952. The behaviour of the great tit (*Parus major*) and some other related species.

Behav Sup. III:1–201.

Hua F, Yong DL, Janra MN, Fitri LM, Prawiradilaga D, Sieving KE. 2016. Functional traits

determine heterospecific use of risk☐related social information in forest birds of tropical

South☐East Asia. Ecol Evol. 6:8485–8494. https://doi.org/10.1002/ece3.2545

Igic B, Ratnayake CP, Radford AN, Magrath RD. 2019. Eavesdropping magpies respond to

the number of heterospecifics giving alarm calls but not the number of species calling. Anim

Behav. 148:133–143. https://doi.org/10.1016/j.anbehav.2018.12.012

Jiang D, Sieving KE, Meaux E, Goodale E. 2020. Seasonal changes in mixed☐species bird

flocks and antipredator information. Ecol Evol. 10:5368–5382.

https://doi.org/10.1002/ece3.6280

Kostan KM. 2002. The evolution of mutualistic interspecific communication: Assessment and

management across species. J Comp Psychol. 116:206–9. https://doi.org/10.1037/0735-

7036.116.2.206

Krams I, Bērziņš A, Krama T, Wheatcroft D, Igaune K, Rantala MJ. 2010. The increased risk

of predation enhances cooperation. Proc Royal Soc B. 277:513–518.

https://doi.org/10.1098/rspb.2009.1614

Kroodsma DE. 1989. Suggested experimental designs for song playbacks. Anim Behav.

37:600–609. https://doi.org/10.1016/0003-3472(89)90039-0

Landsborough B, Wilson DR, Mennill DJ. 2020. Variation in chick-a-dee call sequences, not

in the fine structure of chick-a-dee calls, influences mobbing behaviour in mixed-species

flocks. Behav Ecol. 31:54–62. https://doi.org/10.1093/beheco/arz155

Magrath RD, Haff TM, Fallow PM, Radford AN. 2015. Eavesdropping on heterospecific

alarm calls: from mechanisms to consequences. Biol Rev. 90:560–586.

https://doi.org/10.1111/brv.12122

Morton ES. 1975. Ecological sources of selection on avian sounds. Am Nat. 109:17–34.

Planade B, Lena JP, Li H, Plénet S, Guegan JF, Thomas F, et al. 2009. Tracking a heterosis

effect in the field: tadpole resistance to parasites in the water frog hybridogenetic complex.

Parasitology. 136:1003–1013. https://doi.org/10.1017/S0031182009006489

Poian A, Yorke IM. 1989. Predator Harassment: More Evidence on the Deadly Risk.

Ethology. 83:167–169. https://doi.org/10.1111/j.1439-0310.1989.tb00526.x

R Core Team, 2022. R: A language and environment for statistical computing. Vienna,

Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Salis A, Léna JP, Lengagne T. 2022. Data & Supplemental information for manuscript

"Acoustic cues and season affect mobbing responses in a bird community "_ Review 1 PCI

(Version 2). Zenodo. https://doi.org/10.5281/zenodo.7271850

Salis A, Léna JP, Lengagne T. 2021. Great tits (*Parus major*) adequately respond to both

allopatric combinatorial mobbing calls and their isolated parts. Ethology. 127:213–222.

https://doi.org/10.1111/eth.13111

Salis A, Léna JP, Lengagne T. 2022. Which acoustic parameters modify the great tit's

response to conspecific combinatorial mobbing calls? Behav Ecol Sociobiol, 76:46.

https://doi.org/10.1007/s00265-022-03157-x

Searcy WA, Nowicki S. 2005.The evolution of animal communication: reliability and

deception in signalling systems. Princeton, NJ: Princeton University Press.

Sordahl TA. 1990. The Risks of Avian Mobbing and Distraction Behavior: An Anecdotal

Review. Wilson Bull. 102:349–352.

Sridhar H, Beauchamp G, Shanker K. 2009. Why do birds participate in mixed-species

foraging flocks? A large-scale synthesis. Anim Behav. 78:337–347.

https://doi.org/10.1016/j.anbehav.2009.05.008

Suhonen J, Halonen M, Mappes T. 1993. Predation Risk and the Organization of the Parus

Guild. Oikos. 66: 94–100. https://doi.org/10.2307/3545200

Suzuki TN, Wheatcroft D, Griesser M. 2016. Experimental evidence for compositional syntax

in bird calls. Nat Com. 7:10986. https://doi.org/10.1038/ncomms10986

Templeton CN, Zollinger SA, Brumm H. 2016. Traffic noise drowns out great tit alarm

calls. Cur Biol. 26:R1173-R1174. https://doi.org/10.1016/j.cub.2016.09.058

Templeton CN, Greene E, Davis K. 2005. Allometry of alarm calls: black-capped chickadees

encode   information   about   predator   size. Science. 308:1934–1937.

https://doi.org/10.1126/science.1108841

Van Duyse E, Pinxten R, Eens M. 2002. Effects of Testosterone on Song, Aggression, and

Nestling Feeding Behavior in Male Great Tits, *Parus major*. Hormones and Behavior.

41:178–186. https://doi.org/10.1006/hbeh.2001.1747

Wheatcroft D, Price TD. 2018. Collective action promoted by key individuals. Am Nat.

192:401–414. http://dx.doi.org/10.1086/698874

Wilson J. D. 1992. Correlates of agonistic display by great tits *Parus major*. Behaviour. 121:

168–214.

Wingfield JC, Ramenofsky M. 1985. Testosterone and aggressive behaviour during the

reproductive cycle of male birds. In: Gilles R, Balthazart J, editors. Neurobiology. Berlin,

Germany: Springer. p. 92-104.

Woods RD, Kings M, McIvor GE, Thornton A. 2018. Caller characteristics influence

recruitment to collective anti-predator events in jackdaws. Sci Rep. 8:1–8.

https://doi.org/10.1038/s41598-018-25793-y

Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. 2009. Mixed effects models and

extensions in ecology with R. New York: Springer.