Acoustic cues and season affect mobbing responses in a bird community

Ambre Salis[1*], Jean-Paul Léna[1], Thierry Lengagne[1]

[1]Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622, Villeurbanne, France

[*]Corresponding author: salis.ambre87@gmail.com

AS - ORCID : 0000-0003-0753-8981

JPL- ORCID : 0000-0002-2135-820X

TL- ORCID : 0000-0001-7840-6068

Address: Darwin C & Forel, 3-6 Rue Raphaël Dubois, 69622 Villeurbanne

Acknowledgments – Funding. This work was supported by the French Ministry of Research and Higher Education funding (to A.S. PhD grants 2019-2022). We are grateful to Coline Guignier and Lou Sayd for useful help with fieldwork. We thank the *Ligue de Protection des Oiseaux* (LPO AuRA) for the information on the presence of coal and crested tits in the territory.

Conflict of interest : The authors declare they have no conflict of interest relating to the content of this article.

Ethics. All the authors follow the ASAB/ABS Guidelines for the Use of Animals in Research and with The European Code of Conduct for Research Integrity. This work did not require special permit.

Data statement. Raw data, R code, and Supplementary files are all added in Zenodo. https://doi.org/10.5281/zenodo.7605316

ABSTRACT

Heterospecific communication is common for birds when mobbing a predator. However, joining the mob should depend on the number of callers already enrolled, as larger mobs imply lower individual risks for the newcomer. In addition, some 'community informant' species seem more reliable regarding the information transferred in mobbing calls. Birds should therefore rely on both the number of callers and the species identity of the caller(s) when mobbing. In the present study, we tested the potential interaction between two acoustic cues. In a playback experiment, we modified the number of callers (through an increased number of calling individuals correlated to an increased duty cycle) and the emitter species (crested tits versus coal tits). Overall, we found that soundtracks with three callers triggered more mobbing than soundtracks with one caller and that soundtracks with coal tits' calls triggered more mobbing than soundtracks with crested tits' calls. Our results therefore support the hypothesis that birds consider both the species and the number of callers when joining a mobbing chorus in winter. Finally, we replicated the experiment in spring and did not record the same responses from the bird community. Indeed, only soundtracks with three coal tits triggered a mobbing response, suggesting therefore that the seasonal context can affect the results of studies on heterospecific communication. The potential mechanisms implicated in the varying responses to different acoustic cues and different seasons are discussed and should deserve further investigations.

Keywords: heterospecific communication, mobbing call, community response, cooperation, seasonal effect

INTRODUCTION

Clustering around a predator and actively harassing it instead of fleeing is a widespread phenomenon termed 'mobbing'. Particularly common in birds (Carlson et al. 2018), mobbing encourages the predator to give up hunting and move to another location in both the short and long term (the Move-On Hypothesis, Curio 1978, Flasskamp 1994). Other benefits, such as monitoring the predators and enhancing learning opportunities for offspring, have been proposed (Curio 1978). Costs associated with such behavior are however non-negligible: in addition to the loss of time and energy when responding to an individual calling, the direct confrontation with a predator could result in direct aggression from the predator (Curio and Regelmann 1986, Poian and Yorke 1989, Sordahl 1990). Mobbing efficiency (i.e., the ratio of costs / benefits) can be improved by increasing the number of mobbing individuals (Krams et al. 2010, Wheatcroft and Price 2018). Indeed, larger groups decrease both the individual risk of being targeted by the predator (Hamilton selfish herd or dilution effect, Foster and Treherne 1981), and the overall success of the predator through confusion effect (Carlson et al. 2018). Larger groups also increase the chances of repelling the predator (Hendrichsen et al. 2006). Such an increase of participants can be achieved both with conspecific and heterospecific individuals, and heterospecific mobs are indeed well documented (e.g., Dutour et al. 2017a, Goodale and Kotagama 2005, Hua et al. 2016). Although heterospecific mobbing responses probably emerged as simple by-product mutualism (Kostan 2002), the relationships between species can be complex. Indeed, participation in such mobs is often unequal (Dutour et al. 2017b), with some species risking less by following the group at a distance (Magrath et al. 2015). In opposition, other species

seem particularly active and trustworthy regarding the information conveyed in the calls

(Farine et al. 2015). For such species that are active, reliable, and highly responded to, the

term 'community informant' has been proposed (Carlson et al. 2020).

The rationale to join mobbing birds should therefore depend on two main acoustic

cues: (i) the number of birds already mobbing, as a greater number of birds indicates a lower

risk for new participants, and (ii) the species identity of the caller(s), since some species

convey more reliable and relevant information than others. To test these hypotheses, we built

a set of playback experiments using a factorial design. We broadcast soundtracks of either one

or three coal tits (*Periparus ater*) and one or three crested tits (*Lophophanes cristatus*) to free-

ranging birds of both species, and recorded their behavioral response (calling and

approaching, the most conspicuous signs of mobbing in birds). Following a recent study

(Carlson et al. 2020), coal tits and crested tits contrast in their call reliability (i.e., coal tits

vary their calls when facing different threats) and heterospecific attraction when mobbing a

predator.

Heterospecific communication related to mobbing is prevalent in winter in passerines

communities (Dutour et al. 2019), notably because of an increased tendency to flock with

heterospecifics to increase predator defense and foraging efficiency (Goodale et al. 2015). We

therefore chose to test first and foremost birds during winter. Yet, we also replicated the same

experiment in spring to test whether seasonal context could influence experiments about

heterospecific communication. Indeed, throughout the year, the physical and social

environment of birds varies greatly, possibly impacting their communication (e.g., Clucas et

al. 2004, Jiang et al. 2020). In spring, the increased aggressiveness due to territoriality and

nest defense could affect results on mobbing behavior (Betts et al. 2005, Jiang et al. 2020). By replicating this experiment in a different season, we test how environmental parameters such as season can affect our biological conclusions.

Our experiment therefore aims at determining the relative flexibility of heterospecific relationships and stability of response to acoustic cues throughout birds' seasonal activity. By looking at the mobbing response of both coal and crested tits to each other's calls, as well as the mobbing response of the overall community, we aim at determining how context affects the acoustic cues used by birds when investing in mobbing.

MATERIAL AND METHODS

Study site & Species

The playbacks experiments described below were all done is the Haut-Bugey region, France. This region is a small mountain environment (altitude: ~800m), with mixed deciduous-coniferous forests. Densities of coal and crested tits are high in this area, as shown by the long-term ornithological census in the region: both species were detected in 94% of points, spaced at 150 m from each other (participative database Faune-ain.org administered by the LPO AuRA DT Ain). In this region, small birds are often predated by several predator including the Eurasian pygmy owl *Glaucidium passerinum*. Previous experiments in the region have shown a mobbing response from a large number of species, including the coal and the crested tits (Dutour et al. 2016; Dutour et al. 2017b). When mobbing occurs, birds

approach the predator cue and produce calls often with specific aggressive postures (e.g., wing flicks and frequent hops, but direct attack is rare (Carlson & Griesser, 2022).

General organization

We aimed at testing the mobbing response of free-ranging birds to different soundtracks. To this aim, we established 100 spots for the playback tests in a 10 km² area of coniferous forest in the East of France (46°13'05.0"N 5°41'50.8"E). Each spot was selected along an existing trail but close to a tree allowing birds' approach and concealment of experimenters. All spots were separated by ~ 100 m (mean and standard deviation: 110.9 ± 27.2 m) since this distance is sufficient to degrade bird sounds (Morton 1975). In addition, we performed a complementary subset of experiments (n = 22 birds tested, 9 crested tits and 13 coal tits) to verify that birds do not follow the observer between successive spots. For this purpose, we followed the same methodology than the one used by Salis et al. on great tits (2022). More specifically, both observers were equipped with the acoustic material and binoculars, and after each test, while one observer was launching the playback experiments on a subsequent location, the other was following the birds from the previous location. We found that from one test to the next one, no bird followed us, and no bird moved farther than 50 meters from their original position (see details in Supplementary File 1). While birds can travel large distances in a short period, it is unlikely that we tested the same birds in consecutive tests in the present experiment given the absence of human following and the absence of attraction from the subsequent playbacks.

We created a factorial design in which our four different treatments (different emitter

species and number of callers, see paragraph Playbacks for details) were broadcast on each

spot. These experiments were first carried out in winter, and then replicated in spring. Each

spot consequently received eight playback tests. We avoided spatial and temporal

autocorrelation by (i) alternating the four treatments at consecutive spots, and (ii) doing the

same number of tests of each treatment, each day. The 400 tests in each season were done in a

short period (two weeks) to avoid a potential intra-seasonal effect, and each consecutive test

spaced by at least five minutes (each consecutive test was at a different spot, so that each spot

was tested only once per day). We changed the order in which the spots were tested each day

(different beginning point each day and different directions in the trails). Post hoc analyses

(Sup. File 2) show no effect of order of playback treatment nor of the repeated presentation of

playbacks on our results.

Playbacks

We created four treatments: soundtrack with only one calling coal tit (1CO), three coal

tits calling simultaneously (3CO), only one calling crested tit (1CR), and lastly, three crested

tits (3CR). We did not use a negative control (e.g., heterospecific song or background noise)

since we were interested in the difference between our treatments. Moreover, background

noise has been used in several studies (Dutour et al. 2019, Salis et al. 2022, Suzuki et al.

2016) and never triggered a response from Parids. To prepare our soundtracks, we elicited

mobbing calls from wild crested tit and coal tit by broadcasting a mobbing chorus of various

birds (including coal and crested tits, Dutour et al. 2016). Once birds arrived to mob they were

recorded with a ME-67 Sennheiser microphone connected to a K6 basis and a Fostex FR2LE

recorder (recording distance of 5 m to 15 m). At last, the recordings were then cleared of any

other bird call, their amplitude homogenized at 50% on the entire file with AvisoftSasLab

(Avisoft Bioacoustics, Glienicke, Germany), and saved as WAV files. We selected recordings

with a number of calls around the mean ($\pm$ 1 SD) of previous recordings obtained by our team

(coal tit: 82 ± 26 notes per min, $N = 30$, crested tit: 134 ± 44 notes per min, $N = 10$). For the

treatments with three birds (trio treatments), we superimposed recordings of three different

birds calling to simulate a chorus. As a result, the final duty cycle (i.e., the amount of signal

present in the playbacks) was higher for the three-birds treatment ($\sim$ 9 seconds) than for the

one-bird treatments ($\sim$ 6.5 seconds, details in Sup. File. 3). Nevertheless, the calls

substantially overlapped, reducing the risk for the focal birds to consider the three-birds

treatments as only one bird calling intensely. For each treatment, we built five different

soundtracks to circumvent the idiosyncrasy of recorded subjects (Kroodsma 1989).

Test procedure

One test consisted in playing 30 sec of a mobbing call sequence at each spot with a

Bose Soundlink Revolve loudspeaker perched on a tripod (H: 1m), put near a tree and at an

amplitude of 84.01 ± 2.70 dB (calculated at 1 m with Lutron SL-4001, C weighting, slow

settings, re. 20 µPa, Templeton et al., 2016). 30 sec is enough to trigger a mobbing response

from nearby birds (previous recordings were obtained with such a stimulation), who can

approach and call as a response, sometimes with additional aggressive behavior (e.g., wing

flicking, Salis et al. 2021). A stimulation of only 30 seconds also limited the influence of the

first birds to call on the following birds recruited. The two observers positioned themselves at

10 m from the tripod at vantage points before launching the soundtrack with an MP4 player

(NW-A45 Sony). Before launching any test, we made sure that no bird was already in the

vicinity nor uttering mobbing calls in a distance. If a bird was detected, we waited only it left

the area (~10m around the loudspeaker). We observed the area with binoculars and all birds

either calling and/or approaching from the beginning of the test to 15 sec after the end of the

soundtrack. One bird was considered as approaching if it came in the 10 m radius around the

tripod (Dutour et al. 2017b). Only birds uttering specific and known mobbing calls (see Sup.

File 4 for spectrograms) were noted as calling. If a bird displayed the complete sequence of

mobbing behavior (i.e., simultaneously calling and approaching the loudspeaker), it was then

considered as giving a mobbing response. The two observers agreed on the highest number of

birds seen simultaneously by both experimenters.

Statistical analyses

All statistical analyses were done with R studio (R v.4.1.1, R core team 2022).

Since social conditions for our study species differ between winter and spring and

factors influencing rates of response presumably therefor differ, the analysis was done

separately for each season. We used three count response variables: the number of responding

birds of any species ("community level"), the number of responding coal tits, and the number

of responding crested tits. Given the high densities of both species in the study area, we

considered that the absence of responding birds is due to the absence of response (i.e.

structural zero) rather than the absence of bird (i.e. sampling zero). We therefore used Hurdle

mixed models which are more convenient than zero inflation models to handle an excess of

zeros of count data in such a situation (Zuur et al. 2009, Feng 2021). More specifically,

Hurdle models are two stage models using a Bernoulli probability mass function to treat the

zero outcomes as the result of a first process driving the occurrence of response (in our case,

the mobbing occurrence), and a left truncated probability mass function to treat the positive

outcomes as the result of a second process driving the response intensity (in our case the

intensity of mobbing). For each count variable, we first constructed an initial full hurdle

model implemented in the package *glmmTMB* (v.1.1.2.3*,* Brooks et al. 2017), with the effect

of the emitter species, the effect of the number of callers, and their interactive effect in both

parts of the model (occurrence and intensity). Moreover, both the spot location and the

soundtracks' ID were introduced as random effects as an intercept in the model. All models

were constructed with a quasi-Newton optimization method ('BFGS') to circumvent

convergence failure. Nevertheless, the random effects were discarded from the model when

analyzing the response of crested tits because of a general lower response precluding the

correct estimation of the random effects. In order to control for potential overdispersion in our

positive count data, we first selected between two alternative left truncated probability mass

functions to handle positive counts, a truncated Poisson distribution and a truncated negative

binomial one allowing the variance to increase more rapidly than the Esperance (note that we

tested both nbinom1 and nbinom2, the former having a linear parameterization and the second

having a quadratic parameterization, (Hardin & Hilbe 2007). For this purpose, both models

were constructed and compared using Bayesian Information Criterion (BIC) and AIC. Since

BIC is more sensitive to the sample size but less sensitive to the unobserved heterogeneity

than AIC (Brewer et al 2016), we only reported BIC. For the community response, a truncated negative binomial distribution led to the lowest BIC and was therefore chosen. Indeed, the dispersion parameter Θ (i.e., the inflation factor associated to the truncated negative binomial distribution: when $\Theta \to 0$, the distribution is closer to a Gamma distribution, while when $\Theta \to \infty$, the distribution is closer to a Poisson distribution) was 1.16 for the community model in winter and 0.79 for the community model in spring. For the isolated response of coal tits and crested tits, the truncated Poisson distribution led to the lowest BIC and was therefore chosen to analyze these responses. The fit of the structure selected for the initial model was then checked by the inspection of the its residuals using the package DHARMa (v 0.4.5, Hartig and Hartig 2017).

For each of the three response variables (at the level of community, crested tits and coal tits) and for each season, we then created four candidate models, each of them with all the explanatory terms of interest (number of callers and emitter species), but for which the interaction term was kept or not, in the occurrence part and the intensity part of the model. Weighted BIC (wBIC) was then computed for the four candidate models and used to assess and compare their relative support using evidence ratios (i.e., ratio of wBIC between two models, Anderson and Burnham 2002). Effects sizes of the differences between treatments were calculated with odds ratios (OR).

Ethical note

We used a sample size that is higher than in other recent studies (commonly around 20-30 tests per treatment) to circumvent common problems of lack of power in animal behavior studies, and because presence/absence data usually require larger sample sizes

RESULTS

Mobbing responses in winter

Eleven different species were attracted to our soundtracks (Figure 1A), with a

maximum diversity of six species at one test. The two main species, apart from coal and

crested tits, were the Goldcrest *(Regulus regulus,* present in 39% of our tests) and the Marsh

tit (*Poecile palustris*, 23%). As indicated by the best supported model (lowest BIC and an

evidence ratio of 12, Table 1A), mobbing occurrence (the probability that at least one bird

responded the playback), irrespective of the species (i.e., at the community level, Figure 2A)

was affected by an additive effect of both the number of callers in the playback and the caller

species (Table 2A). Indeed, birds mobbed more often the coal tit soundtracks compared to the

crested tit soundtracks, and more to soundtracks with three birds rather than only one bird

calling (1CO: 64%, 3CO: 77%, 1CR: 33%, 3CR: 59%). This additive effect was also detected

when looking at the mobbing intensity (i.e., the number of mobbing birds when mobbing

occurs, Figure 2B, Table 2A). Indeed, the largest mobs were initiated by playbacks with three

coal tits (4.22 ± 3.65 birds, mean ± standard deviation) while the smaller mobs were initiated

by playbacks with one crested tit (1.88 ± 1.24 birds).

When focusing on the occurrence of response of coal tits or the one of the crested tits,

the best supported model comprised an additive effect of the number of callers and the emitter

species (Table 1B and 1C, Table 2B and 2C), resulting in a lower response toward singletons

of crested tits (8% of points attracted coal tits or crested tits), intermediate scores toward trios

of crested tits and singletons of coal tits, and the highest occurrence of response toward

soundtracks with three coal tits (46% triggered a response from coal tits and 41% triggered a

response from crested tits, Figure 2C and 2E). However, for the crested tit, the model with an

interaction between number of callers and emitter species was also well supported (evidence

ratio of 0.51/0.43 = 1.19, Table 1C). Indeed, the difference between 1CR and 3CR was higher

(OR: 4.74, 95%CI: [1.92; 10.40]) than the difference between 1CO and 3CO (OR: 1.35,

95%CI: [0.76; 2.40]). Regarding mobbing intensity (Figure 2D and 2F), for both the coal tits'

and crested tits' response, the additive effect of number of callers and emitter species was less

stringent than for the occurrence of mobbing (the effect of emitter species for the coal tit, and

the effect of number of callers for the crested tits did not reach statistical significance when

reporting the estimates, Table 2B and 2C).

Mobbing responses in spring

In spring, we detected a lower mobbing propensity: the occurrence of response to any

treatment did not reach 25%, compared to almost 50% in winter. 15 different species were

attracted to our soundtracks (Figure 1B), with a maximum diversity of five species at one test.

The two most common species that responded were the Goldcrest (present in 24/% of our

tests) and the Common Chaffinch (*Fringilla coelebs*, 18%).

Regarding mobbing occurrence at the community level (Figure 3A), the model with

the lowest BIC was the one including an interaction between emitter species and number of

callers in the playbacks (Table 1D, Table 2D). Indeed, the effect sizes depict a higher

response towards the 3CO treatment than towards any of the three other playbacks (e.g., 3CO

vs 3CR: 3.03, 95%CI: [1.70; 5.38]), while the three other playbacks triggered a similar

response (e.g., 3CR vs 1CO: 1.14, 95%CI: [0.64; 2.05]). Note however that this interaction is

not strongly supported since the model including only the additive effects of number of callers

and emitter species gave a similar BIC (evidence ratio of 0.54/0.4 = 1.35, Table 1D). When

focusing on the intensity of response (Figure 3B), we detected no difference in the number of

birds recruited to the four different playbacks (Table 2D).

The best supported model regarding the presence of at least one coal tit included the

emitter species of the playback, but no effect of the number of callers (Figure 3C, Table 1E,

Table 2E). For the crested tit's occurrence, we recorded an interaction between the emitter

species and the number of callers in the playbacks (Figure 3E, Table 1F, Table 2F). Indeed,

our playbacks attracted more often crested tits when there were three coal tits in the playbacks

compared to any of the three other types of playbacks. For both species, the number of birds

recruited when mobbing occurred did not differ between the four types of playbacks (Figure

3D and 3F, Table 2E and 2F).

Figure 1. Number of spots (100 per acoustic treatment) in which at least one bird of each species mobbed (i.e., approached and called), when facing each of our four acoustic treatments (3CO: three coal tits, 1CO: one coal tit, 3CR: three crested tits, 1CR: one crested tit). Responses to each of the four treatments are stacked in sequence on each bar so that the entire bar represents the sum of all responses by a given species per treatment. Species taxonomy : Blue tit = *Cyanistes caeruleus,* Carrion crow = *Corvus corone,* Crested tit = *Lophophanes cristatus*, Coal tit = *Periparus ater*, Common blackbird = *Turdus merula,* Common chaffinch = *Fringilla coelebs*, Common chiffchaff = *Phylloscopus collybita*, Eurasian nuthatch = *Sitta europaea*, Eurasian wren = *Troglodytes troglodytes*, European blackcap = *Sylvia atricapilla,* European Robin = *Erithacus rubecula*, Goldcrest = *Regulus regulus*, Great tit = *Parus major*, Long-tailed tit = *Aegithalos caudatus,* Marsh tit = *Poecile palustris*, Mistle thrush = *Turdus viscivorus*, Treecreeper = *Certhia familiaris*, Willow tit = *Poecile montanus.*

Figure 2. Mobbing response of the bird community tested in winter to our four different mobbing soundtracks (1CO: one coal tit, 3CO: three coal tits, 1CR: one crested tit, 3CR: three crested tits). Error bars are 95% confidence intervals. Graphs on the left represent mobbing occurrence: the proportion of spots in which at least one bird mobbed (i.e., approach and called, $N = 100$ per treatment). Graphs on the right represent mobbing intensity: the number of birds that responded when there was a mobbing response (sample sizes are the proportion of the graphs on the left). The upper graphs are the response of the general bird community, middle graphs responses from coal tits, and lower graphs are response from crested tits.

Figure 3. Mobbing response of the bird community tested in a replication of the first experiment, during the reproductive season (spring). Birds' responses are recorded when facing four different mobbing soundtracks (1CO: one coal tit, 3CO: three coal tits, 1CR: one crested tit, 3CR: three crested tits). Error bars are 95% confidence intervals. Graphs on the left represent mobbing occurrence: the proportion of spots in which at least one bird mobbed (i.e., approach and called, $N = 100$ per treatment). Graphs on the right represent mobbing intensity: the number of birds that responded when there was a mobbing response (sample sizes are the proportion of the graphs on the left). The upper graphs are the response of the general bird community, middle graphs responses from coal tits, and lower graphs are response from crested tits.

Table 1. Hurdle models selection tables. For each response variable (number of responding birds at the community level, number of responding coal tits, number of responding crested tits), we first constructed a full Hurdle model with the effect of the emitter species, the effect of the number of callers as well as their interactive effect in both parts of the model (see material and method for details).   We compare this full model to models without the interaction in both the occurrence part and the intensity part of the model. We provide the Bayesian Information Criterion (BIC) and the weighted BIC (wBIC) to represent the relative support of each model. With wBIC we can calculate evidence ratio between two models (e.g., the first model is 0.86 / 0.07 ≈ 12 times more supported than the second model).

Occurrence (Presence of mobbers)	Intensity (Number of mobbers)	BIC	wBIC
WINTER			
A. Community			
Emitter Species + Number of callers	Emitter Species + Number of callers	1471,13	0,86
Emitter Species x Number of callers	Emitter Species + Number of callers	1476,03	0,07
Emitter Species + Number of callers	Emitter Species x Number of callers	1476,56	0,06
Emitter Species x Number of callers	Emitter Species x Number of callers	1481,46	0,004
B. Coal tit			
Emitter Species + Number of callers	Emitter Species + Number of callers	758,07	0,81
Emitter Species + Number of callers	Emitter Species x Number of callers	762,03	0,11
Emitter Species x Number of callers	Emitter Species + Number of callers	762,99	0,07
Emitter Species x Number of callers	Emitter Species x Number of callers	766,95	0,01
C. Crested tit			
Emitter Species + Number of callers	Emitter Species + Number of callers	731,95	0,51
Emitter Species x Number of callers	Emitter Species + Number of callers	732,3	0,43
Emitter Species + Number of callers	Emitter Species x Number of callers	737,25	0,04
Emitter Species x Number of callers	Emitter Species x Number of callers	737,61	0,03

	SPRING			

D. Community	Emitter Species x Number of callers	Emitter Species + Number of callers	964,13	0,54
	Emitter Species + Number of callers	Emitter Species + Number of callers	964,72	0,4
	Emitter Species x Number of callers	Emitter Species x Number of callers	969,86	0,03
	Emitter Species + Number of callers	Emitter Species x Number of callers	970,45	0,02
E. Coal tit	Emitter Species + Number of callers	Emitter Species + Number of callers	480,37	0,61
	Emitter Species + Number of callers	Emitter Species x Number of callers	482,59	0,2
	Emitter Species x Number of callers	Emitter Species + Number of callers	483,37	0,14
	Emitter Species x Number of callers	Emitter Species x Number of callers	485,6	0,05
F. Crested tit	Emitter Species x Number of callers	Emitter Species + Number of callers	289,76	0,55
	Emitter Species + Number of callers	Emitter Species + Number of callers	290,42	0,4
	Emitter Species x Number of callers	Emitter Species x Number of callers	295,66	0,03
	Emitter Species + Number of callers	Emitter Species x Number of callers	296,33	0,02

Table 2. Outputs of the models selected in bold in Table 1. Each Hurdle model is a two-stage model, the first one examining the effect of explanatory terms on the occurrence of response (mobbing occurrence) and the second one examining the effect of explanatory terms on the positive counts (mobbing intensity), see material and method for details. We provide the estimates with their standard error (the intercept is the 1CO treatment), the z value and the associated p-value. The detailed outputs of the three other models in Table 1 are added as supplementary material 5.

WINTER				

A. Community

Occurrence

	Estimate	SE	z	p
(Intercept)	-0,94	0,17	-5,5	< 0,0001
Emitter Species	1,11	0,22	4,95	< 0,0001
Number of Callers	0,63	0,16	4,04	< 0,0001

Intensity

	Estimate	SE	z	p
(Intercept)	0,97	0,12	7,83	< 0,0001
Emitter Species	-0,65	0,16	-3,94	< 0,0001
Number of Callers	-0,38	0,11	-3,3	0,0009

B. Coal tit

Occurrence

	Estimate	SE	z	p
(Intercept)	0,53	0,16	3,32	0,001
Emitter Species	1,28	0,26	4,96	< 0,0001
Number of Callers	0,63	0,18	3,57	0,0003

Intensity

	Estimate	SE	z	p
(Intercept)	0,4	0,11	3,47	0,001
Emitter Species	-0,44	0,23	-1,87	0,06
Number of Callers	-0,31	0,15	-2,03	0,04

C. Crested tit

Occurrence

	Estimate	SE	z	p
(Intercept)	0,53	0,15	3,55	0,0004
Emitter Species	1,03	0,24	4,33	< 0,0001
Number of Callers	0,51	0,17	3,06	0,002

Intensity

	Estimate	SE	z	p
(Intercept)	0,35	0,12	3,04	0,002
Emitter Species	-0,475	0,24	-2,01	0,04
Number of Callers	-0,276	0,15	-1,8	0,07

SPRING				

D. Community

Occurrence

	Estimate	SE	z	p
(Intercept)	0,09	0,15	0,59	0,56
Emitter Species	0,57	0,21	2,65	0,008
Number of Callers	0,9	0,22	4,16	< 0,0001
Emitter Species: Number of Callers	-0,81	0,3	-2,65	0,008

Intensity

	Estimate	SE	z	p
(Intercept)	-0,2	0,41	-0,5	0,62
Emitter Species	-12,2	933,47	-0,013	0,99
Number of Callers	-0,58	0,39	-1,49	0,14

E. Coal tit

Occurrence

	Estimate	SE	z	p
(Intercept)	1,36	0,18	7,74	< 0,0001
Emitter Species	0,9	0,3	3,02	0,003
Number of Callers	0,23	0,2	1,13	0,26

Intensity

	Estimate	SE	z	p
(Intercept)	-0,4	0,36	-1,11	0,27
Emitter Species	-0,97	0,63	-1,53	0,13
Number of Callers	-0,13	0,31	-0,41	0,69

F. Crested tit

Occurrence

	Estimate	SE	z	p
(Intercept)	3,09	0,52	5,95	< 0,0001
Emitter Species	-0,56	0,59	-0,95	0,34
Number of Callers	2,13	0,74	2,9	0,004
Emitter Species: Number of Callers	-1,82	0,83	-2,19	0,03

Intensity

	Estimate	SE	z	p
(Intercept)	-0,06	0,48	-0,12	0,9
Emitter Species	-1,3	0,88	-1,048	0,14
Number of Callers	0,51	0,61	0,84	0,4

DISCUSSION

In winter, coal tits' soundtracks triggered more mobbing response from conspecifics and heterospecifics than crested tits' soundtracks; and soundtracks with three callers triggered more mobbing response from the bird community than soundtracks with only one caller. However, when replicating the experiment in spring, we found a lower general response but also differences between playbacks, with increased responses only toward the 3 coal tits' playbacks. This interaction between context and acoustic cues demonstrates the flexible nature of heterospecific communication.

In winter, both the number of caller and emitter species influence mobbing responses

Birds usually modulates their mobbing responses depending on the threat they perceive. For example, different predators are mobbed with different levels of intensity (Curio et al. 1983, Templeton et al. 2005). Individuals can also change their mobbing response depending on the distance of the threat, the movement of the predator, or other cues surrounding the predator (Book & Freeberg 2015, Carlson et al. 2017). In this study, we recorded a higher mobbing response towards soundtracks with three individuals than towards soundtracks with only one individual calling. This result is congruent with the hypothesis that birds will use acoustic cues to gain information on the threat. Indeed, a larger number of birds may indicate a more significant predator, as larger mobs are produced in front of more important predators (Dutour et al. 2017b, Sandoval & Wilson 2012). In addition, joining a group instead of a lone caller increases the dilution effect, hence reducing risk for the

newcomer (Sridhar et al. 2009). Alternatively, the increased response to the playbacks with
more birds may be unrelated to an assessment of risk by birds, but rather be a simple
mechanical threshold reached when the call is more salient to receivers (by reaching a specific
threshold and/or being easier to detect). One solution to test the risk assessment hypothesis
could be to create a similar experiment but based only on visual cues. The idea would be to
test the mobbing response of birds in front of a predator model accompanied with either one
or three models of conspecifics. This kind of experiment should be done in large aviaries for
which we can control what visual cues the birds receive. If the focal bird approach and mob
more a when a group is already present, then the risk hypothesis would be more supported.

The mechanisms implicated in the differentiation between playbacks of one and three
callers can be various. In natural settings, birds can consider the number of spatially different
acoustic sources (Bradbury and Vehrencamp 2011). In our study, we launched the
soundtracks with only one loudspeaker whatever the treatment to suppress this effect.
Therefore, in our tests, the acoustic criteria that remain available are the duty cycle (i.e., the
proportion of the calling sequence when the signal is present), and the count of calling
individual through individual signatures. Our experiment does not add any insights on which
criteria was used by birds. Based on the current literature, the duty cycle is probably one
major coding strategy for increased risk in Parids (Landsborough et al. 2020, Salis et al.
2022), and Parids modify their response to unknown non-Parids calls with different duty
cycles (Dutour et al. 2022). Yet, great tits can also recognize caller identity, as they increased
their mobbing response toward soundtracks made with calls of several individuals compared
to soundtracks with only one individual calling (Dutour et al. 2021). In this latter experiment,

the duty cycles of the different treatments were strictly equal. This result was however not

replicated when testing the response to different number of heterospecifics (chaffinches

Fringilla coelebs, Dutour and Randler 2021). In our experiments, we believe that the

overlapping of the calls in the three birds treatments avoid the risk of interpreting these

treatments as only one bird calling intensely. Further experiments exploring the response of

each species to conspecific and heterospecific calls with controlled duty cycle may enlighten

whether individual recognition can also be used in heterospecific communication.

A mobbing response occurred more often when broadcasting coal tits' mobbing calls

compared to crested tits' calls, but also more birds responded to it. Unexpectedly, even

crested tits responded more to coal tits' mobbing calls than to calls from their own species.

Coal tits therefore appear to be listened to and heavily responded to, leading to larger (and

possibly more efficient) mobs. This is in line with the hypothesis that species from the same

community show different levels of reliability (Magrath et al. 2015). The notion of

"community informant" was developed for Parids in Carlson et al. (2020). They investigated

whether the birds possessed a reliable way of encoding predator information, and if several

heterospecifics relied on these calls. They showed that the great tit (*Parus major*) best fitted

the definition of community informant. The coal tit approached the definition, with only one

caveat: the dunnock (*Prunella modularis*) did not respond to it. As the authors suggested, the

lack of response from one species does not mean that other species from the community do

not respond to it (Carlson et al. 2020). Indeed, in our study, 14 species responded to coal tits'

soundtracks. In contrast, the crested tit did not meet any of the criteria set by Carlson and

colleagues. Coal tits appear therefore to be one important species regarding predator

information in the community, and this is congruent with their increased sensibility to

predation by pygmy owls (*Glaucidium passerinum*) in winter (Suhonen et al. 1993).

Replicating the experiment in spring: A lower general response

In winter, Parids living in temperate regions often flock with heterospecifics,

sometimes leading to impressive mobs (up to 20 birds in the present experiment). In

opposition, during the reproductive period (May-July), Parids nest and defend their territory

with intensity (Hinde 1952). For this reason, we first explore the mobbing response of birds in

winter, as this is the season in which interactions and cooperative mobbing with

heterospecifics makes more sense. However, we replicated the experiment in spring to

explore whether seasonal context of the experiment could impact our results. We did not test

the same birds and cannot control the changes in environment and community between the

first tests in winter and the replicate in spring. For these reasons, we did not statistically

compare the two seasons, but will nonetheless discuss the differences found between the

original experiment and the replication.

In spring, the number of birds mobbing to the four different types of playbacks was

lower than in winter and did not differ between playback types. We here propose that in

spring, when all birds defend their territory, the number of birds that can respond is restricted

to the neighbors. Moreover, in spring, aggressiveness toward conspecifics is high and may

therefore reduce the number of potential birds responding to mobbing calls. This

aggressiveness may also explain why not so many birds responded to conspecific mobbing

calls in spring (coal tits to coal tits and crested tits to crested tits).

Additionally, not only did fewer individuals respond in spring than in winter, but in

spring, the proportion of locations with any response was lower than in winter. This

difference must be taken with cautiousness, as the community and the density of the

populations may vary with the seasons: a decrease in mobbing response may simply be

related to fewer individuals in the territory. An order effect due to tests in winter being done

before the tests in spring is unlikely given the absence of order effect in our experiment at a

short time scale (see Sup. Mat. 2 For details). In addition, in spring, we were able to hear coal

tits singing at the 100 spots studied. We are therefore confident that, in spring, each spot could

have recorded one coal tit's mobbing response. This suggests that at least for the coal tit, the

response to conspecific and heterospecific mobbing calls decreases in spring. This result is

consistent with Dutour et al. (2019) who detected in Parids a higher mobbing response toward

heterospecific calls in winter compared to summer. The proximal reasons for such a decrease

can be various. Increased territoriality and aggression in spring may very well limit

cooperative communication, since the mobbing calls may resemble intra-specific

aggression/territoriality calls, leading to a lower relevance for heterospecifics. Other factors

such as decreased predator pressure in spring (Dutour et al. 2017b) could also result in a lower

investment in mobbing in spring. The ratio cost/benefits in responding to distanced mobbing

calls is therefore probably flexible through different times of the year. Given that most of

these factors are intercorrelated, determining which one is responsible for the difference in

mobbing is unfeasible in natural conditions.

478   **Replicating the experiment in spring: Almost no response to crested mobbing**

479   **calls**

480   In addition to a general lower mobbing response in spring, the differences between

481   treatments were also impacted by the season. Indeed, while we selected similar models for the

482   community, coal tits', and crested tits' response in winter (additive effect of number of caller

483   and emitter species), we found different models in spring. A general tendency was detected,

484   with only the playbacks with three coal tits triggering more response than the three other

485   playbacks. This suggests that the crested tit is not considered as informative in spring, even

486   when mobbing in groups, and unexpectedly, even to conspecifics. Several explanations can be

487   proposed. First, a group of three crested tits in spring may be too rare to bear meaning, as they

488   are in pairs and defending their nest. However, this hypothesis does not stand as this is also

489   the case for the coal tit, but that the difference between one and three callers still stands in

490   spring for this species. Alternatively, the contact with crested tits may be reduced in spring if

491   crested tits densities are lower during this season, hence decreasing learning opportunities for

492   heterospecifics. However, crested tits stay on the same territory throughout the year (Ekman

493   1979) making this hypothesis unlikely despite the fact that our experiments do not allow us to

494   formally rule out it. We rather suggest that this lack of mobbing response may emerge from

495   reduced reliability of the calls. To be efficient, an acoustic signal needs to be easily

496   distinguishable from other signals (Bradbury and Vehrencamp 2011). The song and mobbing

497   calls of the crested tits are extremely similar (Cramp and Perrins 1993, Hailman 1989). As

498   crested tits produce both songs and mobbing calls in spring, we can hypothesize that the

499   global vocal production of crested tits therefore becomes less reliable from an external

individual, hence leading to a decreased response to such calls. In contrast, the coal tit appears

to be reliable and responded to in both seasons. This result is consistent with Jiang et al.

(2020) who also found that between seasons differences in playback responses did not affect

the nuclear status of some particular species (in their case, David's Fulvetta *Alcippe davidi*).

The difference between the response of the bird community to coal tits' playbacks compared

to crested tits' playbacks may also be due to a higher aggressiveness from crested tits. Crested

tits are known to be more aggressive during spring (Campbell 1958), and crested tits, larger

than coal tits, have higher rank dominance status (Suhonen et al. 1993). We have, however,

little data on whether the heterospecific aggressiveness is higher than coal tits' aggressiveness

since dominance status is not necessarily linked to increased aggressiveness (Wilson 1992).

Finally, difference in nest predation may impact the reliability of the information produced,

but to our knowledge, nest predators are similar between Parid species (Cramp and Perrins

1993).

To conclude, birds from a community respond differently to acoustic situations with

varying emitter species and number of callers. The number of callers may be recognized

either with caller identity and/or changes in duty cycles. Those acoustic cues are not

responded to in the same way throughout the year, possibly because of changes in

territoriality and reliance on heterospecific calls. These results emphasize the importance of

seasons in studies investigating the complexity of heterospecific communication.

REFERENCES

Anderson DR, Burnham KP. 2002. Avoiding pitfalls when using information-theoretic

methods. J Wildl Manag. 66:912–918.

Betts MG, Hadley AS, Doran PJ. 2005. Avian Mobbing Response is Restricted by Territory

Boundaries : Experimental Evidence from Two Species of Forest Warblers. Ethology.

111:821–835. https://doi.org/10.1111/j.1439-0310.2005.01109.x

Book DL, Freeberg TM. 2015. Titmouse calling and foraging are affected by head and body

orientation of cat predator models and possible experience with real cats. Animal Cognition.

18:1155-1164. https://doi.org/10.1007/s10071-015-0888-7

Bradbury JW, Vehrencamp SL. 1998. Principles of animal communication. Sunderland, MA:

Sinauer Associates.

Brewer MJ, Butler A, Cooksley SL. 2016. The relative performance of AIC, AICC and BIC

in the presence of unobserved heterogeneity. Methods Ecol Evol. 7: 679–692.

Brooks ME, Kristensen K, van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ,

Maechler M, Bolker BM (2017). glmmTMB Balances Speed and Flexibility Among Packages

for Zero-inflated Generalized Linear Mixed Modeling. The R Journal. 9:378–400.

https://journal.r-project.org/archive/2017/RJ-2017-066/index.html

Campbell, B. 1958. The crested tit. Forestry commission leaflet, Her Majesty's stationery

office.

Carlson NV, Healy SD, Templeton CN. 2018. Mobbing. Cur Biol. 28:R1081–R1082.

doi:10.1016/j.cub.2018.06.025

Carlson NV, Healy SD, Templeton CN. 2020. What makes a 'community informant'?

Reliability and anti-predator signal eavesdropping across mixed-species flocks of tits. Anim

Behav Cogn. 7:214–246. https://doi.org/10.26451/abc.07.02.13.2020

Carlson NV, Griesser M. 2022. Mobbing in animals: A thorough review and proposed future

directions. Adv Study Behav. 54:1-41.

Carlson NV, Pargeter HM, Templeton CN. 2017. Sparrowhawk movement, calling, and

presence of dead conspecifics differentially impact blue tit (Cyanistes caeruleus) vocal and

behavioral mobbing responses. Behav Ecol Sociobiol. 71:1-11.

https://doi.org/10.1007/s00265-017-2361-x

Clucas BA, Freeberg TM., Lucas JR. 2004. Chick-a-dee call syntax, social context, and

season affect vocal responses of Carolina chickadees (Poecile carolinensis). Behav Ecol

Sociobiol, 57:187-196. https://doi.org/10.1007/s00265-004-0847-9

Cramp S, Perrins CM, Brooks DJ. 1993. Vol. VII: Flycatchers to shrikes. Oxford, UK:

Oxford University Press.

Curio E. 1978. The adaptive significance of avian mobbing: I Teleonomic hypotheses and

predictions. Z Tierpsychol. 48:175–183. https://doi.org/10.1111/j.1439-0310.1978.tb00254.x

Curio E, Klump G, Regelmann K. 1983. An anti-predator response in the great tit (Parus

major) : Is it tuned to predator risk? Oecologia, 60: 83-88.

https://doi.org/10.1007/BF00379324

Curio E, Regelmann K. 1986. Predator harassment implies a real deadly risk: A reply to

Hennessy Ethology. 72:75–78. https://doi.org/10.1111/j.1439-0310.1986.tb00607.x

Dutour M, Lena JP, Lengagne T. 2016. Mobbing behaviour varies according to predator

dangerousness and occurrence. Anim Behav. 119:119–124.

https://doi.org/10.1016/j.anbehav.2016.06.024

Dutour M, Léna JP, Lengagne T. 2017a. Mobbing calls: a signal transcending species

boundaries. Anim Behav. 131:3–11. https://doi.org/10.1016/j.anbehav.2017.07.004

Dutour M, Lena JP, Lengagne T. 2017b. Mobbing behaviour in a passerine community

increases with prevalence in predator diet. Ibis. 159:324–330.

https://doi.org/10.1111/ibi.12461

Dutour M, Cordonnier M, Léna JP, Lengagne T. 2019. Seasonal variation in mobbing

behaviour of passerine birds. J Ornithol. 160:509–514. https://doi.org/10.1007/s10336-019-

01630-5

Dutour M, Fernández GJ, Randler C. 2022. How great tits respond to urgency-based

information in allopatric Southern house wren mobbing calls. Ethology. 128: 676-683.

https://doi.org/10.1111/eth.13329

Dutour M, Kalb N, Salis A, Randler C. 2021. Number of callers may affect the response to

conspecific mobbing calls in great tits (*Parus major*). Behav Ecol Sociobiol. 75:1–8.

https://doi.org/10.1007/s00265-021-02969-7

Dutour M, Randler C. 2021. Mobbing responses of great tits (*Parus major*) do not depend on

the   number   of   heterospecific   callers.   Ethology.   127:379-384.

https://doi.org/10.1111/eth.13138

Ekman J. 1979. Coherence, composition and territories of winter social groups of the willow

tit *Parus montanus* and the crested tit *P cristatus*. Ornis Scand. 10:56–68.

https://doi.org/10.2307/3676345

Farine DR, Aplin LM, Sheldon BC, Hoppitt W. 2015. Interspecific social networks promote

information transmission in wild songbirds. Proc Royal Soc B. 282:20142804.

https://doi.org/10.1098/rspb.2014.2804

Feng CX. 2021. A comparison of zero-inflated and hurdle models for modeling zero-inflated

count data. J Stat Distrib App, 8: 8. https://doi.org/10.1186/s40488-021-00121-4

Flasskamp A. 1994. The adaptive significance of avian mobbing V An experimental test of

the 'move on'hypothesis. Ethology. 96:322–333. https://doi.org/10.1111/j.1439-

0310.1994.tb01020.x

Foster WA, Treherne JE. 1981. Evidence for the dilution effect in the selfish herd from fish

predation on a marine insect. Nature. 293:466–467. https://doi.org/10.1038/293466a0

Goodale E, Kotagama SW. 2005. Testing the roles of species in mixed-species bird flocks of

a Sri Lankan rain forest. J Trop Ecol. 21:669–676.

https://doi.org/10.1017/S0266467405002609

Goodale E, Ding P, Liu X, Martínez A, Si X, Walters M, et al. 2015. The structure of mixed-

species bird flocks, and their response to anthropogenic disturbance, with special reference to

East Asia. Avian Res. 6:14. https://doi.org/10.1186/s40657-015-0023-0

Hailman JP. 1989. The organization of major vocalizations in the Paridae. Wil Bull. 101:305–

343.

Hardin JW, Hilbe JM. 2007. Generalized linear models and extensions. TX, USA: Stata press.

Hartig F, Hartig MF. 2017. Package 'DHARMa'. Vienna, Austria: R Development Core

Team.

Hendrichsen D, Christiansen PK, Nielsen E, Dabelsteen T, Sunde P. 2006. Exposure affects

the risk of an owl being mobbed–experimental evidence. J Avian Biol. 37:13–18.

https://doi.org/10.1111/j.2005.0908-8857.03658.x

Hinde, RA. 1952. The behaviour of the great tit (*Parus major*) and some other related species.

Behav Sup. III:1–201.

Hua F, Yong DL, Janra MN, Fitri LM, Prawiradilaga D, Sieving KE. 2016. Functional traits

determine heterospecific use of risk-related social information in forest birds of tropical

South-East Asia. Ecol Evol. 6:8485–8494. https://doi.org/10.1002/ece3.2545

Jennions MD. 2003. A survey of the statistical power of research in behavioral ecology and

animal behavior. Behav Ecol; 14:438-445. https://doi.org/10.1093/beheco/14.3.438

Jiang D, Sieving KE, Meaux E, Goodale E. 2020. Seasonal changes in mixed-species bird

flocks   and   antipredator   information.   Ecol   Evol.   10:5368–5382.

https://doi.org/10.1002/ece3.6280

Kostan KM. 2002. The evolution of mutualistic interspecific communication: Assessment and

management across species. J Comp Psychol. 116:206–9. https://doi.org/10.1037/0735-

7036.116.2.206

Krams I, Bērziņš A, Krama T, Wheatcroft D, Igaune K, Rantala MJ. 2010. The increased risk

of predation enhances cooperation. Proc Royal Soc B. 277:513–518.

https://doi.org/10.1098/rspb.2009.1614

Kroodsma DE. 1989. Suggested experimental designs for song playbacks. Anim Behav.

37:600–609. https://doi.org/10.1016/0003-3472(89)90039-0

Landsborough B, Wilson DR, Mennill DJ. 2020. Variation in chick-a-dee call sequences, not

in the fine structure of chick-a-dee calls, influences mobbing behaviour in mixed-species

flocks. Behav Ecol. 31:54–62. https://doi.org/10.1093/beheco/arz155

Magrath RD, Haff TM, Fallow PM, Radford AN. 2015. Eavesdropping on heterospecific

alarm calls: from mechanisms to consequences. Biol Rev. 90:560–586.

https://doi.org/10.1111/brv.12122

Morton ES. 1975. Ecological sources of selection on avian sounds. Am Nat. 109:17–34.

Poian A, Yorke IM. 1989. Predator Harassment : More Evidence on the Deadly Risk.

Ethology. 83:167–169. https://doi.org/10.1111/j.1439-0310.1989.tb00526.x

R Core Team, 2022. R: A language and environment for statistical computing. Vienna,

Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Salis A, Léna JP, Lengagne T. 2022. Data & Supplemental information for manuscript

"Acoustic cues and season affect mobbing responses in a bird community "_ Review 1 PCI

(Version 3). Zenodo. https://doi.org/10.5281/zenodo.7605316

Salis A, Léna JP, Lengagne T. 2021. Great tits (*Parus major*) adequately respond to both

allopatric combinatorial mobbing calls and their isolated parts. Ethology. 127:213–222.

https://doi.org/10.1111/eth.13111

Salis A, Léna JP, Lengagne T. 2022. Which acoustic parameters modify the great tit's

response to conspecific combinatorial mobbing calls? Behav Ecol Sociobiol, 76:46.

https://doi.org/10.1007/s00265-022-03157-x

Sandoval L, Wilson DR. 2012. Local predation pressure predicts the strength of mobbing

responses in tropical birds. Cur Zool. 58:781-790. https://doi.org/10.1093/czoolo/58.5.781

Sordahl TA. 1990. The Risks of Avian Mobbing and Distraction Behavior: An Anecdotal

Review. Wilson Bull. 102:349–352.

Sridhar H, Beauchamp G, Shanker K. 2009. Why do birds participate in mixed-species

foraging   flocks?   A   large-scale   synthesis.   Anim   Behav. 78:337–347.

https://doi.org/10.1016/j.anbehav.2009.05.008

Suhonen J, Halonen M, Mappes T. 1993. Predation Risk and the Organization of the Parus

Guild. Oikos. 66: 94–100. https://doi.org/10.2307/3545200

Suzuki TN, Wheatcroft D, Griesser M. 2016. Experimental evidence for compositional syntax

in bird calls. Nat Com. 7:10986. https://doi.org/10.1038/ncomms10986

Templeton CN, Zollinger SA, Brumm H. 2016. Traffic noise drowns out great tit alarm

calls. Cur Biol. 26:R1173-R1174. https://doi.org/10.1016/j.cub.2016.09.058

Templeton CN, Greene E, Davis K. 2005. Allometry of alarm calls: black-capped chickadees

encode   information   about   predator   size. Science. 308:1934–1937.

https://doi.org/10.1126/science.1108841

Wheatcroft D, Price TD. 2018. Collective action promoted by key individuals. Am Nat.

192:401–414. http://dx.doi.org/10.1086/698874

Wilson J. D. 1992. Correlates of agonistic display by great tits *Parus major*. Behaviour. 121:

168–214.

Woods RD, Kings M, McIvor GE, Thornton A. 2018. Caller characteristics influence

recruitment   to   collective   anti-predator   events   in   jackdaws.   Sci   Rep.   8:1–8.

https://doi.org/10.1038/s41598-018-25793-y

Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. 2009. Mixed effects models and

extensions in ecology with R. New York: Springer.