Parental care allows genetic variation to accumulate by relaxing selection

5 Sonia Pascoal[1], Rahia Mashoodh[1], Hideyasu Shimadzu[2,3], Rebecca M. Kilner[1*]

 [1]Department of Zoology, University of Cambridge, Cambridge, CB2 3EJ, U.K.

 [2]Department of Mathematical Sciences, Loughborough University, Leicestershire, LE11 3TU,

 UK

 [3]Graduate School of Public Health, Teikyo University, Tokyo, Japan

 *Corresponding author: rmk1002@cam.ac.uk

 Keywords: social evolution, mutation load, extinction rate, *Nicrophorus vespilloides,* standing

 genetic variation, parent-offspring

Benign social environments are thought to relax selection against deleterious mutations, enabling them to accumulate in the genome. However, it has not proved straightforward to demonstrate causality in this relationship. We allowed burying beetle populations to evolve with and without the benign effects of parental care for over 20 generations, before inbreeding lineages that were seeded from these experimental populations to expose any differences in their mutation load. We found that inbred lineages with a history of parental care went extinct most quickly, indicating that they harboured more recessive deleterious alleles.

Classical population genetics models imagine that populations attain an equilibrium level of genetic variation, at which the rate of input of new genetic variants is balanced by the rate of their elimination by natural selection (known as mutation-selection balance[1-4]). The concept of mutation-selection balance has long been used to understand the effects of genetic mutation rate on the health of human populations, partly because it is recognised that humans can modify their own environment and so change the forces of natural selection to which they are exposed[2-4]. Benevolent social activities, such as the widely available provision of medical care, are suggested to have been particularly influential in preventing natural selection from purging deleterious mutations[2-4]. Consistent with this suggestion, recent comparative genomic analyses have revealed a greater incidence of genetic pathologies in western industrialised populations than in more traditional human societies[3-7] - though it is impossible to demonstrate that benevolent social behaviours have caused this difference.

Benevolent social behaviours are relatively commonplace in other animals too, where they are more formally defined as acts of cooperation[8]. Here the effect on mutation-selection balance has received relatively little empirical attention[9], though similar effects are expected in theory[1]. Parental care is a widespread example of cooperation[10]. By building protective

nests, defending their brood from attack and nourishing them, animal parents shield their

young from environmental stressors[11] and weaken the correlation between the phenotypic

variation seen by selection and the underlying genetic variation[1,12]. In these ways, parents

relax selection on the offspring phenotype[1, 11-14], theoretically allowing mildly deleterious

alleles to accumulate[1,11]. Previous experimental work with insects has shown that parental

care enables new mutants[11] and inbred offspring[8] to survive for one generation. However, it is

not yet known whether parental care also enables mildly deleterious mutations to accumulate

and persist over multiple generations.

We tested this suggestion by evolving replicate laboratory populations of burying

beetles *Nicrophorus vespilloides* under sharply contrasting levels of parental care, for more

than 20 generations. Burying beetles breed on the body of a small dead vertebrate[15], which the

parents jointly convert into a carrion nest by removing the fur or feathers, rolling the flesh

into a ball, covering it with anti-microbial anal exudates, and burying it. This is pre-hatching

parental care[16]. Parents also guard and feed larvae after hatching, though larvae can survive in

the lab with no post-hatching care at all[17]. In two of our evolving populations, larvae were

able to receive both pre-hatching and post-hatching parental care (these were called the 'Full

Care' lines) while in two other populations we prevented parents from supplying any post-

hatching care by removing them before the larvae hatched, after the carrion nest was complete

(these were called the 'No Care' lines). During the first 20 or so generations of experimental

evolution, No Care lines rapidly adapted to a life without parental care[18], through divergent

phenotypic change in both larval[19] and parental traits[16].

To determine whether parental care causes deleterious genetic variation to accumulate

over the generations, we inbred sub-populations, each derived from the replicate experimental

evolving populations, for 8 successive generations (we called this The Evolutionary History

Experiment). For these 8 generations, we measured the extent to which inbreeding reduced

measures of reproductive success in comparison with control outbred populations. To

determine whether parental care could temper the rate of extinction (as implied by [9]), in half

of all our treatments parents were allowed to provide care after their offspring hatched, while

in the remainder they were prevented from supplying post-hatching care. This generated 8

different treatments in total (see Figure 1 in Supplementary Information, for the design of the

Evolutionary History Experiment).

We used the data from the Evolutionary History Experiment, to test three predictions:

(1) that the harsh conditions of the No Care environment eliminates any genetic variants that

are not adapted to survive under these conditions. (2) That the more benign conditions of the

Full Care environment relaxes selection and promotes the survival of more genetic variants.

(3) That inbred populations from the Full Care lines should exhibit greater inbreeding

depression than inbred populations from the No Care lines (due to a greater accumulation of

deleterious recessive mutations under relaxed selection). The outbred populations acted as a

control treatment for tests of all three predictions.

To test predictions (1) and (2) we initially focused on the data collected from the first

generation of breeding in the Evolutionary History Experiment. The results supported both

predictions in showing that the No Care is harsher than the Full Care environment, and that a

Full Care environment can relax selection against genetic variants that would otherwise be

eliminated in the harsher No Care environment. We found that inbreeding reduced

reproductive success, regardless of the evolutionary history of the lineage (Figure 1,

Supplementary Table 1). However, a supply of post-hatching care enabled more broods to

survive, even if they were inbred - replicating the findings of [9]. In addition, we found that

families that were allowed to provide care for their offspring were more likely to produce

surviving offspring, whether or not the parents were inbred, and regardless of the evolutionary

history of their lineage (Figure 1, Supplementary Table 1).

To test prediction (3), we continued to examine inbred families in the first generation of breeding in the Evolutionary History Experiment. We found that the chance of brood survival depended on the levels of current care supplied as well as on the evolutionary history of the lineage (Supplementary Table 2). To examine the effect of evolutionary history in more detail, we split the dataset by the current level of care supplied. In support of prediction (3), inbred families derived from the Full Care populations had lower brood survival than families drawn from the No Care populations (log(OR) =1.12 [0.49-1.80], z=3.42, $p<0.001$) – in a No Care current environment. No equivalent differences were observed in the Full Care current environment (log(OR)=0.20 [-1.4,1.9], z=0.25, $p=0.80$). For the outbred families, the evolutionary history of the lineage had no effect on breeding success, though broods were in general less successful when they received no post-hatching care (Supplementary Table 2).

To further test prediction (3), we calculated the extent of inbreeding depression at each generation. Inbreeding depression was greater in Generation 1 for families descended from the Full Care evolving populations than the No Care evolving populations, in the No Care current environment. We found the same pattern in the Full Care current environment – though here the differences between lineages were seen at Generation 3 (Figure 2A).

Finally, we compared the survival of all lines in the Evolutionary History Experiment, by fitting multivariate Cox proportional hazard models (Figure 2B). A lineage was considered extinct if none of its members survived to reproduce in the subsequent generation. Whilst all inbred lineages in our experiments eventually went extinct, outbred lineages were still reproducing successfully at the point at which the experiment was terminated (Supplementary Figure 3A, B). For the inbred lineages a No Care current environment resulted in particularly rapid extinction (Figure 2B). Of these, lineages seeded from the Full Care evolving populations had significant lower median survival than inbred lineages seeded from the No Care evolving populations (non-parametric Kruskal Wallis test: $H(1)=4.59$, $p=0.03$). The

lineages seeded from the Full Care evolving populations reached 50% extinction one

generation sooner than those seeded from the No Care evolving populations (Supplementary

Table 3). We could detect no equivalent difference in lineage survival between the evolved

populations in the Full Care current environment ($H(1)=0.06$, $p=0.81$).

Together, these analyses support all three of our predictions, and demonstrate that the

benign social environment created by parental care can weaken selection sufficiently to

enable the accumulation of genetic mutations within populations. In the short-term, such

genetic mutants are overwhelmingly likely to be recessive and mildly deleterious [1-7]. It would

be interesting to determine in future work whether this sets in motion a positive evolutionary

feedback loop between the extent of cooperation and mutation load, due to a greater reliance

on cooperative behaviour to compensate for any genetic deficiencies. In the longer-term

though, benevolent social interactions could build up genetic diversity that might turn out to

be beneficial and cause rapid evolution, especially if environmental conditions change

suddenly, or if mutations promote novelty through compensatory evolution[20].

METHODS

***Nicrophorus vespilloides* natural history.** The common burying beetle breeds on a small

dead vertebrate (like a songbird or mouse). The larvae hatch from eggs laid nearby in the soil

and crawl to their carrion nest, which they can feed upon themselves[15]. Once at the carcass,

larvae receive post-hatching biparental care. Parents supply fluids to their offspring through

oral trophallaxis, and defend their brood and the carrion nest from attack by predators,

microbes and rival beetles[15]. The duration and extent of post-hatching care are highly

variable, however. For example, when wild beetles are brought into the lab to breed, roughly

5% of larvae receive no post-hatching care at all, yet larvae can still survive to become

reproductively competent adults (see[19] and references therein). Within roughly a week of

hatching, the larvae complete development and at this point (which we refer to as 'dispersal),

they start to crawl away from the scant remains of the carcass to pupate in the soil. The

parents, meanwhile, fly off in search of a new carcass.

Experimental evolution. The experimental lines used in this work have been described in

detail elsewhere[16, 19]. In brief, we established a large founding population of *N. vespilloides* by

interbreeding wild-caught individuals from four different woodlands. This was then divided

into four experimental lines. In two lines, larvae experienced 'Full Care' at each generation,

with both parents staying in the breeding box throughout the breeding bout and able to

provide post-hatching care as well as pre-hatching care. In the other two 'No Care' lines,

parents engaged in pre-hatching care but at each generation they were then removed from the

breeding box, around 53 hours after they were paired, so that they never interacted with their

larvae. The work reported here began when these lines had been exposed to 20 generations of

experimental evolution under these contrasting regimes of care.

Evolutionary History Experiment (see Supplementary Figure 1 for overview of design).

Preparatory common garden generation The experiment began by exposing individuals

drawn from the four lines (Full Care replicated twice and No Care replicated twice) to a

common garden Full Care environment for one generation (N = 60 pairs for each No Care line

(to counter-balance the slightly lower breeding success caused by the No Care environment)

and N = 50 pairs for each Full Care line). In this way, we minimised any potentially

confounding transgenerational effects prior to starting the Evolutionary History Experiment.

Overview. Broods from the common garden generation seeded lineages in the experimental

treatments: broods derived from the Full Care lines founded lineages that were Inbred or

Outbred, in either a Full Care or No Care environment, and the same was true for broods

derived from the No Care lines. Thus, for each line of origin, individuals in the different

treatments came from a similar genetic pool. From Generation 1 onwards, half of the beetles

drawn from each line were exposed to continuous inbreeding (full-sibling crosses) for up to 8

generations (by which point all the inbred lineages had gone extinct) (N = c. 45 crosses per

treatment at Generation 1. The remaining beetles were outbred in identical conditions to

provide a control baseline for comparison with the inbred lineages (N = c. 35-40 crosses per

treatment, per generation). Half of all inbred lineages, and half of the outbreeding

populations, were allowed to provide post-hatching care for their young (Full Care

environment), while the remaining beetles were only allowed to provide pre-hatching care

(No Care environment). The experiment therefore had a 2 x 2 x 2 design, with 8 treatments in

all (Full Care versus No Care line of origin; Inbred versus Outbred; Full Care environment

versus No Care environment), with each treatment replicated twice due to replicate Full Care

and No Care lines (Supplementary Figure 1).

Detailed methods: Beetle maintenance was carried out following standard protocols [18].

Briefly, adult beetles were kept individually in plastic boxes (12 x 8 x 6cm) filled with moist

soil and fed twice a week with raw beef mince. Adults were bred at 2-3 weeks post-eclosion

in a breeding box (17 x 12 x 6cm) with soil and a mouse carcass (11-13 g for all treatments

except for the individuals derived from the Full Care lines, that were outbred under Full Care

conditions (8-14 g)). To ease the considerable burden of work, data for broods in this

treatment were collected from the ongoing experimental evolution lines in the laboratory, and

so received mice in the size range assigned to that treatment for breeding. Carcass size was

included, where appropriate, as a factor in the statistical analyses (see below).

For the inbreeding treatments, we paired full siblings (one pair per family) whereas for the

outbreeding treatments we paired males and females at random and did not pair siblings or

cousins. Each pair was given a breeding box with a dead mouse sitting on soil, and the

breeding boxes were placed in a dark cupboard to simulate natural underground conditions.

For broods assigned to a No Care environment, parents were removed around 53 hours after

pairing. Eight days after pairing (which is when the larvae have completed their development

and start to disperse away from the carcass) we scored two standardly used measures of

reproductive success in burying beetles[16]: brood success (fail = no larvae produced; success =

some larvae produced) and brood size at dispersal. Larvae were then placed into cells (2 x 2 x

2cm) in an eclosion box (10 x 10 x 2cm), with one eclosion box per brood, which was filled

with soil until larvae had developed into sexually immature adults (about 18 days after

dispersal). At this point, adults were transferred to individual boxes until they reached sexual

maturity roughly 2 weeks later. Both the eclosion boxes and the individual boxes were kept

on shelves in the laboratory at 21°C on a 16L:8D hour light cycle.

Statistical Analyses

All statistical tests were conducted in R version 3.5.1[21]. Data handling and visualisation were

carried out using the 'tidyverse'[24] and 'survminer'[23] R packages. All data and code presented

in the manuscript is available through: https://github.com/r-mashoodh/nves_MutationLoad.

Testing predictions (1) and (2)

To test predictions (1) and (2) we focused on the data collected from Generation 1. Using a

binomial generalised linear model (GLM) in the base 'statistics' package in R, we tested the

effect of evolutionary history (*i.e.* derived from a No Care evolving population or from a Full

Care evolving population), current care environment (*i.e.* experienced No Care or Full Care

during Generation 1), and inbreeding (*i.e.* inbred or outbred) on brood success. We defined

brood success at dispersal in the following way: broods that produced at least one surviving

larva were defined as successful (following [16,18]) whereas those that did not produce any

surviving young were classified as failures.

We subsequently ran analyses separately for the inbreeding and outbreeding conditions to

examine any interactions between evolutionary history (*i.e.* derived from a No Care evolving

population or from a Full Care evolving population) and the current environment (*i.e.*

experienced No Care or Full Care during Generation 1) dropping non-significant interaction

terms where appropriate. We included block and carcass weight as covariates to ensure any

effects we detected occurred over and above any variation in these variables.

Testing prediction (3)

Calculation of inbreeding depression.

For direct comparison with [9], we calculated the inbreeding depression rate: $\delta = (w_o -$

$w_i)/w_o$, where w_o and w_i are respectively the number of surviving outbred and inbred

offspring at dispersal. We combined data from both blocks to derive the measures shown in

Figure 2A.

Survival analysis across generations

To determine the effect of evolutionary history (*i.e.* derived from a No Care evolving

population versus a Full Care evolving population), current care environment (*i.e.*

experienced No Care or Full Care) on the survival of the different lineages in the

Evolutionary History Experiment (Supplementary Figure 1), we fit multivariate Cox

proportional hazard models using the 'survival' R package [24], again adding carcass weight

and block as covariates. A lineage was considered to be extinct if it did not survive to

reproduce in the subsequent generation. The outbred lineages did not go extinct and so were

analysed separately from the inbred lineages. We additionally used the non-parametric

Kruskal Wallis test to determine if median survival times of each inbred lineage differed, by

comparing the effect of evolutionary history (*i.e.* derived from a No Care evolving population

or from a Full Care evolving population) in separate analyses, one for each current care

environment (No Care versus Full Care). Model diagnostics were checked visually and using

the cox.zph function.

Figures

Figure 1

Figure 1. Breeding success of evolved populations in Generation 1 of the Evolutionary

History Experiment. Predicted means ± S.E survival probabilities of populations that have

evolved under Full Care (red lines and datapoints, FC_{POP}) or No Care (blue lines and

datapoints NC_{POP}) are shown, based on the current care environment (Full Care (FC) *vs* No

Care (NC)) under inbred (left) and outbred (right) conditions.

Figure 2

Figure 2. Survival of inbred populations across generations. A) inbreeding depression

values for population lineages from the Evolutionary History Experiment, shown in their

native ($FC_{POP}FC_{ENV}$ and $NC_{POP}NC_{ENV}$) and reciprocal current environments ($FC_{POP}NC_{ENV}$

and $NC_{POP}FC_{ENV}$). (B) Survival curves for inbred population lineages from the Evolutionary

History Experiment and associated 95% confidence intervals for the inbred lines.

REFERENCES

1. Linksvayer, T. A. & Wade, M. J. Genes with social effects are expected to harbor more
sequence variation within and between species. *Evolution* **63,** 1685-1696 (2009).

2. Wright, S. On the number of self-incompatibility alleles maintained in equilibrium by a
given mutation rate in a population of a given size: a reexamination. *Biometrics* **16**,
61-85 (1960).

3. Lynch, M. Mutation and human exceptionalism: Our future genetic load. *Genetics* **202**,
869-875. (2016).

4. Kondrashov, A. *Crumbling Genome: The impact of deleterious mutations on human*
populations. (Wiley Online Library 2017).

5. You, W & Henneberg, H. Cancer incidence increasing globally: The role of relaxed natural
selection. Evolutionary Applications **11,** 140-152 (2017).

6. Aris-Brosou, S. Direct evidence of an increasing mutational load in humans. *Mol Biol Evol*
36, 2823-2829 (2019).

7. Saniotis, A., Hennberg, M. & Mohammadi, K. Genetic load and biological changes to
extant humans. *Journal of Biosocial Science* **53,** 639-642. (2021).

8. Davies, N. B., Krebs, J. R. & West, S. A. *An Introduction to Behavioural Ecology* (Wiley-
Blackwell, 2012).

9. Pilakouta, N., Jamieson, S., Moorad J. A. & Smiseth, P. T. Parental care buffers against
inbreeding depression in burying beetles. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 8031-8035
(2015).

10. Royle, N. J., Smiseth, P. T. & Kölliker, M. *The Evolution of Parental Care* (Oxford
University Press, 2012).

11. Snell-Rood, E. C., Burger, M., Hutton, Q. & Moczek, A. P. Effects of parental care on the
accumulation and release of cryptic genetic variation: review of mechanisms and a
case study of dung beetles. *Evolutionary Ecology.* **30**, 251-265 (2016).

12. Wade, M. J. in Mousseau, T. & Fox, C. *Maternal Effects as Adaptations* (Oxford
University Press, 1998).

13. Whitlock, M. C. The red queen beats the jack-of-all trades: the limitations on the
evolution phenotypic plasticity and niche breadth. *American Naturalist* **148,** S65-S77.
(1996).

14. Van Dyken, J. D. & Wade, M. J. The genetic signature of conditional expression.
Genetics. **184**, 557–570 (2010).

15. Scott, M. P. The ecology and behavior of burying beetles. *Annu. Rev. Entomol.* **43**, 595–
618 (1998).

16. Duarte, A., Rebar, D., Hallett, A. C., Jarrett, B. J. M. & Kilner, R. M. Evolutionary
change in the construction of the nursery environment when parents are prevented
from caring for their young directly. *Proc. Natl. Acad. Sci. U.S.A.* **48,** e2102450118
(2021).

17. A. K. Eggert, M. Reinking, J. K Müller. Parental care improves offspring survival and
growth in burying beetles. *Anim. Behav*. **55**, 97–107 (1998).

18. M. Schrader, B. J. M. Jarrett, R. M. Kilner. Using experimental evolution to study
adaptations for life within the family. *Am. Nat.* **185**, 610-619 (2015).

19. D. Rebar, N. W. Bailey, B. J. M. Jarrett, R. M. Kilner. An evolutionary switch from
sibling rivalry to sibling cooperation, caused by a sustained loss of parental care. *Proc.*
Natl. Acad. Sci. U.S.A. **117**, 2544-2550 (2020).

20. Farkas, Z., Kovács, K., Sarkadi, Z., Kalapis, D., Fekete, G., Birtyik, F., Ayaydin, F.,
Molnár, C., Horváth, P., Pál, C. & Papp, B. Gene loss and compensatory evolution
promotes the emergence of morphological novelties in budding yeast. *Nature Ecol*
Evol https://doi.org/10.1038/s41559-022-01730-1 (2022)

21. R Core Team. R: A Language and Environment for Statistical Computing [Internet].
Vienna, Austria: R Foundation for Statistical Computing; 2019. Available from:
https://www.R-project.org/

22. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to
the Tidyverse. JOSS. 2019 Nov 21;4(43):1686.

23. Alboukadel Kassambara, Marcin Kosinski and Przemyslaw Biecek (2021). survminer:
Drawing Survival Curves using 'ggplot2'. R package version 0.4.9. https://CRAN.R-
project.org/package=survminer

24. Therneau T (2021). _A Package for Survival Analysis in R_. R package version 3.2-13,
<URL: https://CRAN.R-project.org/package=survival>.

ACKNOWLEDGEMENTS:

We thank Sue Aspinall and Chris Swannack for helping with beetle maintenance. We also

thank Benjamin Jarrett and Darren Rebar for collecting data from the experimental evolution.

Funding: This project was supported by a Consolidator's Grant from the European Research

Council (310785 Baldwinian_Beetles), by a Wolfson Merit Award from the Royal Society,

The Leverhulme Trust (RPG-2018-232) and The Isaac Newton Trust (18.23(q)), each to

RMK. RM was supported by a Biotechnology and Biological Sciences Research Council

Future Leaders Fellowship (BB/R01115X/1). HS was partially supported by the Japan Society

for the Promotion of Science (KAKENHI Grant Number: JP19K21569).

Authors contribution: Conceived the idea and designed the Evolutionary History

Experiment: RMK, SP; Performed the Evolutionary History Experiment: SP; Collected data

from Experimentally evolving populations: RM; Analysed the data: RM, HS; Wrote the

manuscript: RMK, RM, SP. All authors discussed the results and commented on the

manuscript.

Competing interests: The authors declare no competing financial or nonfinancial interests.

Data and materials availability: Data and code needed to evaluate the conclusions in the

paper are available from https://github.com/r-mashoodh/nves_MutationLoad.

SUPPLEMENTARY FIGURES AND TABLES

Supplementary Figure 1. Overview of the Evolutionary History Experiment. Beetles that

had evolved in the Full Care (FC) or No Care (NC) lines for 20 generations were used to seed

new experimental populations in the lab. Before inbreeding began, all beetles from these lines

experienced one generation of a FC common garden environment to minimise potentially

confounding transgenerational effects. Sequential inbreeding or outbreeding was then applied

for up to eight generations under both FC and NC environments. n: number of families per

population in generation one of inbreeding (*i.e.* the generation immediately after the common

garden FC generation). The design includes two replicate populations organised into Blocks

(Block 1 and Block 2) whose breeding was staggered by one week (to ease the workload of

maintaining them). Grey boxes: data for these two populations were collected from the

evolving populations.

Supplementary Figure 2. Survival analysis in inbred lineages. Forest plots of Cox

proportional hazard survival model estimates (with 95% confidence intervals) for CurrentEnv

(*i.e.* whether families experienced No Care or Full Care in the current generation) and

Evolved history (*i.e.* whether families were derived from the No Care or Full Care evolving

populations). Block was included as a covariate for inbred lineages (see Supplementary

Figure 1 for experimental design).

Supplementary Figure 3. Survival of outbred lineages across generations. (A) Survival

curves for outbred population lineages from the Evolutionary History Experiment shown in

their native ($FC_{POP}FC_{ENV}$ and $NC_{POP}NC_{ENV}$) and reciprocal current environments

($FC_{POP}NC_{ENV}$ and $NC_{POP}FC_{ENV}$) and associated 95% confidence intervals (FC: Full Care; NC:

No Care). (B) Forest plots of Cox proportional hazard survival model estimates (with 95%

confidence intervals) for CurrentEnv (*i.e.* whether families experienced No Care or Full Care

in the current generation) and Evolved (*i.e.* whether families were derived from the No Care

or Full Care evolving populations). Block was included as a covariate for inbred lineages (see

Supplementary Figure 1 for experimental design).

Supplementary Table 1. Summary of binomial generalised linear model estimates for brood

success in Generation 1 of the Evolutionary History Experiment, predicted by Evolution

History (*i.e.* whether families were derived from the No Care or Full Care evolving

populations), Current Environment (*i.e.* whether families experienced No Care or Full Care in

the current generation) and Breeding Condition (*i.e.* whether families were inbred or outbred).

Carcass weight and Block were included as a covariate for inbred lineages (see

Supplementary Figure 1 for experimental design).

Factor	log(OR)[1]	95% CI[1]	p-value
Evolution History			
Full Care	—	—	
No Care	-0.06	-1.2, 1.1	>0.9
Current Environment			
Full Care	—	—	
No Care	**-3.0**	**-3.9, -2.2**	**<0.001**
Breeding Condition			
Inbred	—	—	
Outbred	**0.82**	**0.37, 1.3**	**<0.001**
Block	-0.18	-0.63, 0.26	0.4
Carcass.weight	0.30	-0.02, 0.63	0.068
Evolved * CurrentEnv			
N * N	0.80	-0.46, 2.0	0.2

[1]OR = Adjusted Odds Ratio, CI = Confidence Interval

Supplementary Table 2 Summary of binomial generalised linear model estimates for brood

success in Generation 1 of the Evolutionary History Experiment, predicted by Evolutionary

History (*i.e.* whether families were derived from the No Care or Full Care evolving

populations) and Current Environment (*i.e.* whether families experienced No Care or Full

Care in the current generation). Carcass weight and Block were included as covariates (see

Supplementary Figure 1 for experimental design). Separate models were run for Inbred and

Outbred families.

	Inbred			Outbred		
Characteristic	log(OR)[1]	95% CI[1]	p-value	log(OR)[1]	95% CI[1]	p-value
Evolutionary History						
Full Care	—	—		—	—	
No Care	**1.0**	**0.39, 1.6**	**0.001**	0.10	-0.61, 0.81	0.8
Current Environment						
Full Care	—	—		—	—	
No Care	**-3.0**	**-3.9, -2.3**	**<0.001**	**-2.0**	**-3.1, -1.1**	**<0.001**
Block	-0.48	-1.1, 0.10	0.11	0.29	-0.41, 1.0	0.4
Carcass.weight	0.28	-0.18, 0.74	0.2	0.13	-0.32, 0.59	0.6

[1]OR = "Adjusted" Odds Ratio, CI = Confidence Interval

404 **Supplementary Table 3** Median survival times of each lineage (measured in generations) by
405 Breeding Condition (Inbred versus Outbred), Evolutionary History (*i.e.* whether families were
406 derived from the No Care or Full Care evolving populations) and Current Environment (*i.e.*
407 whether families experienced No Care or Full Care in the current generation) with 95%
408 confidence intervals (CI). (NA = populations did not go extinct).

Breeding Condition	Evolutionary History	Current Environment	median	lower CI	upper CI
Inbred	Full Care	Full Care	3	3	3
Inbred	Full Care	No Care	1	1	1
Inbred	No Care	Full Care	3	3	3
Inbred	No Care	No Care	2	1	2
Outbred	Full Care	Full Care	NA	NA	NA
Outbred	Full Care	No Care	NA	NA	NA
Outbred	No Care	Full Care	NA	NA	NA
Outbred	No Care	No Care	NA	NA	NA